



Calving fronts and where to find them: a benchmark dataset and methodology for automatic glacier calving front extraction from synthetic aperture radar imagery

Nora Gourmelon¹, Thorsten Seehaus², Matthias Braun², Andreas Maier¹, and Vincent Christlein¹

¹Pattern Recognition Lab, Computer Science Department, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany

²Institute of Geography, Department of Geography and Geosciences, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany

Correspondence: Nora Gourmelon (nora.gourmelon@fau.de)

Received: 25 April 2022 – Discussion started: 2 May 2022

Revised: 11 July 2022 – Accepted: 23 July 2022 – Published: 22 September 2022

Abstract. Exact information on the calving front positions of marine- or lake-terminating glaciers is a fundamental glacier variable for analyzing ongoing glacier change processes and assessing other variables like frontal ablation rates. In recent years, researchers started implementing algorithms that can automatically detect the calving fronts on satellite imagery. Most studies use optical images, as calving fronts are often easy to distinguish in these images due to the sufficient spatial resolution and the presence of different spectral bands, allowing the separation of ice features. However, detecting calving fronts on synthetic aperture radar (SAR) images is highly desirable, as SAR images can also be acquired during the polar night and are independent of weather conditions (e.g., cloud cover), facilitating year-round monitoring worldwide. In this paper, we present a benchmark dataset (Gourmelon et al., 2022b) of SAR images from multiple regions of the globe with corresponding manually defined labels providing information on the position of the calving front (<https://doi.org/10.1594/PANGAEA.940950>). With this dataset, different approaches for the detection of glacier calving fronts can be implemented, tested, and their performance fairly compared so that the most effective approach can be determined. The dataset consists of 681 samples, making it large enough to train deep learning segmentation models. It is the first dataset to provide long-term glacier calving front information from multi-mission data. As the dataset includes glaciers from Antarctica, Greenland, and Alaska, the wide applicability of models trained and tested on this dataset is ensured. The test set is independent of the training set so that the generalization capabilities of the models can be evaluated. We provide two sets of labels: one binary segmentation label to discern the calving front from the background, and one label for multi-class segmentation of different landscape classes. Unlike other calving front datasets, the presented dataset contains not only the labels but also the corresponding preprocessed and geo-referenced SAR images as PNG files. The ease of access to the dataset will allow scientists from other fields, such as data science, to contribute their expertise. With this benchmark dataset, we enable comparability between different front detection algorithms and improve the reproducibility of front detection studies. Moreover, we present one baseline model for each kind of label type. Both models are based on the U-Net, one of the most popular deep learning segmentation architectures. In the following two post-processing procedures, the segmentation results are converted into 1-pixel-wide front delineations. By providing both types of labels, both approaches can be used to address the problem. To assess the performance of different models, we suggest first reviewing the segmentation results using the recall, precision, F_1 score, and the Jaccard index. Second, the front delineation can be evaluated by calculating the mean distance error to the labeled front. The presented vanilla models provide a baseline of $150\text{ m} \pm 24\text{ m}$ mean distance error for the Mapple Glacier in Antarctica and $840\text{ m} \pm 84\text{ m}$ for the Columbia Glacier in Alaska, which has a more complex

calving front, consisting of multiple sections, compared with a laterally well constrained, single calving front of Mapple Glacier.

1 Introduction

Ice mass loss from ice sheets and glaciers is one of the major contributors to current global sea level rise (Frederikse et al., 2020; Zemp et al., 2019; Sheperd et al., 2018; Khan et al., 2015). Alongside surface mass balance, ice losses at the calving fronts of marine- or lake-terminating glaciers are the major causes of ice depletion (Khan et al., 2015; Sheperd et al., 2018). Hence, frontal ablation, defined as glacier mass loss due to calving and submarine melt (Dryak and Enderlin, 2020), is an essential parameter of total glacier mass balances. This is all the more true because ice discharge at the calving fronts of marine-terminating glaciers is a self-reinforcing system: calving events lead to an expansion of the ice mélange triggered by the jamming wave, and the breakup and thinning of the mélange subsequently weakens the buttressing forces stabilizing the glacier (Robel, 2017). Some estimates of frontal ablation at glaciers outside the large ice sheets exist (Burgess et al., 2013; McNabb et al., 2015; Minowa et al., 2021). Neglecting the frontal ablation can significantly bias the numerical glacier models. For example Recinos et al. (2019) reported an underestimation of ice thickness in Alaska in the range of 19 % on regional scales and up to 30 % on glacier scales when omitting frontal ablation. A spatiotemporal quantification of frontal ablation can, therefore, provide much needed reference data for glacier model parametrization (Recinos et al., 2019, 2021). Frontal ablation has successfully been parameterized for individual glaciers (Åström et al., 2014; Ultee and Bassis, 2016; Todd and Christoffersen, 2014; Nick et al., 2010). These models, however, rely on data that are hard to obtain for entire glaciated regions (Recinos et al., 2019). An automated process to determine the frontal ablation based on easily accessible data is required. Changes in the position of the glaciers' calving fronts are crucial information for estimating frontal ablation. Satellite imagery facilitates the mapping of calving front positions in remote areas and on large spatial scales. Calving front positions can be acquired from optical as well as synthetic aperture radar (SAR) satellite imagery. Both imaging modalities have advantages and disadvantages for calving front delineation (Baumhoer et al., 2018). Optical satellite imagery has higher spatial accuracy and often higher resolution compared with SAR imagery. Moreover, different spectral bands allow the separation of ice types in optical images (Tedesco, 2014; Gao et al., 1998), whereas the backscatter in SAR images might be similar for different ice types. However, the water–ice boundary in SAR images has a high contrast (Liu and Jezek, 2004). Additionally, SAR can penetrate clouds and thin snow cover and is

independent of solar illumination, allowing recordings during the night and polar night. This leads to higher temporal availability, in particular in polar regions, for SAR images than for optical images. The majority of the calving glaciers are situated in polar regions (Hugonnet et al., 2021). Consequently, we decided to use SAR imagery for calving front delineation due to the better temporal coverage. Until recently, ice shelf fronts and glacier termini in related work were usually manually delineated (Baumhoer et al., 2018). However, this approach is no longer feasible with the rapidly growing satellite image archives, as manual delineation is a highly time-consuming, tedious, and expensive task. Recent studies (Baumhoer et al., 2019; Cheng et al., 2021; Davari et al., 2022, 2021; Hartmann et al., 2021; Heidler et al., 2021; Holzmann et al., 2021; Marochov et al., 2021; Mohajerani et al., 2019; Periyasamy et al., 2022; Zhang et al., 2019, 2021) have focused on deep learning to extract the calving front and have shown great success. However, the evaluations of these studies have generally been based on different datasets and are, therefore, not comparable. To bridge this gap, we introduce a benchmark dataset for calving front delineation of marine-terminating glaciers located in the Arctic, Greenland, and Antarctica. It is the first dataset to provide long-term calving front information from multiple missions. Time series from multiple missions introduce new challenges for calving front segmentation, such as different spatial resolutions; different penetration depths or sensitivity to surface changes; different signal-to-noise ratios; and different geometries, topographic effects, shading, and overlay effects. Deep learning models can be trained and tested on this publicly available dataset, and the baseline models also presented in this paper provide an initial reference point for the performance of future models.

To automatically delineate the calving front in SAR images, our baseline models perform a segmentation of the SAR image. In segmentation, each pixel is assigned to a specific class. In related works, two different segmentation approaches have been used for the delineation: the first approach performs a binary segmentation between land and ocean (including ice mélange) and extracts the calving front from the boundary between the two classes in a post-processing step, whereas the second approach directly assigns the pixels of the SAR image to the calving front and background classes. Both segmentation approaches are supervised learning tasks. Hence, ground truth segmentation images are needed for the algorithm to learn to distinguish the classes. As the classes in the two approaches differ (front and background versus land and ocean), different ground truth masks are needed for the two segmentation tasks. Like-

wise, our presented benchmark dataset has two ground truth annotations for each SAR image, one showing the segmentation front versus the background and the other showing a segmentation into the ocean (including ice mélange), rock, glacier, and “no information available” (hereafter referred to as “NA”) classes, the latter of which consists of SAR shadows, layover regions, and areas outside the swath. The second set does not represent a binary pixel classification task but rather a multi-class segmentation task. The calving front delineation is then carried out during post-processing using the boundary between the glacier and ocean classes.

In this study, we present the first publicly available annotated dataset for quantifying the performance of glacier calving front delineation algorithms on SAR images. Besides the calving front labels, the dataset also contains the corresponding preprocessed and geo-referenced SAR images, enabling a direct application of deep learning segmentation models. Our training and testing datasets include both ice-covered images (where the ocean in front of the glacier front is covered by ice mélange) and open water images (where mainly open water is present). This ensures the generalizability of tested algorithms to all ocean surface settings. Two vanilla baseline models are presented: one for the multi-class segmentation into glacier, ocean, rock outcrop, and NA areas, and one for the binary segmentation into front and background. Future models can use these models as a basis and compare them against their model’s performance using the presented dataset. For the evaluation of the algorithms, metrics are introduced that assess the segmentation performance of the presented convolutional neural network (CNN) models and the result of the front delineation after post-processing.

The next section of this paper (Sect. 2) gives an overview of related work, including other datasets and algorithms for front delineation. In Sect. 3, we explain the details of our own dataset, and Sect. 4 introduces the baseline methods. The evaluation of these methods on our dataset is given in Sect. 5. Finally, we draw some conclusions in Sect. 7.

2 Related work

Due to the high demand for information on the position of ice shelf fronts and glacier termini, several related datasets have been published, and studies have been carried out that have aimed to automatically extract these positions from satellite imagery.

2.1 Datasets

Existing datasets can be divided into SAR imagery, optical imagery, and datasets that are based on both imaging modalities. Most datasets have been constructed using solely optical images (Lippl, 2019; King and Howat, 2020; Fausto et al., 2019; Schild and Hamilton, 2013; Cheng et al., 2020). The dataset constructed by Lippl (2019) contains manually delineated calving front locations throughout

James Ross Island that were extracted using panchromatic Landsat 8 data. The remaining optical datasets feature delineations of Greenlandic glaciers. The calving front positions of the Helheim and Kangerdlugssuaq glaciers are given by Schild and Hamilton (2013) based on Moderate Resolution Imaging Spectroradiometer (MODIS) imagery. Fausto et al. (2019) delineate 47 of the largest outlet glaciers in Greenland annually at the end of the melt season between 1999 and 2018 based on Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER) and Landsat imagery. Front position changes of an even larger number (234) of Greenlandic glaciers are displayed over more than 3 decades by King and Howat (2020). Their delineations are based on ASTER and Landsat 4–8 imagery. Another extensive database (Cheng et al., 2020) comprises 22 678 calving front lines across Greenlandic marine-terminating glaciers. Parts of the calving fronts were manually delineated using optical Landsat near-infrared (NIR) band imagery, whereas the deep learning Calving Front Machine (CALFIN) algorithm (Cheng et al., 2021) produced the other part of the provided calving fronts. CALFIN was trained and tested on the manually mapped fronts included in the dataset as well as additional SAR images from Antarctica. Only two datasets (both for Jakobshavn Glacier, Sermeq Kujalleq, in Greenland) were constructed solely using SAR imagery. Zhang (2019a) comprises manually delineated calving fronts based on TerraSAR-X imagery that Zhang et al. (2019) used to train a neural network to automatically extract calving fronts from SAR images, and Zhang (2019b) includes the calving fronts that the trained network extracted from additional TerraSAR-X images. The ESA Greenland Ice Sheet CCI project team (2019) provides the calving front positions of 28 major outlet glaciers in Greenland. The manual delineation is carried out based on both SAR and optical imagery taken from European Remote-Sensing Satellite (ERS) 1 and 2 (ERS-1 and ERS-2, respectively); Envisat; Sentinel-1; and Landsat 5, 7, and 8. The dataset is continuously updated; we refer to version 3 here. Optical and SAR imagery from Landsat 8, Sentinel-2, Envisat, Advanced Land Observing Satellite (ALOS) 1 (ALOS-1), TerraSAR-X, Sentinel-1, and ALOS-2 was also used by Zhang et al. (2020a) to manually delineate the Jakobshavn Glacier, Kangerlussuaq, and Helheim glaciers. This dataset was again used to train and test a deep learning network that was subsequently used to produce a second dataset (Zhang et al., 2020b). Two massive databases that need to be mentioned for the sake of completeness are the Global Land Ice Measurements from Space (GLIMS) (Raup et al., 2018) and the Scientific Committee on Antarctic Research (SCAR) Antarctic Digital Database (ADD) (Gerrish et al., 2021). ADD includes Antarctica’s coastline south of 60° S, including grounding lines and ice shelf fronts. To date, GLIMS provides 546 300 glacier outlines from glaciers around the world with an ingestion rate of 7529 outlines per month. Their outlines are based on optical imagery.

An overview of all datasets can be seen in Table 1. Every described dataset only provides the glacier outlines or polygons, not the corresponding preprocessed satellite imagery, thereby hindering the direct application of deep learning algorithms.

2.2 Algorithms

Since 2019, several studies have applied deep learning techniques to delineate the calving front of marine-terminating glaciers in satellite imagery. The first studies were all based on the U-Net (Ronneberger et al., 2015), which is still the basis of many state-of-the-art networks in image segmentation. Mohajerani et al. (2019) employed this encoder–decoder network to segment multispectral Landsat images into calving front and background. Baumhoer et al. (2019) and Zhang et al. (2019) performed the first analyzes based on SAR imagery. Zhang et al. (2019) employed the dataset from Zhang (2019a) for training and testing, but the database from Baumhoer et al. (2019) is not publicly available. In a subsequent study, Baumhoer et al. (2021) modified their U-Net and used it to delineate the entire Antarctic coastline for 2018.

The U-Net architecture was replaced with DeepLabv3 (Chen et al., 2018b) by both Zhang et al. (2021) and Cheng et al. (2021). Both studies segment optical and SAR imagery into land and sea, including sea ice. Zhang et al. (2021) performs a comparison between the U-Net and DeepLabv3+ with different backbones. They published their manual delineations in Zhang et al. (2020a). In contrast to Zhang et al. (2019), Cheng et al. (2021) employs the Xception model (Chollet, 2017) as a backbone, like in the original DeepLabv3 paper (Chen et al., 2018b). The network of Cheng et al. (2021), called CALFIN, outputs two probability masks: land versus sea and coastline versus background. They evaluate CALFIN not only on their own published dataset (Cheng et al., 2020) but also on images from Mohajerani et al. (2019), Zhang et al. (2019), and Baumhoer et al. (2019). Another study that learns two tasks simultaneously is conducted by Heidler et al. (2021). Their network is based on the U-Net architecture but features two output heads: one for the segmentation into sea and land and one for delineating the coastline. They compare their model against other deep learning approaches including the U-Net model from Baumhoer et al. (2019). While their code is open source, their dataset is not publicly available. A completely different approach to front delineation is taken by Marochov et al. (2021). Instead of directly segmenting the entire images into the desired classes, Marochov et al. (2021) use classification networks to determine the class of every single pixel in each image separately. In their paper, Marochov et al. (2021) compare their model's results with the results given in Baumhoer et al. (2019), Cheng et al. (2021), Mohajerani et al. (2019), and Zhang et al. (2019). However, this is not a valid comparison because the models were trained and tested on different datasets, so the differences in performance can not solely be

attributed to the models but may also be due to the data utilized. Five more studies performed their experiments solely on SAR imagery. Holzmann et al. (2021) incorporate attention gates in the skip connections of the U-Net to improve the segmentation performance. Hartmann et al. (2021) implement a Bayesian U-Net to quantify the uncertainty of the model and use the uncertainty as an additional input to a second U-Net to improve its prediction of the calving front position. Davari et al. (2021) use a distance map-based loss to train their U-Net. Periyasamy et al. (2022) further optimize the feature extraction of the U-Net. Lastly, Davari et al. (2022) formulate the front segmentation as a regression task, letting a U-Net predict the distance of each pixel to the front. All five studies use the U-Net from Zhang et al. (2019) as a baseline, but they do not publish their dataset.

About half of the presented studies make their code publicly available (Cheng et al., 2021; Davari et al., 2021; Heidler et al., 2021; Marochov et al., 2021; Mohajerani et al., 2019; Zhang et al., 2021). Only two-thirds of the described studies make an effort to compare their method with other existing methods by either testing on the same dataset or by retraining and testing the other's model on the own dataset (Cheng et al., 2021; Davari et al., 2022, 2021; Hartmann et al., 2021; Heidler et al., 2021; Holzmann et al., 2021; Periyasamy et al., 2022; Zhang et al., 2021). Furthermore, testing on another's dataset alone without retraining one's own model on the other's training dataset beforehand is not a strong comparison, as one's own training dataset might form a better learning basis. The low rate of informative comparisons is understandable, as most datasets and codes are not easily accessible nor employable, satellite imagery is not provided, and preprocessing steps like geo-referencing need to be repeated. Therefore, a benchmark dataset and an easily reusable baseline code are highly needed. The dataset and code will benefit not only the comparability of future studies but also their reproducibility and the broad applicability of future models.

3 Dataset

3.1 Study sites

Seven tidewater glaciers situated in Greenland, Alaska, and on the Antarctic Peninsula (AP) were selected to generate labels to train and evaluate automated calving front mapping approaches for SAR imagery (see Fig. 1). The glaciers were selected considering existing datasets of calving fronts, available SAR coverage, and reports on calving front variability.

Along the AP, five former ice shelf tributary glaciers were selected. The AP is a hot spot of global warming, and a significant temperature increase was observed during the 20th century (Oliva et al., 2017; Turner et al., 2016). As a result, Cook and Vaughan (2010) reported the rapid retreat and even disintegration of various ice shelves along the AP. In 1995, the Prince Gustav and Larsen A ice shelves broke up, fol-

Table 1. Overview of publicly available front line datasets. “Type” indicates what types of glacial features are provided. The different types are abbreviated as line (full-line delineation), center (centerline delineation), poly (polygon-bounded), fjord (fjord boundaries), coast (coastline), glacier (glacial outline), and zones (landscape zones, as described in Sect. 3.2). “Annotation” refers to how the delineations were produced (i.e., manually, by a network, or by a model). “No. glaciers” gives the number of presented glaciers, and “No. mapped fronts” gives the number of glacier front delineations over all glaciers inherent in a dataset. The “Resolution” indicates the spatial resolution of images used for the mapping of the glacier fronts. The datasets are as follows: Lippl (Lippl, 2019); King (King and Howat, 2020); Fausto (Fausto et al., 2019); Schild (Schild and Hamilton, 2013); Cheng (Cheng et al., 2020); ADD (Gerrish et al., 2021); GLIMS (Raup et al., 2018); Zhang19a (Zhang, 2019a); Zhang19b (Zhang, 2019b); Zhang20a (Zhang et al., 2020a); Zhang20b (Zhang et al., 2020b); ESA (ESA Greenland Ice Sheet CCI project team, 2019); and the dataset presented in this paper (Gourmelon et al., 2022b), denoted as “Ours”.

Modality	Dataset	Type	Annotation	Area	No. glaciers	No. mapped fronts	Time span	Resolution
Optical	Lippl	Line	Manually	Antarctic	26	656	2014–2018	15 m
	King	Center	Manually	Greenland	234	128 442	1985–2018	30 m
	Fausto	Line	Manually	Greenland	47	1180	1999–2018	10–30 m
	Schild	Poly	Manually	Greenland	2	1862	2001–2010	250 m
	Cheng	Line and fjord	Manually and network	Greenland	66	22 678	1972–2019	30 m
	ADD	Coast	Manually and semiautomatic	Antarctic			Since 1843	
	GLIMS	Glacier	Manually and semiautomatic	Global	~ 200 000	546 300	Since 1750	
SAR	Zhang19a	Line	Manually	Greenland	1	159	2009–2015	3 m
	Zhang19b	Line	Network	Greenland	1	159	2009–2015	3 m
	Ours	Line and zones	Manually	Alaska, Antarctic, and Greenland	7	681	1996–2020	6–20 m
Optical and SAR	Zhang20a	Line	Manually	Greenland	3	2087	2002–2019	3–40 m
	Zhang20b	Line	Network	Greenland	3	2087	2002–2019	3–40 m
	ESA	Line	Manually	Greenland	28	1089	1990–2016	10–30 m

lowed by the disintegration of the Larsen B ice shelf in 2002 (Cooper, 1997; Scambos et al., 2004; Skvarca et al., 1998). Consequently, the former ice shelf tributary glaciers reacted with increased ice discharge and further frontal recession due to the loss of buttressing forces by the ice shelves (e.g., Rott et al., 2014; Seehaus et al., 2015, 2016). For this benchmark database, we selected the Dinsmoor–Bombardier–Edgeworth (DBE) and Sjøgren Inlet (SI) glacier systems, which were major tributaries of the Larsen A and Prince Gustav ice shelves, respectively. At the Larsen B embayment, the former Crane, Mapple, and Jorum ice shelf tributaries were chosen. Similar reaction patterns were observed at these glaciers. A significant rise in ice flux after the ice shelf breakups, followed by a long-term decrease, was measured. Concurrently, the glacier fronts retreated strongly after the disintegration of the ice shelf and partially stabilized or showed a readvance again after a few years (e.g., Rott et al., 2014, 2018; Wuite et al., 2015; Seehaus et al., 2015, 2016).

In Greenland, we incorporated Jakobshavn Glacier (JAC), Sermeq Kujalleq, in our database. It is located on the west coast and drains the Greenland ice sheet. For the last few decades, pronounced ice flow and calving front variabilities have been reported (Joughin et al., 2008, 2012). A frontal retreat of 16 km between 2002 and 2008 was revealed by Rosenau et al. (2013). Correlations between changes in the calving front positions and variations in ice discharge and the

formation of ice mélange in the glacier fjord have been observed in various analyses (Amundson et al., 2010; Joughin et al., 2008, 2012).

In Alaska, we selected Columbia Glacier. It is a large marine-terminating glacier and has strongly retreated since the early 1980s (McNabb et al., 2015; Krimmel, 2001). It split into two branches – the main and the west branch – in 2010 (Vijay and Braun, 2017). Between 1957 and 2007, an average ice thinning rate of 10 m a^{-1} was found (McNabb et al., 2012), and pronounced seasonal and interannual variability in the ice flow and calving front position was reported by Vijay and Braun (2017).

At all of the selected glaciers, a good temporal coverage exists for SAR imagery. In particular, TerraSAR-X and TanDEM-X strip map imagery has high temporal coverage because most glaciers are part of the so-called TanDEM-X “super test sites”. For the DBE and SI glacier systems, a detailed analysis of the calving front evolution, based on manual picking of the front positions on multi-mission SAR imagery, exists (Seehaus et al., 2015, 2016) and was incorporated in this benchmark database. At the other glaciers, the glacier fronts were also manually picked on SAR intensity imagery.




Figure 1. Overview of sampled glaciers in Alaska, Greenland, and on the Antarctic Peninsula. Pink polygons highlight the subset areas used for the data generation. The background maps are Esri satellite imagery © Esri.

3.2 Dataset generation

We used SAR imagery from the ERS-1/2, Envisat, RADARSAT-1, ALOS Phased Array L-band Synthetic Aperture Radar (ALOS PALSAR), TerraSAR-X (TSX), TanDEM-X (TDX), and Sentinel-1 satellite missions, covering the period from 1995 to 2020. The SAR data were provided by the German Aerospace Center (DLR), the European Space Agency (ESA), and the Alaska Satellite Facility (ASF). The SAR imagery was provided as Single Look Complex (SLC) data, except for some RADARSAT-1 imagery on the AP, which was provided in Precision Image (PRI) format (similar to a multi-looked intensity image). At first, the SAR images were calibrated and multi-looked to obtain approximately squared pixel sizes and to reduce speckle noise. The applied multi-look factors are provided in Table 2. They were selected based on a trade-off between loss of spatial resolution and speckle noise reduction. Subsequently, the SAR intensity imagery was geocoded and orthorectified.

On the AP, the enhanced ASTER digital elevation model (DEM) of the AP (Cook et al., 2012) was employed; for Columbia Glacier, the Shuttle Radar Topography Mission (SRTM) DEM was utilized; and for Jakobshavn Glacier, the global TanDEM-X DEM was used. The specifications and parameters of the SAR sensors and imagery are provided in Table 2.

The SAR data processing was done using the GAMMA Remote Sensing (RS) software. The manually picked glacier front positions employed for the studies by Seehaus et al. (2015, 2016) were used for the DBE and SI glacier systems. Additionally, glacier front positions were manually mapped at the Larsen B embayment and for Columbia Glacier. At Jakobshavn Glacier, Zhang et al. (2019) provided only spatial lines of their manually picked calving front positions. Thus, we ordered TSX/TDX SLC strip map imagery at DLR from the same orbits and dates as those used by Zhang et al. (2019) and applied the same SAR processing as mentioned above in

Table 2. Summary of the SAR satellites and specifications of the used imagery. All imagery was provided in Single Look Complex (SLC) format; however, for RADARSAT-1, some data takes were provided in Precision Image (PRI) format.

Platform	Sensor	Mode	SAR band	Repetition cycle [d]	Time interval	Multi-looking factor	Approx. slant range \times azimuth res. [m]	Ground range res. [m]
ERS-1/2	SAR	IM	C band	35/1	13 November 1995– 2 April 2010	1 \times 5	8 \times 4	20
RADARSAT 1	SAR	ST	C band	24	10 September 2000– 20 January 2008	1 \times 4	12 \times 5 (SLC)	20 (SLC) 12.5 (PRI)
Envisat	SAR	IM	C band	35	5 December 2003– 3 July 2010	1 \times 5	8 \times 4	20
ALOS	PALSAR	FBS	L band	46	18 May 2006– 3 March 2011	2 \times 5	5 \times 4	16.7
TerraSAR-X and TanDEM-X	SAR	SM	X band	11	13 October 2008– 20 May 2016	3 \times 3	1.4 \times 2	6.7
Sentinel-1A/B	SAR	IW	C band	6/12	18 December 2015– 12 June 2020	5 \times 1	4 \times 20	20

The abbreviations used in the table are as follows: IM – image mode, ST – standard beam, FBS – fine-beam single polarization, SM – stripmap, and IW – interferometric wide swath.

Table 3. Quality factors of manual calving front mapping and the related horizontal uncertainty of the mapped glacier front position.

Quality factor	1	2	3	4	5	6
[m]	70	130	200	230	450	> 450

order to have the same imagery setting for each satellite at all of our study sites. As Zhang et al. (2019) applied an additional geo-referencing step using manually defined ground control points by means of Google Earth imagery, which we could not completely replicate, there was still a spatial offset between our SAR imagery and their front positions. Thus, we manually remapped (i.e., redrew) all glacier fronts at Jakobshavn Glacier. A quality factor ranging from 1 to 6 was assigned to each calving position. The quality factors are a subjective measure of the reliability of the picked front position depending on the similarity of the ice mélange and the glacier. Table 3 shows the quality factors and the respective uncertainty values perpendicular to the glacier front.

Next, the geo-referenced SAR intensity imagery of all glaciers was cropped to the areas of interest (see Fig. 1) and converted to 16 bit single-channel images. In order to define different surface types in the SAR imagery, the manually mapped calving fronts (spatial line) were combined with a stable glacier outline dataset. At the AP, the “ice feature catchments” and “rock outcrops” polygons from the ADD were taken. At Jakobshavn Glacier, we manually generated the relevant layers. At Columbia Glacier, we relied on the Randolph Glacier Inventory (RGI 6.0), which was slightly manually refined because Columbia Glacier retreated and thinned strongly and new rock outcrops were formed, which are not included in the inventory. The 8 bit single-channel

images identifying four “zones” – “ocean”, “rock outcrops”, “glacier area”, and “no information available” (with the latter including the SAR shadow, layover regions, and areas outside the SAR swath) – in the SAR imagery were generated based on these spatial features. Additionally, two-class imagery was generated solely using the picked calving front locations. The categorized images have the same extent, dimensions, and spatial resolution as the respective SAR image subsets, allowing the definition of two sets of labels for the SAR image subsets that can be used for machine learning methods: one binary set for direct front segmentation and one set (including the four-class images) for multi-class zone segmentation. For the second set of labels, four zones are chosen instead of only two zones (ocean versus non-ocean). This is done for two reasons: first, the rock outcrop is needed to distinguish between the coastline and calving front; second, SAR shadows, overlay regions, and areas outside the swath are simply displayed as zero values in the satellite imagery. Labeling the same pattern (patches of zero values) once as ocean and once as non-ocean may hinder the learning ability of the neural networks. To support the post-processing of the calving front detection, we provide additional bounding box information on the maximum glacier front extents at each glacier for each label (see Sect. 4.3).

The resulting database, comprising the preprocessed SAR image subsets and both label sets, was split into training and test samples. The samples from the DBE, SI, Crane and Jorum glaciers as well as JAC are used to train the calving front detection algorithms. This training dataset comprises samples of different glacier geometries (see Fig. 1) and data from all incorporated SAR sensors. The samples from the Mapple and Columbia glaciers are used to evaluate the performance of the neural networks. At both glaciers, the samples contain

imagery with open ocean and ice-covered sea surface next to the calving front. Mapple Glacier has a relatively simple glacier geometry, comprising a single calving front that is well constrained by the fjord valley walls. The geometry of Columbia Glacier is much more complex. It consists of multiple branches, and the strong glacier retreat causes the split of the calving front into various sections. The number of images per glacier, sensor, and label set is summarized in Table 4.

4 Baseline methods

Alongside our benchmark dataset, we present two models, one for direct front segmentation and one for zone segmentation, which can serve as baseline methods for future studies. The complete workflow comprises preprocessing, processing with the neural network, and post-processing, including front extraction. An overview of this workflow is given in Fig. 2.

4.1 Preprocessing

As most of the standard preprocessing has already been done for the benchmark dataset (see Sect. 3), only preprocessing techniques related to the specific architecture of the neural network need to be applied. In the front segmentation model, we thicken the front labels by morphological dilation by employing a rectangular 5 pixel \times 5 pixel structuring element. Thickening the front mitigates the class imbalance inherent in the calving front dataset. A problem due to class imbalance arises when the class distributions are highly imbalanced, leading to low prediction accuracy for the rare class (Ling and Sheng, 2010). In our case, the front is only a 1-pixel-wide line and, therefore, has notably fewer pixels than the background. Hence, we use this morphological dilation and a specialized loss function for the training, which is described in Sect. 5.2. For the evaluation, the 1-pixel-wide fronts are used.

Another preprocessing step that we apply is patch extraction, where each image is divided into tiles of the same size. Neural networks usually process several images in one forward pass. This group of images is called a batch, and a batch size of more than one allows for faster training (Bengio, 2012). As the images in our dataset are large and have different sizes, there are two ways to achieve a sufficient batch size that still fits in the GPU: resizing and patch extraction. We choose patch extraction, as information is lost during resizing; however, the downside to patch extraction is the loss of the global context of the patches. Before the images can be divided into patches, each image must be padded with zeros so that no remainder is left that is smaller than the size of the patch. Next, a sliding-window approach (see Fig. 3) is used to extract 256 pixel \times 256 pixel patches. The step size for the validation and test images is set to 128 so that the patches overlap and stitching artifacts can be reduced when the images are reassembled (see Sect. 4.3). For training im-

ages, the stride is chosen such that the patches do not overlap, as this reduces the computational load considerably, and the network still sees the complete dataset once.

4.2 Deep learning models

Both our models are based on the U-Net (Ronneberger et al., 2015), an encoder–decoder CNN, which is used in most state-of-the-art networks for image segmentation. We adapt the standard U-Net architecture to our needs and refine it by inserting Atrous Spatial Pyramid Pooling (ASPP) (Chen et al., 2018a) in the bottleneck layer. ASPP applies atrous convolutions with different dilation rates in parallel, resulting in differently sized fields of view, and then fuses the obtained feature maps, allowing robust segmentation of objects at multiple scales. As calving fronts, like any other geologic structure, do not have a fixed size, this specialized layer is advantageous for our models. Our chosen dilation rates are 1, 2, 4, and 8. Other changes to the original U-Net architecture are the choice of Leaky Rectified Linear Unit (ReLU) with a negative slope of 0.1 as the activation layer, 32 as the number of start feature layers, and “same” instead of “valid” padding. Our complete architecture can be seen in Fig. 4. The output of the model depends on the type of segmentation: in direct front segmentation, we receive a probability map where each pixel value indicates the predicted probability (although not normalized to $\in [0, 1]$ – the sigmoid function still needs to be applied) that this pixel belongs to the front; in zone segmentation, we get such a probability map for each zone. For the delineation of the calving front, these probability maps are further processed during post-processing.

4.3 Post-processing for front delineation

During post-processing, the output probability patches are first merged into complete images. For merging, we do not simply use the average of the values where the patches overlap but apply Gaussian importance weighting before taking the average, which reduces stitching artifacts. Gaussian importance weighting gives higher weights to pixels close to the center of patches and lower weights to pixels near the edge of the patches (Isensee et al., 2021). This is done by element-wise multiplication of each patch prediction by a Gaussian kernel of similar size. During patch merging, the sum of the respective pixel values is normalized by the sum of the corresponding Gaussian kernel’s elements. Afterwards, the area padded with zeros during preprocessing in the SAR image is removed from our output.

4.3.1 Zone segmentation

For zone segmentation, we obtain the zone prediction for each pixel by looking at all the probability maps and selecting the zone whose probability map has the highest value at the given pixel. To fill gaps in the ocean zone, which

Table 4. Summary of labels for each glacier and sensor as well as for training and test datasets.

Platform	DBE	SI	Jorum	Crane	Mapple	Columbia	Jakobshavn	Training	Test	All
ERS-1/2	16	28	3	5	2			52	2	54
RADARSAT	27	27						54		54
Envisat	26	27	10	9	10			72	10	82
ALOS	20	7	6	7	8			40	8	48
TerraSAR-X and TanDEM-X	44	32	43	48	22	47	159	326	69	395
Sentinel-1A/B			15		15	18		15	33	48
Sum	133	121	77	69	57	65	159	559	122	681




Figure 2. Workflow for both front and zone segmentation. The enhancement of the ocean prediction includes filling gaps in the ocean zone and selecting only the largest connected ocean component as the ocean region.

is required for further steps, we do not apply morphological filtering like Baumhoer et al. (2019, 2021). Instead, we perform a connected component analysis (CCA) on the inverted predicted ocean region to receive all connected non-ocean regions and mark all but the largest connected component in the original prediction that belong to the ocean region. In this way, we can guarantee the filling of small (and larger) gaps inside the ocean zone without altering the zone’s outer boundary. This might eliminate islands (which were not present in our training data); however, this is unimportant for the front delineation task. Performing CCA on the ocean zone and omitting all but the largest connected component leaves us with one ocean area. The boundary between this ocean area and all adjacent predicted glacier zones yields the 1-pixel-wide glacier termini. We mask the glacier

termini with a bounding box specified separately for each glacier to obtain only the calving front of the observed dynamic glacier. Finally, independent front segments shorter than 750 m are removed because they likely belong to static ice–ocean boundaries that are only partially excluded by the bounding box and that we are not interested in mapping. The shortest front in our dataset is 1.5 km long. Thus, we choose the minimum reasonable length of a predicted calving front to be half of this length. However, be aware that this parameter might need to be adjusted for new glaciers.

4.3.2 Front segmentation

To receive the front prediction, we first apply the sigmoid function in a pixel-wise manner on the probability map to




Figure 3. The sliding-window approach for patch extraction. A window with the desired patch size is slid over the image with a certain stride or step size. SAR imagery is provided by DLR, ESA, and ASF.

receive predicted probability values between zero and one. Next, we use a threshold of 0.12 to binarize the prediction. If the probability value is higher than the threshold, the pixel belongs to the front; otherwise, it belongs to the background. The threshold was empirically chosen based on results using the validation set. The obtained prediction is not yet a 1-pixel-wide line. Therefore, we use skeletonization to extract a network of 1-pixel-wide lines from our prediction. As these lines still show branches, we search for the longest path in each separate skeleton and delete the remaining branches to obtain filaments. We employ the method of Koch and Rosolowsky (2015) to receive filaments from skeletons. As with zone segmentation, we use the specified bounding boxes to hide static ice–ocean interfaces and discard fronts shorter than 750 m.

5 Evaluation

5.1 Evaluation metrics

For different tasks, different evaluation metrics are needed. Therefore, this section presents appropriate evaluation metrics for zone and front segmentation as well as the main objective – front delineation. Segmentation evaluation metrics can only tell how well the segmentation itself works, not how close the resulting delineated front is to the actual front after post-processing. Hence, our segmentation metrics only assess the quality of the intermediate result (i.e., the segmentation). If the intermediate result is not sufficient, post-processing will likely not fix the problem. Therefore, an evaluation of segmentation performance is also necessary, primarily because it directly evaluates the neural network’s performance and can provide insight into whether perfor-

mance problems lie with the neural network or with the post-processing of its results.

5.1.1 Segmentation

Performance metrics for supervised classification and segmentation tasks are calculated using the confusion matrix, which gives the number of true positives, false positives, true negatives, and false negatives (Fawcett, 2006). In binary segmentation, like our front segmentation, the pixels that belong to the desired class are referred to as positive; the remaining pixels are referred to as negative. The prefixes “true” and “false” indicate whether the division into positive and negative pixels is consistent with the prediction stated in the label or contradicts the label. In multi-class segmentation, like our zone segmentation, each class is first evaluated separately by considering the other classes as background and, thus, negative. Separately evaluating each class allows multi-class problems to be split into several binary tasks, and a separate confusion matrix can be determined for each class. Thus, a value for the metric is obtained for each class. To get a collective value for the metric, one can average the class-wise metrics. We use both the class-wise and multi-class metrics to evaluate the zone model because the individual class-wise metrics provide valuable insight into how good the predictions are for each class. The averaged multi-class metric gives us an overview of how well the segmentation is working in general, but only the class-wise metrics allow us to analyze individual classes (e.g., that the rock outcrop segmentation is accurate, but the glacier segmentation is poor).

We apply four common metrics (Pedregosa et al., 2011): recall, precision, the F_1 score, and the Jaccard index (Jaccard, 1912), the latter of which is also called “Intersection over Union” (IoU). The recall is the percentage of all positive pixels predicted to be positive, while precision is the percentage of positively predicted pixels among all positive pixels (Lewis, 1990). The F_1 score is the harmonic mean of recall and precision (Fawcett, 2006). As the name infers, the IoU divides the intersection of two sets by their union, where these two sets are the pixels of the predicted class and the pixels of the actual target for binary segmentation. A visual interpretation of the IoU is shown in Fig. 5. The Eqs. (1)–(4) provide the calculations of the four metrics from true positives (TP), false positives (FP), true negatives (TN), and false negatives (FN). We calculate all metrics on the binarized predictions.

$$\text{Recall} = \frac{\text{TP}}{\text{TP} + \text{FN}} \quad (1)$$

$$\text{Precision} = \frac{\text{TP}}{\text{TP} + \text{FP}} \quad (2)$$

$$F_1 \text{ score} = 2 \cdot \frac{\text{Precision} \cdot \text{Recall}}{\text{Precision} + \text{Recall}} \quad (3)$$

$$\text{IoU} = \frac{\text{TP}}{\text{TP} + \text{FP} + \text{FN}} \quad (4)$$




Figure 4. The architecture of both presented segmentation models. Different blocks denote the output of the transformations indicated by the color of each block. Note that the sizes of the blocks are not true to scale. The numbers beneath the blocks indicate the number of feature maps. The size of the features maps is indicated inside or next to the blocks. The number of classes, given by “# Classes” in the figure, is one for front segmentation and four for zone segmentation. SAR imagery is provided by DLR, ESA, and ASF.

$$\text{IoU} (A, B) = \frac{|A \cap B|}{|A \cup B|}$$

Figure 5. A visualization of the Jaccard index or Intersection over Union (IoU). The two ellipses, *A* and *B*, signify two sets.

where \mathcal{I} is the set of all evaluated images, \mathcal{P} is the set of ground truth front pixels of one specific image, \mathcal{Q} is the set of corresponding predicted front pixels in that image, and $|\cdot|$ denotes the cardinality of a set.

5.1.2 Front delineation

As an evaluation metric for the front delineation, we employ the mean distance error (MDE), which is adapted from Baumhoer et al. (2019), Cheng et al. (2021), Heidler et al. (2021), Mohajerani et al. (2019, 2021), and Zhang et al. (2019, 2021). Each predicted front is compared to its ground truth. The Euclidean distance to the closest pixel in the ground truth front is calculated for each pixel in the predicted front. To make the metric symmetric, the distance to the closest pixel in the predicted front is also calculated for each pixel of the ground truth front. These distances are stored for all images, and the average is taken over all images, forming the mean distance error. The analytical formula of the MDE is as follows:

$$\text{MDE}(\mathcal{I}) = \frac{1}{\sum_{(\mathcal{P}, \mathcal{Q}) \in \mathcal{I}} (|\mathcal{P}| + |\mathcal{Q}|)} \cdot \sum_{(\mathcal{P}, \mathcal{Q}) \in \mathcal{I}} \left(\sum_{p \in \mathcal{P}} \min_{q \in \mathcal{Q}} \|p - q\|_2 + \sum_{q \in \mathcal{Q}} \min_{p \in \mathcal{P}} \|p - q\|_2 \right), \quad (5)$$

5.2 Experimental protocol

For our experiments, we further split the training dataset presented in Sect. 3 into two parts: a part used for training, and a part used for validation. A 9 : 1 ratio is applied for training versus validation. All hyperparameter optimizations were done, comparing the results on the validation set. Using the validation set allows us to frequently check the performance of our model during implementation while keeping the final evaluation on the test set unbiased. Like the test set, the validation set is preprocessed with overlapping patch extraction, which increases the amount of validation data relative to the training data. On-the-fly augmentations are applied to the training set only. These random augmentations include rotations, horizontal flips, brightness adjustments, Gaussian noise, and a transform called “wrap” that acts like many simultaneous elastic transforms with Gaussian sigmas set at various harmonics (Nicolaou et al., 2022). A visualization of these augmentations is presented in Fig. 6. The probabilities with which these augmentations are applied to the input were determined empirically and depend on the segmentation task. For front segmentation, all augmentations are applied with a probability of 0.65; for zone segmentation, brightness adjustment and wrap are applied with a probability of 0.1, Gaussian noise and rotation are applied with a probability of 0.5, and flipping is applied with a probability of 0.3.




Figure 6. Applied augmentations. From left to right, the panels show the original image/label, horizontal flip, rotation (here 180°), brightness adjustment, Gaussian noise, and wrap. The first row shows the SAR images, the second row gives the label for the front segmentation, and the third row shows the label for the zone segmentation. For brightness adjustment, Gaussian noise, and wrap augmentations, the intensities of the transforms have been enhanced such that they are more visible. SAR imagery is provided by DLR, ESA, and ASF.

Next, all patches (including the validation and test sets) are z score normalized using the mean and standard deviation of the training dataset. The patches are fed into the neural network with a batch size of 16, as this was the maximum amount that could fit into the GPU using a 256×256 patch size. With a larger batch size, the patches would need to be smaller. However, we chose a 256×256 patch size to ensure a sufficiently large global context and adjust the batch size accordingly. To train the models, we employ a cyclic learning rate scheduler (Smith, 2017) in combination with the Adam optimizer (Kingma and Ba, 2014) and apply different loss functions for each segmentation task. For the zone segmentation model, the base learning rate for the scheduler is chosen to be 4×10^{-5} , and the maximum learning rate is 2×10^{-4} . For the front segmentation model, the base learning rate is 1×10^{-4} , and the maximum learning rate is 5×10^{-4} . We use the rule of thumb given in Smith (2017) to select a proper step size for both models. Hence, the step size is set to 30 000, which roughly equals the number of training patches divided by the batch size times 10. Moreover, we perform gradient clipping to avoid exploding gradients. The global norm of the gradients (i.e., the norm calculated over all model parameters) is truncated to 1.0 for both models. The values for gradient clipping, base, and maximum learning rates were determined empirically using the Optuna hyperparameter optimization framework (Akiba et al., 2019). The loss function

for zone segmentation is a weighted combination of Dice (Sudre et al., 2017) and cross-entropy loss (Bishop, 1995). With Optuna, the weighting was determined to be optimal if both parts of the combined loss function were weighted equally. For front segmentation, an improved distance map loss (Davari et al., 2021) is employed. The improved distance map loss from Davari et al. (2021) multiplies the network's prediction by weights calculated based on the front label before computing the cross-entropy of the prediction and the front label. We replace the cross-entropy with the Dice loss, as the Dice loss is better suited for class-imbalanced problems; otherwise, we keep the loss from Davari et al. (2021) as is. In order to get the weights for the prediction, the front label is first dilated (see Eq. 6, where δ_w indicates the morphological dilation with a $w \times w$ rectangular structuring element). Second, the Euclidean distance transform (EDT) is applied on the dilated label, which is then divided by a relaxation factor R and fed into a sigmoid function, which is denoted by σ (see Eq. 7). The final weight map is the sum of the output of the sigmoid function and the dilated front label, which is first inverted and then scaled by a factor k (see Eq. 8). The improved distance map loss is visualized in Fig. 7. The hyperparameters w , R , and k were set to 5, 1, and 0.1, respectively, after empirical evaluations.

$$\mathbf{x}_{\text{dil}} = \delta_w(\mathbf{x}) \quad (6)$$

$$\mathbf{x}_{\text{edt}} = \sigma\left(\frac{\text{EDT}(\mathbf{x}_{\text{dil}})}{R}\right) \quad (7)$$

$$\mathbf{weights} = \mathbf{x}_{\text{edt}} + k \cdot (1 - \mathbf{x}_{\text{dil}}) \quad (8)$$

Both models were trained five times for 150 epochs using early stopping with a patience of 30 epochs. For the zones model, the early-stopping criterion is the mean validation multi-class IoU, while the criterion is the mean validation loss for the front model. The weights of the best epoch of each training round are used to evaluate the models' performance on the test set. The best epoch is determined by the highest mean IoU on the validation set for zone segmentation and by the lowest validation loss for front segmentation. Finally, the mean and standard deviation of the two models' performances over the five training rounds are calculated.

5.3 Results

In this section, we evaluate the performance of our vanilla models, present the results quantitatively and qualitatively, and discuss these results objectively. First and foremost, we present our results on the test set. As our test set is out of sample (i.e., the study sites in the test set were not covered in the training set), the test set is entirely independent of the training set and, hence, qualifies for estimating the generalizability of our trained models to new unseen glaciers. Some previous calving front delineation studies (Cheng et al., 2021; Davari et al., 2022, 2021; Hartmann et al., 2021; Holzmann et al., 2021; Marochov et al., 2021; Periyasamy et al., 2022; Zhang et al., 2019) used in-sample test sets (i.e., the test set includes images from the same glaciers as covered in the training set but from different time points). Deep learning models will produce more precise front delineations on in-sample test sets compared with out-of-sample test sets (Marochov et al., 2021), as the generalization gap between the training and test set is smaller for in-sample test sets (Quinero-Candela et al., 2008). If a model is meant only to delineate fronts from glaciers covered in the training set, it is legitimate to evaluate the model on an in-sample test set. However, if the model shall also be applied to new glaciers, it is crucial to evaluate it on an out-of-sample test set. To approximate an evaluation on an in-sample test set, we also present our results on the validation set (see Appendix A). Bear in mind that we used this validation set to optimize our hyperparameters; hence, the results on the validation set do not accurately reflect the performance of our model on in-sample test data and likely lead to somewhat biased results.

5.3.1 Zone segmentation

We begin our evaluation with the segmentation results of the zones model. Therefore, we calculate the segmentation metrics presented in Sect. 5.1.1. The resulting values for the test set are given in Table 5. All metrics show the highest values

for the NA area; this was to be expected, as areas outside the swath, lay-over regions, and SAR shadows are relatively easy to distinguish in the images compared with the other classes. Overall, the precision is higher than the recall except for the glacier class. The rock outcrop class has the highest drop between precision and recall, implying that a considerable fraction of the predicted rock pixels is actually rock, but only a smaller fraction of the real rock outcrop was predicted to be rock. The rock outcrop class also has the lowest F_1 score and lowest IoU. The evaluation on the validation set results in a precision of 94.5 ± 0.2 , a recall of 91.8 ± 0.5 , an F_1 score of 92.9 ± 0.3 , and an IoU of 88.0 ± 0.3 . The complete evaluation on the validation set is given in Table A1. All metrics are considerably higher for the validation set than for the out-of-sample test set.

5.3.2 Front segmentation

For the evaluation of the front segmentation model, we do not distinguish between multi-class and class-wise metrics, as the front segmentation is a binary task. The precision, recall, F_1 score, and IoU for the test set are given in Table 6. The recall is the highest metric (57.3 ± 5.4), signifying that more than half of the ground truth front pixels are covered by the predicted front. However, all values are considerably lower than for the zone segmentation. The low values are explained by the composition and structure of the label, as the front label shows a high class imbalance (i.e., the background occupies significantly more pixels than the front, which is just a 1-pixel-wide line). The 1-pixel-wide structure of the front also leads to problems with the segmentation metrics. If, for example, the predicted front lies right next to the ground truth front separated by only 1 pixel, the IoU will be zero, as it measures the fraction of overlap between the prediction and ground truth. Therefore, even though the prediction would be very close to the ground truth, the IoU would not reflect the quality of this prediction. Hence, these results should not be compared directly to the zone segmentation metrics but only to other front segmentation results. Instead, the proposed MDE given in Sect. 5.1.2 should be preferred for comparisons between the different segmentation models. The front segmentation metrics on the validation set are shown in Table A2.

5.3.3 Front delineation

After post-processing, the MDE can be calculated. Table 7 gives the MDEs for the zone and front models and additionally breaks down the metric by glacier and season. Overall, the zone segmentation model leads to front predictions that are, on average, 134 m closer to the front than the front segmentation model. Interestingly, however, the latter yields more accurate predictions for the Mapple Glacier than the zone segmentation model. Qualitative results of the front models are shown in Fig. 8. Figure 8a and b are examples




Figure 7. Visualization of the improved distance map loss, showing (a) the weights, (b) the network's prediction, and (c) the front label. \circ denotes the Hadamard product.

Table 5. Segmentation results on the test set for the zones model (in percent). The scope implicates the multi-class or respective class-wise metric.

Scope	Precision \uparrow	Recall \uparrow	F_1 score \uparrow	IoU \uparrow
NA area	99.5 \pm 0.1	91.2 \pm 1.3	94.8 \pm 0.8	90.9 \pm 1.3
Rock outcrop	82.0 \pm 0.5	59.6 \pm 1.3	67.9 \pm 0.8	53.5 \pm 0.7
Glacier	74.5 \pm 0.7	89.5 \pm 1.1	80.9 \pm 0.3	68.5 \pm 0.5
Ocean and ice mélange	80.9 \pm 2.2	78.3 \pm 3.1	76.8 \pm 1.6	66.0 \pm 1.5
All zones	84.2 \pm 0.5	79.6 \pm 0.9	80.1 \pm 0.5	69.7 \pm 0.6

An upward arrow next to the metric's name indicates that the model's performance improves as the metric's value increases, whereas a downward arrow indicates that the model's performance improves as the metric's value decreases.

Table 6. Segmentation results on the test set for the front model (in percent).

Precision \uparrow	Recall \uparrow	F_1 score \uparrow	IoU \uparrow
2.4 \pm 0.3	57.3 \pm 5.4	4.3 \pm 0.5	2.2 \pm 0.3

An upward arrow next to the metric's name indicates that the model's performance improves as the metric's value increases, whereas a downward arrow indicates that the model's performance improves as the metric's value decreases.

of accurate front delineations. Figure 8c shows an inaccurate front delineation: firstly, part of the coastline is confused as calving front; secondly, only one of the three calving fronts of the Columbia Glacier is recognized. The two unidentified calving fronts contribute significantly to the MDE. As the MDE is symmetric, for each pixel in the two unidentified ground truth fronts, the Euclidean distance to the closest pixel in the predicted front contributes to the MDE, and, obviously, the closest pixels are far away from the closest identified calving front pixels. Figure 8d gives an example where sea ice was confused with a calving front. In general, predictions for images from summer are more accurate than images from winter, as can be seen in Table 7. This seasonal gap can be explained by sea ice forming in front of the calving front during winter, with similar back-scattering properties as the glacier. Therefore, sea ice makes an accurate calving front delineation more difficult. Qualitative results of the zone segmentation model are given in Fig. 9. Figure 9a and b are examples of accurate front delineations, whereas Fig. 9c and d show confusions with sea ice again.

Table 8 breaks the MDEs down by glacier and satellite, showing that predictions for Sentinel-1 images are by far the least accurate over the complete test set. The reason for the high MDEs for Sentinel-1 is that only one of the three calving fronts is identified in most images at Columbia Glacier, and, as explained earlier, the other two fronts negatively impact the MDE. Front model predictions for Sentinel-1 images of Mapple Glacier only have a marginally higher MDE than images from other satellites, while zone model predictions for Sentinel-1 images of Mapple Glacier show the lowest MDE across all satellites. In summary, this shows that the sensor influences the quality of the front delineation, but the glacier geometry has a greater impact.

A breakdown of the MDEs by glaciers and the sensor spatial resolution is shown in Table 9. A spatial resolution of 20 m² is associated with the highest MDEs. When the images with this resolution are split up into glaciers, it becomes apparent that the high MDEs result from images of the Columbia Glacier. The lowest average MDEs are observed for a spatial resolution of 17 m². This can be explained by the fact that the test set only includes images of the Mapple Glacier at a resolution of 17 m². Hence, we can conclude that the glacier geometry has a higher impact on the front delineation performance than the spatial resolution of the image.

The predictions on the validation set receive an average MDE of 391 m \pm 94 m for the front model and 449 m \pm 31 m for the zone model, which is considerably lower than on the test set. Figures A1 and A2 show that the front delineation is very accurate, but part of the coastlines is predicted as calving front. Hence, with a post-processing scheme that can eliminate calving front predictions on the coastline, the MDE

Table 7. Mean distance errors (MDEs) on the test set (in meters), also differentiated by glacier and season. \emptyset stands for the number of images for which no front was predicted. The number after \in denotes how many images of the category (given glacier and season) were present in the test set.

Network		MDE↓	\emptyset ↓	Summer		Winter	
				MDE↓	\emptyset ↓	MDE↓	\emptyset ↓
Front	All	887 ± 189	7 ± 3 ∈ 122	738 ± 111	4 ± 1 ∈ 68	1054 ± 308	4 ± 2 ∈ 54
	Columbia	1032 ± 227	2 ± 1 ∈ 65	907 ± 131	0 ± 0 ∈ 28	1157 ± 350	2 ± 1 ∈ 37
	Mapple	150 ± 24	6 ± 2 ∈ 57	140 ± 26	2 ± 1 ∈ 40	173 ± 33	2 ± 1 ∈ 17
Zones	All	753 ± 76	1 ± 1 ∈ 122	732 ± 93	1 ± 1 ∈ 68	776 ± 65	0 ± 0 ∈ 54
	Columbia	840 ± 84	0 ± 0 ∈ 65	854 ± 111	0 ± 0 ∈ 28	826 ± 66	0 ± 0 ∈ 37
	Mapple	287 ± 48	0 ± 1 ∈ 57	262 ± 29	0 ± 1 ∈ 40	340 ± 93	0 ± 0 ∈ 17

An upward arrow next to the metric’s name indicates that the model’s performance improves as the metric’s value increases, whereas a downward arrow indicates that the model’s performance improves as the metric’s value decreases.




Figure 8. Visualization of the front segmentation models’ performance on the test set. Blue represents the ground truth, yellow represents the prediction, and pink represents the overlap of the ground truth and prediction. The turquoise rectangle is the bounding box explained in Sect. 4.3. Panel (a) is an image of the Columbia Glacier acquired on the 19 March 2012 by the TDX satellite, panel (b) is an image of the Mapple Glacier acquired on the 2 November 2009 by the TSX satellite, panel (c) is an image of the Columbia Glacier acquired on the 6 January 2018 by the Sentinel-1 satellite, and panel (d) is an image of the Mapple Glacier acquired on the 30 June 2013 by the TSX satellite. To enhance the visibility of the region of interest, the front prediction and ground truth in panels (a), (b), and (d) are dilated with a 9×9 kernel, and the front prediction and ground truth in panel (c) are dilated with a 3×3 kernel. The images of the Columbia Glacier are cropped to the region of interest for visualization purposes. SAR imagery is provided by DLR, ESA, and ASF.

would even be smaller. The complete evaluations on the validation set are given in Tables A3, A4, and A5.

5.4 Discussion and outlook

In Sect. 5.3, major performance differences among results using different satellites, resolutions, and glaciers became apparent, which can mostly be attributed to the complexity gap between the Mapple and Columbia glaciers in the test set. The Columbia Glacier consists of three separate calv-




Figure 9. Visualization of the zone segmentation models' performance on the test set. Blue represents the ground truth, yellow represents the prediction, and pink represents the overlap of the ground truth and prediction. The turquoise rectangle is the bounding box explained in Sect. 4.3. Panel (a) is an image of the Columbia Glacier acquired on the 19 March 2012 by the TDX satellite, panel (b) is an image of the Mapple Glacier acquired on the 5 November 2010 by the TSX satellite, panel (c) is an image of the Columbia Glacier acquired on the 17 April 2016 by the TDX satellite, and panel (d) is an image of the Mapple Glacier acquired on the 21 November 2007 by the PALSAR satellite. To enhance the visibility of the region of interest, the front prediction and ground truth in panels (a), (b), and (c) are dilated with a 9×9 kernel, and the front prediction and ground truth in panel (d) are dilated with a 3×3 kernel. The images of the Columbia Glacier are cropped to the region of interest for visualization purposes. SAR imagery is provided by DLR, ESA, and ASF.

Table 8. Mean distance errors (MDEs) on the test set (in meters), differentiated by glacier and satellite. \emptyset stands for the number of images for which no front was predicted. The number after \in denotes how many images of the category (given glacier and satellite) were present in the test set.

Network	Glacier		Sentinel-1	Envisat	ERS-1/2	PALSAR	TSX/TDX
Front	All	MDE↓	2806 ± 300	191 ± 32	127 ± 38	197 ± 41	663 ± 188
		\emptyset ↓	$2 \pm 1 \in 33$	$2 \pm 2 \in 10$	$0 \pm 0 \in 2$	$3 \pm 2 \in 8$	$0 \pm 0 \in 69$
	Columbia	MDE↓	3537 ± 422	–	–	–	744 ± 218
		\emptyset ↓	$2 \pm 1 \in 18$	–	–	–	$0 \pm 0 \in 47$
	Mapple	MDE↓	206 ± 33	191 ± 32	127 ± 38	197 ± 41	129 ± 33
		\emptyset ↓	$0 \pm 0 \in 15$	$2 \pm 2 \in 10$	$0 \pm 0 \in 2$	$3 \pm 2 \in 8$	$0 \pm 0 \in 22$
Zones	All	MDE↓	2201 ± 246	493 ± 119	404 ± 172	437 ± 42	547 ± 61
		\emptyset ↓	$0 \pm 0 \in 33$	$0 \pm 0 \in 10$	$0 \pm 0 \in 2$	$0 \pm 0 \in 8$	$0 \pm 0 \in 69$
	Columbia	MDE↓	2587 ± 299	–	–	–	587 ± 67
		\emptyset ↓	$0 \pm 0 \in 18$	–	–	–	$0 \pm 0 \in 47$
	Mapple	MDE↓	141 ± 29	493 ± 119	404 ± 172	437 ± 42	246 ± 57
		\emptyset ↓	$0 \pm 0 \in 15$	$0 \pm 0 \in 10$	$0 \pm 0 \in 2$	$0 \pm 0 \in 8$	$0 \pm 0 \in 22$

An upward arrow next to the metric's name indicates that the model's performance improves as the metric's value increases, whereas a downward arrow indicates that the model's performance improves as the metric's value decreases.

Table 9. Mean distance errors (MDEs) on the test set (in meters), differentiated by glacier and resolution. \emptyset stands for the number of images for which no front was predicted. The number after \in denotes how many images of the category (given glacier and resolution) were present in the test set.

Network		20 m ²		17 m ²		7 m ²	
		MDE↓	\emptyset ↓	MDE↓	\emptyset ↓	MDE↓	\emptyset ↓
Front	All	2436 ± 289	4 ± 2 ∈ 45	197 ± 41	3 ± 2 ∈ 8	663 ± 188	0 ± 0 ∈ 69
	Columbia	3537 ± 422	2 ± 1 ∈ 18	–	–	744 ± 218	0 ± 0 ∈ 47
	Mapple	192 ± 26	2 ± 2 ∈ 27	197 ± 41	3 ± 2 ∈ 8	129 ± 33	0 ± 0 ∈ 22
Zones	All	1939 ± 220	0 ± 0 ∈ 45	437 ± 42	0 ± 0 ∈ 8	547 ± 61	0 ± 0 ∈ 69
	Columbia	2587 ± 299	0 ± 0 ∈ 18	–	–	587 ± 67	0 ± 0 ∈ 47
	Mapple	323 ± 69	0 ± 0 ∈ 27	437 ± 42	0 ± 0 ∈ 8	246 ± 57	0 ± 0 ∈ 22

An upward arrow next to the metric's name indicates that the model's performance improves as the metric's value increases, whereas a downward arrow indicates that the model's performance improves as the metric's value decreases.

ing fronts in contrast to Mapple, which only features one calving front. In addition, the Columbia Glacier's calving fronts exhibit greater variability in their structure and curvature than Mapple's calving front. We, therefore, conclude that the complexity of the glacier, including the number and shape of calving fronts, has a significant impact on the delineation performance of a deep learning model.

Performance differences between zone and front models are also remarkable. The zone models' predictions lie mainly close to the calving front or in the ocean and only rarely far away on the other side of the image between, e.g., rock outcrop and NA area or rock outcrop and glacier. On the other hand, the front models' predictions are not that constrained in the region, but if they hit the correct calving front region, the prediction lies more accurately on the true calving front than the zone models' prediction. Therefore, a multi-task learning approach could be beneficial for future work, as the two tasks (zone segmentation and front segmentation) excel in different aspects and can learn from each other here.

From the class-wise segmentation metrics presented in Sect. 5.3.1, it is notable that the rock outcrop class was the least accurately predicted. One reason for this is the fact that the rock outcrop is static for each glacier (i.e., no adjustments were made for the individual image labels even if part or the whole rock outcrop was covered by snow or new rocks became visible as the ice receded throughout time). Hence, this drop in performance comes from the labels themselves. Nevertheless, we decided to use static rock outcrops, as an accurate rock outcrop prediction is not the aim of our work and only contributes to an accurate front delineation; moreover, manually labeling the rock as done for the calving front would have been beyond the scope of this paper.

Two weaknesses that can be addressed in the future include low performance on images with sea ice and confusion of the coastline with the calving front. Weak performance on images with sea ice might be compensated for by augmenting winter images more frequently than summer images or giving higher weight to the loss of winter images. The confusion

of the coastline with the calving front could be remedied by improved post-processing, such as masking with a rock outcrop, which, of course, would have to be available for each new glacier.

All in all, the vanilla models presented in this paper provide a starting point for accurate segmentation of calving fronts, and the introduced dataset enables training, testing, and comparison of deep learning calving front delineation models.

6 Code and data availability

The code is publicly available from GitHub at https://github.com/Nora-Go/Calving_Fronts_and_Where_to_Find_Them (Gourmelon et al., 2022c). The exact version of the model used to produce the results presented in this paper is archived on Zenodo: <https://doi.org/10.5281/zenodo.6469519> (Gourmelon et al., 2022a). The dataset is publicly available from PANGAEA at <https://doi.org/10.1594/PANGAEA.940950> (Gourmelon et al., 2022b).

7 Conclusions

In this paper, we introduce a benchmark dataset for calving front delineation of marine-terminating glaciers using SAR imagery. The dataset is intended to be used for the training, testing, and comparison of deep learning calving front extraction models. It comprises 681 samples and is the first to provide long-term calving front information from multi-mission data. Furthermore, besides the provided segmentation labels, the dataset contains the corresponding preprocessed and geo-referenced SAR images as PNG files. The dataset's ease of access allows scientists to focus on the deep learning model and invites researchers from related scientific fields, such as data science, to contribute to this urgent task. The calving front position is an indicator of frontal ablation, which is a key parameter of the total glacier mass balance.

With this dataset, we aim to enable the training of deep-learning models capable of accurately delineating calving fronts of glaciers around the world at any time, permitting a spatiotemporal quantification of calving rates. The dataset includes SAR imagery of glaciers from Antarctica, Greenland, and Alaska. With its light independency and cloud penetration ability, SAR imagery permits continuous front tracking across all seasons. The test set includes the Mapple Glacier located in Antarctica and the Columbia Glacier in Alaska. Glaciers from different regions in the test set guarantee that, during testing, the wide applicability of the model is verified. Moreover, no images from these two glaciers are included in the training set, making it an out-of-sample test set. Hence, the models' generalizability to unseen glaciers and regions also outside Antarctica and Greenland can be ensured during testing. Additionally, with the Columbia Glacier, the test set includes challenging samples, as Columbia Glacier has three calving fronts with highly variable geometry. Therefore, a low MDE on these samples indicates the robustness of the trained model to variations in the shape and number of calving fronts in the given images. The two labels in the dataset, zone and front segmentation, supply comprehensive information on the calving front delineation task and allow for the task to be approached in different ways. The well-documented and easy-to-use code of the two vanilla baseline models introduced in this paper ensures the reproducibility of all experiments and provides a starting point for an accurate front delineation method that can easily be extended. The performance of different models can be quantified and compared to other models consistently with the presented evaluation metrics. The resulting MDE of $150\text{ m} \pm 24\text{ m}$ for the Mapple Glacier and $840\text{ m} \pm 84\text{ m}$ for the Columbia Glacier build a sound baseline for future calving front delineation models.

Appendix A: Validation results

This section depicts the vanilla models' results on the in-sample validation set. The validation set was used to optimize the models' hyperparameters; hence, the results are biased.

A1 Zone segmentation

Table A1 presents the evaluation of the zone models' segmentation results on the validation set. The segmentation metrics introduced in Sect. 5.1.1 are employed for the evaluation.

Table A1. Segmentation results on the validation set for the zones model (in percent). The scope implicates the multi-class or respective class-wise metric.

Scope	Precision↑	Recall↑	F_1 score↑	IoU↑
All zones	94.5 ± 0.2	91.8 ± 0.5	92.9 ± 0.3	88.0 ± 0.3
NA area	96.0 ± 1.0	91.2 ± 2.3	92.9 ± 1.3	89.3 ± 1.4
Rock outcrop	89.4 ± 0.8	82.6 ± 1.1	85.7 ± 0.3	75.8 ± 0.4
Glacier	95.6 ± 0.1	97.5 ± 0.2	96.5 ± 0.1	93.3 ± 0.1
Ocean and ice mélange	97.3 ± 0.3	96.0 ± 0.4	96.5 ± 0.2	93.5 ± 0.3

An upward arrow next to the metric's name indicates that the model's performance improves as the metric's value increases, whereas a downward arrow indicates that the model's performance improves as the metric's value decreases.

A2 Front segmentation

Table A2 gives the precision, recall, F_1 score, and IoU of the front models' segmentation results on the validation set.

Table A2. Segmentation results on the validation set for the front model (in percent).

Precision↑	Recall↑	F_1 score↑	IoU↑
2.3 ± 0.2	67.5 ± 1.4	4.5 ± 0.4	2.3 ± 0.2

An upward arrow next to the metric's name indicates that the model's performance improves as the metric's value increases, whereas a downward arrow indicates that the model's performance improves as the metric's value decreases.

A3 Front delineation

The MDEs on the post-processed validation set results are given in Table A3, which also differentiates between glaciers and seasons. Table A4 breaks these results down by glacier and satellite, and Table A5 breaks them down by glacier and resolution. Visualizations of the prediction results on the validation set can be seen in Figs. A1 and A2.

Table A3. Mean distance errors (MDEs) on the validation set (in meters), also differentiated by glacier and season. \emptyset stands for the number of images for which no front was predicted. The number after \in denotes how many images of the category (given glacier and season) were present in the validation set.

Network		MDE↓	\emptyset ↓	Summer		Winter	
				MDE↓	\emptyset ↓	MDE↓	\emptyset ↓
Front	All	391 ± 94	0 ± 0 ∈ 56	466 ± 89	0 ± 0 ∈ 36	315 ± 106	0 ± 0 ∈ 20
	Crane	593 ± 123	0 ± 0 ∈ 8	695 ± 128	0 ± 0 ∈ 6	368 ± 135	0 ± 0 ∈ 2
	DBE	561 ± 144	0 ± 0 ∈ 10	717 ± 185	0 ± 0 ∈ 6	225 ± 66	0 ± 0 ∈ 4
	JAC	222 ± 21	0 ± 0 ∈ 16	221 ± 16	0 ± 0 ∈ 6	222 ± 25	0 ± 0 ∈ 10
	Jorum	276 ± 99	0 ± 0 ∈ 8	276 ± 99	0 ± 0 ∈ 8	–	–
	SI	741 ± 383	0 ± 0 ∈ 14	730 ± 255	0 ± 0 ∈ 10	774 ± 652	0 ± 0 ∈ 4
Zones	All	449 ± 31	0 ± 0 ∈ 56	563 ± 43	0 ± 0 ∈ 36	318 ± 26	0 ± 0 ∈ 20
	Crane	784 ± 79	0 ± 0 ∈ 8	852 ± 106	0 ± 0 ∈ 6	580 ± 170	0 ± 0 ∈ 2
	DBE	1043 ± 85	0 ± 0 ∈ 10	1095 ± 121	0 ± 0 ∈ 6	907 ± 30	0 ± 0 ∈ 4
	JAC	150 ± 16	0 ± 0 ∈ 16	169 ± 25	0 ± 0 ∈ 6	138 ± 12	0 ± 0 ∈ 10
	Jorum	402 ± 73	0 ± 0 ∈ 8	402 ± 73	0 ± 0 ∈ 8	–	–
	SI	737 ± 73	0 ± 0 ∈ 14	777 ± 48	0 ± 0 ∈ 10	669 ± 124	0 ± 0 ∈ 4

An upward arrow next to the metric's name indicates that the model's performance improves as the metric's value increases, whereas a downward arrow indicates that the model's performance improves as the metric's value decreases.

Table A4. Mean distance errors (MDEs) on the validation set (in meters), differentiated by glacier and satellite. \emptyset stands for the number of images for which no front was predicted. The number after \in denotes how many images of the category (given glacier and satellite) were present in the validation set.

Network			RADARSAT-1	Sentinel-1	Envisat	ERS-1/2	PALSAR	TSX/TDX
Front	All	MDE↓	545 ± 142	205 ± 25	539 ± 158	840 ± 134	1427 ± 239	341 ± 89
		\emptyset ↓	0 ± 0 ∈ 4	0 ± 0 ∈ 1	0 ± 0 ∈ 9	0 ± 0 ∈ 5	0 ± 0 ∈ 2	0 ± 0 ∈ 35
	Crane	MDE↓	–	–	1147 ± 625	303 ± 81	–	569 ± 109
		\emptyset ↓	–	–	0 ± 0 ∈ 1	0 ± 0 ∈ 1	–	0 ± 0 ∈ 6
	DBE	MDE↓	358 ± 145	–	366 ± 137	–	614 ± 168	663 ± 211
		\emptyset ↓	0 ± 0 ∈ 1	–	0 ± 0 ∈ 4	–	0 ± 0 ∈ 1	0 ± 0 ∈ 4
	JAC	MDE↓	–	–	–	–	–	222 ± 21
		\emptyset ↓	–	–	–	–	–	0 ± 0 ∈ 16
	Jorum	MDE↓	–	205 ± 25	442 ± 236	–	–	252 ± 91
		\emptyset ↓	–	0 ± 0 ∈ 1	0 ± 0 ∈ 2	–	–	0 ± 0 ∈ 5
SI	MDE↓	575 ± 153	–	600 ± 261	916 ± 143	1780 ± 291	725 ± 590	
	\emptyset ↓	0 ± 0 ∈ 3	–	0 ± 0 ∈ 2	0 ± 0 ∈ 4	0 ± 0 ∈ 1	0 ± 0 ∈ 4	
Zones	All	MDE↓	900 ± 118	164 ± 45	776 ± 136	846 ± 157	1229 ± 215	365 ± 26
		\emptyset ↓	0 ± 0 ∈ 4	0 ± 0 ∈ 1	0 ± 0 ∈ 9	0 ± 0 ∈ 5	0 ± 0 ∈ 2	0 ± 0 ∈ 35
	Crane	MDE↓	–	–	393 ± 356	154 ± 27	–	837 ± 73
		\emptyset ↓	–	–	0 ± 0 ∈ 1	0 ± 0 ∈ 1	–	0 ± 0 ∈ 6
	DBE	MDE↓	812 ± 78	–	938 ± 140	–	1434 ± 164	1065 ± 126
		\emptyset ↓	0 ± 0 ∈ 1	–	0 ± 0 ∈ 4	–	0 ± 0 ∈ 1	0 ± 0 ∈ 4
	JAC	MDE↓	–	–	–	–	–	150 ± 16
		\emptyset ↓	–	–	–	–	–	0 ± 0 ∈ 16
	Jorum	MDE↓	–	164 ± 45	155 ± 55	–	–	441 ± 80
		\emptyset ↓	–	0 ± 0 ∈ 1	0 ± 0 ∈ 2	–	–	0 ± 0 ∈ 5
SI	MDE↓	912 ± 135	–	801 ± 209	939 ± 186	1118 ± 331	605 ± 82	
	\emptyset ↓	0 ± 0 ∈ 3	–	0 ± 0 ∈ 2	0 ± 0 ∈ 4	0 ± 0 ∈ 1	0 ± 0 ∈ 4	

An upward arrow next to the metric's name indicates that the model's performance improves as the metric's value increases, whereas a downward arrow indicates that the model's performance improves as the metric's value decreases.

Table A5. Mean distance errors (MDEs) on the validation set (in meters), differentiated by glacier and resolution. \emptyset stands for the number of images for which no front was predicted. The number after \in denotes how many images of the category (given glacier and resolution) were present in the validation set.

Network	Glacier		20	17	12	7	6
Front	All	MDE↓	636 ± 130	1427 ± 239	545 ± 142	605 ± 264	222 ± 21
		\emptyset ↓	0 ± 0 ∈ 15	0 ± 0 ∈ 2	0 ± 0 ∈ 4	0 ± 0 ∈ 19	0 ± 0 ∈ 16
	Crane	MDE↓	754 ± 337	–	–	569 ± 109	–
		\emptyset ↓	0 ± 0 ∈ 2	–	–	0 ± 0 ∈ 6	–
	DBE	MDE↓	366 ± 137	614 ± 168	358 ± 145	663 ± 211	–
		\emptyset ↓	0 ± 0 ∈ 4	0 ± 0 ∈ 1	0 ± 0 ∈ 1	0 ± 0 ∈ 4	–
	JAC	MDE↓	–	–	–	–	222 ± 21
		\emptyset ↓	–	–	–	–	0 ± 0 ∈ 16
	Jorum	MDE↓	369 ± 165	–	–	252 ± 91	–
		\emptyset ↓	0 ± 0 ∈ 3	–	–	0 ± 0 ∈ 5	–
	SI	MDE↓	774 ± 185	1780 ± 291	575 ± 153	725 ± 590	–
		\emptyset ↓	0 ± 0 ∈ 6	0 ± 0 ∈ 1	0 ± 0 ∈ 3	0 ± 0 ∈ 4	–
Zones	All	MDE↓	782 ± 134	1229 ± 215	900 ± 118	718 ± 62	718 ± 62
		\emptyset ↓	0 ± 0 ∈ 15	0 ± 0 ∈ 2	0 ± 0 ∈ 4	0 ± 0 ∈ 19	0 ± 0 ∈ 16
	Crane	MDE↓	270 ± 179	–	–	837 ± 73	–
		\emptyset ↓	0 ± 0 ∈ 2	–	–	0 ± 0 ∈ 6	–
	DBE	MDE↓	938 ± 140	1434 ± 164	812 ± 78	1065 ± 126	–
		\emptyset ↓	0 ± 0 ∈ 4	0 ± 0 ∈ 1	0 ± 0 ∈ 1	0 ± 0 ∈ 4	–
	JAC	MDE↓	–	–	–	–	150 ± 16
		\emptyset ↓	–	–	–	–	0 ± 0 ∈ 16
	Jorum	MDE↓	158 ± 46	–	–	441 ± 80	–
		\emptyset ↓	0 ± 0 ∈ 3	–	–	0 ± 0 ∈ 5	–
	SI	MDE↓	874 ± 170	1118 ± 331	912 ± 135	605 ± 82	–
		\emptyset ↓	0 ± 0 ∈ 6	0 ± 0 ∈ 1	0 ± 0 ∈ 3	0 ± 0 ∈ 4	–

An upward arrow next to the metric's name indicates that the model's performance improves as the metric's value increases, whereas a downward arrow indicates that the model's performance improves as the metric's value decreases.




Figure A1. Visualization of the front segmentation models' performance on the validation set. Blue represents the ground truth, yellow represents the prediction, and pink represents the overlap of the ground truth and prediction. The turquoise rectangle is the bounding box explained in Sect. 4.3. Panel (a) is an image of the Crane Glacier acquired on the 8 December 2010 by the TSX satellite, panel (b) is an image of the DBE Glacier acquired on the 21 July 2004 by the Envisat satellite, panel (c) is an image of the Jakobshavn Glacier acquired on the 3 October 2012 by the TSX satellite, panel (d) is an image of the Jorum Glacier acquired on the 19 December 2010 by the TSX satellite, and panel (e) is an image of the SI Glacier acquired on the 19 October 2013 by the TSX satellite. The images are cropped to the region of interest for visualization purposes. To enhance the visibility, the front prediction and ground truth in panels (a), (c), (d), and (e) are dilated with a 9×9 kernel, and the front prediction and ground truth in panel (b) are dilated with a 3×3 kernel. SAR imagery is provided by DLR, ESA, and ASF.




Figure A2. Visualization of the zone segmentation models' performance on the validation set. Blue represents the ground truth, yellow represents the prediction, and pink represents the overlap of the ground truth and prediction. The turquoise rectangle is the bounding box explained in Sect. 4.3. Panel (a) is an image of the Crane Glacier acquired on the 8 December 2010 by the TSX satellite, panel (b) is an image of the DBE Glacier acquired on the 11 July 2013 by the TSX satellite, panel (c) is an image of the Jakobshavn Glacier acquired on the 15 August 2009 by the TSX satellite, panel (d) is an image of the Jorum Glacier acquired on the 1 September 2012 by the TSX satellite, and panel (e) is an image of the SI Glacier acquired on the 19 August 2011 by the TSX satellite. The images are cropped to the region of interest for visualization purposes. To enhance the visibility, the front predictions and ground truths are dilated with a 9×9 kernel. SAR imagery is provided by DLR, ESA, and ASF.

Author contributions. NG was responsible for conceptualizing the study, developing the methodology and software, project administration, and writing and preparing the manuscript. TS curated the data and prepared and wrote the manuscript. MB and AM supervised the study and reviewed and edited the manuscript. VC supervised the study, validated data, and prepared and wrote the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Special issue statement. This article is part of the special issue "Benchmark datasets and machine learning algorithms for Earth system science data (ESSD/GMD inter-journal SI)". It is not associated with a conference.

Acknowledgements. We acknowledge the free provision of the SAR data via various proposals from ESA, ASF, and DLR.

Financial support. This research has been supported by the Friedrich-Alexander-Universität Erlangen-Nürnberg (Emerging Fields Initiative "Tapping the Potential of Earth Observation") and the STAEDTLER Foundation.

Review statement. This paper was edited by David Carlson and reviewed by two anonymous referees.

References

- Akiba, T., Sano, S., Yanase, T., Ohta, T., and Koyama, M.: Op-tuna: A Next-generation Hyperparameter Optimization Framework, in: Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, Anchorage, AS, USA, 4–8 August 2019, 2623–2631, <https://doi.org/10.1145/3292500.3330701>, 2019.
- Amundson, J. M., Fahnestock, M., Truffer, M., Brown, J., Lüthi, M. P., and Motyka, R. J.: Ice mélange dynamics and implications for terminus stability, Jakobshavn Isbræ, Greenland, *J. Geophys. Res.*, 115, F01005, <https://doi.org/10.1029/2009JF001405>, 2010.
- Åström, J. A., Vallot, D., Schäfer, M., Welty, E. Z., O'Neel, S., Bartholomäus, T. C., Liu, Y., Riikilä, T. I., Zwinger, T., Timonen, J., and Moore, J. C.: Termini of calving glaciers as self-organized critical systems, *Nat. Geosci.*, 7, 874–878, <https://doi.org/10.1038/ngeo2290>, 2014.
- Baumhoer, C., Dietz, A., Dech, S., and Kuenzer, C.: Remote Sensing of Antarctic Glacier and Ice-Shelf Front Dynamics – A Review, *Remote Sens.*, 10, 1445, <https://doi.org/10.3390/rs10091445>, 2018.
- Baumhoer, C. A., Dietz, A. J., Kneisel, C., and Kuenzer, C.: Automated Extraction of Antarctic Glacier and Ice Shelf Fronts from Sentinel-1 Imagery Using Deep Learning, *Remote Sens.*, 11, 2529, <https://doi.org/10.3390/rs11212529>, 2019.
- Baumhoer, C. A., Dietz, A. J., Kneisel, C., Paeth, H., and Kuenzer, C.: Environmental drivers of circum-Antarctic glacier and ice shelf front retreat over the last two decades, *The Cryosphere*, 15, 2357–2381, <https://doi.org/10.5194/tc-15-2357-2021>, 2021.
- Bengio, Y.: Practical Recommendations for Gradient-Based Training of Deep Architectures, in: *Neural Networks: Tricks of the Trade*, 2 edn., Springer, Berlin, Heidelberg, 437–478, https://doi.org/10.1007/978-3-642-35289-8_26, 2012.
- Bishop, C. M.: *Neural networks for pattern recognition*, 1 edn., edited by: Jordan, M., Kleinberg, J., and Schölkopf, B., Clarendon Press, ISBN-10 0-387-31073-8, ISBN-13 978-0387-31073-2, 1995.
- Burgess, E. W., Forster, R. R., and Larsen, C. F.: Flow velocities of Alaskan glaciers, *Nat. Commun.*, 4, 2146, <https://doi.org/10.1038/ncomms3146>, 2013.
- Chen, L.-C., Papandreou, G., Kokkinos, I., Murphy, K., and Yuille, A. L.: DeepLab: Semantic Image Segmentation with Deep Convolutional Nets, Atrous Convolution, and Fully Connected CRFs, *IEEE T. Pattern. Anal.*, 40, 834–848, <https://doi.org/10.1109/TPAMI.2017.2699184>, 2018a.
- Chen, L.-C., Zhu, Y., Papandreou, G., Schroff, F., and Adam, H.: Encoder-Decoder with Atrous Separable Convolution for Semantic Image Segmentation, in: Proceedings of the European Conference on Computer Vision (ECCV), Munich, Germany, 8–14 September 2018, 801–818, 2018b.
- Cheng, D., Hayes, W., and Larour, E.: CALFIN: Calving front dataset for East/West Greenland, 1972–2019, Dryad [data set], <https://doi.org/10.7280/D1FH5D>, 2020.
- Cheng, D., Hayes, W., Larour, E., Mohajerani, Y., Wood, M., Velicogna, I., and Rignot, E.: Calving Front Machine (CALFIN): glacial termini dataset and automated deep learning extraction method for Greenland, 1972–2019, *The Cryosphere*, 15, 1663–1675, <https://doi.org/10.5194/tc-15-1663-2021>, 2021.
- Chollet, F.: Xception: Deep Learning With Depthwise Separable Convolutions, in: Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition, Honolulu, HI, USA, 21–26 July 2017, 1251–1258, <https://doi.org/10.1109/CVPR.2017.195>, 2017.
- Cook, A. J. and Vaughan, D. G.: Overview of areal changes of the ice shelves on the Antarctic Peninsula over the past 50 years, *The Cryosphere*, 4, 77–98, <https://doi.org/10.5194/tc-4-77-2010>, 2010.
- Cook, A. J., Murray, T., Luckman, A., Vaughan, D. G., and Barand, N. E.: A new 100-m Digital Elevation Model of the Antarctic Peninsula derived from ASTER Global DEM: methods and accuracy assessment, *Earth Syst. Sci. Data*, 4, 129–142, <https://doi.org/10.5194/essd-4-129-2012>, 2012.
- Cooper, A.: Historical observations of Prince Gustav Ice Shelf, *Polar Rec.*, 33, 285–294, <https://doi.org/10.1017/S0032247400025389>, 1997.
- Davari, A., Islam, S., Seehaus, T., Hartmann, A., Braun, M., Maier, A., and Christlein, V.: On Mathews Correlation Coefficient and Improved Distance Map Loss for Automatic Glacier Calving

- Front Segmentation in SAR Imagery, *IEEE T. Geosci. Remote.*, 60, 5213212, <https://doi.org/10.1109/TGRS.2021.3115883>, 2021.
- Davari, A., Baller, C., Seehaus, T., Braun, M., Maier, A., and Christlein, V.: Pixel-wise Distance Regression for Glacier Calving Front Detection and Segmentation, *IEEE T. Geosci. Remote.*, 60, 5224610, <https://doi.org/10.1109/TGRS.2022.3158591>, 2022.
- Dryak, M. C. and Enderlin, E. M.: Analysis of Antarctic Peninsula glacier frontal ablation rates with respect to iceberg melt-inferred variability in ocean conditions, *J. Glaciol.*, 66, 457–470, <https://doi.org/10.1017/jog.2020.21>, 2020.
- ESA Greenland Ice Sheet CCI project team: ESA Greenland Ice Sheet Climate Change Initiative (Greenland_Ice_Sheet_cci): Greenland Calving Front Locations, v3.0, Centre for Environmental Data Analysis [data set], <https://catalogue.ceda.ac.uk/uuid/8889dfe3de45406e815bce13ae8a0c92> (last access: 20 September 2022), 2019.
- Fausto, R. S., Andersen, J., Hansen, K., Box, J. E., Andersen, S. B., Ahlstrøm, A. P., van As, D., Citterio, M., Colgan, W., Karlsson, N. B., Kjeldsen, K. K., Korsgaard, N. J., Larsen, S. H., Mankoff, K. D., Pedersen, A. Ø., Shields, C. L., Solgaard, A., and Vandecrux, B.: Programme for monitoring of the Greenland ice sheet (PROMICE): Calving front line, 1999–2018, Arctic Data Center [data set], https://doi.org/10.22008/promice/data/calving_front_lines, 2019.
- Fawcett, T.: An introduction to ROC analysis, *Pattern Recogn. Lett.*, 27, 861–874, <https://doi.org/10.1016/j.patrec.2005.10.010>, 2006.
- Frederikse, T., Landerer, F., Caron, L., Adhikari, S., Parkes, D., Humphrey, V. W., Dangendorf, S., Hogarth, P., Zanna, L., Cheng, L., and Wu, Y.-H.: The causes of sea-level rise since 1900, *Nature*, 584, 393–397, <https://doi.org/10.1038/s41586-020-2591-3>, 2020.
- Gao, B.-C., Han, W., Tsay, S. C., and Larsen, N. F.: Cloud Detection over the Arctic Region Using Airborne Imaging Spectrometer Data during the Daytime, *J. App. Meteorol. Clim.*, 37, 1421–1429, [https://doi.org/10.1175/1520-0450\(1998\)037<1421:CDOTAR>2.0.CO;2](https://doi.org/10.1175/1520-0450(1998)037<1421:CDOTAR>2.0.CO;2), 1998.
- Gerrish, L., Fretwell, P., and Cooper, P.: High resolution vector polylines of the Antarctic coastline (7.4), UK Polar Data Centre, Natural Environment Research Council, UK Research & Innovation [data set], <https://doi.org/10.5285/e46be5bc-ef8e-4fd5-967b-92863fbe2835>, 2021.
- Gourmelon, N., Seehaus, T., Braun, M. H., Maier, A., and Christlein, V.: Calving Fronts and Where to Find Them, Zenodo [code], <https://doi.org/10.5281/zenodo.6469519>, 2022a.
- Gourmelon, N., Seehaus, T., Braun, M. H., Maier, A., and Christlein, V.: CaFFe (CALving Fronts and where to Find thEm: a benchmark dataset and methodology for automatic glacier calving front extraction from sar imagery), PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.940950>, 2022b.
- Gourmelon, N., Seehaus, T., Braun, M. H., Maier, A., and Christlein, V.: Vincent: Calving Fronts and Where to Find Them, GitHub [code], https://github.com/Nora-Go/Calving_Fronts_and_Where_to_Find_Them, last access: 20 September 2022c.
- Hartmann, A., Davari, A., Seehaus, T., Braun, M., Maier, A., and Christlein, V.: Bayesian U-Net for Segmenting Glaciers in SAR Imagery, in: 2021 IEEE International Geoscience and Remote Sensing Symposium IGARSS, Brussels, Belgium, 11–16 July 2021, 3479–3482, <https://doi.org/10.1109/IGARSS47720.2021.9554292>, 2021.
- Heidler, K., Mou, L., Baumhoer, C., Dietz, A., and Zhu, X. X.: HED-UNet: Combined Segmentation and Edge Detection for Monitoring the Antarctic Coastline, *IEEE T. Geosci. Remote.*, 2021, 4300514, <https://doi.org/10.1109/TGRS.2021.3064606>, 2021.
- Holzmann, M., Davari, A., Seehaus, T., Braun, M., Maier, A., and Christlein, V.: Glacier Calving Front Segmentation Using Attention U-Net, in: 2021 IEEE International Symposium on Geoscience and Remote Sensing (IGARSS), Brussels, Belgium, 11–16 July 2021, 3483–3486, <https://doi.org/10.1109/IGARSS47720.2021.9555067>, 2021.
- Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussaillant, I., Brun, F., and Käab, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- Isensee, F., Jaeger, P. F., Kohl, S. A. A., Petersen, J., and Maier-Hein, K. H.: nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation, *Nat. Methods*, 18, 203–211, <https://doi.org/10.1038/s41592-020-01008-z>, 2021.
- Jaccard, P.: The Distribution of the Flora in the Alpine Zone, *New Phytol.*, 11, 37–50, <https://doi.org/10.1111/j.1469-8137.1912.tb05611.x>, 1912.
- Joughin, I., Das, S. B., King, M. A., Smith, B. E., Howat, I. M., and Moon, T.: Seasonal Speedup Along the Western Flank of the Greenland Ice Sheet, *Science*, 320, 781–783, <https://doi.org/10.1126/science.1153288>, 2008.
- Joughin, I., Smith, B. E., Howat, I. M., Floricioiu, D., Alley, R. B., Truffer, M., and Fahnestock, M.: Seasonal to decadal scale variations in the surface velocity of Jakobshavn Isbrae, Greenland: Observation and model-based analysis, *J Geophys. Res.-Earth.*, 117, F02030, <https://doi.org/10.1029/2011JF002110>, 2012.
- Khan, S. A., Aschwanden, A., Bjørk, A. A., Wahr, J., Kjeldsen, K. K., and Kjær, K. H.: Greenland ice sheet mass balance: a review, *Rep. Prog. Phys.*, 78, 046801, <https://doi.org/10.1088/0034-4885/78/4/046801>, 2015.
- King, M. and Howat, I.: Data from: Dynamic ice loss from the Greenland Ice Sheet driven by sustained glacier retreat, Dryad [data set], <https://doi.org/10.5061/dryad.qrfj6q5cb>, 2020.
- Kingma, D. P. and Ba, J.: Adam: A Method for Stochastic Optimization, arXiv [preprint], <https://doi.org/10.48550/arXiv.1412.6980>, 22 December 2014.
- Koch, E. W. and Rosolowsky, E. W.: Filament identification through mathematical morphology, *Mon. Not. R. Astron. Soc.*, 452, 3435–3450, <https://doi.org/10.1093/mnras/stv1521>, 2015.
- Krimmel, R. M.: Photogrammetric Data Set, 1957–2000, and Bathymetric Measurements for Columbia Glacier, Alaska, Tech. Rep., U.S. Geological Survey, <https://doi.org/10.3133/wri20014089>, 2001.
- Lewis, D. D.: Representation Quality in Text Classification: An Introduction and Experiment, in: Proceedings of the Workshop on Speech and Natural Language, Hidden Valley, Pennsylvania, USA, 24–27 June 1990, 288–295, <https://doi.org/10.3115/116580.116681>, 1990.

- Ling, C. X. and Sheng, V. S.: Class Imbalance Problem, in: *Encyclopedia of Machine Learning*, Springer US, Boston, MA, USA, 171–171, https://doi.org/10.1007/978-0-387-30164-8_110, 2010.
- Lipl, S.: Glacier Surface Velocities and Outlet Areas from 2014–2018 on James Ross Island, Northern Antarctic Peninsula, PAN-GAEA [data set], <https://doi.org/10.1594/PANGAEA.907062>, 2019.
- Liu, H. and Jezek, K. C.: Automated extraction of coastline from satellite imagery by integrating Canny edge detection and locally adaptive thresholding methods, *Int. J. Remote Sens.*, 25, 937–958, <https://doi.org/10.1080/0143116031000139890>, 2004.
- Marochov, M., Stokes, C. R., and Carbonneau, P. E.: Image classification of marine-terminating outlet glaciers in Greenland using deep learning methods, *The Cryosphere*, 15, 5041–5059, <https://doi.org/10.5194/tc-15-5041-2021>, 2021.
- McNabb, R., Hock, R., O’Neel, S., Rasmussen, L., Ahn, Y., Braun, M., Conway, H., Herreid, S., Joughin, I., Pfeffer, W., Smith, B., and Truffer, M.: Using surface velocities to calculate ice thickness and bed topography: a case study at Columbia Glacier, Alaska, USA, *J. Glaciol.*, 58, 1151–1164, <https://doi.org/10.3189/2012JoG11J249>, 2012.
- McNabb, R. W., Hock, R., and Huss, M.: Variations in Alaska tidewater glacier frontal ablation, 1985–2013, *J. Geophys. Res.-Earth*, 120, 120–136, <https://doi.org/10.1002/2014JF003276>, 2015.
- Minowa, M., Schaefer, M., Sugiyama, S., Sakakibara, D., and Skvarca, P.: Frontal ablation and mass loss of the Patagonian icefields, *Earth Planet. Sc. Lett.*, 561, 116811, <https://doi.org/10.1016/j.epsl.2021.116811>, 2021.
- Mohajerani, Y., Wood, M., Velicogna, I., and Rignot, E.: Detection of Glacier Calving Margins with Convolutional Neural Networks: A Case Study, *Remote Sens.*, 11, 74, <https://doi.org/10.3390/rs11010074>, 2019.
- Mohajerani, Y., Jeong, S., Scheuchl, B., Velicogna, I., Rignot, E., and Milillo, P.: Automatic delineation of glacier grounding lines in differential interferometric synthetic-aperture radar data using deep learning, *Sci. Rep.*, 11, 4992, <https://doi.org/10.1038/s41598-021-84309-3>, 2021.
- Nick, F. M., van der Veen, C. J., Vieli, A., and Benn, D. I.: A physically based calving model applied to marine outlet glaciers and implications for the glacier dynamics, *J. Glaciol.*, 56, 781–794, <https://doi.org/10.3189/002214310794457344>, 2010.
- Nicolaou, A., Christlein, V., Riba, E., Shi, J., Vogeler, G., and Seuret, M.: TorMentor: Deterministic dynamic-path, data augmentations with fractals, *arXiv [preprint]*, <https://doi.org/10.48550/arxiv.2204.03776>, 7 April 2022.
- Oliva, M., Navarro, F., Hrbáček, F., Hernández, A., Nývlt, D., Pereira, P., Ruiz-Fernández, J., and Trigo, R.: Recent regional climate cooling on the Antarctic Peninsula and associated impacts on the cryosphere, *Sci. Total Environ.*, 580, 210–223, <https://doi.org/10.1016/j.scitotenv.2016.12.030>, 2017.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E.: Scikit-learn: Machine Learning in Python, *J. Mach. Learn. Res.*, 12, 2825–2830, <http://jmlr.org/papers/v12/pedregosa1a.html> (last access: 20 September 2022), 2011.
- Periyasamy, M., Davari, A., Seehaus, T., Braun, M., Maier, A., and Christlein, V.: How to Get the Most Out of U-Net for Glacier Calving Front Segmentation, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 15, 1712–1723, <https://doi.org/10.1109/JSTARS.2022.3148033>, 2022.
- Quinonero-Candela, J., Sugiyama, M., Schwaighofer, A., and Lawrence, N.: *Dataset Shift in Machine Learning*, Neural Information Processing series, MIT Press, ISBN 978-0-262-17005-5, 2008.
- Raup, B., Racoviteanu, A., Khalsa, S., Helm, C., Armstrong, R., and Arnaud, Y.: GLIMS and NSIDC (2005, updated 2018): Global Land Ice Measurements from Space glacier database. Compiled and made available by the international GLIMS community and the National Snow and Ice Data Center, Boulder CO, USA, GLIMS [data set], <https://doi.org/10.7265/N5V98602>, 2018.
- Recinos, B., Maussion, F., Rothenpieler, T., and Marzeion, B.: Impact of frontal ablation on the ice thickness estimation of marine-terminating glaciers in Alaska, *The Cryosphere*, 13, 2657–2672, <https://doi.org/10.5194/tc-13-2657-2019>, 2019.
- Recinos, B., Maussion, F., Noël, B., Möller, M., and Marzeion, B.: Calibration of a frontal ablation parameterisation applied to Greenland’s peripheral calving glaciers, *J. Glaciol.*, 67, 1177–1189, <https://doi.org/10.1017/jog.2021.63>, 2021.
- Robel, A. A.: Thinning sea ice weakens buttressing force of ice-berg mélange and promotes calving, *Nat. Commun.*, 8, 14596, <https://doi.org/10.1038/ncomms14596>, 2017.
- Ronneberger, O., Fischer, P., and Brox, T.: U-Net: Convolutional Networks for Biomedical Image Segmentation, in: *Proceedings of the 18th International Conference on Medical Image Computing and Computer Assisted Interventions*, Munich, Germany, 5–9 October 2015, 234–241, https://doi.org/10.1007/978-3-319-24574-4_28, 2015.
- Rosenau, R., Schwalbe, E., Maas, H.-G., Baessler, M., and Dietrich, R.: Grounding line migration and high-resolution calving dynamics of Jakobshavn Isbræ, West Greenland, *J. Geophys. Res.-Earth*, 118, 382–395, <https://doi.org/10.1029/2012JF002515>, 2013.
- Rott, H., Floricioiu, D., Wuite, J., Scheiblauer, S., Nagler, T., and Kern, M.: Mass changes of outlet glaciers along the Nordsjököld Coast, northern Antarctic Peninsula, based on TanDEM-X satellite measurements, *Geophys. Res. Lett.*, 41, 8123–8129, <https://doi.org/10.1002/2014GL061613>, 2014.
- Rott, H., Abdel Jaber, W., Wuite, J., Scheiblauer, S., Floricioiu, D., van Wessel, J. M., Nagler, T., Miranda, N., and van den Broeke, M. R.: Changing pattern of ice flow and mass balance for glaciers discharging into the Larsen A and B embayments, Antarctic Peninsula, 2011 to 2016, *The Cryosphere*, 12, 1273–1291, <https://doi.org/10.5194/tc-12-1273-2018>, 2018.
- Scambos, T. A., Bohlander, J., Shuman, C. A., and Skvarca, P.: Glacier acceleration and thinning after ice shelf collapse in the Larsen B embayment, Antarctica, *Geophys. Res. Lett.*, 31, L18402, <https://doi.org/10.1029/2004GL020670>, 2004.
- Schild, K. and Hamilton, G.: Terminus position time series: Helheim and Kangerdlugssuaq glaciers, Greenland, Arctic Data Center [data set], <https://doi.org/10.18739/A2W93G>, 2013.
- Seehaus, T., Marinsek, S., Helm, V., Skvarca, P., and Braun, M.: Changes in ice dynamics, elevation and mass discharge of Dinsmoor–Bombardier–Edgeworth glacier system,

- Antarctic Peninsula, *Earth Planet. Sc. Lett.*, 427, 125–135, <https://doi.org/10.1016/j.epsl.2015.06.047>, 2015.
- Seehaus, T. C., Marinsek, S., Skvarca, P., van Wessem, J. M., Reijmer, C. H., Seco, J. L., and Braun, M. H.: Dynamic Response of Sjøgren Inlet Glaciers, Antarctic Peninsula, to Ice Shelf Breakup Derived from Multi-Mission Remote Sensing Time Series, *Front. Earth Sci.*, 4, 66, <https://doi.org/10.3389/feart.2016.00066>, 2016.
- Sheperd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., and Joughin, I.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- Skvarca, P., Rack, W., Rott, H., and Y Donángelo, T. I.: Evidence of recent climatic warming on the eastern Antarctic Peninsula, *Ann. Glaciol.*, 27, 628–632, <https://doi.org/10.3189/S0260305500018164>, 1998.
- Smith, L. N.: Cyclical learning rates for training neural networks, in: 2017 IEEE winter conference on applications of computer vision (WACV), Santa Rosa, CA, USA, 24–31 March 2017, 464–472, <https://doi.org/10.1109/WACV.2017.58>, 2017.
- Sudre, C. H., Li, W., Vercauteren, T., Ourselin, S., and Jorge Cardoso, M.: Generalised Dice Overlap as a Deep Learning Loss Function for Highly Unbalanced Segmentations, in: *Deep Learning in Medical Image Analysis and Multimodal Learning for Clinical Decision Support*, Québec City, QC, Canada, 14 September 2017, 240–248, https://doi.org/10.1007/978-3-319-67558-9_28, 2017.
- Tedesco, M.: Remote sensing of the cryosphere, in: *The Cryosphere Science Series*, 1st edn., edited by: Knight, P. G., Wiley Blackwell, Hoboken, NJ, USA, <https://doi.org/10.1002/9781118368909>, 2014.
- Todd, J. and Christoffersen, P.: Are seasonal calving dynamics forced by buttressing from ice mélange or undercutting by melting? Outcomes from full-Stokes simulations of Store Glacier, West Greenland, *The Cryosphere*, 8, 2353–2365, <https://doi.org/10.5194/tc-8-2353-2014>, 2014.
- Turner, J., Lu, H., White, I., King, J. C., Phillips, T., Hosking, J. S., Bracegirdle, T. J., Marshall, G. J., Mulvaney, R., and Deb, P.: Absence of 21st century warming on Antarctic Peninsula consistent with natural variability, *Nature*, 535, 411–415, <https://doi.org/10.1038/nature18645>, 2016.
- Ultee, L. and Bassis, J.: The future is Nye: an extension of the perfect plastic approximation to tidewater glaciers, *J. Glaciol.*, 62, 1143–1152, <https://doi.org/10.1017/jog.2016.108>, 2016.
- Vijay, S. and Braun, M.: Seasonal and Interannual Variability of Columbia Glacier, Alaska (2011–2016): Ice Velocity, Mass Flux, Surface Elevation and Front Position, *Remote Sens.*, 9, 635, <https://doi.org/10.3390/rs9060635>, 2017.
- Wuite, J., Rott, H., Hetzenecker, M., Floricioiu, D., De Rydt, J., Gudmundsson, G. H., Nagler, T., and Kern, M.: Evolution of surface velocities and ice discharge of Larsen B outlet glaciers from 1995 to 2013, *The Cryosphere*, 9, 957–969, <https://doi.org/10.5194/tc-9-957-2015>, 2015.
- Zemp, M., Huss, M., Thibert, E., Eckert, N., McNabb, R., Huber, J., Barandun, M., Machguth, H., Nussbaumer, S. U., Gärtner-Roer, I., Thomson, L., Paul, F., Maussion, F., Kutuzov, S., and Cogley, J. G.: Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016, *Nature*, 568, 382–386, <https://doi.org/10.1038/s41586-019-1071-0>, 2019.
- Zhang, E.: The ground truth of the calving fronts in Jakobshavn Isbræ, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.897065>, 2019a.
- Zhang, E.: The calving fronts delineated by the network in Jakobshavn Isbræ, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.897064>, 2019b.
- Zhang, E., Liu, L., and Huang, L.: Automatically delineating the calving front of Jakobshavn Isbræ from multitemporal TerraSAR-X images: a deep learning approach, *The Cryosphere*, 13, 1729–1741, <https://doi.org/10.5194/tc-13-1729-2019>, 2019.
- Zhang, E., Liu, L., Huang, L., and Ng, K. S.: Manually delineated calving fronts at Jakobshavn Isbræ, Kangerlussuaq, and Helheim, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.923270>, 2020a.
- Zhang, E., Liu, L., Huang, L., and Ng, K. S.: Network delineated calving fronts at Jakobshavn Isbræ, Kangerlussuaq, and Helheim, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.923272>, 2020b.
- Zhang, E., Liu, L., Huang, L., and Ng, K. S.: An automated, generalized, deep-learning-based method for delineating the calving fronts of Greenland glaciers from multi-sensor remote sensing imagery, *Remote Sens. Environ.*, 254, 112265, <https://doi.org/10.1016/j.rse.2020.112265>, 2021.