



GWL_FCS30: a global 30 m wetland map with a fine classification system using multi-sourced and time-series remote sensing imagery in 2020

Xiao Zhang^{1,2}, Liangyun Liu^{1,2,3}, Tingting Zhao^{1,4}, Xidong Chen⁵, Shangrong Lin⁶, Jinqing Wang^{1,2,3}, Jun Mi^{1,2,3}, and Wendi Liu^{1,2,3}

¹International Research Center of Big Data for Sustainable Development Goals, Beijing 100094, China

²Key Laboratory of Digital Earth Science, Aerospace Information Research Institute, Chinese Academy of Sciences, Beijing 100094, China

³School of Electronic, Electrical and Communication Engineering, University of Chinese Academy of Sciences, Beijing 100049, China

⁴College of Geomatics, Xi'an University of Science and Technology, Xi'an 710054, China

⁵North China University of Water Resources and Electric Power, Zhengzhou 450046, China

⁶School of Atmospheric Sciences, Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Sun Yat-sen University, Zhuhai 519082, Guangdong, China

Correspondence: Liangyun Liu (liuly@radi.ac.cn)

Received: 24 May 2022 – Discussion started: 19 August 2022

Revised: 21 December 2022 – Accepted: 27 December 2022 – Published: 17 January 2023

Abstract. Wetlands, often called the “kidneys of the earth”, play an important role in maintaining ecological balance, conserving water resources, replenishing groundwater and controlling soil erosion. Wetland mapping is very challenging because of its complicated temporal dynamics and large spatial and spectral heterogeneity. An accurate global 30 m wetland dataset that can simultaneously cover inland and coastal zones is lacking. This study proposes a novel method for wetland mapping by combining an automatic sample extraction method, existing multi-sourced products, satellite time-series images and a stratified classification strategy. This approach allowed for the generation of the first global 30 m wetland map with a fine classification system (GWL_FCS30), including five inland wetland sub-categories (permanent water, swamp, marsh, flooded flat and saline) and three coastal tidal wetland sub-categories (mangrove, salt marsh and tidal flats), which was developed using Google Earth Engine platform. We first combined existing multi-sourced global wetland products, expert knowledge, training sample refinement rules and visual interpretation to generate large and geographically distributed wetland training samples. Second, we integrated the Landsat reflectance time-series products and Sentinel-1 synthetic aperture radar (SAR) imagery to generate various water-level and phenological information to capture the complicated temporal dynamics and spectral heterogeneity of wetlands. Third, we applied a stratified classification strategy and the local adaptive random forest classification models to produce the wetland dataset with a fine classification system at each $5^\circ \times 5^\circ$ geographical tile in 2020. Lastly, GWL_FCS30, mosaicked by 961 $5^\circ \times 5^\circ$ regional wetland maps, was validated using 25 708 validation samples, which achieved an overall accuracy of 86.44 % and a kappa coefficient of 0.822. The cross-comparisons with other global wetland products demonstrated that the GWL_FCS30 dataset performed better in capturing the spatial patterns of wetlands and had significant advantages over the diversity of wetland sub-categories. The statistical analysis showed that the global wetland area reached 6.38 million km², including 6.03 million km² of inland wetlands and 0.35 million km² of coastal tidal wetlands, approximately 72.96 % of which were distributed poleward of 40° N. Therefore, we can conclude that the proposed method is suitable for large-area wetland mapping and that the GWL_FCS30 dataset is an accurate wetland mapping product that has the potential to provide vital support for wetland manage-

ment. The GWL_FCS30 dataset in 2020 is freely available at <https://doi.org/10.5281/zenodo.7340516> (Liu et al., 2022).

1 Introduction

The Ramsar Convention defines wetlands as “areas of marsh, fen, peatland or water, whether natural or artificial, permanent or temporary, with water that is static or flowing, fresh, brackish or salt, including areas of marine water the depth of which at low tide does not exceed six meters” (Gardner and Davidson, 2011). Wetlands not only provide humans with a large amount of food, raw materials and water resources (Ludwig et al., 2019; Z. Zhang et al., 2022) but also play an important role in maintaining ecological balance, conserving water resources, replenishing groundwater and controlling soil erosion (Hu et al., 2017a; Mao et al., 2021; Wang et al., 2020; Zhu and Gong, 2014). Therefore, they are also called the “kidneys of the earth” (Guo et al., 2017). However, due to increasing human activities, including agriculturalization, industrialization and urbanization (McCarthy et al., 2018; Xi et al., 2020), and climatic changes, such as sea-level rise and coastal erosion (Cao et al., 2020; Wang et al., 2021), wetlands have been seriously degraded and threatened over the past few decades (Mao et al., 2020). Thus, having access to timely and accurate wetland mapping information is pivotal for protecting biodiversity and supporting the sustainable development goals.

Along with the rapid development of remote sensing techniques and computing abilities, a variety of regional and global wetland datasets have been produced with spatial resolutions ranging from 30 m to 1° (~ 112 km) (Chen et al., 2022; Gumbrecht et al., 2017; Lehner and Döll, 2004; Mao et al., 2020; Matthews and Fung, 1987; Tootchi et al., 2019). Recently, Tootchi et al. (2019) and Hu et al. (2017a) have systematically reviewed the generation process of global wetland datasets with various spatial and temporal resolutions and wetland categories and found significant uncertainties and inconsistencies among these datasets. For example, the global total wetland area reviewed by Hu et al. (2017a) ranged from 2.12 to 7.17 million km² based on remote sensing products. Therefore, great uncertainties among global wetland datasets directly hindered wetland applications and analysis. Furthermore, from the perspective of spatial resolution, although many wetland products have been produced at regional or global scales using various remote sensing imagery and different methods (Guo et al., 2017; Tootchi et al., 2019), most of them were coarse-spatial-resolution datasets, ranging from 100 m to 25 km. Recently, with the improvement in computing power and storage abilities, three global 30 m land-cover products (including GlobeLand30; Chen et al., 2015; FROM_GLC; Gong et al., 2013; and GLC_FCS30; Zhang et al., 2021b) and several 10 m land-cover products

(WorldCover; Zanaga et al., 2021; Dynamic World; Brown et al., 2022; and FROM_GLC10; Gong et al., 2019), containing an independent wetland layers, were produced, but their classification algorithms were not specifically designed for the wetland environment, so wetlands usually suffered from low accuracy in these products. In addition, several global coastal tidal wetland products have been developed, including the global mangrove extent (Bunting et al., 2018; Hamilton and Casey, 2016) and global 30 m tidal flat datasets from 1984 to 2016 (Murray et al., 2019), but these only covered the intertidal zones. Thus, an accurate global 30 m thematic wetland dataset, with fine wetland categories and covering both inland and coastal zones, is still lacking.

One of the largest challenges of current state-of-the-art methods for large-area wetland mapping is to collect a massive number of training samples (Liu et al., 2021; Ludwig et al., 2019). Zhang et al. (2021b) mentioned two options for collecting training samples, including the visual interpretation method and deriving training samples from pre-existing products. First, since the visual interpretation method had significant advantages over the confidence of training samples, it was widely used for local or regional wetland mapping (Amani et al., 2019; Wang et al., 2020). However, collecting accurate and sufficient training samples is usually a time-consuming process and involves a large amount of manual work, so it was impractical and nearly impossible to use the visual interpretation for collecting global wetland samples. Comparatively, the process of deriving training samples from existing products and applying some rules or refinement methods to identify these high-confidence samples from existing products shows promise (Zhang et al., 2021b). So this approach is practical in that it could quickly produce a large and geographically diverse distribution of training samples without much manual effort. Thus, the second option attracted increasing attention and has been successfully used for large-area land-cover mapping (Zhang and Roy, 2017; Zhang et al., 2020, 2021b). For example, Zhang et al. (2021b) used global training samples derived from the combination of the CCI_LC and MCD43A4 Distribution Function Adjusted Reflectance (NBAR) datasets to produce a global 30 m land-cover product with a fine classification system in 2015 and 2020 (GLC_FCS30) with an overall accuracy of 82.5 %. Therefore, if we take effective measures to fuse these existing products and then derive high-confidence training samples using some refinement rules, the deriving approach would hold great potential for global wetland mapping.

Another major challenge inherent to wetland mapping is the complicated temporal dynamics and the spatial and spectral heterogeneity. The spectral characteristics of the wet-

lands would quickly change with the seasonal or daily water levels of the underlying surface (Ludwig et al., 2019; Mahdianpari et al., 2020). Therefore, many studies proposed to combine multi-sourced, time-series remote sensing imagery to capture the spatial extent and temporal dynamics of wetlands (LaRocque et al., 2020; Ludwig et al., 2019; Murray et al., 2019; Wang et al., 2021; Z. Zhang et al., 2022). For example, Z. Zhang et al. (2022) and Murray et al. (2019) used the Landsat time-series imagery to generate tidal-level and phenological features for identifying coastal tidal wetlands and successfully produced the coastal tidal wetlands in China with an overall accuracy of 97.2 % (Z. Zhang et al., 2022) and global trajectory tidal flats with the overall map accuracy of 82.3 % (Murray et al., 2019). Except for optical imagery, synthetic aperture radar (SAR) data, which were sensitive to soil moisture, vegetation structure and inundation, enabled data acquisition regardless of solar illumination, clouds or haze and were also widely used for wetland mapping, especially after the Sentinel-1 data became open-access (Li et al., 2020; Slagter et al., 2020; Zhang et al., 2018). For example, Li et al. (2020) used the Sentinel-1 time-series imagery to discriminate wetlands with and without trees and achieved an overall accuracy of 86.0 ± 0.2 %. Therefore, the fusion of multi-sourced and time-series remote sensing imagery is vital for accurate wetland mapping.

Due to the complicated temporal dynamics and the spatial and spectral heterogeneity of wetlands, there are very few global thematic wetland datasets covering both inland and coastal regions with a fine classification system and high spatial resolution, which also cause global 30 m wetland mapping with a fine classification system to remain a challenging task. In this study, we combined several existing wetland products and multi-sourced time-series remote sensing imagery to (1) derive large and geographically distributed wetland training samples from pre-existing multi-sourced global wetland products to minimize the manual participation; (2) develop a robust method to capture the temporal dynamics of wetlands and then produce the first global 30 m wetland dataset with a fine classification system (GWL_FCS30); and (3) quantitatively analyze the spatial distribution of different wetland categories and assess the accuracy of GWL_FCS30 in 2020.

2 Datasets

2.1 Multi-sourced remote sensing imagery

Three types of remote sensing imagery were collected to capture the temporal dynamics and spatial and spectral heterogeneity of wetlands. These include Landsat optical data and Sentinel-1 SAR and ASTER Global Digital Elevation Model (GDEM) topographical data. First, all available Landsat imagery, including Landsat 7 Enhanced Thematic Mapper Plus (ETM+) and Landsat 8 Operational Land Imager (OLI) missions, during 2019–2021 was obtained for the nominal year

of 2020 via the Google Earth Engine platform for minimizing the influence of frequent cloud contamination in the tropics and snow and ice in the high latitudes. To minimize the effect of atmosphere, each Landsat image was atmospherically corrected to the surface reflectance by the United States Geological Survey using the Land Surface Reflectance Code (LaSRC) method (Vermote et al., 2016) and then archived on the Google Earth Engine (GEE) platform. These “bad-quality” observations (shadow, cloud, snow and saturated pixels) in Landsat imagery were masked using the CFmask cloud detection method, which built a series of decision rules, using temperature, spectral variability, brightness and geometric relationship between cloud and shadow, to identify these “poor-quality” pixels and achieved the overall accuracy of 96.4 % (Zhu et al., 2015; Zhu and Woodcock, 2012). In this study, six optical bands, including blue, green, red, NIR (near infrared), SWIR1 (shortwave infrared 1) and SWIR2 (shortwave infrared 2) bands, were used for wetland mapping. In total, 764 239 Landsat scenes were collected to capture various water-level and phenological features according to the spectral characteristics of various land-cover types, presented in Sect. 4. Figure 1a illustrates the spatial distribution of all clear-sky observations for all Landsat scenes, and it can be seen that there were more than 10 clear observations after masking these “poor-quality” observations in each region and even in the tropics.

Then, the Sentinel-1 SAR data, which were demonstrated to be sensitive to the soil moisture, vegetation structure and inundation information (Li et al., 2020), used dual-polarization C-band backscatter coefficients to measure the incident microwave radiation scattered by the land surface (Torres et al., 2012). This study obtained the Sentinel-1 time-series imagery archived on the GEE platform in 2020 in Interferometric Wide swath mode with a dual-polarization of VV and VH. Notably, all Sentinel-1 SAR imagery on the GEE platform has been pre-processed by the Sentinel-1 Toolbox with thermal noise removal, radiometric calibration and terrain correction using 30 m elevation data (Veci et al., 2014). Figure 1b also illustrates the spatial distribution of all available Sentinel-1 SAR imagery; there were enough Sentinel-1 SAR observations in each area to capture the water-level dynamics of wetlands because it was immune to cloud and shadow and had a revisit time of 6 d after launching the Sentinel-1B mission. Lastly, as many studies have demonstrated that the topography would directly affect the spatial distribution of wetlands, which are mainly distributed in low-lying areas (Hu et al., 2017b; Ludwig et al., 2019; Tootchi et al., 2019), the ASTER GDEM elevation and derived slope and aspect were used as auxiliary information for wetland mapping. It had a spatial resolution of 30 m and covered the entire global land area (Tachikawa et al., 2011a). Quantitative assessment indicated that the GDEM achieved an absolute vertical accuracy of 0.7 m over bare areas and 7.4 m over forested areas (Tachikawa et al., 2011b).




Figure 1. The spatial distribution of clear observations after masking these “poor-quality” observations during 2019–2021 (a) and availability of Sentinel-1 SAR time-series observations in 2020 (b).

2.2 Prior global wetland datasets

To achieve the goal of deriving a large and geographically diverse distribution of training samples with minimum manual labor, we propose combining various prior global wetland datasets for generating high-confidence training samples. Table 1 lists the characteristics of several global wetland datasets. Specifically, we collected five global mangrove forest products with different spatial resolutions and time spans, and all of them achieved desirable accuracy. For example, the Global Mangrove Watch (GMW) was validated to reach an overall accuracy of 95.25 %, and the user and producer accuracies of mangrove forest were 97.5 % and 94.0 %, respectively (Thomas et al., 2017). Furthermore, to derive the samples of salt marsh and tidal flats, we collected the global 30 m tidal flat time-series products from 1984 to 2016

with an interval of 3 years, achieving an overall map accuracy of 82.3 % (Murray et al., 2019). The global salt marsh dataset, containing 350 985 individual occurrence polygon shapefiles, helped generate the global salt marsh estimation (McOwen et al., 2017).

Except for the coastal tidal wetland products, two thematic wetland products (TROP-SUBTROP Wetland and Global Lakes and Wetlands Database (GLWD) contained various wetland sub-categories), three global land-cover products (GlobeLand30, GLC_FCS30 and CCI_LC contained an independent layer) and the 30 m water dynamic time-series dataset (JRC_GSW) were combined to determine the inland maximum wetland extents and generate the wetland training samples after using a series of refinement rules given in Sect. 3. Specifically, the TROP-SUBTROP was produced by combining the hydrological model and annual time series

of soil moisture, mainly covering the tropics and subtropics (40° N–60° S) with a resolution of 231 m (Gumbricht, 2015). The GLWD, combining the GIS functionality and a variety of existing maps and information, was developed with 12 wetland sub-categories at a resolution of 1 km (Lehner and Döll, 2004). The JRC_GSW dynamic water dataset achieved a producer accuracy of 98.5 % for these seasonal waters (Pekel et al., 2016) and was used to identify inundated pixels. Furthermore, three global land-cover products, simultaneously containing wetland and non-wetland land-cover types, were used to determine the non-wetland samples and then served as the auxiliary datasets to improve the confidence of inland wetland samples.

2.3 Global 30 m tree cover product

The global 30 m forest cover change in tree cover (GFCC30TC) data in 2015 was produced by downscaling the 250 m MODIS VCF (Vegetation Continuous Fields) tree cover product using Landsat imagery and then incorporating the MODIS cropland layer to guarantee the tree cover accuracy in agricultural areas (Sexton et al., 2016, 2013). This product was used to accurately distinguish between inland swamp and marsh wetlands because both of them reflected obvious vegetation spectral characteristics. It was validated to achieve an overall accuracy of 91 %; the average producer and user accuracies for stable forests were 92.5 % and 95.4 %, respectively (Sexton et al., 2016; Townshend et al., 2012).

2.4 National wetland products

Three national wetland products, including NLCD (National Land Cover Database) (Homer et al., 2020), NWI (National Wetlands Inventory) (Wilén and Bates, 1995) and CLC (CORINE Land Cover) (Büttner, 2014), were used as the comparative datasets to analyze the performance of developed global wetland maps in Sect. 6.2. Specifically, the NLCD contained open water, woody wetlands and emergent herbaceous wetlands, the NWI contained eight sub-categories (estuarine and marine deepwater, estuarine and marine wetland, freshwater emergent wetland, freshwater forest/shrub wetland, freshwater pond, lake, other and riverine), and the CLC identified the wetlands in 10 sub-categories: inland marshes, peat bogs, salt marshes, saline, intertidal flats, water courses, waterbodies, coastal lagoons, estuaries, and sea and oceans.

3 Collecting training samples and determining maximum wetland extent

In this study, after considering the applicability of moderate resolution (10–30 m) imagery, their practical use for ecosystem management and the available pre-existing global wetland dataset, the fine wetland classification system, contain-

ing eight sub-categories (three coastal tidal sub-categories and five inland sub-categories), was proposed to comprehensively depict the spatial patterns of global wetlands (Table 2). Specifically, the sub-categories of coastal tidal wetlands consist of mangroves, salt marshes and tidal flats. By importing the vegetation and water cover information associated with this land cover, these categories were widely recognized in many previous studies (Wang et al., 2021; Z. Zhang et al., 2022). The inland wetland types shared similar characteristics and were grouped into swamp, marsh and flooded flat. Meanwhile, in order to capture saline soils and halophytic plant species along saline lakes, the inland saline wetland, inherited from the GLWD (Lehner and Döll, 2004), was also imported. Lastly, the permanent water, including lakes, rivers and streams that are always flooded, was widely identified as a wetland layer in previous studies (Davidson, 2014; Dixon et al., 2016; Hu et al., 2017b) and was also added into our fine wetland classification system.

Many studies have explained that the quality and confidence of training samples directly affected the classification performance (Zhang et al., 2021b; Zhu et al., 2016). The previously mentioned process of collecting sufficient training samples via visual interpretation was time-consuming and involved a lot of manual labor. Fortunately, a variety of regional and global wetland products have been developed and released over the past few decades (Table 1), and many studies have demonstrated that deriving training samples from existing products could be used for large-area classification and mapping (Huang et al., 2021; Zhang et al., 2021b). Therefore, we propose to combine existing global wetland datasets to independently derive coastal and inland wetland training samples and their maximum distribution extents (Fig. 2).

3.1 Deriving coastal tidal wetland training samples and maximum extents

This study divided the coastal tidal wetlands into three sub-categories: mangrove forest, salt marsh and tidal flat. The previously existing products have been collected in Table 1. For the mangrove training samples, we collected five global mangrove products with different spatiotemporal resolutions, all of which achieved good performances. For example, Hamilton and Casey (2016) stated that their continuous mangrove forest cover (CGMFC) dataset could cover 99 % of all mangrove forests from 2000 to 2012, and Thomas et al. (2017) validated their Global Mangrove Watch (GMW) products from 1996 to 2016 and reached an overall accuracy of 95.25 %. Therefore, we first measure the temporal consistency of the three mangrove forest time-series products (CGMFC, GMW and GBTM mangroves), and only these temporally stable mangrove forest pixels were selected as the primary candidate points ($P_{\text{mangrove}}^{\text{Tstable}}$). Meanwhile, to minimize the influence of classification error in each mangrove forest product, the cross-consistency of five mangrove prod-

Table 1. The characteristics of 13 global wetland products with various spatiotemporal resolutions (unit of area: million km²).

Dataset name and reference	Wetland categories	Year	Resolution	Total area	Coverage
World Atlas of Mangroves (WAM) Spalding (2010)	Mangrove	2010	1 : 1 000 000	0.152	Global
Global Mangrove Watch (GWM) Thomas et al. (2017)		1996–2016	~ 25 m	~ 0.136	Global
A global biophysical typology of mangroves (GBTM) Worthington et al. (2020)		1996–2016	~ 25 m	~ 0.136	Global
Continuous global mangrove forest cover (CGMFC) Hamilton and Casey (2016)		2000–2010	30 m	0.083	Global
Global Distribution of Mangroves USGS (GDM_USGS) Giri et al. (2011)		2011	30 m	~ 0.138	Global
Global distribution of tidal flat ecosystems Murray et al. (2019)	Tidal flat	1984–2016	30 m	0.124–0.132	60° S–60° N
Global distribution of salt marsh McOwen et al. (2017)	Salt marsh	1973–2015	1 : 10 000	~ 0.05	Global
Tropical and Subtropical Wetland Distribution (CIFOR) Gumbrecht (2015)	Open water, mangrove, swamps, fens, riverine, floodplains, marshes	2011	~ 231 m	4.7	60° S–40° N
Global Lakes and Wetlands Database (GLWD) Lehner and Döll (2004)	Lake, reservoir, river, marsh, swamps, coastal tidal wetland, saline wetland and peatland	2004	~ 1 km	10.7–12.7	Global
JRC_GSW Pekel et al. (2016)	Water	1984–2021	30 m	~ 4.46	Global
ESA CCI_LC Defourny et al. (2018)	Swamps, mangrove, and shrub or herbaceous covered wetlands	1992–2020	300 m	6.1	Global
GlobeLand30 Chen et al. (2015)	Wetland	2000–2020	30 m	7.01–7.17	Global
GLC_FCS30 Zhang et al. (2021b)	Wetland	2015, 2020	30 m	6.36	Global

ucts was analyzed, and only the pixel simultaneously identified as mangrove forest in all five products was labeled as stable and consistent candidate points ($P_{\text{mangrove}}^{\text{Tstable, Scons}}$). Furthermore, considering that there was a temporal interval between prior mangrove products and our study and that mangrove deforestation usually followed the pattern of edge-to-center contraction, a morphological erosion filter with a local window of 3×3 was applied to the $P_{\text{mangrove}}^{\text{Tstable, Scons}}$ points to further ensure the confidence of mangrove training samples. Lastly, as for the maximum mangrove forest extents


($\text{MaxExtent}_{\text{mangrove}}$), the union operation was applied to five global mangrove products as shown in Eq. (1).

$$\text{MaxExtent}_{\text{mangrove}} = M_{\text{WAM}} \cup M_{\text{GMW}} \cup M_{\text{GBTM}} \cup M_{\text{CGMFC}} \cup M_{\text{GDM_USGS}}, \quad (1)$$

where M_{WAM} , M_{GMW} , M_{GBTM} , M_{CGMFC} and $M_{\text{GDM_USGS}}$ are the spatial distributions of five global mangrove forest products listed in Table 1. It should be noted that these prior mangrove products were demonstrated to cover almost all

Table 2. The description of the wetland classification system in this study.

Category I	Category II	Description
Tidal wetland	Mangrove	The forest or shrubs which grow in the coastal brackish or saline water
	Salt marsh	Herbaceous vegetation (grasses, herbs and low shrubs) in the upper coastal intertidal zone
	Tidal flat	The tidal flooded zones between the coastal high and low tide levels including mudflats and sandflats.
Inland wetland	Swamp	The forest or shrubs which grow in the inland freshwater
	Marsh	Herbaceous vegetation (grasses, herbs and low shrubs) grows in the freshwater
	Flooded flat	The non-vegetated flooded areas along the rivers and lakes
	Saline	Characterized by saline soils and halophytic (salt tolerant) plant species along saline lakes
	Permanent water	Lakes, rivers and streams that are always flooded

**Figure 2.** The flowchart of deriving coastal and inland wetland samples from multiple pre-existing datasets.

mangroves over the world, so the $\text{MaxExtent}_{\text{mangrove}}$ can be used as the boundary for mangrove mapping; namely, only the pixel within the maximum mangrove extent was labeled as mangrove forest.

Regarding the collection of tidal flat samples, the prior global 30 m tidal flat time-series products (Gtidalflat) from 1984 to 2016 were validated to achieve an overall map accuracy of 82.3 % and user accuracies for the non-tidal and tidal flat of 83.3 % and 81.1 %, respectively (Murray et al., 2019). To ensure the accuracy of tidal flat samples, we first applied temporal consistency analysis to the time series of tidal flat datasets from 2000 to 2016 and identified the temporally stable tidal flat pixels ($P_{\text{tidal}}^{\text{Tstable}}$) during 16 consecutive years. The reason why we discarded the tidal flat datasets before 2000 was that the available Landsat imagery was sparse and could not accurately capture the high-tidal and low-tidal information and suffered lower monitoring accuracy. Next, Radoux et al. (2014) found that transition zones between

two different land-cover types are likely to be misclassified; therefore, the candidate tidal flat samples $P_{\text{tidal}}^{\text{Tstable}}$ were further refined by the morphological erosion filter with a local window of 3×3 . Furthermore, as a tidal flat is a non-vegetated coastal tidal wetland, we combined the empirical rule (enhanced vegetation index (EVI) ≥ 0.1 , normalized difference vegetation index (NDVI) ≥ 0.2 and land surface water index (LSWI) > 0) proposed by Wang et al. (2020) and Landsat time-series imagery in 2020 (approximately 142 000 Landsat scenes) to exclude all vegetated pixels from tidal flat training samples. Lastly, to derive the maximum tidal flat extents ($\text{MaxExtent}_{\text{tidalflat}}$), the union operation was applied to the tidal flat time-series products from 1984 to 2016. It should be noted that Murray's global 30 m tidal flat datasets only covered the regions of 60°N – 60°S (Murray et al., 2019); therefore, we used the coastal shorelines ($\text{Line}_{\text{coastal}}$) to create a 50 km buffer (applied by Wang et al., 2020, and Murray et al., 2019) as the potential tidal flat zones in the

high-latitude regions ($> 60^\circ \text{N}$) as in Eq. (2). It should be noted that we only identified and then retained these tidal flat pixels within the maximum extents by using the classification models in Sect. 4.2.

$$\text{MaxExtent}_{\text{tidalflat}} = \left\{ \begin{array}{l} \bigcup_{t=1984}^{2016} \text{Gtidalflat}_{t,s}, \quad s \in [60^\circ \text{S}, 60^\circ \text{N}] \\ \text{Line}_{\text{coastal}} \pm 50 \text{ km}, \quad s \in [60^\circ \text{N}, 90^\circ \text{N}] \end{array} \right. \quad (2)$$

Compared with mangrove forest and tidal flat, the pre-existing global or regional salt marsh products were relatively sparse. The global distribution of the salt marsh dataset contained 350 985 individual vector polygons and was the most complete dataset on salt marsh occurrence and extent at the global scale (McOwen et al., 2017). However, after careful review, we found some mislabeled salt marsh polygons, so this dataset cannot be used directly to derive training samples. This study first used the random sampling method to generate 35 099 salt marsh points (approximately 10 % of the total polygons) based on prior datasets. We combined the visual interpretation method and high-resolution imagery to check each salt marsh point. After discarding the incorrect and uncertain samples, a total of 32 712 salt marsh points were retained. However, the prior dataset only captured the extent of salt marshes in 99 countries worldwide (McOwen et al., 2017), further noting that the distribution of salt marshes was spatially correlated with tidal flat and mangrove forest (Wang et al., 2021). Consequently, the maximum extents of tidal flat and mangrove forest, in addition to the prior salt marsh extent, were used for salt marsh mapping. Meanwhile, as the wetland layer in the global land-cover products (GLC_FCS30, GlobeLand30 and CCI_LC) also covered some coastal tidal wetlands, the saline-water wetland layer in the CCI_LC and the wetland data in the other two products close to the coastal shorelines were also imported as supplementary material when determining the maximum coastal tidal wetland extents.

3.2 Deriving inland wetland training samples and maximum extents

The pre-existing inland wetland datasets usually suffered from lower accuracy compared to coastal tidal wetland products; for example, the wetland layer in GlobeLand30-2010 and GLC_FCS30-2015 was validated to achieve a user accuracy of 74.9 % (Chen et al., 2015) and 43.4 % (Zhang et al., 2021b), respectively. Therefore, we first generated high-confidence inland wetland samples and then determined their sub-categories (swamp, marsh, inland flat, saline wetland and permanent water). Specifically, the consistency analysis of five global wetland datasets (TROP-SUBTROP Wetland, GLWD, CCI_LC, GlobeLand30 and GLC_FCS30) and the temporal stability checking for CCI_LC (1992–2020), GlobeLand30 (2000–2020) and GLC_FCS30 (2015–2020) were applied to identify these temporally stable and high-cross-consistency wetland points ($P_{\text{inlandWet}}^{\text{Tstable,Scns}}$). It should be noted that the coarse wetland products (GLWD, TROP-SUBTROP

and CCI_LC) were resampled to 30 m using the nearest-neighbor method on the GEE platform, and the coastal tidal wetland layers in these products were excluded. Namely, only the pixel identified as inland wetland in all five products was retained. Then, the morphological erosion filter with a local window of 3×3 was also used to decrease the sampling uncertainty over these land-cover transition areas because the transition zones between two different land-cover types are likely to be misclassified (Lu and Wang, 2021; Radoux et al., 2014).

Afterward, to determine the wetland sub-category for each inland wetland sample, we first used the empirical vegetation rule ($\text{EVI} \geq 0.1$, $\text{NDVI} \geq 0.2$ and $\text{LSWI} > 0$) proposed by Wang et al. (2020) and Landsat time-series imagery to split candidate samples into two parts: vegetated wetland samples (swamp and marsh) and non-vegetated wetland samples (flooded flat, saline and permanent water). Then, as the swamp was defined as the forest or shrubs which grow in the inland freshwater, the global 30 m tree cover dataset (GFCC30TC) was adopted to distinguish the swamp and marsh from vegetated wetland samples. Specifically, if the tree cover of the sample was greater than 30 % (Hansen et al., 2013), it was labeled as swamp, and the remaining vegetated wetland samples were labeled as marsh. Furthermore, to distinguish between the inland flat, saline samples and permanent water, the saline blocks in the prior GLWD products were first checked by visual interpretation and then imported as the reference dataset to identify all saline wetland samples. The remaining non-vegetated wetland samples were further refined using the time series of the JRC_GSW datasets; only the remaining samples whose water probability was less than the threshold of 0.95 (suggested by Wang et al., 2020) were labeled as flooded flat. Lastly, regarding the permanent water samples, the JRC_GSW water dynamic dataset was validated and achieved producer and user accuracies of 99.7 % and 99.1 % for permanent water (Pekel et al., 2016). The permanent water training samples were directly derived from the JRC_GSW dataset without any refinement rules.

Lastly, as for determining the maximum inland wetland extents ($\text{Mextent}_{\text{inWet}}$), the union operation was conducted with six pre-existing global wetland datasets as in Eq. (3).

$$\text{Mextent}_{\text{inWet}} = W_{\text{TROP-SUBTROP}} \cup W_{\text{GLWD}} \cup W_{\text{CCI_LC}} \cup W_{\text{GLC_FCS30}} \cup W_{\text{GlobeLand30}} \cup W_{\text{JRC_GSW}} \quad (3)$$

Here $W_{\text{TROP-SUBTROP}}$, W_{GLWD} , $W_{\text{CCI_LC}}$, $W_{\text{GLC_FCS30}}$ and $W_{\text{GlobeLand30}}$ are wetland distributions of five pre-existing global wetland products, and $W_{\text{JRC_GSW}}$ is the JRC_GSW water dynamic time-series datasets, which identified the inundated probability at a monthly scale during 1984–2021 (Pekel et al., 2016). It should be noted that the omission error can be ignored for derived maximum inland wetland extents ($\text{Mextent}_{\text{inWet}}$) because the GLWD and TROP-SUBTROP

Wetland datasets captured almost all potential wetlands using compilation and model simulation methods (Gumbrecht, 2015; Lehner and Döll, 2004).

3.3 Deriving non-wetland training samples from prior land-cover products

Except for inland and coastal tidal wetland samples, the non-wetland samples were also necessary because some non-wetland land-cover types were shown to have a similar spectrum to wetlands. For example, swamp and forest or shrubs exhibited the same vegetation reflectance characteristics in optical imagery, and marsh and grassland shared similar spectral curves during the growing season (Z. Zhang et al., 2022). Except for eight fine wetland sub-categories training samples, we also divided the non-wetlands into forest/shrubland, grassland, cropland and others (bare land, impervious surfaces and snow). To automatically derive these non-wetland samples, the multi-epoch GlobeLand30, GLC_FCS30 and CCI_LC global land-cover products were integrated. Specifically, the temporal stability and cross-consistency analyses were applied to three land-cover products to identify temporally stable forest/shrubland, grassland, cropland and other candidate samples. Furthermore, the morphological erosion filter with the local window of 3×3 was also adopted to decrease the sampling uncertainty over land-cover transition areas.

3.4 Determining the sample size and distributions using stratified random sampling strategy

Except for the confidence of training samples, many studies also found that the size and distribution of training samples also affected classification performances (Jin et al., 2014; Zhu et al., 2016). As this study aimed to identify wetlands instead of all land-cover types, the equal allocation sample distribution would perform better than the proportional distribution (the sample size determined by the area) (Jin et al., 2014; Zhang et al., 2020). Namely, the approximate proportion of inland wetland, coastal tidal wetland and non-wetland samples was 5 : 3 : 4 in the coexisting areas because the classification system was composed of five inland and three coastal tidal wetland sub-categories and four non-wetland land-cover types. Regarding the sample size, Zhu et al. (2016) had analyzed the quantitative relationships of sample size and the mapping accuracy and found that the mapping accuracies slowly increased and then remained stable with any further increase in the number of samples and suggested using a total of 20 000 samples in the Landsat scene. In this study, we used the stratified random sampling strategy to collect the training samples (excluding salt marsh because it was collected globally using visual interpretation in Sect. 3.1) at each $5^\circ \times 5^\circ$ geographical grid (corresponding to the local adaptive modeling in Sect. 4.2) using an approximate sample size of 2000 for each category. According to

our statistics, this study derived more than 20 million training samples for mapping global fine wetlands.

4 Mapping wetland using the stratified classification strategy and the water-level and phenological features

Figure 3 illustrates the flowchart of the proposed method for generating the global 30 m fine wetland maps. First, we combined the Landsat 8 and Sentinel-1 SAR time-series observations and ASTER DEM topographical images to derive multi-sourced and multi-temporal features, including three topographical and various water-level and phenological features. Then, the training samples (coastal tidal, inland wetlands and non-wetlands) and derived multi-sourced and multi-temporal features were combined to train the stratified random forest classifiers (a classic and widely used machine learning classification model; Breiman, 2001) at each local region. Next, using the trained random forest models and derived multi-sourced and multi-temporal features, we could develop corresponding coastal tidal wetland and inland wetland maps. Finally, the post-processing step was used to generate the global 30 m fine wetland map in 2020.

4.1 Generating various water-level and phenological composites

Before generating various water-level and phenological features, four spectral indices, including normalized difference water index (NDWI), LSWI, NDVI and EVI, were imported because many studies have demonstrated that they were of great help in wetland mapping (Mao et al., 2020; Wang et al., 2020).

$$\begin{aligned} \text{LSWI} &= \frac{\rho_{\text{nir}} - \rho_{\text{swir1}}}{\rho_{\text{nir}} + \rho_{\text{swir1}}}, & \text{NDWI} &= \frac{\rho_{\text{green}} - \rho_{\text{swir1}}}{\rho_{\text{green}} + \rho_{\text{swir1}}}, \\ \text{NDVI} &= \frac{\rho_{\text{nir}} - \rho_{\text{red}}}{\rho_{\text{nir}} + \rho_{\text{red}}}, \\ \text{EVI} &= 2.5 \times \frac{\rho_{\text{nir}} - \rho_{\text{red}}}{\rho_{\text{nir}} + 6 \times \rho_{\text{red}} - 7.5 \times \rho_{\text{blue}} + 1}, \end{aligned} \quad (4)$$

where ρ_{blue} , ρ_{green} , ρ_{red} , ρ_{nir} and ρ_{swir1} are the blue, green, red, near-infrared and shortwave infrared bands of Landsat imagery, respectively.

Then, the spectral characteristics of the wetlands would quickly change along with the seasonal or daily water levels of the underlying surface. For example, the tidal flat was the status of seawater at the high tidal stage and mudflats or sandflats at low tidal stages (Wang et al., 2021); therefore, it was necessary to extract the highest- and lowest-water-level composites to completely capture these inundated wetlands. Over the past several years, the time-series compositing strategy has been widely used to capture phenological and cloud-free composites (Jia et al., 2020; Ludwig et al., 2019; Murray et al., 2019; Zhang et al., 2021a). In this study, considering that NDWI was sensitive to open surface water and that




Figure 3. The flowchart of wetland mapping using water-level, phenological and topographical features and a stratified classification strategy.

Z. Zhang et al. (2022) found a positive relationship between tidal height and NDWI using field survey data, the maximum NDWI compositing was applied to the clear-sky Landsat time-series imagery to capture the highest-water-level optical composites illustrated in Fig. 4b. As for the lowest water-level features, considering that the tidal and flooded flat or marsh usually reflected higher NDVI and EVI values than waterbodies and that Z. Zhang et al. (2022) also used the field data to demonstrate that there was a negative relationship between tidal-level height and NDVI, the maximum NDVI composite was applied to capture the lowest-water-level optical information illustrated in Fig. 4a. Considering that optical observations were usually contaminated by clouds, especially during the rainy seasons, and that the SAR back coefficients had a great advantage in the presence of cloud coverage and were found to be sensitive to the soil moisture, vegetation structure and inundation information, the Sentinel-1 SAR time-series imagery could be used as a complementary dataset for capturing the highest- and lowest-water-level composites (DeVries et al., 2020; Li et al., 2020; Mahdianpari et al., 2018). Specifically, as the SAR active transmitting signals were heavily absorbed when they reached the waterbody, the corresponding SAR back coefficients in the waterbody had lower values compared to other land-cover types. To capture the high-water-level features from the Sentinel-1 time-series imagery, the percentile compositing method using the 5th percentile was applied, as illustrated in Fig. 4d. Conversely, the 95th percentiles of Sentinel-1 VV and VH were generated to capture the lowest-water-level information (Fig. 4c). It should be noted that the minimum and maximum percentiles were not used because the Sentinel-1 time-series imagery still contained the residual errors caused by the quantitative processing.

Many studies also demonstrated that a multi-temporal phenology was also essential for classifying the vegetated wetlands and excluding these non-wetland land-cover types (Li et al., 2020; Ludwig et al., 2019). There were usually two options for capturing phenological features from Landsat time-series imagery. These included seasonal-based compositing (Zhang et al., 2021a, 2022) and percentile-based compositing (Hansen et al., 2014; Zhang and Roy, 2017; Zhang et al., 2021b). The former used the phenological calendar for selecting time-matched imagery. It then adopted the compositing rule to capture the seasonal features, while the latter directly used the statistical distributions to select various percentiles. Azzari and Lobell (2017) quantitatively analyzed the performance of two compositing methods and found that both of them had similar mapping accuracy for land-cover mapping. Meanwhile, the seasonal-based compositing method needed the prior phenological calendar, while the percentile compositing method did not require any prior knowledge or explicit assumptions regarding the timing of the season; therefore, the percentile compositing method was more suitable for generating phenological features. This study compositing Landsat reflectance time-series bands and four spectral indices into five percentiles (15th, 30th, 50th, 70th and 85th) because we wanted to capture as many of the phenological changes in wetlands as possible when comparing with the four seasonal composites (Zhang and Roy, 2017). It should be noted that the minimum and maximum percentiles were excluded because they were usually affected by residual clouds, shadows and saturated observations.

Lastly, the topographical variables were also important factors for determining the spatial distribution of wetlands (Ludwig et al., 2019; Tootchi et al., 2019). For example, the widely used topographical wetness index (TWI) uses the local slope to reveal soil wetness, which improves wetland




Figure 4. The lowest- and highest-water-level features derived from (a–b) Landsat optical reflectance time-series data and (c–d) the Sentinel-1 SAR imagery using the time-series compositing method in Poyang Lake, China.

classification performance and reduces commission errors within upland areas (Ludwig et al., 2019). Therefore, the elevation, aspect and slope, calculated from the ASTER GDEM dataset, were included in the multi-sourced features. In summary, there are a total of 77 multi-sourced training features (listed in Table 3), including 70 optical features from Landsat imagery, 4 SAR features from Sentinel-1 imagery and 3 topographical features from ASTER GDEM.

4.2 The stratified classification strategy for wetland mapping

Since we have simultaneously extracted the maximum coastal and inland wetland extents when deriving training samples from prior wetland datasets, the stratified classification strategy was adopted to fully use the maximum extent constraint. If a pixel was classified as a coastal tidal wetland outside the maximum coastal tidal wetland extents, it would be identified as a misclassification. Furthermore, there were two ideas for the large-area land-cover mapping, including global classification modeling (using one universal model for entire areas) and local adaptive modeling (using various models for different local zones) (Zhang et al., 2020). For example, Zhang and Roy (2017) demonstrated that local adaptive modeling outperformed the global classification modeling strategy. Therefore, the global land surface was first divided into 961 $5^\circ \times 5^\circ$ geographical tiles illustrated in Fig. 5, which were inherited from the global 30 m land-cover mapping by Zhang et al. (2021b). Then, we trained the local adaptive classification models using derived training samples in Sect. 3 and multi-sourced and multi-temporal features (the highest water level, lowest water level, phenological composites and topographical variables) at each $5^\circ \times 5^\circ$ geographical tile. It should be noted that we used the training samples from neighboring 3×3 geographical tiles to train the classification model and classify the central tile for guaranteeing the spatially continuous transition over adjacent regional wetland maps. Namely, we trained 961 local adaptive classification models and then produced 961 $5^\circ \times 5^\circ$ wetland maps.


Finally, we spatially mosaicked these 961 regional wetland maps into the global 30 m wetland map in 2020.

Afterward, the random forest (RF) classifier was demonstrated to have obvious advantages, including dealing with high-dimensional data, robustness for training noise and feature selection, as well as achieving higher classification when compared to other widely used machine learning classifiers (e.g., support vector machines, neural networks, decision trees) (Belgiu and Drăguț, 2016; Gislason et al., 2006). Therefore, the RF classifier was selected for mapping inland and coastal tidal wetlands using multi-sourced features on the GEE platform. It should be noted that the RF classifier had two key parameters: the number of selected prediction variables (M_{try}) and the number of decision trees (N_{tree}). Belgiu and Drăguț (2016) and Z. Zhang et al. (2022) have demonstrated the quantitative relationship of N_{tree} against classification accuracy and found that the classification accuracy stabilized when N_{tree} was greater than 100. Meanwhile, Belgiu and Drăguț (2016) suggested that M_{try} should take its default value of the square root of the number of all input features. Therefore, N_{tree} and M_{try} took 100 and the square root of the number of all input features, respectively.

The inland and coastal tidal wetland maps were produced by combining water-level and phenological features, the stratified classification strategy, local adaptive modeling, and the derived wetland and non-wetland training samples. As the inland and coastal tidal wetlands were independently produced, some pixels in the overlapping area of maximum inland and coastal tidal wetland extents were simultaneously labeled as inland wetlands and coastal tidal wetlands. However, as the final global wetland map was a hard classification, these pixels should be post-processed into one label. As the random forest classifier could provide the posterior probability for each pixel, we determined the labels of the confused pixels by comparing the posterior probabilities. In addition, as the tidal flats were demonstrated to overestimate some coastal ponds as tidal flats, the global lake and reservoir dataset, developed by Khandelwal et al. (2022), was applied to optimize the tidal flat.

Table 3. The multi-sourced and multi-temporal training features for wetland mapping.

Data	Derived training features from multi-sourced remote sensing imagery
Landsat	Water-level features: the lowest and highest composites with blue, green, red, NIR, SWIR1, SWIR2, LSWI, NDWI, NDVI and EVI bands Phenological features: 15th, 30th, 50th, 70th and 85th percentiles with blue, green, red, NIR, SWIR1, SWIR2, LSWI, NDWI, NDVI and EVI bands
Sentinel-1 SAR	Water-level features: the lowest and highest composites using 5th and 95th percentiles for VV and VH bands.
ASTER GDEM	Topographical features: elevation, slope and aspect.

**Figure 5.** The spatial distribution of 961 $5^\circ \times 5^\circ$ geographical tiles used for local adaptive modeling, which was inherited from the global 30 m land-cover mapping by Zhang et al. (2021b). The background imagery came from the National Aeronautics and Space Administration (<https://visibleearth.nasa.gov>, last access: 10 November 2022).

4.3 Accuracy assessment

To quantitatively analyze the performance of our GWL_FCS30 wetland map, a total of 25 709 validation samples (illustrated in Fig. 6), including 10 558 non-wetland points and 15 151 wetland points, were collected. Firstly, as the wetland was a sparse land-cover type compared to the non-wetlands (forest, cropland, grassland and bare land), the stratified random strategy was applied to randomly derive validation points at each stratum as

$$n_i = n \times \frac{W_i \times p_i (1 - p_i)}{\sum W_i \times p_i (1 - p_i)},$$

$$n = \frac{(\sum W_i \sqrt{p_i (1 - p_i)})^2}{V + \sum W_i p_i (1 - p_i) / N}, \quad (5)$$

where W_i and p_i are the area proportion and expected accuracy of class i , n_i and n are the sample size of class i and total sample size, V is the standard error of the estimated overall accuracy, and N is the number of pixel units in the study region. Then, as the wetlands had a significant correlation with the water levels (Z. Zhang et al., 2022), the time-series optical observations archived on the GEE cloud platform were used as the auxiliary dataset to interpret these water-level-sensitive wetlands, such as tidal flat and flooded flat. It should be noted that the visual interpretation was implemented on the GEE cloud platform because it archives a large number of satellite images with various time spans and spatiotemporal resolution (X. Zhang et al., 2022). Meanwhile, each validation point is independently interpreted by five experts for minimizing the effect of an expert's subjective knowledge,

and only these complete agreement points were retained; otherwise, they were discarded. Then, we employed four metrics typically used to evaluate accuracy, which include the kappa coefficient, overall accuracy, user accuracy (measuring the commission error) and producer accuracy (measuring the omission error) (Gómez et al., 2016; Olofsson et al., 2014), for calculations using 25 709 global wetland validation samples.

5 Results

5.1 The reliability analysis of derived training samples

This study proposed combining pre-existing multi-sourced wetland products, refinement rules and expert knowledge to automatically derive these massive inland and coastal tidal wetland training samples globally. To demonstrate the reliability of the derived training samples for wetland mapping, we randomly selected approximately 10 000 points from the sample pool and checked their confidence using visual interpretation. It should be noted that we cannot check all the training samples because the number of derived samples was massive (exceeding 20 million training samples in Sect. 3). After a point-to-point inspection, these selected training samples achieved an overall accuracy of 91.53 % in 2020. Meanwhile, we also used 10 000 selected wetland training samples and many non-wetland samples to analyze the overall and producer accuracies of coastal and inland wetlands vs. number of erroneous training samples. Specifically, we gradually increased the “contaminated” samples by randomly altering the label of a certain percentage of training samples in steps of 0.01 and then used these “contaminated” samples to build the RF classification model. After repeating the process 100 times, the quantitative relationship between mapping accuracies and erroneous samples is illustrated in Fig. 7. Obviously, the overall accuracy and producer accuracy of wetlands (merging seven sub-categories into one wetland) were insensitive to the erroneous training samples when the percentage of erroneous samples was controlled within 20 %. Beyond this threshold, the accuracies slowly decreased along with the increase in erroneous training samples. Similarly, previous studies by Zhang et al. (2021b, 2022) quantitatively analyzed the relationship between overall accuracy and the erroneous training sample size. They found that the overall accuracy stabilized when the percentage of erroneous training samples was controlled within the threshold and then rapidly decreased after exceeding the threshold. Therefore, the derived training samples in Sect. 3 were accurate enough to support large-area fine wetland mapping.

5.2 The importance of multi-sourced phenological features for wetland mapping

The complicated temporal dynamics and spectral heterogeneity caused great uncertainties in wetland mapping be-

cause their spectral characteristics quickly changed with the seasonal or daily water levels of the underlying surface (Ludwig et al., 2019). To quantitatively analyze the importance of these multi-sourced and multi-temporal features, we used the random forest classification model, which calculated the increased mean squared error by permuting the out-of-bag data of a variable while keeping the remaining variables constant (Breiman, 2001; Zhang et al., 2020) in an effort to compute their importance. Figure 8 illustrates the importance of all multi-sourced and phenological features, and it can be found that the phenological features which made the most significant contribution mainly did so because they used the multi-temporal percentiles to comprehensively capture vegetation phenology (EVI and NDVI) and water-level dynamics (NDWI and LSWI) for the various land-cover types. Then, the combination of optical and Sentinel-1 SAR water-level features was ranked as the second-most-important role in distinguishing the fine wetlands and non-wetlands. Based on the lowest- and highest-water-level features in Fig. 4, the highest- and lowest-water-level features greatly contributed to determining these water-sensitive wetlands (marsh, tidal flat and flooded flat). For example, Z. Zhang et al. (2022) quantitatively analyzed the contribution of multi-sourced features to mapping accuracy. They found that importing water-level features significantly improved the ability to separate tidal flats from non-wetlands. Lastly, three topographical variables also contributed to wetland mapping because the spatial distribution of wetlands had a significant relationship with topography and was mainly distributed in low-lying areas (Zhu and Gong, 2014).

5.3 The spatial pattern of global wetlands in 2020

Figure 9 illustrates the spatial distributions of our GWL_FCS30 wetland map and their area statistics in latitudinal and longitudinal directions in 2020. Overall, the GWL_FCS30 map accurately captured the spatial patterns of wetlands. It mainly concentrated on the high-latitude areas in the Northern Hemisphere and the rainforest areas (Congo Basin and Amazon rainforest in South America). Quantitatively, according to the latitudinal statistics, approximately 72.96 % of wetlands were distributed poleward of 40° N (a large number of wetlands are located in Canada and Russia), and 10.6 % of wetlands were located in equatorial areas, between 10° S–10° N, within which the Congo and Amazon rainforest wetlands are located. As for the longitudinal direction, there were mainly four statistical peak intervals: 120–50° W (Canada wetlands and Amazon wetlands), 15–25° E (Congo wetlands), 40–55° E (the Caspian Sea) and 60–90° E (Russia wetlands). Afterward, to more intuitively understand the performance of our GWL_FCS30 wetland map, four local enlargements in Florida, the Congo Basin, Sundarbans and Poyang Lake were also illustrated. All of them comprehensively captured the wetland patterns in these local areas. For example, there was significant consistency be-




Figure 6. The spatial distribution of 25 709 global wetland validation samples using the stratified sampling strategy.




Figure 7. The relationship between mapping accuracies with the percentage of erroneous training samples with a step of 1 %.

tween our results and Hansen's regional wetland maps in the Congo Basin (Bwangoy et al., 2010); both results indicated that the wetlands occurred closer to major rivers and floodplains. Next, according to the lowest- and highest-water-level features derived from Sentinel-1 SAR and Landsat optical imagery in Fig. 4, the inland wetlands, varied with the water levels, were also comprehensively identified in the Poyang wetland map (Fig. 9d). Figure 9c illustrates the spatial distributions of the world's largest mangrove forest in the Sundarbans (Fig. 9c), and the cross-comparison in Fig. 14 also demonstrates the great performance of the GWL_FCS30 dataset. Lastly, the Florida wetlands simultaneously contained six sub-categories (mangrove, tidal flat, salt marsh, marsh, permanent water and swamp). These were distributed along the coastlines and rivers and are accurately captured in Fig. 9a.

Figure 10 illustrates the spatial distribution of eight wetland sub-categories after aggregating to the $0.5^\circ \times 0.5^\circ$ grid cell. Intuitively, permanent waterbody, swamp and marsh accounted for most inland wetlands, while the flooded tidal and inland saline wetlands had obviously lower proportions, and the latter was only distributed along the surroundings of several saline lakes. In terms of the spatial distribution, it can be found that (1) the swamp wetlands mainly were concentrated in the Congo and Amazon rainforests, southern United States, and northern Canada; (2) most marsh wetlands were located in high-latitude areas in the Northern Hemisphere including northern Canada, Russia and Sweden; and (3) there were significant coexistent relationships between flooded flat, permanent water, swamp and marsh wetlands. Then, as for three coastal tidal wetlands, the mangrove forests were only found in coastal areas below 30° N and




Figure 8. The importance of multi-sourced and multi-temporal features derived from the random forest classification model.

were mainly concentrated in regions between 30° N–30° S, including Southeast Asia, West Africa and the east coast of South America. The salt marshes and tidal flats shared similar spatial distributions. They were widely distributed globally and can be observed along most coastlines. In addition, the tidal flat distributions were closely related to the slope of coastlines, tidal ranges and sediment inflows. For example, the tidal flats in Asia and Europe usually were located in the tide-dominated estuaries and deltas. Similarly, Murray et al. (2019) also demonstrated that there were often more tidal flats where the river flowed into the sea.

To quantitatively summarize the distribution of the eight wetland sub-categories, the total area and area percentages of eight fine wetland sub-categories over each continent are calculated in Fig. 11 and Table 4. The total wetland area was 6.38 million km², including 6.03 million km² of inland wetlands and 0.35 million km² of coastal tidal wetlands, and the distribution of wetlands varied across different continents. Intuitively, approximately 60 % of coastal tidal wetlands (tidal flat, salt marsh and mangrove) and 70 % of permanent water, flooded flat and marsh wetlands were distributed in the Northern Hemisphere, especially on the Asian and North American continents. Comparatively, more than 85 % of saline wetlands were located in the Southern Hemisphere, especially the Oceanian continent. Then, in terms of specific wetland sub-categories, most permanent water was concentrated in the Northern Hemisphere and especially in North America (nearly 50 % of the world's permanent waterbodies). The swamp was mainly distributed on the North American, African and South American continents, which contained many rainforest wetlands, with corresponding swamp areas of 0.39, 0.18 and 0.32 million km², respectively. Swamp areas on the Oceanian continent were the smallest, covering only 6572 km² mainly because the forest cover in Oceania was smaller than on other continents. The marsh and flooded flats shared similar areal proportions

on all six continents and were mainly concentrated in the Northern Hemisphere (exceeding 70 %), where many lakes and rivers were distributed. Next, as the mangrove forests only covered regions south of 30° N and were mostly concentrated in tropical regions near the Equator, such as Southeast Asia, East Africa and Central America, this sub-category was absent on the Europe continent and sparse in Oceania.

5.4 Accuracy assessment of global 30 m fine wetland map

Using 25 709 global validation samples, the confusion matrix of the novel GMW_FCS30 wetland map was calculated in Table 5. Overall, our wetland map achieved an overall accuracy of 86.44 % and a kappa coefficient of 0.82 across the fine wetland classification system. In terms of the producer and user accuracies, the non-wetlands achieved the highest producer accuracy of 94.24 % mainly because we combined pre-existing multi-sourced wetland datasets to determine the maximum wetland boundary and further used multi-sourced and time-series imagery to distinguish between wetlands and non-wetlands. The permanent water achieved the highest user accuracy of 95.99 % because the permanent water had unique and stable spectral characteristics, and the training samples were directly from the JRC_GSW database (Pekel et al., 2016). Then, as for the coastal tidal wetlands, mangrove forest and tidal flat achieved higher accuracies than salt marsh, with producer accuracies of 91.43 % and 88.12 % and user accuracies of 95.69 % and 94.81 %, respectively. The salt marsh had a lower producer accuracy of 74.09 % because its reflectance spectra were affected by both water levels and vegetation cover with considerable spatiotemporal heterogeneity, and the sparser prior salt marsh products were adopted. Next, as for inland sub-categories, the swamp and marsh obviously performed better than the flooded flat, with producer accuracies of 72.03 % and 78.09 %, respectively. It can be seen that the confusion between swamp and




Figure 9. The overview of global 30 m fine wetland maps and their area statistics in latitudinal and longitudinal directions in 2020. Four local enlargements in (a) Florida, (b) Congo Basin, (c) Sundarbans and (d) Poyang Lake are also illustrated.

Table 4. The total wetland area (unit: $10^4 \times \text{km}^2$) of eight wetland sub-categories for six continents and the globe.

	Permanent water	Swamp	Marsh	Flooded flat	Saline	Mangrove	Salt marsh	Tidal flat
Asia	90.529	13.227	58.229	7.244	1.215	6.636	1.852	5.347
North America	123.754	39.314	45.350	11.867	0.008	2.590	2.619	2.697
Europe	27.111	7.010	22.513	3.601	0.005	0.000	0.717	1.408
Africa	24.214	18.393	14.926	1.318	1.248	3.105	0.688	0.731
South America	18.310	32.337	21.640	5.242	1.888	2.175	0.520	1.238
Oceania	1.330	0.657	6.151	0.233	4.355	1.219	1.094	0.875
Total	285.247	110.938	168.810	29.504	8.719	15.725	7.491	12.296

marsh was the main source of the misclassification error of swamp and that the marsh was simultaneously confused with non-wetland, swamp and flooded flat because the spectra of marsh changed along with the water levels. For example, the marsh in Poyang Lake, shown in Fig. 4b, was flooded at its highest water levels. Then, the flooded flat achieved a low producer accuracy of 65.83 % because it usually coexisted

with the marsh and shared similar spectral characteristics, so approximately 10.89 % of flooded flat points were labeled as marsh in our wetland map. The saline wetland was mainly concentrated along the edge of salt lakes and demonstrated great performance in our mapping, with producer and user accuracies of 91.96 % and 91.66 %, respectively.




Figure 10. The spatial distributions of the eight wetland sub-categories after aggregating them to a resolution of $0.5^\circ \times 0.5^\circ$.




Figure 11. The area proportions of eight wetland sub-categories over each continent.

6 Discussion

6.1 Cross-comparisons with other global wetland maps

To comprehensively understand the performance of the GWL_FCS30 wetland maps, four existing global wetland datasets (GLC_FCS30, GlobeLand30, CCI_LC and GLWD), listed in Table 1, were selected. Figure 12 quantitatively illustrates the total wetland area of five products over each continent. Specifically, the total wetland area of different wetland products varied. The GLWD obviously overestimated the wetland area on each continent mainly because it was derived from the compilation model instead of actual remote sensing observations (Lehner and Döll, 2004). Namely, the GLWD classified a large amount of non-

wetlands as potential wetlands. The remaining four wetland products, derived from the Landsat and PROBE-V remote sensing imagery, shared a total wetland area of 4.128–7.364 million km², and our GWL_FCS30 wetland dataset had a total area of 6.387 million km² among these datasets. The CCI LC wetland layer contained the smallest wetland area of 4.128 million km², and the estimated area in North America was profoundly lower than the other datasets mainly because the CCI LC heavily underestimated the wetland distribution in Canada after a comparison with the Canadian Wetland Inventory (Amani et al., 2019). Next, the total wetland area in the GlobeLand30 and GLC_FCS30 wetland layer was higher than the developed GWL_FCS30 wetland dataset because some water-level-sensitive non-

Table 5. The confusion matrix of the global 30 m fine wetland map using 25 709 validation points.

	NWT	PW	SWP	MSH	FFT	SAL	MGV	SMH	TFT	Total	P.A.
NWT	9950	17	254	224	39	3	12	33	26	10 588	94.24
PW	69	2251	4	15	63	0	0	8	9	2419	93.06
SWP	272	5	2127	452	74	11	3	9	0	2953	72.03
MSH	546	18	135	3218	149	18	2	34	1	4121	78.09
FFT	145	21	26	95	574	3	1	5	2	872	65.83
SAL	26	1	0	43	5	846	0	0	0	921	91.86
MGV	65	4	11	2	2	1	1109	15	3	1213	91.43
SMH	157	15	6	85	9	30	26	998	22	1347	74.09
TFT	78	13	0	11	7	11	6	29	1150	1305	88.12
Total	11 308	2345	2563	4145	922	923	1159	1131	1213	25 709	
U.A.	87.99	95.99	82.99	79.56	62.26	91.66	95.69	88.24	94.81		
O.A.						86.44					
Kappa						0.822					

Note: NWT: non-wetlands; PW: permanent water; SWP: swamp; MSH: marsh; FFT: flooded flat; SAL: saline; SMH: salt marsh; MGV: mangrove forest; TFT: tidal flat; O.A.: overall accuracy; P.A.: producer accuracy; U.A.: user accuracy.

wetlands (such as irrigated cropland) were also captured in these two datasets.

Figure 13 illustrates the performances of five wetland products for two typical wetland regions (Poyang Lake in China and Pantanal wetland in Brazil). The reasons for choosing these two regions were that the wetlands in Poyang Lake quickly changed with water levels, and the Pantanal wetland was the largest wetland in the world. Intuitively, the GWL_FCS30 wetland maps had the greatest performance in capturing the spatial patterns of various wetland sub-categories. Comparatively, the GLC_FCS30 wetland layer suffered serious underestimation and misclassification problems in these two regions, which obviously misclassified many water-sensitive wetlands (swamp and marsh) as waterbodies in Poyang Lake and also missed a large number of marsh and swamp wetlands in the Pantanal wetland. Zhang et al. (2021b) also stated that the wetland in GLC_FCS30 suffered from low accuracy because of a lack of enough wetland samples and multi-sourced wetland-sensitive features. Then, the GlobeLand30 wetland layer performed better in the Pantanal wetland than in Poyang Lake, which also obviously misclassified many marsh wetlands as waterbodies in the Poyang Lake mainly because the low-water-level features were not captured during the development of GlobeLand30 (Chen et al., 2015). In addition, the wetland layer of GlobeLand30 in Pantanal still suffered from the overestimation problem, and some non-wetlands in Pantanal Wetland Park were mislabeled as wetland, so the wetland layer in GlobeLand30 only achieved a user accuracy of 74.87 % (Chen et al., 2015). The CCI LC was highly consistent with the GWL_FCS30 wetland maps in spatial distribution when comparing with GLC_FCS30 and Globeland30; however, details show that the wetlands in the CCI LC were still underestimated in the Poyang Lake wetland and overestimated

in the Pantanal wetland based on the highest- and lowest-water-level composites. Lastly, the GLWD dataset significantly overestimated the wetlands in two regions; namely, the mapped marsh area was obviously greater than its actual area, and it also misclassified these water-sensitive wetlands as waterbodies near Poyang Lake.

Figure 14 illustrates the comparisons between the GWL_FCS30 map with three widely used global mangrove forest products (World Atlas of Mangroves, GMW_V3 (Global Mangrove Watch Version3) and USGS Global Distribution of Mangroves) listed in Table 1 in two typical mangrove regions (coastal Indonesia and Sundarbans). Overall, there was great consistency over four mangrove datasets because the mangrove forest reflected obvious and strong vegetation reflectance characteristics and was easier to identify than other wetland sub-categories. In detail, the Atlas mangrove dataset suffers from the underestimation problem; namely, the mangrove area in the Atlas mangrove dataset was obviously lower than the other three products, especially in coastal Indonesia (local enlargements). The USGS mangrove product can comprehensively and accurately capture the spatial distribution of mangroves over two regions. Still, it missed small and isolated fragments of mangrove forests in two regions (green rectangle) based on high-resolution imagery. The GMW_V3 dataset was validated to achieve an overall accuracy of 95.25 %, with user and producer accuracies of mangrove forests of 97.5 % and 94.0 %, respectively (Bunting et al., 2018; Thomas et al., 2017), which shows the greatest agreement with our GWL_FCS30 maps in these two regions and enlargements. Using the high-resolution imagery, it can be found that GWL_FCS30 and GWM_V3 accurately identified the spatial patterns of mangrove forest in both regions.




Figure 12. The total wetland area (unit: million km²) of five global wetland products on six continents.

Figure 15 illustrated the comparisons between the GWL_FCS30 tidal flat layer with Murray’s tidal flat V1.1 in 2016 (Murray et al., 2019) and the updated Murray’s tidal flat V1.2 in 2019 (Murray et al., 2022) in two local regions, and the corresponding highest- and lowest-tidal-level composites are also listed. Overall, three products can comprehensively capture the spatial patterns of tidal flats in these two regions, and GWL_FCS30-2020 and Murray’s tidal flat V1.2 performed with a higher spatial consistency, while Murray’s tidal flat V1.1 suffered the obvious omission error in three typical areas (red rectangles). In detail, we can find that Murray’s tidal flat products misclassified some coastal ponds and lakes as tidal flats especially in the first region, while GWL_FCS30-2020 achieved the best performance and accurately excluded these coastal ponds and lakes. In addition, GWL_FCS30 also distinguished between the salt marshes and tidal flats especially in the Yellow River estuary, while Murray’s tidal flat V1.2 database misclassified a lot of salt marshes as tidal flats.

6.2 Comparisons with the national wetland products

Using 1835 validation points (from the global validation points in Sect. 4.3) over the contiguous United States, we quantitatively assessed the accuracy metrics of NLCD (National Land Cover Database) with GWL_FCS30 after merging the wetland sub-categories into four classes in Table 6. Overall, GWL_FCS30 achieved a higher performance than that of the NLCD mainly because a lot of herbaceous wetlands were misclassified as open water in the NLCD, so the user accuracy of herbaceous wetland and producer accuracy of open water in NLCD were lower than those of GWL_FCS30. Then, as the NWI had a different wetland system with the NLCD and GWL_FCS30, we also analyzed the

metrics of the NWI with GWL_FCS30 after merging into five classes. It can be found that the NWI shared similar performances with GWL_FCS30 on the non-wetlands and marine wetlands, but the user accuracies of forest wetland and herbaceous wetland of the NWI were lower than those of GWL_FCS30 mainly because some non-wetlands and open water were overestimated as wetland in the NWI. Similarly, Gage et al. (2020) also demonstrated that the NWI more easily overestimated the wetland areas.

Figure 16 illustrates the comparisons between our GWL_FCS30-2020, the NLCD wetland layer and the NWI in San Francisco and Florida. It should be noted that the ocean was excluded in GWL_FCS30-2020, while the NLCD and the NWI still retained it. Overall, three wetland products performed with a great spatial consistency and accurately captured the spatial patterns of wetlands over two regions. From the perspective of the diversity of the wetland sub-category, GWL_FCS30 and the NWI had obvious advantages over the NLCD, which simply divided the wetlands into open water, woody wetlands and emergent herbaceous wetlands. Afterwards, the NWI had the largest wetland areas in San Francisco because it included the irrigated cropland (red color), while the other two datasets excluded irrigated cropland. Then, the local enlargement showed that GWL_FCS30 and the NWI also had better performance than the NLCD because they comprehensively captured the coastal tidal wetlands, and our GWL_FCS30 further distinguished between the tidal flats and salt marshes, which also demonstrated that GWL_FCS30 performed better than the NWI over the coastal tidal wetlands. In Florida, the NWI and GWL_FCS30 accurately divided the inland and coastal tidal wetlands, and GWL_FCS30 further identified the coastal tidal wetlands as mangrove forest. Meanwhile, the local enlargement also demonstrated the great consistency of three wetland prod-




Figure 13. The cross-comparisons between our GWL_FCS30 wetland maps with four existing wetland products: GLC_FCS30 generated by Zhang et al. (2021b), GlobeLand30 generated by Chen et al. (2015), CCI LC generated by Defourny et al. (2018) and GLWD generated by Lehner and Döll (2004) at Pantanal and Poyang Lake wetland. The false-color composed Landsat imagery (SWIR1, NIR and red bands) at the highest and lowest water levels is also illustrated.

ucts. However, it can be found that there was an obvious difference between GWL_FCS30 and the NWI over the wetland categories, in which GWL_FCS30 classified most inland wetlands as marshes, while the NWI classified them as emergent wetlands and forest/shrub wetlands mainly because of the differences in the definition of the classification system (GWL_FCS30 defined those low shrubs that grow in freshwater as marsh; Table 1).

Table 7 illustrated the accuracy metrics of CLC (CORINE Land Cover) and GWL_FCS30 after merging the wetland categories over the European Union area using 1996 validation points from the global validation points in Sect. 4.3. Overall, GWL_FCS30 performed better than the CLC, and the former mainly had fewer commission errors than that of

the CLC for salt marsh and tidal flat. To intuitively understand the overestimation of tidal flat, Fig. 17 illustrates the comparison between our GWL_FCS30-2020 and the CLC wetland layer in 2018 over the Nordic Sea, in which mainly distributed in tidal flats and open water, and these tidal flats gathered around the coastline. In terms of the specific wetland sub-category, it can be found that the CLC database had a larger tidal flat area than that of GWL_FCS30; however, the lowest-tidal-level composite from Landsat time-series imagery indicated that the CLC overestimated the tidal flats in the region. For example, the local enlargement showed that a lot of permanent ocean pixels were wrongly labeled as tidal flats in CLC and accurately identified as ocean in GWL_FCS30. The comparison also demonstrated why the




Figure 14. The cross-comparisons between our GWL_FCS30 wetland maps with three mangrove products (Atlas mangrove developed by Spalding, 2010; GMW_V3 developed by Bunting et al., 2022; Mangrove USGS developed by Giri et al., 2011) in Sundarbans and coastal Indonesia. The high-resolution imagery comes from the © Google Earth Engine platform (<https://earthengine.google.com>; last access: 16 May 2022).

CLC had a low user accuracy of 62.90 % for tidal flat and producer accuracy of 57.76 % for waterbodies. Then, the local enlargement also indicated that the total area of salt marsh in CLC was lower than that of GWL_FCS30 (green rectangles); namely, some salt marshes were wrongly labeled as tidal flat and waterbody, so the accuracy metrics in Table 7 showed the user accuracy of salt marsh in CLC was 35.86 %.

6.3 The limitations and prospects of our global fine wetland map

It should be noted there were still many uncertainties and limitations to the proposed method and global wetland maps. First, the proposed method used continuous Landsat reflectance and Sentinel-1 SAR imagery to capture various water-level information. Still, it might fail when the avail-




Figure 15. The comparisons between the tidal flat of GWL_FCS30 in 2020, Murray’s tidal flat V1.1 in 2016 (Murray et al., 2019) and Murray’s tidal flat V1.2 in 2019 (Murray et al., 2022) for two local regions. In each case, the highest- and lowest-tidal-level composites, composited by SWIR1, NIR and red bands, are illustrated.

Table 6. The accuracy metrics of NLCD, NWI and GWL_FCS30 using 1835 validation points over the contiguous United States.

(a) NLCD vs. GWL_FCS30											
		NWT	Open water			Woody wetland		Emergent herbaceous wetland		O.A.	Kappa
NLCD	U.A.	96.46	93.98			77.92		61.97		83.58	0.756
	P.A.	88.80	53.65			85.96		87.61			
		NWT	PW	FFT	TFT	SWP	MGV	MSH	SMH	O.A.	Kappa
GWL_FCS30	U.A.	90.55	94.81			69.87			87.61	85.76	0.786
	P.A.	85.99	95.52			77.97			88.36		

(b) NWI vs. GWL_FCS30													
		NWT	FPD	EMD	RVR	LKE	FSSW	FEW	EMW		O.A.	Kappa	
NWI	U.A.	94.45					67.58	60.25	85.71		83.49	0.762	
	P.A.	84.93					86.62	82.76	91.53				
		NWT	PW		SWP		MSH	TFT	MGV	SMH	TFT	O.A.	Kappa
GWL_FCS30	U.A.	90.55	94.74		68.96		80.75		90.08		85.23	0.789	
	P.A.	85.99	95.45		76.76		78.78		94.98				

Note: NWT: non-wetlands; PW: permanent water; SWP: swamp; MSH: marsh; FFT: flooded flat; SMH: salt marsh; MGV: mangrove forest; TFT: tidal flat; FPD: freshwater pond; EMD: estuarine and marine deepwater; RVR: riverine; LKE: lake; FSSW: freshwater forested/shrub wetland; FEW: freshwater emergent wetland; EMW: estuarine and marine wetland; O.A.: overall accuracy; P.A.: producer accuracy; U.A.: user accuracy.

able Landsat observations were sparse and lacked the aid of Sentinel-1 SAR data, especially before 2000. Thus, our future work would focus on combining a richer multi-sourced data source, including MODIS, Sentinel-2, SPOT and PAL-SAR imagery, to develop a more robust wetland mapping method. For example, Chen et al. (2018) integrated Landsat and MODIS observations to successfully monitor the wetland dynamics from 2000 to 2014 using a spatiotemporal adaptive fusion model. Then, in this study, we combined the multi-sourced wetland products and their practical use for ecosystem management to define a fine wetland clas-

sification system containing eight sub-categories; however, there are still many wetland sub-categories, such as submergent vegetation (*Nymphaea*), groundwater-dependent wetlands (karst and cave systems) and seagrass beds (Richardson et al., 2022), that cannot be captured because remote sensing observations usually had poor performance on penetrating waterbodies and then capturing underwater characteristics, and there was currently no prior dataset for global underwater wetlands. Meanwhile, some coastal swamps (except for mangrove), which were usually overlooked in most coastal wetland mapping (Murray et al., 2022; Z. Zhang et al., 2022),




Figure 16. The comparisons between GWL_FCS30 in 2020, National Land Cover Database (NLCD) wetland layer (Homer et al., 2020) and National Wetlands Inventory (NWI, <https://www.fws.gov/program/national-wetlands-inventory>, last access: 12 November 2022) in San Francisco and Florida. The high-resolution imagery comes from the © Google Earth Engine platform (<https://earthengine.google.com/>; last access: 12 November 2022).

Table 7. The accuracy metrics between CLC and GWL_FCS30 after merging the wetland categories.

CLC		GWL_FCS30						CLC					
		NWT	WC	WB	CL	ET	SO	Peat bogs and inland marshes	SMH	TFT	O.A.	Kappa	
U.A.	U.A.	92.94			94.81			68.63	35.86	62.90	80.75	0.706	
	P.A.	82.80			57.76			83.93	91.23	75.00			
GWL_FCS30		GWL_FCS30						CLC					
		NWT	WC	WB	CL	ET	SO	SWP	MSH	FFT	SMH	TFT	O.A.
	U.A.	91.22			88.02			80.98	86.21	94.35	88.10	0.816	
	P.A.	88.54			97.69			80.82	91.91	97.50			

Note: NWT: non-wetlands; WC: water courses; WB: waterbodies; CL: coastal lagoons; ET: estuaries; SO: sea and ocean; PW: permanent water; SWP: swamp; MSH: marsh; FFT: flooded flat; SAL: saline; SMH: salt marsh; MGCV: mangrove forest; TFT: tidal flat; O.A.: overall accuracy; P.A.: producer accuracy; U.A.: user accuracy.

were also missed in GWL_FCS30 mainly because there are no global or large-area coastal swamp datasets that can be imported, and the coastal swamp is also sparser than the mangrove forest in the low and middle latitudes. So, our further work will pay more attention to combine multi-sourced auxiliary datasets, such as hydrological, bathymetric and climate data, to map these special wetland sub-categories in a targeted manner.

We combined the pre-existing global wetland products to derive the training samples and maximum extents; however, the salt marsh and saline samples still used the visual interpretation method to ensure their reliability because of a lack of sufficient pre-existing global products. Additionally, it was found that the producer accuracy of salt marsh and saline in Table 4 was relatively poor compared with other sub-categories mainly because visual interpretation cannot




Figure 17. The comparisons between GWL_FCS30 and CORINE Land Cover (CLC) wetland layer in 2018 (<https://land.copernicus.eu/pan-european/corine-land-cover/clc2018?tab=metadata>, last access: 22 November 2022). The lowest-tidal-level Landsat composite, composed by NIR, red and green bands, is illustrated.

provide massive and geographically distributed salt marsh and saline training samples. Namely, this study cannot comprehensively capture the regional adaptive reflectance characteristics of salt marsh and saline. Fortunately, many studies have built expert knowledge of these sub-categories over recent years. For example Mao et al. (2020) combined multi-scale segmentation, multiple normalized indices and rule-based classification methods to develop a wetland map of China with an overall classification accuracy of 95.1 %. Similarly, Wang et al. (2020) used the four widely used spectral indices to successfully identify three sub-categories within coastal tidal wetlands. Thus, our further work should focus more effort on the spectral characteristics of salt marsh and saline wetlands and build expert knowledge of them for automatically deriving their training samples.

In addition, we used the derived maximum extents as the boundary for identifying inland and coastal tidal wetlands; in other words, we assumed that the derived maximum extents contained all inland and coastal tidal wetlands with zero omission error. Actually, the inland maximum extents in Eq. (3) fulfilled the assumption of zero omission error because the GLWD and TROP-SUBTROP products, produced by the compilation and model simulation method (Gumbrecht, 2015; Lehner and Döll, 2004), can capture most wetland areas at the expense of a higher commission error. For example, Fig. 13 illustrates the cross-comparisons between our GWL_FCS30 wetland maps and four existing wetland products, and the GLWD obviously overestimated the inland wetlands. On the other hand, the union of five global wetland datasets in Eq. (3) also minimized the omission error of each dataset for inland wetland sub-categories. Next, as for

the maximum mangrove forest extents (Eq. 1), as the high producer and user accuracies were achieved by five prior mangrove products (explained in Sect. 2.2) and the mangrove time-series products were integrated in order that these missed mangroves may be complemented by other products or time-series products, the derived maximum extents can also be considered as zero omission error and covered almost all mangrove forests. Recently, Bunting et al. (2022) developed the newest mangrove products covering 1996–2020, it can be used as another important prior dataset in our further works for deriving the maximum mangrove extents. Lastly, the maximum tidal flat extents, derived from Murray’s time-series products from 1985–2016 by using the union operation (Eq. 2), can also contain almost all tidal flats because previous studies demonstrated that they suffered more commission errors than omission errors (Jia et al., 2021; Z. Zhang et al., 2022). The missed tidal flats would concentrate on these newly increased tidal flats during 2016–2020; fortunately, new global tidal flat time-series products during 1999–2019 have been developed (Murray et al., 2022) and can be used as an important supplement in our further work for deriving the maximum tidal flat extent with zero omission error.

7 Data availability

The GWL_FCS30 wetland dataset in 2020 is freely available at <https://doi.org/10.5281/zenodo.7340516> (Liu et al., 2022). It is composed of 961 files of $5^\circ \times 5^\circ$ geographical grid tiles, and each tiled file is stored using the geographical projection system with a spatial resolution of 30 m in the GeoTIFF format. The fine wetland sub-category information is labeled as

0, 180, 181, 182, 183, 184, 185 186 and 187, representing the non-wetland, permanent water, swamp, marsh, flooded flat, saline, mangrove forest, salt marsh and tidal flat, respectively. The validation samples are available upon request.

8 Conclusions

Over the past few decades, many global and regional wetland products have been developed; however, an accurate global 30 m wetland dataset, with fine wetland categories and coverage of both inland and coastal zones, is still lacking. In this study, the Landsat reflectance and Sentinel-1 SAR time-series imagery, together with the stratified classification strategy and local adaptive random forest classification algorithm, was successfully integrated to produce the first global 30 m wetland product with a fine classification system in 2020. The wetlands were classified as four inland wetlands (swamp, marsh, flooded flat and saline) and three coastal tidal wetlands (mangrove, salt marsh and tidal flat). The produced wetland dataset, *GWL_FCS30*, accurately captured the spatial patterns of seven wetland sub-categories with an overall accuracy of 86.44 % and a kappa coefficient of 0.822 for the fine wetland classification system with fewer omission and commission errors compared to other global products. The quantitative statistical analysis showed that the global wetland area reached 6.38 million km², including 6.03 million km² of inland wetlands and 0.35 million km² of coastal tidal wetlands. Approximately 72.96 % of wetlands were distributed poleward of 40° N. Therefore, the proposed method is suitable for large-area fine wetland mapping, and the *GWL_FCS30* dataset can serve as an accurate wetland map that could potentially provide vital support for wetland management.

Author contributions. LL conceptualized the project. XZ performed the investigation. LL and XZ designed the methodology. XZ and XC developed the software. XZ, TZ, SL, WL, JM and JW performed the validation. XZ prepared the original draft of the paper. LL reviewed and edited the paper.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. We gratefully acknowledge all data providers whose data have been used in this study and would like to thank the topical editor and the three anonymous referees for their constructive comments.

Financial support. This research has been supported by the Innovative Research Program of the International Research Center of Big Data for Sustainable Development Goals (grant no. CBAS2022ORP03), the National Natural Science Foundation of China (grant no. 41825002) and National Earth System Science Data Sharing Infrastructure (grant no. 2005DKA32300).

Review statement. This paper was edited by Yuyu Zhou and reviewed by three anonymous referees.

References

- Amani, M., Mahdavi, S., Afshar, M., Brisco, B., Huang, W., Mohammad Javad Mirzadeh, S., White, L., Banks, S., Montgomery, J., and Hopkinson, C.: Canadian Wetland Inventory using Google Earth Engine: The First Map and Preliminary Results, *Remote Sens.-Basel*, 11, 842, <https://doi.org/10.3390/rs11070842>, 2019.
- Azzari, G. and Lobell, D. B.: Landsat-based classification in the cloud: An opportunity for a paradigm shift in land cover monitoring, *Remote Sens. Environ.*, 202, 64–74, <https://doi.org/10.1016/j.rse.2017.05.025>, 2017.
- Büttner, G.: CORINE land cover and land cover change products, in: Land use and land cover mapping in Europe, Springer, https://doi.org/10.1007/978-94-007-7969-3_5, 2014.
- Belgiu, M. and Drăguț, L.: Random forest in remote sensing: A review of applications and future directions, *ISPRS J. Photogramm.*, 114, 24–31, <https://doi.org/10.1016/j.isprsjprs.2016.01.011>, 2016.
- Breiman, L.: Random Forests, *Mach. Learn.*, 45, 5–32, <https://doi.org/10.1023/a:1010933404324>, 2001.
- Brown, C. F., Brumby, S. P., Guzder-Williams, B., Birch, T., Hyde, S. B., Mazzariello, J., Czerwinski, W., Pasquarella, V. J., Haertel, R., Ilyushchenko, S., Schwehr, K., Weisse, M., Stolle, F., Hanson, C., Guinan, O., Moore, R., and Tait, A. M.: Dynamic World, Near real-time global 10 m land use land cover mapping, *Scientific Data*, 9, 1–7, <https://doi.org/10.1038/s41597-022-01307-4>, 2022.
- Bunting, P., Rosenqvist, A., Lucas, R., Rebelo, L.-M., Hilarides, L., Thomas, N., Hardy, A., Itoh, T., Shimada, M., and Finlayson, C.: The Global Mangrove Watch—A New 2010 Global Baseline of Mangrove Extent, *Remote Sens.-Basel*, 10, 1669, <https://doi.org/10.3390/rs10101669>, 2018.
- Bunting, P., Rosenqvist, A., Hilarides, L., Lucas, R. M., Thomas, N., Tadono, T., Worthington, T. A., Spalding, M., Murray, N. J., and Rebelo, L.-M.: Global Mangrove Extent Change 1996–2020: Global Mangrove Watch Version 3.0, *Remote Sens.-Basel*, 14, 3657, <https://doi.org/10.3390/rs14153657>, 2022.
- Bwangoy, J.-R. B., Hansen, M. C., Roy, D. P., Grandi, G. D., and Justice, C. O.: Wetland mapping in the Congo Basin using optical and radar remotely sensed data and derived topographical indices, *Remote Sens. Environ.*, 114, 73–86, <https://doi.org/10.1016/j.rse.2009.08.004>, 2010.
- Cao, W., Zhou, Y., Li, R., and Li, X.: Mapping changes in coastlines and tidal flats in developing islands using the full time series of Landsat images, *Remote Sens. Environ.*, 239, 111665, <https://doi.org/10.1016/j.rse.2020.111665>, 2020.

- Chen, B., Chen, L., Huang, B., Michishita, R., and Xu, B.: Dynamic monitoring of the Poyang Lake wetland by integrating Landsat and MODIS observations, *ISPRS J. Photogramm.*, 139, 75–87, <https://doi.org/10.1016/j.isprsjprs.2018.02.021>, 2018.
- Chen, G., Jin, R., Ye, Z., Li, Q., Gu, J., Luo, M., Luo, Y., Christakos, G., Morris, J., He, J., Li, D., Wang, H., Song, L., Wang, Q., and Wu, J.: Spatiotemporal Mapping of Salt Marshes in the Intertidal Zone of China during 1985–2019, *Journal of Remote Sensing*, 2022, 1–15, <https://doi.org/10.34133/2022/9793626>, 2022.
- Chen, J., Chen, J., Liao, A., Cao, X., Chen, L., Chen, X., He, C., Han, G., Peng, S., Lu, M., Zhang, W., Tong, X., and Mills, J.: Global land cover mapping at 30m resolution: A POK-based operational approach, *ISPRS J. Photogramm.*, 103, 7–27, <https://doi.org/10.1016/j.isprsjprs.2014.09.002>, 2015.
- Davidson, N. C.: How much wetland has the world lost? Long-term and recent trends in global wetland area, *Mar. Freshwater Res.*, 65, 934–941, <https://doi.org/10.1071/mf14173>, 2014.
- Defourny, P., Kirches, G., Brockmann, C., Boettcher, M., Peters, M., Bontemps, S., Lamarche, C., Schlerf, M., and Santoro, M.: Land Cover CCI: Product User Guide Version 2, https://www.esa-landcover-cci.org/?q=webfm_send/84 (last access: 22 November 2022), 2018.
- DeVries, B., Huang, C., Armston, J., Huang, W., Jones, J. W., and Lang, M. W.: Rapid and robust monitoring of flood events using Sentinel-1 and Landsat data on the Google Earth Engine, *Remote Sens. Environ.*, 240, 111664, <https://doi.org/10.1016/j.rse.2020.111664>, 2020.
- Dixon, M. J. R., Loh, J., Davidson, N. C., Beltrame, C., Freeman, R., and Walpole, M.: Tracking global change in ecosystem area: The Wetland Extent Trends index, *Biol. Conserv.*, 193, 27–35, <https://doi.org/10.1016/j.biocon.2015.10.023>, 2016.
- Gómez, C., White, J. C., and Wulder, M. A.: Optical remotely sensed time series data for land cover classification: A review, *ISPRS J. Photogramm.*, 116, 55–72, <https://doi.org/10.1016/j.isprsjprs.2016.03.008>, 2016.
- Gage, E., Cooper, D. J., and Lichvar, R.: Comparison of USACE three-factor wetland delineations to national wetland inventory maps, *Wetlands*, 40, 1097–1105, <https://doi.org/10.1007/s13157-019-01234-y>, 2020.
- Gardner, R. C. and Davidson, N. C.: The Ramsar convention, in: *Wetlands*, Springer, https://doi.org/10.1007/978-94-007-0551-7_11, 2011.
- Giri, C., Ochieng, E., Tieszen, L. L., Zhu, Z., Singh, A., Loveland, T., Masek, J., and Duke, N.: Status and distribution of mangrove forests of the world using earth observation satellite data, *Global Ecol. Biogeogr.*, 20, 154–159, <https://doi.org/10.1111/j.1466-8238.2010.00584.x>, 2011.
- Gislason, P. O., Benediktsson, J. A., and Sveinsson, J. R.: Random Forests for land cover classification, *Pattern Recogn. Lett.*, 27, 294–300, <https://doi.org/10.1016/j.patrec.2005.08.011>, 2006.
- Gong, P., Wang, J., Yu, L., Zhao, Y., Zhao, Y., Liang, L., Niu, Z., Huang, X., Fu, H., Liu, S., Li, C., Li, X., Fu, W., Liu, C., Xu, Y., Wang, X., Cheng, Q., Hu, L., Yao, W., Zhang, H., Zhu, P., Zhao, Z., Zhang, H., Zheng, Y., Ji, L., Zhang, Y., Chen, H., Yan, A., Guo, J., Yu, L., Wang, L., Liu, X., Shi, T., Zhu, M., Chen, Y., Yang, G., Tang, P., Xu, B., Giri, C., Clinton, N., Zhu, Z., Chen, J., and Chen, J.: Finer resolution observation and monitoring of global land cover: first mapping results with Landsat TM and ETM+ data, *Int. J. Remote Sens.*, 34, 2607–2654, <https://doi.org/10.1080/01431161.2012.748992>, 2013.
- Gong, P., Liu, H., Zhang, M., Li, C., Wang, J., Huang, H., Clinton, N., Ji, L., Li, W., Bai, Y., Chen, B., Xu, B., Zhu, Z., Yuan, C., Ping Suen, H., Guo, J., Xu, N., Li, W., Zhao, Y., Yang, J., Yu, C., Wang, X., Fu, H., Yu, L., Dronova, I., Hui, F., Cheng, X., Shi, X., Xiao, F., Liu, Q., and Song, L.: Stable classification with limited sample: transferring a 30 m resolution sample set collected in 2015 to mapping 10 m resolution global land cover in 2017, *Sci. Bull.*, 64, 370–373, <https://doi.org/10.1016/j.scib.2019.03.002>, 2019.
- Gumbricht, T.: Hybrid mapping of pantropical wetlands from optical satellite images, hydrology, and geomorphology, *Remote Sensing of Wetlands*, CRC Press, 435–454, <https://doi.org/10.1201/b18210>, 2015.
- Gumbricht, T., Roman-Cuesta, R. M., Verchot, L., Herold, M., Wittmann, F., Householder, E., Herold, N., and Muriyarsa, D.: An expert system model for mapping tropical wetlands and peatlands reveals South America as the largest contributor, *Glob. Change Biol.*, 23, 3581–3599, <https://doi.org/10.1111/gcb.13689>, 2017.
- Guo, M., Li, J., Sheng, C., Xu, J., and Wu, L.: A Review of Wetland Remote Sensing, *Sensors*, 17, 777, <https://doi.org/10.3390/s17040777>, 2017.
- Hamilton, S. E. and Casey, D.: Creation of a high spatio-temporal resolution global database of continuous mangrove forest cover for the 21st century (CGMFC-21), *Global Ecol. Biogeogr.*, 25, 729–738, <https://doi.org/10.1111/geb.12449>, 2016.
- Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thau, D., Stehman, S. V., Goetz, S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O., and Townshend, J. R.: High-resolution global maps of 21st-century forest cover change, *Science*, 342, 850–853, <https://doi.org/10.1126/science.1244693>, 2013.
- Hansen, M. C., Egorov, A., Potapov, P. V., Stehman, S. V., Tyukavina, A., Turubanova, S. A., Roy, D. P., Goetz, S. J., Loveland, T. R., Ju, J., Kommareddy, A., Kovalsky, V., Forsyth, C., and Bents, T.: Monitoring conterminous United States (CONUS) land cover change with Web-Enabled Landsat Data (WELD), *Remote Sens. Environ.*, 140, 466–484, <https://doi.org/10.1016/j.rse.2013.08.014>, 2014.
- Homer, C., Dewitz, J., Jin, S., Xian, G., Costello, C., Danielson, P., Gass, L., Funk, M., Wickham, J., Stehman, S., Auch, R., and Riitters, K.: Conterminous United States land cover change patterns 2001–2016 from the 2016 National Land Cover Database, *ISPRS J. Photogramm.*, 162, 184–199, <https://doi.org/10.1016/j.isprsjprs.2020.02.019>, 2020.
- Hu, S., Niu, Z., and Chen, Y.: Global Wetland Datasets: a Review, *Wetlands*, 37, 807–817, <https://doi.org/10.1007/s13157-017-0927-z>, 2017a.
- Hu, S., Niu, Z., Chen, Y., Li, L., and Zhang, H.: Global wetlands: Potential distribution, wetland loss, and status, *Sci. Total Environ.*, 586, 319–327, <https://doi.org/10.1016/j.scitotenv.2017.02.001>, 2017b.
- Huang, X., Li, J., Yang, J., Zhang, Z., Li, D., and Liu, X.: 30 m global impervious surface area dynamics and urban expansion pattern observed by Landsat satellites: From 1972 to 2019, *Science China Earth Sciences*, 64, 1922–1933, <https://doi.org/10.1007/s11430-020-9797-9>, 2021.

- Jia, M., Mao, D., Wang, Z., Ren, C., Zhu, Q., Li, X., and Zhang, Y.: Tracking long-term floodplain wetland changes: A case study in the China side of the Amur River Basin, *Int. J. Appl. Earth Observ.*, 92, 102185, <https://doi.org/10.1016/j.jag.2020.102185>, 2020.
- Jia, M., Wang, Z., Mao, D., Ren, C., Wang, C., and Wang, Y.: Rapid, robust, and automated mapping of tidal flats in China using time series Sentinel-2 images and Google Earth Engine, *Remote Sens. Environ.*, 255, 112285, <https://doi.org/10.1016/j.rse.2021.112285>, 2021.
- Jin, H., Stehman, S. V., and Mountrakis, G.: Assessing the impact of training sample selection on accuracy of an urban classification: a case study in Denver, Colorado, *Int. J. Remote Sens.*, 35, 2067–2081, <https://doi.org/10.1080/01431161.2014.885152>, 2014.
- Khandelwal, A., Karpatne, A., Ravirathinam, P., Ghosh, R., Wei, Z., Dugan, H. A., Hanson, P. C., and Kumar, V.: ReaLSAT, a global dataset of reservoir and lake surface area variations, *Scientific Data*, 9, 1–12, <https://doi.org/10.1038/s41597-022-01449-5>, 2022.
- LaRocque, A., Phiri, C., Leblon, B., Pirotti, F., Connor, K., and Hanson, A.: Wetland Mapping with Landsat 8 OLI, Sentinel-1, ALOS-1 PALSAR, and LiDAR Data in Southern New Brunswick, Canada, *Remote Sens.-Basel*, 12, 2095, <https://doi.org/10.3390/rs12132095>, 2020.
- Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296, 1–22, <https://doi.org/10.1016/j.jhydrol.2004.03.028>, 2004.
- Li, Z., Chen, H., White, J. C., Wulder, M. A., and Hermosilla, T.: Discriminating treed and non-treed wetlands in boreal ecosystems using time series Sentinel-1 data, *Int. J. Appl. Earth Observ.*, 85, 102007, <https://doi.org/10.1016/j.jag.2019.102007>, 2020.
- Liu, L., Zhang, X., Gao, Y., Chen, X., Shuai, X., and Mi, J.: Finer-Resolution Mapping of Global Land Cover: Recent Developments, Consistency Analysis, and Prospects, *Journal of Remote Sensing*, 2021, 1–38, <https://doi.org/10.34133/2021/5289697>, 2021.
- Liu, L., Zhang, X., and Zhao, T.: GWL_FCS30: global 30 m wetland map with fine classification system using multi-sourced and time-series remote sensing imagery in 2020, Zenodo [data set], <https://doi.org/10.5281/zenodo.7340516>, 2022.
- Lu, Y. and Wang, L.: How to automate timely large-scale mangrove mapping with remote sensing, *Remote Sens. Environ.*, 264, 112584, <https://doi.org/10.1016/j.rse.2021.112584>, 2021.
- Ludwig, C., Walli, A., Schleicher, C., Weichselbaum, J., and Riffler, M.: A highly automated algorithm for wetland detection using multi-temporal optical satellite data, *Remote Sens. Environ.*, 224, 333–351, <https://doi.org/10.1016/j.rse.2019.01.017>, 2019.
- Mahdianpari, M., Salehi, B., Mohammadimanesh, F., Homayouni, S., and Gill, E.: The First Wetland Inventory Map of Newfoundland at a Spatial Resolution of 10 m Using Sentinel-1 and Sentinel-2 Data on the Google Earth Engine Cloud Computing Platform, *Remote Sens.-Basel*, 11, 43, <https://doi.org/10.3390/rs11010043>, 2018.
- Mahdianpari, M., Jafarzadeh, H., Granger, J. E., Mohammadimanesh, F., Brisco, B., Salehi, B., Homayouni, S., and Weng, Q.: A large-scale change monitoring of wetlands using time series Landsat imagery on Google Earth Engine: a case study in Newfoundland, *GISci. Remote Sens.*, 57, 1102–1124, <https://doi.org/10.1080/15481603.2020.1846948>, 2020.
- Mao, D., Wang, Z., Du, B., Li, L., Tian, Y., Jia, M., Zeng, Y., Song, K., Jiang, M., and Wang, Y.: National wetland mapping in China: A new product resulting from object-based and hierarchical classification of Landsat 8 OLI images, *ISPRS J. Photogramm.*, 164, 11–25, <https://doi.org/10.1016/j.isprs.2020.03.020>, 2020.
- Mao, D., Wang, Z., Wang, Y., Choi, C. Y., Jia, M., Jackson, M. V., and Fuller, R. A.: Remote Observations in China's Ramsar Sites: Wetland Dynamics, Anthropogenic Threats, and Implications for Sustainable Development Goals, *Journal of Remote Sensing*, 2021, 1–13, <https://doi.org/10.34133/2021/9849343>, 2021.
- Matthews, E. and Fung, I.: Methane emission from natural wetlands: Global distribution, area, and environmental characteristics of sources, *Global Biogeochem. Cy.*, 1, 61–86, <https://doi.org/10.1029/GB001i001p00061>, 1987.
- McCarthy, M. J., Radabaugh, K. R., Moyer, R. P., and Muller-Karger, F. E.: Enabling efficient, large-scale high-spatial resolution wetland mapping using satellites, *Remote Sens. Environ.*, 208, 189–201, <https://doi.org/10.1016/j.rse.2018.02.021>, 2018.
- McOwen, C. J., Weatherdon, L. V., Bochove, J. V., Sullivan, E., Blyth, S., Zockler, C., Stanwell-Smith, D., Kingston, N., Martin, C. S., Spalding, M., and Fletcher, S.: A global map of saltmarshes, *Biodivers. Data J.*, 5, e11764, <https://doi.org/10.3897/BDJ.5.e11764>, 2017.
- Murray, N. J., Phinn, S. R., DeWitt, M., Ferrari, R., Johnston, R., Lyons, M. B., Clinton, N., Thau, D., and Fuller, R. A.: The global distribution and trajectory of tidal flats, *Nature*, 565, 222–225, <https://doi.org/10.1038/s41586-018-0805-8>, 2019.
- Murray, N. J., Worthington, T. A., Bunting, P., Duce, S., Hagger, V., Lovelock, C. E., Lucas, R., Saunders, M. I., Sheaves, M., and Spalding, M.: High-resolution mapping of losses and gains of Earth's tidal wetlands, *Science*, 376, 744–749, <https://doi.org/10.1126/science.abm9583>, 2022.
- Olofsson, P., Foody, G. M., Herold, M., Stehman, S. V., Woodcock, C. E., and Wulder, M. A.: Good practices for estimating area and assessing accuracy of land change, *Remote Sens. Environ.*, 148, 42–57, <https://doi.org/10.1016/j.rse.2014.02.015>, 2014.
- Pekel, J. F., Cottam, A., Gorelick, N., and Belward, A. S.: High-resolution mapping of global surface water and its long-term changes, *Nature*, 540, 418–422, <https://doi.org/10.1038/nature20584>, 2016.
- Radoux, J., Lamarche, C., Van Bogaert, E., Bontemps, S., Brockmann, C., and Defourny, P.: Automated Training Sample Extraction for Global Land Cover Mapping, *Remote Sens.-Basel*, 6, 3965–3987, <https://doi.org/10.3390/rs6053965>, 2014.
- Richardson, D. C., Holgerson, M. A., Farragher, M. J., Hoffman, K. K., King, K. B. S., Alfonso, M. B., Andersen, M. R., Cheruvil, K. S., Coleman, K. A., Farruggia, M. J., Fernandez, R. L., Hondula, K. L., Lopez Moreira Mazacotte, G. A., Paul, K., Peierls, B. L., Rabaey, J. S., Sadro, S., Sanchez, M. L., Smyth, R. L., and Sweetman, J. N.: A functional definition to distinguish ponds from lakes and wetlands, *Sci. Rep.-UK*, 12, 10472, <https://doi.org/10.1038/s41598-022-14569-0>, 2022.
- Sexton, J., Feng, M., Channan, S., Song, X., Kim, D., Noojipady, P., Song, D., Huang, C., Annand, A., and Collins, K.: Earth Science Data Records of Global Forest Cover and Change, User guide, 38, https://lpdaac.usgs.gov/documents/1370/GFCC_ATBD.pdf (last access: 22 November 2022), 2016.

- Sexton, J. O., Song, X.-P., Feng, M., Noojipady, P., Anand, A., Huang, C., Kim, D.-H., Collins, K. M., Channan, S., DiMiceli, C., and Townshend, J. R.: Global, 30 m resolution continuous fields of tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based estimates of error, *Int. J. Digit. Earth*, 6, 427–448, <https://doi.org/10.1080/17538947.2013.786146>, 2013.
- Slagter, B., Tsensbazar, N.-E., Vollrath, A., and Reiche, J.: Mapping wetland characteristics using temporally dense Sentinel-1 and Sentinel-2 data: A case study in the St. Lucia wetlands, South Africa, *Int. J. Appl. Earth Observ.*, 86, 102009, <https://doi.org/10.1016/j.jag.2019.102009>, 2020.
- Spalding, M.: *World atlas of mangroves*, Routledge, A collaborative project of ITTO, ISME, FAO, UNEP-WCMC, UNESCO-MAB, UNU-INWEH and TNC, London (UK), Earthscan, London, <https://doi.org/10.4324/9781849776608>, 2010.
- Tachikawa, T., Hato, M., Kaku, M., and Iwasaki, A.: Characteristics of ASTER GDEM Version 2, *Geoscience and Remote Sensing Symposium (IGARSS)*, 24–29 July 2011, Vancouver, 12477285, 3657–3660, <https://doi.org/10.1109/IGARSS.2011.6050017>, 2011a.
- Tachikawa, T., Kaku, M., Iwasaki, A., Gesch, D. B., Oimoen, M. J., Zhang, Z., Danielson, J., Krieger, T., Curtis, B., and Haase, J.: ASTER Global Digital Elevation Model Version 2 – Summary of validation results, *Kim Fakultas Sastra Dan Budaya*, <https://doi.org/10.1093/oxfordjournals.pubmed.a024792>, 2011b.
- Thomas, N., Lucas, R., Bunting, P., Hardy, A., Rosenqvist, A., and Simard, M.: Distribution and drivers of global mangrove forest change, 1996–2010, *PLoS One*, 12, e0179302, <https://doi.org/10.1371/journal.pone.0179302>, 2017.
- Tootchi, A., Jost, A., and Ducharme, A.: Multi-source global wetland maps combining surface water imagery and ground-water constraints, *Earth Syst. Sci. Data*, 11, 189–220, <https://doi.org/10.5194/essd-11-189-2019>, 2019.
- Torres, R., Snoeijs, P., Geudtner, D., Bibby, D., Davidson, M., Attema, E., Potin, P., Rommen, B., Floury, N., Brown, M., Traver, I. N., Deghaye, P., Duesmann, B., Rosich, B., Miranda, N., Bruno, C., L'Abbate, M., Croci, R., Pietropaolo, A., Huchler, M., and Rostan, F.: GMES Sentinel-1 mission, *Remote Sens. Environ.*, 120, 9–24, <https://doi.org/10.1016/j.rse.2011.05.028>, 2012.
- Townshend, J. R., Masek, J. G., Huang, C., Vermote, E. F., Gao, F., Channan, S., Sexton, J. O., Feng, M., Narasimhan, R., Kim, D., Song, K., Song, D., Song, X.-P., Noojipady, P., Tan, B., Hansen, M. C., Li, M., and Wolfe, R. E.: Global characterization and monitoring of forest cover using Landsat data: opportunities and challenges, *Int. J. Digit. Earth*, 5, 373–397, <https://doi.org/10.1080/17538947.2012.713190>, 2012.
- Veci, L., Prats-Iraola, P., Scheiber, R., Collard, F., Fomferra, N., and Engdahl, M.: The sentinel-1 toolbox, <https://sentinels.copernicus.eu/web/sentinel/toolboxes/sentinel-1> (last access: 25 May 2022), 2014.
- Vermote, E., Justice, C., Claverie, M., and Franch, B.: Preliminary analysis of the performance of the Landsat 8/OLI land surface reflectance product, *Remote Sens. Environ.*, 185, 46–56, <https://doi.org/10.1016/j.rse.2016.04.008>, 2016.
- Wang, X., Xiao, X., Zou, Z., Hou, L., Qin, Y., Dong, J., Doughty, R. B., Chen, B., Zhang, X., Chen, Y., Ma, J., Zhao, B., and Li, B.: Mapping coastal wetlands of China using time series Landsat images in 2018 and Google Earth Engine, *ISPRS J. Photogramm.*, 163, 312–326, <https://doi.org/10.1016/j.isprsjprs.2020.03.014>, 2020.
- Wang, X., Xiao, X., Xu, X., Zou, Z., Chen, B., Qin, Y., Zhang, X., Dong, J., Liu, D., Pan, L., and Li, B.: Rebound in China's coastal wetlands following conservation and restoration, *Nature Sustainability*, 4, 1076–1083, <https://doi.org/10.1038/s41893-021-00793-5>, 2021.
- Wilén, B. O. and Bates, M.: The US fish and wildlife service's national wetlands inventory project, in: *Classification and inventory of the world's wetlands*, Springer, <https://doi.org/10.1007/BF00045197>, 1995.
- Worthington, T. A., Zu Ermgassen, P. S., Friess, D. A., Krauss, K. W., Lovelock, C. E., Thorley, J., Tingey, R., Woodroffe, C. D., Bunting, P., and Cormier, N.: A global biophysical typology of mangroves and its relevance for ecosystem structure and deforestation, *Sci. Rep.-UK*, 10, 1–11, <https://doi.org/10.1038/s41598-020-71194-5>, 2020.
- Xi, Y., Peng, S., Ciais, P., and Chen, Y.: Future impacts of climate change on inland Ramsar wetlands, *Nat. Clim. Change*, 11, 45–51, <https://doi.org/10.1038/s41558-020-00942-2>, 2020.
- Zanaga, D., Van De Kerchove, R., De Keersmaecker, W., Souverijns, N., Brockmann, C., Quast, R., Wevers, J., Grosu, A., Paccini, A., and Vergnaud, S.: ESA WorldCover 10 m 2020 v100, Zenodo, <https://doi.org/10.5281/zenodo.5571936>, 2021.
- Zhang, H., Wang, T., Liu, M., Jia, M., Lin, H., Chu, L. M., and Devlin, A.: Potential of Combining Optical and Dual Polarimetric SAR Data for Improving Mangrove Species Discrimination Using Rotation Forest, *Remote Sens.-Basel*, 10, 467, <https://doi.org/10.3390/rs10030467>, 2018.
- Zhang, H. K. and Roy, D. P.: Using the 500 m MODIS land cover product to derive a consistent continental scale 30 m Landsat land cover classification, *Remote Sens. Environ.*, 197, 15–34, <https://doi.org/10.1016/j.rse.2017.05.024>, 2017.
- Zhang, X., Liu, L., Wu, C., Chen, X., Gao, Y., Xie, S., and Zhang, B.: Development of a global 30 m impervious surface map using multisource and multitemporal remote sensing datasets with the Google Earth Engine platform, *Earth Syst. Sci. Data*, 12, 1625–1648, <https://doi.org/10.5194/essd-12-1625-2020>, 2020.
- Zhang, X., Liu, L., Chen, X., Gao, Y., and Jiang, M.: Automatically Monitoring Impervious Surfaces Using Spectral Generalization and Time Series Landsat Imagery from 1985 to 2020 in the Yangtze River Delta, *Journal of Remote Sensing*, 2021, 1–16, <https://doi.org/10.34133/2021/9873816>, 2021a.
- Zhang, X., Liu, L., Chen, X., Gao, Y., Xie, S., and Mi, J.: GLC_FCS30: global land-cover product with fine classification system at 30 m using time-series Landsat imagery, *Earth Syst. Sci. Data*, 13, 2753–2776, <https://doi.org/10.5194/essd-13-2753-2021>, 2021b.
- Zhang, X., Liu, L., Zhao, T., Gao, Y., Chen, X., and Mi, J.: GISD30: global 30 m impervious-surface dynamic dataset from 1985 to 2020 using time-series Landsat imagery on the Google Earth Engine platform, *Earth Syst. Sci. Data*, 14, 1831–1856, <https://doi.org/10.5194/essd-14-1831-2022>, 2022.
- Zhang, Z., Xu, N., Li, Y., and Li, Y.: Sub-continental-scale mapping of tidal wetland composition for East Asia: A novel algorithm integrating satellite tide-level and phenological features, *Remote Sens. Environ.*, 269, 112799, <https://doi.org/10.1016/j.rse.2021.112799>, 2022.

- Zhu, P. and Gong, P.: Suitability mapping of global wetland areas and validation with remotely sensed data, *Science China Earth Sciences*, 57, 2283–2292, <https://doi.org/10.1007/s11430-014-4925-1>, 2014.
- Zhu, Z. and Woodcock, C. E.: Object-based cloud and cloud shadow detection in Landsat imagery, *Remote Sens. Environ.*, 118, 83–94, <https://doi.org/10.1016/j.rse.2011.10.028>, 2012.
- Zhu, Z., Wang, S. X., and Woodcock, C. E.: Improvement and expansion of the Fmask algorithm: cloud, cloud shadow, and snow detection for Landsats 4-7, 8, and Sentinel 2 images, *Remote Sens. Environ.*, 159, 269–277, <https://doi.org/10.1016/j.rse.2014.12.014>, 2015.
- Zhu, Z., Gallant, A. L., Woodcock, C. E., Pengra, B., Olofsson, P., Loveland, T. R., Jin, S., Dahal, D., Yang, L., and Auch, R. F.: Optimizing selection of training and auxiliary data for operational land cover classification for the LCMAP initiative, *ISPRS J. Photogramm.*, 122, 206–221, <https://doi.org/10.1016/j.isprsjprs.2016.11.004>, 2016.