



Supplement of

Multiyear high-temporal-resolution measurements of submicron aerosols at 13 French urban sites: data processing and chemical composition

Hasna Chebaicheb et al.

Correspondence to: Hasna Chebaicheb (hasna.chebaicheb@ineris.fr)

The copyright of individual parts of the supplement might differ from the article licence.

Table S1: Description of the measurement stations from the CARA program used in this paper. A detailed description of these sites is presented at the end of SI, section “Sites description”.

<i>Operator</i>	Station name	City	Airbase ID	Altitude (m a.s.l)	Latitude	Longitude	Site type
<i>IMT Nord Europe</i>	ATOLL	Lille	-	70	50.6111	3.1404	Suburban
<i>LSCE & Ineris</i>	SIRTA	Greater Paris	-	163	48.71181	2.14818	Suburban
<i>Airparif</i>	Gennevilliers	Greater Paris	FR04002	28	48.9298083	2.2946194	Urban background
<i>Airparif</i>	Paris Les Halles	Paris	FR04055	35	48.8627083	2.3446972	Urban background
<i>Airparif</i>	Paris BPEst	Paris	FR04329	49	48.8385167	2.4126242	Urban Traffic
<i>Air Breizh</i>	Rennes Pays-Bas	Rennes	FR19017	45	48.08965	-1.65911	Urban background
<i>Atmo AuRA</i>	Lyon Centre	Lyon	FR20062	160	45.75779	4.85422	Urban background
<i>Atmo Grand Est</i>	Metz Borny	Metz	FR01012	204	49.1102806	6.2233361	Urban background
<i>Atmo Grand Est</i>	Strasbourg Danube	Strasbourg	FR42010	135	48.5062222	7.7511806	Urban background
<i>Atmo Hauts-de-France</i>	Creil Faïencerie	Creil	FR18043	31	49.2597222	2.4744444	Urban background
<i>AtmoSud & LCE</i>	Marseille Longchamp	Marseille	FR03043	73	43.3052889	5.3947056	Urban background
<i>Atmo Nouvelle-Aquitaine</i>	Poitiers Augouard	Poitiers	FR09404	105	46.5839885	0.3455967	Urban background
<i>Atmo Nouvelle-Aquitaine</i>	Talence	Bordeaux	FR31002	23	44.800442	-0.5893941	Urban background

Table S2: Key aspects of AMS/ACSM instruments.

Instrument	type	Advantages	Disadvantages
AMS	cToF-AMS or HR-ToF-AMS	<ul style="list-style-type: none"> - field campaign - size distribution - high-time (~2-5 min) and mass resolution (HR-ToF-AMS) 	<ul style="list-style-type: none"> - require more monitoring and technical maintenance
ACSM	-Q-ACSM	<ul style="list-style-type: none"> - long-term measurements - less maintenance - time resolution of ~30 min 	<ul style="list-style-type: none"> - no size distribution - unit mass resolution
	-ToF-ACSM	<ul style="list-style-type: none"> - long term measurements - time resolution of about 10min - higher mass resolution ($m/\Delta m \approx 300$) 	<ul style="list-style-type: none"> - no size distribution - data treatment procedures relatively complex

Table S3: Percentages of data that fell within the range of $-3 \times DL$ to DL and data below $-3 \times DL$.

Station name	Pourcentages of data below $-3 \times DL$						Percentages of data between $-3 \times DL$ and DL					
	eBC	OA	NH ₄	SO ₄	NO ₃	Cl	eBC	OA	NH ₄	SO ₄	NO ₃	Cl
ATOLL	NaN	NaN	0.004	0.004	0.004	3.70	6.27	0.49	23	0.46	0.28	33
Paris BPEst	NaN	0.016	NaN	0.032	NaN	8.24	NaN	0.16	24	0.66	NaN	32
Creil	NaN	NaN	NaN	0.005	0.005	1.52	6.12	0.66	25	0.81	0.27	18
Gennevilliers	NaN	NaN	NaN	NaN	NaN	0.01	1.78	0.15	25	0.10	0.01	16
Lyon	NaN	NaN	NaN	0.002	NaN	0.00	1.67	0.04	31	0.40	0.02	26
Marseille	NaN	NaN	NaN	NaN	NaN	NaN	1.04	0.49	30	1.10	1.31	37
Metz	NaN	0.055	0.075	0.106	0.068	0.66	6.33	0.42	22	0.36	0.14	26
Paris Les Halles	NaN	NaN	NaN	0.283	NaN	0.87	1.06	0.19	41	3.44	NaN	35
Poitiers	NaN	0.045	0.003	0.154	0.008	4.81	7.35	1.50	33	2.09	0.36	25
Rennes	NaN	0.027	0.018	0.124	0.009	0.24	11.59	1.66	53	2.37	0.84	18
SIRTA	NaN	NaN	0.002	0.008	0.015	0.38	11.56	0.44	31	0.55	0.16	20
Strasbourg	NaN	NaN	NaN	0.526	NaN	0.15	2.18	0.51	40	2.69	0.13	32
Talence	NaN	NaN	NaN	0.036	NaN	5.95	7.37	1.12	36	1.45	0.34	26




Figure S1: $NR-PM_{1} + eBC$ vs. $PM_{2.5}$ for each site during the whole study period.




Figure S2: Box plots of statistical information (mean in circle, median, the percentiles 5, 25, 75, and 95th) of eBC_{fr} and eBC_{wb} for the 13 French sites.




Figure S3: Pollution rose of Chloride at Creil site (Zefir).




Figure S4: Relative contribution of chemical species as a function of PM₁ percentiles by season (continued)




Figure S4: Relative contribution of chemical species as a function of PM₁ percentiles by season (continued).




Figure S4: Relative contribution of chemical species as a function of PM₁ percentiles by season.




Figure S5: Monthly concentrations normalized by site-specific annual averages of the species studied here.




Figure S6: Monthly IQR/median ratios for each component from Fig. 4 from the main text.




Figure S7: Diel concentration profiles, normalized by site-specific daily averages.




Figure S8: Seasonal diel concentration profiles, normalized by site-specific daily averages.






Figure S9: Diel median and IQR for each species across all sites.








Figure S10: Temporal variability from simulations and observations for PM₁ chemical species at each site.




Figure S11: Observed and modeled diel profiles during the spring and autumn of 2018 across nine French sites. The shaded areas correspond to the interquartile range.




Figure S12: Scatterplots of comparison between daily averages of simulations and either online (in red; ACSM or AE33) or offline observations (in green; filter analyses) (in $\mu\text{g m}^{-3}$) for OA, NO₃, NH₄, SO₄, and eBC at four French urban sites (Talence, Poitiers, Lyon, and Marseille).

Sites description

A description of the sites discussed in this paper is presented below. For the ATOLL, SIRTA, and Marseille stations, their details can be found in Chebaicheb et al. 2023, Zhang et al. 2019, and Chazeau et al. 2021 et 2022, respectively.

1. Paris Les Halles

The "Paris Les Halles" is an urban background station, located in the 1st district of Paris, 22 km from SIRTA. Les Halles, in the heart of Paris, is characterized by high local activity, with green spaces, shopping malls, restaurants, and numerous areas dedicated to leisure and culture, as well as the Châtelet - Les Halles RER train station, which is the largest underground station and provides access to the entire Paris region network.

The station is multi-instrumented, with permanent in-situ measurements since October 2019. Samples are taken at heights of between 1.55 and 2 m above ground level, with a 270° clearance to ensure optimum representativeness of measurements.



Figure S12: Map indicating the location of the Airparif Paris 1st-Les Halles station (top); photographs (credit: AirParif) of the sampling heads (bottom left) and the interior of the station (bottom right).

2. Boulevard Périphérique Est (BPEst)

The Boulevard Périphérique Est station, located in the 12th district of Paris, is an urban traffic station that has been in operation since January 2013. The Boulevard Périphérique surrounds the Paris city and is 35 km long. In general, it has four lanes in each direction. The largest traffic flows in the Paris region remain concentrated on the boulevard périphérique, with a traffic record of 180,000 to 250,000 vehicles per day according to the Atelier Parisien d'urbanisme (APUR) in 2019. This road generates significant emissions of gaseous and particulate pollutants into the air.

Samples are taken at heights ranging from 2.8 to 3.1 m above ground level. Sampling distances are 5 m to the road edge and 6.5 m to the center of the first traffic lane. The measurement axis covers 2×4 lanes with a total width of 34 m.

These measurements are used to monitor air quality in the vicinity of Paris's Boulevard Périphérique Est, assessing gaseous pollutants, fine particles, and aerosol levels. By focusing on an environment characterized by dense traffic, this station contributes to understanding the impact of road traffic-related pollution on air quality in this urban area.



Figure S13: Map detailing the location of the BPEST station (left) and a photograph (credit: AirParif) of the sampling heads at the edge of the ring road (right).

3. Gennevilliers

The Gennevilliers station is an urban background station, located 2.1 km northwest of Paris, and has been in operation since January 1986. Gennevilliers is a densely populated urban commune (49,880 inhabitants in 2020; 4,300 inhabitants/km²). It lies at the crossroads of two freeways, the A15 and the A86, and covers an area of 1,200 hectares. Two-thirds of its surface area is devoted to economic and industrial activities. The commune is located in the Hauts-de-Seine department, a community rich in river transport.

Samples are taken at a height of around 15 m above ground level, on the roof of the Collège Louis Pasteur, to obtain representative ambient air samples. What's more, the station enjoys 360° clearance, ensuring unobstructed sampling around the site.



Figure S14: Map specifying the location of the Gennevilliers station (left) and a photograph (credit: AirParif) of the sampling heads on the roof of Louis Pasteur College (right).

4. Creil

The Creil monitoring station is an urban station located in the Oise department, in the Hauts-de-France region. It is specifically located in the center of the commune of Creil, which is the center of an urban unit comprising 22 communes and home to almost 120,000 inhabitants, making it the largest conurbation in the Oise region. Creil is close to major cities such as Paris, Beauvais, and Amiens, at distances of 45.35 and 72 km respectively. The town is crossed by the Oise River and is surrounded by two forests, the Halatte forest to the east and the Chantilly forest to the southeast.

The Creil measuring station is located in an urban environment influenced by both urban traffic and residential emissions. These measurements enable us to monitor air quality in the commune of Creil and assess the impact of emissions from road traffic, residential activities, and other local sources on air pollution in this urban area.



Figure S15: Map specifying the location of the Creil station (left) and a photograph (credit: Atmo Hauts-de-France) of the exterior of the station (right).

5. Lyon

The Lyon Centre urban background site, located in the Bassin Lyonnais/Nord Isère area, has been operational since January 5, 2007. Located in the heart of Lyon city, the monitoring station plays an essential role in collecting data on air quality in this densely populated agglomeration. It is France's third largest city, with 522,969 inhabitants

in 2019 (INSEE, 2019). Lyon is the capital of the Auvergne-Rhône-Alpes region, which covers an area of 69,711 km². The city is known for its strong industrial activity, but also its agricultural and agri-food activities.

This site is representative of the Lyon population's exposure to atmospheric pollution; it characterizes the air breathed by all Lyon residents. In addition, the monitoring station is strategically positioned to capture pollution levels resulting from urban activities such as road traffic, industry, and domestic emissions. It thus provides crucial information for assessing the effectiveness of pollution reduction measures and supporting the implementation of environmental policies towards improving air quality in the region.




Figure S16: Satellite images pinpointing the location of the Lyon Centre station on a large scale (top left) and a small scale (top right), along with photographs (credit: Atmo AuRA) showing an exterior view (bottom left) and an interior view of the measurement station (bottom right).

6. Metz

The Metz measurement station is located in the Borny district, which is situated in the southeast of the city center of Metz. The city is located halfway between Strasbourg (130 km) and Reims (155 km). It is 55 km east from the Paris region. Its population density is 2,866.3 inhabitants/km² in 2020 (120,211 inhabitants). It is known for its strong economic and industrial activity, notably metallurgy, petrochemicals, automotive, logistics, and commerce. This station is considered an urban background site and is located in a residential area. On the same street, there are mainly single-family homes and a nursery school, which means there is plenty of parking for school buses and parents. To the south of the station, there is also a shopping area with restaurants. These environmental features are important to consider when interpreting the measurement data, as they can influence air quality in this particular area. The proximity of homes, schools, and commercial activities can contribute to the exposure of the local population to air pollutants. For this reason, monitoring this urban background station is of particular importance in assessing the impact of human activities on air quality in this area.




Figure S17: Map showing the location of Atmo Grand-Est's measuring station in Metz. (© Google Maps)

7. Strasbourg

The Danube station is located in the Neudorf district, south-east of Strasbourg, and has been in operation from December 2019 to the present day. The Neudorf district, where Danube station is located, is an eco-district currently under construction. It is surrounded by residential areas to the south and north, with a commercial zone including restaurants. To the east of the station, there is also a river port and a major thoroughfare (Avenue du Rhin) less than 100 m away.

Strasbourg is a predominantly urban agglomeration, representing 26% of the Strasbourg Urban Community (CUS), with a strong agricultural base covering 40% of the CUS. It also boasts five major industrial sites. The city is served by two major roads, the A35-A4 freeway (north/south axis) and the Rhine road (east/west axis). There are also rail links, with a TER station and a TGV station in Strasbourg, as well as an international airport in Entzheim and a river port in Strasbourg.

In terms of meteorology and topography, Strasbourg has a semi-continental climate, with low precipitation and limited sunshine. The preferred wind directions are south-southwest and north-northeast, and air masses are predominantly wind-driven, with some influence from Central European air masses.



Figure S18: A map specifying the positions of the two Strasbourg measurement stations.

8. Talence

The Talence station is an urban background station, set back from major traffic arteries. It is a reference station for monitoring particulate matter. Talence is part of the Bordeaux metropolitan area, which comprises 73 communes. It is one of the most densely populated communes in the region, with over 5,312.5 inhabitants/km² (44,359 inhabitants) in 2020. It is a highly urbanized city, with a dominant service sector. On the other hand, there is very little industrial or agricultural activity. It is surrounded by regional nature parks, notably the Landes de Gascogne forest to the southwest and the Parc du Médoc to the northwest.



Figure S19: A map and satellite image precisely indicating the location of the Talence station on a large scale (left) and a small scale (right)

9. Poitiers

The Poitiers Augouard station is an urban background station located on rue Augouard in downtown Poitiers. The Poitiers agglomeration, located halfway between Paris and Bordeaux, is a densely populated urban area with 90,033 inhabitants in 2020. The measurement station plays an important role as a reference station for fine particulate monitoring in the region. It has been equipped with an AE33 since 2014 and an ACSM since 2015 to continuously analyze the chemical composition of fine particles in ambient air. In addition to these measurements, the station also carries out measurements of gaseous pollutants.



Figure S20: A map and satellite image providing precise details on the location of the Poitiers Augouard station on a large scale (left) and a small scale (right).

10. Rennes

The "Pays-Bas" station in Rennes is an urban background station that was established on August 30, 2012. Rennes is located in north-western France and is the capital of the Brittany region. It has a population of 354,000 in 2020. Rennes lies at the junction of the Channel-Atlantic link and is characterized by an oceanic climate. Within the Rennes metropolitan area, road transport and industry make smaller contributions. On the other hand, the agricultural and residential sectors play a greater role in PM₁₀ and PM_{2.5} emissions (AASQA, 2022).

The station is located in the south of Rennes, in a residential area served by the metro, and includes several apartment buildings, some of which are connected to the biomass heating network. Near the station, there is an unpaved footpath and a parking lot to the southeast.




Figure S21: Satellite images providing precise location details of the Rennes Pays-Bas station on a large scale (left) and a small scale (right).

References

AASQA: Rapport Caractérisation des particules à Rennes Bilan 2020, 2022.

APUR:<https://cities.newstank.fr/article/view/148500/paris-180000-250000-vehicules-jour-peripherique-selon-apur.html>, last access: 22 August 2023.

Chazeau, B., Temime-Roussel, B., Gille, G., Mesbah, B., D'Anna, B., Wortham, H., and Marchand, N.: Measurement report: Long-term real-time characterisation of the submicronic aerosol and its atmospheric dynamic in a Mediterranean coastal city: Tracking the polluted events at the Marseille-Longchamp supersite, *Atmospheric Chem. Phys. Discuss.*, 1–44, <https://doi.org/10.5194/acp-2020-1015>, 2020.

Chazeau, B., Temime-Roussel, B., Gille, G., Mesbah, B., D'Anna, B., Wortham, H., and Marchand, N.: Measurement report: Fourteen months of real-time characterisation of the submicronic aerosol and its atmospheric dynamics at the Marseille–Longchamp supersite, *Atmospheric Chem. Phys.*, 21, 7293–7319, <https://doi.org/10.5194/acp-21-7293-2021>, 2021.

Chebaicheb, H., F. de Brito, J., Chen, G., Tison, E., Marchand, C., Prévôt, A. S. H., Favez, O., and Riffault, V.: Investigation of four-year chemical composition and organic aerosol sources of submicron particles at the ATOLL site in northern France, *Environ. Pollut.*, 330, 121805, <https://doi.org/10.1016/j.envpol.2023.121805>, 2023.

Insee: <https://www.insee.fr/fr/statistiques/4481460>, last access: 13 October 2021.

Zhang, Y., Favez, O., Petit, J.-E., Canonaco, F., Truong, F., Bonnaire, N., Crenn, V., Amodeo, T., Prévôt, A. S. H., Sciare, J., Gros, V., and Albinet, A.: Six-year source apportionment of submicron organic aerosols from near-continuous highly time-resolved measurements at SIRTa (Paris area, France), *Atmospheric Chem. Phys.*, 19, 14755–14776, <https://doi.org/10.5194/acp-19-14755-2019>, 2019.