Earth Syst. Sci. Data, 17, 2681–2692, 2025 https://doi.org/10.5194/essd-17-2681-2025 © Author(s) 2025. This work is distributed under the Creative Commons Attribution 4.0 License.



# The SahulCHAR collection: a palaeofire database for Australia, New Guinea, and New Zealand

Emma Rehn<sup>1</sup>, Haidee Cadd<sup>2</sup>, Scott Mooney<sup>3</sup>, Tim J. Cohen<sup>2</sup>, Henry Munack<sup>2</sup>, Alexandru T. Codilean<sup>2</sup>, Matthew Adeleye<sup>4</sup>, Kristen K. Beck<sup>5</sup>, Mark Constantine IV<sup>2</sup>, Chris Gouramanis<sup>6</sup>, Johanna M. Hanson<sup>7</sup>, Penelope J. Jones<sup>8</sup>, A. Peter Kershaw<sup>9</sup>, Lydia Mackenzie<sup>10</sup>, Maame Maisie<sup>3</sup>, Michela Mariani<sup>11</sup>, Kia Matley<sup>12</sup>, David McWethy<sup>13</sup>, Keely Mills<sup>14</sup>, Patrick Moss<sup>15</sup>, Nicholas R. Patton<sup>16,17</sup>, Cassandra Rowe<sup>1</sup>, Janelle Stevenson<sup>18</sup>, John Tibby<sup>19</sup>, and Janet Wilmshurst<sup>20</sup> <sup>1</sup>ARC Centre of Excellence for Australian Biodiversity and Heritage, College of Arts, Society, and Education, James Cook University, Cairns, 4870, Australia <sup>2</sup>ARC Centre of Excellence for Australian Biodiversity and Heritage, School of Earth, Atmospheric, and Life Sciences, University of Wollongong, Wollongong, 2500, Australia <sup>3</sup>ARC Centre of Excellence for Australian Biodiversity and Heritage, School of BEES, University of New South Wales, Sydney, 2052, Australia <sup>4</sup>Department of Geography, University of Cambridge, Cambridge, CB2 1DB, Cambridgeshire, United Kingdom <sup>5</sup>Catchments and Coasts Research Group, Department of Geography, University of Lincoln, Brayford Pool, Lincoln, LN6 7TS, United Kingdom <sup>6</sup>Research School of Earth Sciences, The Australian National University, Canberra, Australia, 0200, Australia <sup>7</sup>School of Earth and Environment, University of Canterbury, Ōtautahi / Christchurch, 8041, Aotearoa / New Zealand <sup>8</sup>Menzies Institute for Medical Research, University of Tasmania, Hobart, 7000, Australia <sup>9</sup>School of Earth, Atmosphere and Environment, Monash University, Clayton, 3800, Australia <sup>10</sup>School of Geography, Planning and Spatial Sciences, University of Tasmania, Hobart, 7001, Australia <sup>11</sup>School of Geography, University of Nottingham, Nottingham, NG72RD, United Kingdom <sup>12</sup>School of BioSciences, The University of Melbourne, Parkville, 3010, Australia <sup>13</sup>Department of Earth Sciences, Montana State University, Bozeman, Montana, 59715, USA <sup>14</sup>British Geological Survey, Keyworth, Nottingham, NG12 5GG, United Kingdom <sup>15</sup>School of Earth & Atmospheric Sciences, Queensland University of Technology, Brisbane, 4072, Queensland, Australia <sup>16</sup>Department of Geosciences, Idaho State University, Pocatello, ID, USA <sup>17</sup>School of Earth and Environment, University of Canterbury, Ōtautahi / Christchurch, 8041, Aotearoa / New Zealand <sup>18</sup>ARC Centre of Excellence for Australian Biodiversity and Heritage, School of Culture, History and Language, The Australian National University, Canberra, 2601, Australia <sup>19</sup>Geography, Environment and Population, University of Adelaide, Adelaide, 5005, Australia <sup>20</sup>Manaaki Whenua – Landcare Research, P.O. Box 69040, Lincoln, 7640, Aotearoa/New Zealand Correspondence: Haidee Cadd (haidee@uow.edu.au)

> Received: 22 August 2024 – Discussion started: 1 October 2024 Revised: 4 March 2025 – Accepted: 6 March 2025 – Published: 16 June 2025

**Abstract.** Recent global fire activity has highlighted the importance of understanding fire dynamics across time and space, with records of past fire (palaeofire) providing valuable insights to inform us on current and future management challenges. New records from the recent increase in palaeofire studies from Australia and surrounds have not been captured in any database for broader comparisons, and Australasia is poorly represented in current international databases used for global modelling of palaeofire trends. These problems are addressed

by SahulCHAR, a new collection of sedimentary charcoal and black carbon records from Sahul (Australia, New Guinea, and offshore islands) and Aotearoa/New Zealand. Data are stored in the OCTOPUS relational database platform, with a structure designed for compatibility with the existing Global Paleofire Database. Metadata are captured at the site level and observation level, with observations including age determinations and charcoal or black carbon data. SahulCHAR version 1 contains 687 records of charcoal or black carbon, including digitized data, unchanged and modified records from the Global Paleofire Database, and original author-submitted data. SahulCHAR is a much-needed update to past regional palaeofire compilations that will also provide greater representation of records from Sahul and Aotearoa/New Zealand in future global syntheses.

## 1 Introduction

Fire is a key ecosystem process with characteristics that vary widely across biomes globally, with "fire-dependent" ecosystems covering around half of the terrestrial globe (Shlisky et al., 2007, p. 6). Recent increases in global fire activity, including extreme fires in Australia and elsewhere, have highlighted the critical importance of understanding fire dynamics across time and space (Duane et al., 2021; Nolan et al., 2021), with more extreme fire weather predicted in the future for southeastern Australia and northern and eastern Aotearoa/New Zealand as a result of climate change (Lawrence et al., 2022).

Palaeofire data (sedimentary charcoal and black carbon, defined as carbonaceous substances of pyrogenic origin) can offer important insights into past fire variability to inform us on current and future challenges, including climatefire-vegetation interactions (Williams and Abatzoglou, 2016; Marlon, 2020). Compilations of palaeofire records have been used to investigate long-term relationships and shifting dynamics between humans, fire, vegetation, and climate in Australia (Lynch et al., 2007; Enright and Thomas, 2008; Williams et al., 2015; Williams, 2011), Aotearoa/New Zealand (McWethy et al., 2010; Perry et al., 2014), and Indonesia and Papua New Guinea (Haberle et al., 2001). Understanding fire regimes over long timescales in Australia and the surrounding region has increasingly become a research priority, reflected in a recent influx of new palaeofire records (for examples from just the past 2 years, see Adeleye et al., 2023; Constantine et al., 2023; Hanson et al., 2022; Laming et al., 2022; Patton et al., 2023; Rowe et al., 2022; Thomas et al., 2022). However, the last major compilation and synthesis of sedimentary charcoal records from Australasia was done by Mooney et al. (2011, 2012), containing 224 sedimentary charcoal records. These records were primarily derived from the Global Paleofire Database version 2 2 (GPDv2, Daniau et al. 2012) and version 2.5 (GPDv2.5. Marlon et al., 2013, 2016). These records are now included in the Global Paleofire Database (GPDv4, International Paleofire Network, 2023). More recent syntheses have been focused on specific regions, such as Mariani et al.'s (2022, 2024) compilation of over 100 charcoal records from southeastern Australia as contained in the GPD to investigate human-fire-vegetation dynamics over the last thousand years. The diverse environments of Australia, New Guinea, and Aotearoa/New Zealand have unique histories of fire-climate-human interactions (Mooney et al., 2011, Fletcher et al., 2021). As identified by Mooney et al. (2011) and Rowe et al. (2023), no individual palaeofire record should be considered to be representative of this vast region; to disentangle long-term influences on fire and potential variations across subregions and ecosystems, a large dataset is required.

Major global databases containing charcoal data such as the GPD, the Reading Palaeofire Database (RPD; Harrison et al., 2022), the Neotoma Paleoecology Database (Neotoma; Williams et al., 2018), and PANGAEA (Feldner et al., 2023) are lacking many palaeofire records from Australasia. The GPD currently contains 179 cores with associated charcoal data from Australia, 23 cores from New Guinea (Papua New Guinea and West Papua), and 10 cores from Aotearoa/New Zealand. These records are also contained in the RPD, and even fewer cores from this region are available in Neotoma (17 cores) and PANGAEA (11 cores, 6 of which are also contained in the GPD). The GPD is a valuable resource for regional and global palaeofire syntheses (e.g. Daniau et al., 2012; Karp et al., 2021; Marlon et al., 2013, 2016: Power et al., 2010), but it requires a significant update to capture the many new palaeofire records now available from Australia, New Guinea, and Aotearoa / New Zealand. As noted by Harrison et al. (2022), the current limitations of the GPD include potential duplicates of sites, missing metadata and age data, and the necessity of updates to incorporate newly published records. The GPD also contains lengthy yet still incomprehensive lists of metadata options, in part due to the array of ways to approach charcoal analysis (e.g. Mooney and Tinner, 2010; Turner et al., 2004), as well as ad hoc user additions and structural constraints (most notably, a single field for measurement units that includes size ranges).

These limitations are addressed for Sahul (Australia, New Guinea, and continental islands) and Aotearoa/New Zealand by the SahulCHAR data collection (Rehn et al. 2024). The purpose of this paper is to introduce the data structure of SahulCHAR and to provide an overview of the data compiled in version 1. In keeping with existing OCTOPUS collections, SahulCHAR was named and intended to have a

## E. Rehn et al.: The SahulCHAR collection: a palaeofire database for Australia, New Guinea, and Aotearoa



**Figure 1.** Graphical representation of the OCTOPUS semantic database model featuring the fully integrated SahulCHAR partner collection. SahulCHAR shares parent and/or lookup tables with the other collections (SahulArch, SahulSED, the IPPD, FosSahul, ExpAge, and CRN) on the global, regional, and bibliographic level.

geographic focus on the Sahul landmass (Australia, New Guinea, and continental islands); during data collection, the geographic scope was extended to include Aotearoa/New Zealand. SahulCHAR data collection was designed to capture new records published since the compilation by Mooney et al. (2011, 2012), to capture older records not previously entered into the GPD, to check (and correct, if required) the details of records in the GPD from this region, and to capture additional metadata wherever possible for records available in the GPD, all data from the GPD were carefully screened during data entry for SahulCHAR (Table S1 in the Supplement). Published sources – including but not limited to sources listed in GPD records – were used to modify and

add to the metadata and charcoal data captured in the GPD wherever possible as part of a greater literature review to identify charcoal records in the region.

#### 2 Data structure and compilation

SahulCHAR is hosted on the OCTOPUS platform (https: //octopusdata.org, last access: 16 March 2025), with data being stored in a relational PostgreSQL database (Fig. 1). For a full description of the OCTOPUS v.2 system architecture, see Codilean et al. (2022), and for the full and up-todate OCTOPUS documentation, see Munack et al. (2023). For compatibility and future integration, the data structure of SahulCHAR is broadly based on the metadata captured in **Table 1.** Table representation of site-level metadata collected for SahulCHAR attributes vs. corresponding GPD fields, where applicable. For a full description of the database tables refer to Munack et al. (2023) and Munack and Codilean (2023).

Metadata field	Description	Field type	Example	Corresponding GPD field
METASITE	Metasite name	Free text	Big Willum Swamp	NA
SITE	Site name	Free text	Big Willum Swamp BWIL2	site_name
COUNTRY	Country where metasite is located	Predefined list	Australia	country_name
SITECODE	Site type, based on primary characteris- tics at the time of collection	Predefined list	Terrestrial, bog For full list, see global-sitecode- fields	site_type_desc
BASIN	Basin size	Predefined list	Large (50.1–500 km <sup>2</sup> ) For full list, see basinsizeid-fields	basin_size_desc
CATCHMENT	Catchment size	Predefined list	Small (<10 km <sup>2</sup> ) For full list, see catchmentsizeid- fields	catchment_size_name
FLOWTYPE	Water flow type	Predefined list	Closed – no inflow or outflow For full list, see flowtypeid-fields	flow_type_name
BIOME	Surrounding biome type	Predefined list	For full list, see global-biomeid-fields	biome_type_name
CORE	Name of collection unit, such as a core or excavation square	Free text	BWIL2	core_name
X_WGS84	Longitude	Numeric (in decimal degrees)	141.998466	longitude
Y_WGS84 ELEVATION	Latitude Elevation above sea level	Numeric (in decimal degrees) Numeric (in metres)	-12.656479 28	latitude elevation
CORDS_ELEV	Source of coordinates and elevation data	Predefined list	INTP_INTP	NA
WATERDEPTH	Water depth at time of sampling	Numeric (in metres)	3.5	water_depth
COREDATE	Sampling date	Date (dd/mm/yyyy)	01/07/2017	coring_date
CORETYPE	Method used to collect the sample	Predefined list	Piston corer	core_type
DEPOS_TYPE	Depositional context type	Predefined list	Alluvial sediment	depo_context

NA: not available.

the Global Paleofire Database (GPD) and structured to comply with the requirements of the OCTOPUS platform. Data in SahulCHAR, according to the OCTOPUS database model, are captured at SITE, UNIT, SAMPLE, and OBSERVATION levels, with not all UNIT types being cores (e.g. archaeological excavations, sediment monoliths). This is shown in Fig. 1, where the shared database structure is kept clearly visible, while the database structure dedicated to other compilations is shadowed out. Following the protocol used by the GPD, unnamed cores are assigned a CORE name consisting of their associated site name and the suffix "\_core1", "\_core2", etc.

# 2.1 Site-level metadata

Site-level metadata fields, descriptions, and examples are presented in Table 1; for the complete documentation. including available options for predefined lists, https://octopussee db.github.io/documentation/data\_tables.html#globalgeoreferencing-tables (last access: 16 March 2025) and https://octopusdb.github.io/documentation/data tables.html#non-

cosmogenics-tables (last access: 16 March 2025). Location data are captured in two forms: metasites and sites. Metasites are area-based (such as a lake) and are stored as polygons, while sites are point-based (such as a specific coring location in a lake) and are stored as coordinates in decimal degrees. Metasites may have multiple associated sites.

Basin and catchment metadata in SahulCHAR (BASIN and CATCHMENT) have been limited to broad categories that do not require numeric values as these data are often not known. Vegetation metadata were limited to broad categories for the major biome surrounding the site (BIOME) as multiple vegetation fields would require extensive list options to be comprehensive. The available options for predefined lists were based on the options available in the GPD, with

2685

Metadata field	Description	Field type	Example or available list	Corresponding GPD field
CORE	Name of collection unit, such as a core or excavation square	Free text	BWIL2	core_name
OBSID1	Unique identifier for observation	Text	BWIL2_0.05_age	NA
SMPID	Unique identifier for sample	Text	BWIL2_0.05	id_sample
DEPTH THICKNESS	Sample depth (mid-point) in metres Sample thickness in centimetres	Numeric (in metres) Numeric (in centimetres)	0.01 1	depth_value NA
LABID	Laboratory ID code for age	Free text	OZX-211	laboratory number
AGE	Age value	Numeric	760	age_value
AGE_ERROR	Age error value	Numeric	20	NA
AGE_UNIT	Measurement unit for age and age error	Predefined list	Radiocarbon years BP	age_units_type
METHOD	Dating method used to generate age	Predefined list	Radiocarbon dating	age_type_name
MATERIAL	Material dated	Predefined list	Bulk sediment, peat	Mat_dated_type
REFDBID1, REFDBID2, REFDBID3	A unique identifier for associated references using the surname of the first author, year of publication, and a keyword (Name:YEARkeyword)	Text	Rehn:2020thesis, Rehn:2021cape	NA

**Table 2.** Table representation of age metadata collected for SahulCHAR attributes vs. corresponding GPD fields, where applicable. For a full description of database tables, refer to Munack et al. (2023) and Munack and Codilean (2023).

NA: not available.

additions where necessary; these changes were informed by author-submitted data.

#### 2.2 Unit- to observation-level metadata

Fields shared across all of the observation-level data are CORE (core or sample name), OBSID1 (internal OCTOPUS identifier, incorporating CORE and identified as "char" or "age"), SMPID (internal OCTOPUS identifier, incorporating CORE and DEPTH), DEPTH, THICKNESS, and references (REFDBID). Observation-level data include ages and charcoal or black carbon records.

## 2.2.1 Age metadata

Age metadata collected in SahulCHAR are presented in Table 2; for the complete documentation, including available options for predefined lists, see https://octopusdb.github.io/documentation/data\_tables.html#sahulchar-

tables (last access: 16 March 2025). The predefined list options are based on the options available in the GPD, with the exception of the METHOD field, which uses an existing OCTOPUS parent table (see https://octopus-db.github.io/documentation/parent\_tables.html#cabah-

methodid-fields, last access: 16 March 2025) to allow for a larger range of options. In line with existing OCTOPUS collections of radiometric ages (such as SahulArch; Saktura et al., 2023), during data entry for version 1, preference was given to uncalibrated rather than calibrated radiocarbon ages where possible to allow for recalibration with future calibration curve updates. Ages reported in calendar years BC/AD or BCE/CE were converted to years BP prior to entry or were entered as AGE\_UNIT = "other" if conversion was not possible. Ages generated from dating methods that are measured as years prior to sample collection and that do not require calibration, such as lead-210 or optically stimulated luminescence, were converted to years BP prior to entry, where possible, or were entered as AGE\_UNIT = "other".

## 2.2.2 Charcoal and black carbon metadata

Charcoal and black carbon metadata collected in SahulCHAR are presented in Table 3; for the documentation, complete including available oplists, tions predefined https://octopusfor see db.github.io/documentation/data\_tables.html#sahulchartables (last access: 16 March 2025). Charcoal and black carbon (hereafter referred to collectively as "char") observations may share the same SMPID as age observations if they are taken from the same depth. Predefined lists are based on the options available in the GPD, except for the

Metadata field	Description	Field type	Example or available list	Corresponding GPD field
CORE	Name of collection unit, such as a core or excavation square	Free text	BWIL2	core_name
OBSID1	Unique identifier for observation	Text	BWIL2_char1_1	
SMPID	Unique sample identifier	Text	BWIL2_0.05	id_sample
DEPTH	Sample depth (mid-point) in metres	Numeric (in metres)	0.01	depth_value
THICKNESS	Sample thickness in centimetres	Numeric (in centimetres)	1	NA
EST_AGE	Estimated age for sample from original publications	Numeric (in years BP)	350	est_age_cal_bp
CALCURVE	Calibration curve used to generate estimated sample age	Predefined list	SHCal20	calibration_curve_version
CALPROGRAM	Calibration programme used to gener- ate estimated sample age	Predefined list	rbacon 2.3.2	calibration_method_type
CHARVALUES	Charcoal or black carbon values	Numeric	0.52	quantity
CHARMETHOD	Preparation method used for charcoal or black carbon analysis	Predefined list	pollen slide For full list, see <i>CHAR method-fields</i>	charcoal_method_name
CHARMEASURE	Measurement units for charcoal or black carbon counts	Predefined list	fragments cm <sup>3</sup> For full list, see <i>Unit</i> <i>measures -fields</i>	charcoal_units_name
CHARMAX	Maximum size for charcoal or black carbon	Numeric	250	NA (included in "charcoal_units_name")
CHARMIN	Minimum size for charcoal or black carbon	Numeric	125	NA (included in "charcoal_units_name")
CHARSIZE_U	Size units for maximum and minimum sizes	Predefined list	Mm For full list, see <i>Unit</i> <i>measures -fields</i>	NA (included in "charcoal_units_name")
CHARSOURCE	Source of charcoal or black carbon data (in field CHARVALUES)	Predefined list	Author For full list, see Data source -fields	data_source_desc
REFDBID1, REFDBID2, REFDBID3	A unique identifier for associated references using the surname of the first author, year of publication, and a keyword (Name:YEARkeyword)	Text	Rehn:2020thesis, Rehn:2021cape	NA

**Table 3.** Table representation of charcoal and black carbon metadata collected for SahulCHAR attributes vs. corresponding GPD fields, where applicable. For a full description of database tables, refer to Munack et al. (2023) and Munack and Codilean (2023).

NA: not available.

2686

CALCURVE and CALPROGRAM fields as the closest corresponding fields in the GPD (calibration\_curve\_version and calibration\_method\_type, respectively) appear as blank drop-downs in the GPD data upload interface and contain no values in terms of data exports.

The structure of SahulCHAR differs from the GPD in terms of its approach to char sizes and measurement units. SahulCHAR specifies the method of charcoal quantification (CHARMETHOD) for each record, with 11 methodologies for measuring charcoal and black carbon particles included (see https://octopusdb.github.io/documentation/parent\_tables.html#cabahcharmethodid-fields, last access: 17 March 2025). Char particle sizes in the GPD are embedded within the field for measurement units (charcoal\_units\_name), resulting in a lengthy (176 options) but incomplete list of available units\_To address this limitation\_char sizes in SabulCHAR

units. To address this limitation, char sizes in SahulCHAR are distinct from measurement units (CHARMEASURE) and are entered separately as maximum (CHARMAX) and minimum values (CHARMIN), along with the measurement



**Figure 2.** Sites with charcoal or black carbon records contained in SahulCHAR version 1, with labels identifying major islands and their national affiliation in square brackets. Sites are coloured by the basal age (ka) of each record. Nation abbreviations: AUS denotes Australia, IDN denotes Indonesia, NZ denotes Aotearoa/New Zealand, and PNG denotes Papua New Guinea.

unit for these size values (CHARSIZE\_U). This allows for a restricted (35 options) yet comprehensive list of measurement units that can be paired with any combination of size values, which may then be merged into a single field during data migration to the GPD. This database structure also allows users to easily separate records by size values for analysis.

While the CHARMEASURE field allows for a wide range of measurement units, volumetric (e.g. fragments per cubic centimetre) rather than influx (e.g. fragments per square centimetre per year) measurements for char were preferred, where possible, during data compilation to allow for recalibration and recalculation of age-depth models when necessary.

SahulCHAR contains char data from the following sources: original data contributed directly by authors; original data, digitized data, and data of unknown origins contained in the GPD; original data contained in non-GPD databases (Neotoma and PANGAEA); original data available in published supplementary materials; and records digitized from published diagrams. Original char data from any source are classified in SahulCHAR as CHARSOURCE = "author". Data sourced from another database, either from digitization data or of unknown origins, are classified as "palaeofire database", and data digitized from published diagrams for SahulCHAR are classified as "digitized". Data were manually digitized for SahulCHAR using WebPlotDigitizer (Rohatgi, 2022). Char data from the GPD were exported via the web interface on 27 February 2023. Char data were

last accessed from PANGAEA on 25 May 2023 and from Neotoma on 11 July 2023. While the RPD contains Australasian char data, these records were derived from the GPD. The methodology applied here involved assessing individual records from the GPD and modifying, updating, or correcting records where necessary based on local knowledge or discussion with the original authors (Table S1). Therefore, even though the RPD includes alternate chronologies and other modifications from the GPD, this was not used for the SahulCHAR data compilation. Due to constantly evolving and updated chronological modelling and calibration techniques, we have not included new chronologies for individual records. Original chronologies produced by the original authors are included, yet it is recommended that recalibration of age-depth models be conducted using the most appropriate and up-to-date methods for records included in SahulCHAR at the time of use.

#### 3 Data summary

SahulCHAR version 1 (V1; Rehn et al. 2024) contains 687 charcoal and black carbon (char) records from 531 cores or samples (hereafter referred to as cores) derived from 425 metasite locations across Sahul (Australia, New Guinea, and the Aru Islands Regency) and Aotearoa/New Zealand (Fig. 2). The majority of metasites are from Australia ( $\sim 64\%$ ), followed by Aotearoa/New Zealand ( $\sim 29\%$ ). Metasites show some geographic clustering, particularly in southeastern Australia and the New Guinea Highlands, with large spatial gaps in central, western, and parts of northern Australia. SahulCHAR is hosted on the OCTOPUS platform (https: //octopusdata.org) and can be accessed directly from the web interface (https://octopus-db.github.io/documentation/usage. html{#}web-interface) or accessed via the Web Feature Service (WFS) (https://octopus-db.github.io/documentation/ usage.html{#}web-feature-service). The WFS data can be accessed directly through GIS or R software (see the Supplement for example code).

Original data were contributed directly to SahulCHAR by 23 authors, totalling 141 records. In cases where authorsubmitted data overlapped with records that already exist in the GPD (27 records), preference was given to the authorsubmitted versions. Approximately 33 % (211) of the records in SahulCHAR are derived from or also exist in the GPD, with 85 records being modified in some way (such as with additional or corrected metadata) with reference to authorsubmitted information or source publications (Fig. 3). Approximately 46 % of records in SahulCHAR are digitized from published diagrams.

The options for SITECODE in SahulCHAR include broad types (e.g. "terrestrial", "lacustrine") and broad types with specific subcategories (e.g. "terrestrial, bog" or "terrestrial, fen") stored in a self-referencing table with subcategories



Figure 3. SahulCHAR V1 (a) sources of char records and (b) broad site types.

linked to their next common denominators; for ease of comparison, sites are grouped into broad types in Fig. 2b. Most sites in SahulCHAR are broadly categorized as terrestrial  $(\sim 49\%, 238 \text{ sites})$ , primarily bogs (107 sites), followed by lacustrine ( $\sim 37 \%$ , 182 sites), primarily classified as lacustrine with no subcategories (154 sites). Categories in SITE-CODE are not exclusive and may overlap (e.g. coastal lakes may be classified as coastal or lacustrine), with these classifications being intended to be a general guide. Site characteristics may also change through time; SITECODE was determined based on site characteristics at the time of sample collection. While archaeological sites were included, these were limited to charcoal quantification undertaken as part of palaeoenvironmental analyses to exclude charcoal potentially associated with archaeological features (e.g. hearths). These archaeological sites were further limited to records where associated depth values were available for char measurements; archaeological sites with char data associated with stratigraphic (SU) or excavation (XU) units without specified depths were excluded.

A total of 3271 ages are contained in SahulCHAR V1. The majority ( $\sim$ 77%) of cores have 1–10 associated ages, and 34 units ( $\sim$ 6%) have no available age data. In instances where no ages are available from a unit with associated charcoal data, other dated units from the same site have been included, where possible (five units, from the metasites of Lake George and Blue Lake in Kosciuszko National Park, both in New South Wales, Australia).

Most UNIT entries have one associated char record (418 cores,  $\sim 79\%$ ) and up to a maximum of six associated records (1 core, MAR2 from the metasite Marura) (Fig. 4). The majority ( $\sim 63\%$ , 432 records) of char records in SahulCHAR are derived from pollen slides, followed by sieved samples ( $\sim 32\%$ , 217 records). Pollen slide charcoal also dominated the dataset compiled by Mooney et al. (2012), although sieved charcoal is slightly better repre-

sented in SahulCHAR (~ 80 % compared to ~ 20 %, respectively, in Mooney et al., 2012, p. 18). Approximately 32 % (220 records) of the char records in SahulCHAR are measured in fragments cm<sup>3</sup>, followed by percentage of pollen sum (~ 14%, 95 records) and fragments cm<sup>-2</sup> yr<sup>-1</sup> (~ 14 %, 94 records). Over half (~ 54 %) of the char records specify a size range for particles, with 54 unique size ranges being specified; this demonstrates both the utility of isolating maximum and minimum particle sizes from measurement units to allow for this variability and the diversity of approaches used to create these records. All char records (~ 67 %) contain 50 entries or fewer. The highest number of entries for any char record is 881 (WL15-2\_char1 from Welsby Lagoon).

#### 4 Data availability

The data in this study are openly available at https://doi.org/10.25900/KKDX-XH23 (Rehn et al., 2024) and via the OCTOPUS database at https://octopusdata.org/ (OCTOPUS, 2025; Codilean et al., 2022). Additional information about the SahulCHAR database collection and the data can be accessed at https://doi.org/10.25900/KKDX-XH23 (Rehn et al., 2024).

# 5 Conclusions and future work

SahulCHAR is the most comprehensive and up-to-date palaeofire database for Sahul and Aotearoa/New Zealand (Rehn et al. 2024) and constitutes an overdue step towards improved representation of Australasia in global syntheses. The latter goal will be addressed through upcoming integration with the GPD as part of the planned conversion of the GPD into a constituent database of the Neotoma Paleoecology Database (Dietze and Vannière, 2022). As an update to the last Australasian compilation (Mooney et al.,



**Figure 4.** SahulCHAR V1 char data summaries: (a) number of char records per UNIT with instances of multiple records from a single core representing different char sizes, analysis methods, or measurement units; (b) sample preparation method for char records; (c) measurement units (for a full list of measurement units and associated abbreviations, see https://octopus-db.github.io/documentation/parent\_tables.html#global-varunitid-fields, last access: 17 March 2025); and (d) number of entries (sample depths) per char record.

2011, 2012, which covered a slightly larger geographic area than SahulCHAR), SahulCHAR triples the number of char records available for the region and incorporates data from numerous new studies produced over the last decade. SahulCHAR follows the FAIR (findability, accessibility, interoperability, and reusability) principles of scientific data management and stewardship (Wilkinson et al., 2016) and the OPEN data requirements of funding agencies, such as the Australian Research Council, to make publicly funded data freely available.

Data creators in the region are encouraged to contribute records either directly to SahulCHAR or to the GPD within Neotoma. Future versions will ideally shift the balance of char sources away from digitized data, with a greater representation of author-contributed original data. Future work relating to SahulCHAR version 1 will provide a synthesis and analysis of the records in the dataset to explore trends in palaeofire regimes across the region and could also explore metadata associated with each record to understand changing approaches to charcoal analysis over time.

Data creators with char records from Australia, New Guinea, or Aotearoa/New Zealand that they would like to contribute can use a SahulCHAR data template (https://doi.org/10.5281/zenodo.10117180; Rehn, 2023) and can contact Dr Haidee Cadd (haidee@uow.edu.au) with enquiries or to submit completed data templates.

**Supplement.** The supplement related to this article is available online at https://doi.org/10.5194/essd-17-2681-2025-supplement.

Author contributions. Conceptualization: ER, HC, SM, TC. Methodology: ER, HC, SM, HM, AC. Data contribution: ER, HC, SM, MA, KB, MC, CG, JH, PJ, APK, LM, MM, MM, KM, DM, KM, PM, NP, CR, JS, JT, JW. Data curation: ER, HM. Visualiza2690

tion: ER, HC. Formal analysis: ER. Writing (original draft): ER, HC, SM, HM. Writing (review and editing): ER, HC, SM, TC, HM, AC, with contributions and approval from all authors. Project administration and supervision: HC. Funding acquisition: ER, HC, SM, TC, AC.

**Competing interests.** The contact author has declared that none of the authors has any competing interests.

**Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. We thank the ARC Centre of Excellence for Australian Biodiversity and Heritage for financial support through their Future Leaders programme. We thank all researchers who contributed data to the SahulCHAR database.

**Financial support.** This research has been supported by the Australian Research Council (grant nos. CE170100015, DP150103875, and DP190102782).

**Review statement.** This paper was edited by Jia Yang and reviewed by Patrick Bartlein and one anonymous referee.

## References

- Adeleye, M. A., Haberle, S. G., and Bowman, D. M. J. S.: Long-term stability of temperate Australian wet forestmoorland mosaics despite recurrent fires associated with late Holocene climate change, Landsc. Ecol., 38, 2675–2685, https://doi.org/10.1007/s10980-023-01738-3, 2023.
- Codilean, A. T., Munack, H., Saktura, W. M., Cohen, T. J., Jacobs, Z., Ulm, S., Hesse, P. P., Heyman, J., Peters, K. J., Williams, A. N., Saktura, R. B. K., Rui, X., Chishiro-Dennelly, K., and Panta, A.: OCTOPUS database (v.2), Earth Syst. Sci. Data, 14, 3695– 3713, https://doi.org/10.5194/essd-14-3695-2022, 2022.
- Constantine IV, M., Williams, A. N., Francke, A., Cadd, H., Forbes, M., Cohen, T., Zhu, X., and Mooney, S. D.: Exploration of the burning question: A long history of fire in eastern Australia with and without people, Fire, 6, 152, https://doi.org/10.3390/fire6040152, 2023.
- Daniau, A.-L., Bartlein, P. J., Harrison, S. P., Prentice, I. C., Brewer, S., Friedlingstein, P., Harrison-Prentice, T. I., Inoue, J., Izumi, K., Marlon, J. R., Mooney, S., Power, M. J., Stevenson, J., Tinner, W., Andrič, M., Atanassova, J., Behling, H., Black, M., Blarquez, O., Brown, K. J., Carcaillet, C., Colhoun, E. A., Colombaroli, D., Davis, B. A. S., D'Costa, D., Dodson, J., Dupont, L., Eshetu, Z., Gavin, D. G., Genries, A., Haberle, S., Hallett, D. J., Hope, G., Horn, S. P., Kassa, T. G., Katamura, F., Kennedy,

L. M., Kershaw, P., Krivonogov, S., Long, C., Magri, D., Marinova, E., McKenzie, G. M., Moreno, P. I., Moss, P., Neumann, F. H., Norström, E., Paitre, c., Rius, D., Roberts, N., Robinson, G. S., Sasaki, N., Scott, L., Takahara, H., Terwilliger, V., Thevenon, F., Turner, R., Valsecchi, V. G., Vannière, B., Walsh, M., Williams, N., and Zhang, Y.: Predictability of biomass burning in response to climate changes, Global Biogeochem. Cy., 26, GB4007, https://doi.org/10.1029/2011GB004249, 2012.

- Dietze, E. and Vannière, B.: The future of the Global Paleofire Database, IPN, https://ipn.paleofire.org/?p=1198 (last access: 31 October 2023), 2022.
- Duane, A., Castellnou, M., and Brotons, L.: Towards a comprehensive look at global drivers of novel extreme wildfire events, Climatic Change, 165, 43, https://doi.org/10.1007/s10584-021-03066-4, 2021.
- Enright, N. J. and Thomas, I.: Pre-European fire regimes in Australian ecosystems, Geogr. Compass, 2, 979–1011, https://doi.org/10.1111/j.1749-8198.2008.00126.x, 2008.
- Feldner, J., Möller, L., Schindler, U., Huber, R., Schumacher, S., Koppe, R., Diepenbroek, M., and Glöckner, F. O.: PANGAEA – Data publisher for Earth & environmental science, Sci. Data, 10, 347, https://doi.org/10.1038/s41597-023-02269-x, 2023.
- Fletcher, M.-S., Hall, T., and Alexandra, A. N.: The loss of an indigenous constructed landscape following British invasion of Australia: An insight into the deep human imprint on the Australian landscape, Ambio, 50, 138–149, https://doi.org/10.1007/s13280-020-01339-3, 2021.
- International Paleofire Network: Global Paleofire Database (GPD), https://paleofire.org/ (last access: 27 February 2023), 2023.
- Haberle, S. G., Hope, G. S., and van der Kaars, S.: Biomass burning in Indonesia and Papua New Guinea: Natural and human induced fire events in the fossil record, Palaeogeogr. Palaeocl., 171, 259– 268, https://doi.org/10.1016/S0031-0182(01)00248-6, 2001.
- Hanson, J. M., VanderGragt, M. L., Welsh, K. J., and Moss, P. T.: Variations in wetland conditions within the Fitzroy Basin, northeastern Australia: A palaeoecological approach, Mar. Freshwater Res., 73, 35–47, https://doi.org/10.1071/MF21082, 2022.
- Harrison, S. P., Villegas-Diaz, R., Cruz-Silva, E., Gallagher, D., Kesner, D., Lincoln, P., Shen, Y., Sweeney, L., Colombaroli, D., Ali, A., Barhoumi, C., Bergeron, Y., Blyakharchuk, T., Bobek, P., Bradshaw, R., Clear, J. L., Czerwiński, S., Daniau, A.-L., Dodson, J., Edwards, K. J., Edwards, M. E., Feurdean, A., Foster, D., Gajewski, K., Gałka, M., Garneau, M., Giesecke, T., Gil Romera, G., Girardin, M. P., Hoefer, D., Huang, K., Inoue, J., Jamrichová, E., Jasiunas, N., Jiang, W., Jiménez-Moreno, G., Karpińska-Kołaczek, M., Kołaczek, P., Kuosmanen, N., Lamentowicz, M., Lavoie, M., Li, F., Li, J., Lisitsyna, O., López-Sáez, J. A., Luelmo-Lautenschlaeger, R., Magnan, G., Magyari, E. K., Maksims, A., Marcisz, K., Marinova, E., Marlon, J., Mensing, S., Miroslaw-Grabowska, J., Oswald, W., Pérez-Díaz, S., Pérez-Obiol, R., Piilo, S., Poska, A., Qin, X., Remy, C. C., Richard, P. J. H., Salonen, S., Sasaki, N., Schneider, H., Shotyk, W., Stancikaite, M., Šteinberga, D., Stivrins, N., Takahara, H., Tan, Z., Trasune, L., Umbanhowar, C. E., Väliranta, M., Vassiljev, J., Xiao, X., Xu, Q., Xu, X., Zawisza, E., Zhao, Y., Zhou, Z., and Paillard, J.: The Reading Palaeofire Database: an expanded global resource to document changes in fire regimes from sedimentary charcoal records, Earth Syst. Sci. Data, 14, 1109-1124, https://doi.org/10.5194/essd-14-1109-2022, 2022.

- Karp, A. T., Fath, J. T., Maron, J. R., and Staver, A. C.: Global response of fire activity to late Quaternary grazer extinctions, Science, 374, 1145–1148, https://doi.org/10.1126/science.abj1580, 2021.
- Laming, A., Fletcher, M.-S., Romano, A., Mullett, R. (on behalf of Gunaikurnai Land and Waters Corporation), Connor, S., Mariani, M., Maezumi, S. Y., and Gadd, P. S.: The curse of conservation: Empirical evidence demonstrating that changes in land-use legislation drove catastrophic bushfires in Southeast Australia, Fire, 5, 175, https://doi.org/10.3390/fire5060175, 2022.
- Lawrence, J., Mackey, B., Chiew, F., Costello, M. J., Hennessy, K., Lansbury, N., Nidumolu, U. B., Pecl, G., Rickards, L., Tapper, N., Woodward, A., and Wreford, A.: Australasia, in: Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Pörtner, H.-O., Roberts, D. C., Tignor, M., Poloczanska, E. S., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A., and Rama, B., Cambridge University Press, Cambridge, UK and New York, NY, USA, https://doi.org/10.1017/9781009325844.013, 1581–1688, 2022.
- Lynch, A. H., Beringer, J., Kershaw, P., Marshall, A., Mooney, S., Tapper, N., Turney, C., and van der Kaars, S.: Using the paleorecord to evaluate climate and fire interactions in Australia, Annu. Rev. Earth Pl. Sc., 35, 215–239, https://doi.org/10.1146/annurev.earth.35.092006.145055, 2007.
- Mariani, M., Connor, S. E., Theuerkauf, M., Herbert, A., Kuneš, P., Bowman, D., Fletcher, M.-S., Head, L., Kershaw, A. P., Haberle, S. G., Stevenson, J., Adeleye, M., Cadd, H., Hopf, F., and Briles, C.: Disruption of cultural burning promotes shrub encroachment and unprecedented wildfires, Front. Ecol. Environ., 20, 292–300, https://doi.org/10.1002/fee.2395, 2022.
- Mariani, M., Wills, A., Herbert, A., Adeleye, M., Florin, A., Cadd, H., Connor, S., Kershaw, P., Theuerkauf, M., Stevenson, J., Fletcher, M. S., Mooney, S., Bowman, D., and Haberle, S.: Shrub cover declined as Indigenous populations expanded across southeast Australia, Science, 386, 567–573, https://doi.org/10.1126/science.adn8668, 2024.
- Marlon, J. R.: What the past can say about the present and future of fire, Quaternary Res., 96, 66–87, https://doi.org/10.1017/qua.2020.48, 2020.
- Marlon, J. R., Bartlein, P. J., Daniau, A.-L., Harrison, S. P., Maezumi, S. Y., Power, M. J., Tinner, W., and Vannière, B.: Global biomass burning: A synthesis and review of Holocene paleofire records and their controls, Quaternary Sci. Rev., 65, 5–25, https://doi.org/10.1016/j.quascirev.2012.11.029, 2013.
- Marlon, J. R., Kelly, R., Daniau, A.-L., Vannière, B., Power, M. J., Bartlein, P., Higuera, P., Blarquez, O., Brewer, S., Brücher, T., Feurdean, A., Romera, G. G., Iglesias, V., Maezumi, S. Y., Magi, B., Courtney Mustaphi, C. J., and Zhihai, T.: Reconstructions of biomass burning from sediment-charcoal records to improve data–model comparisons, Biogeosciences, 13, 3225–3244, https://doi.org/10.5194/bg-13-3225-2016, 2016.
- McWethy, D. B., Whitlock, C., Wilmshurst, J. M., McGlone, M. S., Fromont, M., Li, X., Dieffenbacher-Krall, A., Hobbs, W. O., Fritz, S. C. ad Cook, E. R.: Rapid landscape transformation in South Island, New Zealand, following initial Polynesian settlement, P. Natl. Acad. Sci. USA, 107, 21343–21348, https://doi.org/10.1073/pnas.1011801107, 2010.

- Mooney, S. D. and Tinner, W.: The analysis of charcoal in peat and organic sediments, Mires Peat, 7, 1–18, 2010.
- Mooney, S. D., Harrison, S. P., Bartlein, P. J., Daniau, A.-L., Stevenson, J., Brownlie, K. C., Buckman, S., Cupper, M., Luly, J., Black, M., Colhoun, E., D'Costa, D., Dodson, J., Haberle, S., Hope, G. S., Kershaw, P., Kenyon, C., McKenzie, M., and Williams, N.: Late Quaternary fire regimes of Australasia, Quaternary Sci. Rev., 30, 28–46, https://doi.org/10.1016/j.quascirev.2010.10.010, 2011.
- Mooney, S. D., Harrison, S. P., Bartlien, P. J., and Stevenson, J.: The prehistory of fire in Australasia, in: Flammable Australia: Fire Regimes, Biodiversity and Ecosystems in a Changing World, edited by: Bradstock, R. A., Gill, A. M., and Williams, R. J., CSIRO Publishing, 3–25, ISBN 9780643104822, 2012.
- Munack, H. and Codilean, A. T.: OCTOPUS database schema (online), https://octopus-db.github.io/schema (last access: 16 July 2024), 2023.
- Munack, H., Rehn, E., Saktura, W. M., and Codilean, A. T.: OCTO-PUS database documentation (online), https://octopus-db.github. io/documentation (last access: 16 July 2024), 2023.
- Nolan, R. H., Bowman, D. M. J. S., Clarke, H., Haynes, K., Ooi, M. K. J., Price, O. F., Williamson, G. J., Whittaker, J., Bedward, M., Boer, M. M., Cavanagh, V. I., Collins, L., Gibson, R. K., Griebel, A., Jenkins, M. E., Keith, D. A., Mcilwee, A. P., Penman, T. D., Smson, S. A., Tozer, M. G., and Bradstock, R. A.: What do the Australian Black Summer fires signify for the global fire crisis?, Fire, 4, 97, https://doi.org/10.3390/fire4040097, 2021.
- OCTOPUS: OCTOPUS database v.2.3, https://octopusdata.org/ (last access: 13 June 2025), 2025.
- Patton, N. R., Shulmeister, J., Hua, Q., Almond, P., Rittenour, T. M., Hanson, J. M., Grealy, A., Gilroy, J., and Ellerton, D.: Reconstructing Holocene fire records using dune footslope deposits at the Cooloola Sand Mass, Australia, Quaternary Res., 115, 67– 89, https://doi.org/10.1017/qua.2023.14, 2023.
- Perry, G. L. W., Wilmshurst, J. M., and McGlone, M. S.: Ecology and long-term history of fire in New Zealand, New Zeal. J. Ecol., 38, 157–176, 2014.
- Power, M. J., Marlon, J. R., Bartein, P. J., and Harrison, S. P.: Fire history and the Global Charcoal Database: A new tool for hypothesis testing and data exploration, Palaeogeogr. Palaeocl., 291, 52–59, https://doi.org/10.1016/j.palaeo.2009.09.014, 2010.
- Rehn, E.: SahulCHAR Data Template for Authors (Version 1), Zenodo [data set], https://doi.org/10.5281/zenodo.10117180, 2023.
- Rehn E., Cadd H., Mooney S., Cohen T. J., Munack H., Codilean A. T., Adeleye, M., Beck, K. K., Constantine IV, M., Gouramanis, C., Hanson, J. M., Jones, P. J., Kershaw, A. P., Mackenzie, L., Maisie, M., Mariani, M., Matley, K., McWethy, D., Mills, K., Moss, P., Patton, N. R., Rowe, C., Stevenson, J., Tibby, J., and Wilmshurst, J.: SahulChar: An open database of sedimentary charcoal and black carbon records from Australia, New Guinea, and New Zealand, University of Wollongong [data set], https://doi.org/10.25900/KKDX-XH23, 2024.
- Rohatgi, A.: WebPlotDigitizer, https://automeris.io/ WebPlotDigitizer/ (last access: 16 July 2024), 2022.
- Rowe, C., Rehn, E., Brand, M., Hutley, L. B., Comley, R., Levchenko, V., Zwart, C., Wurster, C., and Bird, M. I.: Holocene climate-fire-vegetation feedbacks in tropical savannas: Insights from the Marura sinkhole, East Arnhem Land, northern Aus-

tralia, J. Veg. Sci., 33, e13158, https://doi.org/10.1111/jvs.13158, 2022.

- Rowe, C., Stevenson, J., Connor, S., and Adeleye, M.: Fire and the transformation of landscapes, in: The Oxford Handbook of the Archaeology of Indigenous Australia and New Guinea, edited by: McNiven, I. J. and David, B., Oxford University Press, 381– 411, ISBN 13:9780190095611, 2023.
- Saktura, W. M., Rehn, E., Linnenlucke, L., Munack, H., Wood, R., Petchey, F., Codilean, A. T., Jacobs, Z., Cohen, T. J., Williams, A. N., and Ulm, S.: SahulArch: A geochronological database for the archaeology of Sahul, Aust. Archaeol., 89, 1– 13, https://doi.org/10.1080/03122417.2022.2159751, 2023.
- Shlisky, A., Waugh, J., Gonzalez, P., Gonzalez, M., Manta, M., Santoso, H., Alvarado, E., Ainuddin Nuruddin, A., Rodriguez-Trejo, D. A., Swaty, R., Schmidt, D., Kaufmann, M., Myers, R., Alencar, A., Kearns, F., Johnson, D., Smith, J., and Zollner, D.: Fire, ecosystems and people: Threats and strategies for global biodiversity conservation, GFI Technical Report 2007-2, The Nature Conservancy, https://www.conservationgateway.org/ Files/Documents/fire\_ecosystems\_and\_people.pdf (last access: 10 September 2024), 2007.
- Thomas, Z., Mooney, S., Cadd, H., Baker, A., Turney, C., Schneider, L., Hogg, A., Haberle, S., Green, K., Weyrich, L. S., Pérez, V., Moore, N. E., Zawadzki, A., Kelloway, S. J., and Khan, S. J.: Late Holocene climate anomaly concurrent with fire activity and ecosystem shifts in the eastern Australian Highlands, Sci. Total Environ., 802, 149542, https://doi.org/10.1016/j.scitotenv.2021.149542, 2022.
- Turner, R., Kelly, A., and Roberts, N.: A critical assessment and experimental comparison of microscopic charcoal extraction methods, in: Charcoals from the past: cultural and palaeoenvironmental implications. Proceedings of the Third International Meeting of Anthracology, edited by: Fiorentino, G. and Magri, D., Archaeopress, Oxford, 265–272, ISBN 978 1 4073 0294 2, 2004.

- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., 't Hoen, P. A. C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., and Mons, B.: The FAIR Guiding Principles for scientific data management and stewardship, Sci. Data, 3, 160018, https://doi.org/10.1038/sdata.2016.18, 2016.
- Williams, A. N., Mooney, S. D., Sisson, S. A., and Marlon, J.: Exploring the relationship between Aboriginal population indices and fire in Australia over the last 20,000 years, Palaeogeogr. Palaeocl., 432, 49–57, https://doi.org/10.1016/j.palaeo.2015.04.030, 2015.
- Williams, A. P. and Abatzoglou, J. T.: Recent advances and remaining uncertainties in resolving past and future climate effects on global fire activity, Curr. Clim. Change Rep., 2, 1–14, https://doi.org/10.1007/s40641-016-0031-0, 2016.
- Williams, J. W., Grimm, E. C., Blois, J. L., Charles, D. F., Davis, E. B., Goring, S. J., Graham, R. W., Smith, A. J., Anderson, M., Arroyo-Cabrales, J., Ashworth, A. C., Betancour, J. L., Bills, B. W., Booth, R. K., Buckland, P. I., Curry, B. B., Giesecke, T., Jackson, S. T., Latorre, C., Nichols, J., Purdum, T., Roth, R. E., Stryker, M., and Takahara, H.: The Neotoma Paleoecology Database, a multiproxy, international, community-curated data resource, Quaternary Res., 89, 156– 177, https://doi.org/10.1017/qua.2017.105, 2018.
- Williams, N.: Late Quaternary fire regimes of Australasia, Quaternary Sci. Rev., 30, 28–46, https://doi.org/10.1016/j.quascirev.2010.10.010, 2011.