



Supplement of

EURO-SUPREME: sub-daily precipitation extremes in the EURO-CORDEX ensemble

Anouk Dierickx et al.

Correspondence to: Hans Van de Vyver (hvijver@meteo.be)

The copyright of individual parts of the supplement might differ from the article licence.

Table S1. Spatial domain and simulation periods of the RCMs that contributed to EURO-SUPREME. Note that RCM-downscaling from MOHC-HadGEM2-ESM ends in 2099 instead of 2100.

RCM	Spatial dimension	Min/max longitude and latitude	Historical period	Future period
ALADIN63	453 x 453	52° W–73° E and 21° N–74° N	1951–2005 ¹	2006–2100
COSMO-crCLIM	424 x 412	45° W–65° E and 22° N–73° N	1950–2005 ²	2006–2100
HadREM3-GA7-05	424 x 412	45° W–65° E and 22° N–73° N	1952–2005 ³	2006–2100
RCA4	424 x 412	45° W–65° E and 22° N–73° N	1970–2005	2006–2100
RegCM4-6	527 x 527	63° W–82° E and 15° N–76° N	1970–2005	2006–2100
REMO2015	424 x 412	52° W–73° E and 21° N–74° N	1950–2005	2006–2100

¹One run starts from 1950.

²Some runs start in 1949 and 1951.

³One run ends in 2004 instead of 2005.

Table S2. National IDF-statistics for the Netherlands, Denmark and Finland. Shown are the 10-year return levels (mm) for various rainfall durations.

Country	1 h	3 h	6 h	12 h	24 h	Reference
The Netherlands	31.0	39.8	46.0	52.9	62.9	[1]
Denmark	24.9	33.3	40.2	46.7	55.2	[2]
Finland	22.9	27.0	34.0	44.0	53.0	[3, 4]

Table S3. Temporal-areal reduction factors to convert return levels of d -hourly precipitation extremes of EURO-SUPREME (i.e. 1-h temporal resolution, 0.11° spatial resolution) to return levels of continuous time sliding point maxima. This is taken from [5, Table 2].

$d = 1 \text{ h}$	$d = 2 \text{ h}$	$d = 3 \text{ h}$	$d = 6 \text{ h}$	$d = 12 \text{ h}$
1.21	1.09	1.06	1.02	1.01

Table S4. Periods associated with global warming levels of 1.5 °C and 3 °C for the GCM runs in the EURO-CORDEX ensemble [6].

GCM run	+1.5°C period	+3°C period
CNRM-CERFACS-CNRM-CM5-r1i1p1	2015 - 2044	2053 - 2082
ICHEC-EC-EARTH-r12i1p1	2012 - 2041	2052 - 2081
ICHEC-EC-EARTH-r1i1p1 ¹	2012 - 2041	
ICHEC-EC-EARTH-r3i1p1	2014 - 2043	2051 - 2080
MOHC-HadGEM2-ES-r1i1p1	2004 - 2033	2037 - 2066
IPSL-IPSL-CM5A-MR-r1i1p1	2008 - 2037	2040- 2069
MPI-M-MPI-ESM-LR-r1i1p1	2014 - 2043	2053 - 2082
MPI-M-MPI-ESM-LR-r2i1p1	2012 - 2041	2051 - 2080
MPI-M-MPI-ESM-LR-r3i1p1	2012 - 2041	2050 - 2079
NCC-NorESM1-M-r1i1p1	2017 - 2046	2058 - 2087

¹For ICHEC-EC-EARTH-r1i1p1, the 3°C period is missing, resulting in 33 usable model runs




Fig. S1. Elevation map of Belgium (m), together with the locations of the (sub-)hourly precipitation stations.

REFERENCES

1. J. Beersma, R. Versteeg, and H. Hakvoort, "Neerslagstatistieken voor korte duren (precipitation statistics for short durations)," Tech. rep., STOWA (2018).
2. H. Madsen, I. B. Gregersen, D. Rosbjerg, and K. Arnbjerg-Nielsen, "Regional frequency analysis of short duration rainfall extremes using gridded daily rainfall data as co-variate," *Water Sci. Technol.* **75**, 1971–1981 (2017).
3. J. Aaltonen, H. Hohti, K. Jylhä, *et al.*, "Rankkasateet ja taajamatulvat (extreme precipitation and urban flooding)," Tech. rep., RATU (2008).
4. A. Venäläinen, S. Saku, T. Kilpeläinen, *et al.*, "Sään ääriilmiöistä suomessa. (aspects about climate extremes in finland. abstract in english.)," Tech. rep., Ilmatieteen laitos Helsinki, Finland (2007).
5. P. Berg, O. B. Christensen, K. Klehmet, *et al.*, "Summertime precipitation extremes in a EURO-CORDEX 0.11° ensemble at an hourly resolution," *Nat. Hazards Earth Syst. Sci.* **19**, 957–971 (2019).
6. M. Hauser, F. Engelbrecht, and E. M. Fischer, "Transient global warming levels for CMIP5 and CMIP6 (v0.3.0)," Zenodo (2022).