



Annual vegetation maps in the Qinghai–Tibet Plateau (QTP) from 2000 to 2022 based on MODIS series satellite imagery

Guangsheng Zhou¹, Hongrui Ren², Lei Zhang², Xiaomin Lv¹, and Mengzi Zhou¹

¹State Key Laboratory of Severe Weather, Chinese Academy of Meteorological Sciences, Beijing 100081, China

²Department of Geomatics, Taiyuan University of Technology, Taiyuan 030024, China

Correspondence: Guangsheng Zhou (zhougs@cma.gov.cn)

Received: 17 May 2024 – Discussion started: 4 July 2024

Revised: 28 December 2024 – Accepted: 7 January 2025 – Published: 28 February 2025

Abstract. The Qinghai–Tibet Plateau (QTP), known as the Third Pole of the Earth and the “water tower of Asia”, plays a crucial role in global climate regulation, biodiversity conservation, and regional socio-economic development. Continuous annual vegetation types and their geographical distribution data are essential for studying the response and adaptation of vegetation to climate change. However, there are very limited data on vegetation types and their geographical distributions on the QTP due to the harsh natural environment. Currently, land cover and surface vegetation data are typically obtained using traditional classification methods for each period’s product based on remote sensing information. These approaches do not consider the temporal continuity of vegetation presence, leading to a gradual increase in misclassified pixels and uncertainty in their locations, consequently decreasing the interpretability of the long-time-series remote sensing products. To address this issue, this study developed a new method for long-time continuous annual vegetation mapping based on reference vegetation maps and annual updates and mapped the vegetation of the QTP from 2000 to 2022 at a 500 m spatial resolution through the MOD09A1 product. The overall accuracy of continuous annual QTP vegetation mapping from 2000 to 2022 reached 83.27 %, with the reference annual 2020 data reaching an accuracy of 83.32 % and a kappa coefficient of 0.82. This study supports the use of remote sensing data for long-term continuous annual vegetation mapping. The 500 m annual vegetation maps are available at <https://doi.org/10.11888/Terre.tpd.301205> (Zhou et al., 2024).

1 Introduction

Vegetation, an integral component of Earth’s ecosystems, plays an irreplaceable role in maintaining climate stability, preserving biodiversity, and supplying vital resources for humans. Vegetation maps not only facilitate a visual comprehension of vegetation types and their geographic distribution, but also provide essential data for natural resource management and environmental protection (Immerzeel et al., 2010). Particularly, long time series of geographical distribution data on vegetation types are crucial for revealing the impacts of climate change and human activities on vegeta-

tion, elucidating the succession processes of vegetation, and scientifically managing vegetation ecosystems.

The availability of large-scale, long-term, and free remote sensing imagery has significantly advanced the development of land cover and surface vegetation data. Notable international products include the European Space Agency (ESA) GlobCover_2005 and GlobCover_2009 (300 m) (Bontemps et al., 2010) as well as the ESA Climate Change Initiative (ESA-CCI) land cover dataset, which offers improved temporal continuity at a 300 m spatial resolution (Defourny et al., 2017). Additionally, the National Aeronautics and Space Administration (NASA) annual MCD12Q1 product that has existed since 2001 (500 m) (Friedl et al., 2010;

Sulla-Menashe et al., 2019) is widely used. In China, representative products include GlobeLand 30 by Chen et al. (2015), Finer-Resolution Observation and Monitoring of Global Land Cover (FROM_GLC)10 and FROM_GLC30 by Tsinghua University (Gong et al., 2013; Gong et al., 2019), Global 30 m Land-cover Classification with a Fine Classification System (GLC_FCS30) and Global 30 m Land-cover Dynamics Monitoring Dataset (GLC_FCS30D) by the Chinese Academy of Sciences (Zhang et al., 2019c; Zhang et al., 2021b, 2024b), and China land cover dataset (CLCD) by Wuhan University (Yang and Huang, 2021). Leveraging the intelligent remote sensing mapping (iMap) concept and framework, there has been rapid progress in developing global-scale seamless daily data cubes and in creating annual and seasonal land surface maps (Feng and Li, 2020; Liu et al., 2021a).

Traditional long-time-series land cover and surface vegetation datasets are often generated by independently classifying each period's product. However, significant spatial distribution differences between products from different periods hinder their direct comparability (Liu et al., 2021b). For instance, NASA's MCD12Q1 v5 product, produced using MODIS data and a decision tree classifier (Friedl et al., 2010), exhibited substantial instability, with an annual land cover label change rate of approximately 11.4 % between 2001 and 2013. Similarly, GlobCover_2009 and GlobCover_2005 show notable spatial distribution differences, making them unsuitable for change detection studies (Bontempo et al., 2010). To address this issue, post-processing methods have been developed and have demonstrated remarkable effectiveness. For example, the hidden Markov model method (Sulla-Menashe et al., 2019), applied to the MCD12Q1.v6 product, reduced the instability of land cover label changes to 1.6 % compared to MCD12Q1 v5. Another study, using spatial–temporal consistency methods (Yang and Huang, 2021), further improved the mapping accuracy of land cover time series datasets. However, these post-processing methods rely on the establishment of numerous subjective rules, leading to complexity and limitations in their application (Zhang et al., 2024b). In recent years, research on continuous and dense change detection methods has made rapid progress, particularly with the continuous change detection and classification (CCDC) algorithm (Zhu and Woodcock, 2014; Zhu et al., 2019). This technique leverages all available remote sensing data to conduct continuous land cover change detection, providing a new approach to address the aforementioned issues. For instance, Xian et al. (2022) used the CCDC method to produce annual land cover products for the United States from 1985 to 2017, while Zhang et al. (2024b) applied it to generate the global GLC-FCS30D product, which demonstrates greater temporal stability compared to GLC-FCS30. Therefore, there is an urgent need to develop long-time-series mapping methods based on change detection to overcome the limitations of traditional classification and post-processing approaches.

The QTP, known as the “roof of the world” and the water tower of Asia, is critical for global climate regulation and regional socio-economic development (Yao et al., 2012). In the 1970s, China carried out the first extensive scientific survey of the QTP. Currently, the second QTP scientific expedition is underway, with vegetation survey being one of the major components (Zhou et al., 2023). Understanding the evolution of vegetation types on the QTP is important for revealing the effects of climate change on vegetation structure and function. Additionally, such insights are essential for elucidating the carbon and water cycles of the QTP and for formulating high-quality, sustainable development strategies for the region amidst global warming (Wang et al., 2022, 2023; Zhang et al., 2024a).

As the Third Pole of the Earth, the QTP has very limited data on vegetation types and geographical distribution due to its harsh natural environment. Currently, the primary vegetation data for the QTP include a vegetation map of the Qinghai–Tibet Plateau in 2020 with a 10 m spatial resolution (Zhou et al., 2022a), a new vegetation map for the Qinghai–Tibet Plateau by integrated classification from multi-source data products (2020) (Zhang et al., 2021a), a vegetation map of the Qinghai–Tibet Plateau in the 1980s (Zhou et al., 2022b), and a vegetation map of the Qinghai–Tibet Plateau permafrost zone (Wang et al., 2016). These vegetation distribution products are single-period maps, which are insufficient for depicting dynamic changes in vegetation, thus limiting the understanding of vegetation evolution trends and mechanisms on the QTP. In addition, Wang et al. (2022) described the distribution of alpine meadows and alpine steppes on the QTP and predicted the distribution trends of these vegetation types over the next century using precipitation data. Wang et al. (2023) analyzed the characteristics of grassland changes on the QTP over the past 40 years from the perspective of plant community structure. Other studies have focused on wetlands, waterbodies, glaciers, and other aspects of the QTP (Zhang et al., 2019a; Hu et al., 2023; Li et al., 2023). However, these long-time-series products, created with traditional classification methods, are primarily focused on specific types, overlooking the need to identify temporal changes across different types, which limits the accuracy of the products. Therefore, in order to provide continuous data support for research on the interaction between vegetation and climate change, there is an urgent need for long-time-series vegetation mapping.

This study aims to develop a new approach to long-time-series vegetation mapping using remote sensing imagery, and to map vegetation of the entire QTP at 500 m from 2000 to 2022, using the MOD09A1 remote sensing data.

2 Materials and methods

2.1 Study area

The QTP, situated between 25°59′30″ N and 40°1′0″ N, 67°40′37″ E and 104°40′57″ E, has an average elevation of about 4320 m and a total area of 3.08 million km² (Zhang et al., 2021c). This study covers the entire QTP, spanning six provinces and regions within China – Tibet, Qinghai, Gansu, Sichuan, Yunnan, and Xinjiang – as well as areas in India, Pakistan, Tajikistan, Afghanistan, Nepal, Bhutan, Myanmar, and Kyrgyzstan. In particular, the area of the QTP within China is about 2.58 million km² (approximately 83.7 % of the QTP), with an average elevation of about 4400 m (Yu et al., 2014; Zhang, 2019; Zhang et al., 2021d, e). The QTP slopes downward from the high northwest to the lower southeast, with a humid and rainy climate in the southeast and arid conditions in the northwest. The vegetation distribution across the QTP is influenced by topography and climatic conditions, revealing a clear horizontal zonation ranging from forests, shrubs, and meadows in the southeast to grasslands and deserts in the northwest. Additionally, as temperatures decrease with increasing altitude, there is a distinct vertical stratification in vegetation, ranging from forests at lower elevations to alpine meadows and alpine vegetation at higher elevations.

2.2 Data sources

2.2.1 Training sample data for vegetation type

The vegetation of the QTP is primarily categorized into 15 types (Editorial Board of the Vegetation Map of China, 2007; Zhou et al., 2023). Given the significance of glaciers and snow cover, this study expanded the vegetation classification to 16 types for the 500 m spatial resolution mapping (Table 1), including evergreen broad-leaved forest (EBF), evergreen coniferous forest (ECF), coniferous and broad-leaved mixed forest (CBMF), deciduous broad-leaved forest (DBF), deciduous coniferous forest (DCF), scrub (SC), alpine scrub meadow (ASM), alpine meadow (AM), alpine grassland (AG), alpine vegetation (AV), alpine desert (AD), cultivated vegetation (CV), wetland (WE), water (WA), non-vegetated area (NVA), and glacier and snow (GS).

The training vegetation samples for the QTP within China were obtained from the vegetation map of the Qinghai–Tibet Plateau in 2020 with a 10 m spatial resolution (Zhou et al., 2022a). This product was created using a regional vegetation mapping method based on terrain–climate–remote sensing information, with a spatial resolution of 10 m and an overall accuracy of 89.5 %. Its classification system is consistent with that of this study, except for the exclusion of GS, and was therefore used to generate the vegetation type training sample data for this research (Zhou et al., 2023). The vegetation map was resampled to 500 m. Subsequently, the dominant vegetation type and its proportional area, derived from

the corresponding 50 × 50 pixels at a 10 m spatial resolution, were determined as the vegetation type categorization and pixel purity (reflected as the percentage of the dominant vegetation type within these 50 × 50 pixels) for the 500 m spatial resolution map. Concurrently, different purity levels and their area proportions were calculated (Table 2). Random sampling was then conducted in areas with purity levels exceeding 70 %, representing approximately 62.34 % of the QTP. For the QTP regions outside of China, vegetation type sample data were obtained from high-resolution Google Earth imagery from 2020. By visually interpreting these images, samples were selected from areas with consistent vegetation types within a 500 m range. Using these methods, a total of 8937 training samples were collected to train the 2020 vegetation classification model (Table 1, Fig. 1).

2.2.2 Validation sample data for vegetation type

As the Third Pole of the Earth, the QTP has high altitudes and sparse human activity, making it difficult to obtain validation samples. The validation samples in this study consist of three parts: field measurements, visual interpretation, and third-party samples.

The field measurement samples include 173 samples collected during field expeditions on the QTP in 2019, 2020, and 2021. These samples were collected during the vegetation-growing season on the QTP, with neighboring samples spaced approximately 50 km apart. It is ensured that each sample has uniform and consistent vegetation distribution within a 1 km × 1 km area.

The visual interpretation samples (1002 in total) were obtained from previous work by Wu et al. (2024) using Google Earth and Google Earth Engine (GEE). The selected samples are from areas with consistent vegetation distribution from 1990 to 2020, and each sample covers an area greater than 500 m. Each sample was independently interpreted by three interpreters using long-time-series Landsat remote sensing images. Conflicting results were resolved through a second interpretation, and inconsistent samples were removed.

The third-party samples include the first all-season sample set (Li et al., 2017), global land cover validation samples (Liu et al., 2019), and validation samples gathered through literature search. The first all-season sample set consists of approximately 140 000 validation samples worldwide, obtained using Landsat 8 data from 2013–2015. The second set includes 44 043 validation samples, primarily from around 2015, and was published as a validation dataset for the GLC-FCS30-2015 product. The literature search samples were obtained from the China National Knowledge Infrastructure (<https://www.cnki.net/>, last access: 19 March 2023) and Web of Science (<https://www.webofscience.com/>, last access: 19 March 2023) databases using keywords such as Qinghai–Tibet Plateau, vegetation, and vegetation cover to collect published validation samples from 2000 to 2022.

Table 1. Vegetation classification system and training and validation sample counts on the QTP.

Types	Definitions	Training samples	Validation samples
Evergreen broad-leaved forest (EBF)	Forest communities composed of evergreen broad-leaved tree species	936	76
Evergreen coniferous forest (ECF)	Forest communities composed of evergreen coniferous tree species	768	102
Coniferous and broad-leaved mixed forest (CBMF)	Forest communities composed of both coniferous and broad-leaved tree species	475	25
Deciduous broad-leaved forest (DBF)	Forest communities composed of broad-leaved tree species that shed leaves in winter and grow in summer	570	62
Deciduous coniferous forest (DCF)	Forest communities composed of coniferous tree species that shed leaves in winter and grow in summer	370	91
Scrub (SC)	Vegetation communities dominated by shrubs	397	79
Alpine scrub meadow (ASM)	Vegetation communities composed of alpine shrubs and alpine meadows, adapted to cold, windy, dry, and alpine climates	213	49
Alpine meadow (AM)	Vegetation communities primarily consisting of herbaceous plants suited to cold climates	1375	310
Alpine grassland (AG)	Vegetation communities composed of alpine grasslands with low biomass and a short growing season	872	217
Alpine vegetation (AV)	Vegetation communities occurring above the treeline or shrub belt and below the permanent snow line, dominated by ice- and cold-tolerant plants	503	97
Alpine desert (AD)	Deserts composed of cold- and drought-tolerant cushion subshrubs	499	325
Cultivated vegetation (CV)	Vegetation communities formed through human cultivation	469	109
Wetland (WE)	Vegetation communities composed of water, bare soil, and herbaceous or woody plants	246	81
Water (WA)	Areas covered by water year-round	377	91
Non-vegetated area (NVA)	Areas composed of natural soil, sand, or rock	204	36
Glacier and snow (GS)	Areas covered by ice and snow year-round	663	439

Table 2. Purity and area proportion of a 500 m resolution pixel samples of vegetation types on the QTP within China.

Purity (%)	0–10	10–20	20–30	30–40	40–50	50–60	60–70	70–80	80–90	90–100
Area proportion (%)	0.02	0.04	1.01	4.20	8.28	12.55	11.56	11.40	12.55	38.39

The third-party validation samples in this study carry certain uncertainties, such as spatial range and homogeneity. According to Li et al. (2017), the first all-season sample set includes samples with varying unit sizes (30, 100, 250, 500, and 1000 m), with those larger than 500 m being rare (only 10.6%). Samples smaller than 500 m are not suitable for validating 500 m vegetation maps. Additionally, the global land

cover validation samples from Liu et al. (2019) and those from literature searches only guarantee accuracy in sampling time and type but not spatial extent. To address these issues, this study applied the method of Feng et al. (2012) to extract spatially homogeneous regions from MOD09A1 data by analyzing the range of values in a 3×3 grid of 500 m pixels. Thresholds for homogeneity were set for each band: 0.03 for




Figure 1. Spatial distribution of training (2020) and validation (2000–2022) samples for vegetation mapping at a 500 m resolution on the QTP.

blue, green, and red; 0.06 for near-infrared; and 0.03 for both shortwave infrared 1 and shortwave infrared 2. If the range of values across all bands fell within these thresholds, the area was considered homogeneous. GS areas on the QTP are mostly classified as heterogeneous areas. Therefore, the samples labeled as GS in the third-party samples are visually interpreted to remove unreasonable ones. After applying this quality control, 1014 third-party validation samples were obtained, including 327 from the global all-season sample library, 499 from the GLC-FCS30–2015 global validation set, and 188 from literature searches.

The validation sample dataset for this product contains 2189 samples from field measurements, visual interpretation, and third-party samples (Table 1, Fig. 1). Among them, 173 field measurement samples are available for validating the 2020 product. There are 1002 samples from visual interpretation, characterized by stable attributes over multiple years, making them suitable for validating annual products from 2000 to 2020. Furthermore, 1014 third-party validation samples can be used to validate annual products from 2000 to 2019.

2.2.3 Remote sensing data

Remote sensing data in this study were sourced from the Moderate Resolution Imaging Spectroradiometer (MODIS) series satellite imagery provided by NASA. As one of the longest-operating Earth observation instruments currently in orbit, MODIS is a multi-spectral observation sensor launched in May 1999 that enables efficient and comprehensive observation of the Earth. The MOD09A1 surface reflectance dataset, obtained through the GEE platform (Gorelick et al.,

2017), is a Level 3 product generated from daily surface reflectance observations using an 8 d compositing algorithm. This algorithm selects the best observation during the 8 d period based on criteria such as minimal cloud cover and highest observation quality (Vermote et al., 2015). The dataset has undergone standardized pre-processing provided by NASA, including radiometric calibration, atmospheric correction, and cloud masking based on quality assurance (QA) flags (Vermote et al., 2015; Vermote, 2021). To ensure the data were suitable for this study, additional processing steps were applied. The original sinusoidal projection of MOD09A1 was converted to the WGS 84 geographic coordinate grid to maintain spatial consistency with other datasets, and the data were spatially subsetted to the QTP region to match the study area. The MOD09A1 dataset provides surface reflectance in seven spectral bands (Table 2) with a spatial resolution of 500 m, and 1051 MOD09A1 images from 1 January 2000 to 31 December 2022 were used to generate annual vegetation maps at a 500 m resolution for the QTP. These pre-processing steps ensure the reliability and quality of the data for vegetation classification and long-term analysis.

2.2.4 Climate and terrain data

Temperature and precipitation are key driving factors for the distribution and dynamic changes in vegetation types on the QTP, and the incorporation of climate data significantly enhances the understanding of vegetation distribution on the QTP (Wang et al., 2022; Zhou et al., 2023; Zhang et al., 2024a). This study utilized climate data, which included annual precipitation (AP) and annual average temperature (AT) across the entire QTP from 2000 to 2022. For

the QTP within China, climate data at 1000 m were obtained from the National Tibetan Plateau Data Center, specifically the 1 km monthly precipitation dataset for China (1901–2023) and the 1 km monthly mean temperature dataset for China (1901–2023) (Peng et al., 2017a, b, 2019; Peng, 2019; Ding and Peng, 2020; Peng, 2020). To derive the precipitation and temperature features required for this study, the monthly precipitation data were summed to calculate annual totals, while the monthly mean temperature data were averaged over the year. In contrast, climate data for areas of the QTP outside China were derived from the Climatic Research Unit (CRU) high-resolution gridded dataset, featuring a spatial resolution of approximately 50 000 m. The digital elevation model (DEM) and derived features are crucial for vegetation mapping. The terrain data was from the Shuttle Radar Topography Mission (SRTM) by the United States Geological Survey (USGS) (Farr et al., 2007), with a spatial resolution of 30 m. At last, this study applied the mean sampling method in GEE to resample data on AP, AT, elevation, slope, and aspect variables derived from SRTM data to a 500 m spatial resolution for integration into vegetation mapping.

2.3 Vegetation mapping of the QTP at a 500 m resolution in 2020

2.3.1 Classification platform and algorithm

The GEE cloud platform offers a variety of machine learning models, such as support vector machine (SVM) and random forest (RF), the latter of which was utilized in this study for vegetation mapping (Gorelick et al., 2017; Zhang et al., 2023). The RF model is an ensemble-learning-based machine learning algorithm that mitigates the overfitting risk inherent in single decision trees by constructing and integrating multiple decision trees (Breiman, 2001). This approach not only enhances the accuracy and stability of predictions, but also has been widely applied in the vegetation mapping.

2.3.2 Construction of vegetation mapping features

The features used in vegetation mapping are divided into four categories (Table 3): terrain (elevation, slope, aspect), climate (AT and AP), surface reflectance (R , N , B , G , M , S_1 , S_2), and 14 index features which are constructed from the single-band surface reflectance. These features were derived from the MOD09A1 remote sensing imagery data spanning from 1 January to 31 December 2020. Additionally, six percentiles – 15 %, 30 %, 45 %, 60 %, 75 %, and 90 % – were calculated for the 7 reflectance bands and the 14 indices, representing the time series characteristics of each pixel. The 15th and 90th percentiles were used as substitutes for the minimum and maximum values of the time series observations, effectively mitigating the influence of extreme values in time series data (Zhang et al., 2021b). The 30 %, 45 %, 60 %, and 75 % features were selected to capture temporal

trends uniformly while avoiding feature redundancy (Sulla-Menashe et al., 2019; Zhang et al., 2024b). A total of 131 features were formulated from these four categories for vegetation mapping on the QTP.

2.3.3 Feature importance evaluation and feature selection

Among the 131 features used for vegetation mapping, 126 optical remote sensing features (excluding topography and climate) were constructed based on surface reflectance. These features are prone to severe collinearity issues, which can lead to model overfitting, increased computational costs, and diminished interpretability. To mitigate the issue of high collinearity among these features, the variance inflation factor (VIF) was employed, providing the ratio of variance in a model with multicollinearity among features to the variance in a model where multicollinearity is absent (James et al., 2013; Ngabire et al., 2022):

$$\text{VIF}_j = \frac{1}{1 - R_j^2}, \quad (1)$$

where VIF_j is the VIF for feature j and R_j^2 is the squared multiple correlation coefficient obtained from the regression of feature j with all other features. A higher VIF value indicates more severe collinearity, with a VIF_j greater than 30 suggesting significant collinearity in feature j . This study utilized the RF classifier to determine the optimal feature combination for vegetation classification. The importance of terrain, climate, and optical remote sensing features was assessed through the RF classifier and filtered using the VIF method (Ramosaj and Pauly, 2019; Zhang et al., 2019b). The features were then ranked according to the calculated importance, and the top-ranking feature, along with combinations such as the top two and top three, was used to construct various RF models for all possible feature combinations. The feature set with the smallest out-of-bag error was chosen as the best feature set for vegetation mapping on the QTP.

2.3.4 Evaluation of mapping accuracy

An optimal combination of terrain, climate, and optical remote sensing data in 2020 was integrated to achieve vegetation mapping of the QTP based on RF model in this study. The mapping accuracy was evaluated using the confusion matrix method, which involved calculating overall accuracy (OA, Eq. 2), kappa coefficient (kappa, Eq. 3), producer's accuracy (PA, Eq. 4), and user's accuracy (UA, Eq. 5).

$$\text{OA} = \frac{\sum_{i=1}^n m_i}{N}, \quad (2)$$

Table 3. Vegetation mapping features at a 500 m resolution on the QTP.

Category	Features	Formula	Description
Terrain	Elevation		The height of the terrain above sea level (m)
	Slope		The steepness of the terrain (°)
	Aspect		The direction the terrain slope faces (°)
Climate	AT		Annual average temperature
	AP		Annual precipitation
Surface reflectance	<i>R</i>	Red	620–670 nm
	<i>N</i>	Near-infrared	841–876 nm
	<i>B</i>	Blue	459–479 nm
	<i>G</i>	Green	545–565 nm
	<i>M</i>	Mid-infrared	1230–1250 nm
	<i>S</i> ₁	Shortwave infrared 1	1628–1652 nm
	<i>S</i> ₂	Shortwave infrared 2	2105–2155 nm
Vegetation index	NDVI	$\frac{N-R}{N+R}$	Normalized difference vegetation index (Tucker, 1979)
	EVI	$2.5 \frac{N-R}{N+6R-7.5B+1}$	Enhanced vegetation index (Huete et al., 1999)
	RVI	$\frac{N}{R}$	Ratio vegetation index (Crippen, 1990)
	DVI	$N - R$	Difference vegetation index (Roujean and Breon, 1995)
	SAVI	$\frac{(N-R) \cdot 1.5}{N+R+0.5}$	Soil adjusted vegetation index (Huete, 1988)
	GCVI	$\frac{N}{G} - 1$	Green chlorophyll vegetation index (Lobell et al., 2015)
	NIRV	$\frac{(N-R) \cdot N}{N+R}$	Near-infrared reflectance of vegetation (Badgley et al., 2017)
Urban index	NDBI	$\frac{S_1-N}{S_1+N}$	Normalized difference built-up index (Zha et al., 2003)
	IBI	$\frac{NDBI-(SAVI+(G-S_1)/(G+S_1))/2}{NDBI+(SAVI+(G-S_1)/(G+S_1))/2}$	Index-based built-up index (Xu, 2008)
Water index	NDWI	$\frac{G-N}{G+N}$	Normalized difference water index (McFeeters, 1996)
	LSWI	$\frac{N-S_1}{N+S_1}$	Land surface water index (Xiao et al., 2004)
Snow index	NDSI	$\frac{G-S_1}{G+S_1}$	Normalized difference snow index (Hall et al., 1995)
	NDGlaI	$\frac{G-R}{G+R}$	Normalized difference glacier index (Keshri et al., 2009)
Soil index	BI	$\frac{(S_1+R)-(N+B)}{(S_1+R)+(N+B)}$	Bare soil index (Chen et al., 2004)

$$\text{kappa} = \frac{N \times \sum_{i=1}^n m_i - \sum_{i=1}^n (G_i \times C_i)}{N^2 - \sum_{i=1}^n (G_i \times C_i)}, \tag{3}$$

$$\text{PA} = \frac{m_i}{G_i}, \tag{4}$$

$$\text{UA} = \frac{m_i}{C_i}, \tag{5}$$

where m_i is the count of correctly classified pixels for category i ; n is the count of categories; N is the overall quantity of classified pixels; C_i and G_i are the total counts of pixels classified as and actually in category i , respectively; OA is the overall accuracy; kappa is the kappa coefficient; PA is the mapping accuracy; and UA is the user accuracy.

2.4 Continuous annual vegetation mapping at a 500 m resolution from 2000 to 2022

The workflow represents the process for generating annual vegetation maps in QTP from 2000 to 2022 (Fig. 2). It begins with breakpoint detection, where the continuous change detection (CCD) algorithm (Eq. 6) is applied to the red, near-infrared, and shortwave infrared 1 bands of MOD09A1 data, identifying breakpoint regions for the QTP (Sect. 2.4.1). These breakpoint regions are then used in the potential vegetation change area identification algorithm to determine potential change areas for 2000–2019 and 2021–2022 (Sect. 2.4.2). Subsequently, the actual vegetation type identification algorithm is applied to these potential change areas, producing preliminary annual vegetation maps for the corresponding years (Sect. 2.4.3). Finally, the spatial–temporal consistency method is used to refine these preliminary maps, generating the annual vegetation maps of the QTP from 2000 to 2022 (Sect. 2.4.4).

2.4.1 Continuous change detection algorithm

The continuous change detection and classification (CCDC) algorithm integrates time series and spectral characteristics to detect temporal “breakpoints” and classify land cover changes. Originally developed for Landsat data (Zhu and Woodcock, 2014; Zhu et al., 2019), it has been extended to MODIS and Sentinel datasets (Shimizu et al., 2019; Tang et al., 2019; Xian et al., 2022). In this study, CCDC was applied to long-term vegetation mapping on the QTP, leveraging its ability to dynamically detect changes and update classifications to improve temporal consistency (Zhang et al., 2024b).

The CCDC algorithm comprises two components: continuous change detection (CCD) and classification (C). The CCD component uses a time series fitting model (Eq. 6) with harmonic and slope terms to analyze remote sensing data, capturing seasonal variations and the interannual changes. When residuals exceed a specified threshold, the corresponding time period is flagged as a “break”. The CCD segments the entire observation sequence into several subsequences, each separated by breaks. Each subsequence is then modeled independently using specific fitting coefficients (Fig. 2). The second part of the CCDC, the C, uses the RF method to classify each subsequence based on its fitting coefficients (Zhu and Woodcock, 2014). Considering the influence of terrain and climate on vegetation distribution in the QTP (Zhou et al., 2023), this study employed a pre-established 2020 classification model (Sect. 2.3) which incorporates terrain, climate, and remote sensing data to identify potential change areas.

$$\hat{\rho}(i, t) = c_{0i} + \sum_{n=1}^N \left(a_{ni} \cos \frac{2\pi n}{T} t + b_{ni} \sin \frac{2\pi n}{T} t \right) + c_{1i} t, \quad (6)$$

where $\hat{\rho}(i, t)$ is the predicted value of the i th band on Julian day t ; T is the average number of days in a year; a_{ni} and

b_{ni} are the harmonic coefficients of the n th order for the i th band; c_{0i} and c_{1i} are the intercept and slope coefficients, respectively; and N represents the highest order of harmonics, which is set to 3 in this study.

The CCDC algorithm was initially developed in MATLAB (Zhu and Woodcock, 2014) and later in Python (Brown et al., 2020). To address its high computational demands for long-term, large-scale monitoring, this study utilized the GEE platform, which provides the CCDC algorithm as a time segmentation tool (ee.Algorithms.TemporalSegmentation.Ccdc) (Arévalo et al., 2020; Pasquarella et al., 2022). This study primarily used GEE’s default parameters, modifying only breakpointBands and dateFormat. The breakpointBands parameter was set to include the red (R), near-infrared (N), and shortwave infrared 1 (S_1) bands, which correspond to chlorophyll content, leaf structure, and water content, respectively (Tucker, 1979; Curran, 1989; Gao, 1996). The dateFormat parameter was set to 2, indicating that all breakpoint times are represented as Unix timestamps. Finally, using the MOD09A1 remote sensing imagery from 2000 to 2022 and the CCD algorithm, breakpoint regions on the QTP during 2000–2022 were identified.

2.4.2 Potential vegetation change area identification algorithm

The annual mapping method in this study involves the baseline year T and the updated year $T - 1$. Therefore, the potential change area for the year $T - 1$ (S_{T-1}) depends on the breakpoint regions of T (B_T) and $T - 1$ (B_{T-1}). The detailed algorithm process is illustrated in Fig. 2. In general, at the same spatial location, if there is at least one breakpoint region in both year T and year $T - 1$ (gray areas), the corresponding location is marked as a potential change area for year $T - 1$ (green areas). This can be expressed as

$$S_{T-1} = (B_{T-1} \cap B_T) \cup (B_{T-1} \cap \overline{B_T}) \cup (\overline{B_{T-1}} \cap B_T), \quad (7)$$

where S_{T-1} is the potential vegetation change area for year $T - 1$; B_{T-1} and $\overline{B_T}$ are the breakpoint regions for year $T - 1$ and T ; and $\overline{B_{T-1}}$ and B_T are the non-breakpoint regions for year $T - 1$ and T , respectively. Finally, using the potential vegetation change area identification algorithm and the breakpoint regions on the QTP during 2000–2022 (Sect. 2.4.1), potential vegetation change areas were identified for 2000–2019 and 2021–2022. The year 2020 was used as the baseline year in this study and did not require updating.

2.4.3 Actual vegetation type identification algorithm

The potential vegetation change areas merely indicate regions where vegetation types might change, thus necessitating further identification of the actual vegetation change areas (Fig. 2). Based on the RF model constructed in 2020 (M_{2020}) and combining the terrain, climate, remote sensing data, and S_{T-1} , the actual vegetation types in the




Figure 2. Workflow for a 500 m resolution vegetation mapping on the QTP from 2000 to 2022.

potential change areas for $T - 1$ (R'_{T-1}) can be determined (Eq. 8). Subsequently, the $T - 1$ vegetation map (R_{T-1}) was obtained by overlaying and analyzing R'_{T-1} and R_T (Eq. 9).

$$R'_{T-1} = M_{2020}(F_{T-1}, S_{T-1}), \tag{8}$$

$$R_{T-1} = \begin{cases} R'_{T-1} & S_{T-1} \text{ and } R'_{T-1} \neq R_T, \\ R_T & \text{otherwise,} \end{cases} \tag{9}$$

where M_{2020} is the RF model for the year 2020; F_{T-1} is the vegetation mapping features for the year $T - 1$; S_{T-1} is the potential vegetation change area for the year $T - 1$; R'_{T-1} is the vegetation classification result for the potential change area of the year $T - 1$; R_T and R_{T-1} are the vegetation classification results for the years T and $T - 1$, respectively. Finally, by integrating the actual vegetation type identification algorithm with the potential vegetation change areas (Sect. 2.4.2), preliminary annual vegetation maps for 2000–2019 and 2021–2022 were generated.

2.4.4 Spatial–temporal consistency

The RF model used in this study, although exhibiting strong classification accuracy and robustness, employs a per-pixel classification strategy, which can result in salt-and-pepper noise (Yang and Huang, 2021). To mitigate this issue, this study employs a spatial–temporal constraint method, which assesses the consistency of each pixel’s label within a $3 \times 3 \times 3$ cube across both spatial and temporal dimensions (Fig. 2). The consistency, $C_{x,y,t}$, is calculated by averaging the agreement of the central pixel’s label with the labels of 27 surrounding pixels (Yang and Huang, 2021; Li et al., 2015):

$$C_{x,y,t} = \frac{1}{27} \sum_{i=x-1}^{x+1} \sum_{j=y-1}^{y+1} \sum_{k=t-1}^{t+1} I(\text{Label}_{x,y,t} = \text{Label}_{i,j,k}), \tag{10}$$

where I is an indicator function returning 1 if the labels match and otherwise 0. If $C_{x,y,t} < 0.5$ and $t > 2000$, the pixel is deemed misclassified, and its label is corrected to match the previous year’s central label. For the year 2000,

where previous year data are unavailable, the label is adjusted to the most frequent label in the spatial 3×3 area. This approach assumes that significant, inconsistent changes are unlikely both spatially and temporally, thereby enhancing the accuracy of the classification. Based on the preliminary annual vegetation maps (Sect. 2.4.3) and the spatial–temporal constraint method, the final product of this study, the annual vegetation maps of the QTP from 2000 to 2022, was produced.

3 Results

3.1 Vegetation mapping of the QTP at a 500 m resolution in 2020

Due to the significant collinearity among the 126 features in MODIS data, this study focused on 13 features with VIF below 30, including various band reflectance features and vegetation indices such as NDGIaI 15 %, NDGIaI 90 %, IBI 30 %, NDBI 90 %, IBI 15 %, IBI 90 %, EVI 90 %, NDVI 90 %, IBI 75 %, IBI 60 %, LSWI 90 %, M 90 %, and IBI 45 %. Combined with two climate factors (AT and AP) and three terrain features (elevation, slope, and aspect), an 18-feature set was created for out-of-bag (OOB) error analysis. The results indicated that the OOB error decreased as the number of features increased, reaching a minimum of 0.135 with 11 features, beyond which there was a slight increase (Fig. 3a). Subsequently, the top 11 features were selected to construct the RF model: AP, elevation, NDVI 90 %, AT, slope, M 90 %, EVI 90 %, NDBI 90 %, NDGIaI 90 %, NDGIaI 15 %, and IBI 75 %. Among them, AP, elevation, AT, and slope ranked first, second, fourth, and fifth in importance score, respectively. Therefore, climate and terrain are significant factors influencing vegetation distribution on the QTP (Fig. 3b). Finally, the 2020 vegetation map of the QTP was generated on the GEE platform at a 500 m spatial resolution (Fig. 4) based on the developed RF model.

The 2020 vegetation map of the QTP includes 16 types (Fig. 4). Among these, four types cover areas larger than $400\,000\text{ km}^2$: AM, AG, AD, and AV. The AM covers the largest area, approximately $798\,000\text{ km}^2$, mainly in the eastern and southern regions of Gansu and Qinghai provinces. The AG covers around $578\,000\text{ km}^2$, predominantly in the central part of the plateau, with a distinct transitional zone adjoining the AM. The AD, covering about $562\,000\text{ km}^2$, is primarily located in the Xinjiang region. The AV covers approximately $474\,000\text{ km}^2$ and is widely distributed from above the treeline or shrub zone to the lower limit of the perennial snow line on the QTP. Among the five forest types in the QTP, ECF cover the largest area, about $154\,000\text{ km}^2$, mainly distributed in the Hengduan Mountains in the south-east of the QTP and near the Indus River in the west. DCF cover the smallest area, approximately $12\,000\text{ km}^2$, primarily located along the northwestern border of the QTP.

3.2 Continuous annual vegetation mapping at a 500 m resolution from 2000 to 2022

This study used the CCD algorithm to detect breaks in the dense time series MODIS data of the QTP from 2000 to 2022, defining these breaks as potential changes (PCs) (Fig. 5a). Subsequently, an RF model incorporating terrain, climate, and remote sensing data was applied to annually update and generate vegetation maps of the QTP, identifying changes between adjacent years as real changes (RCs) (Fig. 5b).

Approximately 31.8 % of the QTP is classified as PC areas. Regions with one or two instances of PC account for 18.2 % and 7.12 % of the area, respectively (Fig. 5g), primarily located at the junctions of AM, AG, and AD in the central part of the plateau (Fig. 5a). Notably, areas near Hala Lake in the northeast (ROI1) and Lake Manasarovar in the southwest Ali region (ROI2) experienced more than five instances of change, covering approximately 0.82 % of the total area of the QTP (Fig. 5c and d). Compared to the PC areas, the RC areas from the annual vegetation maps of the QTP are significantly smaller (Fig. 5b), encompassing only 3.17 % of the total area. As shown in Fig. 5g, regions with one or two instances of RC account for 2.75 % and 0.34 % of the QTP, respectively. Areas with three or more instances of RC make up only 0.08 %, with no regions experiencing five or more instances of change. In all categories of change frequency, the area of RC is consistently smaller than that of PC. In the RC regions of ROI1 and ROI2, only a few areas show one or two instances of change. Notably, in ROI2, no areas with five instances of PC (red) appear in the corresponding RC regions; only a small portion shows a single instance of RC change.

Although the CCD model, which relies on dense spectral data, often detects many false changes – meaning that many breaks detected by CCD do not necessarily represent land type transformations (Zhu et al., 2019) – the RF model based on climate, terrain, and remote sensing data produces annual vegetation maps that effectively reduce these false changes. The frequency and area of detected change regions are significantly smaller compared to the CCD model results.

Figure 6 illustrates the area changes in 16 vegetation types on the QTP from 2000 to 2022. All forest types remained relatively stable, with changes within 0.3 %. Among the five forest types, ECF showed a small decrease, reducing by approximately 400 km^2 (0.28 %), while DCF had a small increase by about 14 km^2 (0.12 %). Except for DBF, SC and ASM have limited distributions on the plateau. ASM decreased by about 300 km^2 (1.02 %), primarily before 2005. AM, AG, AV, and AD are the four most widespread types on the plateau. AM and AV showed significant increases, with AM rising by about 8800 km^2 (1.11 %) and AV by about $12\,600\text{ km}^2$ (2.73 %). In contrast, AD significantly decreased by about $21\,500\text{ km}^2$ (3.69 %). AG fluctuated but notably declined by about 3100 km^2 (0.54 %) after 2016. CV increased




Figure 3. Evaluation of the 2020 vegetation classification model on the QTP.




Figure 4. Vegetation types and spatial distribution at a 500 m resolution on the QTP in 2020.

by approximately 500 km² (1.32 %), mainly after 2018. WA showed the most significant change, increasing by about 4300 km² (8.40 %) between 2000 and 2022. GS decreased by approximately 200 km² (0.17 %), primarily after 2010. Both WE and NVA showed a decreasing trend, with reductions of about 600 and 100 km², respectively, from 2000 to 2022.

As shown in Fig. 7, most RC occurred only once, predominantly in the northern, eastern, and southwestern parts of Selin Co. In 2000, these areas were classified as WE, AD, and AG, respectively. From 2000 to 2022, these regions gradually transformed into WA, as evident in the remote sensing images. Specifically, the northern WE transi-

tioned to WA between 2000 and 2015, the eastern AD between 2000 and 2010, and the southern AG between 2000 and 2005. The WA in this region increased from approximately 43.43 % in 2000 to 48.73 % in 2022, marking a 5.3 % rise. Simultaneously, AG, AD, and WE decreased by 1.09 %, 2.56 %, and 1.65 %, respectively. The eastern AD was the most significant contributor to the expansion of Selin Co’s water.

In the northeastern part of Hala Lake, close to ROI1, the area primarily consists of AG, AV, AD, and GS. Figure 8b shows that most changes in this region from 2000 to 2022 occurred once or twice. The area of GS changed very lit-




Figure 5. Overview of potential change (PC) and real change (RC) areas on the QTP from 2000 to 2022. (a, b) Distribution of PC and RC on the QTP. (c–f) PC and RC areas in ROI1 and ROI2. (g) Area statistics of PC and RC on the QTP.




Figure 6. Annual area changes for 16 vegetation types on the QTP from 2000 to 2022.




Figure 7. Annual vegetation cover changes in Area 1 (31.5002° N, 88.6985° E). (a, c) Remote sensing images from 2000 and 2022, respectively. (b) Distribution and frequency of RC. (d–i) Annual vegetation maps for 2000, 2005, 2010, 2015, 2020, and 2022. (j, k) Area proportions of major vegetation types in 2000 and 2022, respectively.

tle over the past 23 years, increasing slightly from 6.32 % to 6.34 %. In contrast, AG expanded significantly, from 41.67 % to 53.49 %, which is an increase of about 11.82 %. Meanwhile, the areas of AD and AV decreased by 9.51 % and 2.33 %, respectively. These changes mainly occurred in transition zones, particularly between AG and AD, where most conversion from AD to AG happened between 2000 and 2010. The shift from AV to AG mostly occurred between 2000 and 2005. Notably, some areas in Fig. 8b experienced two changes. These changes mainly reflect fluctuations between AD and AG (Fig. 8c and d), likely due to similar climate and spectral characteristics in the transition

zones, causing some pixels to switch between the two types multiple times.

3.3 Accuracy validation

The annual vegetation map of the QTP is dynamically updated using 2020 as the reference year, making its accuracy crucial for long-term product reliability. Due to minimal differences in vegetation distribution between adjacent years, samples from 2019, 2020, and 2021 were used to validate the reference map. This validation set included 1175 samples. The results showed an OA of 83.32 % for 2020, with a kappa




Figure 8. Annual vegetation cover changes in Area 2 (37.6311° N, 95.8644° E). (a, c) Remote sensing images from 2000 and 2022, respectively. (b) Distribution and frequency of RC. (d–i) Annual vegetation maps for 2000, 2005, 2010, 2015, 2020, and 2022. (j, k) Area proportions of major vegetation types in 2000 and 2022, respectively.

coefficient of 0.82 (Table 4). For most vegetation types, the UA for DCF, AM, AV, AD, and WA exceeded 90 %, and the PA for DCF, WE, WA, and GS was also above 90 %. However, the UA and PA for CBMF were only 57.14 % and 48.00 %, respectively. This lower accuracy is likely because CBMF, as a mix of coniferous and broad-leaved forests, is often misclassified by the RF model as EBF, ECF, or DBF due to spatial similarities. Overall, the 2020 vegetation map, validated with 1175 samples, achieved sufficient accuracy to be used as the reference for dynamic updates.

The third-party samples (1014 samples) and visual interpretation samples (1002 samples) were used to validate the

accuracy of annual vegetation maps. The third-party samples mainly include grasslands, AD, and GS. Because the first all-season sample set and the global land cover validation samples did not differentiate between AM and AG, these two types were combined under the grasslands category during the validation process. The third-party samples mainly originate from 2014 (190 samples) and 2015 (666 samples), with smaller numbers from other years. The visual interpretation samples exhibit stable characteristics across multiple years, making them suitable for validating annual vegetation maps from 2000 to 2019.

Table 4. Confusion matrix, PA, and UA of vegetation mapping at a 500 m resolution on the QTP in 2020.

	EBF	ECF	CBMF	DBF	DCF	SC	ASM	AM	AG	AV	AD	CV	WE	WA	NVA	GS	Total	UA (%)
EBF	56	11	8	1	0	0	0	0	0	0	0	0	0	0	0	0	76	73.68
ECF	2	74	3	2	1	1	0	3	0	0	0	0	0	0	0	0	86	86.05
CBMF	0	4	12	4	0	0	0	0	0	0	0	1	0	0	0	0	21	57.14
DBF	5	4	2	51	0	0	0	0	0	0	0	0	0	0	0	0	62	82.26
DCF	0	0	0	0	87	0	0	0	2	0	2	0	0	0	0	0	91	95.60
SC	0	0	0	0	0	46	2	6	1	0	1	2	0	0	0	0	58	79.31
ASM	5	0	0	0	0	8	29	4	1	0	0	1	0	0	0	0	48	60.42
AM	0	0	0	0	0	0	1	54	2	0	0	0	0	0	0	0	57	94.74
AG	0	1	0	0	0	4	0	8	61	0	7	2	0	0	0	0	83	73.49
AV	0	0	0	0	0	0	1	2	5	89	0	0	0	0	0	0	97	91.75
AD	0	0	0	0	5	1	0	1	1	0	85	0	0	0	0	0	93	91.40
CV	0	1	0	1	0	1	0	6	0	0	2	87	0	0	7	0	105	82.86
WE	0	0	0	0	2	0	0	6	1	0	4	3	60	0	0	0	76	78.95
WA	0	0	0	0	0	1	0	3	0	0	0	0	0	83	0	0	87	95.40
NVA	0	2	0	0	0	1	0	2	0	0	0	9	0	0	22	0	36	61.11
GS	0	0	0	0	0	0	0	0	0	16	0	0	0	0	0	83	99	83.84
Total	68	97	25	59	95	63	33	95	74	105	101	105	60	83	29	83	1175	
PA (%)	82.35	76.29	48.00	86.44	91.58	73.02	87.88	56.84	82.43	84.76	84.16	82.86	100.00	100.00	75.86	100.00		83.32




Figure 9. Annual overall accuracy of vegetation maps from 2000 to 2019.

Validation results indicated that accuracy from 2000 to 2019 remained above 80 %, with the lowest being 82.07 % in 2015 (Fig. 9). Upon inspection, most of the misclassified samples were located in the barren land (vegetation cover < 10 %) category in the first all-season sample set, which corresponds to AD in the vegetation maps. However, some samples of this category were classified as AG in the vegetation map. There are definitional differences between barren land and AD. Additionally, AD and AG represent two largest vegetation types on the QTP, and there are significant transitional zones between these two types (Fig. 4). As a result, the accuracy in 2015, when a large number of third-party samples were used, was slightly lower than in other years. Moreover, because there were insufficient validation samples for 2021 and 2022 and 2020 has already been validated as the reference year, the overall accuracy of the annual vegetation maps is estimated to be 83.27 % based on the average accuracy from 2000 to 2019.

4 Discussion

4.1 Evaluating the efficacy of the CCD algorithm in annual vegetation mapping from 2000 to 2022

This study proposed a method for long-time continuous annual vegetation mapping. Specifically, the CCD algorithm was applied to MODIS data from 2000 to 2022 to detect breakpoints. Subsequent processes involved identifying potential change areas, recognizing true vegetation types, and ensuring spatial–temporal consistency. This enabled consistent mapping of vegetation on the QTP annually from 2000 to 2022. The CCD algorithm in this study used harmonic functions to fit long-term remote sensing images, thereby identifying breakpoints and determining the timing of these breakpoints. To ensure the algorithm effectively captures vegetation dynamics, we selected the *R* (red), *N* (near-infrared), and *S*₁ (shortwave infrared 1) bands for breakpoint detection. The *R* band is related to chlorophyll content, the *N* band to canopy structure, and the *S*₁ band to vegetation water content (Tucker, 1979; Curran, 1989; Gao, 1996). Combining these complementary bands enables the CCD al-




Figure 10. Validation of vegetation mapping: consistent change detection in WA sample (35.2425° N, 82.9294° E) by CCD, Landsat, and annual vegetation maps.

gorithm to effectively detect breakpoints in vegetation dynamics on the QTP.

For instance, in Fig. 10, the CCD algorithm was applied to detect changes in the R , N , and S_1 bands of the sampling site from 2000 to 2022. The results indicated that there was a breakpoint in 2011 (highlighted in yellow), dividing the period into Fit1 (from 2000 to 2011) and Fit2 (from 2011

to 2022). The annual amplitude of the three bands exhibited small fluctuations with stable interannual patterns. In contrast, the amplitude of the R and N bands in Fit2 far exceeded that of Fit1, showing significant differences in seasonal patterns within the year. Additionally, for the S_1 band, the overall reflectance in Fit2 was substantially lower than in Fit1, reflecting distinct seasonal fluctuations within the year. Based




Figure 11. Validation of vegetation mapping: consistency of no change detection in forest sample (28.7617° N, 95.2794° E) by CCD, Landsat, and annual vegetation maps.

on the annual Landsat images from 2009 to 2013, there was a noticeable expansion of WA in the selected area, marking a transition from AD prior to 2011 to WA thereafter. The annual vegetation maps accurately captured this transition, with the area represented as brown–yellow (AD) before and up to 2010 and shifted to deep blue (WA) from 2011 onward.

This change was further corroborated by the CCD fitting results and visual interpretation of the Landsat images.

In Fig. 11, the CCD algorithm indicated that there were no breakpoints from 2000 to 2022. The annual and interannual variations in the R , N , and S_1 bands were stable over the years. Combined with the Landsat images, the selected area was consistently classified as EBF without any detected




Figure 12. Validation of vegetation mapping: CV sample (35.2474° N, 100.8570° E) changes detected by CCD not reflected in Landsat and annual vegetation maps.

changes. The long-time annual vegetation maps consistently reflected this characteristic.

However, abrupt changes detected by the CCD algorithm may not always accurately reflect real changes on the ground (Zhu and Woodcock, 2014; Du et al., 2023). In Fig. 12, an area of CV on the QTP was examined. The CCD results indicated a breakpoint in 2018, with data from 2000 to 2018 cat-

egorized as Fit1 and post-2018 as Fit2. Although Fit2 shows greater amplitude in the R , N , and S_1 bands compared to Fit1, the waveform remains similar. Despite a breakpoint being detected, Landsat images from 2016 to 2020 confirm that the area consistently featured CV. The detected changes were likely caused by variations in cultivation practices in 2018 or similar factors rather than actual changes in vegetation. As a




Figure 13. Comparison with the annual vegetation maps, MCD12Q1, CLCD, and GLC-FCS30D. (a) Region dominated by WA and AG. (b) Region dominated by GA and AM.

result, the annual vegetation maps did not capture the change detected by the CCD in 2018.

Given its performance, the CCD algorithm effectively identifies the regions and timings of breakpoints in long-time remote sensing imagery, making it a suitable foundational method for detecting potential change areas in annual vegetation mapping. Although the CCD algorithm is susceptible to false positives due to factors such as changes in cultivated species (Fig. 12), the subsequent methods employed in this

study, including potential area identification, true vegetation type recognition, and spatial–temporal constraints, help mitigate these false positive errors (Fig. 12).

4.2 Inter-comparison with other products

MCD12Q1 (Friedl and Sulla-Menashe, 2022; Sulla-Menashe et al., 2019), CLCD (Yang and Huang, 2021, 2023), and GLC-FCS30D (Liu et al., 2023; Zhang et al., 2024b) were




Figure 14. Comparison with the annual vegetation maps, MCD12Q1, CLCD, and GLC-FCS30D. (a) Region dominated by AM and AG. (b) Region dominated by ECF and AM).

selected for cross-comparison with the annual vegetation maps. MCD12Q1 provides annual global land cover data at a 500 m resolution from 2001 to 2022 using the International Geosphere–Biosphere Programme (IGBP) classification system, which includes 17 categories, such as evergreen needleleaf forest, grassland, and cropland. CLCD offers 30 m land cover data for China from 1990 to 2022, classifying land into nine types: cropland, forest, shrub, grassland, water, snow/ice, barren, impervious, and wetland. GLC-FCS30D

delivers land cover data from 1985 to 2022, with 5-year intervals before 2000 and annual intervals thereafter, covering 35 detailed land cover categories. Based on the temporal coverage of these products, annual vegetation maps from 2001, 2011, and 2021 were selected for cross-validation, with a regional focus on four main types: WA, GS, grassland, and forest.

Landsat remote sensing images reveal significant shrinkage of water in this region from 2001 to 2021 (Fig. 13a).

In 2001, all four products accurately depicted the distribution of WA, with clear boundaries and complete representation. By 2011, GLC-FCS30D identified part of the WA as WE, while the other three products consistently showed it as WA. By 2021, the WA had markedly shrunk compared to their extent in 2001. Compared to the Landsat images, CLCD showed an omission of WA, with most areas transforming into AD, while GLC-FCS30D misclassified part of the WA as WE and identified some non-water areas as WA.

For GS areas, all four products demonstrated good identification in 2001, 2011, and 2021 (Fig. 13b). Uniquely, the annual vegetation maps identified additional alpine vegetation. Consequently, in the vicinity of GS, MCD12Q1 categorized these areas as AD, while CLCD and GLC-FCS30D largely classified them as grassland. The annual vegetation maps not only preserved the distribution details of GS, AD, and AG from the comparison products, but also provided a more detailed classification of AV.

However, the three comparison products classified this region simply as grassland without further differentiation (Fig. 14a). Grassland is the dominant vegetation type on the QTP, covering approximately 57 % of the QTP. According to Zhou et al. (2022a), grasslands on the QTP can be divided into AG, AM, ASM, and AD. Therefore, it is essential to differentiate between AG, AM, ASM, and AD. In Fig. 14a, the overall distribution of AG and AM has remained consistent over time, though changes have occurred in their transition zones. For example, in the southwestern AM region of Fig. 14a, some AG areas have gradually transitioned to AM over the past 23 years.

Forests are sparsely distributed on the QTP, found mainly on the southern slopes of the Himalayas and in the southeastern region (Fig. 14b). Their distribution has remained stable over the years with minimal change. Among the four products, CLCD did not differentiate forest types and thus represents them under the general category of forest. In contrast, the other three products classified the forested areas as ECF. MCD12Q1 shows a smaller forest extent and includes a notable proportion of CBMF within this range. The forest extent in the annual vegetation maps, CLCD, and GLC-FCS30D shows high consistency with the Landsat images. However, the annual vegetation maps identified some ASM in the meadow areas adjacent to forests.

5 Data availability

The 500 m annual vegetation maps of QTP from 2000 to 2022 are available at <https://doi.org/10.11888/Terre.tpdc.301205> (Zhou et al., 2024). The vegetation maps are stored in TIFF format, with the file name QTP_Vegetation_Map_XXXX.tif, where XXXX represents the year. All files can be opened and reprocessed using software such as ArcGIS, QGIS, and ENVI. Each TIFF dataset contains values from 0 to 16,

where 0 represents invalid values and 1 to 16 correspond to the 16 vegetation types listed in Table 1. We used MOD09A1 (<https://doi.org/10.5067/MODIS/MOD09A1.061>, Vermote, 2021), MCD12Q1 (<https://doi.org/10.5067/MODIS/MCD12Q1.061>, Friedl and Sulla-Menashe, 2022), and SRTM (Farr et al., 2007) data used in this study. The 1 km monthly precipitation dataset for China (1901–2023), the 1 km monthly mean temperature dataset for China (1901–2023), the vegetation map of the Qinghai–Tibet Plateau in 2020 with a 10 m spatial resolution, and QTP boundary dataset were provided by the National Tibetan Plateau/Third Pole Environment Data Center (<https://data.tpdc.ac.cn>, National Tibetan Plateau/Third Pole Environment Data Center, 2024). The first all-season sample set can be downloaded from <https://data-starcloud.pcl.ac.cn/resource/54> (Peng Cheng Laboratory, 2024). The global land cover validation samples can be downloaded from <https://doi.org/10.5281/zenodo.3551995> (Liu et al., 2019). The CLCD can be downloaded from <https://doi.org/10.5281/zenodo.8176941> (Yang and Huang, 2023), and the GLC-FCS30D can be download from <https://doi.org/10.5281/zenodo.8239305> (Liu et al., 2023).

6 Conclusions

long time series of annual regional vegetation types and geographic distribution data are vital for examining the impact of climate change on vegetation and its evolutionary trends. In this study, annual vegetation of the QTP from 2000 to 2022 at a 500 m spatial resolution was mapped through the MOD09A1 product together with a reference year vegetation classification map and a breakpoint detection algorithm. The study achieved an overall accuracy of 83.27 % for continuous annual vegetation mapping at a 500 m resolution from 2000 to 2022. The study supports the use of remote sensing data to mapping long-term continuous annual vegetation. Furthermore, it facilitates the elucidation of the spatial and temporal evolution of regional and global vegetation against the background of global warming.

Author contributions. GZ and HR conceived the study. LZ and HR analyzed the data and drafted the paper. XL and MZ supported the generation of the dataset and the analysis of the results.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes ev-

ery effort to include appropriate place names, the final responsibility lies with the authors.

Acknowledgements. We would like to express our sincere gratitude to all the data providers for their valuable contributions to this research. Special thanks are extended to the reviewers for their insightful comments and suggestions, which significantly improved the quality of this work. We also acknowledge the support of our colleagues and collaborators who have contributed to various aspects of this study.

Review statement. This paper was edited by Peng Zhu and reviewed by Shudong Wang and seven anonymous referees.

Financial support. This work was supported by the Second Tibetan Plateau Scientific Expedition and Research Program (grant no. 2019QZKK0106) and the National Natural Science Foundation of China (grant no. 42141007).

References

- Arévalo, P., Bullock, E. L., Woodcock, C. E., and Olofsson, P.: A suite of tools for continuous land change monitoring in Google Earth Engine, *Front. Clim.*, 2, 111051, <https://doi.org/10.3389/fclim.2020.576740>, 2020.
- Badgley, G., Field, C. B., and Berry, J. A.: Canopy near-infrared reflectance and terrestrial photosynthesis, *Sci. Adv.*, 3, e1602244, <https://doi.org/10.1126/sciadv.1602244>, 2017.
- Bontemps, S., Defourny, P., Bogaert, E. V., Arino, O., Kalogirou, V., and Perez, J. R.: GLOBCOVER 2009 Products Description and Validation Report, http://due.esrin.esa.int/files/GLOBCOVER2009_Validation_Report_2.2.pdf (last access: 29 September 2024), 2010.
- Breiman, L.: Random forests, *Mach. Learn.*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.
- Brown, J. F., Tollerud, H. J., Barber, C. P., Zhou, Q., Dwyer, J. L., Vogelmann, J. E., Loveland, T. R., Woodcock, C. E., Stehman, S. V., Zhu, Z., Pengra, B. W., Smith, K., Horton, J. A., Xian, G., Auch, R. F., Sohl, T. L., Saylor, K. L., Gallant, A. L., Zelenak, D., Reker, R. R., and Rover, J.: Lessons learned implementing an operational continuous United States national land change monitoring capability: The Land Change Monitoring, Assessment, and Projection (LCMAP) approach, *Remote Sens. Environ.*, 238, 111356, <https://doi.org/10.1016/j.rse.2019.111356>, 2020.
- Chen, J., Chen, J., Liao, A., Cao, X., Chen, L., Chen, X., He, C., Han, G., Peng, S., Lu, M., Zhang, W., Tong, X., and Mills, J.: Global land cover mapping at 30 m resolution: A POK-based operational approach, *ISPRS J. Photogram. Remote Sens.*, 103, 7–27, <https://doi.org/10.1016/j.isprsjprs.2014.09.002>, 2015.
- Chen, W., Liu, L., Zhang, C., Wang, J., Wang, J., and Pan, Y.: Monitoring the seasonal bare soil areas in Beijing using multitemporal TM images, in: *IGARSS 2004: 2004 IEEE International Geoscience and Remote Sensing Symposium*, 20–24 September 2004, Anchorage, AK, 3379–3382, <https://doi.org/10.1109/IGARSS.2004.1370429>, 2004.
- Crippen, R. E.: Calculating the vegetation index faster, *Remote Sens. Environ.*, 34, 71–73, [https://doi.org/10.1016/0034-4257\(90\)90085-z](https://doi.org/10.1016/0034-4257(90)90085-z), 1990.
- Curran, P. J.: Remote sensing of foliar chemistry, *Remote Sens. Environ.*, 30, 271–278, [https://doi.org/10.1016/0034-4257\(89\)90069-2](https://doi.org/10.1016/0034-4257(89)90069-2), 1989.
- Defourny, P., Bontemps, S., Lamarche, C., Brockmann, C., Boettcher, M., Wevers, J., and Kirches, G.: Land cover CCI: product user guide version 2.0, https://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf (last access: 18 November 2024), 2017.
- Ding, Y. and Peng, S.: Spatiotemporal trends and attribution of drought across China from 1901–2100, *Sustainability*, 12, 477, <https://doi.org/10.3390/su12020477>, 2020.
- Du, Z., Yu, L., Li, X., Zhao, J., Chen, X., Xu, Y., Yang, P., Yang, J., Peng, D., Xue, Y., and Gong, P.: Integrating remote sensing temporal trajectory and survey statistics to update land use/land cover maps, *Int. J. Digit. Earth*, 16, 4428–4445, <https://doi.org/10.1080/17538947.2023.2274422>, 2023.
- Editorial Board of the Vegetation Map of China: Chinese Academy of Sciences, Vegetation Map of the People's Republic of China (1 : 1 000 000), Geology Publishing House, Beijing, China, ISBN 978-7116045132, 2007.
- Farr, T. G., Rosen, P. A., Caro, E., Crippen, R. E., Duren, R., Hensley, S., Kobrick, M., Paller, M., Rodriguez, E., Roth, L., Seal, D., Shaffer, S., Shimada, J., Umland, J., Werner, M., Oskin, M., Burbank, D., and Alsdorf, D.: The shuttle radar topography mission, *Rev. Geophys.*, 45, RG2004, <https://doi.org/10.1029/2005RG000183>, 2007.
- Feng, M. and Li, X.: Land cover mapping toward finer scales, *Sci. Bull.*, 65, 1604–1606, <https://doi.org/10.1016/j.scib.2020.06.014>, 2020.
- Feng, M., Huang, C., Channan, S., Vermote, E. F., Masek, J. G., and Townshend, J. R.: Quality assessment of Landsat surface reflectance products using MODIS data, *Comput. Geosci.*, 38, 9–22, <https://doi.org/10.1016/j.cageo.2011.04.011>, 2012.
- Friedl, M. and Sulla-Menashe, D.: MODIS/Terra+Aqua Land Cover Type Yearly L3 Global 500 m SIN Grid V061, NASA EOSDIS Land Processes Distributed Active Archive Center [data set], <https://doi.org/10.5067/MODIS/MCD12Q1.061>, 2022.
- Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, X.: MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets, *Remote Sens. Environ.*, 114, 168–182, <https://doi.org/10.1016/j.rse.2009.08.016>, 2010.
- Gao, B. C.: NDWI – A normalized difference water index for remote sensing of vegetation liquid water from space, *Remote Sens. Environ.*, 58, 257–266, [https://doi.org/10.1016/S0034-4257\(96\)00067-3](https://doi.org/10.1016/S0034-4257(96)00067-3), 1996.
- Gong, P., Wang, J., Yu, L., Zhao, Y., Zhao, Y., Liang, L., Niu, Z., Huang, X., Fu, H., Liu, S., Li, C., Li, X., Fu, W., Liu, C., Xu, Y., Wang, X., Cheng, Q., Hu, L., Yao, W., Zhang, H., Zhu, P., Zhao, Z., Zhang, H., Zheng, Y., Ji, L., Zhang, Y., Chen, H., Yan, A., Guo, J., Yu, L., Wang, L., Liu, X., Shi, T., Zhu, M., Chen, Y., Yang, G., Tang, P., Xu, B., Giri, C., Clinton, N., Zhu, Z., Chen, J., and Chen, J.: Finer resolution observation and monitoring of global land cover: First mapping results with Landsat TM and ETM+ data, *Int. J. Remote Sens.*, 34, 2607–2654, <https://doi.org/10.1080/01431161.2012.748992>, 2013.

- Gong, P., Liu, H., Zhang, M., Li, C., Wang, J., Huang, H., Clinton, N., Ji, L., Li, W., Bai, Y., Chen, B., Xu, B., Zhu, Z., Yuan, C., Suen, H., Guo, J., Xu, N., Li, W., Zhao, Y., Yang, J., Yu, C., Wang, X., Fu, H., Yu, L., Dronova, I., Hui, F., Cheng, X., Shi, X., Xiao, F., Liu, Q., and Song, L.: Stable classification with limited sample: Transferring a 30-m resolution sample set collected in 2015 to mapping 10-m resolution global land cover in 2017, *Sci. Bull.*, 64, 370–373, <https://doi.org/10.1016/j.scib.2019.03.002>, 2019.
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, R.: Google Earth Engine: Planetary-scale geospatial analysis for everyone, *Remote Sens. Environ.*, 202, 18–27, <https://doi.org/10.1016/j.rse.2017.06.031>, 2017.
- Hall, D. K., Riggs, G. A., and Salomonson, V. V.: Development of methods for mapping global snow cover using moderate resolution imaging spectroradiometer data, *Remote Sens. Environ.*, 54, 127–140, [https://doi.org/10.1016/0034-4257\(95\)00137-P](https://doi.org/10.1016/0034-4257(95)00137-P), 1995.
- Hu, M., Zhou, G., Lv, X., Zhou, L., Wang, X., He, X., and Tian, Z.: Warming Has Accelerated the Melting of Glaciers on the Tibetan Plateau, but the Debris-Covered Glaciers Are Rapidly Expanding, *Remote Sens.*, 15, 132, <https://doi.org/10.3390/rs15010132>, 2023.
- Huete, A., Justice, C., and Van Leeuwen, W.: MODIS vegetation index (MOD13), Algorithm theoretical basis document, <https://www.cen.uni-hamburg.de/en/icdc/data/land/docs-land/modis-collection6-vegetation-index-atbd-mod13-v03-1.pdf> (last access: 18 November 2024), 1999.
- Huete, A. R.: A soil-adjusted vegetation index (SAVI), *Remote Sens. Environ.*, 25, 295–309, [https://doi.org/10.1016/0034-4257\(88\)90106-X](https://doi.org/10.1016/0034-4257(88)90106-X), 1988.
- Immerzeel, W. W., Van Beek, L. P. H., and Bierkens, M. F. P.: Climate change will affect the Asian water towers, *Science*, 328, 1382–1385, <https://doi.org/10.1126/science.1183188>, 2010.
- James, G., Witten, D., Hastie, T., and Tibshirani, R.: An introduction to statistical learning, Springer, <https://doi.org/10.1007/978-1-0716-1418-1>, 2013.
- Keshri, A. K., Shukla, A., and Gupta, R. P.: ASTER ratio indices for supraglacial terrain mapping, *Int. J. Remote Sens.*, 30, 519–524, <https://doi.org/10.1080/01431160802385459>, 2009.
- Li, C., Peng, G., Wang, J., Zhu, Z., Biging, G. S., Yuan, C., Hu, T., Zhang, H., Wang, Q., Li, X., Liu, X., Xu, Y., Guo, J., Liu, C., Hackman, K. O., Zhang, M., Cheng, Y., Yu, L., Yang, J., Huang, H., and Clinton, N.: The first all-season sample set for mapping global land cover with Landsat-8 data, *Sci. Bull.*, 62, 508–515, <https://doi.org/10.1016/j.scib.2017.03.011>, 2017.
- Li, X., Gong, P., and Liang, L.: A 30-year (1984–2013) record of annual urban dynamics of Beijing City derived from Landsat data, *Remote Sens. Environ.*, 166, 78–90, <https://doi.org/10.1016/j.rse.2015.06.007>, 2015.
- Li, Y., Hou, Z., Zhang, L., Song, C., Piao, S., Lin, J., Peng, S., Fang, K., Yang, J., Qu, Y., Wang, Y., Li, J., Li, R., and Yao, X.: Rapid expansion of wetlands on the Central Tibetan Plateau by global warming and El Niño, *Sci. Bull.*, 68, 485–488, <https://doi.org/10.1016/j.scib.2023.02.021>, 2023.
- Liu, H., Gong, P., Wang, J., Wang, X., Ning, G., and Xu, B.: Production of global daily seamless data cubes and quantification of global land cover change from 1985 to 2020-iMap World 1.0, *Remote Sens. Environ.*, 258, 112364, <https://doi.org/10.1016/j.rse.2021.112364>, 2021a.
- Liu, L., Gao, Y., Zhang, X., Chen, X., and Xie, S.: A Dataset of Global Land Cover Validation Samples, Zenodo [data set], <https://doi.org/10.5281/zenodo.3551995>, 2019.
- Liu, L., Zhang, X., Gao, Y., Chen, X., Shuai, X., and Mi, J.: Finer Resolution Mapping of Global Land Cover: Recent Developments, Consistency Analysis, and Prospects, *J. Remote Sens.*, 2021, 5289697, <https://doi.org/10.34133/2021/5289697>, 2021b.
- Liu, L., Zhang, X., and Zhao, T.: GLC_FCS30D: the first global 30-m land-cover dynamic monitoring product with fine classification system from 1985 to 2022, Zenodo [data set], <https://doi.org/10.5281/zenodo.8239305>, 2023.
- Lobell, D. B., Thau, D., Seifert, C., Engle, E., and Little, B.: A scalable satellite-based crop yield mapper, *Remote Sens. Environ.*, 164, 324–333, <https://doi.org/10.1016/j.rse.2015.04.021>, 2015.
- McFeeters, S. K.: The use of the Normalized Difference Water Index (NDWI) in the delineation of open water features, *Int. J. Remote Sens.*, 17, 1425–1432, <https://doi.org/10.1080/01431169608948714>, 1996.
- National Tibetan Plateau/Third Pole Environment Data Center: Data bank for Tibetan Plateau research projects, <https://data.tpdc.ac.cn> (last access: 29 September 2024), 2024.
- Ngabire, M., Wang, T., Xue, X., Liao, J., Sahbeni, G., Huang, C., Duan, H., and Song, X.: Soil salinization mapping across different sandy land-cover types in the Shiyang River Basin: A remote sensing and multiple linear regression approach, *Remote Sens. Appl. Soc. Environ.*, 28, 100847, <https://doi.org/10.1016/j.rsase.2022.100847>, 2022.
- Pasquarella, V. J., Arévalo, P., Bratley, K. H., Bullock, E. L., Gorelick, N., Yang, Z., and Kennedy, R. E.: Demystifying LandTrendr and CCDC temporal segmentation, *Int. J. Appl. Earth Obs.*, 110, 102806, <https://doi.org/10.1016/j.jag.2022.102806>, 2022.
- Peng, S.: 1-km monthly mean temperature dataset for China (1901–2023), TPDC [data set], <https://doi.org/10.11888/Meteoro.tpdc.270961>, 2019.
- Peng, S.: 1-km monthly precipitation dataset for China (1901–2023), TPDC [data set], <https://doi.org/10.5281/zenodo.3114194>, 2020.
- Peng, S., Gang, C., Cao, Y., and Chen, Y.: Assessment of climate change trends over the Loess Plateau in China from 1901 to 2100, *Int. J. Climatol.*, <https://doi.org/10.1002/joc.5331>, 2017a.
- Peng, S., Ding, Y., Wen, Z., Chen, Y., Cao, Y., and Ren, J.: Spatiotemporal change and trend analysis of potential evapotranspiration over the Loess Plateau of China during 2011–2100, *Agr. Forest Meteorol.*, 233, 183–194, <https://doi.org/10.1016/j.agrformet.2016.11.129>, 2017b.
- Peng, S., Ding, Y., Liu, Z., and Li, Z.: 1 km monthly temperature and precipitation dataset for China from 1901 to 2017, *Earth Syst. Sci. Data*, 11, 1931–1946, <https://doi.org/10.5194/essd-11-1931-2019>, 2019.
- Peng Cheng Laboratory: Star Cloud Data Service Platform, <https://data-starcloud.pcl.ac.cn/resource/54> (last access: 29 September 2024), 2024.
- Ramosaj, B. and Pauly, M.: Consistent estimation of residual variance with random forest Out-Of-Bag errors, *Stat. Probabil. Lett.*, 151, 49–57, <https://doi.org/10.1016/j.spl.2019.03.017>, 2019.
- Roujean, J. L. and Breon, F. M.: Estimating PAR absorbed by vegetation from bidirectional reflectance measurements, *Remote Sens. Environ.*, 51, 375–384, [https://doi.org/10.1016/0034-4257\(94\)00114-3](https://doi.org/10.1016/0034-4257(94)00114-3), 1995.

- Shimizu, K., Ota, T., and Mizoue, N.: Detecting Forest Changes Using Dense Landsat 8 and Sentinel-1 Time Series Data in Tropical Seasonal Forests, *Remote Sens.*, 11, 1899, <https://doi.org/10.3390/rs11161899>, 2019.
- Sulla-Menashe, D., Gray, J. M., Abercrombie, S. P., and Friedl, M. A.: Hierarchical mapping of annual global land cover 2001 to present: The MODIS Collection 6 Land Cover product, *Remote Sens. Environ.*, 222, 183–194, <https://doi.org/10.1016/j.rse.2018.12.013>, 2019.
- Tang, X., Bullock, E. L., Olofsson, P., Estel, S., and Woodcock, C. E.: Near real-time monitoring of tropical forest disturbance: New algorithms and assessment framework, *Remote Sens. Environ.*, 224, 202–218, <https://doi.org/10.1016/j.rse.2019.02.003>, 2019.
- Tucker, C. J.: Red and photographic infrared linear combinations for monitoring vegetation, *Remote Sens. Environ.*, 8, 127–150, [https://doi.org/10.1016/0034-4257\(79\)90013-0](https://doi.org/10.1016/0034-4257(79)90013-0), 1979.
- Vermote, E.: MODIS/Terra Surface Reflectance 8-Day L3 Global 500 m SIN Grid V061, NASA EOSDIS Land Processes Distributed Active Archive Center [data set], <https://doi.org/10.5067/MODIS/MOD09A1.061>, 2021.
- Vermote, E. F., Roger, J. C., and Ray, J. P.: MODIS Surface Reflectance User's Guide, USGS, https://lpdaac.usgs.gov/documents/306/MOD09_User_Guide_V6.pdf (last access: 18 November 2024), 2015.
- Wang, Y., Sun, J., He, W., Ye, C., Liu, B., Chen, Y., Zeng, T., Ma, S., Gan, X., Miao, C., Zhou, H., and Tsunekawa, A.: Migration of vegetation boundary between alpine steppe and meadow on a century-scale across the Tibetan Plateau, *Ecol. Indic.*, 136, 108599, <https://doi.org/10.1016/j.ecolind.2022.108599>, 2022.
- Wang, Y., Xue, K., Hu, R., Ding, B., Zeng, H., Li, R., Xu, B., Pang, Z., Song, X., Li, C., Du, J., Yang, X., Zhang, Z., Hao, Y., Cui, X., Guo, K., Gao, Q., Zhang, Y., Zhu, J., Sun, J., Li, Y., Jiang, L., Zhou, H., Luo, C., Zhang, Z., Gao, Q., Chen, S., Ji, B., Xu, X., Chen, H., Li, Q., Zhao, L., Xu, S., Liu, Y., Hu, L., Wu, J., Yang, Q., Dong, S., He, J., Zhao, X., Wang, S., Piao, S., Yu, G., and Fu, B.: Vegetation structural shift tells environmental changes on the Tibetan Plateau over 40 years, *Sci. Bull.*, 68, 1928–1937, <https://doi.org/10.1016/j.scib.2023.07.035>, 2023.
- Wang, Z., Wang, Q., Zhao, L., Wu, X., Yue, G., Zou, D., Nan, Z., Liu, G., Pang, Q., Fang, H., Wu, T., Shi, J., Jiao, K., Zhao, Y., and Zhang, L.: Mapping the vegetation distribution of the permafrost zone on the Qinghai–Tibet Plateau, *J. Mt. Sci.*, 13, 1035–1046, <https://doi.org/10.1007/s11629-015-3485-y>, 2016.
- Wu, F., Ren, H., and Zhou, G.: The 30 m vegetation maps from 1990 to 2020 in the Tibetan Plateau, *Sci. Data*, 11, 804, <https://doi.org/10.1038/s41597-024-03649-7>, 2024.
- Xian, G. Z., Smith, K., Wellington, D., Horton, J., Zhou, Q., Li, C., Auch, R., Brown, J. F., Zhu, Z., and Reker, R. R.: Implementation of the CCDC algorithm to produce the LCMAP Collection 1.0 annual land surface change product, *Earth Syst. Sci. Data*, 14, 143–162, <https://doi.org/10.5194/essd-14-143-2022>, 2022.
- Xiao, X., Hollinger, D., Aber, J., Goltz, M., Davidson, E. A., Zhang, Q., and Moore III, B.: Satellite-based modeling of gross primary production in an evergreen needleleaf forest, *Remote Sens. Environ.*, 89, 519–534, <https://doi.org/10.1016/j.rse.2003.11.008>, 2004.
- Xu, H.: A new index for delineating built-up land features in satellite imagery, *Int. J. Remote Sens.*, 29, 4269–4276, <https://doi.org/10.1080/01431160802039957>, 2008.
- Yang, J. and Huang, X.: The 30 m annual land cover dataset and its dynamics in China from 1990 to 2019, *Earth Syst. Sci. Data*, 13, 3907–3925, <https://doi.org/10.5194/essd-13-3907-2021>, 2021.
- Yang, J. and Huang, X.: The 30 m annual land cover datasets and its dynamics in China from 1985 to 2022, Zenodo [data set], <https://doi.org/10.5281/zenodo.8176941>, 2023.
- Yao, T., Thompson, L. G., Mosbrugger, V., Zhang, F., Ma, Y., Luo, T., Xu, B., Yang, X., Joswiak, D. R., Wang, W., Joswiak, M. E., Devkota, L. P., Tayal, S., Jilani, R., and Fayziev, R.: Third pole environment (TPE), *Environ. Dev.*, 3, 52–64, <https://doi.org/10.1016/j.envdev.2012.04.002>, 2012.
- Yu, B., Lv, C., Lv, T., Yang, A., and Liu, C.: Datasets of the boundary and area of the Tibetan Plateau, *Acta Geogr. Sin.*, 69, 65–68, <https://doi.org/10.11821/dlxb2014S012>, 2014.
- Zha, Y., Gao, J., and Ni, S.: Use of normalized difference built-up index in automatically mapping urban areas from TM imagery, *Int. J. Remote Sens.*, 24, 583–594, <https://doi.org/10.1080/01431160304987>, 2003.
- Zhang, C., Zhang, H., and Tian, S.: Phenology-assisted supervised paddy rice mapping with the Landsat imagery on Google Earth Engine: Experiments in Heilongjiang Province of China from 1990 to 2020, *Comput. Electron. Agric.*, 212, 108105, <https://doi.org/10.1016/j.compag.2023.108105>, 2023.
- Zhang, G., Luo, W., Chen, W., and Zheng, G.: A robust but variable lake expansion on the Tibetan Plateau, *Sci. Bull.*, 64, 1306–1309, <https://doi.org/10.1016/j.scib.2019.07.018>, 2019a.
- Zhang, H., Zhao, C., and Zhu, W.: A new vegetation map for Qinghai–Tibet Plateau by integrated classification from multi-source data products (2020), TPDC [data set], <https://doi.org/10.11888/Terre.tpd.271997>, 2021a.
- Zhang, J., Okin, G. S., and Zhou, B.: Assimilating optical satellite remote sensing images and field data to predict surface indicators in the Western US: Assessing error in satellite predictions based on large geographical datasets with the use of machine learning, *Remote Sens. Environ.*, 233, 111382, <https://doi.org/10.1016/j.rse.2019.111382>, 2019b.
- Zhang, L., Shen, M., Yang, Z., Wang, Y., and Chen, J.: Spatial variations in the difference in elevational shifts between greenness and temperature isolines across the Tibetan Plateau grasslands under warming, *Sci. Total Environ.*, 906, 167715, <https://doi.org/10.1016/j.scitotenv.2023.167715>, 2024a.
- Zhang, X., Liu, L., Chen, X., Gao, Y., Xie, S., and Mi, J.: GLC_FCS30: global land-cover product with fine classification system at 30 m using time-series Landsat imagery, *Earth Syst. Sci. Data*, 13, 2753–2776, <https://doi.org/10.5194/essd-13-2753-2021>, 2021b.
- Zhang, X., Liu, L., Chen, X., Xie, S., and Gao, Y.: Fine land-cover mapping in China using Landsat datacube and an operational SPECLib-based approach, *Remote Sens.*, 11, 1056, <https://doi.org/10.3390/rs11091056>, 2019c.
- Zhang, X., Zhao, T., Xu, H., Liu, W., Wang, J., Chen, X., and Liu, L.: GLC_FCS30D: the first global 30 m land-cover dynamics monitoring product with a fine classification system for the period from 1985 to 2022 generated using dense-time-series Landsat imagery and the continuous change-detection method, *Earth Syst. Sci. Data*, 16, 1353–1381, <https://doi.org/10.5194/essd-16-1353-2024>, 2024b.
- Zhang, Y.: Integration dataset of Tibet Plateau boundary, TPDC [data set], <https://doi.org/10.11888/Geogra.tpd.270099>, 2019.

- Zhang, Y., Li, B., Liu, L., and Zheng, D.: Redetermine the region and boundaries of Tibetan Plateau, *Geogr. Res.*, 40, 1543–1553, <https://doi.org/10.11821/dlyj020210138>, 2021c.
- Zhang, Y., Liu, L., Li, B., and Zheng, D.: Boundary Data of the Tibetan Plateau (2021 Version), *Digital Journal of Global Change Data Repository* [data set], <https://doi.org/10.3974/geodb.2021.07.10.V1>, 2021d.
- Zhang, Y., Liu, L., Li, B., and Zheng, D.: Comparison of boundary datasets covering Tibetan Plateau between 2021 and 2014 versions, *J. Global Change Data Discov.*, 5, 322–332, <https://doi.org/10.3974/geodp.2021.03.10>, 2021e.
- Zhou, G., Ren, H., Liu, T., Zhou, L., Ji, Y., Song, X., and Lv, X.: Vegetation map of Qinghai Tibet Plateau in 2020 with 10 m spatial resolution, TPDC [data set], <https://doi.org/10.11888/Terre.tpd.272408>, 2022a.
- Zhou, G., Ren, H., Liu, T., Zhou, L., Ji, Y., Song, X., and Lv, X.: A new regional vegetation mapping method based on terrain-climate-remote sensing and its application on the Qinghai-Xizang Plateau, *Sci. China Earth Sci.*, 66, 237–246, <https://doi.org/10.1007/s11430-022-1006-1>, 2023.
- Zhou, G., Ren, H., Zhang, L., Lv, X., and Zhou, M.: 500 m annual vegetation maps of Qinghai Tibet Plateau (2000–2022), TPDC [data set], <https://doi.org/10.11888/Terre.tpd.301205>, 2024.
- Zhou, J., Zheng, Y., Song, C., Cheng, C., Gao, P., Shen, S., and Ye, S.: Vegetation map of Qinghai Tibet Plateau in 1980s, TPDC [data set], <https://doi.org/10.11888/Terre.tpd.272385>, 2022b.
- Zhu, Z. and Woodcock, C. E.: Continuous change detection and classification of land cover using all available Landsat data, *Remote Sens. Environ.*, 144, 152–171, <https://doi.org/10.1016/j.rse.2014.01.011>, 2014.
- Zhu, Z., Zhang, J., Yang, Z., Aljaddani, A. H., Cohen, W. B., Qiu, S., and Zhou, C.: Continuous monitoring of land disturbance based on Landsat time series, *Remote Sens. Environ.*, 238, 111116, <https://doi.org/10.1016/j.rse.2019.03.009>, 2019.