



Extending the late 1963 to 1964 Mt Agung rescued searchlight aerosol profiles dataset at 32°N, from early 1963 to 1975

Juan Carlos Antuña-Marrero^{1,2,3}, Abel Calle², Juan Antonio Añel³, Victoria Cachorro², Laura de la Torre³, David Barriopedro¹, Ricardo García Herrera^{1,4}, and Javier Pacheco²

¹Instituto de Geociencias (IGEO), CSIC-UCM, Madrid 28040, Spain

²Grupo de Óptica Atmosférica (GOA-UVA), Universidad de Valladolid, Valladolid 47011, Spain

³EPhysLab, CIM-UVigo, Universidad de Vigo, 32004, Ourense, Spain

⁴Departamento de Física de la Tierra y Astrofísica, Universidad Complutense de Madrid, 28040, Madrid, Spain

Correspondence: Juan Carlos Antuña-Marrero (jcam45@gmail.com)

Received: 10 February 2026 – Discussion started: 2 April 2026

Revised: 12 June 2026 – Accepted: 18 June 2026 – Published: 6 July 2026

Abstract. A set of 11 aerosol turbidity profiles (ATP) and 2 aerosol extinction profiles (AEP) at $\lambda = 0.55 \mu\text{m}$, observed with searchlight in New Mexico at 32°N, has been digitized from plots in scientific articles. They cover the period February to June 1963 and September 1965 to May 1975, complementing the already rescued and previously published 105 individual AEP, corresponding to 36 days, between December 1963 and December 1964. Eleven AEP are calculated (AEPc) from the ATP, and the corresponding stratospheric aerosol optical depth (sAOD) between 12 and 25 km is also derived. Estimates of the digitization errors for the AEPc and the sAOD are also calculated using information available in the literature. The combined set of rescued AEP reported here and the earlier rescued set of AEP from searchlight observations, are the only AEP dataset covering the period between the 1963 Mt Agung and the 1974 Fuego eruptions at northern midlatitudes. In this regard two relevant features identified in the AEP and the sAOD are described here. The first, using AEPc from March and April 1963 identified what could be the date of arrival of the stratospheric aerosols from the Mt. Agung first eruption on 17 March 1963. This would challenge the accepted criteria that the stratospheric aerosols from Mt Agung arrived at the northern hemisphere midlatitudes in the second half of 1963. The second feature evidences two anomalous increases of the sAOD during a period supposed to be the decay of the sAOD from Mt. Agung eruption. They show our limited knowledge and understanding of the 1963 Mt Agung volcanic stratospheric aerosol transport. Finally, we describe evidences found in the literature pointing to the possible existence of the original searchlight raw signals and its processing software. The dataset described in this work is available at: <https://doi.org/10.1594/PANGAEA.992616> (Antuña-Marrero et al., 2026a).

1 Introduction

The 1963 Mt Agung eruptions, Bali, Indonesia at 8.3° S, the third biggest volcanic eruption of the second half of the 20th century, consisted in a series of eruptions (Fontijn et al., 2015). Two of these eruptions, on 17 March and 16 May, injected 4.7 and 2.3 Tg SO₂ into the stratosphere respectively (Self and Rampino, 2012). Before 1979, the so-called pre-satellite era, the only atmospheric observational dataset of

stratospheric aerosols (SA) used by the Coupled Model Intercomparison Project (CMIP) is derived using an aerosol microphysical model with the input ice core data (Gao et al., 2008) and two datasets of SA optical depth (sAOD) (Arfeuille et al., 2014). The two sAOD observations sets are the one from Sato et al. (1993), mainly based on astronomical observations and summarized by Dyer and Hicks (1968), and the one from Stothers (2001), based on observations of atmospheric attenuation of starlight and direct sunlight.

The mentioned datasets are the only used; however, other relevant data sources exist and could be relevant for research purposes. Since the later years of the 50's of the 20th century, aircrafts and balloons conducted observations of the particle size distributions and chemical composition of the upper troposphere and SA (Antuña-Marrero et al., 2026b). Furthermore, vertical profiles of the upper tropospheric and SA backscattering/extinction were observed with lidars/searchlight (e.g., Fiocco and Grams, 1964; Clemesha et al., 1966; Elterman, 1975). Although the results of the analysis of information generated by those observational datasets were published long time ago, and can be found in the literature, the datasets have not been available until the present.

This issue has been one of the focuses of attention and discussions by the Stratospheric Sulfur and its Role in Climate (SSiRC), recently renamed Stratospheric Aerosol Activity, an established APARC (Atmospheric Processes And their Role in Climate) activity, with APARC being a core project within the World Climate Research Program (WCRP). The implementation of the now Stratospheric Aerosol Data Rescue Activity (https://www.aparc-stratosphericaerosol.org/activities/data_rescue, last access: 18 March 2025) in 2020 began addressing this issue. This activity has already rescued five datasets: four stratospheric lidar aerosol backscattering profiles (ABP) datasets and one searchlight aerosol extinction profile (AEP). The first four lidar ABP consist of two lidar datasets from two northern hemisphere lidar sites at Lexington, MA, USA, at 42° N, and Fairbanks, AK, USA, at 64° N after the 1963 Mt Agung, Indonesia, eruptions (Antuña-Marrero et al., 2021). The other two have been observed by lidars onboard two Soviet research ships conducting transects of the Atlantic Ocean after the 1991 Mt Pinatubo eruption (Antuña-Marrero et al., 2020a). Those four stratospheric ABP and their calculated stratospheric AEP have been stored respectively at the PANGAEA open access data repository (Antuña-Marrero et al., 2020b, c). The fifth dataset consists of searchlight tropospheric and stratospheric AEP from White Sands, NM, USA, at 32° N, after the Mt Agung eruption, covering the period December 1963 to December 1964 (both inclusive) that have been rescued and recalibrated (Antuña-Marrero et al., 2024). The original and the recalibrated tropospheric and stratospheric searchlight AEP have been stored at PANGAEA (Antuña-Marrero et al., 2023) like the other four stratospheric lidar AEP datasets cited above.

Two recent studies illustrate rescued lidar data applications. The first conducted by NASA scientists developing the Global Space-based Stratospheric Aerosol Climatology (GloSSAC), reported the recovering and archiving the SA backscatter profiles from five airborne lidar missions flown between July 1982 and January 1984 (https://asdc.larc.nasa.gov/project/NASA-Airborne-Lidar/NASA_Airborne_Lidar_Flights_1, last access: 29 May 2025). They were used, together with the SA profiles from the NASA 48-inch ground-based lidar data

and aerosol extinction profiles from SAM II, to fill the SAGE I/II gap from 1982 to 1984 (Thomason et al., 2018). The NASA Langley airborne missions data from July 1991 and May 1992, were also archived (https://cmr.earthdata.nasa.gov/virtual-directory/collections/C3880594343-LARC_CLOUD/temporal, last access: 29 June 2025) for future GloSSAC improvements (Thomason et al., 2018). The second application, used the January 1964 to August 1965 ground-based lidar datasets from Lexington (Antuña-Marrero et al., 2020c; 2021) to evaluate the vertical extent of the Agung simulated volcanic SA (Dhomse et al., 2020). The article describing the development of GloSSAC also mention the existence of historical unexploited sources of data and considers worthwhile the potential efforts for its recovery (Thomason et al., 2018). A recent article about the stratospheric aerosol forcing datasets to be provided by the Coupled Model Intercomparison Project-phase 7 (CMIP7) to the participating modelling teams, mention that the five rescued datasets by the APARC Stratospheric Aerosol Activity Data Rescue could be considered to be incorporated to the aerosol optical properties dataset (Aubry et al., 2026).

Here we report the rescue of the available information to complete the New Mexico searchlight tropospheric and stratospheric AEP dataset with the available information from the scientific literature. The new searchlight AEP rescued dataset covers the periods February to June 1963 and September 1965 to May 1975, before and after the December 1963 to December 1964 already rescued searchlight dataset (Antuña-Marrero et al., 2023, 2024). As a result, the combination of the earlier rescued searchlight dataset and the one we report here, produces the only AEP dataset covering February 1963 to May 1975, providing unique information on the SA transported to the northern hemisphere midlatitudes from the 1963 Mt Agung as well as the Fuego eruption in Guatemala (14.5° N), on 14 and 17 October 1974.

2 The searchlight tropospheric and stratospheric AEP observations at New Mexico 1963 to 1975

The US Air Force Cambridge Research Laboratories (AFCRL) carried out a program of tropospheric and stratospheric AEP observations with searchlight between 1963 and 1975 at White Sands (32° N), New Mexico. The cited program included the development and implementation of the processing algorithm to calculate the AEP (Elterman, 1966a, b). This series of AEPs include profiles under both SA background and under perturbed volcanic conditions, particularly from the eruptions of 1963 Mt Agung and Fuego in Guatemala in October 1974 (Elterman, et al., 1973; Elterman, 1966a, 1975, 1976).

Between February 1963 and May 1975, at least 359 individual tropospheric and stratospheric AEPs were collected, in many cases more than one in the same night (listed in Table S1 in the Supplement). One hundred five of them, carried

on between December 1963 to December 1964, were made public in form of individual AEP plots at an AFCRL report (Elterman, 1966b). This subset of searchlight AEPs has already been rescued (Antuña-Marrero et al., 2024). The other, at least 254, single AEPs are still missing, and they consist of 9 AEPs collected between February and June 1963 and 245 AEPs collected between September 1965 and May 1975.

The source of the rescued AEPs presented here, are the aerosol turbidity profiles (ATPs) reported by Elterman et al. (1973) and Elterman (1975, 1976). $ATP(z)$ were calculated using Eq. (1) where $AEP(z)$ is the original aerosol extinction profile and $REP(z)$ is the Rayleigh (molecular) extinction profile at $\lambda = 0.55 \mu\text{m}$, tabulated in Elterman (1964). The ATP averages at monthly, bi-monthly and multi-monthly temporal scales, were calculated and published in plots in the above cited articles.

$$ATP(z) = \frac{AEP(z)}{REP(z)} \quad (1)$$

2.1 ATP and AEP plots: its digitization and ATP conversion to AEP

In total 16 profiles, fourteen ATPs and two AEPs, were reported in Elterman et al. (1973) and in Elterman (1975, 1976). Within the current study 11 ATP and 2 AEP have been digitized from plots contained in the three cited papers, covering several temporal scales: three single daily profiles, three monthly, two bi-monthly and three multi-monthly averaged profiles. Three of the ATPs have not been digitized, by reasons listed in Table S2 in the Supplement. The 11 ATPs and 2 AEPs plots, and consequently the corresponding digitized profiles, mainly contain information between 10 to 26 km, (Table S3). The plot digitization was conducted using the WebPlotDigitizer software (Rohatgi, 2024), at the 0.5 km resolution, producing the digitized aerosol transmission profiles (ATPd) and digitized aerosol extinction profiles (AEPd) from the plotted ATP and AEP respectively. Then the ATPd were interpolated to 1 km resolution to match the resolution of the $REP(z)$ from Table 5.11 in Elterman (1964), followed by the calculation of the corresponding AEP using Eq. (1). The final step was interpolating those calculated AEP to produce the final AEP (AEPc) at the searchlight instrumental vertical resolution of 58 levels in the altitude range from 2.76 to 35.28 km (Elterman, 1966b).

The AEPc are listed in Table 1, where the first column contains the date and, for the single day profiles, the hour the observation was conducted, in Mountain Standard Time (MST), UTC-7. The second column reports the numbers of single raw searchlight profiles used in the respective calculations. The superscripts in the date column identify the profiles having both digitized ATP and AEP: a single daily profile (11 November 1974) and a bi-monthly mean profile (October and November 1970). Those digitized AEP and the corresponding AEPc were used to estimate the AEPc uncertainties, attributed to its determination from the ATP, described

Table 1. Listing of the original eleven ATP plots, the number of single profiles reported for each one and its respective source article, and the number of the Figure source on it. In the case of existing several profiles in the same Figure, the individual profile identifiers are included after the figure number. Superscripts ¹ and ² identify a single daily and a bi-monthly mean profile that have both their AEP and ATP plotted and were digitized.

Single Daily (Hour MST)	Profiles	Source
28 March 1963 (02:36)	1	Elterman et al. (1973); Fig. 3
20 April 1963 (23:45)	1	Elterman et al. (1973); Fig. 3
11 November 1974 ¹ (22:17)	1	Elterman (1975); Fig. 2
Monthly mean		
November 1974	12	Elterman (1976); Fig. 1b
January 1975	10	Elterman (1976); Fig. 1c
May 1975	19	Elterman (1976); Fig. 1d
Bi-Monthly mean		
February–March 1963	3	Elterman et al. (1973); Fig. 2G1
October–November 1970 ²	41 (9)	Elterman et al. (1973); Fig. 2G7
Multi-Monthly mean		
April–June 1963	6	Elterman et al. (1973); Fig. 2G2
September 1965–January 1966	50	Elterman et al. (1973); Fig. 2G6
September 1973–July 1974	113	Elterman (1976); Fig. 1a

Table 2. Idem Table 1, but for the two AEP plotted. Superscript ¹ and ² identify a single daily and a bi-monthly mean profile that have both their AEP and ATP plotted and were digitized.

Single Daily (Hour MST)	Profiles	Source
11 November 1974 ¹ (22:17)	1	Elterman (1975); Fig. 1
Bi-Monthly mean profile		
October–November 1970 ²	41 (9)	Elterman et al. (1973); Fig. 1

in the following section. The number in brackets after the number of profiles for the bi-monthly mean profile October–November 1970 is the number of days the observations were conducted, nine days. It is the only case where that information is available.

In Table 2 the two AEP digitized (AEPd), complementing the ATP information for the same periods in Table 1. They will be used below to validate the results of the AEP calculation from the digitized ATPd.

2.2 Estimated errors

The estimated errors for the originally determined AEP, between 15 % and less than 25 %, were reported by Wells (1968). Three error sources were only considered for this estimate: neglecting both the multiple scattering from the searchlight beam and the ozone absorption, and for use of the Reeger–Siedentopf aerosol phase function, measured at the surface in Germany (at 50° N) during a period when no volcanic aerosols were in the stratosphere (Reeger and Sieden-

topf, 1946). The error associated to the first two sources were determined to be negligible, with the aerosol phase function accounting for the all the error. A recent error estimation considered all the significant error sources, conducted for same originally determined AEP, rescued from tables in Elterman (1966b) for the period December 1963 to December 1964, increase the former range to between 69 % and 84 %. The same estimate was conducted for the recalibrated AEP dataset resulted in ranges between 54 % and 69 % for the recalibrated tropospheric AEP and between 40 % and 55 % (Antuña-Marrero et al., 2024).

2.2.1 Estimated errors of the ATPd and the calculated sAOD

Table 3 shows the quantitative information reported in Tables 1 and 2 in Elterman et al. (1973). The first column list date (period) from a daily profile (average for the period). The second reports the maximum (mean maximum) turbidity value for the respective ATP. The third shows the sAOD (mean sAOD) in the 12 to 25 km layer calculated from the corresponding AEP. Column four reports the maximum (mean maximum) ATPd and the five the sAOD (mean sAOD) in the 12 to 25 km layer, calculated from the AEPc reported by this study. Columns two and three are the only tabulated available information about some of the ATP and sAOD that we have rescued. We use it to get an estimate of the errors associated to the digitization of the ATP plot. To do it, we used the mean relative error (MRE), for each of the variables defined by the equation:

$$\text{MRE} = \frac{1}{N} \sum_1^N \frac{V_c - V_r}{V_r} \times 100 \quad (2)$$

Where V_c is the the ATPd value in column 2 (sAOD value in column 3), and V_r is the reference ATP in column 3 (sAOD value in column 4). The MRE for the turbidity maximum is in the order of 2 %, providing information about the error introduced by the digitization procedure. In the case of the sAOD the MRE is 5 %, a reasonable value considering the magnitude of the digitization error and the fact that the sAOD is the integral of the AEPc between 12 and 25 km. Those values are in the same order of magnitude than the 1.2 % reported in the digitization of the plots of the AEP from December 1963 to December 1964, thus making the rescued dataset of AEP reported here consistent with the AEP reported in Antuña-Marrero et al. (2024).

2.2.2 Estimated error of the AEPc

To estimate the MRE in the current set of 11 AEPc calculated from the ATPd with Eq. (1), we use the 11 November 1974 at 22:17 MST digitized AEPd and ATPd, from Figs. 1 and 2 respectively in Elterman (1975). We also use the October and November 1970 average AEPd and ATPd, from Figs. 1 and 2, respectively, in Elterman et al. (1973). Figure 1 shows




Figure 1. Plot of the digitized and calculated AEP from the 11 November 1974 at 22:17 MST and October–November 1970 average profiles, reported in plots in Elterman (1975) and Elterman et al. (1973) respectively. Both pairs of digitized AEPd and AEPc show good agreements.

in the left panel the plot of the daily AEPc and AEPd for 11 November 1974 and in the right panel for the AEPc and AEPd averages for the period October–November 1970. In both cases a very good match is appreciated, a feature supported by the MRE values of -1.03% and -1.05% respectively. Those values are in the same order of magnitude than the MRE associated to the digitization of the ATP. The very low magnitude of the MRE allows to consider that the magnitudes of the AEPc using Eq. (1) match the original AEP, that was the primary source to calculate the ATP reported both in Elterman et al. (1973) and Elterman (1976).

3 Some volcanic SA features present in the 1963–1975 rescued searchlight AEP and sAOD series

The discussion and analysis of the two features described in this section are beyond the purpose of this paper focussed on describing the rescued dataset, the methods applied for that purpose and guarantee the dataset storage in an open access data repository. However, they demonstrate applications of the rescued dataset we report. The first, showing the cross hemispheric of volcanic SA across the equator, provides information and at the same motivates future research to address our current limited knowledge and understanding of the 1963 Mt Agung volcanic stratospheric aerosol transport. The second, provides unique and consistent AEP and sAOD information from two volcanic events in the pre – satellite era, with the same ground-based instrument and processing algorithm.

Table 3. Values of the mean maximum turbidity and the sAOD between 12 and 25 km, reported by Elterman et al. (1973) in columns 2 and 3 respectively. Columns 4 and 5 report the maximum turbidity value and the sAOD derived from the digitized ATP and the calculated AEP respectively.

Date/Period	Elterman et al. (1973)		ATPd and AEPc	
	Mean Max. Turbidity	Mean sAOD	Maximum Turbidity	sAOD
February–March 1963	1.4	1.8×10^{-2}	1.5	2.0×10^{-2}
28 March 1963	1.4	1.9×10^{-2}	1.4	1.9×10^{-2}
20 April 1963	4.5	3.9×10^{-2}	4.3	4.0×10^{-2}
April–June 1963	3.4	3.1×10^{-2}	3.1	3.3×10^{-2}
September 1965–January 1966	1.9	2.2×10^{-2}	1.8	2.3×10^{-2}
October–November 1970	1.5	2.0×10^{-2}	1.5	1.9×10^{-2}




Figure 2. Plots of the digitized ATP (left panel) and the calculated AEP (right panel) for 28 March and 20 April 1963.

3.1 The transport of the SA from the 1963 Mt Agung eruption from 8°S to 32°N

The ATPs derived from searchlight observations on 28th and April 20th 1963, reported in Fig. 3 in Elterman et al. (1973), identified the arrival of the Mt Agung SA to White Sands, NM. Figure 2 shows on panel (a) both ATPs digitized and in panel (b) the AEPc from the former ones. Those two, as well as the other two AEPs, consisting in the average of 3 profiles taken in January and February 1963 and the average of 6 profiles between April and June 1963 provided unique information about the time, vertical distribution and stratospheric AOD of the SA from the first, 17 March 1963, Mt. Agung eruption.

The two daily AEP from 1963 in Fig. 2 allowed Elterman et al. (1973) to specify with a margin of error of $\sim \pm 10$ d the date of arrival at White Sands of the SA cloud from the Agung eruption. Furthermore, both profiles allowed to quantify the values of the sAOD of 0.019 and 0.040 before and after the arrival of the SA cloud respectively, listed in Table 3 above.

3.2 Series of sAOD observations at Sacramento Peak, New Mexico, 1963 to 1975

The series of sAOD calculated between 12 and 25 km, combining the 36 d observations between December 1963 and December 1964 already rescued (Antuña-Marrero et al., 2024), and the observations of the sAOD calculated from the 11 rescued AEPc reported here are shown in Fig. 3. The 4 AEPc from February to June 1963, depict the arrival to 32° N of the SA from the first, 17 March, Mt Agung eruption. It was identified and reported by Elterman et al. (1973). The magnitude of 0.040 of the sAOD in 20 April is among the highest registered in the joint sAOD series, with only three observations showing higher sAOD magnitudes, all of them in the first half of 1964: 14 and 16 February, with sAOD of 0.048 and 0.045 respectively and 0.050 on 12 June 1964. It should be emphasized that the cited sAOD value on 12 June, together with the increase in the magnitudes of the four daily AEPs observed from 9 to 12 June 1964, represents an anomaly in the expected decay state of the SA from the Mt Agung eruptions on 17 March and 16 May 1963. It could be associated to one of the eruptions registered in the northern hemisphere with a Volcanic Explosive Index (VEI) equal or higher than 3. A similar situation is present also for Mt Agung 1963 eruption, but in the rescued lidar record at Lexington, MA, at 42° N. There the sAOD from lidar AEPs shows peaks with magnitudes inconsistent with the SA decay in November and December 1964 and between March and July 1965 (Antuña-Marrero et al., 2021). Further research is necessary to identify the sources of SA in all those cases.

The final section of the AEP and the sAOD series provides information about almost a year before the 14 and 16 October 1974 Fuego eruptions, and also after it. The volcanic signal is present in the AEP and sAOD on 11 November 1974, as well as in the monthly mean values of the AEP and sAOD for November 1974, January and May 1975. The high sAOD magnitude between 1959 and 1962 were reported on contemporary observations. In the tropics, positive anomalies of the Linke turbidity coefficient between 1958 and 1962 were reported at Mauna Loa Observatory, before the 1963 Agung




Figure 3. Joint SA optical depth (sAOD) of the 36 daily AEP from December 1963 to December 1964 already rescued (blue stars), and the 11 profiles (in red). The different markers used for the rescued sAOD represent their variable temporal resolutions. The red asterisks are the sAOD from daily single profiles: 28 March 1963, 20 April 1963 and 11 November 1974. The red diamonds are the sAOD of the monthly averaged AEP profiles for November 1974, January and May 1975. The sections of discontinuous line represent the sAOD of the multi-monthly averaged AEP profiles, whose length is determined by the 15th days of the beginning and ending months of the averages for February–March and April–June 1963, September 1965–January 1966, October–November 1970 and September 1973–July 1974.

eruption (Ellis and Pueschel, 1971). At northern midlatitudes, the ratio of observed scattering to Rayleigh scattering at 660 nm from twilight measurements at Blue Hill Observatory, conducted during the winter of 1959–1960, are higher than 3 between 10 and 20 km in Fig. 2 from Volz (1964). The high sAOD values were attributed to the number of nuclear explosions in the atmosphere during that period that peaked between 1960 and 1962 (Kondratyev et al., 1974).

4 Data availability

Data described in this work is available at <https://doi.org/10.1594/PANGAEA.992616> (Antuña-Marrero et al., 2026a).

5 Summary and outlook

The present work is an important step forward to determine and to understand the temporal evolution of the SA's optical properties from the 1963 Mt Agung, the 1974 El Fuego volcanic eruptions and the background periods around them, in the northern hemisphere midlatitudes. However, as we pointed out above, the number of AEPs rescued represent less than half of the AEP searchlight observations conducted

between 1963 and 1975. Those still missing AEPs will contribute to a deeper understanding of the SA evolution.

The search for the original searchlight ABPs and its original processing algorithm continues. The search in the available literature, revealed that the searchlight ABPs were processed by computer (Elterman, 1966a), using the processing output on punched cards to produce AEPs and ATPs plots and few partial tabulations that were published in articles and reports. Four software applications for the searchlight ABP processing, were reported to be implemented and stored at the Analysis and Simulation Branch of the Computation Centre at the AFCRL; at the time all the US government computer facilities conducted the transition from the storage of data and software in punched cards to magnetic tapes (Cronin, 1972). So far, the search for those digital records has been unsuccessful, but we will not give up.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/essd-18-4669-2026-supplement>.

Author contributions. JCAM, AC and VC contributed designing and leading the data rescue procedure. JP conducted the digitalization of the turbidity/aerosol profiles from the plots and their preliminary processing. All co-authors contributed to either advising the data recovery, the processing and analysis of the rescued data. RGH and JAA conducted detailed reviews of the paper final version.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. To the memory of Louis Elterman and the team he led for their extraordinary achievements both technologically and scientifically. We also thank the support from the Library of the Universidade de Vigo and the Library of Faculty of Physics of the Complutense University of Madrid for their contributions in the search, location and acquisition of scientific literature. We thank Jeannette van den Bosch and the NASA JSC White Sands Test Facility for their effort in establishing and maintaining AERONET White_Sands_HELSTF site. Juan Carlos Antuña Marrero recognizes the support from the Optics Atmospheric Group, the Theoretical, Atomic, and Optical Physics Department, and the Applied Physics Department of University of Valladolid, Spain. Juan Carlos Antuña Marrero also recognizes the support from the EPhysLab of the University of Vigo at Ourense.

Financial support. This research work was supported by the Ministry for the Ecological Transition and the Demographic Challenge (MITECO) and the European Commission NextGenerationEU (Regulation EU 2020/2094), through CSIC's Interdisciplinary Thematic Platform Clima (PTI-Clima) /Development of Operational Climate Services (Ref. CSC2304000), by the Grant PID2024-158326NB-I00 funded by MICIU/AEI/10.13039/501100011033. The EPhysLab is supported by the Government of Galicia (Grant: ED431C 2025/37).

The article processing charges for this open-access publication were covered in part by the CSIC Open Access Publication Support Initiative through its Unit of Information Resources for Research (URICI).

Review statement. This paper was edited by Graciela Raga and reviewed by Sandip Dhomse and two anonymous referees.

References

- Antuña-Marrero, J.-C., Mann, G. W., Keckhut, P., Avdyushin, S., Nardi, B., and Thomason, L. W.: Shipborne lidar measurements showing the progression of the tropical reservoir of volcanic aerosol after the June 1991 Pinatubo eruption, *Earth Syst. Sci. Data*, 12, 2843–2851, <https://doi.org/10.5194/essd-12-2843-2020>, 2020a.
- Antuña-Marrero, J. C., Mann, G. W., Keckhut, P., Avdyushin, S. I., Nardi, B., and Thomason, L. W.: Ship borne lidar measurements in the Atlantic of the 1991 Mt Pinatubo aerosol cloud, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.912770>, 2020b.
- Antuña-Marrero, J.-C., Mann, G. W., Barnes, J., Rodríguez-Vega, A., Shallcross, S., Dhomse, S., Fiocco, G., and Grams, G. W.: The first ever multi-year lidar dataset of the stratospheric aerosol layer, from Lexington, MA, and Fairbanks, AK, January 1964 to July 1965, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.922105>, 2020c.
- Antuña-Marrero, J.-C., Mann, G. W., Barnes, J., Rodríguez-Vega, A., Shallcross, S., Dhomse, S. S., Fiocco, G., and Grams, G. W.: Recovery of the first ever multi-year lidar dataset of the stratospheric aerosol layer, from Lexington, MA, and Fairbanks, AK, January 1964 to July 1965, *Earth Syst. Sci. Data*, 13, 4407–4423, <https://doi.org/10.5194/essd-13-4407-2021>, 2021.
- Antuña-Marrero, J.-C., Mann, G. W., Barnes, J., Calle, A., Dhomse, S., Cachorro Revilla, V. E., Deshler, T., Zhengyao, L., Sharma, N., and Elterman, L.: Tropospheric and stratospheric aerosol extinction and stratospheric aerosol optical depth from searchlight measurements conducted at White Sands, New Mexico, US between December 1963 and December 1964, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.949377>, 2023.
- Antuña-Marrero, J.-C., Mann, G. W., Barnes, J., Calle, A., Dhomse, S. S., Cachorro, V. E., Deshler, T., Li, Z., Sharma, N., and Elterman, L.: The Recovery and Re-Calibration of a 13-Month Aerosol Extinction Profiles Dataset from Searchlight Observations from New Mexico, after the 1963 Agung Eruption, *Atmosphere*, 15, 635, <https://doi.org/10.3390/atmos15060635>, 2024.
- Antuña-Marrero, J.-C., Calle, A., Añel, J. A., Cachorro, V. E., de la Torre, L., Barriopedro, D., García-Herrera, R., and Pacheco, J.: Rescued searchlight aerosol profiles dataset at New Mexico (32° N), completing the period from early 1963 to 1975, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.992616>, 2026a.
- Antuña-Marrero, J.-C., Calle, A., Cachorro, V. E., Deshler, T., Barriopedro, D., García-Herrera, R., and Muriel, M.: Rescued, digitized and homogenized UTLS aerosol particle size distributions, measured by US projects monitoring radioactive debris from nuclear weapons tests, 1960 to 1964, in preparation, 2026b.
- Arfeuille, F., Weisenstein, D., Mack, H., Rozanov, E., Peter, T., and Brönnimann, S.: Volcanic forcing for climate modeling: a new microphysics-based data set covering years 1600–present, *Clim. Past*, 10, 359–375, <https://doi.org/10.5194/cp-10-359-2014>, 2014.
- Aubry, T. J., Toohey, M., Khanal, S., Chim, M. M., Verkerk, M., Johnson, B., Schmidt, A., Kovilakam, M., Sigl, M., Nicholls, Z., Thomason, L., Naik, V., Rieger, L., Stiller, D., Ziegler, E., Durack, P., and Smith, I. H.: Stratospheric aerosol forcing for CMIP7 – Part 1: optical properties for pre-industrial, historical, and scenario simulations, *Geosci. Model Dev.*, 19, 3725–3756, <https://doi.org/10.5194/gmd-19-3725-2026>, 2026.
- Clemesha, B. R., Kent, G. S., and Wright, R. W. H.: Laser probing of the lower atmosphere, *Nature*, 209, 184–185, <https://doi.org/10.1038/209184a0>, 1966.
- Cronin, E. C.: A Systems Analysis, A Functional Organization; A Customer Users Library (CUL), AFCRL-72-035, 626 pp., <https://apps.dtic.mil/sti/tr/pdf/AD0755389.pdf> (last access: 26 May 2025), 1972.
- Dhomse, S. S., Mann, G. W., Antuña Marrero, J. C., Shallcross, S. E., Chipperfield, M. P., Carslaw, K. S., Marshall, L., Abraham, N. L., and Johnson, C. E.: Evaluating the simulated radiative forcings, aerosol properties, and stratospheric warmings from the 1963 Mt Agung, 1982 El Chichón, and 1991 Mt Pinatubo volcanic aerosol clouds, *Atmos. Chem. Phys.*, 20, 13627–13654, <https://doi.org/10.5194/acp-20-13627-2020>, 2020.
- Dyer, A. J. and Hicks, B. B.: Global spread of volcanic dust from the Bali eruption of 1963, *Q. J. Roy. Meteor. Soc.*, 94, 545–554, <https://doi.org/10.1002/qj.49709440209>, 1968.
- Ellis, H. T. and Pueschel, R. F.: Solar radiation: absence of air pollution trends at Mauna Loa, *Science*, 172, 845–847, <https://10.1126/science.172.3985.845>, 1971.
- Elterman, L.: Atmospheric Attenuation Model in the Ultraviolet, Visible, and Infrared Regions for Altitudes to 50 km, AFCRL-64-740, <https://apps.dtic.mil/sti/pdfs/AD0607859.pdf> (last access: 20 August 2025), 1964.
- Elterman, L.: Aerosol Measurements in the Troposphere and Stratosphere, *Appl. Optics*, 5, 1769–1776, <https://doi.org/10.1364/AO.5.001769>, 1966a.
- Elterman, L.: An Atlas of Aerosol Attenuation and Extinction Profiles for the Troposphere and Stratosphere, Report AFCRL-66-828, AFCRL, Bedford, MA, USA, 128 pp., <https://apps.dtic.mil/sti/pdfs/AD0649778.pdf> (last access: 5 June 2024), 1966b.
- Elterman, L.: Stratospheric aerosol parameters for the Fuego volcanic incursion, *Appl. Optics*, 14, 1262–1263, <https://doi.org/10.1364/AO.14.001262>, 1975.

- Elterman, L.: Aerosol measurements since 1973 for normal and volcanic stratospheres, *Appl. Optics*, 15, 1113–1114, <https://doi.org/10.1364/AO.15.001113>, 1976.
- Elterman, L., Toolin, R. B., and Essex, J. D.: Stratospheric Aerosol Measurements with Implications for Global Climate, *Appl. Optics*, 12, 1262–1263, <https://doi.org/10.1364/AO.12.000330>, 1973.
- Fiocco, G. and Grams, G.: Observations of the aerosol layer at 20 km by optical radar, *J. Atmos. Sci.*, 21, 323–324, [https://doi.org/10.1175/1520-0469\(1964\)021<0323:OOTALA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1964)021<0323:OOTALA>2.0.CO;2), 1964.
- Fontijn, K., Costa, F., Sutawidjaja, I., Newhall, C. G., and Herrin, J. S.: A 5000-year record of multiple highly explosive mafic eruptions from Gunung Agung (Bali, Indonesia): implications for eruption frequency and volcanic hazards, *B. Volcanol.*, 77, 59, <https://doi.org/10.1007/s00445-015-0943-x>, 2015.
- Gao, C., Robock, A., and Ammann, C.: Volcanic forcing of climate over the past 1500 years: an improved ice core-based index for climate models, *J. Geophys. Res.*, 113, D23111, <https://doi.org/10.1029/2008JD010239>, 2008.
- Kondratyev, K. Ya., Ivlev, L. S., and Nikolov, G. A.: Investigations of the stratospheric aerosols, Proc. 3rd Conf. on the Climatic Impact Assessment Program, 26 February–1 March, 143–152, <https://rosap.nsl.bts.gov/view/dot/10975> (last access: 24 May 2025), 1974.
- Reeger, E. and Siedentopf, H.: Die Streufunktion des Atmosphärischen Dunstes nach Scheinwerfer Messungen, *Optik*, 1, 15–41, 1946.
- Rohatgi, A.: WebPlotDigitizer, Version 5, <https://automeris.io> (last access: 23 November 2024), 2024.
- Sato, M., Hansen, J. E., McCormick, M. P., and Pollack, J. B.: Stratospheric aerosol optical depths, *J. Geophys. Res.*, 98, 22987, <https://doi.org/10.1029/93JD02553>, 1993.
- Self, S. and Rampino, M. R.: The 1963–1964 eruption of Agung volcano (Bali, Indonesia), *B. Volcanol.*, 74, 1521–1536, <https://doi.org/10.1007/s00445-012-0615-z>, 2012.
- Stothers, R. B.: Major optical depth perturbations to the stratosphere from volcanic eruptions: Stellar extinction period, 1961–1978, *J. Geophys. Res.-Atmos.*, 106, 2993–3003, <https://doi.org/10.1029/2000JD900652>, 2001.
- Thomason, L. W., Ernest, N., Millán, L., Rieger, L., Bourassa, A., Vernier, J.-P., Manney, G., Luo, B., Arfeuille, F., and Peter, T.: A global space-based stratospheric aerosol climatology: 1979–2016, *Earth Syst. Sci. Data*, 10, 469–492, <https://doi.org/10.5194/essd-10-469-2018>, 2018.
- Volz, F. E.: Twilight Phenomena Caused by the Eruption of Agung Volcano., *Science*, 144, 1121–1122, <https://www.jstor.org/stable/1713387> (last access: 2 February 2025), 1964.
- Wells, M. B.: Monte Carlo Analysis of Searchlight Scattering Measurements, AFCRL-68-0311, AD0675153, Air Force Cambridge Research Laboratories, Office of Aerospace Research, United States Air Force: Bedford, MA, USA, 1968, 23 pp., <https://apps.dtic.mil/sti/pdfs/AD0675153.pdf> (last access: 26 May 2025), 1968.