



A 40-year high-resolution gridded meteorological dataset derived from station observations in the Reynolds Creek Experimental Watershed

Andrew R. Hedrick¹, Brandon Stairs¹, C. Jason Williams¹, Joachim Meyer², James P. McNamara², and Patrick Kormos³

¹Northwest Watershed Research Center, US Department of Agriculture –
Agricultural Research Service, Boise, Idaho 83702, USA

²Department of Geosciences, Boise State University, Boise, Idaho 83712, USA

³Idaho Snow Survey, US Department of Agriculture – Natural Resource Conservation Service,
Boise, Idaho 83709, USA

Correspondence: Andrew R. Hedrick (andrew.hedrick@usda.gov)

Received: 26 September 2025 – Discussion started: 17 April 2026

Revised: 8 July 2026 – Accepted: 12 July 2026 – Published: 28 July 2026

Abstract. A 40-year gridded meteorological forcing dataset spanning the water years from 1 October 1983 to 30 September 2023 was compiled for the Reynolds Creek Experimental Watershed (RCEW) in southwest Idaho, USA. This Reynolds Creek Long-Term (RCLT) dataset consists of hourly, 10 m resolution grids of air temperature, vapor pressure, precipitation mass and phase, incoming shortwave and longwave radiation, visible and infrared snow albedo, and wind speed and direction. These variables were interpolated and calculated from hourly measurements from the dense meteorological station network within the mountainous 239 km² RCEW, which contains elevations that span the historical winter rain-to-snow transition. The observations are foundational for many ecological and hydrological Land Surface Models (LSMs) used in research and operational applications. Additionally, an example use case is presented in which we show how the snow-dominated area of the basin has evolved over the data record. This 13 TB dataset, stored in cloud-optimized Zarr format, enables future model development, benchmarking, uncertainty analyses of existing models, independent validation of gridded atmospheric reanalysis datasets, and novel investigations of hydroclimatic variability across snow-dominated semi-arid environments. Data access is available via the following repository: <https://doi.org/10.15482/USDA.ADC/30199954> (Hedrick et al., 2025).

1 Introduction

Hydrologic models are important tools for examining relationships between hydrologic response and interannual weather variability, yet modeling studies investigating these relationships require accurate forcing data over suitably long periods of record. Studies examining long-term hydrologic trends have been based on data from measurement networks such as the US Department of Agriculture SNOTEL Network (Daly et al., 1994; Mote et al., 2005; Serreze et al., 1999; Trujillo and Molotch, 2014; Zeng et al., 2018) and the US Geological Survey (USGS) stream gauge net-

work (McCabe and Clark, 2005; Stewart et al., 2005). However, these station networks only represent individual locations and are unable to infer changes across broader spatial scales, especially in complex mountain topography. More recently, long-term reanalysis datasets such as the Modern-Era Retrospective Analysis for Research and Applications, version 2 (MERRA-2) (Gelaro et al., 2017), the North American Land Data Assimilation System, version 2 (NLDAS-2) (Xia et al., 2012), the US National Weather Service (NWS) Analysis of Record for Calibration (AORC) (Fall et al., 2023), the National Center for Atmospheric Research (NCAR)–USGS collaborative CONUS404 dataset (Rasmussen et al., 2023),

and the Gridded Surface Meteorological (gridMET) dataset (Abatzoglou, 2013), among others, have been developed to overcome this lack of spatial support for sparse weather measurements. These datasets, often referred to as “high-resolution” for grid spacings ranging between 1 and 4 km², are produced by downscaling global climate models (GCMs) using either statistical methods or Numerical Weather Prediction (NWP) models such as the Weather Research and Forecasting (WRF) model (Skamarock et al., 2021). Some datasets go one step further by assimilating local station measurements to increase accuracy and reduce bias. However, many widely used reanalysis datasets remain daily products, precluding any ability to discern precipitation phase on an individual storm basis due to diurnal air temperature variability.

Although many long-term atmospheric reanalysis datasets exist for regional- and continental-scale applications, there is a lack of long-term, high spatial and temporal resolution datasets geared toward basin-scale applications and evaluating biases in reanalysis datasets. Specifically, distributed snow energy balance models such as iSnobal (Marks et al., 1999), SnowModel (Liston and Elder, 2006), Crocus (Brun et al., 1989), and Alpine3D (Lehning et al., 2006), originally developed for research applications, are becoming more common for operational applications (Meyer et al., 2023; Morin et al., 2020; Mott et al., 2023). The need for real-time forcing data to execute these models in water resource applications now necessitates the use of the NWP models such as the National Oceanic and Atmospheric Administration’s High-Resolution Rapid Refresh (HRRR) model (Meyer et al., 2023). Rigorous evaluations of uncertainty and bias of NWP model forcing data are crucial since forcing data uncertainty is the largest contributor to model uncertainty (Raleigh et al., 2015; Voordendag et al., 2021).

Here we present an hourly, 10 m spatially distributed dataset of all meteorological variables required to force a snow energy balance or land surface model for the Reynolds Creek Experimental Watershed (RCEW) in southwestern Idaho, USA along with an example use case. This dataset extends and updates existing records, enabling long-term analysis over a forty-year record, and highlights the importance of maintaining public datasets while sharing them openly with the greater science community to maximize utility.

2 Site description

In 1960, the United States Congress allocated funding for an experimental research watershed to advance hydrologic research in western US rangelands and to develop new solutions for sustainable land management within the Great Basin ecosystem. Since then, the US Department of Agriculture (USDA) Agricultural Research Service (ARS) Northwest Watershed Research Center (NWRC) has managed the scientific infrastructure in the 239 km²

Reynolds Creek Experimental Watershed (RCEW) in southwest Idaho (43.205, −116.75), which is characteristic of semiarid snow-dominated mountain environments found throughout the Great Basin. The RCEW spans an elevation gradient of 1100 to 2242 m a.s.l. (above sea level) and experiences a significant elevational and directional precipitation gradient due to the prevailing northeast-trending storms during the winter and spring, when most of the annual precipitation occurs (Hanson, 2001). Since 1984, average annual precipitation ranged from 228 mm at site RC.057 in the lowest elevations to 1086 mm at site RC.163 in the highest elevations of the watershed (Fig. 1). The NWRC has a long history of publishing station-based hydrometeorological datasets, including Slaughter et al. (2001), Reba et al. (2011) and Godsey et al. (2018). This dataset updates the gridded air temperature, relative humidity, and precipitation dataset reported by Kormos et al. (2018), includes additional wind and radiation data, and appends nine more recent years (2015–2023) that have experienced a wider range of weather variability (Monteiro and Morin, 2023).

3 Dataset description

The gridded dataset, called the Reynolds Creek Long-Term (RCLT) dataset, was derived using the dense RCEW station network (one station per 6 km²) maintained by the USDA-ARS-NWRC. Dataset variables include air temperature, vapor pressure, precipitation mass, wet bulb temperature-derived precipitation phase, density of new fallen snow, *U*- and *V*-components of wind, incoming short-wave and longwave radiation, and modeled visible and infrared spectral albedo for snow-covered surfaces (Fig. 2). The temporal record spans 40 water years from 1 October 1983 to 30 September 2023 (a water year beginning on 1 October of the previous year and ending on 30 September of the current year).

4 Instrumentation and variable distribution

The Spatial Modeling for Resources Framework (SMRF v0.11.7) (Havens et al., 2017) was employed to distribute each of the 12 land-surface meteorological variables to a 10 m regular grid. Each forcing variable has either been empirically derived or directly interpolated from hourly station measurements across the catchment domain. Owing to the considerable length of time encompassed by the dataset, many different sensors have been deployed in the watershed over the 40-year data record with differing levels of accuracy, which are not reported here but are available upon request. Invalid data were preliminarily removed for all measured variables besides precipitation (which necessitated a unique approach detailed in Sect. 4.6), and temporal interpolation was performed for data gaps of two hours or less. Gaps lasting longer than two hours were




Figure 1. Map of Reynolds Creek Experimental Watershed overlaid on a 10 m digital elevation model with site locations of the meteorological measurements used to produce this dataset. Hourly site measurements include precipitation mass (Ppt), air temperature (T_a), relative humidity (RH), wind speed and direction (u), and incoming shortwave radiation (S_{in}). Also, the six primary sub-watersheds are delineated by their outlet weir locations where streamflow (Q) is measured. Weir sites are denoted by the solid single-color circles corresponding to the basin boundary colors.

RCLT Hourly Variables

Air Temperature [°C] (air_temp)	Precipitation [mm] (precip)
Vapor Pressure [Pa] (vapor_pressure)	Snow Fraction of Precipitation [0-1] (percent_snow)
East-West Wind Speed [m/s] (u_component)	Density of New Snow [kg/m³] (snow_density)
North-South Wind Speed [m/s] (v_component)	Wet Bulb Temperature [°C] (precip_temp)
Incident Shortwave Radiation [W/m²] (incoming_solar)	Visible Band Snow Albedo [0-1] (albedo_vis)
Incident Longwave Radiation [W/m²] (thermal)	Infrared Band Snow Albedo [0-1] (albedo_ir)

Figure 2. The 12 meteorological variables and their units produced for the high-resolution Reynolds Creek Long-Term (RCLT) dataset. Variable filenames in the Zarr dataset are listed in parentheses.

left empty with the foreknowledge that spatial interpolation from nearby sites in the high-density network would act as surrogate data for those time steps. It is worth noting that the NWRC has always employed a full-time staff of technicians tasked with the calibration and servicing of each sensor deployed in the RCEW. The gridded interpolation methods for all 12 modeled variables from the six measured variables are described in Hedrick et al. (2018) and further elaborated upon in the following subsections.

4.1 Air temperature

Hourly measurements of air temperature (T_a) were made at 35 individual sites over the 40-year period (Fig. 1), with the number of sites significantly increasing after water year 2000 (Fig. 3). Measurements are currently made using various incarnations of the widely used Vaisala HMP series of temperature and humidity sensors with ventilated radiation shields.

A modified inverse distance weighting (IDW) approach was used to distribute T_a across the 10 m grid. In this process, the elevational trend is calculated at each time step, constrained to be negative due to the general relationship between elevation and T_a , then subtracted from the station measurements to produce a temperature residual. These residuals are distributed using standard IDW and added to each grid cell's position on the elevation gradient slope line. This approach for distributing T_a improves station representativeness for areas with complex mountain topography (Havens et al., 2017). Between 1984 and 2023, the mean annual distributed air temperature across the entire RCEW was 8.4 °C and the all-time maximum and minimum hourly temperatures were 42.4 and −31.5 °C, respectively.

4.2 Vapor pressure

Gridded actual vapor pressure (e_a) values were interpolated from measurements of relative humidity (RH) at the same 35 sites as T_a over the 40-year data record, using the same Vaisala HMP instruments referred to in Sect. 4.1. The empirical Tetens equation was used for deriving e_a from RH and T_a (in degrees Celsius) when temperatures are above freezing:

$$e_a = \text{RH} \cdot 0.6108 \cdot e^{\left(\frac{17.3 \times T_a}{237.3 + T_a}\right)}, \quad (1)$$

while the Magnus–Tetens equation is used when temperatures are below freezing and the ground surface is assumed to be ice:

$$e_a = \text{RH} \cdot 0.6108 \cdot e^{\left(\frac{21.9 \times T_a}{265.5 + T_a}\right)}. \quad (2)$$

Station-derived e_a was then distributed to the 10 m grid using the same modified IDW detrending approach described in Sect. 4.1. Dew point temperature was also calculated for the wet bulb temperature calculation (Sect. 4.6.2 below) but was not stored in the RCLT dataset because it can be calculated from the vapor pressure and air temperature using the existing empirical relationships (e.g., the Clausius–Clapeyron equation). Over the dataset record, the mean distributed e_a across the RCEW was 584 Pa, with a maximum value of 2551 Pa during an intense summer thunderstorm and a threshold minimum of 20 Pa that only occurred during a handful of very cold and dry periods.

4.3 Wind

Hourly average wind speed (u_s) and hourly resultant-mean-wind direction (u_{dir}) were measured at a total of 29 sites over




Figure 3. Annual water year measurement record used in SMRF gridded interpolation from the 40 individual sites within RCEW from 1 October 1983 to 30 September 2023. The horizontal bars for each variable indicate station observation data present for over half of a water year, however smaller data gaps are not shown. Some sites measure all six variables required by SMRF to produce the 12 gridded hydrometeorological forcing variables, though many are instrumented to measure fewer variables. For example, five stations solely measure precipitation (RC.049, RC.057, RC.116C, RC.147, and RC.155).

the data record, though only three sites were available prior to 1994 and four sites prior to 2002 (Fig. 3). The sparseness of wind measurements through the early years of this dataset is likely a source of uncertainty in the distributed wind grids for that period, though we should note that the pre-2002 measurements captured the full elevation gradient in RCEW at low (RC.076), mid (RC.127), and high elevation sites (RC.176) (Fig. 1). Minimum wind speeds were set to a lower threshold of 0.447 m s^{-1} (1 mph) to match the lower accuracy reported by most cup anemometer manufacturers. Similarly, maximum wind speeds were capped at 35 m s^{-1}

to avoid numerical instability issues when calculating turbulent heat fluxes in physically based energy balance models.

Station measurements of wind were distributed to the 10 m grid using the maximum upwind slope (maxus) terrain parameter described in Winstral et al. (2002, 2009). In short, the underlying digital elevation model (DEM) was used to calculate a maxus value (in degrees) over a user-defined upwind distance (here 300 met) for all possible upwind directions (0 to 360°) in 5° increments. The resulting 72 layers of maxus grids were stored in a lookup library. Then, for each station, the measured wind speed was adjusted to sim-

ulate what the wind speed would have been on a flat surface (“flatwind”) using the maxus value for the measured wind direction at the site. Once the adjusted “flatwind” speeds and wind direction components had been distributed across the entire grid using standard IDW, the distributed wind directions were used to find the maxus value for each grid cell, and the distributed “flatwind” speeds were converted back to actual wind speeds.

For the gridded dataset, wind speed and direction were converted into U - and V -components to match the conventions of NWP models such as the WRF and HRRR models. The U -component represents the East–West wind speed, with positive values indicating wind out of the west, while the V -component represents the North–South wind speed, with positive values indicating wind out of the south. Over the dataset record, the mean hourly wind speed was 4.1 m s^{-1} and the mean resultant wind direction was out of the southwest (227).

4.4 Shortwave radiation

Gridded incoming shortwave radiation (S_{in}) was measured at 23 sites across the RCEW (Fig. 3) but cannot be directly spatially interpolated from measurements due to the complex terrain and the variable vegetation canopy present across the catchment. Rather, a three-step process produced the hourly gridded S_{in} product.

1. Hourly clear sky atmospheric S_{in} was modeled following Dozier (1980) and then corrected for surrounding terrain in each 10 m grid cell following Dubayah (1994).
2. Station measurements of S_{in} were divided by modeled clear sky radiation to derive a cloud factor (C_{fac}) at each station pixel ($C_{\text{fac}} = 1$ represents cloud-free conditions), which was then distributed using standard IDW across the domain.
3. Canopy-corrected S_{in} values were estimated using empirical relationships presented by Link and Marks (1999), where direct beam shortwave radiation under canopy (R_{b}) can be represented by:

$$R_{\text{b}} = S_{\text{b},\text{in}} \cdot e^{-\frac{\mu h}{\cos \theta}}. \quad (3)$$

In Eq. (2), $S_{\text{b},\text{in}}$ is the above canopy cloud corrected direct beam radiation, μ is a canopy extinction coefficient, h is the height of the canopy, and θ is the solar zenith angle. Diffuse shortwave radiation under canopy (R_{d}) was computed by adjusting the above canopy cloud corrected diffuse radiation ($S_{\text{d},\text{in}}$) by the canopy optical transmissivity (τ):

$$R_{\text{d}} = \tau S_{\text{d},\text{in}}. \quad (4)$$

The terrain-, cloud-, and canopy-corrected S_{in} presented in this dataset is the sum of Eqs. (3) and (4).

When a comparison was performed between modeled and measured S_{in} across the 40-year record, we discovered that the values for τ and μ presented in Link and Marks (1999), which were derived in the Canadian Boreal forests, led to an overestimation of S_{in} at open sites by up to 40 %, and an underestimation of S_{in} at forested sites by 30 % or more. Therefore, manual adjustments of τ , μ , and height parameters were performed to produce S_{in} values that more closely matched the station observations (Fig. 4). Importantly, the scale difference between point measurements of S_{in} and averaged S_{in} across a 10 m by 10 m grid cell precludes direct agreement since shortwave radiation at the ground surface varies over very short spatial scales. However, the general trends in modeled S_{in} magnitude as a function of cloud cover were well represented in the spatial dataset. Over the dataset record, the mean distributed S_{in} over the RCEW was 204 W m^{-2} (including nighttime hours) and the maximum reported value was $1,401 \text{ W m}^{-2}$. Though this maximum value exceeds average top-of-atmosphere incoming shortwave irradiance (1340 W m^{-2}), cloud enhancement (CE) and albedo enhancement (AE) events have been shown to result in pyranometer measurements of this magnitude at snow-dominated sites (Gueymard, 2017).

Calculating net shortwave radiation from the RCLT incoming solar product requires an estimate of land surface reflectance, or albedo, and there are many ways to derive an albedo product from both models and remote sensing products. Users of the RCLT dataset are encouraged to use their own methods, but for simplicity we also include here visible and infrared bands of snow albedo for when snow is present. These albedo estimates use a time decay approach to capture albedo change as a function of springtime snow metamorphism, terrain factors, and solar zenith angle (Marshall and Warren, 1987). Importantly, these albedo estimates do not apply for snow-free conditions and would need to be masked for general land surface modeling applications.

4.5 Longwave radiation

Incoming longwave radiation (L_{in}) was also not directly interpolated from observations since the network lacked upward looking pyrgeometers for measuring thermal radiation during long periods throughout the RCLT time domain. Instead, clear sky L_{in} was empirically estimated from distributed air temperature and humidity using the methods of Brutsaert (1975), then corrected for surrounding terrain using equations presented in Marks and Dozier (1979). Next, cloud-corrected L_{in} was computed using the empirical relationship described by Garen and Marks (2005) and the same cloud factor calculated from the incoming shortwave radiation. The final step was an adjustment to the cloud-corrected L_{in} using vegetation maps derived from the LANDFIRE 2016 dataset (LANDFIRE, 2016) and empirically derived transmissivity (τ) reported by Link and Marks (1999) then modified as described in Sect. 4.4.




Figure 4. For an example water year (2009), observed versus modeled maximum daily incoming solar radiation (S_{in}) at three sites with different surrounding canopy coverage. The teal model traces were derived using canopy transmissivity values and extinction coefficients derived by Link and Marks (1999), while the yellow marker traces resulted from new adjusted values to more closely match pyranometer measurements at sites throughout the watershed.

Over the dataset record, mean L_{in} across the RCEW was 290 W m^{-2} , with maximum and minimum hourly values of 600 and 83 W m^{-2} , respectively.

4.6 Precipitation

To satisfy the standard input requirements of an energy and mass balance snow model, the RCLT dataset contains four distinct variables related to precipitation in the basin. These variables of precipitation mass, temperature of the falling hydrometeor, initial density of newly fallen snow, and the snow proportion of precipitation are described in the following subsections.

4.6.1 Precipitation mass

Despite being the foundation upon which hydrologic models rely, precipitation measurements are often the largest source of predictive hydrologic uncertainty (Bárdossy et al., 2022). Across the wide spectrum of snow-dominated watersheds in the Western US, the majority of in situ measurements are made with weighing buckets fitted with alter shields that reduce wind speeds above the bucket orifice and thus increase the gauge catch efficiency (CE), or the ratio of measured precipitation to a “true” value (Thériault et al., 2021). However, many sites lack co-located wind speed measurements for applying the necessary World Meteorological Organization (WMO) transfer functions for undercatch correction

(Kochendorfer et al., 2018), which can lead to low biases in regional and basin estimates of precipitation. Additionally, precipitation exhibits high spatial heterogeneity in complex terrain, which cannot be captured by a single measurement site in a large mountain basin.

To overcome the issue of spatial representativeness, the RCEW measurement network was initially planned to contain one gauge for every square mile with 110 stations in the watershed. By the beginning of this dataset in water year 1984, the number of sites had been reduced to the 25 stations (~ 1 measurement per 10 km^2) used here to produce the hourly gridded precipitation fields.

To address the undercatch issue, a unique dual-gauge approach (Hamon and Cox, 1972) was used for all sites in the RCEW except for RC.124B. This method requires two co-located Belfort-type weighing buckets with one existing unshielded and the other fitted with a single-alter shield. The combination of the shielded and unshielded measurements allows an empirical extrapolation of more accurate “actual” precipitation data compared with single shielded gauges employing a WMO transfer function (Hanson et al., 2004). Site RC.124B is the only single shielded gauge in the basin and was corrected using the WMO transfer function. Hourly measurements of precipitation mass were distributed across the 10 m grid using a Detrended Kriging interpolation method (Garen et al., 1994), which is identical to the approach by Kormos et al. (2018). Over the dataset record, the mean annual precipitation normalized over the RCEW was 461 mm , with a maximum of 686 mm (water year 1984) and a minimum of 279 mm (water year 1994). However, due to the large elevational gradient in precipitation, the mean for the high elevation Reynolds Mountain East sub-catchment (0.4 km^2) was 913 mm while conversely the mean for the low elevation Nancy Gulch sub-catchment (0.3 km^2) was only 325 mm .

4.6.2 Precipitation temperature

The precipitation temperature variable is represented by the hourly computed ice or wet bulb temperature calculated with a widely used Newton-Raphson iterative solution to the psychrometric equation (Campbell and Norman, 1998). This approach requires air temperature (Sect. 4.1), dew point temperature (from calculated vapor pressure in Sect. 4.2) and estimated atmospheric pressure from elevation. Previous research has demonstrated that wet bulb temperature is the most suitable precipitation phase partitioning method in a semiarid watershed such as the RCEW (Marks et al., 2013).

4.6.3 Density of new snow

Hourly estimates of new snow density are included in the RCLT dataset for energy and mass balance snow models that may require it. For this long-term application, we computed new snow density from a lookup table based on previous

work (pg. 2007, Susong et al., 1999) using the calculated precipitation temperature (see Sect. 4.6.2) and precipitation mass (see Sect. 4.6.1) in each grid cell. These lookup table values are distributed in a stepwise fashion between -5 and $+0.5^\circ\text{C}$ with values ranging between 75 and 250 kg m^{-3} , respectively. Snowfall occurring below -5°C , which rarely occurs in the RCEW, is assigned a new snow density of 75 kg m^{-3} .

4.6.4 Snow fraction of precipitation

In addition to new snow density, the lookup table from Susong et al. (1999) was also used to determine the amount of precipitation that fell as snow across the catchment. When time step precipitation temperatures fell between -0.5 and $+0.5^\circ\text{C}$, the precipitation was defined as mixed phase. The relationship between precipitation phase and wet bulb temperature is linear such that 0°C results in a 0.5 snow fraction, -0.5°C is 100% snow, and $+0.5^\circ\text{C}$ is 100% rain.

4.6.5 Example dataset use case

This dataset allows a detailed analysis of the storm-by-storm precipitation amount, spatial extent, and phase over the 40-year record. As an example of the utility of this unique dataset, we examined how the annual average storm-based rain/snow transition elevation (RS_z) has changed over time in the snow-dominated 55 km^2 Tollgate sub-watershed in the southern portion of the RCEW. We began by designating storm hours as any time steps in which any pixel in the catchment area received above a precipitation threshold of 0.5 mm . Next, storm periods were defined for consecutive hours with a maximum storm hour threshold of 6 h . For each storm hour, the mean elevation of mixed phase precipitation (e.g. grid cells where $-0.5^\circ\text{C} < T_{\text{wb}} < 0.5^\circ\text{C}$) was computed and mass-weighted by precipitation amount, then the hourly mean elevation was mass-weighted across the storm period. Lastly, those elevations were further mass-weighted by storm total precipitation to arrive at water year and accumulation season mean RS_z (Fig. 5a). The accumulation season was defined as the period from 1 October to peak snow water equivalent (SWE) accumulation at the Reynolds Mountain Snow Pillow site (RC.RMSP). Overall, the snow-dominated portion of the catchment has retreated toward higher elevations. When considering all storms throughout each water year, the RS_z has risen from 1830 m in 1984 to 1991 m in 2023, resulting in a decrease in snow-dominated basin area from 52% to 24% (Fig. 5b). For storms during the accumulation season, the RS_z has risen from 1580 to 1708 m between 1984 and 2023, with the snow-dominated basin area decreasing from 91% to 74% (Fig. 5c).




Figure 5. (a) The annual precipitation mass-weighted mean rain/snow transition elevation for water years 1984 to 2023 in the Tollgate sub-watershed for all storms over the course of a full year (orange diamonds and orange line of best fit) and the snow accumulation season from 1 October to peak SWE date (blue stars and blue line of best fit). Horizontal dotted lines represent the upper and lower elevations of the Tollgate sub-watershed. (b) The RS_z in 1984 (blue) and 2023 (red) when considering all storms in each year. (c) Same as (b) but only considering storms occurring from 1 October to the date of peak SWE at the Reynolds Mountain snow pillow (denoted by the black star).

5 Data availability

The gridded dataset in its published form is at hourly temporal and 10 m spatial resolution, which was determined to be optimal for users requiring high resolution forcing data while also being suitable for disk storage. NetCDF files, though standard format for meteorological data, can become unwieldy when file sizes become too large, which is mitigated by the cloud-optimized Zarr format (Miles et al., 2023) that can be read by programming libraries in python, R, C++, and Java. Nevertheless, users that desire a coarser spatiotemporal RCLT dataset have two options for decreasing resolution. The simplest approach would be to read the provided

high resolution files, resample to desired resolution using mean or sum values across space and time, and save to a new NetCDF file. A slightly more accurate approach would be to install SMRF locally, create a model setup file at the desired spatial resolution, edit the configuration files to define the coarser temporal resolution, and run the distribution code. However, this approach will require detailed knowledge of the software and is not recommended for most users. An example SMRF configuration file for a single water year and all station data CSV files organised by water year are provided in the Supplementary Materials.

Accessing temporal and spatial slices of this 13 TB dataset is straightforward as it is stored as

cloud-optimized Zarr-formatted files and hosted by the open-access Ag Data Commons data repository (<https://doi.org/10.15482/USDA.ADC/30199954>, Hedrick et al., 2025) maintained by the USDA National Agricultural Library (<https://agdatacommons.nal.usda.gov>, last access: 22 July 2026). Users may download either the entire 40-year record or by individual water year ($\sim 325 \text{ GB yr}^{-1}$) through the linked Globus web interface. It is also recommended that users work with the dataset on a High-Performance Computing (HPC) or Cloud environment with parallel processing capabilities. An example jupyter notebook script for loading the Zarr-formatted data using the python Xarray package is provided in the repository. The coordinates are stored as projected coordinates in UTM Zone 11 using the WGS84 geodetic reference system. At 10 m spatial resolution, the model domain shape is 3010 (north–south) by 1602 (east–west) pixels. The time domain is 350 640 total time steps, resulting in 1.7 trillion total stored values per variable, or 20.3 trillion stored values across the entire dataset. A tagged release for the SMRF source code used to create this dataset is available at <https://github.com/iSnobal/smrf/releases/tag/20250926> (last access: 22 July 2026; <https://doi.org/10.5281/zenodo.21514687>, Meyer et al., 2025).

6 Summary

The RCLT dataset comprises all meteorological forcing variables required to simulate the accumulation and melt of the seasonal snow cover using a mass and energy balance snow model as well as the fundamental land surface variables estimated by Numerical Weather Prediction models and historical atmospheric reanalysis products. The RCLT provides unique opportunities for comparison of various physically based models, evaluation of long-term atmospheric reanalysis datasets, and new studies on agroecosystem response to differing weather patterns. By publicly sharing this dataset in a cloud optimized format, we support modern software workflows, analysis, and open science best practices to support the broader science community. Maintaining and regular releasing long-term in situ network data is an important contribution to better understand and predict future water availability in snow-dominated watersheds.

Supplement. The supplement related to this article is available online at <https://doi.org/10.5194/essd-18-5531-2026-supplement>.

Author contributions. Conceptualization: ARH, PK. Methodology: ARH, BS, and JM. Data curation: ARH and BS. Writing (initial): ARH. Writing (review and editing): ARH, BS, CJW, JM, and JPM. Supervision: CJW.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgements. This research used resources provided by the SCINet project of the USDA Agricultural Research Service, ARS project number 0201-88888-003-000D. The open-access Spatial Modeling for Resources Framework can be accessed at <https://github.com/iSnobal/smrf/tree/20250926> (last access: 22 July 2026). This research was supported by the USDA-ARS CRIS Project, Ecohydrology of Mountainous Terrain in a Changing Climate (2052-13610-012-00D). This research is a contribution from the Long-Term Agroecosystem Research (LTAR) network. LTAR is supported by the USDA. The Reynolds Creek Experimental Watershed is part of the USDA Agricultural Research Service, Great Basin LTAR site within the LTAR Network. USDA is an equal opportunity employer.

Financial support. This research has been supported by two US Department of Agriculture – Agricultural Research Service projects entitled Ecohydrology of Mountainous Terrain in a Changing Climate (2052-13610-012-00D) and Disturbance Mitigation and Adaptive Restoration of Sagebrush-Steppe Ecosystems (2052-21500-001-000-D).

Review statement. This paper was edited by John Pomeroy and reviewed by two anonymous referees.

References

- Abatzoglou, J. T.: Development of gridded surface meteorological data for ecological applications and modelling, *Int. J. Climatol.*, 33, 121–131, <https://doi.org/10.1002/joc.3413>, 2013.
- Bárdossy, A., Kilsby, C., Birkinshaw, S., Wang, N., and Anwar, F.: Is Precipitation Responsible for the Most Hydrological Model Uncertainty? *Front. Water*, 4, <https://doi.org/10.3389/frwa.2022.836554>, 2022.
- Brun, E., Martin, E., Simon, V., Gendre, C., and Coleou, C.: An Energy and Mass Model of Snow Cover Suitable for Operational Avalanche Forecasting, *J. Glaciol.*, 35, 333–342, <https://doi.org/10.3189/s0022143000009254>, 1989.
- Brutsaert, W.: On a derivable formula for long-wave radiation from clear skies, *Water Resour. Res.*, 11, 742–744, <https://doi.org/10.1029/WR011i005p00742>, 1975.
- Campbell, G. S. and Norman, J. M.: *An Introduction To Environmental Biophysics*, in: 2nd Edn., Springer-Verlag, ISBN 978-0-387-94937-6, 1998.

- Daly, C., Neilson, R. P., and Phillips, D. L.: A Statistical-Topographic Model for Mapping Climatological Precipitation over Mountainous Terrain, *J. Appl. Meteorol.*, 33, 140–158, [https://doi.org/10.1175/1520-0450\(1994\)033<0140:ASTMFM>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<0140:ASTMFM>2.0.CO;2), 1994.
- Dozier, J.: A clear-sky spectral solar radiation model for snow-covered mountainous terrain, *Water Resour. Res.*, 16, 709–718, <https://doi.org/10.1029/WR016i004p00709>, 1980.
- Dubayah, R. C.: Modeling a solar radiation topoclimatology for the Rio Grande River Basin, *J. Veg. Sci.*, 5, 627–640, <https://doi.org/10.2307/3235879>, 1994.
- Fall, G., Kitzmiller, D., Pavlovic, S., Zhang, Z., Patrick, N., St. Laurent, M., Trypaluk, C., Wu, W., and Miller, D.: The Office of Water Prediction's Analysis of Record for Calibration, version 1.1: Dataset description and precipitation evaluation, *J. Am. Water Resour. Assoc.*, 59, 1246–1272, <https://doi.org/10.1111/1752-1688.13143>, 2023.
- Garen, D. C. and Marks, D.: Spatially distributed energy balance snowmelt modelling in a mountainous river basin: estimation of meteorological inputs and verification of model results, *J. Hydrol.*, 315, 126–153, <https://doi.org/10.1016/j.jhydrol.2005.03.026>, 2005.
- Garen, D. C., Johnson, G. L., and Hanson, C. L.: Mean Areal Precipitation for Daily Hydrologic Modeling in Mountainous Regions, *J. Am. Water Resour. Assoc.*, 30, 481–491, <https://doi.org/10.1111/j.1752-1688.1994.tb03307.x>, 1994.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Parityka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Godsey, S. E., Marks, D., Kormos, P. R., Seyfried, M. S., Enslin, C. L., Winstral, A. H., McNamara, J. P., and Link, T. E.: Eleven years of mountain weather, snow, soil moisture and streamflow data from the rain–snow transition zone – the Johnston Draw catchment, Reynolds Creek Experimental Watershed and Critical Zone Observatory, USA, *Earth Syst. Sci. Data*, 10(3), 1207–1216, <https://doi.org/10.5194/essd-10-1207-2018>, 2018.
- Gueymard, C. A.: Cloud and albedo enhancement impacts on solar irradiance using high-frequency measurements from thermopile and photodiode radiometers. Part 1: Impacts on global horizontal irradiance, *Solar Energy*, 153, 755–765, <https://doi.org/10.1016/j.solener.2017.05.004>, 2017.
- Hamon, W. R. and Cox, L. M.: Measurement of atmospheric and ground level precipitation, *Water Resour. Res.*, 8, 272–272, <https://doi.org/10.1029/WR008i001p00272-01>, 1972.
- Hanson, C. L.: Long-Term Precipitation Database, Reynolds Creek Experimental Watershed, Idaho, United States, *Water Resour. Res.*, 37, 2831–2834, <https://doi.org/10.1029/2001WR000415>, 2001.
- Hanson, C. L., Pierson, F. B., and Johnson, G. L.: Dual-Gauge System for Measuring Precipitation: Historical Development and Use, *J. Hydrol. Eng.*, 9, 350–359, [https://doi.org/10.1061/\(ASCE\)1084-0699\(2004\)9:5\(350\)](https://doi.org/10.1061/(ASCE)1084-0699(2004)9:5(350)), 2004.
- Havens, S., Marks, D., Kormos, P. R., and Hedrick, A. R.: Spatial Modeling for Resources Framework (SMRF): A modular framework for developing spatial forcing data for snow modeling in mountain basins, *Comput. Geosci.*, 109, 295–304, <https://doi.org/10.1016/j.cageo.2017.08.016>, 2017.
- Hedrick, A. R., Marks, D., Havens, S., Robertson, M., Johnson, M., Sandusky, M., Marshall, H. P., Kormos, P. R., Bormann, K. J., and Painter, T. H.: Direct Insertion of NASA Airborne Snow Observatory-Derived Snow Depth Time Series Into the iSnobal Energy Balance Snow Model, *Water Resour. Res.*, 54, 8045–8063, <https://doi.org/10.1029/2018WR023190>, 2018.
- Hedrick, A. R., Stairs, B., Williams, C. J., Meyer, J., and McNamara, J. P.: The Reynolds Creek Long-Term Dataset: A long-term meteorological dataset derived from station observations in the Reynolds Creek Experimental Watershed, Ag Data Commons, United States Department of Agriculture, National Agricultural Library [data set], <https://doi.org/10.15482/USDA.ADC/30199954>, 2025.
- Kochendorfer, J., Nitu, R., Wolff, M., Mekis, E., Rasmussen, R., Baker, B., Earle, M. E., Reverdin, A., Wong, K., Smith, C. D., Yang, D., Roulet, Y.-A., Meyers, T., Buisan, S., Isaksen, K., Brækkan, R., Landolt, S., and Jachcik, A.: Testing and development of transfer functions for weighing precipitation gauges in WMO-SPICE, *Hydrol. Earth Syst. Sci.*, 22, 1437–1452, <https://doi.org/10.5194/hess-22-1437-2018>, 2018.
- Kormos, P. R., Marks, D., Seyfried, M. S., Havens, S. C., Hedrick, A. R., Lohse, K. A., Sandusky, M., Kahl, A., and Garen, D.: 31 years of hourly spatially distributed air temperature, humidity, and precipitation amount and phase from Reynolds Critical Zone Observatory, *Earth Syst. Sci. Data*, 10, 1197–1205, <https://doi.org/10.5194/essd-10-1197-2018>, 2018.
- LANDFIRE: Existing Vegetation Type Layer, LANDFIRE 2.0.0, <https://www.landfire.gov/viewer> (last access: 22 July 2026), 2016.
- Lehning, M., Völksch, I., Gustafsson, D., Nguyen, T. A., Stähli, M., and Zappa, M.: ALPINE3D: A detailed model of mountain surface processes and its application to snow hydrology, *Hydrol. Process.*, 20, 2111–2128, <https://doi.org/10.1002/hyp.6204>, 2006.
- Link, T. E. and Marks, D.: Distributed simulation of snow-cover mass- and energy-balance in the boreal forest, *Hydrol. Process.*, 13, 2439–2452, [https://doi.org/10.1002/\(SICI\)1099-1085\(199910\)13:14/15<2439::AID-HYP866>3.0.CO;2-1](https://doi.org/10.1002/(SICI)1099-1085(199910)13:14/15<2439::AID-HYP866>3.0.CO;2-1), 1999.
- Liston, G. E. and Elder, K. J.: A Distributed Snow-Evolution Modeling System (SnowModel), *J. Hydrometeorol.*, 7, 1259–1276, <https://doi.org/10.1175/JHM548.1>, 2006.
- Marks, D. and Dozier, J.: A clear-sky longwave radiation model for remote alpine areas, *Arch. Meteorol. Geophys. Bioklimatol. B*, 27, 159–187, <https://doi.org/10.1007/BF02243741>, 1979.
- Marks, D., Domingo, J., Susong, D., Link, T., and Garen, D.: A spatially distributed energy balance snowmelt model for application in mountain basins, *Hydrol. Process.*, 13, 1935–1959, [https://doi.org/10.1002/\(SICI\)1099-1085\(199909\)13:12/13<1935::AID-HYP868>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1099-1085(199909)13:12/13<1935::AID-HYP868>3.0.CO;2-C), 1999.

- Marks, D., Winstral, A., Reba, M., Pomeroy, J., and Kumar, M.: An evaluation of methods for determining during-storm precipitation phase and the rain/snow transition elevation at the surface in a mountain basin, *Adv. Water Resour.*, 55, 98–110, <https://doi.org/10.1016/j.advwatres.2012.11.012>, 2013.
- Marshall, S. E. and Warren, S. G.: Parameterization of snow albedo for climate models, in: Large-scale effects of seasonal snow cover, edited by: Goodison, B. E., Barry, R. G., and Dozier, J. E., International Association of Hydrological Sciences Publication 166, Wallingford: IAHS Press, 43–50, 1987.
- McCabe, G. J. and Clark, M. P.: Trends and Variability in Snowmelt Runoff in the Western United States, *J. Hydrometeorol.*, 6, 476–482, <https://doi.org/10.1175/JHM428.1>, 2005.
- Meyer, J., Horel, J., Kormos, P., Hedrick, A., Trujillo, E., and Skiles, S. M.: Operational water forecast ability of the HRRR-iSnobal combination: an evaluation to adapt into production environments, *Geosci. Model Dev.*, 16, 233–250, <https://doi.org/10.5194/gmd-16-233-2023>, 2023.
- Meyer, J., Robledano, A., Hu, M., Wilder, B., Ammatelli, J., Hedrick, A., Kormos, P., Ragar, D., Havens, S., Johnson, M., and Sandusky, M.: iSnobal/smr: RCLT (Version 20250926), Zenodo [code], <https://doi.org/10.5281/zenodo.21514687>, 2025.
- Miles, A., Kirkham, J. A., Bussonnier, M., Moore, J., Papadopoulos Orfanos, D., Bourbeau, J., Fulton, A., Lee, G., Patel, Z., Bennett, D., Rocklin, M., Awa, B., Chopra, S., Abernathy, R., Kristensen, M. R. B., de Andrade, E. S., Durant, M., Schut, V., Dussin, R., Verma, S., Chaudhary, S., Barnes, C., Hamman, J., Nunez-Iglesias, J., Williams, B., Mohar, B., Noyes, C., and Bolari- inwa., E.: zarr-developers/zarr-python: v2.15.0, Zenodo [code], <https://doi.org/10.5281/zenodo.8039103>, 2023.
- Monteiro, D. and Morin, S.: Multi-decadal analysis of past winter temperature, precipitation and snow cover data in the European Alps from reanalyses, climate models and observational datasets, *The Cryosphere*, 17, 3617–3660, <https://doi.org/10.5194/tc-17-3617-2023>, 2023.
- Morin, S., Horton, S., Techel, F., Bavay, M., Coléou, C., Fierz, C., Gobiet, A., Hagenmuller, P., Lafaysse, M., Ližar, M., Mitterer, C., Monti, F., Müller, K., Olefs, M., Snook, J. S., van Herwijnen, A., and Vionnet, V.: Application of physical snowpack models in support of operational avalanche hazard forecasting: A status report on current implementations and prospects for the future, *Cold Reg. Sci. Technol.*, 170, 102910, [doi:10.1016/j.coldregions.2019.102910](https://doi.org/10.1016/j.coldregions.2019.102910), 2020.
- Mote, P. W., Hamlet, A. F., Clark, M. P., and Lettenmaier, D. P.: Declining mountain snowpack in western north America, *B. Am. Meteorol. Soc.*, 86, 39–49, <https://doi.org/10.1175/BAMS-86-1-39>, 2005.
- Mott, R., Winstral, A., Cluzet, B., Helbig, N., Magnusson, J., Mazzotti, G., Quéno, L., Schirmer, M., Webster, C., and Jonas, T.: Operational snow-hydrological modeling for Switzerland, *Front. Earth Sci.*, 11, <https://doi.org/10.3389/feart.2023.1228158>, 2023.
- Raleigh, M. S., Lundquist, J. D., and Clark, M. P.: Exploring the impact of forcing error characteristics on physically based snow simulations within a global sensitivity analysis framework, *Hydrol. Earth Syst. Sci.*, 19, 3153–3179, <https://doi.org/10.5194/hess-19-3153-2015>, 2015.
- Rasmussen, R. M., Chen, F., Liu, C., Ikeda, K., Prein, A., Kim, J., Schneider, T., Dai, A., Gochis, D., Dugger, A., Zhang, Y., Jaye, A., Dudhia, J., He, C., Harrold, M., Xue, L., Chen, S., Newman, A., Dougherty, E., Abolafia-Rosenzweig, R., Lybarger, N. D., Viger, R., Lesmes, D., Skalak, K., Brakebill, J., Cline, D., Dunne, K., Rasmussen, K., and Miguez-Macho, G.: CONUS404: The NCAR-USGS 4-km Long-Term Regional Hydroclimate Reanalysis over the CONUS, *B. Am. Meteorol. Soc.*, 104, E1382–E1408, <https://doi.org/10.1175/BAMS-D-21-0326.1>, 2023.
- Reba, M. L., Marks, D., Seyfried, M., Winstral, A., Kumar, M., and Flerchinger, G.: A long-term data set for hydrologic modeling in a snow-dominated mountain catchment, *Water Resour. Res.*, 47, <https://doi.org/10.1029/2010WR010030>, 2011.
- Serreze, M. C., Clark, M. P., Armstrong, R. L., McGinnis, D. A., and Pulwarty, R. S.: Characteristics of the western United States snowpack from snowpack telemetry (SNOTEL) data, *Water Resour. Res.*, 35, 2145–2160, <https://doi.org/10.1029/1999wr900090>, 1999.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J., Duda, M. G., Barker, D. M., and Huang, X.-Y.: A Description of the Advanced Research WRF Model Version 4.3, UCAR, <https://doi.org/10.5065/1dfh-6p97>, 2021.
- Slaughter, C. W., Marks, D., Flerchinger, G. N., Van Vactor, S. S., and Burgess, M.: Thirty-five years of research data collection at the Reynolds Creek Experimental Watershed, Idaho, United States, *Water Resour. Res.*, 37, 2819–2823, <https://doi.org/10.1029/2001wr000413>, 2001.
- Stewart, I., Cayan, D. R., and Dettinger, M.: Changes toward Earlier Streamflow Timing across Western North America, *J. Climate*, 18, 1136–1155, <https://doi.org/10.1175/JCLI3321.1>, 2005.
- Susong, D., Marks, D., and Garen, D.: Methods for developing time-series climate surfaces to drive topographically distributed energy- and water-balance models, *Hydrol. Process.*, 13, 2003–2021, [https://doi.org/10.1002/\(SICI\)1099-1085\(199909\)13:12/13<2003::AID-HYP884>3.0.CO;2-K](https://doi.org/10.1002/(SICI)1099-1085(199909)13:12/13<2003::AID-HYP884>3.0.CO;2-K), 1999.
- Thériault, J. M., Leroux, N. R., and Rasmussen, R. M.: Improvement of Solid Precipitation Measurements Using a Hot-plate Precipitation Gauge, *J. Hydrometeorol.*, 22, 877–885, <https://doi.org/10.1175/jhm-d-20-0168.1>, 2021.
- Trujillo, E. and Molotch, N. P.: Snowpack regimes of the Western United States, *Water Resour. Res.*, 50, 5611–5623, <https://doi.org/10.1002/2013WR014753>, 2014.
- Voordendag, A., Réveillet, M., MacDonell, S., and Lhermitte, S.: Snow model comparison to simulate snow depth evolution and sublimation at point scale in the semi-arid Andes of Chile, *The Cryosphere*, 15, 4241–4259, <https://doi.org/10.5194/tc-15-4241-2021>, 2021.
- Winstral, A., Elder, K., Davis, R. E.: Spatial Snow Modeling of Wind-Redistributed Snow Using Terrain-Based Parameters, *J. Hydrometeorol.*, 3, 524–538, [https://doi.org/10.1175/1525-7541\(2002\)003<0524:SSMOWR>2.0.CO;2](https://doi.org/10.1175/1525-7541(2002)003<0524:SSMOWR>2.0.CO;2), 2002.
- Winstral, A., Marks, D., and Gurney, R.: An efficient method for distributing wind speeds over heterogeneous terrain, *Hydrol. Process.*, 23, 2526–2535, <https://doi.org/10.1002/hyp.7141>, 2009.
- Xia, Y., Mitchell, K., Ek, M., Sheffield, J., Cosgrove, B., Wood, E., Luo, L., Alonge, C., Wei, H., Meng, J., Livneh, B., Lettenmaier, D., Koren, V., Duan, Q., Mo, K., Fan, Y., and Mocko, D.: Continental-scale water and energy flux analysis and validation for the North American Land Data Assimilation System project phase 2 (NLDAS-2): 1. Intercomparison and applica-

tion of model products, *J. Geophys. Res.-Atmos.*, 117, D03109, <https://doi.org/10.1029/2011JD016048>, 2012.

Zeng, X., Broxton, P., and Dawson, N.: Snowpack Change From 1982 to 2016 Over Conterminous United States, *Geophys. Res. Lett.*, 45, 12940–12947, <https://doi.org/10.1029/2018GL079621>, 2018.