



Supplement of

History of anthropogenic Phosphorus inputs (HaPi) to the terrestrial biosphere from 1860 to 2020

Zihao Bian et al.

Correspondence to: Zihao Bian (zhbian@nynu.edu.cn), Hao Shi (haoshi@rcees.ac.cn), and Hanqin Tian (hanqin.tian@bc.edu)

The copyright of individual parts of the supplement might differ from the article licence.

Contents:

Figure S1. Comparison of manure N:P ratio maps derived from three datasets in 1961 (a, d, g), 1980 (b, e, h), and 2010 (c, f, i). The first, second, and third rows show the GLW3, AGLW, and merged datasets, respectively. Columns correspond to 1961, 1980, and 2010.

Figure S2. Global patterns of P fertilizer application on pasture in the 1960s, 1980s, 2000s, and 2010s.

Figure S3. Global patterns of manure P application on pasture in the 1860s, 1910s, 1960s, and 2010s.

Figure S4. Global patterns of manure P deposition on pasture in the 1860s, 1910s, 1960s, and 2010s.

Figure S5. Comparisons of P fertilizer inputs of the HaPi dataset with national statistics in (a) China and (b) the U.S.

Figure S6. Changes in annual P inputs (excluding atmospheric P deposition) and P input rates with growth of GDP per capita across cropland, pasture, and grassland in 18 different regions.




Figure S1. Comparison of manure N:P ratio maps derived from three datasets in 1961 (a, d, g), 1980 (b, e, h), and 2010 (c, f, i). The first, second, and third rows show the GLW3, AGLW, and merged datasets, respectively. Columns correspond to 1961, 1980, and 2010.




Figure S2. Global patterns of P fertilizer application on pasture in the 1960s, 1980s, 2000s, and 2010s.




Figure S3. Global patterns of manure P application on pasture in the 1860s, 1910s, 1960s, and 2010s.




Figure S4. Global patterns of manure P deposition on pasture in the 1860s, 1910s, 1960s, and 2010s.




Figure S5. Comparisons of P fertilizer inputs of the HaPi dataset with national statistics in (a) China and (b) the U.S.




Figure S6. Changes in annual P inputs (excluding atmospheric P deposition) and P input rates with growth of GDP per capita across cropland, pasture, and grassland in 18 different regions. The colored trend lines are fitted by generalized additive models. The 18 regions include the USA, Canada (CAN), Central America (CAM), northern South America (NSA), Brazil (BRA), southwestern South America (SSA), Europe (EU), Northern Africa (NAF), Equatorial Africa (EQAF), Southern Africa (SAF), Russia (RUS), Central Asia (CAS), the Middle East (MIDE), China (CHN), Korea and Japan (KAJ), South Asia (SAS), Southeast Asia (SEAS), and Oceania (OCE). For visual clarity, not all region names are shown in each panel.