



A review of Last Interglacial sea-level proxies in South Asia and the Indian Ocean

Mubashir Ali^{1,2}, Ciro Cerrone^{1,3,4}, and Alessio Rovere^{1,5}

¹Department for Environmental Sciences, Informatics and Statistics (DAIS),
Ca' Foscari University of Venice, Via Torino 155, 30172 Venice, Italy

²Department of Earth and Environmental Sciences, University of Milano-Bicocca, 20126 Milano, Italy

³International Centre for Climate Change Research and Studies (CSRCC),
Franchetti Palace, San Marco 2847, 30124 Venice, Italy

⁴Department of Mathematics, Informatics and Geosciences, University of Trieste, 34128 Trieste, Italy

⁵MARUM, Center for Marine Environmental Sciences, University of Bremen,
Leobener Str. 8, 28359 Bremen, Germany

Correspondence: Alessio Rovere (alessio.rovere@unive.it)

Received: 24 February 2026 – Discussion started: 22 June 2026

Revised: 17 September 2026 – Accepted: 3 October 2026 – Published: 20 October 2026

Abstract. In this work, we present a standardized database of Last Interglacial (LIG, ~ 132 –80 ka) sea-level proxies across the South Asian and Indian Ocean region, including the coasts of Iran, Oman, India, the Maldives, and the Cocos (Keeling) Islands. The database stems from a systematic review of the data included in 17 peer-reviewed studies, structured according to the World Atlas of Last Interglacial Shorelines (WALIS), and is available as Ali et al. (2026, <https://doi.org/10.5281/zenodo.18512769>). Within this framework, a total of 54 sea-level index points were re-evaluated to recalculate paleo-RSL elevation and associated uncertainties. This was achieved through updated determinations of their indicative meaning (i.e., reference water level and indicative range), considering the original stratigraphic context, elevation measurement techniques, depositional environments, and chronological constraints. The chronological framework relies on U-Th, luminescence, electron spin resonance, and cosmogenic nuclide dating. These data are characterized by regional differences in the elevation of past relative sea-level proxies that are driven primarily by tectonics, dynamic topography, and glacio-hydro-isostatic forcings. On the active margins of southern Iran and northern Oman, marine terraces dated to Marine Isotope Stage (MIS) 5e (~ 125 ka), 5c (~ 100 ka), and 5a (~ 80 ka) are preserved at 20–60 m a.s.l., reflecting uplift linked to the Zagros foreland and Makran subduction system at least since the Last Interglacial. On the coasts of western and southern India, MIS 5e relative sea-level proxies are found at 1–4 m a.s.l., more in line with near-zero vertical land motion. On Indian Ocean islands, such as the Maldives and Cocos (Keeling) Islands, MIS 5e reefs are now submerged at depths of 10–17 m b.s.l., indicating subsidence since the LIG.

1 Introduction

This study reviews evidence of relative sea-level (RSL) proxies from the Last Interglacial (LIG), broadly corresponding to the Marine Isotope Stage (MIS) 5 (~ 132 –80 ka), across the South Asia Indian Ocean region, including the Maldives, India, Oman, and the southern coasts of Iran (Fig. 1). This study focuses on a region that, in previous global compilations of LIG sea-level indicators, was a geographic gap

(Rovere et al., 2023) and serves as an update and extension to earlier datasets, such as that presented by Pedoja et al. (2014). Regionally, this work complements a recent World Atlas of Last Interglacial Shorelines (WALIS) compilation along East Africa and the Western Indian Ocean (Boyden et al., 2021).

The database was assembled by systematically extracting, standardising, and re-evaluating data and metadata from




Figure 1. Sea-level index points re-evaluated and standardized in South Asia and the Indian Ocean region. Colored dots show sea-level datapoints included in the WALIS database, with colors indicating different elevation ranges. Red lines represent tectonic plate boundaries from Bird (2003), modified by H. Hugo Ahlenius (Nordpil, GitHub: <https://github.com/fraxen/tectonicplates>, last access: 17 October 2026). Basemap assembled from datasets provided by the Global Multi-Resolution Topography (GMRT) (Ryan et al., 2009).

peer-reviewed publications providing stratigraphic records and chronological constraints related to LIG sea-level variations in the region. All sea-level indicators and associated chronological data were harmonised using the WALIS platform (Garzón and Rovere, 2024; Rovere et al., 2020, available at: <https://warmcoasts.eu/walis.html>, last access: 17 October 2026). The database was exported from the WALIS interface in spreadsheet format and is available as an Excel file via Zenodo under a CC-BY license (Ali et al., 2026, <https://doi.org/10.5281/zenodo.18512769>). Table 2, included in the text, provides a summary of the elevation, chronological constraints, references, and quality assessments for each RSL data point included in the investigated area.

Altogether, data from 17 peer-reviewed studies were compiled, yielding 54 relative sea level index points (SLIPs). The database compilation was finalized in September 2025; accordingly, data from papers published after this date were not

included. Each SLIP was assigned one or more chronological constraints. The age constraints for all SLIPs were taken from previously published original papers. In particular, we inserted in the database data and metadata related to samples dated with luminescence ($n=7$), U-Th ($n=20$ on corals, $n=4$ on mollusks), electron spin resonance (ESR; $n=22$), and one cosmogenic radionuclide (^{10}Be) techniques.

A brief description of the types of sea-level indicators within this study is provided in the following section. This is followed by a discussion of elevation measurement methods, sea-level reference datums, and main dating techniques used by the original authors. Finally, the data points are described and organized according to national and regional boundaries.

2 Types of sea-level indicators, positioning and sea-level datums

In this study, seven main types of SLIPs were identified (Table 1). These include three depositional (beach deposit or beachrock, beach ridge, beachrock), two biological (coral reef terrace, shallow or intertidal marine fauna) and two geomorphological indicators (marine terrace and tidal notch). To establish a connection between these indicators and calculate paleo RSL, we employed the concept of indicative meaning, a methodological framework that quantifies how specific sea-level markers relate to tidal datums (Khan et al., 2019). This framework defines that each indicator must be assigned an indicative meaning, which is decomposed into two values: the indicative range (IR), which captures the vertical range at which a SLIP can occur according to its modern analog (Khan et al., 2019), and the reference water level (RWL), which is the midpoint of the IR. Originally developed for Holocene studies (Shennan, 1982, 1986, 1989, 2015; Shennan et al., 1983), the indicative meaning approach has more recently been applied to Pleistocene sea-level proxies (Rovere et al., 2016b), allowing for consistent interpretation of SLIPs across different time periods. To quantify the indicative meaning, here we used the IMCalc software (Lorscheid and Rovere, 2019), which generates *ex situ* estimates of RWL and IR using global tidal and wave datasets. This approach is particularly useful when direct, site-specific modern analog data from primary sources are not available, as is the case for the works published in our study area. While elevation measurement techniques are reported in most of the studies included in our review, a small subset of works lacks this information (Brückner, 1989; Reyss et al., 1999; Normand et al., 2019; Goswami et al., 2019). Specifically, for exactly 5 RSL datapoints in Iran, the elevation measurement technique was not reported in the original papers (Rovere et al., 2016b). Among the six measurement techniques recorded in this database, the vast majority of sea-level index point elevations were determined using either barometric altimeters ($n = 25$) or tape and Abney (a handheld surveying instrument used to measure slope angles, gradients, and vertical heights) levelling ($n = 11$). Although these relatively older surveying techniques, along with the undocumented methods, are inherently associated with lower resolution and potentially substantial measurement uncertainties, they represent the only historical data available for these areas. The database also includes modern and higher-precision techniques. These consist of: (i) differential GPS (dGPS) alone ($n = 2$) in the Sur lagoon area (Oman; Mauz et al., 2015; Falkenroth et al., 2020), which offers the highest operational accuracy; (ii) a combination of dGPS, hand level, and tape ($n = 2$) in Oman (Hoffmann et al., 2020; Decker et al., 2024; Falkenroth et al., 2019); (iii) handheld GPS ($n = 6$) in Iran and the Maldives (Pirazzoli et al., 2004; Gischler, 2008; Gischler et al., 2008, 2018; Normand et al., 2019; Goswami et al., 2019); and (iv) the distance calculated from the top of

a drill core ($n = 3$). A couple of works (Brückner, 1989; Goswami et al., 2019) do not report any elevation measurement technique for the (i) Porbandar and Tunkra and (ii) Tamil Nadu areas, respectively. Since most studies did not report exact site coordinates, we estimated them using place names via geolocation tools (e.g., Google Maps) or by georeferencing original maps in QGIS.

3 Relative sea-level data

In the following sections we describe the data we revised, organized by country in a clockwise order starting from Iran, and, where applicable, by subordinate administrative units such as states, regions, or provinces. Throughout the text, the elevation of SLIPs is expressed as “a.s.l.” (above present sea level) or “b.s.l.” (below present sea level). A summary of sites, associated paleo RSL, and chronological constraints associated to each site are detailed in Table 2. Sea-level indicators compiled in the database are referenced by their WALIS RSL ID (hereafter RSL_NUMBER). This identifier, displayed in the first column of the “RSL proxies” spreadsheet, is automatically generated and uniquely assigned to each record within WALIS.

3.1 Iran

3.1.1 Southern Mainland Iran

The Northern margin of the Persian Gulf in southern Iran hosts a broad coastal sector that provides important archives for Quaternary sea-level reconstructions (Fig. 2). The Haleh coastal sector (Fig. 1B), situated along the southern Iranian mainland, provides important archives of Quaternary sea-level history. This sector comprises a series of stepped marine terraces that record repeated interglacial highstand phases, including the Last Interglacial (LIG) and older interglacials, separated by distinct relative elevation differences (e.g., Reyss et al., 1999). Similarly, the Jazeh coastal area (Fig. 2) preserves a well-defined sequence of Quaternary marine terraces, offering an additional, complementary record for reconstructing relative sea-level evolution in the region (Reyss et al., 1999). Moving eastward, the Konarak, Ramin, and Lipar areas, along the southern coastal margin of Iran (Fig. 2), also host late Quaternary RSL indicators (e.g., Normand et al., 2019). Together, these coastal sites provide critical datasets for understanding the timing, magnitude, and variability of past sea-level changes along this area.

Haleh

Along the Haleh coastal sector, a staircase of well-preserved marine terraces is exposed, reaching elevations exceeding 100 m a.s.l. (Reyss et al., 1999). The terrace sequence includes a highly elevated coral-bearing unit at 100–110 m a.s.l., a narrower abrasion terrace at 85 m a.s.l., and a

Table 1. Summary of sea-level proxies included in the South Asian and Indian Ocean database. Abbreviations: MLLW – Mean Lower Low Water; MHHW – Mean Higher High Water; HAT – Highest Astronomical Tide.

SLIP	Description of RSL indicator	Indicative range	Indicator reference(s)	Number of occurrences in the database
Beach deposit or beachrock	Definition by Mauz et al. (2015): “Beachrocks are lithified coastal deposits that are organized in sequences of slabs with seaward inclination generally between 5° and 15°.”	Ordinary berm to breaking depth	Mauz et al. (2015), Rovere et al. (2016b)	3
Beach ridge	Definition by Otvos (2000): “[beach ridges are] stabilized, relict intertidal and supratidal, eolian and wave-built shore ridges that may consist of either siliciclastic or calcareous clastic matter of a wide range of clasts dimensions, from fine sand to cobbles and boulders.”	Storm wave swash height to ordinary berm	Otvos (2000), Rovere et al. (2016b)	1
Beachrock	Cement: isopachous or miniscus of HMC or aragonite; granular, bladed or fibrous crystals Bedding: symmetrical ripples or horizontal plane-parallel laminae or planar and trough cross-beds	HAT (or spray zone) to MLLW	Mauz et al. (2015)	2
Coral reef terrace (general definition)	Coral-built flat surface, corresponding to shallow-water reef terrace to reef crest.	MLLW to breaking depth	Rovere et al. (2016b), Lorscheid and Rovere (2019)	32
Marine Terrace	Definition by Pirazzoli (2005): “Any relatively flat surface of marine origin.”	Storm wave swash height to breaking depth	Pirazzoli (2005), Rovere et al. (2016b)	10
Shallow or intertidal marine fauna	Marine fauna usually associated with very shallow water or intertidal environments	Based on the upper and lower limits of living modern analog faunas		5
Tidal notch	Definition by Antonioli et al. (2015): “Indentations or undercuttings cut into rocky coasts by processes acting in the tidal zone (such as tidal wetting and drying cycles, bioerosion, or mechanical action).”	MHHW to MLLW	Antonioli et al. (2015), Rovere et al. (2016b)	1

broad reef platform that gently slopes from 33 to 23 m a.s.l., before merging seaward into a low, cliffed bench situated between 5 and 2.5 m a.s.l. Among these four terrace levels, the third terrace (33–23 m) is particularly well constrained in terms of its elevation, morphology, stratigraphic context and age. Near the seaward edge of this terrace, which represents a previous reef platform, coral colonies preserved in growth position have been sampled and dated using U-Th techniques, yielding an age of 111 ± 4 ka (Reyss et al., 1999). This age corresponds to MIS 5e, with an uplift rate calculated from Reyss et al. (1999) of $\sim 0.2 \text{ mm yr}^{-1}$ since the LIG and the associated SLIP has been included in WALIS as RSL_4527. The lithofacies assemblages of the LIG terrace

are typical of shallow-marine reef-flat environments, including coral framestones interbedded with lumachelle-bearing limestones and bioclastic sands, which rest unconformably over Mio-Pliocene marls.

Jazeh

Three major terrace levels are exposed along the coastal area of Jazeh (Fig. 2): (i) a higher terrace at 60–65 m, (ii) a broad intermediate terrace sloping between 35 and 25 m, and (iii) a narrow low bench near the modern shoreline recorded by Reyss et al. (1999). The RSL indicators constrained to the Last Interglacial are preserved within the intermediate ter-

Table 2. Summary of SLIPs and dated samples/chronostratigraphic constraints for the South Asia and Indian Ocean region. Legend for SLIP types: CRT = Coral Reef Terrace; MT = Marine Terrace; BD = Beach deposit or beachrock; SIMF = Shallow or intertidal marine fauna; TN = Tidal notch; and BR = Beach Ridge. * represent original reported corrected age.

Walis ID	Site name	Latitude	Longitude	SLIP type	Elevation (m)	Paleo RSL (m)	MIS	Age (ka)	Original dating ID	References
RSL_4527	Haleh	27.396944	52.593889	CRT	23 ± 5	23.63 ± 5	MIS 5e	111 ± 4	HA7	Reyss et al. (1999)
RSL_4528	Jazeh	26.826667	53.528056	MT	9 ± 5	8.77 ± 5	MIS 5e	130 ± 13	JA8	
RSL_4529	Jazeh	26.826667	53.528056	MT	21 ± 5	20.76 ± 5	MIS 5e	118 ± 10	JA9	
RSL_4507	Kish Island	26.535278	54.025556	CRT	5.2 ± 2	5.89 ± 2	MIS 5e	126 ± 14	OKI-08	Pirazzoli et al. (2004), Preusser et al. (2003)
RSL_4508	Kish Island	26.535278	54.025556	CRT	6.7 ± 2	7.39 ± 2	MIS 5e	122 ± 10	OKI-10	
RSL_4509	Kish Island	26.535278	54.025556	CRT	8.1 ± 2	8.79 ± 2	MIS 5e	117 ± 18	OKI-11	
RSL_4510	Kish Island	26.535278	54.025556	CRT	11.6 ± 2	12.29 ± 2	MIS 5e/MIS 5c	100 ± 9	OKI-14	
RSL_4511	Kish Island	26.506111	53.999444	CRT	18 ± 2	18.69 ± 2	MIS 5e/MIS 5d	115 ± 15	OKI-15	
RSL_4512	Kish Island	26.506111	53.999444	CRT	18 ± 2	18.69 ± 2	MIS 5e	126 ± 12	OKI-16	
RSL_4513	Kish Island	26.506111	53.999444	CRT	18 ± 2	18.69 ± 2	MIS 5e/MIS 5d	114 ± 16	OKI-17	
RSL_4514	Kish Island	26.536389	53.958889	CRT	32 ± 1	32.69 ± 1	MIS 5	148 ± 20	OKI-23	
RSL_4515	Kish Island	26.536389	53.958889	CRT	32 ± 1	32.69 ± 1	MIS 5e	128.5 ± 4	OKI-24	
RSL_4515	Kish Island	26.536389	53.958889	CRT	32 ± 1	32.69 ± 1	MIS 5e	119 ± 10	OKI-24	
RSL_4516	Kish Island	26.499167	53.966944	CRT	6 ± 2	6.69 ± 2	MIS 5e	116 ± 33	OKI-03	Pirazzoli et al. (2004)
RSL_4516	Kish Island	26.535278	54.025556	CRT	6 ± 2	6.69 ± 2	MIS 5e	248 ± 15	OKI-03	
RSL_4536	RSL_4536	26.535278	54.025556	CRT	8.4 ± 2	9.08 ± 2	MIS 5e	143 ± 12	OKI-12	
RSL_4517	Qesham	26.926111	56.211111	CRT	3 ± 2	3.79 ± 2	MIS 5e	133 ± 10	OQE-29	Pirazzoli et al. (2004)
RSL_4518	Qesham	26.926111	56.211111	CRT	12 ± 5	12.79 ± 5	MIS 5	139 ± 6	QE100	Reyss et al. (1999), Pirazzoli et al. (2004)
RSL_4519	Qesham	26.991667	56.217222	CRT	20 ± 1	20.8 ± 1	MIS 5e	133 ± 13	OQE-51	Pirazzoli et al. (2004)
RSL_4520	Qesham	26.990556	56.216667	CRT	21 ± 5	21.8 ± 5	MIS 5e	127 ± 6	QE102	Reyss et al. (1999) Pirazzoli et al. (2004)
RSL_4521	Qesham	26.928056	56.216944	CRT	26 ± 2	26.79 ± 2	MIS 5e	126 ± 11	OQE-40	Pirazzoli et al. (2004)
RSL_4522	Qesham	26.893333	56.197778	CRT	26 ± 5	26.78 ± 5	MIS 5c	104 ± 5	QE54	Reyss et al. (1999) Pirazzoli et al. (2004)
RSL_4523	Qesham	26.929444	56.238056	CRT	30 ± 2	30.79 ± 2	MIS 5e	123 ± 18	OQE-33	Pirazzoli et al. (2004)
RSL_4524	Qesham	26.991667	56.217222	CRT	30 ± 2	30.8 ± 2	MIS 5e	115 ± 9	OQE-53	
RSL_4525	Qesham	26.929444	56.238056	CRT	30 ± 2	30.79 ± 2	MIS 5e	137 ± 11	OQE-34	
RSL_4526	Qesham	26.991667	56.217222	CRT	25 ± 2	25.8 ± 2	MIS 5e	138 ± 12	OQE-52	
RSL_4532	Konarak T3	25.343285	60.305565	MT	32.3 ± 2	31.76 ± 2.7	MIS 5e	141.3 ± 12	RN17-48	Normand et al. (2019)
RSL_4531	Ramin T1	25.261995	60.827053	MT	59.5 ± 2	58.93 ± 2.8	MIS 5e	140.21 ± 12	RN17-43	
RSL_4530	Lipar T3	25.249691	60.841739	MT	58.5 ± 2	57.93 ± 2.8	MIS 5e	130.28 ± 10	RN17-30	
RSL_4501	Oman (Terrace T3)	22.911695	59.205471	BD	36.7 ± 1	36.44 ± 1.3	MIS 5	97 ± 6	DEL4-Q	Hoffmann et al. (2020)

Table 2. Continued.

Walis ID	Site name	Latitude	Longitude	SLIP type	Elevation (m)	Paleo RSL (m)	MIS	Age (ka)	Original dating	References
RSL_4501	Oman (Terrace T3)	22.911695	59.205471	BD	36.7 ± 1	36.44 ± 1.3	MIS 5	118 ± 8	DEL5-Q	Decker et al. (2024) Falkenroth et al. (2019)
RSL_4502	Sur Lagoon	22.5821	59.4518	TN	3.93 ± 0.12	4.04 ± 0.94	MIS 5	< 371 > 80	TD13-009	Falkenroth et al. (2020)
RSL_4541	Sur Lagoon	22.567297	59.507797	BD	3.73	3.76 ± 1.3	MIS 5	80 ± 3	LV565	Mauz et al. (2015)
RSL_4540	SAURASHTRA COAST	21.634782	69.610289	SIMF	4	5.53 ± 0.4	MIS 5	95 (72.6–123.2 ka*)		Por 1c-M/1:1
RSL_4540	SAURASHTRA COAST	21.634782	69.610289	SIMF	4	5.53 ± 0.4	MIS 5	97.6 ± 10	Tun 2d (D-1292)	Brückner (1989)
RSL_4540	SAURASHTRA COAST	21.634782	69.610289	SIMF	4	5.53 ± 0.4	MIS 5	94.9 ± 10	Por 1c-S (D-1278)	
RSL_4540	SAURASHTRA COAST	21.634782	69.610289	SIMF	4	5.53 ± 0.4	MIS 5	105 ± 10	Por 1b (D-1277)	
RSL_4492	Cape Comorin	8.085205	77.496911	BD	1.2 ± 0.5	1.53 ± 1.7	MIS 5e/5c	112 ± 10	Cape C	Banerjee (2000)
RSL_4493	Idindakarai	8.182555	77.755555	SIMF	3.4 ± 0.5	3.38 ± 5	MIS 5e	124 ± 11	Vija52	Brückner (1989)
RSL_4494	Manappad	8.373333	78.066666	BD	2.5 ± 0.5	2.44 ± 1.6	MIS 5e	120 ± 8	Tun 2b	
RSL_4489	Rameswaram	9.313888	79.333333	CRT	1.1 ± 0.5	1.83 ± 0.7	MIS 5e	125.6 ± 8	PKB 8a	
RSL_4490	Rameswaram	9.313888	79.333333	CRT	2.9 ± 0.5	3.63 ± 0.7	MIS 5c	104 ± 10	R5	Banerjee (2000)
RSL_4491	Rameswaram	9.313888	79.333333	CRT	2.9 ± 0.5	3.63 ± 0.7	MIS 5c	116 ± 10	R3	
RSL_4503	Rameswaram	9.313888	79.333333	CRT	2.9 ± 0.5	3.63 ± 0.7	MIS 5c	92 ± 6	R6	
RSL_4504	Rameswaram	9.313888	79.333333	CRT	2.4 ± 0.5	3.13 ± 0.7	MIS 5e/MIS 5d	112 ± 8	Vadakkadu 4	Banerjee (2000)
RSL_4504	Rameswaram	9.313888	79.333333	CRT	2.4 ± 0.5	3.13 ± 0.7	MIS 5e	139 ± 10	Vadakkadu 4	Brückner (1989)
RSL_4505	Rameswaram	9.313888	79.333333	CRT	2.9 ± 0.5	3.63 ± 0.7	MIS 5c	104 ± 10	R7	Banerjee (2000)
RSL_4506	Rameswaram	9.313888	79.333333	CRT	2.4 ± 0.5	3.13 ± 0.7	MIS 5c	108 ± 3	01/97	
RSL_4539	Vetalkaranippu	10.553467	79.83545	BR	−13.5	12.59 ± 0.2	MIS 5e	121 ± 11	VK112F	Goswami et al. (2019)
RSL_4495	Rasdhoo Atoll	4.291922	72.940327	CRT	−15.2	−14.18 ± 0.6	MIS 5e	136.9 ± 2	1–14.2	Gischler (2008),
RSL_4537	Rasdhoo Atoll	4.291922	72.940327	CRT	−16.3	−15.27 ± 0.6	MIS 5e	114.3 ± 2	1–15.3	Gischler et al. (2008, 2018)
RSL_4538	Rasdhoo Atoll	4.291922	72.940327	CRT	−17	−15.97 ± 0.6	MIS 5e	117.3 ± 3	1–16.0	
RSL_4496	Keeling	−12.222222	96.999999	CRT	−10.5 ± 1	−9.19 ± 1	MIS 5e	123 ± 7	sample without cement	Woodroffe et al. (1991),
RSL_4496	Keeling	−12.222222	96.999999	CRT	−10.5 ± 1	−9.19 ± 1	MIS 5e	118 ± 7	Bulk sample	Woodroffe (2005)




Figure 2. Map of the southern coast of Iran and northern Oman, showing the locations of SLIPs from this region. Basemap assembled from datasets provided by the Global Multi-Resolution Topography (GMRT).

race, where in situ *Porites* corals were sampled and dated by U-Th techniques. Two of these in situ corals yielded U-Th ages of 121 ± 6 and 117 ± 6 ka (WALIS IDs RSL_4528 and RSL_4529, respectively), consistent with deposition during MIS 5e, which allow Reyss et al. (1999) to calculate an uplift rate of $\sim 0.2 \text{ mm yr}^{-1}$. The dated corals are associated with framestone lithofacies interbedded with lumachelle-rich limestones and bioclastic sands, suggesting shallow reef-flat to nearshore depositional environments.

Konarak

The Konarak T3 terrace, located on the southeastern Makran coastline (Fig. 2), is characterized by flat-topped, bedrock deposits interbedded with various-sized boulders. In WALIS, this terrace is catalogued as RSL_4532, with a shoreline angle elevation at 32 m a.s.l. Optically stimulated luminescence (OSL) dating of feldspar from the terrace T3 yielded a fading-corrected age of 141.3 ± 12.1 ka, which falls within the broader range of MIS 5e (Normand et al., 2019). In addition, the combined morphological expression, facies, and chronological evidence provide strong grounds for assigning T3 to the Last Interglacial highstand. The Konarak T3 terrace elevation allows Normand et al. (2019) to calculate an uplift rate for this terrace between of 0.27 mm yr^{-1} since the Last Interglacial.

Ramin

The Ramin coastal site, situated east of Chabahar on the Makran margin (Fig. 2), features a prominent marine terrace (T1) preserved at an elevation of 59.5 ± 2.0 m a.s.l., recorded in the WALIS database as RSL_4531. At this site, terrace T1 is composed of trough-cross-stratified marine deposits. Luminescence dating of sediments associated with the T1 platform yielded an age of 140.2 ± 12.5 ka, placing it within the MIS 5e (Normand et al., 2019). Conversely, a lower inset bench at 3 m a.s.l. yielded an OSL of 78 ka, attributing this lower level to MIS 5a (Normand et al., 2019), which was already added in WALIS by Thompson and Crevelling (2021) with label RSL_3534. The WALIS record referred to T1 (RSL_4531) and the geomorphic, chronological, and stratigraphic evidence at Ramin T1 together support its use as a significant but tectonically upward-displaced RSL indicator of the Last Interglacial.

Lipar

The Lipar coastal plain, located along the eastern Makran margin of southeastern Iran, host isolated, flat-topped, marine terrace deposits composed of beachrocks, providing key Late Quaternary sea-level records for the Oman Sea-Arabian Sea region (Fig. 2). Three main terrace levels have been described at this site by Normand et al. (2019). The lowest two

levels (T1 and T2, RSL_3532 and RSL_3533 by Thompson and Creveling, 2021) occur between 20 and 50 m a.s.l. and consist of beachrock deposits interbedded with various-sized boulders and shell fragments. The sedimentary structures are dominated by high- to low-angle trough cross-stratification, which shows that the deposition of Lipar T1 and T2 deposits occurs in high-energy marine environments. OSL dating provides the age constraints for T1 and T2, yielding ages between 73 and 80 ka and correlating them with MIS 5a (Normand et al., 2019; Thompson and Creveling, 2021). Further inland and higher on the coastal slope, the most prominent surface is the Lipar T3 terrace (RSL_4530), occurring up to 58.5 m a.s.l. The lithological composition is similar to that of Lipar T1 and T2, and is mainly composed of beach rock deposits interbedded with various-sized boulders and shell fragments. The sedimentary structures include high- to low-angle trough cross-stratification, which characterizes the deposition of Lipar T3 and indicates a high-energy marine environment. The OSL-based chronology yielded an age of 130.28 ± 10 ka, corresponding to MIS 5e. Consistent ages make it one of the clearest RSL indicators of the Last Interglacial highstand along the Makran coast, archived in the WALIS database as RSL_4530. Based on the combined elevations and chronological constraints of MIS 5e (T3) and MIS 5a (T1–T2) terraces, Normand et al. (2019) calculated an uplift rate of approximately $0.25\text{--}0.57$ mm yr^{−1} for the Lipar area since the Last Interglacial.

3.1.2 Iran's islands

Most of the late Quaternary RSL indicators reported in the literature for the southern coast of Iran originate from its islands, including Kish and Qeshm islands, which are in the vicinity of the Hormuz Strait (Fig. 2). Together, both Kish and Qeshm islands hold the best documented record of LIG sea level fluctuations from the southern Iranian region. A brief description of these areas is provided below.

Kish Island

Kish Island, situated in the southeastern Persian Gulf about 20 km off the Iranian mainland, preserves a staircase of raised coral reef terraces. Kish Island, rising to c. 32 m a.s.l., is composed predominantly of Pleistocene reef limestones that have been systematically mapped and chronologically studied in several studies (Preusser et al., 2003; Pirazzoli et al., 2004). At least two reef terrace levels occur. In particular, the lower one occurs between 5 and 12 m a.s.l. (RSL_4507, RSL_4508, RSL_4509, RSL_4510, RSL_4516, and RSL_4536), while the higher one is located between 18 and 32 m a.s.l. (WALIS IDs 4511–4515). Importantly, both terrace levels contain sea-level indicators dated to MIS 5. These terraces are composed of coral framestones and boundstones dominated by *Platygyra*, *Favia*, *Porites*, and *Goniastrea*. These coral colonies are preserved in growth

position and are overlain by reef debris and cemented bioclastic sands. Such lithofacies are characteristic of shallow subtidal to intertidal reef accretion during stable highstands. Chronological constraints on the lower terrace are primarily based on ESR and U-Th dating of corals. The ESR results consistently fall between 100 and 126 ka, with most values clustering at 122–126 ka, consistent with deposition during the Last Interglacial (Preusser et al., 2003). At an elevation of 6 m a.s.l., one U-Th date on the coral species *Platygyra* yielded an anomalous age of 248 ± 15 ka, likely due to re-worked material from higher (and older) terraces. However, the same sample produced an ESR age of 116 ± 33 ka, which is consistent with deposition during MIS 5e (Pirazzoli et al., 2004).

The coral reef terraces occurring between 18 and 32 m a.s.l. (WALIS IDs 4511–4515) yielded ESR and U-Th ages in the range of 114–126 ka, again strongly corresponding to MIS 5e, but some samples from these terraces also give older ages consistent with MIS 7 (200–240 ka), showing that Kish Island may host a polycyclic reef record in which different interglacial reef surfaces have been preserved at similar elevations. The estimated uplift rates of $0.13\text{--}0.24$ mm yr^{−1} for the island are determined from the combined MIS 5 and MIS 7 ages by Pirazzoli et al. (2004) and Preusser et al. (2003).

Qeshm Island

Qeshm Island, the largest island in the Persian Gulf, located within the Strait of Hormuz (Fig. 2), preserves a prominent sequence of Quaternary marine terraces comprising up to 18 distinct terrace levels, with the highest reaching an elevation of 220 m a.s.l. Previous works have mapped and dated all these terraces (Reyss et al., 1999; Pirazzoli et al., 2004). The most reliable RSL indicators of the Last Interglacial on Qeshm Island are reef terraces between 10 and 26 m a.s.l. (from RSL_4517 to RSL_4526). These deposits consist of reefal limestones dominated by *Porites* and *Favia* corals, interbedded with lumachelle-rich limestones, barnacle-bearing beachrocks, and cemented bioclastic sands. The coral assemblages are typically preserved in situ. Such lithofacies are consistent with shallow subtidal to intertidal reef accretion and abrasion platforms.

Chronological control is obtained from a combination of U-Th and ESR dating across Qeshm Island. U-Th ages from the terraces at 12–21 m a.s.l., yielded results of 139 ± 6 and 127 ± 6 ka, consistent with MIS 5e deposition. One terrace on Qeshm Island at 26.5 m a.s.l. yielded a U-Th age of 104 ± 4 ka from corals in growth position, placing it in the MIS 5c age and positioning that terrace above the MIS 5e record observed by other studied sites on the island. This may be due to local tectonic activities (RSL_4522; Pirazzoli et al., 2004; Reyss et al., 1999). The overall ages at different elevations are further complemented by ESR dating (Pirazzoli et al., 2004), which confirmed age ranges 110–133 ka

for corals at 10–21 m a.s.l., thereby reinforcing attribution of these reef terraces to MIS 5e. RSL indicators, such as corals in Qeshm, which are above 21 m a.s.l., were dated using ESR, and yielded 115–138 ka with an average age uncertainty of 12 ka, indicating that they may also correspond to MIS 5.

In addition, ESR results highlighted diagenetic complexities in some samples, but the overlap between U-Th and ESR datasets provides robust chronological evidence for Last Interglacial reef growth on Qeshm, with an estimated average uplift rates of 0.2 mm yr^{-1} since MIS 5 for that area (Reyss et al., 1999). Thus, Qeshm Island provides a well-documented chronological control along a stratigraphically coherent record of the Last Interglacial in the Persian Gulf. Terraces at approximately 10–20 m a.s.l., constrained by overlapping U-Th and ESR ages of 110–139 ka, record reef accretion during MIS 5e.

3.2 Oman

3.2.1 Northern Oman T3

The northeastern coast of Oman, between Quriyat and Qalhat (Fig. 2), exposes laterally extensive and morphologically well-preserved marine terraces. This sequence was systematically documented by Hoffmann et al. (2013a, b, 2020), who recognized at least twelve levels, among which the lowest three (T1–T3) are the most prominent and best dated. The T3 terrace, recorded in WALIS as RSL_4501, is correlated with the Last Interglacial and occurs consistently at elevations of 36 m a.s.l. Chronological control for T3 is provided by two OSL dating (samples DEL4-Q and DEL5-Q, yielding 118 and 97 ka, respectively), whereas cosmogenic nuclide exposure dates yield significantly older ages. Sedimentologically, T3 comprises eight lithofacies grouped into six facies associations, ranging from foreshore cross-bedded sandstones and upper shoreface coarse conglomerates to mollusk-rich cemented beachrocks and carbonate-cemented gravels, deposited in a high-energy environment (Falkenroth et al., 2019).

Two lower terraces, T2 (2 to 4 m a.s.l.) and T1 (1 to 2 m a.s.l.), are tentatively attributed to MIS 5c and MIS 5a, respectively. Together with T3, this terrace sequence provides a detailed record of Last Interglacial sea-level oscillations in Oman, within a tectonically uplifted setting. The lithofacies confirm deposition in shallow coastal environments, providing precise sea-level index points.

3.2.2 Sur lagoon

Sur Lagoon is located on the northeastern coast of Oman, just below the Qalhat area (Fig. 2), and presents a laterally continuous bioerosional notch associated with the Last Interglacial. The notch, developed in fanglomerate bedrock, encircles the lagoon at a consistent elevation. It hosts an assemblage of biological markers, including mussel borings

(*Gastrochaenolites* sp.), sponge borings (*Entobia* sp.), barnacle colonies, and oysters, that provide constraints on paleo sea level (Falkenroth et al., 2020). Differential GPS surveys place the notch apex at $3.93 \pm 0.12 \text{ m a.s.l.}$, indicating a MIS 5 highstand of about 4 m a.s.l. The attribution to MIS 5 is supported by cosmogenic nuclide ^{10}Be exposure dating of the underlying fanglomerate but it is not reliable due to unknown precise values (Falkenroth et al., 2020). This notch is archived in WALIS as RSL_4541, with its indicative meaning tied to the intertidal zone and only minimal vertical uncertainty due to clear biological zonation.

A second indicator in the area consists of cemented beach deposits (beachrock). These deposits, composed of intertidal sands cemented during early diagenesis, preserve bedding features that fix their indicative range to the intertidal zone. OSL dating by Mauz et al. (2015) yielded an age of $80 \pm 3 \text{ ka}$, correlating the beachrock with MIS 5a. This record is archived in WALIS as RSL_4502. Sur Lagoon offers one of the clearest Last Interglacial sea-level records in the Arabian Sea region, with reliable chronological control provided by Mauz et al. (2015) and methodological refinement by Falkenroth et al. (2020).

3.3 India

3.3.1 The Saurashtra Coast of Kathiwar Peninsula

The Saurashtra coast of western India (Figs. 1 and 3) preserves Late Quaternary shorelines, with the coastal *miliolite* belt representing the most important evidence of marine transgression. These carbonate-rich deposits, locally known as *Porbandar Stone*, have been investigated by Brückner (1989) and later reassessed by Bhatt and Bhonde (2006).

At Porbandar and Tunkra (Figs. 1 and 3), fossiliferous deposits crop out at 4 m a.s.l. (RSL_4540). The lithofacies is characterized by fossiliferous miliolite, a shallow-marine limestone containing mollusks assemblages (*Tellina*, *Cerithium*, *Ostrea*, *Area*), some of which are preserved in situ position. The skeletal assemblages include benthic foraminifera, ostracods, and bryozoans, which confirms deposition in intertidal to shallow-subtidal environments. These deposits occur at the base of the coastal miliolite succession and are overlain by aeolian dune carbonates, marking the regression that followed the Last Interglacial (Brückner, 1989).

Chronological control is provided by U-Th and ESR dating of shells of *Tellina*, which yielded ESR ages of 94.9 ± 10 , 97.6 ± 10 , and $105.0 \pm 10 \text{ ka}$, consistent with attribution to MIS 5 (Brückner, 1989). U-Th dating attempts were less reliable due to diagenetic alteration of *Tellina* shells. The inland miliolite formations, sometimes rising above 30–60 m a.s.l. are dominantly aeolian in origin and not directly linked to sea level (Bhatt and Bhonde, 2006). The coastal miliolite belt preserves two distinct marine phases. The younger, marine-influenced generation (Miliolite-I) corresponds to the Last Interglacial transgression, while older




Figure 3. Map of the Indian Peninsula Last Interglacial SLIPs. Colored dots represent sea-level data points included in the WALIS database, with colors indicating elevation. Basemap assembled from datasets provided by the Global Multi-Resolution Topography (GMRT).

miliolite formations (Miliolite-II and beyond) likely predate MIS 5 (Brückner, 1989).

The coastal miliolite deposits of Saurashtra indicate that during the Last Interglacial, sea level reached 4 m a.s.l. Thus, Saurashtra provides the western Indian reference site for the Last Interglacial sea level record consisting of faunal assemblages, marine lithofacies, and chronology.

3.3.2 Cape Comorin

At the southern extremity of the Indian peninsula, Cape Comorin (Fig. 3) has been investigated in detail by Brückner (1989), followed by Banerjee (2000). A site is located 6 km west of Cape Comorin, and the preserved Last Interglacial shoreline deposit consists of a cemented beach conglomerate (RSL_4492) that outcrops at an elevation of 1.2 m above low-tide level (LTL). The lithofacies is characterized by coarse beachrock facies, composed of clast-supported conglomerates cemented by sparry calcite. The deposit contains abundant shallow-water bioclasts, including marine shell fragments and foraminiferal tests. The skeletal assemblage is dominated by intertidal foraminifera (*Ammonia*, *Elphidium*, *Cibicides*, *Globigerinoides*, *Amphistegina*, and *Epinoides*), which indicates deposition in a high-energy foreshore environment. The presence of iron-oxide coatings

on many bioclasts suggests brief episodes of subaerial exposure during or after deposition (Banerjee, 2000).

Dating of this unit was performed by using U-Th analysis of marine shells. One specimen yielded a U-Th age of $112 \text{ ka} \pm 16 / - 14$ (Brückner, 1989; Banerjee, 2000). Although mollusks are generally less reliable due to uranium exclusion and susceptibility to diagenesis, the age is stratigraphically matched with the regional Last Interglacial patterns. Radiocarbon attempts on the same facies produced anomalously young ages, attributed to secondary carbonate overprinting (Banerjee, 2000). The Cape Comorin beachrock (RSL_4492) provides a Last Interglacial index point combining lithofacies evidence of intertidal deposition with a U-Th chronology of $112 \pm 16 / - 14 \text{ ka}$.

3.3.3 Idindakarai

This site is located 35 km northeast of Cape Comorin on the Gulf of Mannar margin and preserves one of the most prominent Last Interglacial shoreline successions described along the Tamil Nadu coast (Figs. 1 and 3). The site was first documented by Brückner (1989), who identified a marine terrace elevated between 3 and 5 m above LTL. The re-investigations by Banerjee (2000) provided a more detailed stratigraphy and depositional environment, confirming its correlation with the Last Interglacial (RSL_4493).

The lower, 1.5 m thick, cross-laminated grainstone sequence is characterized by ripple bedding sealed with bioturbated contact, which is interpreted as foreshore to shoreface in origin. This is overlain by a 0.6–1.0 m thick grainstone facies containing abundant mollusk bioclasts, peloids, and borings produced by intertidal organisms such as *Balanus*. This facies indicates deposition in a shallow, high-energy intertidal environment during MIS 5e.

Chronological control is determined from U-Th analysis of marine shells. Brückner (1989) reported an age of approximately $124 \pm 14 / -8$ ka, and Banerjee (2000) confirmed the assignment of this terrace to MIS 5e. However, mollusks are problematic substrates for U-Th dating, as they generally incorporate little uranium during life and are prone to diagenetic alteration (Kaufman et al., 1971). Despite these uncertainties, the stratigraphic position, elevation (3.4 m a.s.l.; RSL_4493), and faunal evidence strongly support attribution to the Last Interglacial.

3.3.4 Manappad

The Manappad site is located about 40 km northeast of Idindakarai on the southeast Indian coast (Figs. 1 and 3) and preserves a well-defined sequence of marine terraces formed during the Late Pleistocene. Brückner (1989) described a series of stepped marine terraces in this area, with the major focus on the marine terrace level (T2) at 2–3 m above LTL. Later, using the U-Th dating method, Banerjee (2000) re-examined the same succession and suggested that the T2 terrace belongs to MIS 5e. The T2 terrace is formed by consolidated beach deposits and beachrock facies. The lithofacies consist of well-cemented, grain-supported conglomerates containing marine shell fragments and barnacle encrustations. A key indicator in situ *Balanus* sp., confirming the intertidal origin of the deposit (Banerjee, 2000). The elevation of marine terrace T2 is reported at 2.15 m above LTL (RSL_4494).

Chronological control was attempted by using U-Th dating on the marine shell *Balanus* sp. Although the analytical results were not always internally consistent due to diagenetic alteration of the shells, Banerjee (2000) correlated this terrace with MIS 5e based on its stratigraphic position, elevation, and comparison with nearby dated sites at Idindakarai and Cape Comorin. Brückner (1989) had already suggested the same correlation. The Manappad terrace, therefore, provides additional evidence for a Last Interglacial recorded at 2 m above LTL along the Tamil Nadu passive margin.

3.3.5 Rameshwaram

The island of Rameswaram, located in the Gulf of Mannar (Tamil Nadu, India, Figs. 1 and 3), provides a rare archive of Last Interglacial sea-level indicators along the passive southeastern continental margin of the Indian Peninsula. Situated at the northern end of Adam's Bridge, Rameswaram

is geomorphologically characterized by low-lying marine terraces, fossil coral colonies, and associated carbonate facies. This locality was first examined by Brückner (1989) using chronological methods that include U-Th dating and ESR. Later, this locality was further investigated by Banerjee (2000), who presented additional U-Th chronology, confirming the presence of multiple Last Interglacial reefal sea level indicators at a maximum elevation of 2.9 m above LTL. Later, Loveson and Nigam (2019) discussed Rameswaram's Last Interglacial coral-bearing sea level proxies and their chronological constraints in their review. These sites are now archived in WALIS (RSL_4489, RSL_4490, RSL_4491, RSL_4503, RSL_4504, RSL_4505, and RSL_4506).

On the northern coast at Narikulam (Rameswaram), an *Acropora* colony of 1 m² in situ (RSL_4489) is present at an elevation of 1.1 m above LTL. This coral yielded a U-Th (alpha count) age of $125.6 \pm 8.5 / 7.5$ ka, closely corresponding to global MIS 5e. However, minor detrital contamination and diagenetic alteration may affect the precision of the dating (Banerjee, 2000). The 1 m² *Acropora* colony is absent 100 m to the south, where a large coral terrace (T2), approximately 10 m² in area, is present. This coral terrace is dominated by the coral genus *Porites* and is associated with *Diploastrea*, *Cycloseris*, and *Goniopora* (RSL_4490, RSL_4491, RSL_4503, RSL_4504, RSL_4505, and RSL_4506). The elevation of this coral terrace (T2) ranges between 2.4 and 2.9 m above LTL. Several *Porites* samples reflect U-Th (Alpha count) ages ranging from 92 to 116 ka. While some specimens show a recrystallization into coarse-fibrous aragonite and localized calcite replacement, the stratigraphic coherence and clustering of ages strongly support their correlation with MIS 5.

The coral terraces dominated by *Porites* extend laterally for several kilometers, covering 10 km² in the case of the Pisasu Munai complex, and form a prominent flat surface dissected by seasonal monsoonal flooding. The repeated monsoon flooding accelerated diagenetic alteration in these corals, explaining the prevalence of secondary mineralization (Brückner, 1989; Plaziat et al., 2008). However, the presence of primary aragonite fractions in several samples confirms that the ages are reliable (Gischler et al., 2008, 2018; Henderson et al., 1993).

The Rameswaram documents a Last Interglacial high-stand between 2 and 3 m a.s.l., constrained by multiple in situ coral colonies dated between 92 and 125 ka (RSL_4489, RSL_4490, RSL_4491, RSL_4503, RSL_4504, RSL_4505, and RSL_4506), and stands as a reference locality for passive margin sea-level studies of the Last Interglacial.

3.3.6 Vettaikaraniruppu

The Vettaikaraniruppu site is situated on the southeast coast of India (Tamil Nadu, India). The Last Interglacial sea level record was first reported by Goswami et al. (2019), who investigated the beach ridge through a drill core (VKI core).

The core provided detailed subsurface stratigraphy of the Last Interglacial marine transgression from the Kaveri delta. The coring site is located at 7 m a.s.l. (RSL_4539).

The VKI core revealed a complete succession of silty sands, organic-rich clays, and carbonate-rich grainstones. The basal sandy layers displayed cross-laminated and ripple bedding, typical of foreshore to shoreface facies deposited under high-energy wave and swash processes. At a depth of 21–21.5 m b.s.l., the organic-rich sandy clays were recovered, representing low-energy conditions, likely associated with lagoonal or back-barrier settings that developed during the landward shift of environments in the marine transgression.

The upper part of the core consists of carbonate-rich grainstones with abundant marine shell debris and benthic foraminifera that include shallow-marine skeletal assemblages. These faunal assemblages confirm deposition in an intertidal to shallow subtidal setting during the peak of the highstand.

Chronological constraints are determined for the VKI core sandy clay layer at 21–21.5 m b.s.l., using OSL dating, and yield an age of 121 ± 11 ka, which correlates the sequence with the Last Interglacial. The stratigraphy of the core demonstrates a marine transgression during MIS 5e, beginning with foreshore sands, passing through lagoonal clays, and culminating in shallow-marine carbonate grainstones.

Thus, the VKI core at Vettaikaraniruppu provides one of the clearest stratigraphic archives of the MIS 5e transgression along the southeast Indian margin, combining geomorphological evidence of beach ridges with subsurface lithofacies and an OSL chronology.

3.4 Indian Ocean

3.4.1 Maldives archipelago

Rasdho Atoll, situated in the central Maldives archipelago (Fig. 1), represents one of the very few documented sites in the equatorial Indian Ocean where Last Interglacial reefal deposits have been identified and dated. The succession, described by Gischler et al. (2008) and revisited in later works (Gischler, 2008; Gischler et al., 2018), is preserved at 14.5 m b.s.l. and is archived in the WALIS database as RSL_4495, RSL_4498, RSL_4537, and RSL_4538. The deposits consist of coralline grainstones dominated by *Isopora palifera* and *Acropora gr. robusta*, with additional contributions from encrusting *Porites*, *Leptoseris*, coralline algae (*Porolithon onkodes*), and vermetids, indicating formation in a shallow back-reef environment with paleo-water depths of less than 10 m.

Chronological control is provided by U-Th dating of acroporid corals. A well-preserved specimen yielded a reliable age of 136.9 ± 2.0 ka, while two further ages of 114.3 ± 1.5 and 117.3 ± 2.7 ka are considered moderately reliable due to partial diagenetic alteration (Gischler, 2008; Gischler et

al., 2008). Later work reported two additional ages close to 151 ka, but these are regarded as unreliable owing to anomalous $^{234}\text{U}/^{238}\text{U}$ signatures (Gischler et al., 2018). Despite these limitations, the stratigraphic and clustering of ages firmly link the Rasdhoo deposits to MIS 5e. The coralline deposits which are dated with U-Th ages and yielded between 114 and 137 ka, are positioned at 14.5 m b.s.l. As archived in WALIS (RSL_4495, RSL_4498, RSL_4537, RSL_4538), these data provide an essential reference point for reconstructing subsidence-corrected relative sea-level histories in low-lying carbonate platforms.

3.4.2 Cocos (Keeling) Islands

The Cocos (Keeling) Islands, located on the Cocos Rise in the eastern Indian Ocean (Fig. 1), provide one of the few well-dated archives of Last Interglacial reef deposits from an isolated oceanic atoll setting. This locality is geomorphologically characterized by lithified reef limestones overlain by unconsolidated Holocene sands and coralline limestones beneath the modern reef islands. The site was described by Woodroffe et al. (1991) based on borehole drilling and later explained within a broader Indian Ocean context by Woodroffe (2005). These deposits are registered in WALIS under RSL_4496.

Drilling across the atoll rim revealed a sharp discontinuity at 8–11 m b.s.l., separating poorly consolidated Holocene sediments from a well-lithified reef limestone. Within this older reef, a massive *Faviid* coral from 10.5 m b.s.l. yielded U-Th ages of 118 ± 7 ka (bulk sample) and 123 ± 7 ka (secondary cement removed), confirming the formation during the Last Interglacial (Woodroffe et al., 1991). The deposits consist mainly of coralline limestones with abundant massive *Faviid* corals and other reefal taxa, consistent with a shallow-water reef crest environment.

The Cocos (Keeling) Islands record documents a Last Interglacial reefal unit now situated 8–11 m b.s.l., constrained by U-Th ages of 118–123 ka. The deposits, represented in WALIS as RSL_4496, provide key evidence of both MIS 5e reef development and sustained atoll subsidence in the eastern Indian Ocean.

4 Discussion

The standardized WALIS dataset compiled in this study provides a comprehensive synthesis of Last Interglacial relative sea-level indicators across the northern and central Indian Ocean, comprising a total of 54 SLIPs. Overall, the SLIPs in our database reveal significant departures from global eustatic estimates (Fig. 4), driven by differential vertical land motions in three main contrasting geodynamic settings: an active convergent margin (Iran and Oman), a stable passive continental margin (India), and an intraplate oceanic setting undergoing subsidence (Maldives and Cocos Keeling Islands). Across the study region, these 54 SLIPs display




Figure 4. MIS 5 RSL elevation as indicated by SLIPs in our database across four major regions of the northern Indian Ocean: southern Iran, northern Oman, peninsular India, the Maldives, and Cocos (Keeling) Islands.

a pronounced spatial elevation gradient. The uplifted active margins of Iran and Oman record RSL elevations reaching up to 60 m a.s.l., whereas the stable passive margin of India preserves SLIPs at 1–4 m a.s.l. In contrast, the tectonic subsidence of the oceanic reef platforms in the Maldives and Cocos Islands is reflected in submerged RSL indicators at depths down to 17 m b.s.l.

4.1 Uplifted active margins of Iran and Oman

In Southern Iran, notably high MIS 5 marine terraces strongly suggest the influence of regional tectonic uplift; for instance, the MIS 5e coral terraces reach up to 23 m a.s.l. in the Haleh area and up to 21 m a.s.l. in the Jazeh area. This setting is coherent with the deformation pattern of the Zagros foreland fold-and-thrust belt, where an uplift rate of 0.2 mm yr^{-1} has been estimated since the last interglacial period (Reyss et al., 1999). This is consistent with estimates derived from other localities in southern Iran (Reyss et al., 1999; Pirazzoli et al., 2004; Preusser et al., 2003; Talebian and Jackson, 2002; Regard et al., 2005). Consequently, relative sea-level positions in this part of the Persian Gulf cannot be explained by eustasy alone and must be understood within a framework of steady Quaternary uplift combined with local salt diapirism processes.

Similarly, the MIS 5 sea-level records from the Kish and Qeshm islands in southern Iran are significantly elevated up to 32 m a.s.l. These high elevations reflect a steady regional uplift rate attributed to active salt doming and neotectonic deformation within the southern Zagros foreland, with an uplift rate of 0.2 mm yr^{-1} since the last interglacial period (Pirazzoli et al., 2004). This mechanism also explains why coral

terraces representing different MIS 5 substages occur at elevations significantly above global eustatic benchmarks for the MIS 5e highstand (i.e., 2 to 9 m a.s.l.; e.g., Kopp et al., 2009; Dutton and Lambeck, 2012). On Kish Island, the lowest reef terraces (5–12 m a.s.l.) yield consistent ESR ages, providing the clearest evidence of the Last Interglacial. The higher coral-bearing terraces reflect the combined effect of MIS 5 reef accretion and subsequent tectonic uplift. Together, the Kish and Qeshm Islands provide stratigraphically coherent archives of MIS 5 sea level history in the Persian Gulf, underscoring the dominant role of regional tectonism in shaping Quaternary coastal geomorphology (Preusser et al., 2003; Pirazzoli et al., 2004; Hearty et al., 2007; Dutton and Lambeck, 2012).

Moving eastward to the Makran coast, MIS 5e terraces reach up to 60 m a.s.l., consistent with uplift rates of 0.27 mm yr^{-1} for Konarak, 0.47 mm yr^{-1} for Ramin, and 0.55 mm yr^{-1} since the LIG (Normand et al., 2019; Regard et al., 2005). These values reflect an eastward increase in forearc deformation linked to the Makran Subduction Zone, where the Arabian Plate is subducting beneath Eurasia (Byrne et al., 1988; Ellouz-Zimmermann et al., 2007). The preservation of lower MIS 5c and 5a terraces below the higher MIS 5e level reflects the interaction between glacio-eustatic sea-level highstands and steady, continuous crustal uplift.

In northern Oman, MIS 5e sea level indicators occur up to 37 m a.s.l. following a similar uplift pattern of Kish and Qeshm islands, with an uplift rate up to 0.2 mm yr^{-1} (Hoffmann et al., 2020; Decker et al., 2024). Local change near the Qualhat Fault, where terraces tilt up to 55 m over short distances, demonstrates active late Quaternary defor-

mation associated with the forebulge of the Makran subduction. Despite this uplift, the Sur Lagoon notch (3.9 m a.s.l.; RSL_4541) and 80 ± 3 ka beachrock (RSL_4502) correspond closely to the global MIS 5e and 5a highstands, respectively, confirming near-eustatic conditions locally. These records form an uplifted but stratigraphically consistent archive of Late Quaternary sea-level change, underscoring the requirement for uplift correction when comparing to global curves (Dutton et al., 2015; Rovere et al., 2016a).

4.2 Peninsular India: a tectonically stable to slightly subsiding passive margin

The Indian coastal region displays a completely different tectonic setting compared to southern Iran and the northern Oman region (Fig. 2). Along the Saurashtra coast of western India and the southeastern Tamil Nadu coast, which includes Cape Comorin Idindakarai, Manapad, and Rameswaram, MIS 5e indicators lie at 1–4 m a.s.l. (Brückner, 1989; Banerjee, 2000). These elevations, in line with GMSL estimates, suggest a tectonically stable passive margin, possibly subjected only to minor uplift/subsidence due to the tectonics of glacial isostatic adjustment response to loading and unloading of the Indian Ocean basin (Banerjee, 2000; Milne et al., 2009). Hydro-isostatic models predict that continental shelves near large oceanic basins subside by several meters in response to the relaxation of peripheral forebulges following deglaciation (Farrell and Clark, 1976; Lambeck et al., 2014). The uniform low-elevation MIS 5e indicators across southern India fit this model, implying minimal tectonic perturbation during the Last Interglacial. Stratigraphic and sedimentological evidence, like beach ridges, marine terraces and shallow-marine fauna, support a marine transgression during MIS5e in the southern Indian region (Banerjee, 2000). Thus, peninsular India provides a regional baseline against which uplifted or subsiding settings in the Indian Ocean can be calibrated. In addition, a substantial geographical gap in LIG sea-level records still exists along much of the eastern and western coasts of India, primarily due to the limited availability of previously published data. A similar data gap exists along the approximately 1050 km-long coastline of Pakistan, highlighting the need for further investigation and the development of well-constrained LIG sea-level indicators across the northern Indian Ocean region.

4.3 Subsiding oceanic platforms: Maldives and Cocos (Keeling)

The Maldives Archipelago and Cocos (Keeling) Islands define the subsiding LIG shoreline record of Indian Ocean. Last Interglacial coral frameworks from the Maldives occur up to 17 m b.s.l., suggesting subsidence rates of $0.09\text{--}0.16\text{ mm yr}^{-1}$ (Gischler et al., 2018). These rates exceed simple thermal contraction values for 60–80 Ma oceanic lithosphere (Parsons and Sclater, 1977), suggesting addi-

tional tectonic or flexural loading components (Woodroffe, 2005). Submergence of MIS 5e reefs also reflects limited accommodation space and continuous platform downwarping (Purdy and Bertram, 1993; Perry et al., 2008). At Cocos (Keeling), submerged coral heads at 10 m b.s.l. yield ages of 123 ka, consistent with subsidence rates of $0.1\text{--}0.12\text{ mm yr}^{-1}$ (Woodroffe et al., 1991; Woodroffe, 2005). These rates, higher than expected for pure lithospheric cooling, may record lithospheric flexure associated with the Wharton Ridge or long-term dynamic topography of the Indian Ocean basin (Müller et al., 2008; Hoggard et al., 2016). Hence, both the Maldives and Cocos records confirm that the apparent depth of MIS 5e reefs is dominated by subsidence.

5 Data availability

The South Asia and the Indian Ocean database is available at <https://doi.org/10.5281/zenodo.18512769> (Ali et al., 2026). The description of the database fields can be found at <https://doi.org/10.5281/zenodo.3961544> (Rovere et al., 2020).

6 Conclusions

This review presents the first standardized Last Interglacial relative sea-level dataset spanning South Asia and the Indian Ocean region, extending from Iran and Oman to India, the Maldives, and the Cocos (Keeling) Islands. By integrating 54 sea-level index points from 17 peer-reviewed studies into the WALIS framework, we reassessed stratigraphic context, depositional environments, elevation accuracy, and chronological control using U-Th, ESR, luminescence, and cosmogenic dating. The resulting dataset reveals strong regional contrasts driven primarily by different tectonic settings. Active margins in southern Iran and northern Oman preserve MIS 5e, 5c, and 5a marine terraces, with elevations up to 60 m a.s.l., reflecting uplift associated with the Zagros fold-and-thrust belt, salt tectonics, and the Makran subduction system. On the other hand, the passive margin of western and southern India shows MIS 5e sea level indicators at 1–4 m a.s.l., consistent with tectonic stability, while the MIS 5e reefs in the Maldives and Cocos (Keeling) Islands lie 10–17 m b.s.l., showing a late Quaternary subsidence of oceanic platforms. Furthermore, the identification of MIS 5c and MIS 5a RSL indicators in uplifted settings, such as Oman and the Makran coast, demonstrates that South Asia and the northern Indian Ocean preserve key evidence of the youngest MIS 5 sea-level oscillations.

Overall, this study provides standardized information for reconstructing Last Interglacial sea level changes in a region that has been historically underrepresented in global databases. However, significant data gaps remain along the Pakistan margin and some areas of India, where the lack of detailed and consistently documented studies currently pre-

vents the inclusion of reliable, standardized sea-level data-points, highlighting the need for targeted future investigations. The standardized dataset provides an essential reference for future modeling of glacio-hydro-isostatic processes, vertical land motion, and Indian Ocean tectonics studies. Importantly, the reviewed sea level records offer valuable analogues for understanding how coastlines with varied tectonic and geomorphic settings may respond to future sea-level rise.

Author contributions. MA and CC: Investigation, Conceptualization, writing the original draft, visualization. AR: Conceptualization, writing the original draft, Supervision, and visualization.

Competing interests. At least one of the (co-)authors is a member of the editorial board of *Earth System Science Data*. The peer-review process was guided by an independent editor, and the authors also have no other competing interests to declare.

Disclaimer. This paper reflects only the author's view and that the EU is not responsible for any use that may be made of the information it contains.

Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. The authors bear the ultimate responsibility for providing appropriate place names. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Special issue statement. This article is part of the special issue “WALIS – the World Atlas of Last Interglacial Shorelines”. It is not associated with a conference.

Acknowledgements. This study is an outcome of the WARMCOAST project, funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 802414). The authors are particularly grateful to the Department of Environmental Sciences, Informatics and Statistics (Ca' Foscari University of Venice) for technical support during the project. The authors are also thankful to the WALIS database contributors.

Financial support. This paper is a result of the WARMCOAST project, funded by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 802414).

Review statement. This paper was edited by Xingchen (Tony) Wang and reviewed by Kim Cohen and Woo Hun Ryang.

References

- Ali, M., Cerrone, C., and Rovere, A.: A review of last interglacial sea-level proxies in South Asia and the Indian Ocean, Zenodo [data set], <https://doi.org/10.5281/zenodo.18512769>, 2026.
- Antonioli, F., Lo Presti, V., Rovere, A., Ferranti, L., Anzidei, M., Furlani, S., Mastronuzzi, G., Orru, P. E., Scicchitano, G., Sannino, G., Spampinato, C. R., Pagliarulo, R., Deiana, G., de Sabata, E., Sansò, P., Vacchi, M., and Vecchio, A.: Tidal notches in Mediterranean Sea: a comprehensive analysis, *Quaternary Sci. Rev.*, 119, 66–84, <https://doi.org/10.1016/j.quascirev.2015.03.016>, 2015.
- Banerjee, P.: Holocene and Late Pleistocene relative sea level fluctuations along the east coast of India, *Mar. Geol.*, 167, 243–260, [https://doi.org/10.1016/S0025-3227\(00\)00028-1](https://doi.org/10.1016/S0025-3227(00)00028-1), 2000.
- Bhatt, N. and Bhonde, U.: Geomorphic expression of late Quaternary sea level changes along the southern Saurashtra coast, western India, *J. Earth Syst. Sci.*, 115, 395–402, <https://doi.org/10.1007/BF02702868>, 2006.
- Bird, P.: An updated digital model of plate boundaries, G-cubed, 4, <https://doi.org/10.1029/2001GC000252>, 2003.
- Boyden, P., Weil-Accardo, J., Deschamps, P., Oppo, D., and Rovere, A.: Last interglacial sea-level proxies in East Africa and the Western Indian Ocean, *Earth Syst. Sci. Data*, 13, 1633–1651, <https://doi.org/10.5194/essd-13-1633-2021>, 2021.
- Brückner, H.: Late quaternary shorelines in India, in: Late Quaternary Sea-Level Correlation and Applications: Walter S. Newman Memorial Volume, Springer Netherlands, Dordrecht, 169–194, https://doi.org/10.1007/978-94-009-0873-4_9, 1989.
- Byrne, D. E., Davis, D. M., and Sykes, L. R.: Loci and maximum size of thrust earthquakes and the mechanics of the shallow region of subduction zones, *Tectonics*, 7, 833–857, <https://doi.org/10.1029/tc007i004p00833>, 1988.
- Decker, V., Falkenroth, M., and Hoffmann, G.: Sedimentological evidence of Late Pleistocene shorelines in Oman, *Mar. Geol.*, 477, 107407, <https://doi.org/10.1016/j.margeo.2024.107407>, 2024.
- Dutton, A. and Lambeck, K.: Ice volume and sea level during the Last Interglacial, *Science*, 337, 216–219, <https://doi.org/10.1126/science.1205749>, 2012.
- Dutton, A., Webster, J. M., Zwart, D., Lambeck, K., and Wohlfarth, B.: Tropical tales of polar ice: evidence of Last Interglacial polar ice sheet retreat recorded by fossil reefs of the granitic Seychelles islands, *Quaternary Sci. Rev.*, 107, 182–196, <https://doi.org/10.1016/j.quascirev.2014.10.025>, 2015.
- Ellouz-Zimmermann, N., Deville, E., Müller, C., Lallemand, S., Subhani, A. B., and Tabreez, A. R.: Impact of sedimentation on convergent margin tectonics: Example of the Makran accretionary prism (Pakistan), Thrust belts and foreland basins, *Springer*, 327–350, https://doi.org/10.1007/978-3-540-69426-7_17, 2007.
- Falkenroth, M., Schneider, B., and Hoffmann, G.: Beachrock as sea-level indicator – A case study at the coastline of Oman (Indian Ocean), *Quaternary Sci. Rev.*, 206, 81–98, <https://doi.org/10.1016/j.quascirev.2019.01.003>, 2019.
- Falkenroth, M., Adolphs, S., Cahnbley, M., Bagci, H., Kázmér, M., Mechernich, S., and Hoffmann, G.: Biological indicators reveal small-scale sea-level variability during MIS 5e (Sur, Sultanate of Oman), *Open Quatern.*, 6, <https://doi.org/10.5334/oq.72>, 2020.

- Farrell, W. E. and Clark, J. A.: On postglacial sea level, *Geophys. J. Roy. Astron. Soc.*, 46, 647–667, <https://doi.org/10.1111/j.1365-246X.1976.tb01252.x>, 1976.
- Garzón, S. and Rovere, A.: WALIS dashboard: An online tool to explore a global paleo sea-level database, *Open Res. Europe*, 3, 114, <https://doi.org/10.12688/openreseurope.16183.2>, 2024.
- Gischler, E.: Corrigendum to “Late Quaternary reef growth and sea level in the Maldives (Indian Ocean)” by E. Gischler, J. H. Hudson and A. Pisera. [*Marine Geology* 250 (2008) 104–113], *Mar. Geol.*, 255, 102, <https://doi.org/10.1016/j.margeo.2008.07.002>, 2008.
- Gischler, E., Hudson, J. H., and Pisera, A.: Late Quaternary reef growth and sea level in the Maldives (Indian Ocean), *Mar. Geol.*, 250, 104–113, <https://doi.org/10.1016/j.margeo.2008.01.004>, 2008.
- Gischler, E., Humblet, M., Braga, J. C., and Eisenhauer, A.: Last interglacial reef facies and late Quaternary subsidence in the Maldives, Indian Ocean, *Mar. Geol.*, 406, 34–41, <https://doi.org/10.1016/j.margeo.2018.09.001>, 2018.
- Goswami, K., Krishnan, S., Kumerasan, A., Sadasivam, S. K., Kumar, P., and Jaiswal, M. K.: Luminescence chronology of fluvial and marine records from subsurface core in Kaveri delta, Tamil Nadu: Implications to sea level fluctuations, *Geochronometria*, 46, 125–137, <https://doi.org/10.1515/geochr-2015-0112>, 2019.
- Hearty, P. J., Hollin, J. T., Neumann, A. C., O’Leary, M. J., and McCulloch, M.: Global sea-level fluctuations during the Last Interglaciation (MIS 5e), *Quaternary Sci. Rev.*, 26, 2090–2112, <https://doi.org/10.1016/j.quascirev.2007.06.019>, 2007.
- Henderson, G. M., Cohen, A. S., and O’Nions, R. K.: $^{234}\text{U}/^{238}\text{U}$ ratios and ^{230}Th ages for Hateruma Atoll corals: implications for coral diagenesis and seawater $^{234}\text{U}/^{238}\text{U}$ ratios, *Earth Planet. Sc. Lett.*, 115, 65–73, [https://doi.org/10.1016/0012-821X\(93\)90213-S](https://doi.org/10.1016/0012-821X(93)90213-S), 1993.
- Hoffmann, G., Reicherter, K., Wiater, T., Grützner, C., and Rausch, T.: Block and boulder accumulations along the coastline between Fins and Sur (Sultanate of Oman): tsunamigenic remains?, *Nat. Hazards*, 65, 851–873, <https://doi.org/10.1007/s11069-012-0399-7>, 2013a.
- Hoffmann, G., Rupprechter, M., and Mayrhofer, C.: Review of the long-term coastal evolution of North Oman-subsidence versus uplift, *Z. Deutsch. Gesell. Geowiss.*, 164, 237–252, <https://doi.org/10.1127/1860-1804/2013/0002>, 2013b.
- Hoffmann, G., Schneider, B., Mechernich, S., Falkenroth, M., Dunai, T., and Preusser, F.: Quaternary uplift along a passive continental margin (Oman, Indian Ocean), *Geomorphology*, 350, 106870, <https://doi.org/10.1016/j.geomorph.2019.106870>, 2020.
- Hoggard, M. J., White, N., and Al-Attar, D.: Global dynamic topography observations reveal limited influence of large-scale mantle flow, *Nat. Geosci.*, 9, 456–463, <https://doi.org/10.1038/ngeo2709>, 2016.
- Kaufman, A., Broecker, W. S., Ku, T. L., and Thurber, D. L.: The status of U-series methods of mollusk dating, *Geochim. Cosmochim. Ac.*, 35, 1155–1183, [https://doi.org/10.1016/0016-7037\(71\)90031-7](https://doi.org/10.1016/0016-7037(71)90031-7), 1971.
- Khan, N. S., Horton, B. P., Engelhart, S., Rovere, A., Vacchi, M., Ashe, E. L., Törnqvist, T. E., Dutton, A., Hijma, M. P., and Shennan, I.: Inception of a global atlas of sea levels since the Last Glacial Maximum, *Quaternary Sci. Rev.*, 220, 359–371, <https://doi.org/10.1016/j.quascirev.2019.07.016>, 2019.
- Kopp, R. E., Simons, F. J., Mitrovica, J. X., Maloof, A. C., and Oppenheimer, M.: Probabilistic assessment of sea level during the last interglacial stage, *Nature*, 462, 863–867, <https://doi.org/10.1038/nature08686>, 2009.
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambridge, M.: Sea level and global ice volumes from the Last Glacial Maximum to the Holocene, *P. Natl. Acad. Sci. USA*, 111, 15296–15303, <https://doi.org/10.1073/pnas.1411762111>, 2014.
- Lorscheid, T. and Rovere, A.: The indicative meaning calculator quantification of paleo sea-level relationships by using global wave and tide datasets, *Open Geospat. Data Softw. Stand.*, 4, <https://doi.org/10.1186/s40965-019-0069-8>, 2019.
- Loveson, V. J. and Nigam, R.: Reconstruction of Late Pleistocene and Holocene sea level curve for the east coast of India, *J. Geol. Soc. India*, 93, 507–514, <https://doi.org/10.1007/s12594-019-1211-z>, 2019.
- Mauz, B., Vacchi, M., Green, A., Hoffmann, G., and Cooper, A.: Beachrock: a tool for reconstructing relative sea level in the far-field, *Mar. Geol.*, 362, 1–16, <https://doi.org/10.1016/j.margeo.2015.01.009>, 2015.
- Milne, G. A., Gehrels, W. R., Hughes, C. W., and Tamisiea, M. E.: Identifying the causes of sea-level change, *Nat. Geosci.*, 2, 471–478, <https://doi.org/10.1038/ngeo544>, 2009.
- Müller, R. D., Sdrolias, M., Gaina, C., and Roest, W. R.: Age, spreading rates and spreading asymmetry of the world’s ocean crust, *Geochem. Geophys.*, 9, <https://doi.org/10.1029/2007GC001743>, 2008.
- Normand, R., Simpson, G., Herman, F., Biswas, R. H., Bahroudi, A., and Schneider, B.: Dating and morpho-stratigraphy of uplifted marine terraces in the Makran subduction zone (Iran), *Earth Surf. Dynam.*, 7, 321–344, <https://doi.org/10.5194/esurf-7-321-2019>, 2019.
- Otvos, E. G.: Beach ridges – definitions and significance, *Geomorphology*, 32, 83–108, [https://doi.org/10.1016/S0169-555X\(99\)00075-6](https://doi.org/10.1016/S0169-555X(99)00075-6), 2000.
- Parsons, B. and Sclater, J. G.: An analysis of the variation of ocean floor bathymetry and heat flow with age, *J. Geophys. Res.*, 82, 803–827, <https://doi.org/10.1029/JB082i005p00803>, 1977.
- Pedoja, K., Husson, L., Johnson, M. E., Melnick, D., Witt, C., Pochat, S., Nexer, M., Delcaillau, B., Pinegina, T., Poprawski, Y., Authemayou, C., Elliot, M., Regard, V., and Garestier, F.: Coastal staircase sequences reflecting sea-level oscillations and tectonic uplift during the Quaternary and Neogene, *Earth Sci. Rev.*, 132, 13–38, <https://doi.org/10.1016/j.earscirev.2014.01.007>, 2014.
- Perry, C. T., Spencer, T., and Kench, P. S.: Carbonate budgets and reef production states: a geomorphic perspective on the ecological phase-shift concept, *Coral Reefs*, 27, 853–866, <https://doi.org/10.1007/s00338-008-0418-z>, 2008.
- Pirazzoli, P. A.: Marine Terraces, in: *Encyclopedia of Coastal Science*, Springer, 632–633, https://doi.org/10.1007/1-4020-3880-1_209, 2005.
- Pirazzoli, P. A., Reyss, J.-L., Fontugne, M., Haghypour, A., Hilgers, A., Kasper, H. U., Nazari, H., Preusser, F., and Radtke, U.: Quaternary coral-reef terraces from Kish and Qeshm Islands, Persian Gulf: new radiometric ages and tectonic implications, *Quatern. Int.*, 120, 15–27, <https://doi.org/10.1016/j.quaint.2004.01.003>, 2004.
- Plaziat, J. C., Reyss, J. L., Choukri, A., and Cazala, C.: Diagenetic rejuvenation of raised coral reefs and precision of dating. The

- contribution of the Red Sea reefs to the question of reliability of the Uranium-series datings of middle to late Pleistocene key reef-terraces of the world, *Carnets de Géologie/Notebooks on Geology*, A04, 1–35, <https://doi.org/10.4267/2042/16743>, 2008.
- Preusser, F., Radtke, U., Fontugne, M., Haghipour, A., Hilgers, A., Kasper, H. U., Nazari, H., and Pirazzoli, P. A.: ESR dating of raised coral reefs from Kish Island, Persian Gulf, *Quaternary Sci. Rev.*, 22, 1317–1322, [https://doi.org/10.1016/s0277-3791\(03\)00038-6](https://doi.org/10.1016/s0277-3791(03)00038-6), 2003.
- Purdy, E. G. and Bertram, G. T.: Carbonate concepts from the Maldives, Indian ocean, *AAPG Studies in Geology*, 1–56, <https://doi.org/10.1306/St34568>, 1993.
- Regard, V., Bellier, O., Thomas, J.-C., Bourlès, D., Bonnet, S., Abbassi, M. R., Braucher, R., Mercier, J., Shabanian, E., Soleymani, S., and Feghhi, K.: Cumulative right-lateral fault slip rate across the Zagros–Makran transfer zone: role of the Minab–Zendan fault system in accommodating Arabia–Eurasia convergence in southeast Iran, *Geophys. J. Int.*, 162, 177–203, <https://doi.org/10.1111/j.1365-246X.2005.02558.x>, 2005.
- Reyss, J.-L., Pirazzoli, P. A., Haghipour, A., Hatté, C., and Fontugne, M.: Quaternary marine terraces and tectonic uplift rates on the south coast of Iran, in: *Coastal Tectonics, Special Publications*, 146, edited by: Stewart, I. S. and Vita-Finzi, C., Geological Society, London, 225–237, <https://doi.org/10.1144/GSL.SP.1999.146.01.13>, 1999.
- Rovere, A., Stocchi, P., and Vacchi, M.: Eustatic and relative sea level changes, *Curr. Clim. Change Rep.*, 2, 221–231, <https://doi.org/10.1007/s40641-016-0045-7>, 2016a.
- Rovere, A., Raymo, M. E., Vacchi, M., Lorscheid, T., Stocchi, P., Gomez-Pujol, L., Harris, D. L., Casella, E., O’Leary, M. J., and Hearty, P. J.: The analysis of Last Interglacial (MIS 5e) relative sea-level indicators: Reconstructing sea-level in a warmer world, *Earth Sci. Rev.*, 159, 404–427, <https://doi.org/10.1016/j.earscirev.2016.06.006>, 2016b.
- Rovere, A., Ryan, D., Murray-Wallace, C., Simms, A., Vacchi, M., Dutton, A., Lorscheid, T., Chutcharavan, P., Brill, D., Bartz, M., Jankowski, N., Mueller, D., Cohen, K., and Gowan, E.: Documentation of the World Atlas of Last Interglacial Shorelines (WALIS) (Version v1.0), Zenodo [data set], <https://doi.org/10.5281/zenodo.3961544>, 2020.
- Rovere, A., Ryan, D. D., Vacchi, M., Dutton, A., Simms, A. R., and Murray-Wallace, C. V.: The World Atlas of Last Interglacial Shorelines (version 1.0), *Earth Syst. Sci. Data*, 15, 1–23, <https://doi.org/10.5194/essd-15-1-2023>, 2023.
- Ryan, W. B. F., Carbotte, S. M., Coplan, J. O., O’Hara, S., Melkonian, A., Arko, R., Weissel, R. A., Ferrini, V., Goodwillie, A., Nitsche, F., Bonczkowski, J., and Zemsky, R.: Global Multi-Resolution Topography synthesis, *Geochem. Geophys. Geosy.*, 10, <https://doi.org/10.1029/2008GC002332>, 2009.
- Shennan, I.: Interpretation of Flandrian sea-level data from the Fenland, England, *P. Geol. Assoc.*, 93, 53–63, [https://doi.org/10.1016/S0016-7878\(82\)80032-1](https://doi.org/10.1016/S0016-7878(82)80032-1), 1982.
- Shennan, I.: Flandrian sea-level changes in the Fenland. I: The geographical setting and evidence of relative sea-level changes, *J. Quaternary Sci.*, 1, 119–153, <https://doi.org/10.1002/jqs.3390010204>, 1986.
- Shennan, I.: Holocene crustal movements and sea-level changes in Great Britain, *J. Quaternary Sci.*, 4, 77–89, <https://doi.org/10.1002/jqs.3390040109>, 1989.
- Shennan, I.: Handbook of sea-level research, in: *Handbook of Sea-Level Research*, Wiley, 3–25, <https://doi.org/10.1002/9781118452547.ch2>, 2015.
- Shennan, I., Tooley, M. J., Davis, M. J., and Haggart, B. A.: Analysis and interpretation of Holocene sea-level data, *Nature*, 302, 404–406, <https://doi.org/10.1038/302404a0>, 1983.
- Talebian, M. and Jackson, J.: Offset on the Main Recent Fault of NW Iran and implications for the late Cenozoic tectonics of the Arabia–Eurasia collision zone, *Geophys. J. Int.*, 150, 422–439, <https://doi.org/10.1046/j.1365-246X.2002.01711.x>, 2002.
- Thompson, S. B. and Creveling, J. R.: A global database of marine isotope substage 5a and 5c marine terraces and paleoshoreline indicators, *Earth Syst. Sci. Data*, 13, 3467–3490, <https://doi.org/10.5194/essd-13-3467-2021>, 2021.
- Woodroffe, C. D.: Late Quaternary sea-level highstands in the central and eastern Indian Ocean: A review, *Global Planet. Change*, 49, 121–138, <https://doi.org/10.1016/j.gloplacha.2005.06.002>, 2005.
- Woodroffe, C. D., Veeh, H. H., Falkland, A. C., McLean, R. F., and Wallensky, E.: Last interglacial reef and subsidence of the Cocos (Keeling) Islands, Indian Ocean, *Mar. Geol.*, 96, 137–143, [https://doi.org/10.1016/0025-3227\(91\)90210-U](https://doi.org/10.1016/0025-3227(91)90210-U), 1991.