

Earth System Science Data Discussions is the access reviewed
discussion forum of *Earth System Science Data*

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Bjerknes Centre for Climate Research, UNIFOB AS, Allegaten 55, 5007 Bergen, Norway

Department of Chemistry, University of Gothenburg, 41296, Göteborg, Sweden

Received: 1 July 2009 – Accepted: 9 July 2009 – Published: 5 August 2009

Correspondence to: A. Olsen (are.olsen@gfi.uib.no)

Published by Copernicus Publications.

ESSDD

2, 181–203, 2009

**Nordic Seas total
dissolved inorganic
carbon data in
CARINA**

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Water column data of carbon and carbon relevant hydrographic and hydrochemical parameters from 188 previously non-publicly available cruises in the Arctic, Atlantic, and Southern Ocean have been retrieved and merged into a new database: CARINA (CARbon IN the Atlantic). The data have been subject to rigorous quality control (QC) in order to ensure highest possible quality and consistency. The data for most of the parameters included were examined in order to quantify systematic biases in the reported values, i.e. secondary quality control. Significant biases have been corrected for in the data products, i.e. the three merged files with measured, calculated and interpolated values for each of the three CARINA regions; the Arctic Mediterranean Seas (AMS), the Atlantic (ATL) and the Southern Ocean (SO). With the adjustments the CARINA database is consistent both internally as well as with GLODAP (Key et al., 2004) and is suitable for accurate assessments of, for example, oceanic carbon inventories and uptake rates and for model validation. The Arctic Mediterranean Seas includes the Arctic Ocean and the Nordic Seas, and the quality control was carried out separately in these two areas. This contribution presents an account of the quality control of the total dissolved inorganic carbon (TCO_2) data from the Nordic Seas in CARINA. Out of the 35 cruises from the Nordic Seas included in CARINA, 25 had TCO_2 data. The data from 7 of these were found to be of low quality and should not be used, thus the final CARINA data product contains TCO_2 data from 18 cruises from the Nordic Seas. These data appear consistent to at least $4 \mu\text{mol kg}^{-1}$.

ESSDD

2, 181–203, 2009

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Data coverage and parameter measured

Repository-Reference: doi: will be provided by CDIAC upon submission of data

Available at: http://cdiac.ornl.gov/oceans/CARINA/Carina_inv.html

Coverage: 59.60° N–82.35° N, 35.23° W–20° E

5 Location Name: Nordic Seas

Date/Time Start: 1991-07-15

Date/Time End: 2003-10-13

Date/Time End: 2003-10-13

Data Product Parameter Name	Data Product Flag name	Exchange File Parameter Name	Exchange File Flag Name	Units
station		STANBR		
day		DATE		
month		DATE		
year		DATE		
latitude		LATITUDE		decimal degrees
longitude		LONGITUDE		decimal degrees
cruiseno				
depth				meters
temperature		CTDTMP		°C
salinity	sf	SALNTY	SALNTY_FLAG_W	
ctdsal	ctdsf	CTDSAL	CTDSAL_FLAG_W	
pressure		CTDPRS		decibars
tco2	tco2f	TCARBN	TCARBN_FLAG_W	$\mu\text{mol kg}^{-1}$

10 For a complete list of all parameters available in CARINA see Key et al. (2009). Note the different names for the parameters in the Exchange files (the individual cruise files) and the merged data product.

ESSDD

2, 181–203, 2009

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

[Title Page](#)

[Abstract](#)

[Instruments](#)

[Data Provenance & Structure](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

1 Introduction

CARINA (CARbon In the Atlantic) is a database of carbon and carbon-relevant data from hydrographic cruises in the Arctic, Atlantic and Southern Oceans. The project started as an essentially informal, unfunded project in Delmenhorst, Germany, in 1999 during the workshop on “CO₂ in the North Atlantic”, with the main goal to create a uniformly formatted database of carbon relevant variables in the ocean to be used for accurate assessments of oceanic carbon inventories and uptake rates. The collection of data and the quality control (QC) of the data have been a main focus of the CARINA project. Both primary and secondary QC of the data has been performed.

The CARINA database consists of two parts: the first part is the set of the individual cruise files where all the data reported by the measurement teams are stored. Quality flags are accompanying the data, in many cases those are the flags originally reported, in others they were assigned by R. M. Key, Princeton University. These files are in WHP (WOCE Hydrographic Program) exchange format where the first lines consist of the condensed metadata. There are essentially no calculated or interpolated values in the individual cruise files, with the exceptions of pressure calculated from depth and some bottle salinities that were taken from ctdsal. No adjustments have been applied to any of these values with the exception that all pH measurements were converted to the seawater pH scale at 25°C.

The second part of CARINA consists of three merged, quality controlled and adjusted data files; one each for the Atlantic Ocean, Arctic Mediterranean Seas and Southern Ocean regions. These files contain all the CARINA data and also include: 1) interpolated values for nutrients, oxygen and salinity if those data were missing and the interpolation could be made according to certain criteria, as described in Key et al. (2009); 2) calculated carbon parameters; e.g. if total dissolved inorganic carbon (TCO₂) and total alkalinity (TA) were measured, pH was calculated; and 3) instances where bottle salinity was missing or bad and were replaced with CTD salinity. Calculated or interpolated values have been given the quality flag “0”. In many cases there are additional

ESSDD

2, 181–203, 2009

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

parameters in the individual cruise files, which have not been included in the secondary QC, such as $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and SF_6 . Most of these are included in the merged data files as well.

This report provides an overview of the TCO_2 data in the Nordic Seas data in CARINA and describes the secondary QC of these data. The data are part of the Arctic Mediterranean Seas subset of CARINA (CARINA-AMS). This subset includes data from the Nordic Seas and the Arctic Ocean. Because of the very different data coverage in these two areas, and because they are more or less separate, they were analysed separately. The TCO_2 data from the Arctic Ocean are described by Jütterström et al. (2009), and the other Nordic Seas data are described in Jeansson et al. (2009), Falck and Olsen (2009), Olafsson and Olsen (2009), Olsen et al. (2009a), and Olsen (2009b). A more comprehensive description of the complete CARINA database can be found in Key et al. (2009), as well as in the other, more specialised, papers in this special issue.

2 Data Provenance and Structure

The Nordic Seas was loosely defined as the region closed by the Fram Strait to the North, Greenland to the west, the Greenland-Scotland Ridge to the South, and Norway, the Barents Sea Opening, and Spitsbergen to the east. Out of the 188 CARINA cruises, 62 are included in the CARINA-AMS, and 35 of these considered as Nordic Seas cruises. Five of these are in common with the Atlantic Ocean subset of CARINA (Tanhua et al., 2009a), in order to ensure consistency between the regions. Out of the 35 cruises included in the Nordic Seas CARINA, 25 had TCO_2 data. Except for the Iceland Sea time series, the Nordic Seas had only been the subject to sporadic sampling for TCO_2 until 1993. That year was the start of a period of quite intense chemical sampling in this region through the ESOP and CARDEEP projects (e.g. Miller et al., 1999). These projects were followed by ESOP2 in 1996, which continued until 1999 (e.g. Chierici et al., 1999), later, between 2001 and 2003, TCO_2 sampling was

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

carried out as part of the TRACTOR project (e.g. Olsen et al., 2006). The major part of the TCO₂ data from the Nordic Seas in CARINA was obtained through these projects. In addition the collection includes data from the aforementioned repeated visits to the Iceland Sea (approx. 68° N, 13° E) carried out by the Marine Research Institute, Reykjavik, data from the two Nordic WOCE expeditions of 1992 and 1993 and data from the Northeast Water Polynya expeditions of 1992 and 1993 (Yager et al., 1995). The TCO₂ data from the Nordic Seas cruise of 1982 onboard CSS Hudson (Chen et al., 1990) and the 1992 Meteor cruise were also subject to secondary QC, but were found to be of too poor quality to be included in the data product, as will be dealt with later.

Table 1 provides an overview of the data that were included in the secondary quality controls described in this paper, and station positions are shown in Fig. 1. Most of the data were obtained using a coulometer, connected either to a SOMMA (single-operator multiparameter metabolic analyzer, Johnson et al., 1993) or similar gas extraction units, in accordance with the standard operating protocols (Dickson et al., 2007). However, data from two of the earlier cruises were obtained using potentiometric techniques. Certified Reference Material (CRM) was used on almost all cruises.

3 Quality control

The secondary QC included an assessment of the precision of the data from each individual cruise as well as a consistency analysis. The consistency analysis was carried out using the crossover and inversion approach (Gouretski and Jancke, 2001; Johnson et al., 2001; Tanhua et al., 2009b), customised as described by Olsen et al. (2009a). Another possible approach would be multiparameter linear regressions (MLRs), similar to Wanninkhof et al. (2003), however, at the time of our analyses the consistency of the nutrient data had not been evaluated and since any inconsistencies would propagate into the MLRs, this method was not used.

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.1 Individual cruise assessments

The metadata contained little or no information on the precision of the TO₂ data obtained at the different cruises, and replicate samples, which would enable us to determine this, were virtually non-existent. However, distribution of chemical properties in the deep Nordic Seas is fairly homogenous (Blindheim and Østerhus, 2005, supplementary material of this paper, <http://www.earth-syst-sci-data-discuss.net/2/181/2009/essdd-2-181-2009-supplement.pdf>). With this in mind the standard deviation of samples from deeper than 1900 dbar between 65° and 80° N and 10° W to 10° E on each cruise was estimated in order to obtain an impression of the data spread on each cruise. These numbers are shown in Table 2. Obviously they must be interpreted with two effects in mind: (1) natural spatial variability and (2) precision of measurements. As regards the first effect, consider the 316N20020530 cruise, which has the largest areal coverage of all cruises with data from almost the full region included in this exercise (cnf. suppl. Fig. 1, <http://www.earth-syst-sci-data-discuss.net/2/181/2009/essdd-2-181-2009-supplement.pdf>). The standard deviation of the deep TCO₂ data obtained on this cruise is 3.10 μmol kg⁻¹. Since all of the other cruises had smaller areal coverage their standard deviation should be less. This is not the case, most cruises show larger spread, indicating poorer precision. In particular 58JH19930730 with a standard deviation of 25.5 μmol kg⁻¹ and 58JH19951108 with 10.6 μmol kg⁻¹. The samples of 58JH19930730 were collected in 50 ml serum bottles, poisoned with HgCl₂ and analysed ashore (Miller et al., 1999). This is not in accordance with standard operating procedures (Dickson et al., 2007), and considering the large spread these data appear to be of poor quality and I recommend that they are flagged questionable. Regarding the 58JH19951108, Miller et al. (1999) do note that that there were “instrumental difficulties” at this cruise and given this information and the high standard deviation of the deep samples, it is recommended that these data are flagged questionable.

ESSDD

2, 181–203, 2009

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Other than these cruises, the TCO₂ data of 18HU19820228 show high standard deviation. However, this is the oldest cruise included in the synthesis and may have special significance as a reference point when accounting for changes due to anthropogenic carbon input. These data were therefore included in the consistency analyses for further evaluation.

3.2 Consistency analyses

The consistency of the Nordic Seas TCO₂ data was evaluated through a crossover and inversion analysis (Gouretski and Jancke, 2001; Johnson et al., 2001), carried out using the cnaX routines described by Tanhua et al. (2009b). This approach compares the deep data (>1900 m for the Nordic Seas (Olsen et al., 2009)), from two cruises that have stations in vicinity and determines the offset between the data within the crossover of the two cruises, and its standard deviation. Thus each cruise gets a set of offsets compared to the other cruises. This set of offsets are then inverted using the Weighted Least Squares (WLSQ) approach of Johnson et al. (2001) to find the corrections which maximizes the consistency of the data from the different cruises.

For the Nordic Seas data two crossover and inversion analyses were carried out: (1) using only data in CARINA and (2) including the Nordic Seas data from the Transient Tracers in the Ocean-North Atlantic Study (TTO-NAS). The latter analysis was carried out because all evidence suggest that CO₂ concentrations in the deep Nordic Seas have increased during the last two decades following the anthropogenic CO₂ invasion (Olsen et al., 2006; Skjelvan et al., 2008) and the consistency of the data can not be evaluated without regard for this increase. The TTO-NAS data that were used had been corrected according to Tanhua and Wallace (2005). The data from two Northeast Water Polynya cruises (32L919920715 and 32I919930718) were not included in the analyses because there were no data from deeper than 1900 dbar and the special sampling area. The TCO₂ data from the Iceland Sea time series was not included either, because the bottom depth at the stations is less than 1900 m. The Iceland Sea time series data was, nevertheless, compared with data from other cruises, this is

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

described in Sect. 4.7.

The corrections determined by the WLSQ inversion of the crossover offsets are shown in Fig. 2, the crossover results in themselves can be accessed at the CARINA website (http://cdiac.ornl.gov/oceans/CARINA/Carina_inv.html). The inclusion of the TTO-NAS data in the analysis had two effects on the results: (1) A correction is suggested for the TTO-NAS data and (2) the suggested correction for 58JH19920712 is reduced. Otherwise the suggested corrections for the different cruises were virtually the same whether TTO-NAS was included or not (Fig. 2).

Note that the cruises have been sorted with time in Fig. 2, and the inversion suggests that the deep TTO-NAS TCO_2 data are nearly $4 \mu\text{mol kg}^{-1}$ lower than those measured at the 316N20020530 and 58GS20030922 cruises (i.e. they should receive a positive correction of $4 \mu\text{mol kg}^{-1}$). In Fig. 2 a straight line has been drawn to emphasise this. This increase is similar to the deep Greenland Sea anthropogenic CO_2 increase determined by Olsen et al. (2006), which may not seem too surprising considering that the same cruises were compared in this study. However, Olsen et al. (2006) did not take any of the data measured in-between into account, whereas the crossover and inversion analyses do. For a meaningful interpretation of the corrections determined by the inversion we have no other option than to assume that this trend is real. Given this, the following features are noteworthy: (1) the 18HU19820228 TCO_2 data appears to be almost $20 \mu\text{mol kg}^{-1}$ too high, (2) the 58AA19950217 TCO_2 data appears around $5 \mu\text{mol kg}^{-1}$ lower than the data from the other cruises from the same period, and (3) the three cruises 58AA19940203, 58JH19940723, and 58JH19950427 fall in a “high group” (i.e. they should receive a negative correction). Finally, note that no results were obtained for 06MT19920701 because there were too few samples from deeper than 1900 dbar.

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4 Recommendations

Given the outcome of the crossover and inversion analyses we believe that the following cruises should be considered further: 18HU19820228, 58AA19940203, 58JH19940723, 58AA19950217, and 58JH19950427. 06MT19920712 is analysed in Sect. 4.6 and the Iceland Sea time series data are analysed in Sect. 4.7.

4.1 18HU19820228

The TCO_2 data of this cruise appear to be almost $20 \mu\text{mol kg}^{-1}$ higher than expected. They were analysed using potentiometric techniques and not referenced to analysis of CRMs, which were not available at the time these data were acquired. Given the existence of the TTO-NAS data from the previous year one possible solution would be to determine the correction for the 18HU19820228 TCO_2 data that would bring them into consistency with the TTO-NAS data. Pursuing this approach, an MLR analysis was carried out. Following Olsen et al. (2006) the TTO-NAS TCO_2 data obtained north of 62°N and deeper than 50 m were fitted as a function of theta, salinity, nitrate, and silicate:

$$\text{TCO}_2 = 67.79S - 6.53\theta - 0.096\text{NO}_3 + 0.65\text{Si} - 222.14 \quad (1)$$
$$r^2 = 0.74, \text{rms} = 4.2$$

Figure 3 shows deviations of the MLR estimates from the measured 18HU19820228 TCO_2 data, versus station number. The offsets fall into three groups and decreases with increasing station number. There are also trends in the three sub-groups, in particular in the last. Student T tests showed that these three groups were different at the 95% significance level. This cruise is thus not only offset, but the magnitude of this offset varies. No robust adjustment factors can be applied and the recommendation for these data is to flag them questionable.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.2 58AA19940203

These data appear too high, around $5 \mu\text{mol kg}^{-1}$ when compared to data from the same period (Fig. 2). This cruise focused on hydrography and only one chemical oceanographer participated. There are few data, only 85 samples distributed over 16 stations.

- 5 The alkalinity data from this cruise are of limited quality as well, and it is recommended that the TCO_2 data are flagged questionable.

4.3 58JH19940723

The TCO_2 data from cruise appears approximately $9 \mu\text{mol kg}^{-1}$ too high considering the other cruises from the same period and the expected trend (Fig. 2). It took place, however, mainly in the Nordic Seas-North Atlantic overflow regions, outside of the Greenland Sea. Considering the variability of this region and the limited overlap with the other cruises, evidence does not appear strong enough to recommend an adjustment for these data. The label NC is therefore applied. For consistency, the other cruise with carbon data in this region, 58JH19920712, is labelled NC as well. This is consistent with the decisions made for the other parameters measured at these cruises (Olsen et al., 2009; Olafsson and Olsen, 2009; Falck and Olsen, 2009).

4.4 58AA19950217

Not only do the data appear too low (Fig. 2), but also the standard deviation of deep samples was quite high: $6.16 \mu\text{mol kg}^{-1}$, twice that of 316N20020530 which covered a much larger region (Table 2). The data are thus not only offset, but precision is limited as well. Given this, it is recommended that the TCO_2 data from this cruise are flagged questionable.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.5 58JH19950427

The data from this cruise appear approximately $9 \mu\text{mol kg}^{-1}$ too high compared to the expected trend and other data from the same period (Fig. 2). In addition to this, the spread of deep samples was twice that of the deep samples measured during the 316N20020530 cruise, which covered a much larger region (Table 2). Given these issues with precision and accuracy, it is recommended that the TCO_2 data from this cruise are flagged questionable.

4.6 06MT19920701

No results were obtained with the crossover and inversion analysis as there were very few data from deeper than 1900 dbar – the minimum sampling depth for data to be considered in the crossovers. Figure 4 shows the TCO_2 data from 3 stations of the 06MT19920701 cruise, along with more recent as well as earlier (TTO-NAS) data from the same area, north of Jan Mayen. The 06MT19920701 TCO_2 data appears too low, between 5 and $10 \mu\text{mol kg}^{-1}$. A similar offset was evident in the TCO_2 data obtained in the other regions covered by this cruise. Gathering information regarding these data was not trivial, as the group that collected them has ceased to exist. The cruise report does not mention whether CRMs were used or not, but given the age we consider it unlikely. It is also not clear whether these samples were measured using potentiometric or coulometric titration, as both types of analytical systems were used on the cruise. However, the latter was carried for testing and we consider it more likely that the reported values are from the potentiometric system. An offset in these data is thus not unlikely. From Fig. 4 it appears to be between 5 and $10 \mu\text{mol kg}^{-1}$. It cannot be further constrained given the possible anthropogenic change, the scarcity of deep-water samples from this cruise (a total of only 3 samples from below 2000 m), and the lack of nutrients data, which prohibits use of MLR. Therefore, rather than possibly introducing errors it is recommended that these data are flagged questionable.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.7 Iceland Sea

These data are collected at two stations northeast of Iceland and given that the bottom depth is less than 1900 m, they were not included in the crossover and inversion analyses, which require sampling depths larger than 1900 dbar. Using another approach to evaluate these data, Fig. 5 compares TCO₂ and nitrate (NO₃) data from four different years of the Iceland Sea time series with data from four different cruises which had stations nearby. The nitrate data from 58JH19940826 have been adjusted by a factor of 1.1 as recommended by Olafsson and Olsen (2009). Evaluated from these figures, the Iceland Sea time series data appear reasonable.

5 Consistency test of final Nordic Seas TCO₂ data

In order to evaluate the consistency of the final Nordic Seas TCO₂ data, a cnaX crossover and inversion analysis (Tanhua et al., 2009b) was carried out on the set of data after the recommendations had been implemented. The TTO-NAS data were included in this analysis. Corrections determined by the WLSQ inversion of the crossover offsets are shown in Fig. 2. Assuming that the TCO₂ concentrations have increased by approximately 4 μmol kg⁻¹ over the two decades, and taking into account that 58JH19940723 has been labelled Not Considered, the data appears consistent to at least 4 μmol kg⁻¹. The individual cruise tracks and TCO₂ profiles of the Nordic Seas data included in the CARINA product are shown in the supplementary material. Note that the recommendations of this paper have been implemented in the merged data product, and the data that were found to be questionable or bad have not been included. They have been retained in the exchange files, however. When the TCO₂ data from a whole cruise were found to be questionable as reported here, they were not necessarily flagged as such in the exchange files. The flags in these, are the ones assigned during the primary QC.

ESSDD

2, 181–203, 2009

**Nordic Seas total
dissolved inorganic
carbon data in
CARINA**

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Acknowledgements. This effort would not have been possible without the support of the Norwegian Research Council through A-CARB (188167), the EU through IP CARBOOCEAN (511176) and the International Ocean Carbon Coordination Project. Additional support from the International Ocean Carbon Coordination Project (IOCCP) and the Hanse Institute for Advanced Study (HWK, Delmenhorst, Germany) have been very valuable.

References

- Blindheim, J. and Østerhus, S.: The Nordic Seas, main oceanographic features, in: The Nordic Seas: an integrated perspective, edited by: Drange, H., Dokken, T., Furevik, T., Gerdes, R., and Berger, W., AGU Geophysical monograph 158, AGU Washington DC, 2005.
- Chen, A., Jones, E. P., and Lin, K.: Wintertime total carbon dioxide measurements in the Norwegian and Greenland Seas, *Deep-Sea Res.*, 37, 1455–1473, 1990.
- Chierici, M., Drange, H., Anderson, L. G., and Johannessen, T.: Inorganic carbon fluxes through the boundaries of the Greenland Sea Basin based on in situ observations and water transport estimates, *J. Mar. Sys.*, 22, 295–309.
- Dickson, A. G., Sabine, C. L., and Christian, J. R. (Eds.): Guide to best practices for ocean CO₂ measurements, PICES Special Publication 3, 191 pp., 2007.
- Falck, E. and Olsen, A.: Nordic Seas oxygen data in CARINA, *Earth Syst. Sci. Data*, in preparation, 2009.
- Gouretski, V. V. and Jancke, K.: Systematic errors as the cause for an apparent deep water property variability: global analysis of the WOCE and historical hydrographic data, *Prog. Oceanog.*, 48, 337–402, 2001.
- Jeansson, E., Olsson, K. A., Tanhua, T., and Bullister, J. L.: Nordic Seas and Arctic Ocean CFC Data in CARINA, *EEarth Syst. Sci. Data*, in preparation, 2009.
- Johnson, G. C., Robbins, P. E., and Hufford, G. E.: Systematic adjustments of hydrographic sections for internal consistency, *J. Atmos. Ocean. Tech.*, 18, 1234–1244, 2001.
- Johnson, K. M., Wills, K. M., Butler, D. B., Johnson, W. K., and Wong, C. S.: Coulometric total carbon dioxide analysis for marine studies: maximizing the performance on an automated gas extraction system and coulometric detector, *Mar. Chem.*, 44, 167–187, 1993.
- Jutterström, S., Anderson, L. G., Bates, N. R., Bellerby, R., Johannessen, T., Jones, E. P., Key,

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- R. M., Lin, X., Olsen, A., and Omar, A. M.: Arctic Ocean data in CARINA, Earth Syst. Sci. Data, submitted, 2009.
- Kasajima, Y., Olsson, K. A., Johannessen, T., Messias, M.-J., Jeansson, E., Bellerby, R. G. J., and Skjelvan, I.: A submesoscale coherent eddy in the Greenland Sea in 2003, J. Geophys. Res., 111, C07013, doi:10.1029/2005JC003130, 2006.
- 5 Key, R. M. Tanhua, T., Olsen, A., Hoppema, M., Jutterström, S., Schirnack, C., van Heuven, S., Lin, X., Wallace, D. W. R., and Mintrop, L.: The CARINA data synthesis project: Introduction and overview, Earth Syst. Sci. Data, in preparation, 2009.
- 10 Miller, L. A., Chierici, M., Johannessen, T., Nojiri, T., Rey, F., and Skjelvan, I.: Seasonal dissolved inorganic carbon variations in the Greenland Sea and implications for atmospheric CO₂ exchange, Deep-Sea Res. II, 46, 1473–1496, 1999.
- Olafsson, J. and Olsen, A.: Nordic Seas nutrients data in CARINA, Earth Syst. Sci. Data, in preparation, 2009.
- 15 Olsen, A., Key, R. M., Jeansson, E., Falck, E., Olafsson, J., van Heuven, S., Skjelvan, I., Omar, A. M., Olsson, K. A., Anderson, L. G., Jutterström, S., Rey, F., Johannessen, T., Bellerby, R. G. J., Blindheim, J., Bullister, J. L., Pfeil, B., Lin, X., Kozyr, A., Schirnack, C., Tanhua, T., and Wallace, D. W. R.: Overview of the Nordic Seas CARINA data and salinity measurements, Earth Syst. Sci. Data Discuss., 2, 1–25, 2009a, <http://www.earth-syst-sci-data-discuss.net/2/1/2009/>.
- 20 Olsen, A.: Nordic Seas total alkalinity data in CARINA, Earth Syst. Sci. Data, in preparation, 2009b.
- Olsen, A., Omar, A. M., Bellerby, R. G. J., Johannessen, T., Ninnemann, U., Brown, K. R., Olsson, K. A., Olafsson, J., Nondal, G., Kivimäe, C., Kringstad, S., Neill, C., and Olafsdottir, S.: Magnitude and origin of the anthropogenic CO₂ increase and ¹³C Suess effect in the Nordic seas since 1981, Global Biogeochem. Cy., 20, GB3027, doi:10.1029/2005GB002669, 2006.
- 25 Skjelvan, I., Falck, E., Rey, F., and Kringstad, S. B.: Inorganic carbon time series at Ocean Weather Station M in the Norwegian Sea, Biogeosciences, 5, 549–560, 2008, <http://www.biogeosciences.net/5/549/2008/>.
- 30 Tanhua, T., Steinfeldt, R., Key, R. M., Brown, P., Gruber, N., Wanninkhof, R., Perez, F., Körtzinger, A., Velo, A., Schuster, U., van Heuven, S., Bullister, J. L., Stando, I., Hoppema, M., Olsen, A., Kozyr, A., Pierrot, D., Schirnack, C., and Wallace, D. W. R.: Overview of the Atlantic Ocean CARINA data and salinity measurements, Earth Syst. Sci. Data, submitted,

Nordic Seas total dissolved inorganic carbon data in CARINAA. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2009a.

Tanhua, T., van Heuven, S., Key, R. M., Velo, A., Olsen, A., and Schirnack, C.: Quality control procedures and methods used for the CARINA data set, *Earth Syst. Sci. Data*, submitted, 2009b.

5 Tanhua, T. and Wallace, D. W. R.: Consistency of TTO-NAS inorganic carbon data with modern measurements, *Geophys. Res. Lett.*, 32, L14618, doi:10.1029/2005GL023248, 2005.

Wanninkhof, R., Peng, T.-H., Huss, B., Sabine, C. L., and Lee, K.: Comparison of inorganic carbon system parameters measured in the Atlantic Ocean from 1990 to 1990 and recommended adjustments, ORNL/CDIAC-140, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, Tenn., U.S., 43 pp., 2003.

10 Yager, P. L., Wallace, D. W. R., Johnson, K. M., Smith Jr., W. O., Minnett, P. J., and Deming, J. W.: The Northeast Water Polynya as an atmospheric CO₂ sink: A seasonal rectification hypothesis, *J. Geophys. Res.*, 100, 4389–4398, 1995.

ESSDD

2, 181–203, 2009

**Nordic Seas total
dissolved inorganic
carbon data in
CARINA**

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. CARINA cruises with TCO₂ data in the Nordic Seas.

No ^a	EXPOCODE	Country/Institute	PI	CRM ^b	Method	Rec. ^c
10	06MT19920701 ^d	Germany/IBM Hamburg	K. Pegler	Not known	Not known	Flag 3
36	18HU19820228	Canada/BiO	E. P. Jones	no	potentiometric titration	Flag 3
67	316N20020530	Norway/UoB	R. G. . Bellerby	yes	SOMMA	OK
78	32L919920715	USA/Brookhaven	D. W. R. Wallace	yes	SOMMA ^e	NC
79	32L919930718	USA/Brookhaven	D. W. R. Wallace	yes	SOMMA	NC
116	58AA19940203	Norway/UoB	I. Skjelvan	yes	SOMMA	Flag 3
117	58AA19940224	Norway/UoB, Sweden/UGOT	T. Johannessen, L. G. Anderson	yes	SOMMA	OK
118	58AA19940826	Norway/UoB	T. Johannessen	yes	SOMMA	OK
119	58AA19950217	Norway/UoB, Sweden/UGOT	T. Johannessen, L. G. Anderson	yes	SOMMA	Flag 3
120	58AA19961121	Norway/UoB	T. Johannessen	yes	SOMMA	OK
121	58AA19970225	Norway/UoB	T. Johannessen	yes	SOMMA	OK
122	58AA19980308	Norway/UoB	T. Johannessen	yes	SOMMA	OK
125	58AA20010527	Norway/UoB	A. Olsen, A. Omar	yes	SOMMA	OK
128	58GS20030922	Norway/UoB	A. Olsen, T. Johannessen	yes	coulometer	OK
130	58JH19920712	Iceland/MRI	J. Olafsson	yes	coulometer ^f	NC
132	58JH19930730	Norway/UoB	T. Johannessen	yes	SOMMA ^e	Flag 3
134	58JH19940525	Norway/UoB	T. Johannessen	yes	SOMMA	OK
135	58JH19940723	Iceland/MRI	J. Olafsson	yes	coulometer ^e	NC
137	58JH19950427	Norway/UoB	T. Johannessen	yes	SOMMA	Flag 3
138	58JH19951108	Norway/UoB	T. Johannessen	yes	SOMMA	Flag 3
141	58JH19970414	Sweden/UGOT	L. G. Anderson	yes	coulometer	OK
142	58JH19980801	Sweden/UGOT	L. G. Anderson	yes	coulometer	OK
176	74JC19960720	Norway/UoB	T. Johannessen	yes	SOMMA	OK
179	77DN20020420	Sweden/UGOT	L. G. Anderson	yes	SOMMA	OK
184	Iceland Sea	Iceland/MRI	J. Olafsson	yes	coulometer	OK

^a Carina cruisenumber, is cruiseidentifier in merged file.

^b Indicates whether Certified Reference Material (CRM) provided by A. G. Dickson was used on the cruise or not.

^c Recommendations from the secondary QC. Flag 3 = data are questionable and are not included in the final data product; OK = data can be used as they are, NC = These data are probably OK, but could not be fully evaluated.

^d This was the 5th and 6st leg of the 1992 Meteor cruise, only leg 5 samples were used in analyses described in this paper. The method and use of CRMs could not be 100% confirmed. Most likely a potentiometric technique was used. It is very unlikely that CRMs were measured.

^e Analysed ashore.

^f First half of samples measured onboard, the rest ashore.

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Standard deviation of TCO₂ samples collected between 65 and 80° N, 10° W and 10° E, and deeper than 1900 dbar.

No	EXPOCODE	std. dev.
10	06MT19920701	2.92
36	18HU19820228	7.91
67	316N20020530	3.10
79	32L919930718	NA ^a
78	32L919920715	NA ^a
116	58AA19940203	3.15
117	58AA19940224	2.41
118	58AA19940826	2.83
119	58AA19950217	6.16
120	58AA19961121	2.74
121	58AA19970225	2.01
122	58A19980308	3.51
125	58AA20010527	5.06
128	58GS20020922_noEddy ^b	1.11
130	58JH19920712	NA ^a
132	58JH19930730	25.5
134	58JH19940525	1.84
135	58JH19940723	1.41
137	58JH19950427	6.56
138	58JH19951108	10.6
141	58JH19970414	4.51
142	58JH19980801	3.38
176	74JC19960720	3.91
179	77DN20020420	2.92
184	Iceland Sea	NA ^a

^a Cruise had no samples within area and depths under consideration.

^b 58GS20030922 had several stations in a submesoscale coherent eddy (Kasajima et al., 2006), the “_noEddy” signifies that these data were removed prior to this analysis.

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen




Fig. 1. Nordic Seas with TCO_2 sampling positions, includes only data retained after secondary quality control.

[Title Page](#)

[Abstract](#)

[Instruments](#)

[Data Provenance & Structure](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen




Fig. 2. Corrections from WLSQ inversion of crossover offsets for (1) the CARINA data, (2) The CARINA and TTO-NAS data, and (3) the CARINA and TTO-NAS data after completed secondary QC. The straight line has been drawn to emphasise the change between TTO-NAS and 58GS20030922. “no_Eddy” signifies that the stations in the submesoscale coherent eddy encountered at this cruise (Kasajima et al., 2006) were excluded from the analyses.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Nordic Seas total
dissolved inorganic
carbon data in
CARINA

A. Olsen




Fig. 3. Difference between measured 18HU19820228 TCO₂ data and those determined using MLR (Eq. 1), measured – estimated.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen




Fig. 4. TCO₂ data obtained to the north of Jan Mayen at the 06MT19920701 cruise, along with data from the TTO-NAS cruise as well as from the other, more recent, cruises with TCO₂ data in CARINA.

[Title Page](#)

[Abstract](#)

[Instruments](#)

[Data Provenance & Structure](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Nordic Seas total dissolved inorganic carbon data in CARINA

A. Olsen




Fig. 5. Iceland Sea TCO₂ vs. nitrate, along with data from 4 cruises which had stations in the vicinity.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

