

This discussion paper is/has been under review for the journal Earth System Science Data (ESSD). Please refer to the corresponding final paper in ESSD if available.

EPOCA/EUR-OCEANS data-mining compilation on the impacts of ocean acidification

A.-M. Nisumaa^{1,2}, **S. Pesant**³, **R. G. J. Bellerby**^{4,5}, **B. Delille**⁶, **J. Middelburg**^{7,8},
J. C. Orr⁹, **U. Riebesell**¹⁰, **T. Tyrrell**¹¹, **D. Wolf-Gladrow**¹², and **J.-P. Gattuso**^{1,2}

¹CNRS-INSU, Laboratoire d'Océanographie de Villefranche (UMR7093), B.P. 28, 06234 Villefranche-sur-Mer Cedex, France

²Université Pierre et Marie Curie-Paris 6, Observatoire Océanologique de Villefranche, 06230 Villefranche-sur-Mer, France

³MARUM, Center for Marine Environmental Sciences, Leobener Strasse, 28359 Bremen, Germany

⁴Bjerknes Centre for Climate Research, University of Bergen, Bergen, Allégaten 55, 5007 Bergen, Norway

⁵Geophysical Institute, University of Bergen, Bergen, Allégaten 70, 5007 Bergen, Norway

⁶Unité d'Océanographie Chimique, Université de Liège, 4000 Liège, Belgium

⁷Netherlands Institute of Ecology, Centre for Estuarine and Marine Ecology, Korringaweg 7, P.O. Box 140, 4400 AC Yerseke, The Netherlands

EPOCA/EUR-OCEANS data-mining compilation

A.-M. Nisumaa et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



⁸Faculty of Geosciences, Utrecht University, P.O. Box 80021, 3508 TA Utrecht, The Netherlands

⁹LSCE/IPSL, Laboratoire des Sciences du Climat et de l'Environnement, CEA/CNRS/UVSQ, Orme des Merisiers, Bat. 712, 91191 Gif-sur-Yvette cedex, France

¹⁰Leibniz Institute of Marine Sciences, IFM-GEOMAR, Düsternbrooker Weg 20, 24105 Kiel, Germany

¹¹School of Ocean and Earth Science, University of Southampton, National Oceanography Centre Southampton, European Way, Southampton, Hants, SO14 3ZH, UK

¹²Alfred Wegener Institute for Marine and Polar Research, Am Handelshafen 12, 27570 Bremerhaven, Germany

Received: 17 March 2010 – Accepted: 21 March 2010 – Published: 30 March 2010

Correspondence to: A.-M. Nisumaa (nisumaa@obs-vlfr.fr) and J.-P. Gattuso (gattuso@obs-vlfr.fr)

Published by Copernicus Publications.

Abstract

The uptake of anthropogenic CO₂ by the oceans has led to a rise in the oceanic partial pressure of CO₂, and to a decrease in pH and carbonate ion concentration. This modification of the marine carbonate system is referred to as ocean acidification.

5 Numerous papers report the effects of ocean acidification on marine organisms and communities but few have provided details concerning full carbonate chemistry and complementary observations. Additionally, carbonate system variables are often reported in different units, calculated using different sets of dissociation constants and on different pH scales. Hence the direct comparison of experimental results has
10 been problematic and often misleading. The need was identified to (1) gather data on carbonate chemistry, biological and biogeochemical properties, and other ancillary data from published experimental data, (2) transform the information into common framework, and (3) make data freely available. The present paper is the outcome of an effort to integrate ocean carbonate chemistry data from the literature which has been
15 supported by the European Network of Excellence for Ocean Ecosystems Analysis (EUR-OCEANS) and the European Project on Ocean Acidification (EPOCA). A total of 166 papers were identified, 86 contained enough information to readily compute carbonate chemistry variables, and 67 datasets were archived at PANGAEA – The Publishing Network for Geoscientific & Environmental Data. This data compilation is
20 regularly updated as an ongoing mission of EPOCA.

Data access: <http://doi.pangaea.de/10.1594/PANGAEA.735138>

ESSDD

3, 109–130, 2010

EPOCA/EUR-OCEANS data-mining compilation

A.-M. Nisumaa et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

In 2008 fossil fuel use, cement production and changes in land use together have emitted ~ 8.7 PgC to the atmosphere. About half remains in the atmosphere, while the rest is removed in roughly equal parts by ocean and land vegetation (Le Quéré et al., 2009). The world ocean absorbs anthropogenic carbon dioxide (CO_2) at a rate of approximately 24 million tons per day. The uptake of anthropogenic CO_2 by the oceans leads to a rise in the oceanic partial pressure of carbon dioxide ($p\text{CO}_2$), a decrease in pH, and a reduction in carbonate ion concentration (Orr et al., 2005). These changes result in a drop of calcium carbonate saturation state, making seawater less favourable to precipitation of calcareous shells and skeletons and more favourable to their dissolution (Gattuso et al., 1999). The future rate of change is unprecedented in the past 65 Myr (Ridgwell and Schmidt, 2010). The impacts of ocean acidification on the structure and functioning of marine ecosystems, including their services to our society, is the focus of several international research initiatives such as the European Network of Excellence for Ocean Ecosystems Analysis (EUR-OCEANS), the European Project on Ocean Acidification (EPOCA) and a number of national initiatives.

Numerous studies addressed the effects of ocean acidification on organisms and communities (e.g., Gattuso et al., 1999; Riebesell et al., 2000; Delille et al., 2005; Orr et al., 2005), however, the resulting carbonate chemistry and other essential ancillary data are sometimes split in several publications and/or part of the data remains unpublished. Additionally, variables of the carbonate system are reported on different pH scales (N.B.S./N.I.S.T., total, free, or sea water; Dickson, in press), using different sets of equilibrium constants, which make data comparison difficult. The main goals of the EPOCA/EUR-OCEANS data rescue and transformation project are to (1) obtain data from papers already published and archive them as supplementary datasets at PANGAEA – the Publishing Network for Geoscientific & Environmental Data, giving full credit to the original authors, (2) compute a consistent set of carbonate chemistry variables, and (3) safeguard the consolidated dataset at PANGAEA, thus ensuring open access to this unique data.

ESSDD

3, 109–130, 2010

EPOCA/EUR-OCEANS data-mining compilation

A.-M. Nisumaa et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Data on seawater carbonate chemistry and biological processes from experiments and observations from the world ocean were compiled. The project was initiated by EUR-OCEANS in January 2008, with the aim to gather data on the effects of ocean acidification on marine organisms and EPOCA took over in August 2008. This is an ongoing effort that will be continued at least during the lifetime of EPOCA (until May 2012).

2 Data provenance and collection of relevant papers

A literature search was conducted and data from selected papers were obtained directly from the authors as numerical files or, alternatively, data were extracted from tables in the original paper. Papers were selected with the following criteria: (1) only data measured under $p\text{CO}_2$ or pH levels that are relevant to past or for future scenarios (i.e. $p\text{CO}_2$ values ranging from about 100 to 1700 μatm and pH_7 values of 7.5 to 8.5) were considered. (2) Salinity and temperature, at least two variables of the carbonate system were needed (Table 1). Once papers were identified, requests for data were sent to the corresponding authors. When there was no reply after three months, a reminder was sent, and another reminder was sent if there was still no reply after 6 months. If after 9 months from the first request there was still no reply, the data were considered unavailable.

A total of 166 papers were identified in the literature search but 50 did not meet the criteria mentioned above. Of the remaining 116 papers, the authors provided data from 86 papers, 17 did not reply, and 13 were unfortunately unable to locate the data (Fig. 1). In some cases, papers shared data from multidisciplinary studies, for example six papers included data from the 2001 Pelagic Ecosystem CO_2 Enrichment Study (PeECE I). In such cases, data from the related papers were combined into a single dataset, which explains why out of 86 papers, there are 67 datasets archived.

The data collected were from laboratory and mesocosm experiments, as well as field measurements (Fig. 2). Laboratory experiments included the seawater $p\text{CO}_2$ or pH perturbation experiments conducted in different laboratories. It is the category,

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which comprises most papers (44). Mesocosm experiments (26 papers) included three large PeECE projects that were carried out near Bergen (Norway) in 2001, 2003 and 2005, together comprising 23 publications. The database also includes datasets from three publications on mesocosm experiments performed in Hawaii and South Korea.

- 5 The category “field measurements” comprises all datasets reporting natural seawater carbonate chemistry measurements and relevant biological and biogeochemical processes (16 publications).

3 Geographical coverage

The geographical coverage of the studied organisms from the EPOCA/EUR-OCEANS data rescue and transformation project on the impacts of ocean acidification data includes all the world oceans except the Southern Ocean. The investigated organisms were living or collected in these locations, including the phytoplankton and bacteria cultured in laboratories after isolation for several years. In recent years there has been some discussion as to whether phytoplankton and bacteria strains collected several years ago, hundreds of generations later may not be genetically identical to the natural populations.

The areas investigated are divided into two groups. The first group (shown in black in Fig. 3) includes studies of organisms in their natural habitat, organisms that were collected shortly before the experiment and corals that have been maintained in aquaria for several years but are genetically identical to their field counterparts. The second group (shown in red in Fig. 3) comprises phytoplankton and bacteria species that have been collected several years ago but they have gone through hundreds of generations.

Most of the natural community studies were conducted in the Northeast Atlantic (33%), including 1 study above 78°N. About 11% of the studies were performed in the central Western Pacific, 10%, Northwest Pacific and 9% in the central Eastern Pacific. Even though the Mediterranean and Black Sea area represents a relatively small surface area of the world ocean (0.8%) it has contributed to a relatively large percent-

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



age of papers (7%). Studies from the Western Indian Ocean area make up about 6% of the publications comprised in the database. Studies from the Central Western Atlantic and Eastern Indian Ocean only represent 5% and 4% of the papers.

The largest amount of studies was performed with phytoplankton and bacteria species collected from the Northeast Atlantic (10% of the papers). Only 2% of the papers on ocean acidification have been written about studies conducted on organisms collected from the Southeast Atlantic Ocean. The Mediterranean and Black sea area and the central Western Pacific area both represent 1% of the papers (Fig. 3).

In general the Southern Hemisphere and higher latitudes are relatively poorly represented in the EPOCA/EUR-OCEANS data rescue and transformation compilation. The geographical coverage of ocean acidification impact studies is similar to that of coastal biogeochemistry studies (Gattuso et al., 2005). Moreover, there is no direct relationship with the research intensity and the systems vulnerability. Model results suggest that the surface waters of the polar oceans will be the first to become undersaturated with respect to aragonite. It may already occur in 2050 in the Southern Ocean (Orr et al., 2005; McNeil and Matear, 2007) and as early as 2016 in the Arctic Ocean (Steinacher et al., 2009). However there are very few studies performed in these areas. This raises great concern if research efforts are well distributed with respect to the vulnerability of ecosystems.

4 Harmonising the carbonate chemistry data

In order to harmonise data the R package seacarb (Lavigne and Gattuso, 2010) was used to compute a complete and consistent set of carbonate system variables (Table 2). Original data of temperature, salinity, pressure, any pair of carbonate system variables (Table 1) and, if available, the concentrations of silicate and phosphate were used. When the original values were expressed in mol l^{-1} , they were converted to mol kg^{-1} by dividing by density calculated from salinity and temperature as indicated in Dickson et al. (2007). pH, can be measured using different scales, these include the

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



total hydrogen ion, seawater, free and N.B.S./N.I.S.T. (National Bureau of Standards, currently National Institute of Standards and Technology) scales. When pH was reported on a scale other than the total scale, the dissolved inorganic carbon (C_T) and total alkalinity (A_T) were first calculated with CO2SYS (Lewis and Wallace, 1998) and subsequently used in seacarb (which does not handle the N.B.S. scale) to estimate pH on the total scale as well as the other carbonate system variables. The pair of carbonate system variables that was used for computations in seacarb and was CO2SYS is identified by a flag (Table 1). For the sake of traceability, that flag was archived in the datasets as a separate variable called “Carbonate System Computation flag” or “CSC flag”. The dissociation constants used for calculations with seacarb are from Lueker et al. (2000) for carbonic acid (K_1 and K_2), Perez and Fraga (1987) for hydrogen fluoride and those recommended by Dickson et al. (2007) for all other constants.

Each value was archived at PANGAEA (<http://www.pangaea.de>) with its corresponding metadata, including citation, sampling and analysis methods, variable description and units. Once the data were imported in the database, the data providers were requested to quality control their data. All EPOCA/EUR-OCEANS data rescue and transformation project data sets are publicly available.

5 Taxonomic coverage

Studies were mostly carried out on phytoplankton (38 papers published, including 31 on the coccolithophore *Emiliana huxleyi*), scleractinian corals (28 papers), bacteria (24) and zooplankton (16). Few studies were performed on viruses, macroalgae, molluscs, crustaceans, fish and echinoderms (Fig. 4).

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



6 Measured parameters

All data sets include all of the carbonate system variables (Table 2), but the original measurements were made on different sets of variables. The carbonate system variable most measured has been total alkalinity (A_T), which was reported in about 90% of the papers (Fig. 5). The second most measured variable was dissolved inorganic carbon (C_T ; 49%). In 32% of the papers, pH was reported on the N.B.S./N.I.S.T. scale. Out of all data sets, 17% included $p\text{CO}_2$ as one of the original variables, and 6% had measured pH on the free and seawater scale (Fig. 5). The relatively large number of papers that reported pH on the N.B.S./N.I.S.T. scale could be surprising. However, it is only recently that the ocean acidification community has been made aware that this scale should not be used to measure the difference in the concentration constants of the calculation of the hydrogen ion (Dickson, 1993). Only 10% of the papers included in the database reported pH on the total scale (Fig. 5), the scale that is currently recommended (Dickson et al., 2007, 2010) for seawater.

In physiological studies the carbonate system is typically calculated from measurement of pH_{NBS} (calibrated in dilute N.B.S./N.I.S.T. buffers) and a single point titration of alkalinity (A_T). This approach has been widespread appeal because it is simple, inexpensive, and requires very low sample volumes. However, its continued application ignores advances in analytical and theoretical oceanic CO_2 chemistry that suggest fundamental flaws in the assumption $\text{pH}_{\text{NBS}} = -\log[\text{H}^+]$ as a consequence of the high ionic strength of seawater and liquid residual junction errors in the electrode (Crawford and Harrison, 1997).

The biological process most measured was calcification (32% of the papers; Fig. 6). Other variables that are often measured that can potentially be affected by ocean acidification, also include primary production (13%), growth rate (12%), respiration (8%), photosynthesis (7%), reproductive processes (3%) and survival (3%). Primary production here comprises several production variables such as net production rate, particulate organic carbon, inorganic carbon, nitrogen and phosphorus production, community

ESSDD

3, 109–130, 2010

EPOCA/EUR-
OCEANS data-mining
compilation

A.-M. Nisumaa et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



production and net carbon production. The database also comprises 21% other biological variables that were included in single papers. This category comprises variables such as shell length, width and linear extension of molluscs, bleaching, invasion, orientation, different blood cell concentration, concentration in the tissues of organisms, rate of nitrogen fixation.

About 85% of the compiled papers have manipulated only the carbonate chemistry (Fig. 7). Few publications have reported the combined effects of changed carbonate chemistry and light (9%). Some studies have also manipulated temperature (4% of papers) and nutrients (3%) together with carbonate chemistry.

7 Data access

The EPOCA/EUR-OCEANS data rescue and transformation on the impact of ocean acidification is available freely and with no restrictions in the PANGAEA data library at <http://doi.pangaea.de/10.1594/PANGAEA.735138>.

8 Recommendations

During the compilation of the EPOCA/EUR-OCEANS data rescue and transformation several obstacles were met. To streamline future archiving, it is recommended that all data and metadata should be provided with a high level of detail (including the scale on which pH measurements were made) and that seawater carbonate chemistry should be reported in mole per kg of seawater instead of mole per liter of seawater. Other technical recommendations are provided by Pesant et al. (2010).

Some geographical areas have received much more attention than others. It seems essential that data on the response to ocean acidification be gathered in high latitude areas as well as in areas that exhibit naturally low pH, such as the North and Eastern Pacific and upwelling areas. Also more studies are necessary to investigate the

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



combined effects of changes in the carbonate chemistry and changes of other variables such as temperature, irradiance, concentration of nutrients. Some taxonomic groups are rather poorly studied including species of commercial interest such as fish, crustaceans and echinoderms.

5 *Acknowledgements.* We thank all authors who have graciously provided data. N. Taalba contributed to a preliminary data compilation. This work is a contribution to the European Project on Ocean Acidification (EPOCA; which received funding from the European Community's Seventh Framework Programme (FP7/2007–2013) under grant agreement no. 211384) and to the European Network of Excellence for Ocean Ecosystems Analysis (EUR-OCEANS).



The publication of this article is financed by CNRS-INSU.

References

- 15 Crawford, D. W. and Harrison, P. J.: Direct measurement of pCO₂ in cultures of marine phytoplankton: how good is the estimate from pH_{NBS} and single point titration of alkalinity?, *Mar. Ecol. Prog. Ser.*, 158, 61–74, 1997.
- Delille, B., Harlay, J., Zondervan, I., Jacquet, S., Chou, L., Wollast, R., Bellerby, R. G. J., Frankignoulle, M., Borges, A. V., Riebesell, U. and Gattuso J.-P.: Response of primary production and calcification to changes of pCO₂ during experimental blooms of the coccolithophorid *Emiliana huxleyi*, *Global Biogeochem. Cy.*, 19, GB2023, doi:10.129/2004GB002318, 2005.
- 20 Dickson, A. G., Sabine, C. L., and Christian J. R.: Guide to best practices for ocean CO₂ measurements, *PICES Special Publication*, 3, 1–191, 2007.

ESSDD

3, 109–130, 2010

EPOCA/EUR-OCEANS data-mining compilation

A.-M. Nisumaa et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dickson, A. G.: The measurement of sea water pH, *Mar. Chem.*, 44, 131–142, 1993.
- Dickson, A. G.: The carbon dioxide system in sea water: equilibrium chemistry and measurements, in: Guide to best practices in ocean acidification research and data reporting, edited by: Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J.-P., in press, 2010.
- 5 Gattuso, J.-P., Allemand, D., and Frankignoulle, M.: Photosynthesis and calcification at cellular, organismal and community levels in coral reefs: a review on interactions and control by carbonate chemistry, *Am. Zool.*, 39, 160–183, 1999.
- Gattuso, J.-P., Dawson, N., Duarte, C. M., and Middelburg, J. J.: Patterns of publication effort in coastal biogeochemistry: a bibliometric survey (1971–2003), *Mar. Ecol. Prog. Ser.*, 294, 9–22, 2005.
- 10 Lavigne, H. and Gattuso, J.-P.: seacarb: seawater carbonate chemistry with R. R package version 2.3.3, <http://CRAN.R-project.org/package=seacarb> (last access: 29 March 2010), 2010.
- Le Quéré, C., Raupach, M. R., Canadell, J. G., Marland, G., Bopp, L., Ciais, P., Conway, T. J., Doney, S. C., Feely, R., Foster, P., Friedlingstein, P., Gurney, K., Houghton, R. A., House, J. I., Huntingford, C., Levy, P. E., Lomas, M. R., Majkut, J., Metzl, N., Ometto, J. P., Peters, G. P., Prentice, I. C., Randerson, J. T., Running, S. W., Sarmiento, J. L., Schuster, U., Sitch, S., Takahashi, T., Viovy, N., van der Werf, G. R., and Woodward, F. I.: Trends in the sources and sinks of carbon dioxide, *Nature Geosci.*, 2, 831–836, 2009.
- 20 Lewis, E. and Wallace, D. W. R.: Program Developed for CO₂ System Calculations. ORNL/CDIAC-105. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee, 1998.
- Mc Neil, B. I. and Matear, R. J.: Climate change feedbacks on future oceanic acidification, *Tellus B*, 59, 191–198, 2007.
- 25 Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean $p\text{CO}_2$ calculated from dissolved inorganic carbon, alkalinity, and equations for K_1 and K_2 : validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium, *Mar. Chem.*, 70, 105–119, 2000.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G.-K., Rodgers, K. B., Sabine, C. L., Sarmiento, L. J., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M.-F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005.
- 30

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Perez, F. F. and Fraga, F.: Association constant of fluoride and hydrogen ions in seawater, *Mar. Chem.*, 21, 161–168, 1987.

Pesant, S., Hook, L. A., Moncoiffe, G., Nisumaa, A.-M., and Pfeil, B.: Data reporting and data usage, in: *Guide to best practices in ocean acidification research and data reporting*, edited by: Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J.-P., in press, 2010.

Ridgwell, A. and Schmidt, D.A.: Past constraints on the vulnerability of marine calcifiers to massive carbon dioxide release, *Nature Geosci.*, 3, 196–200, 2010.

Riebesell, U., Zondervan, I., Rost, B., Tortell, P. D., and Morel, F. M. M.: Reduced calcification of marine plankton in response to increased atmospheric CO₂, *Nature*, 407, 364–367, 2000.

10 Steinacher, M., Joos, F., Frölicher, T. L., Plattner, G.-K., and Doney, S. C.: Imminent ocean acidification in the Arctic projected with the NCAR global coupled carbon cycle-climate model, *Biogeosciences*, 6, 515–533, 2009, <http://www.biogeosciences.net/6/515/2009/>.

ESSDD

3, 109–130, 2010

**EPOCA/EUR-
OCEANS data-mining
compilation**

A.-M. Nisumaa et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Carbonate System Computation (CSC) flags describing which pair of variables (in addition to salinity and temperature) were used to compute the carbonate system variables listed in Table 2.

CSC flag	Software	Input variables
1	seacarb	pH_T and CO_2
2	seacarb	CO_2 and HCO_3
3	seacarb	CO_2 and CO_3
4	seacarb	CO_2 and A_T
5	seacarb	CO_2 and C_T
6	seacarb	pH_T and HCO_3
7	seacarb	pH_T and CO_3
8	seacarb	pH_T and A_T
9	seacarb	pH_T and C_T
10	seacarb	HCO_3 and CO_3
11	seacarb	HCO_3 and A_T
12	seacarb	HCO_3 and C_T
13	seacarb	CO_3 and A_T
14	seacarb	CO_3 and C_T
15	seacarb	A_T and C_T
21	seacarb	ρCO_2 and pH_T
22	seacarb	ρCO_2 and HCO_3
23	seacarb	ρCO_2 and CO_3
24	seacarb	ρCO_2 and A_T
25	seacarb	ρCO_2 and C_T
26	Step 1. CO2SYS	pH (other scale) and A_T
	Step 2. seacarb	A_T and C_T (obtained with CO2SYS)
27	Step 1. CO2SYS	pH (other scale) and C_T
	Step 2. seacarb	A_T and C_T (obtained with CO2SYS)
28	Step 1. CO2SYS	pH (other scale) and $f\text{CO}_2$
	Step 2. seacarb	A_T and C_T (obtained with CO2SYS)
29	Step 1. CO2SYS	pH (other scale) and ρCO_2
	Step 2. seacarb	A_T and C_T (obtained with CO2SYS)

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. R package seacarb (Lavigne and Gattuso, 2010) computes the following carbonate system variables.

Short variable name	Long variable name and unit
<i>S</i>	Salinity
<i>T</i>	Temperature (°C)
<i>P</i>	Pressure (bar)
pH	pH _T (total scale)*
CO ₂	Carbon dioxide concentration (mol kg ⁻¹)
<i>p</i> CO ₂	Carbon dioxide partial pressure (µatm)
<i>f</i> CO ₂	Carbon dioxide fugacity (µatm)
HCO ₃	Bicarbonate concentration (mol kg ⁻¹)
CO ₃	Carbonate ion concentration (mol kg ⁻¹)
<i>C</i> _T	Dissolved inorganic carbon concentration (mol kg ⁻¹)
<i>A</i> _T	Total alkalinity (mol kg ⁻¹)
OmegaAragonite	Aragonite saturation state
OmegaCalcite	Calcite saturation state

* The definition of pH is implicitly based on a concentration unit for hydrogen ion, although the pH value itself has the dimension 1. The concentration unit for hydrogen ion used here in mol kg⁻¹.

EPOCA/EUR-OCEANS data-mining compilation

A.-M. Nisumaa et al.




Fig. 1. Summary of the data compilation. The number of papers successfully included in the database (archived) versus those in which only one carbonate system variable was reported (NE), papers for which data access was not possible (ND), or for which the data have been lost (Lost).

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



EPOCA/EUR-OCEANS data-mining compilation

A.-M. Nisumaa et al.




Fig. 2. Total number of publications (Total) included in the database, including field measurements (FM), as well as laboratory (LE) and mesocosm experiments (ME).

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Regional distribution of the organisms and communities investigated, broken down following the Food and Agriculture Organization of the United Nations (FAO) major fishing areas. AS, Arctic Sea; ANW, Atlantic, Northwest; ANE, Atlantic, Northeast; AWC, Atlantic, Western Central; ANC, Atlantic, Eastern Central; MBS, Mediterranean and Black Sea; ASW, Atlantic, Southwest; ASE, Atlantic, Southeast; AA, Atlantic, Antarctic; IOW, Indian Ocean, Western; IOE, Indian Ocean, Eastern; IOAS, Indian Ocean, Antarctic and Southern; PNW, Pacific, Northwest; PNE, Pacific, Northeast; PWC, Pacific, Western Central; PEC, Pacific, Eastern Central; PSW, Pacific, Southwest; PSE, Pacific, Southeast; PA, Pacific, Antarctic. Areas of natural communities, organisms that were collected in these areas shortly before the experiment and coral sampling sites are shown in black while organisms that have gone through many generations are shown in red.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 4. Taxonomic coverage of the papers included in the database.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 5. Variables of the seawater carbonate system reported in the papers.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Biological processes reported in the papers: G, calcification rate; OBP, other biological variables; PP, primary production; μ , growth rate; R, respiration; PS, photosynthesis; RP, reproductive processes; and S, survival.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. Percent of the papers that have manipulated the carbonate chemistry alone and those that have manipulated the carbonate chemistry as well as another variable. CC, carbonate chemistry; T, temperature.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

