

Earth Syst. Sci. Data Discuss., 3, 55–78, 2010
www.earth-syst-sci-data-discuss.net/3/55/2010/
© Author(s) 2010. This work is distributed under
the Creative Commons Attribution 3.0 License.

This discussion paper is/has been under review for the journal Earth System Science Data (ESSD). Please refer to the corresponding final paper in ESSD if available.

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Nordic Seas nutrients data in CARINA

J. Olafsson^{1,2} and A. Olsen^{3,4}

¹Marine Research Institute, Skulagata 4, IS 121 Reykjavik, Iceland

²Institute of Earth Sciences, University of Iceland, Sturlugata 7, IS 101 Reykjavik, Iceland

³Bjerknes Centre for Climate Research, UNI Research, Allégaten 55, 5007 Bergen, Norway

⁴Department of Chemistry, University of Gothenburg, 41296 Göteborg, Sweden

Received: 23 February 2010 – Accepted: 26 February 2010 – Published: 3 March 2010

Correspondence to: J. Olafsson (jon@hafro.is)

Published by Copernicus Publications.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Water column data of carbon and carbon relevant hydrographic and hydrochemical parameters from 188 previously non-publicly available cruises in the Arctic, Atlantic, and Southern Ocean have been retrieved and merged into a new database: CARINA (CARbon IN the Atlantic). The data have been subject to rigorous quality control (QC) in order to ensure highest possible quality and consistency. The data for most of the parameters included were examined in order to quantify systematic biases in the reported values, i.e. secondary quality control. Significant biases have been corrected for in the data products, i.e. the three merged files with measured, calculated and interpolated values for each of the three CARINA regions; the Arctic Mediterranean Seas (AMS), the Atlantic (ATL) and the Southern Ocean (SO). With the adjustments, the CARINA database is consistent both internally as well as with GLODAP (Key et al., 2004) and is suitable for accurate assessments of, for example, oceanic carbon inventories and uptake rates, and for model validation. The Arctic Mediterranean Seas is the collective term for the Arctic Ocean and the Nordic Seas, and the quality control was carried out separately in these two areas. This contribution presents an account of the quality control of the nutrients (nitrate, phosphate, and silicate) data from the Nordic Seas in CARINA. Out of the 35 cruises from the Nordic Seas included in CARINA, 33 had nutrients data. The nitrate data from 4 of these appeared to be of so poor quality that they should not be used, for phosphate this number is 7 and for silicate it is 3. We also recommend that the nitrate data from 4 of the cruises should be adjusted, for phosphate and silicate only data from one cruise should be adjusted. The final data appears consistent to 5% based on evaluation of deep data. For nitrate this corresponds to $0.6 \mu\text{mol kg}^{-1}$, and for phosphate and silicate it corresponds to 0.04 and $0.6 \mu\text{mol kg}^{-1}$, respectively.

ESSDD

3, 55–78, 2010

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Data coverage and parameter measured

Repository-Reference: doi: 10.3334/CDIAC/otg.CARINA.AMS.V1.2

Available at: <http://cdiac.ornl.gov/oceans/CARINA/Carina.inv.html>

Coverage: 59.60–82.35° N, 35.23° W–28° E

5 Location Name: Nordic Seas

Date/Time Start: 28-02-1982

Date/Time End: 10-02-2006

10 Parameter list with parameters directly relevant for the nutrient analysis. For a complete list of parameters for the CARINA data base, see (Key et al., 2009). Note the different names for the parameters in the Exchange files (the individual cruise files) and the merged data product.

Data Product Parameter Name	Data Product Flag Name	Exchange File Parameter Name	Exchange File Flag Name	Units
station		STANBR		
day		DATE		
month		DATE		
year		DATE		
latitude		LATITUDE		decimal degrees
longitude		LONGITUDE		decimal degrees
cruiseno				
depth				meters
temperature		CTDTMP		°C
salinity	sf	SALNTY	SALNTY_FLAG_W	
pressure		CTDPRS		decibars
nitrate	no3f	NITRAT	NITRAT_FLAG_W	micromole kg ⁻¹
nitrite	no2f	NITRIT	NITRIT_FLAG_W	micromole kg ⁻¹
silicate	sif	SILCAT	SILCAT_FLAG_W	micromole kg ⁻¹
phosphate	po4f			micromole kg ⁻¹

ESSDD

3, 55–78, 2010

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

CARINA (CARbon In the Atlantic) is a database of carbon and carbon-relevant data from hydrographic cruises in the Arctic, Atlantic and Southern Oceans. The project started as an essentially informal, unfunded project in Delmenhorst, Germany, in 1999 during the workshop on “CO₂ in the North Atlantic”, with the main goal to create a uniformly formatted database of carbon relevant variables in the ocean to be used for accurate assessments of oceanic carbon inventories and uptake rates. The collection and quality control (QC) of the data have been the main focus of the CARINA project. Both primary and secondary QC of the data has been performed.

The CARINA database consists of two parts: the first part is the set of the individual cruise files where all the data reported by the measurement teams are stored. Quality flags are accompanying the data, in many cases these are the flags originally reported, in others they were assigned by R. M. Key, Princeton University (Tanhua et al., 2010) These files are in WHP (WOCE Hydrographic Program) exchange format where the first lines consist of the condensed metadata. There are essentially no calculated or interpolated values in the individual cruise files, with the exceptions of pressure calculated from depth and some bottle salinities that were taken from ctdsal. No adjustments have been applied to any of the data in the exchange files, with the exception that all pH measurements were converted to the seawater pH scale at 25 °C (Velo et al., 2009).

The second part of CARINA consists of three merged, quality controlled and adjusted data files; one for each of the three CARINA regions the Atlantic Ocean (ATL), Arctic Mediterranean Seas (AMS) and Southern Ocean (SO). These files contain all the CARINA data and also include: 1) interpolated values for nutrients, oxygen and salinity if those data were missing and the interpolation could be made according to certain criteria, as described in (Key et al., 2009); 2) calculated carbon parameters; e.g. if total dissolved inorganic carbon (TCO₂) and total alkalinity (TA) were measured, pH was calculated; and 3) instances where bottle salinity was missing or bad and was replaced with CTD salinity. Calculated or interpolated values have been given the qual-

ESSDD

3, 55–78, 2010

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ity flag “0”. In many cases there are additional parameters in the individual cruise files, which have not been included in the secondary QC, such as $\Delta^{14}\text{C}$, $\delta^{13}\text{C}$ and SF_6 . Many of these are included in the merged data files as well.

This report provides an overview of the nutrient (nitrate, phosphate and silicate) data from the Nordic Seas in CARINA and describes the secondary QC of these data. These data are part of the Arctic Mediterranean Seas subset of CARINA (CARINA-AMS). This subset contains data from the Nordic Seas and the Arctic Ocean. Not one of the cruises of CARINA covered both the Arctic Ocean and the Nordic Seas to the extent that it could be included in the secondary quality control in both regions. In addition, the differences in data density in the two regions enforced the use of different QC methods for the data collected in them. The data from these regions were therefore analysed separately. The data from the Arctic Ocean are described by Jutterström et al., 2010 while the other Nordic Seas data are described in Jeansson et al., 2010; Falck and Olsen, 2009; Olsen et al., 2009; and Olsen, 2009a, b. The data acquisition and long term QC procedures for two time series stations, one of which is in the Iceland Sea and included in CARINA-AMS, are described in Olafsson et al. (2009). A more comprehensive description of the complete CARINA database can be found in Key et al. (2009), as well as in the other, more specialised, papers in this special issue.

2 Data Provenance and Structure

The Nordic Seas were loosely defined as the region within the Fram Strait to the north, Greenland to the west, the Greenland-Scotland Ridge to the south, and Norway, the Barents Sea Opening, and Spitsbergen to the east. Out of the 188 CARINA cruises, 62 are included in the CARINA-AMS, and 35 of these considered as Nordic Seas cruises. Five of these are in common with the Atlantic Ocean subset of CARINA (Tanhua et al., 2009b), in order to ensure consistency between the regions. Out of the 35 cruises included in the Nordic Seas CARINA, 33 had nutrient data.

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1 provides an overview of the data that were included in the secondary quality controls described in this paper, and station positions are shown in Fig. 1.

For reasons of analytical methodology, nitrate is detected along with nitrite. In some of the files these were not reported separately, but as nitrate + nitrite. In such cases it was far from trivial, if not impossible to extract the nitrate and nitrite concentrations. However, given that nitrite is normally only a trace constituent or absent, particularly in oxygenated deep water, we follow here the example for CARINA cruises in the Atlantic and report these data as nitrate, ignoring the possible contribution of nitrite (Tanhua et al., 2009b).

3 Quality control methods

Two methods were used to quality control the data : crossover and inversion analysis and evaluation of phosphate-nitrate (P-N) relationships, these are described in the following two sections.

3.1 Crossover and inversion analyses

The crossover and inversion analysis (Gouretski and Jancke, 2001; Johnson et al., 2001) used to evaluate the consistency of the Nordic Seas nutrients data, was carried out using the version of the cnaX routines (Tanhua et al., 2010) described by Olsen et al. (2009). Briefly, the crossover points were found by identifying all station pairs, from different cruises, within 300 km from each other. Each cruise pair's crossover stations were then further partitioned into at most 6 clusters; groups of data within a circle with a radius of at most 175 km, using a k-means clustering routine (Seber, 1984), each containing at least two stations from each cruise. The data from each station in the cluster were then interpolated into 50-m intervals, and the weighted mean offset and standard deviation for each cruise pair was determined using observations deeper than 1900 dbar. Note that this is deeper than in the other CARINA regions (i.e. 1500 dbar),

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in order to avoid effects of ventilation which reached down to 1600 dbar during the 1990s (Ronski and Budeus, 2005). Also note that whereas the crossover analysis is normally carried out using temperature or density as ordinate, depth was used in the Nordic Seas because of the small gradients of both temperature and density in this region.

The offsets and uncertainties from all cruises were used to generate the set of corrections that maximized the consistency between the cruises using a least square model inversion, the weighted least square (WLSQ) inversion of (Johnson et al., 2001), which weights each crossover offset by its standard deviation so that offsets with large uncertainty are less influential than crossovers with small uncertainty. The data from 32L919920715, OMEX1NS and Iceland Sea time series were not included in this analysis because of lack of samples from deeper than 1900 dbar. The data from 32L919930718 were not included either, because of the special region covered by this cruise (East Greenland Shelf and Shelf edge, see Fig. 3 of Olsen et al., 2009 for station positions of each individual cruise). The data from these cruises have been labelled NC (Not considered) in Table 1.

3.2 Phosphate-nitrate relationships

The nutrients are not conservative constituents of sea water. In addition to mixing their concentrations are also affected by biological processes (Redfield et al., 1963). The regional changes in phosphate and nitrate generally follow linear trends in a ratio of $\Delta P:\Delta N=1:16$, the Redfield ratio. The phosphate-nitrate relationships therefore provide an additional tool in the evaluation of nutrient data quality and data adjustments. In the Nordic Seas an account has also to be taken of the nutrient concentration variability associated with water masses. The sea water fluxes through the Nordic Seas include warm northward flowing Atlantic Water, cold and low salinity Polar Water from the Arctic Ocean and, at depth, Arctic- and Arctic Intermediate Waters which flow over the Greenland-Scotland Ridges and contribute to the global ocean thermohaline circulation (Hansen et al., 2008). These different sources result in wide ranges

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in sea water temperature and salinity and affect also the nutrient relationships. The biochemical changes in the low salinity Polar Water follow a linear Redfield slope pattern, $\text{NO}_3 = a\text{PO}_4 + b$, where the slope, a , is similar as for Atlantic Water and near the Redfield value, but the intercept, b , is significantly lower (Stefansson, 1968; Jones et al., 2003). As an additional QC analysis we therefore examined the nutrients in the CARINA Nordic Seas data set after excluding significant Polar Water influences, select salinity >34.7 , and select data with quality flags 2. With this subset of the data the slope of the P-N regression line should be between 15 and 17 and the intercept on the Y-axis (nitrate axis) should be small and possibly slightly negative.

4 Results

The crossover results in themselves can be accessed at the CARINA website (http://cdiac.ornl.gov/oceans/CARINA/Carina_inv.html). Here we only present the corrections suggested by the WLSQ inversion of the crossover offsets (Fig. 2), and the slope and intercept of the P-N relationships are listed in Table 1. The suggested corrections are multiplicative. The CARINA-wide threshold for considering application of corrections to nutrient data were 2% (Tanhua et al., 2009b). However, in the deep Nordic Seas, 2% corresponds to $\sim 0.3 \mu\text{mol kg}^{-1}$ for nitrate, $\sim 0.02 \mu\text{mol kg}^{-1}$ for phosphate and $\sim 0.3 \mu\text{mol kg}^{-1}$ for silicate. These limits are similar to the measurement precision for the nutrients. It was therefore considered more appropriate for the Nordic Seas to set the limits at 5%. These limits are shown in the figure, and for nitrate the following 5 cruises are outside the $\pm 5\%$ limits, 58AA199940203, 58AA19940224, 58AA19940826, 58AA19950217 and 74JC19960720. For phosphate following 6 cruises are outside the 5% limits: 58AA19940203, 58AA19940224, 58JH19940723, 58AA19940826, 34AR19970805, 58AA19980308 and 58GS2000309. And for silicate the following 4 cruises fell outside the 5% limit: 58JH19911105, 58JH19931106, 58AA19940826 and 34AR19970805. These cruises are further investigated in the following.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.1 Recommendations

4.1.1 Nitrate

For nitrate the following 5 cruises are outside the 5% limits, 58AA199940203, 58AA19940224, 58AA19940826, 58AA19950217 and 74JC19960720.

Cruise 58AA19940203

This cruise had only few stations in the northern Greenland Sea. The inversion suggests that these data should be adjusted by a factor of 1.09. Both TCO₂ and TA have been suggested removed from the synthesis product (Olsen, 2009a, b). We see little reason to salvage the nitrate values and suggest that these are flagged 3.

Cruise 58AA19940224

The inversion suggests that these data are 12% too low, and the intercept of the PN regression of -4.115 also suggest low nitrate data. The exact value of the adjustment factor was determined by comparing the nitrate data of 58AA19940224 with data from previous and subsequent good cruises (Fig. 3). As evaluated from this figure an adjustment factor of 1.12 appears very reasonable.

Cruise 58AA19940826

The inversion suggests that these data are 10% too low. The P-N regression slope is within Redfieldian expectations but the intercept, -5.47 , is very low suggesting systematically low nitrate or high phosphate (Table 1). The value of the adjustment factor was further evaluated by comparing the nitrate data of 58AA19940826 with data from previous and subsequent good cruises (Fig. 4). Data selection have been limited to Norwegian Sea where most of the 58AA19940826 cruise was carried out. There is

ESSDD

3, 55–78, 2010

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



not much data, but the 10% adjustment appears entirely reasonable.

Cruise 58AA19950217

- 5 The inversion suggested that this cruise was 13% too low and the P-N regression slope, 10.6, is unrealistic (Table 1). The data show considerable scatter in addition to the offset. Since both TCO₂ and TA have been suggested removed from the synthesis (Olsen, 2009a, b), we see very little reason to salvage these data and suggest that they are flagged 3.

10

Cruise 74JC19960720

- The inversion suggested that this cruise was 20% too high. This result is supported by the rather high P-N slope with its slightly positive intercept (Table 1). The value of the adjustment factor was further evaluated by comparing the nitrate data of 74JC19960720 with data from previous and subsequent good cruises. Based on this analysis the adjustment factor of 0.8 appeared somewhat large (deep mean hitting the -5% limit) , whereas 0.83 appeared reasonable (Fig. 5) Thus 0.83 is the recommended correction factor for these data.

20

The recommended nitrate adjustments are summarized in Table 1.

4.1.2 Phosphate

For phosphate the following 6 cruises are outside the 5% limits: 58AA19940203, 58AA19940224, 58JH19940723, 58AA19940826, 34AR19970805, 58AA19980308 and 58GS2000309.

ESSDD

3, 55–78, 2010

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Cruise 58JH19940723

This cruise was carried out across the Greenland-Scotland Ridge (see Fig. 3 of Olsen et al., 2009 for station positions of each cruise), a highly variable water mass region and outside the core area of the crossover analyses. Given this we have normally chosen to label the data from this cruise (as well as from cruise 58JH19920712 which covered the same area) Not Considered (NC). However, based on the analyses of the Atlantic group (Tanhua et al., 2009a) the phosphate data of 58JH19940723 were flagged questionable (3).

Cruise 34AR19970805

This cruise took place mainly in the Denmark Strait which is outside our main crossover area, and for this cruise we adopt the corrections suggested by the Atlantic group (Tanhua et al., 2009a) for phosphate, as well as for the other nutrients.

As for the other offset cruises, the initial assessment of the phosphate data (Table 1) and the distribution of suggested adjustments (Fig. 2) indicate that the offset cruises have too high phosphate data (i.e. significant adjustment factors are less than one). In addition, the depth profiles (not shown) suggested that the errors are affecting the low concentration range in shallow water samples to a greater extent than deep water samples, particularly in spring and summer. The deep waters are without seasonal variations and the errors in those samples are therefore for other reasons. The causes of the errors are most probably related to sample handling, e.g. filtration of photic layer samples or not, analytical procedures such as blank evaluation or instrumentation. We find therefore no justifiable method to adjust these data. Thus, to get a consistent set of data we suggest that the cruises with high values are flagged 3. This applies to the cruises: 58AA19940203, 58AA19940224, 58AA19940826, and 58AA19980308. Other cruises are not adjusted. The recommended phosphate flags are summarized in Table 1.

ESSDD

3, 55–78, 2010

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.1.3 Silicate

For silicate the following 4 cruises fell outside the 5% limit: 58JH19911105, 58JH19931106, 58AA19940826 and 34AR19970805.

For silicate a complicating factor is the increasing fraction of deep waters from the Arctic Ocean which has raised silicate concentrations in the deep Greenland Sea (Blindheim and Rey, 2004). This trend can be recognised in the inversion results, there is a slight tendency for lower correction factors with time. Taking this into consideration we are reluctant to recommend that adjustments for 58JH19911105 and 58JH19930730 are implemented.

Cruise 34AR19970805

This cruise took place mainly in the Denmark Strait which is outside our main crossover area, and for this cruise we adopt the corrections suggested by the Atlantic group (Tanhua et al., 2009a).

Cruise 58AA19940826

The inversion suggested adjustment factor of 0.75. To further evaluate the adjustment factor, the silicate data of 58AA19940826 was compared with data from other cruises. Figure 6 shows these silicate profiles and reveals that there are clear issues with the data from 58AA19940826. The high regime is very high, silicate around $18 \mu\text{mol kg}^{-1}$, the low range seems reasonable. We suggest that data from this cruise are are flagged 3.

The recommended silicate adjustments are summarized in Table 1.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Consistency of final Nordic Seas nutrient dataset

In order to evaluate the consistency of the final Nordic Seas nutrients data, a crossover and inversion analysis was carried out on the data with the recommendations implemented. The corrections suggested by the WLSQ inversion are shown in Fig. 2.

5 The dataset appears fairly consistent, except for the phosphate data of 58JH19920712 which appear low, this cruise took place over the Greenland Scotland Ridge area and can not be expected to be consistent with the others which were located mainly in the Greenland Sea and the Norwegian Sea, and has been labelled are “NC”. The inversion suggest differences within 5% for all variables. This corresponds to $\sim 0.7 \mu\text{mol kg}^{-1}$ in nitrate, $0.05 \mu\text{mol kg}^{-1}$ in phosphate and $0.6 \mu\text{mol kg}^{-1}$ in silicate.

10 In this work we have concentrated on the differences or anomalies within the collective data. It should also be emphasized, what is apparent from Table 1, that there exists good quality data from the Nordic Seas in different seasons. However, it is clear that in this collection of data there is least consensus on phosphate. Majority of the adjustments suggested for phosphate would result in lower concentrations. Two cruises, in 15 2002 and 2003, 316N20020530 and 58GS20030922, appear to have good quality data from the Greenland and Iceland Seas. There are, however, indications of systematic differences in phosphate concentrations from these two cruises. The P-N regressions, Table 1, show the former one to have $0.06 \mu\text{mol/kg}$ higher deep water phosphate than the latter. The intercept value for this cruise, -1.072 , suggests a positive error in the phosphate determination. On the other hand, the 58GS20030922 cruise has overall 20 the lowest phosphate concentrations of the “good” cruises. However, the differences between these two cruises can not be resolved. Both laboratories involved apply quality control procedures in their routines and have participated in interlaboratory exercises such as QUASIMEME (Wells et al., 1997).

25 It is clear that there are still uncertainties with nutrient data quality that need to be addressed urgently. This issue requires international effort and coordination. In the meantime some measures can be suggested to individual laboratories to improve the

ESSDD

3, 55–78, 2010

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



precision and accuracy of nutrient analyses, e.g.:

- Subscribe and work according to a laboratory performance programme, e.g. QUASIMEME AQ-1 Nutrients in sea water, <http://www.quasimeme.org/>.
- Use Certified Reference Materials, CRM, made up in sea water.
- A laboratory reference material, LRM (Aminot and K erouel, 1998), can be regularly produced and compared to CRMs and QUASIMEME materials. The LRM should be run with each batch of samples.

6 Data access

The whole CARINA database set is published at http://cdiac.ornl.gov/oceans/CARINA/Carina_inv.html. It contains 188 individual cruise files in comma-separated, WHPO exchange format. Condensed metadata are contained in the header of each data file, the data in these files have not been adjusted according to the recommendations presented here. They have, however, been adjusted in the three merged, comma-separated data files with the data products, one for each of the CARINA regions, AMS, ATL and SO. Also note that when the nutrient data from a whole cruise were found to be questionable as reported here, they were not necessarily flagged as such in the exchange files. The flags in these, are the ones assigned during the primary QC.

These files are divided into the three geographical regions of CARINA. No special software is needed to access the data, but software for MATLAB users is offered to facilitate reading of the data.

Acknowledgements. The authors appreciate support from Norwegian Research Council through A-CARB (188167) and CARBON-HEAT (185093), from EU IP CARBOOCEAN (511176) and the International Ocean Carbon Coordination Project (IOCCP). We would also like to express our gratitude to R. M. Key at Princeton University, T. Tanhua at Leibniz-Institut f ur Meereswissenschaften and S. van Heuven at University of Groningen whose extensive efforts made the CARINA project possible. Finally we would like to thank the investigators who

ESDD

3, 55–78, 2010

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



collected the data at sea, often during adverse conditions, and contributed their data to the CARINA project. They are identified, whenever possible, in the metadata included in the exchange files. This is contribution no XXX of the Bjerknes Centre for Climate Research.

References

- 5 Aminot, A. and K  rouel, R.: Pasteurization as an alternative method for preservation of nitrate and nitrite in sea water samples, *Mar. Chem.*, 61, 203–208, 1998.
- Blindheim, J. and Rey, F.: Water-mass formation and distribution in the nordic seas during the 1990s, *ICES J. Mar. Sci.*, 61, 846–863, 2004.
- Falck, E. and Olsen, A.: Nordic Seas dissolved oxygen data in CARINA, *Earth Syst. Sci. Data Discuss.*, 2, 537–553, 2009,
10 <http://www.earth-syst-sci-data-discuss.net/2/537/2009/>.
- Gouretski, V. V. and Jancke, K.: Systematic errors as the cause for an apparent deep water property variability: Global analysis of the woce and historical hydrographic data, *Prog. Oceanogr.*, 48, 337–402, 2001.
- 15 Hansen, B.,   sterhus, S., Turrell, W. R., J  nsson, S., Valdimarsson, H., H  t  n, H., and Olsen, S. M.: The inflow of atlantic water, heat and salt to the nordic seas across the greenland-scotland ridge, in: *Arctic-subarctic ocean fluxes*, edited by: Dickson, R. R., Meincke, J., and Rhines, P., Springer, Dordrecht, 15-43, 2008.
- Jeansson, E., Olsson, K. A., Tanhua, T., and Bullister, J. L.: Nordic Seas and Arctic Ocean CFC
20 data in CARINA, *Earth Syst. Sci. Data*, 2, 79–97, 2010,
<http://www.earth-syst-sci-data.net/2/79/2010/>.
- Johnson, G. C., Robbins, P. E., and Hufford, G. E.: Systematic adjustments of hydrographic sections for internal consistency, *J. Atmos. Ocean. Tech.*, 18, 1234–1244, 2001.
- Jones, E. P., Swift, J. H., Anderson, L. G., Lipizer, M., Civitarese, G., Falkner, K. K., Kattner, G.,
25 and McLaughlin, F.: Tracing pacific water in the north atlantic ocean, *J. Geophys. Res.*, 108, 3116, doi:3110.1029/2001JC001141, 2003.
- Jutterstr  m, S., Anderson, L. G., Bates, N. R., Bellerby, R., Johannessen, T., Jones, E. P., Key, R. M., Lin, X., Olsen, A., and Omar, A. M.: Arctic Ocean data in CARINA, *Earth Syst. Sci. Data*, 2, 71–78, 2010,
30 <http://www.earth-syst-sci-data.net/2/71/2010/>.

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nordic Seas nutrients
data in CARINA

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Kasajima, Y., Olsson, K. A., Johannessen, T., Messias, M.-J., Jeansson, E., Bellerby, R. G. J., and Skjelvan, I.: A submesoscale coherent eddy in the Greenland Sea in 2003, *J. Geophys. Res.*, 111, C07013, doi:10.1029/2005JC003130, 2006.

Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy, C., and Peng, T. H.: A global ocean carbon climatology: Results from global data analysis project (glodap), *Global Biogeochem. Cy.*, 18, GB4031, doi:10.1029/2004GB002247, 2004.

Key, R. M., Tanhua, T., Olsen, A., Hoppema, M., Jutterström, S., Schirnack, C., van Heuven, S., Kozyr, A., Lin, X., Velo, A., Wallace, D. W. R., and Mintrop, L.: The CARINA data synthesis project: introduction and overview, *Earth Syst. Sci. Data Discuss.*, 2, 579–624, 2009, <http://www.earth-syst-sci-data-discuss.net/2/579/2009/>.

Olafsson, J., Olafsdottir, S. R., Benoit-Cattin, A., and Takahashi, T.: The Irminger Sea and the Iceland Sea time series measurements of sea water carbon and nutrient chemistry 1983–2006, *Earth Syst. Sci. Data Discuss.*, 2, 477–492, 2009, <http://www.earth-syst-sci-data-discuss.net/2/477/2009/>.

Olsen, A.: Nordic Seas total dissolved inorganic carbon data in CARINA, *Earth Syst. Sci. Data*, 1, 35–43, 2009a, <http://www.earth-syst-sci-data.net/1/35/2009/>.

Olsen, A.: Nordic Seas total alkalinity data in CARINA, *Earth Syst. Sci. Data*, 1, 77–86, 2009b, <http://www.earth-syst-sci-data.net/1/77/2009/>.

Olsen, A., Key, R. M., Jeansson, E., Falck, E., Olafsson, J., van Heuven, S., Skjelvan, I., Omar, A. M., Olsson, K. A., Anderson, L. G., Jutterström, S., Rey, F., Johannessen, T., Bellerby, R. G. J., Blindheim, J., Bullister, J. L., Pfeil, B., Lin, X., Kozyr, A., Schirnack, C., Tanhua, T., and Wallace, D. W. R.: Overview of the Nordic Seas CARINA data and salinity measurements, *Earth Syst. Sci. Data*, 1, 25–34, 2009, <http://www.earth-syst-sci-data.net/1/25/2009/>.

Redfield, A. C., Ketchum, B. H., and Richards, F. A.: The influence of organisms on the composition of sea-water, in: *The sea*, edited by: Hill, M. N., Interscience Publishers, 26–77, 1963.

Ronski, S. and Budeus, G. T.: Time series of winter convection in the Greenland Sea, *J. Geophys. Res.*, 110, C04015, doi:10.1029/2004JC002318, 2005.

Seber, C. A. F.: *Multivariate observations*, John Wiley & Sons, Inc., NJ, 1984.

Stefansson, U.: Dissolved nutrients, oxygen and water masses in the northern Irminger Sea,

- Deep-Sea Res., 15, 541–575, 1968.
- Tanhua, T., Brown, P. J., and Key, R. M.: CARINA: nutrient data in the Atlantic Ocean, Earth Syst. Sci. Data, 1, 7–24, 2009a, <http://www.earth-syst-sci-data.net/1/7/2009/>.
- 5 Tanhua, T., Steinfeldt, R., Key, R. M., Brown, P., Gruber, N., Wanninkhof, R., Perez, F., Krtzinger, A., Velo, A., Schuster, U., van Heuven, S., Bullister, J. L., Stendardo, I., Hoppema, M., Olsen, A., Kozyr, A., Pierrot, D., Schirnick, C., and Wallace, D. W. R.: Atlantic Ocean CARINA data: overview and salinity adjustments, Earth Syst. Sci. Data, 2, 17–34, 2010b, <http://www.earth-syst-sci-data.net/2/17/2010/>.
- 10 Tanhua, T., van Heuven, S., Key, R. M., Velo, A., Olsen, A., and Schirnick, C.: Quality control procedures and methods of the CARINA database, Earth Syst. Sci. Data, 2, 35–49, 2010, <http://www.earth-syst-sci-data.net/2/35/2010/>.
- Velo, A., Pérez, F. F., Lin, X., Key, R. M., Tanhua, T., de la Paz, M., van Heuven, S., Jutterström, S., and Ríos, A. F.: CARINA data synthesis project: pH data scale unification and cruise
15 adjustments, Earth Syst. Sci. Data Discuss., 2, 421–475, 2009, <http://www.earth-syst-sci-data-discuss.net/2/421/2009/>.
- Wells, D. E., Aminot, A., De Boer, J., Cofino, W., Kirkwood, D., and Pedersen, B.: A review of the achievements of the EU project “QUASIMEME” 1993–1996, Mar. Pollut. Bull., 35, 3–17, 1997.

**Nordic Seas nutrients
data in CARINA**

J. Olafsson and A. Olsen

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Table 1. CARINA cruises with nutrient data from the Nordic Seas, P-N regression coefficients and recommended multiplicative adjustments for the nutrients.

No.	Expocode	Intercept	Slope	<i>r</i>	<i>n</i>	Nitrate adj.	Phosphate adj.	Silicate adj.
36	18HU19820228	0.36	14.5	0.98	1754	1	1	1
67	316N20020530	-1.6	15.9	0.99	2060	1	1	1
78	32L919920715	2.4	11.0	0.45	122	NC	NC	NC
79	32L919930718	7.8	5.25	0.34	83	NC	NC	NC
91	34AR19970805	-2.6	17.7	0.97	99	1.05 ^a	1.04 ^a	1.15 ^a
116	58AA19940203	5.1	6.91	0.73	76	Flag 3	Flag 3	No data
117	58AA19940224	7.6	3.86	0.11	63	1.12	Flag 3	No data
118	58AA19940826	-5.5	16.5	0.89	130	1.10	Flag 3	Flag 3
119	58AA19950217	1.7	10.7	0.75	286	Flag 3	1	1
120	58AA19961121	4.0	11.2	0.98	348	Flag 3	Flag 3	Flag 3
121	58AA19970225	NA ^b	NA ^b	NA ^b	0 ^b	Flag 3 ^b	Flag 3	Flag 3
122	58AA19980308	-0.48	13.4	0.92	268	1	Flag 3	1
128	58GS20030922	-0.12	15.7	0.98	261	1	1	1
129	58JH19911105	-0.30	15.5	0.95	347	1	1	1
130	58JH19920712	0.72	14.2	0.83	52	NC ^c	NC ^c	NC ^c
131	58JH19921105	-0.55	15.2	0.96	523	1	1	1
132	58JH19930730	-4.4	19.1	0.97	418	1	1	1
133	58JH19931106	-0.32	15.2	0.97	653	1	1	1
134	58JH19940525	-4.1	17.7	0.99	452	1	1	1
135	58JH19940723	-4.7	18.4	0.92	431	NC ^c	Flag 3 ^a	NC ^c
136	58JH19941028	-0.77	15.8	0.98	1119	1	1	1
137	58JH19950427	-1.7	15.0	0.96	942	1	1	1
138	58JH19951108	-0.72	15.3	0.99	450	1	1	1
139	58JH19960720	-1.8	16.5	0.99	1324	1	1	1
140	58JH19961030	-0.74	15.6	0.99	330	1	1	1
141	58JH19970414	-0.13	15.0	0.99	1140	1	1	1
142	58JH19980801	-2.7	17.4	0.97	632	1	1	1
143	58JH19990615	-1.7	16.7	0.99	568	1	1	1
144	58JH20000527	-1.4	16.1	0.99	319	1	1	1
176	74JC19960720	0.17	17.5	0.92	431	0.83	1	1
179	77DN20020420	-2.8	17.8	0.98	216	1	1	1
184	Iceland Sea TS	-0.37	15.3	0.93	984	1	1	1
186	OMEX1NS	-5.6	20.2	0.99	57	NC	NC	NC

^a These adjustment factors were derived by the Atlantic group (Tanhua et al., 2009 – nuts paper). ^b Nitrate data were flagged 3 during primary QC (Key et al., 2009), hence no regression analysis could be carried out. ^c Labelled NC because of we felt that the data could not be soundly QC'd given the high hydrographic variability of the area where these data were obtained, the Nordic Seas- North Atlantic overflow regions. See (Olsen et al., 2009) for map of station positions of each individual cruise.

Nordic Seas nutrients data in CARINA

J. Olafsson and A. Olsen




Fig. 1. Nordic Seas with nutrient sampling positions, includes only data retained in CARINA after secondary quality control.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Results of automated crossover and inversion scripts (cna with WLSQ inversion) applied on all cruises in the collection. Dashed lines show 5% limits. Note that the cruises have been sorted by time in this figure. 58GS20030922 (128) had several stations in a submesoscale coherent eddy (Kasajima et al., 2006), the “noEddy” signifies that these data were removed prior to this analysis.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Greenland Sea (73–80° N, 10° W to 5° E) nitrate profiles from 58AA19940224 with (red) and without (blue) corrections applied, along with data from the previous and subsequent cruises. The thick vertical bars at bottom show mean values deeper than 1900 dbar. For reference, red solid and dashes lines have been drawn at drawn at $15 \mu\text{mol kg}^{-1} \pm 5\%$.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 4. Norwegian Sea (61–70° N, 5° W to 10° E) nitrate profiles from 58AA19940826 with (red) and without (blue) corrections applied, along with data from the previous and subsequent cruises. The thick vertical bars at bottom show mean values deeper than 1900 dbar. For reference, red solid and dashes lines have been drawn at drawn at $15 \mu\text{mol kg}^{-1} \pm 5\%$.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 5. Greenland Sea (73–89° N, 25° W to 3° E) nitrate profiles from 74JC19960720 with (red) and without (blue) corrections applied, along with data from the previous and subsequent cruises. The thick vertical bars at bottom show mean values deeper than 1900 dbar. For reference, red solid and dashes lines have been drawn at drawn at $15 \mu\text{mol kg}^{-1} \pm 5\%$.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Silicate profiles from 58AA19940826 and two other cruises.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

