

**Juneau Icefield Mass
Balance Program
1946–2011**

M. Pelto et al.

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto¹, J. Kavanaugh², and C. McNeil³

¹Department of Environmental Sciences, Nichols College, Dudley, MA 01571, USA

²Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, AB T6G 2R3, Canada

³USGS, 4210 University Dr., Anchorage, AK 99508-4626, USA

Received: 30 March 2013 – Accepted: 22 April 2013 – Published: 24 May 2013

Correspondence to: M. Pelto (mspelto@nichols.edu)

Published by Copernicus Publications.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The mass balance records of the Lemon Creek Glacier and Taku Glacier observed by the Juneau Icefield Research Program are the longest continuous glacier mass balance data sets in North America. On Taku Glacier annual mass balance averaged $+0.40 \text{ m a}^{-1}$ from 1946–1985 and -0.08 m a^{-1} from 1986–2011. The recent mass balance decline has resulted in the cessation of the long term thickening of the glacier. Mean annual mass balance on Lemon Creek Glacier has declined from -0.30 m a^{-1} for the 1953–1985 period to -0.60 m a^{-1} during the 1986–2011 period. The overall mass balance change is -26.6 m water equivalent, a 29 m of ice thinning over the 55 yr. Probing transects above the transient snow line (TSL) indicate a consistent balance gradient from year to year. Observations of the rate of summer TSL rise on Lemon Creek and Taku Glacier indicate a comparatively consistent rate of 3.8 to 4.1 m d^{-1} . The relationship between TSL on Lemon Creek and Taku Glacier to other Juneau Icefield glaciers, Norris, Mendenhall, Herbert, and Eagle, is strong with correlations exceeding 0.82 in all cases.

doi:10.5065/D6NZ85N3

1 Introduction

The Juneau Icefield Research Program (JIRP) is the longest ongoing program of its kind in North America, facilitating arctic and alpine education and expeditionary training in the fields of climate science, glaciology and glacial geology. JIRP has examined the mass balance of the Juneau Icefield since 1946, with principal efforts focused on Lemon Creek Glacier and Taku Glacier. This database is the longest glacier field measurement mass balance data set in North America. The data are reported to the World Glacier Monitoring Service (WGMS) annually and is available through the Advanced Cooperative Arctic Data and Information Service (ACADIS). This paper reports

ESSDD

6, 119–151, 2013

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in Alaska at 1375 m (Nolan et al., 1995). Taku Glacier is noteworthy for its positive mass balance from 1946–1988, which resulted from the cessation of calving around 1950 (Pelto and Miller, 1990). The positive mass balance resulting from this dynamic change with calving cessation gives the glacier an unusually high AAR (accumulation area ratio: percentage of glacier in accumulation zone at end of hydrologic year) for a non-calving glacier and makes the glacier relatively insensitive to climate change (Miller and Pelto, 1990; Pelto et al, 2008; Criscitiello et al., 2010).

2.2 Lemon Creek Glacier

Lemon Creek Glacier, Alaska was chosen as a representative glacier for the 1958 IGY global glacier network. This choice was based on its sub-arctic latitude and on the ongoing mass balance program of (JIRP) that had begun in 1948 (Miller, 1972; Pelto and Miller, 1990). JIRP has continued annual balance measurements on Lemon Creek Glacier through the present (Fig. 2). In 1957 Lemon Creek Glacier was 6.4 km long and had an area of 12.67 km² (Heusser and Marcus, 1964). In 1998 the glacier was 5.6 km long and had an area of 11.8 km² (Marcus et al., 1995). From the head of the glacier at 1450 m to the mean ELA at 1050–1100 m the glacier flows northward, in the ablation zone the glacier turns westward terminating at 600 m. The glacier can be divided into four sections: (1) Steep peripheral northern and western margins draining into the main valley portion of the glacier. (2) A low slope (4⁰) upper accumulation zone from 1220 m to 1050 m. (3) A steeper section (6⁰) in the ablation zone as the glacier turns west from 1050–850 m. (4) An icefall (18⁰) leading to the two fingered termini at 600 m. The maximum thickness exceeding 200 m is 1 km above the icefall (Miller, 1972). Lemon Creek Glacier has retreated 1200 m since 1948 and 800 m since 1957, an average of 10–13 m a⁻¹ between 1998 and 2009.

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Mass balance methods

JIRP has relied on applying consistent mass balance methods at standard measurement sites (Pelto and Miller, 1990; Miller and Pelto, 1999; Pelto, 2011). The key annual measurements are: (1) Snowpits at fixed locations on Taku Glacier and Lemon Creek Glacier ranging in elevation from 950 to 1800 m by which the snow water equivalent (SWE) through the entire annual snowpack profile is directly measured. (2) Ablation measurements at survey stakes along survey profiles, along with repeat height measurements of the stakes. (3) Observations of the TSL (transient snow line; snowline at the time of observation) and ELA (equilibrium line altitude: snowline at the end of the melt season) that allow ablation adjustments to snowpack.

3.1 Snowpits

The standard snowpack measurement method used by JIRP is the snowpit. Each snowpit is excavated down to the previous summer surface, identified by a dirty and/or ice horizon and density discontinuity. The most important aspect of the snowpit is accurate identification of the depth. The previous annual layer is well developed during each ablation season on the Lemon Creek Glacier and below 1500 m on Taku Glacier (LaChapelle, 1954). Typically, snowpits hit the previous summer's firn surface, rather than blue ice. The summer's surface is a laterally continuous surface that typically had undergone several freeze/thaw cycles and often has a low density depth hoar just above it. In mass balance the key element is the snow water equivalent (SWE) of the snowpack. This is the mass per unit area of water that would be yielded were the snowpack melted and is the product of depth and snow density. The density of the snowpack is quite variable during the winter and spring, but by July the mean density is generally consistent from year to year and location to location on the Juneau Icefield. LaChapelle (1954) noted that one of the first characteristics apparent upon examination of the snowpit profiles is the remarkable uniformity of firn density in a vertical profile, and in distribution over the glacier, and with time during the summer. LaChapelle

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(1954) found that snowpack density is consistently 540–565 kg m⁻³ after early July on the Taku Glacier; this has been observed by many other detailed studies since (Pelto and Miller, 1990). Despite this fact, density is measured in all snowpit profiles, in part because this is an excellent training tool.

Snow depth can be verified in shallower snowpacks, those less than 3.5 m, using probing or crevasse stratigraphy. Measurements of retained accumulation in the snowpits are completed during late July and August, and are adjusted to end of the balance year values based on the variations of the TSL, observed ablation and the measured balance gradient (Pelto and Miller, 1990; Miller and Pelto, 1999).

For each of the 17 locations where snow pits are utilized on Taku Glacier and 5 locations on Lemon Creek Glacier, a snow pit is dug at a fixed location, verified using GPS. Once onsite, the southern wall of the snow pit is marked off in order to prevent contaminating any density measurements that will be taken; the south wall of the pit is selected for density measurements in order to mitigate any error that may come from ablation caused by direct sunlight on the snow pit wall. Snow pits are always dug at least 50cm into the previous year's firn pack in order to ensure continuity of the layer. Once the snow pit is dug down into the previous year's layer, the southern wall of the snow pit is shaved back to expose a flat, clean face from the top of the snow pit down into the previous year's layer, this face is used to take density measurements of the snow pack. Using a 500 cc snow corer, samples are taken every ten centimeters down the vertical profile of the snow pit into the previous year's layer and the density measured. The final step of the snow pit survey is to record all ice lenses present in the vertical profile of the snow pack (Fig. 3). An ice lens is a horizontal layer of ice formed when water percolates through the snow pack, hits a denser and/or colder layer of snow, spreads out laterally, and refreezes. The depth, thickness, and continuity of all ice lenses are recorded. Due to the small size of these features the density of the lens is assumed at 0.9 g cm⁻³.

On Taku Glacier, six of the snowpit sites are near the ELA ranging from 950–1200 m, six are located at 1200–1400 m and five are located at 1400–1800 m altitude.

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Retained accumulation thickness has been observed at up to 300 points in a single summer season on Lemon Creek Glacier (1998) and 450 measurements on Taku Glacier (1998). Probing is not effective at depths greater than 5 m. The probe used is a 3/4 inch steel probe that easily penetrates ice lenses within the most recent winter snowpack, because the ice layers have comparatively soft unconsolidated firn beneath them. The previous summer surface cannot be penetrated, because the entire layer was melted and refrozen many times, raising its density and cohesion.

Annual layers in the walls of crevasses are often quite obvious. It is similar to reading tree ring width for climate analysis. Crevasse stratigraphy provides a means to view the two dimensional nature of the annual layer, in contrast a point measurement that is yielded by probing or the small scale view provided by a snowpit. Only vertically walled crevasses can be used for these observations. The key to identification of the annual layer in crevasses is the lateral continuity of the ice layer, as no other feature will be continuous. Crevasse stratigraphy is not a standard method used on the Taku Glacier, but has been used since the beginning of the program for validating snow pit snow depth observations in specific regions of the glacier, where snow depths are large and probing cannot be used to validate the snowpits.

4 Results

4.1 Annual mass balance record

On Taku Glacier, the annual ELA has risen 60 m from the 1946–1985 period to the 1986–2010 period. B_a during the two periods were +0.40 and $-0.08 \text{ m.w.e. a}^{-1}$, respectively, indicative of the snowline rise resulting in cessation of the long term thickening of the glacier (Table 1). This overall mass balance change from 1946–2011 is +13.7 m.w.e. (Fig. 4). The long term positive mass balance is continuing to drive its advance (Pelto and Miller, 1990; Post and Motyka, 1995; Pelto et al., 2008). All other outlet glaciers of the Juneau Icefield are retreating, and are thus consistent with

the dominantly negative alpine glacier mass balance that has been observed globally (Zemp et al., 2009).

Ba on Lemon Creek Glacier has declined from $-0.3 \text{ m.w.e. a}^{-1}$ for the 1953–1985 period to $-0.60 \text{ m.w.e. a}^{-1}$ during the 1986–2011 period. The overall mass balance change is -26.6 m , a 29 m ice thickness loss over the 55 yr (Table 1).

4.2 Mass balance record validation

Possible errors in the mass balance record include the sparse nature of measurement points (1 per 37 km^2), extrapolation to the end of the balance year, infrequent measurements of melting in the ablation zone, and measurements carried out by many different investigators. However, Pelto and Miller (1990), suggest that these sources of error are mitigated by: (1) measuring the same locations at the same time using the same methods each year; (2) using nine years of ablation data to extrapolate mass balance in the ablation zone; (3) validation of snow depth variation using probing transects (see next section). The principal error is due to the lack of data from the ablation zone.

The Taku Glacier mass balance record has been validated with geodetic balance information from independent observation of glacier surface elevation change using the ongoing laser altimetry by the University of Alaska, Fairbanks (Sapiano et al., 1998; Arendt et al., 2002; Larsen et al., 2007). This was accomplished from a centerline profile providing a mean glacier thickness change. Surface elevation change is not strictly a measure of mass balance, though it is reported as such (Arendt, 2006). Surface dynamics can also play a role. On Taku Glacier in the vicinity of the ELA annual velocity surveys indicate a consistent ice dynamics from 1950–2006, indicating that surface elevation change should mostly reflect surface mass balance (Pelto et al, 2008). The observed change in Taku Glacier surface elevation was $+0.69 \text{ m a}^{-1}$ from 1948–1993 and -0.28 m a^{-1} from 1993–1997 (Arendt, 2006). The observed mass balance for these periods from field observations is $+0.38 \text{ m a}^{-1}$ for 1948–1993 and -0.60 m a^{-1} for 1993–1997. The surface record includes the large negative mass balance of 1997, while the laser altimetry only includes part of the 1997 ablation season and would tend

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to underestimate thinning by a small amount (Sapiano et al., 1998). Repeat laser altimetry profiling indicates a Ba of -0.21 m a^{-1} for the 1993–2007 period, compared to the JIRP mean Ba of -0.16 m a^{-1} . A comparison of surface elevations from the 2000 Shuttle Radar Topography Mission and a DEM derived from the 1948 USGS mapping indicates a mean Ba of $+0.45 \text{ m a}^{-1}$ versus the JIRP record of $+0.27 \text{ m a}^{-1}$ for the 1948–2000 period (Larsen et al., 2007). The long term observed ice surface elevation changes taken over varied periods using different techniques validates the accuracy of the mass balance record of the Taku Glacier.

On Lemon Creek Glacier the mass balance record determined from field measurements yields a cumulative mass balance of -26.9 mm from 1953–2011. The Ba record of -12.7 m w.e. (13.9 m of ice thickness) from 1957–1989 compares well to the thinning identified from geodetic methods of 1957–1989 of -13.2 (Marcus et al, 1995). The annual balance record of -17.1 m w.e. (-19.0 m of ice thickness) from 1957–1995 compared to an observed ice thickness change of -16.4 m (Sapiano et al., 1998). Airborne surface profiling by the University of Alaska-Fairbanks (Sapiano et al., 1998; Larsen et al., 2007) noted an additional -13.1 m surface elevation change, compared to a surface mass balance of -10 m from 1995–2007. The error in both geodetic programs is less than 1.5 m . In each of the three time intervals using two different ice thickness assessment techniques the annual balance record is confirmed.

5 Probing transects for determination of balance gradient

In late July of 1984, 1998 (Mauri Pelto), 2004, 2005 (Matt Beedle) and 2010, 2011 (Chris McNeil), measured the mass balance along transects from near the TSL at 900 to 1150 m in late July on Taku Glacier by probing at a horizontal interval of 200 m. This method allows direct identification of the mass balance gradient at this elevation (Table 2).

The balance gradient determined from probing between 925 and 1100 m ranges from $3.3\text{--}3.8 \text{ mm m}^{-1}$, with a mean of 3.5 mm m^{-1} . The balance gradient has been

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



consistent on Taku Glacier for each year observed regardless of the respective mass balance (Fig. 5). Standard deviation of accumulation along the transect from the best fit linear regression is 40 mm. The balance gradient can be compared to that determined directly from snowpit measurements at TKGTP5 and DGTP1 at 1000 m on the Taku Glacier (Pelto, 2011). On the date that each snowpit is excavated the difference between measured SWE at each test pit and the TSL elevation on that date provides a direct balance gradient measurement. The balance gradient derived from the snowpits was 2.6–3.5 mm m⁻¹. This is less than that for probing, but the TSL on the date of snowpit excavation is almost always lower than the lowest elevation of the probing transects and represents a lower elevation band of 800 to 1000 m.

On Lemon Creek Glacier the balance gradient generated annually from the ELA and snowpit elevations illustrates the similarity of the balance gradient from year to year. Above the balance gradient parallels the probed gradient in 1998. The balance gradient ranges from 4.6 to 5.1 mm m⁻¹ (Fig. 6).

Balance gradient from TSL variation

The TSL is readily identifiable on 34 scenes acquired from 1995–2011, and was delineated using the software package US Geological Survey Globalization Viewer. The Juneau Icefield falls in path/row 58/19 and 57/19; all images are false color RGB composites, bands 3, 4, and 5, with a 2% linear stretch applied. The spatial resolution of 30 m, combined with mean surface gradients of 0.04–0.08 m m⁻¹, yields an error of less than ±5 m in TSL elevation. The exception is when the TSL rises to 1200 m a.s.l. or is below 900 m a.s.l. Lemon Creek Glacier; in both cases the surface slopes increases leading to higher error margins. The satellite images were georeferenced in ArcMap 9.3 using five topographically unique reference points. The data frame containing imagery and base map was transformed to NAD_1983_UTM_Zone_8N to ensure spatial accuracy for measurements. For years with multiple images, the rate of rise of the TSL is determined. This rate of rise is only calculated for periods of longer than 15 days.

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For example, in 2006 the TSL was identified in five Landsat images on Taku Glacier. The TSL in 2006 rose from 370 m on 26 May, to 575 m on 10 June, 730 m on 5 July, 800 m on 29 July, and finally 980 m on 15 September (Fig. 7; Table 3). The TSL rise ranged from 3.1 to 6.2 m d⁻¹. Mean rise of the TSL for 15 period's averages 4.1 ± 0.9 m d⁻¹ during the July–September period, for the elevation range between 750–1100 m (Table 3; Fig. 7).

The TSL for Lemon Creek Glacier has been observed for 34 dates from 1998–2011 these observations define 18 time periods for which satellite observations were at least 15 days apart (Table 3; Fig. 8). For Lemon Creek Glacier the observed positive TSL migration rates varied from 3.0 to 5.2 m d⁻¹ with a mean of 4.0 ± 0.6 m d⁻¹. The mean TSL migration rate on Lemon Creek Glacier of 3.8 m d⁻¹ compares well with the mean migration rate of 3.7 m d⁻¹ on nearby Taku Glacier (Pelto, 2011). This suggests a consistency in the rate of rise of the TSL from glacier to glacier and year to year on the Juneau Icefield.

6 ELA-TSL observations

Observations of the TSL and ELA can now be reliably made each year using a combination of less frequent Landsat Imagery, and the daily MODIS images (Pelto, 2011; Hock et al., 2007). Use of the latter insures that observations within a short period of the end of the ablation season. The last usable Landsat image for the ablation season is used to assess the TSL for six glaciers of the Juneau Icefield from 1995–2011 (Table 4; Fig. 9). The observed TSL are highly correlated for these dates on all of the glaciers. This paper presents only a single late season TSL from each year; additional analysis is required to determine the rate of change of TSL in September for each glacier. The ELA can be reasonably estimated from the late season TSL observation on each glacier, once both the rate of rise and the date of the end of the ablation season are known. The ELA in turn is a good indicator of mass balance (Mernild et al., 2013). The World Glacier Monitoring Service derives plots of ELA versus *Ba* each year.

The plots generated for the WGMS (2011) from Lemon Creek Glacier are below. The fit is not good for Taku Glacier (Fig. 10). The fit for Lemon Creek Glacier is excellent for the ELA-*Ba* (Fig. 11).

7 Conclusions

5 The mass balance record from Lemon Creek Glacier and Taku Glacier illustrate a decline in mass balance for both glaciers after 1985. Independent geodetic observations of glacier mass balance validate the long term changes quantified by the *Ba* for both glaciers. The balance gradient of the Taku Glacier and Lemon Creek Glacier is consistent in the region near the ELA from year to year. The rate of rise of the TSL is
10 relatively consistent from year to year and glacier to glacier on the Juneau Icefield. The ELA provides a reasonable first estimate of *Ba* on Taku Glacier and Lemon Creek Glacier, and as such determination of this relationship for other Juneau Icefield glacier utilizing simultaneous TSL variations with Lemon Creek and Taku Glacier has value.

References

- 15 Arendt, A. A., Echelmeyer, K. A., Harrison, W. D., Lingle, C. S., and Valentine, V.B.: Rapid wastage of Alaska glaciers and their contribution to rising sea level, *Science*, 297, 382–286, 2002.
- Criscitiello, A., Kelly, M., and Tremblay, B.: The Response of Taku and Lemon Creek Glaciers to Climate, *Arct. Antarct. Alp. Res.*, 42, 34–44, 2010.
- 20 Echelmeyer, K., Nolan, M., Motyka, R., and Trabant, D.: Ice thickness Measurements of Taku Glacier, Alaska, USA, and their Relevance to it Recent Behavior, *J. Glaciol.*, 41, 541–552, 1995.
- Hock, R., Koostra, D., and Reijmeer, C.: Deriving glacier mass balance from accumulation area ratio on Storglaciären, Sweden, in: *Glacier Mass Balance Changes and Meltwater Discharge* IAHS, 318, 163–170, 2007.
- 25

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Heusser, C. E. and Marcus, M. G.: Surface movement, hydrological change and equilibrium flow on Lemon Creek Glacier, Alaska, *J. Glaciol.*, 5, 61–75, 1964.
- LaChapelle, E. R.: Snow studies on the Juneau Icefield, in: 1589: Variability of glacier mass American Geographical Society, edited by: Walters, R. A. and Meier, M. F., JIRP Report no. 9, 1954.
- Larsen, C. F., Motyka, R. J., Arendt, A. A., Echelmeyer, K. A., and Geissler, P. E.: Glacier changes in southeast Alaska and northwest British Columbia and contribution to sea level rise, *J. Geophys. Res.*, 112, F01007, doi:10.1029/2006JF000586, 2007.
- March, R. S. and Trabant, D. C.: Mass balance, meteorological, ice motion, surface altitude, and runoff data at Gulkana Glacier, Alaska, 1992 balance year: U.S. Geological Survey Water-Resources Investigations Report 95-4277, 1996.
- Marcus, M. G., Chambers, F. B., Miller, M. M., and Lang, M.: Recent trends in The Lemon Creek Glacier, Alaska, *Phys. Geogr.*, 16, 150–161, 1995.
- Mayo, L. R., Trabant, D. C., and March, R. S.: A 30-Year Record of Surface Mass Balance (1966–95), and Motion and Surface Altitude (1975–95) at Wolverine Glacier, Alaska, U.S. Geological Survey Open-File Report 2004–1069, 2004.
- Mernild, S. H., Pelto, M. S., Malmros, J. K., Yde, J. C., Knudsen, N. T., and Hanna, E.: Identification of ablation rate, ELA, AAR, and net mass balance using transient snowline variations on two Arctic glaciers, *J. Glaciol.*, 215, in press, 2013.
- Miller, M. M.: A principles study of factors affecting the hydrological balance of the Lemon Creek Glacier system and adjacent sectors of the Juneau Icefield, SE Alaska, 1965–1969, Institute of Water Research, Michigan State University and the US Federal Office of Water Resources Research, Department of Interior, Publication No. 33, 1–295, 1972.
- Miller, M. M. and Pelto, M. S.: Mass Balance measurements on the Lemon Creek Glacier, Juneau Icefield, AK 1953–1998, *Geogr. Ann.*, 81A, 671–681, 1999.
- Motyka, R. J. and Echelmeyer, K. A.: Taku Glacier (Alaska, U.S.A.) on the move again: active deformation of proglacial sediments, *J. Glaciol.*, 49, 50–59, 2003.
- Pelto, M.: Utility of late summer transient snowline migration rate on Taku Glacier, Alaska, *The Cryosphere*, 5, 1127–1133, doi:10.5194/tc-5-1127-2011, 2011.
- Pelto, M. and Miller, M. M.: Mass Balance of the Taku Glacier, Alaska from 1946 to 1986, *Northwest Sci.*, 64, 121–130, 1990.

Pelto, M. S., Miller, M. M., Adema, G. W., Beedle, M. J., McGee, S. R., Sprenke, K. F., and Lang, M.: The equilibrium flow and mass balance of the Taku Glacier, Alaska 1950–2006, *The Cryosphere*, 2, 147–157, doi:10.5194/tc-2-147-2008, 2008.

5 Post A. and Motyka, R.: Taku and Le Conte Glaciers, Alaska: Calving speed control of late-Holocene asynchronous advances and retreats, *Phys. Geogr.*, 16, 59–82, 1995.

Ramage, J. M., Isacks, B. L., and Miller, M. M.: Radar Glacier zones in southeast Alaska, U.S.A.: field and satellite observations, *J. Glaciol.*, 46, 287–296, 2000.

Sapiano, J. J., Harrison, W. D., and Echelmeyer, K. A.: Elevation, volume and terminus changes of nine glaciers in North America, *J. Glaciol.*, 44, 119–135, 1998.

10 Zemp, M., Hoelzle, M., and Haerberli, M.: Six decades of glacier mass-balance observations: a review of the worldwide monitoring network, *Ann. Glaciol.*, 50, 101–111, 2009.

ESSDD

6, 119–151, 2013

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Mass balance record of the Taku and Lemon Creek Glacier including annual mass balance (*Ba*), ELA, AAR and cumulative *Ba*.

Year	Taku <i>Ba</i> (mm)	LC <i>Ba</i> (mm)	LC cumulative <i>Ba</i> (mm)	Taku cumulative <i>Ba</i> (mm)	Taku ELA (m)	LC ELA (m)	LC AAR	Taku AAR
1946	-40			-40	980			85
1947	360			320	900			88
1948	510			830	870			89
1949	930			1760	800			90
1950	-180			1580	1010			84
1951	-340			1240	1160			68
1952	160			1400	950			86
1953	-150	-560	-560	1250	1010	1080	40	84
1954	-70	-180	-740	1180	1160	1025	58	68
1955	970	1120	380	2150	950	810	90	86
1956	-130	-640	-260	2020	1010	1075	40	84
1957	-40	0	-260	1980	980	1000	62	85
1958	210	-580	-840	2190	780	1040	52	91
1959	350	-900	-1740	2540	1000	1150	33	84
1960	160	-820	-2560	2700	1010	1130	35	84
1961	480	-240	-2800	3180	930	1080	40	87
1962	390	-690	-3490	3570	915	1110	37	88
1963	570	170	-3320	4140	950	970	66	86
1964	1130	1040	-2280	5270	885	885	80	90
1965	790	80	-2200	6060	900	980	65	90
1966	80	-490	-2690	6140	875	1100	38	89
1967	250	-600	-3290	6390	750	1130	36	91
1968	460	-220	-3510	6850	810	1060	58	90
1969	1170	210	-3300	8020	965	1000	63	86
1970	760	-90	-3390	8780	930	1060	58	91
1971	630	-400	-3790	9410	885	1110	39	88
1972	420	-650	-4440	9830	730	1140	34	92
1973	520	-520	-4960	10350	825	1110	39	90
1974	580	-370	-5330	10930	850	1090	42	89
1975	850	290	-5040	11780	800	1010	63	90
1976	660	-250	-5290	12440	850	1080	48	89
1977	470	-480	-5770	12910	885	1110	40	88
1978	310	-800	-6570	13220	915	1150	35	87

**Juneau Icefield Mass
Balance Program
1946–2011**

M. Pelto et al.

Title Page

Abstract Instruments

Data Provenance & Structure

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Continued.

Year	Taku <i>Ba</i> (mm)	LC <i>Ba</i> (mm)	LC cumulative <i>Ba</i> (mm)	Taku cumulative <i>Ba</i> (mm)	Taku ELA (m)	LC ELA (m)	LC AAR	Taku AAR
1979	140	-630	-7200	13360	950	1110	40	86
1980	540	-270	-7470	13900	870	1100	42	89
1981	120	-810	-8280	14020	980	1120	40	85
1982	150	-430	-8710	14170	950	1070	51	86
1983	-420	-1620	-10330	13750	1085	1220	17	79
1984	640	-250	-10580	14390	875	1010	65	89
1985	1400	330	-10250	15790	600	965	75	93
1986	1200	-510	-10760	16990	720	1070	51	92
1987	390	-840	-11600	17380	910	1100	43	88
1988	600	110	-11490	17980	890	1000	69	88
1989	-810	-1240	-12730	17170	1115	1130	40	74
1990	-450	-1110	-13840	16720	1080	1125	42	79
1991	380	-380	-14220	17100	900	1050	60	88
1992	170	-660	-14880	17270	940	1075	54	86
1993	-40	-980	-15860	17230	980	1130	43	85
1994	90	-760	-16620	17320	970	1100	46	85
1995	-760	-1310	-17930	16560	1050	1150	38	81
1996	-960	-1580	-19510	15600	1150	1370	5	68
1997	-1340	-1810	-21320	14260	1225	1400	5	60
1998	-980	-1460	-22780	13280	1120	1300	7	73
1999	400	200	-22580	13680	900	1020	68	88
2000	1030	650	-21930	14710	750	900	82	91
2001	880	400	-21530	15590	850	950	77	89
2002	100	-250	-21780	15690	975	1025	67	88
2003	-900	-1900	-23680	14790	1100	1400	5	77
2004	-830	-1250	-24930	13960	1100	1150	59	86
2005	-720	-470	-25400	13240	1050	1050	61	86
2006	230	-170	-25570	13470	975	1025	68	86
2007	480	150	-25420	13950	930	1000	72	87
2008	750	800	-24620	14700	800	920	80	90
2009	-310	-700	-25320	14390	960	1060	51	86
2010	-120	-580	-25900	14270	1000	1075	55	83
2011	-550	-720	-26620	13720	1025	1100	47	82

ESSDD

6, 119–151, 2013

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

[Title Page](#)

[Abstract](#) [Instruments](#)

[Data Provenance & Structure](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 2. Accumulation measurement on probing transect above the TSL on Taku Glacier in July 1998, 2004, 2005 and 2010.

Pt. Name	Easting	Northing	Elevation	2004	2005	1998	2010
P1	541911	6501411	1099	1001	1030	910	
P2	542013	6501311	1097	938	980		
P3	542115	6501200	1094	954	960	850	
P4	542217	6501111	1092	925	1221		
P5	542319	6501000	1090	893	1177	770	
P6	542454	6500911	1089	882	1148	800	
P7	542568	6500800	1087	873	960		
P8	542669	6500712	1085	859	962	850	
P9	542783	6500601	1084	815	1075		
P10	542885	6500501	1082	871	1112	910	
P11	542998	6500412	1081	916	1110		
P12	543100	6500312	1078	936	1030		1370
P13	543202	6500201	1077	889	1143	880	1434
P14	543304	6500101	1074	902	1156		1419
P15	543395	6500012	1073	992	1171		1435
P16	543519	6499901	1070	1001	1218	770	1443
P17	543599	6499801	1068	1035	1302		1413
P18	543712	6499701	1066	1004	1307	750	1385
P19	543814	6499612	1063	963	1312		1316
P20	543905	6499512	1060	981	1318		1327
P21	543996	6499412	1057	959	1344	850	1365
P22	544086	6499312	1056	945	1278		1396
P23	544189	6499201	1052	983	1278		1383
P24	544291	6499113	1050	950	1259	830	1350
P25	544393	6499001	1047	927	1164		1350
P26	544495	6498901	1044	920	1161	830	1303
P27	544574	6498813	1042	866	1116		1238
P28	544688	6498713	1039	844	1142		1295
P29	544790	6498602	1036	814	1078	740	1226
P30	544903	6498513	1033	796	1092		1202
P31	544994	6498391	1031	774	1016		1225
P32	545074	6498302	1030	734	1052	740	1208
P33	545187	6498202	1027	707	1041		1213
P34	545289	6498102	1025	689	1022	660	1180

**Juneau Icefield Mass
Balance Program
1946–2011**

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Continued.

Pt. Name	Easting	Northing	Elevation	2004	2005	1998	2010
P35	545380	6498002	1022	684	968		1154
P36	545482	6497891	1021	697	963	660	1120
P37	545562	6497791	1019	666	975		1141
P38	545642	6497713	1016	670	957	610	1118
P39	545733	6497613	1013	639	941		1073
P40	545846	6497513	1010	661	922	550	1071
P41	545948	6497425	1006	668	938		1090
P42	546039	6497313	1003	621	901	560	1109
P43	546130	6497213	1000	659	889		1090
P44	546232	6497102	996	639	858	490	1099
P45	546323	6497025	994	605	879		1079
P46	546414	6496914	992	565	834	550	1040
P47	546516	6496803	989	617	843		1025
P48	546607	6496714	988	590	841	440	1066
P49	546698	6496603	985	617	815		1070
P50	546800	6496503	981	646	726	410	1135
P51	546891	6496403	977	576	789		1096
P52	547005	6496314	972	574	675	380	1135
P53	547107	6496214	967	601	725		1116
P54	547198	6496114	963	583	688		1116
P55	547300	6496003	960	587	694	360	1077
P56	547391	6495903	956	405	680		1025
P57	547494	6495792	953	330	606	310	1036
P58	547596	6495715	949		780		1034
P59	547709	6495604	944		757		945
P60	547789	6495504	940		680		913
P61	547914	6495393	935		744		905
P62	547994	6495304	930		606		911

Table 3. Transient snowline observation on Lemon Creek Glacier (LCG) and Taku Glacier (TG), and the respective rates of rise between image dates at least 16 days apart. The date listed is the final date of the measurement interval.

Date	LC TSL (m)	Lemon Creek TSL rate of rise (m d^{-1})	Taku TSL (m)	Taku TSL rate of rise (m d^{-1})
7/11/1998	950		850	
7/30/1998	1025	3.95	880	
8/20/1998	1100	3.57	980	4.76
9/16/1998	1200	3.85	1075	4.42
8/31/1999	950		850	
8/29/2000	900		775	
8/15/2001	900		800	
10/3/2002	1025		950	
7/12/2003	1040		915	
8/5/2003	1110	3.33	950	
8/22/2003	1170	3.53	1075	3.90
7/15/2004	950		850	
8/8/2004	1075	5.21	930	3.48
8/16/2004	1100	4.69	950	3.13
8/24/2004	1150	4.69	980	3.13
9/1/2004	1100	3.13	1050	6.25
8/10/2005	1050	3.92	920	
9/11/2005	1050		1000	4.06
7/29/2006	935		760	
9/15/2006	1025	3.04	975	4.48
8/8/2007	875		850	
8/16/2007	925		900	
9/2/2007	1000	4.41	965	3.82
7/2/2008	800		400	
8/19/2008	900	3.38	775	
7/13/2009	900		750	
8/5/2009	975	3.57	825	3.26
8/29/2009	1050	3.13	900	3.13
9/14/2009	1050		950	3.13
7/8/2010	925		580	
8/3/2010	1000	3.85	775	
8/14/2010	1050	3.57	800	
8/28/2010	1075	3.26	900	5.00

**Juneau Icefield Mass
Balance Program
1946–2011**

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4. TSL observation on the same date from Landsat Images on six Juneau Icefield glaciers.

Date	Norris	Taku	Lemon Creek	Mendenhall	Herbert	Eagle
8/29/1995	975	1025	1125	1075	1100	1050
9/4/1996	1050	1075	1150	1125	1150	1175
9/6/1997	1100	1125	1300	1200	1200	1175
9/16/1998	1050	1075	1200	1175	1150	1150
8/31/1999	750	850	950	900	925	900
8/29/2000	750	775	900	875	925	900
8/15/2001	800	800	900	900	925	900
10/3/2002	925	950	1025	1050	1075	1025
10/1/2003	1000	1075	1300	1150	1175	1150
9/1/2004	1075	1050	1100	1150	1200	1200
9/11/2005	1000	1000	1050	1050	1100	1100
9/16/2006	1025	975	1025	1125	1150	1150
9/22/2007	925	925	1000	1050	1050	1025
8/19/2008	700	775	900	850	900	900
9/14/2009	925	950	1050	1050	1050	1050
9/18/2010	950	975	1075	1075	1050	1025
9/11/2011	1000	975	1100	1150	1150	1125




Fig. 1. Base map of the Juneau Icefield, indicating the glaciers examined in this study. LC = Lemon Creek Glacier, bold black line = Probing transect on Taku Glacier. Black dots = snowpit locations on Taku Glacier.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Map of Lemon Creek Glacier indicating primary snow pit locations, and the TSL location on specific dates in 2003.

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Fig. 3. Snowpit on Taku Glacier in 2011. Note the ice lenses. Density measurements are being taken from the south wall; the tape measure aids recording ice lens and sample depths.

ESSDD

6, 119–151, 2013

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ESSDD

6, 119–151, 2013

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.

[Title Page](#)

[Abstract](#)

[Instruments](#)

[Data Provenance & Structure](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 4. Cumulative annual mass balance of Taku (red) and Lemon Creek Glacier (blue).

Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.




Fig. 5. Taku Glacier balance gradient determined from probing in various years. Note the similar gradient in this elevation range of the Taku Glacier in July near the ELA.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.




Fig. 6. Annual balance gradient on Lemon Creek Glacier for 1998 and 2003–2011, and probing gradient from 1998.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. TSL identification on Taku Glacier in 2006 Landsat image from 9/14/2006. A = 5/26/2006, B = 7/5/2006, C = 7/28/2006, D = 9/14/2006.

**Juneau Icefield Mass
Balance Program
1946–2011**

M. Pelto et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.




Fig. 8. TSL elevation rise rate on Taku Glacier and Lemon Creek Glacier.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.




Fig. 9. Transient snowline elevation on the same date on six Juneau Icefield glaciers 1995–2011.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Juneau Icefield Mass Balance Program 1946–2011

M. Pelto et al.




Fig. 10. Relationship of Taku Glacier annual mass balance and the ELA.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Juneau Icefield Mass
Balance Program
1946–2011**

M. Pelto et al.




Fig. 11. Relationship of Lemon Creek Glacier annual mass balance and the ELA.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)