

Use of Various Remote Sensing Land Cover Products for PFT Mapping over Siberia

C. Ottlé¹, J. Lescure¹, F. Maignan¹, B. Poulter¹, T. Wang¹, N. Delbart²

1- LSCE-IPSL, UMR8212, CNRS-CEA-UVSQ, Orme des Merisiers, Gif-sur-Yvette, France

2- PRODIG, UMR8586, Université Paris-Diderot, Paris, France

Corresponding author: C. Ottlé (catherine.ottle@lsce.ipsl.fr)

Abstract:

High-latitude ecosystems play an important role in the global carbon cycle and in regulating the climate system and are presently undergoing rapid environmental change. Accurate land cover datasets are required to both document these changes as well as to provide land-surface information for benchmarking and initializing earth system models. Earth system models also require specific land cover classification systems based on Plant Functional Types (PFT), rather than species or ecosystems, and so post-processing of existing land cover data is often required. This study compares over Siberia, multiple land cover datasets against one another and with auxiliary data to identify key uncertainties that contribute to variability in PFT classifications that would introduce errors in earth system modeling. Land cover classification systems from GLC 2000, GlobCover 2005 and 2009, and MODIS collections 5 and 5.1 are first aggregated to a common legend, and then compared to high-resolution land cover classification systems, continuous vegetation fields (MODIS-VCF) and satellite-derived tree heights (to discriminate against sparse, shrub, and forest vegetation). The GlobCover dataset, with a lower threshold for tree cover and taller tree heights and a better spatial resolution, tends to have better distributions of tree cover compared to high-resolution data. It has therefore been chosen to build new PFTs maps for the ORCHIDEE land surface model at 1 km scale. Compared to the original PFT dataset, the new PFT maps based on GlobCover 2005 and an updated cross-walking approach mainly differ in the characterization of forests and degree of tree cover. The partition of grasslands and bare soils now appears more realistic compared with ground-truth data. This new vegetation map provides a framework for further development of new PFTs in the ORCHIDEE model like shrubs, lichens and mosses, to better represent the water and carbon cycles in northern latitudes. Updated land cover datasets are critical for improving and maintaining the relevance of earth system models for assessing climate and human impacts on biogeochemistry and biophysics.

The new PFT map at 5 km scale is available for download from the PANGAEA website, at: <http://doi.pangaea.de/10.1594/PANGAEA.810709>

Keywords: Land cover, PFTs mapping, Siberia, remote sensing, vegetation

1. Introduction

The Siberian region has been a focus of research attention in recent years because it is considered as a hot spot for climate change studies (see for example Lenton et al., 2008). The region is currently undergoing a warming trend with impacts already visible in the environment, its vegetation and soils (Lucht et al., 2002). Pronounced climatic warming in Siberia (Chapin et al., 2005) has had large implications on vegetation, changes which have been already confirmed by numerous studies at various scales. For example, Tape et al. (2006) demonstrated using aerial photography, an expansion of deciduous shrubs in tundra areas in northern Alaska during the last 50 years. Satellite datasets and especially NDVI products have also documented landscape-scale greening signals and/or phenological changes, in relation with air temperature (see for example, Forbes et al., 2010, Huttich et al., 2007, Delbart et al., 2005, 2007, Myneni et al., 2001). However, the response of continental-scale vegetation shifts due to climate warming is not simple because different processes and feedbacks linked to snow, permafrost, soil moisture, albedo, and species competition (Chapin et al., 2005, Lorant and Goetz, 2012), lead to large uncertainties in predicting and attributing ecosystems and land-cover change dynamics.

One approach to better understand the role of interacting processes and how the various species compete for water, light and nutrients, is the use of ecosystem models. Ecosystem models are now able to represent the main high latitudes physical and biogeochemical processes and especially, permafrost and snow modeling and vegetation interactions, as well as vegetation dynamics, but these models require a correct representation of current land coverage as initial conditions or for benchmarking dynamic global vegetation models (Quaife et al., 2008).

In Northern Eurasia, the main challenge for ecosystem modelers is to be able to differentiate short from high-statured vegetation, as well as deciduous from evergreen phenology. Even at this coarse thematic resolution, very different energy, water and carbon cycling processes are represented. For example, vegetation height is directly related to surface roughness and consequently affects turbulent fluxes; in addition, vegetation height can alter the effects of snow on ecosystem energy budgets with implications for surface albedo and related feedbacks. The deciduous character of shrubs or trees is also very important for the calculation of spring and autumn water and carbon fluxes and their seasonal variations.

Improved mapping of current land cover is a high priority for representation within earth system models, yet there are several challenges that need to be considered. Remote sensing instruments provide regular data at global scales, with increasing spatial resolution, and have been used for years to map land cover. Thus, a number of global products have been derived over the last 20 years. They are used for a wide range of environmental studies and especially in climate models to characterize the land surface and its physical and biogeochemical properties and to determine the energy and matter transfers to the atmosphere. In such models, for simplification, to reduce the computer time, and to develop testable hypotheses, the various ecosystems are grouped in Plant Functional Types (PFT), with a limited number of types, usually around 10 to 15. As an example, the ORCHIDEE Dynamic Global Vegetation Model (DGVM) (Krinner et al., 2005), part of the Institut Pierre Simon Laplace (IPSL) Earth system model (LMDZ, Hourdin et al., 2006, Dufresne et al., 2013), distinguishes 12 PFTs to represent the global land surface. Moreover, the reclassification in PFTs is done with constant, but qualitative, rules defined across climate zones (Poulter et al., 2011), which can lead to significant uncertainty in the class fractions.

The current ORCHIDEE PFT map is based both on the International Geosphere-Biosphere Program (IGBP) 1-km global land cover map (Belward et al. 1999) reduced by a dominant-type method to 5 km spatial resolution, and on the Olson classification (96 types) (Olson et al. 1983). This spatial resolution is clearly not sufficient for future local-scale studies focused on the environmental impacts of global warming and land use in Siberia and for development perspectives in terms of biogeochemical processes parameterization. Therefore, our objective in this study is to develop a new map at 1 km resolution based on recent land cover products, suitable for earth system modeling which could be further refined if new PFTs are developed.

For that purpose, different remote sensing land cover products are available. They have been developed from multispectral and multitemporal imagery, in order to separate the various ecosystems presenting different spectral properties and seasonal variations. At medium resolution (hundred meters to kilometer), the most popular and most recent products are the GLC 2000 land cover database (Bartholomé and Belward, 2005) based on SPOT 4-VEGETATION instrument, the GlobCover land cover products (Arino et al., 2005, 2012) derived from Envisat/MERIS radiometer and the MODIS land cover datasets (Friedl et al., 2002, 2010), based on Terra and Aqua MODIS instruments.

These products have been compared in previous works and for some of them over Siberia. For example, Jung et al., 2006 developed the SYNMAP product dedicated to earth system modeling, based on the merging of GLC 2000 and MODIS 4.0 products. The final map, which separates 48 classes, is available at 30" scale (~1km). Frey and Smith, 2007 inter-compared AVHRR and MODIS products at 1km scale on West Siberia and highlighted the weaknesses of global LC products in northern wetland environments. Urban et al., 2010, focused on pan-arctic land cover mapping, and combine GlobCover, SYNMAP, MODIS LC and VCF and additional regional products like fire products, to create a new harmonized map separating 4 classes: trees, shrubs, herbaceous and barren areas. Sulla-Meneshe et al., 2011 developed the Northern Eurasia Land Cover (NELC) database from supervised classification of MODIS data which allows to separate 15 land cover classes including land use (urban, agriculture), wetlands and tundra classes. Meanwhile, Pflugmacher et al., 2011 cross-compared GLC 2000, GlobCover and MODIS products as well as Landsat-based reference maps on northern Eurasia. The map legends were converted to a common classification on the basis of dominant Life Form Type (LFT). The results show regional disagreements among products and the difficulties to map shrubs and herbaceous vegetation types. More recently, Shepaschenko et al., 2011 produced a highly detailed land cover/land use dataset for Russia essentially based on GLC 2000 dataset at 1km resolution combined with Vegetation Continuous Fields (VCF) from MODIS, soil and vegetation databases and different inventories and statistics.

All these studies found significant differences between datasets and highlighted strengths and weaknesses of each product but none concluded on the superiority of one compared to the other. Moreover, since for most of these works, the final objective was not PFT mapping, the methodology developed for the cross-comparison and the final mapping could not be used directly for our study. Further, no comparison to date has included the MODIS 5.1 product, which benefits from a reprocessing of the complete MODIS archive, with an up-to-date training database, and an extension of the land cover data to 2011. Therefore, it was necessary to develop a new comparison of the most recent land cover products available on Siberia, to build dedicated aggregation rules for ORCHIDEE PFT mapping and to generate a new PFT map over Siberia at 1km scale.

This paper presents the methodology used to compare medium-resolution remote sensing land cover products for Siberia. The evaluation was performed after aggregating the different land-cover datasets to the same spatial scale and under the same harmonized legend. A comparison of thematic differences was conducted to highlight areas of disagreement and we developed a methodology to generate the PFT distributions. Our results are presented in terms of product comparison and final PFT mapping, with discrepancies explicitly addressed.

2. Methods

We acquired recent land-cover satellite products available at medium spatial resolution (300 to 1000 meters) and focused our comparison on Siberia. The datasets are presented in Table 1 with their specifications in terms of spatial resolution, time of acquisition, geographic projection, and thematic information, including the number of land-cover classes with the respective classification legends in Table2. The global products include the GLC 2000 product (Bartholomé and Belward, 2005) updated on Northern Eurasia by Bartalev et al., (2003), the GlobCover 2005 and 2009 products (Bicheron et al., 2006, Arino et al., 2005, 2012) and the MODIS Land Cover Type Collection 5.0 and 5.1 (Friedl et al., 2002, 2010). The first 4 products have been already compared and evaluated in various regions and at different scales, using ground truth measurements and have shown strengths and weaknesses (See and Fritz, 2006, Jung et al., 2006, Frey and Smith, 2007, Kaptué et al., 2010, 2011, Pflugmacher et al., 2011, Schepaschenko et al., 2011).

Our first goal was to identify the most suitable product for further PFT mapping. To achieve this, we assessed the land-cover classification methodology and land-cover class definition in terms of their capacity to represent the spatial heterogeneity and in terms of the spatial agreement between products. To assess these criteria, we compared various datasets at different temporal and spatial scales, including high-resolution optical images like Landsat-TM products.

Table 1, about here

List of products used and their characteristics

2.1. Land cover products

The GLC 2000 land cover map was developed for different parts of the world with regional experts before applying a generalized legend to create a global land-cover map. In this work, we used the regional product over Northern Eurasia developed by the European Commission's Joint Research Center and the Russian Academy of Science's Center for Forest Ecology and Productivity (Bartalev et al., 2003, Bartholomé and Belward, 2005). The land cover map was produced from daily observations provided by the SPOT4-VEGETATION instrument for the year 2000, at 1/112° ground sampling distance (GSD), corresponding to a ~1km spatial resolution. The automated classification process allows separating 22 land cover types based on local expert opinion following the Land Cover Classification System (LCCS, Di Gregorio and Jensen, 1998) of the Food Agricultural Organization (FAO). The map is available from the JRC Land Resource Management Unit website (<http://bioval.jrc.ec.europa.eu>), in equal area projection (Plate Carrée) with map datum WGS84.

The GlobCover land cover products (GlobCover 2005 and GlobCover 2009) were developed within the framework of European Space Agency (ESA) projects (Bicheron et al., 2004, Arino et al., 2005, 2012). They are both based on Envisat/MERIS data acquired on years 2005 and 2009 respectively and available from the ESA GlobCover Project website (<http://ionia1.esrin.esa.int>). The maps are available in Plate Carrée (WGS84) projection with a 300m spatial resolution (1/360° GSD), under the same class definition as GLC 2000, i.e. LCCS but with a larger number of classes (40) for the regional product available on eastern Eurasia in 2005, whereas GlobCover 2009 product is available only with a global legend which separates 22 classes, fully compatible with the GLC 2000 one. GlobCover uses a fully automated unsupervised classification approach using GLC 2000 as main auxiliary dataset for the class interpretation.

Finally, the MODIS land cover products developed by the Boston University Department of Geography and Center for Remote Sensing (<https://lpdaac.usgs.gov>) are based on NASA MODIS instruments onboard AQUA and TERRA platforms. They are available at annual time step, from 2001 until 2007 for the C5.0 product (Friedl et al., 2002, 2010) with a spatial resolution of 500m (1/240° GSD). The most recent product (C5.1) is available for the 2001-2010 period with the same spatial resolution of 500m. The 2 products are available on an Integerized Sinusoidal Grid (ISG) projection, with a legend based on the IGBP classification

system which separates 17 classes. The new product C5.1 is an update of C5.0 and an extension in time period. The same classification methodology was used but significant errors in the training dataset were adjusted as noted in the User Guide for the MODIS Land Cover Type Product, MCD12Q1 (which is available at http://www.bu.edu/lcsc/files/2012/08/MCD12Q1_user_guide.pdf).

2.2. Auxiliary datasets

Several auxiliary products representing different features of the land surface were used to assist in the evaluation of the global products (see Table 1). These products first helped us in the interpretation of the different legends and later assisted in the merging process, which permitted us to build the harmonized legend. Among them, two land cover maps based on aerial photointerpretation and ground truth data have been used to better understand the class significance and evaluate product accuracy and spatial variability representation. For these 2 products, the digital database was not accessible and only graphical maps have been used. The first one is the Circumpolar Arctic Vegetation Map (CAVM, Walker et al., 2005) which was developed within the National Science Foundation Arctic Transitions in the Land-Atmosphere System (ATLAS) project, and is presently the most precise mapping of the Arctic tundra. It is available at 1:7,500,000 scale at <http://www.geobotany.uaf.edu/cavm>, in Lambert azimuthal projection, and separates 18 classes describing very precisely the various tundra ecosystems. This dataset is based on photo-interpretation by vegetation experts of nine Arctic regions which allowed delineating the various biomes onto an NOAA-AVHRR image database.

The second one is the Yakutsk region land cover mapping provided by A. Fedorov (personal communication), which was derived from Landsat images acquired in 2002 combined with ground truth data. In this map, 12 land cover classes were separated including 6 different types of forest like larch and birch in different states and 3 types of grasslands (Alas, wet and dry valley meadows). These 2 maps have mostly been used in the following to evaluate the ability of the various land cover products to separate the shrubs / herbaceous classes and the broadleaf / needleleaf forests.

Lastly, the MODIS Global Vegetation Continuous Fields (VCF, Hansen et al., 2003, 2006) and the forest canopy height map proposed by Simard et al., 2011, complemented all these datasets. The VCF product

derived from MODIS sensors is provided at 500m spatial resolution on an annual basis (2000-2010) and at global scales. The VCF is proportional estimates of vegetative cover types: woody vegetation, herbaceous vegetation, and bare ground. The Collection 4 (version 3) was downloaded at <http://glcf.umiacs.umd.edu/data/vcf> , where it is available in the same projection as the land cover product. Finally, the forest height product based on 2005 data from the Geoscience Laser Altimeter System (GLAS) onboard ICESat (Ice Cloud and Land Elevation satellite), is available globally at 1km spatial resolution and provides an estimation of the canopy height. These two last products provided an independent mapping of the forested areas and were mostly used in the PFT maps generation. The data were obtained through the website <http://lidarradar.jpl.nasa.gov> in GeoTiff format.

2.3. Harmonized legend approach

Because these land cover products did not have the same spatial resolution and more importantly, did not use the same classification system, a harmonization procedure was developed. As already discussed by all the works dedicated to land cover map cross comparison (to cite but a few: See and Fritz, 2006; Frey and Smith, 2007; Urban et al., 2010; Sulla-Menashe et al., 2011; Pflugmacher et al., 2011; Kaptué et al., 2011), the classification method, the original data, the number of thematic classes chosen, etc..., can highly bias the classification results and the overall regional biogeographic characteristics.

Table 2 about here

(Aligning the legends of global maps: dominant life form types (LFT) and corresponding land cover classes from GLC-2000, GLOBCOVER & MODIS IGBP)

For example, GLC 2000 and GlobCover legends give more weight to the dominant tree species than to the density character, compared to MODIS. This is probably the result of the classification methodology applied for the MODIS product, which is based on the combined use of surface reflectance and land-surface temperature (LST), contrary to the other products, which use only surface reflectance. The addition of LST, which is known to be highly sensitive to vegetation fraction, could have increased the weight of the tree

coverage in the class separation. Therefore, the LCCS legend used for GLC 2000 and GlobCover defines forest as greater than 15% tree cover with trees defined as woody plants larger than 5m, whereas IGBP used in the MODIS product, defines forest as greater than 60% tree cover with trees defined as woody plants larger than 2m. Two other IGBP classes of 8 and 9 (woody savannas/savannas) are then used to represent more open canopies with the same height thresholds but different cover thresholds down to 10%. In the same way, for shrublands, LCCS distinguishes between evergreen and deciduous species, whereas IGBP considers open and closed types. Further, for barren lands, IGBP merges bare and sparsely vegetated soils, when LCCS separates sparse herbaceous, sparse shrubs and bare areas. Given all these features, a comparison work could not be performed before having converted all the study products in a common legend. For our final purpose of water and carbon cycle modeling, this common classification requires first, to be based on PFT features and secondly, to discriminate trees, shrubs, water and barren but also leaf type and senescence. This choice leads us to merge the IGBP and LCCS classes under the 16 classes listed in Table 3, which are in close agreement with the GlobCover 2009 legend. The merging rules and the allocation of the ambiguous classes have been driven by the comparison of the spatial distribution of the land cover classes and the help of the auxiliary products, especially the high resolution maps (CAVM map and the various Landsat images acquired on different subregions of Eurasia).

Table 3 about here:

Harmonized legend used and correspondence with original products classes

Afterwards, in order to allow the comparison, all the data have been re-projected to the WGS84 Plate Carrée projection with square pixel size ($1/112^\circ$ about 1km scale), using a majority class criterion, since GLC 2000 and GlobCover are already available in these projection and grid.

3. Results

The various products have been compared over Siberia, with a focus on central and south-west Siberia (over Yakutia and around Omsk respectively, where high resolution data were available). The coordinates of these 2 domains are 55°N - 75°N / 104°E - 163°E and 50°N - 58°N / 56°E - 96.5°E , respectively. These 2

regions were chosen because they cover almost all the variety of Siberian ecosystems. The first region, Yakutia (Sakha Federal Republic of Russia, capital Yakutsk) covers a large area of about 3M square kilometers, with 40% above the Arctic Circle. The region is one of the coldest continental regions in the world (outside Antarctica) with large annual temperature amplitude varying between -60°C to +40°C. It is all covered by permafrost and mainly drained by the Lena River and its tributaries. The vegetation is driven by these extreme climate conditions, which limits the extent of Arctic tundra, composed of lichens and mosses in the north, and the taiga forest mostly composed of deciduous trees (especially larch) in the south. The other region studied is situated in the South-West part of Siberia. It represents an area of about 2M square kilometers and is part of the Irtysh river catchment. The vegetation is mostly composed of croplands (wheat, barley, potatoes ...) and deciduous forests with the predominant species being larch in the north taiga and birch and aspen in the south.

3.1. Comparison on Central Siberia

Figure 1 presents the land cover maps extracted from the 5 global products (GLC 2000, GlobCover 2005 and 2009, MODIS 5.0 and 5.1) at 1 km scale under the harmonized legend discussed previously. The 5 products were compared considering their respective representative time period. Thus, GlobCover 2005 was compared with MODIS 2005 dataset, GlobCover 2009 with MODIS 2009 product and GLC 2000 with MODIS 2001.

Figure 1 about here (Land cover maps with the harmonized legend on Yakutia)

The maps clearly show the latitudinal distribution of tree cover, with forested areas between the mountains of Verkhoyansk and Chersky on the East side and Stanovoi in the South, sparse vegetation in the northern latitudes and bare soils in the mountainous areas. In this large region, the main land cover classes are deciduous needleleaf forest (mostly larch) covering the middle latitudes, and shrubs often mixed with forests and sparse vegetation. Although present in the 5 products, the fraction and spatial distribution of forest differ significantly among them. Table 4 presents the fraction of each land cover class for each study product. The fractions were calculated excluding the water pixels in order to avoid the Arctic Ocean pixels

which could have biased the statistics. In the Table, values greater than 10% have been highlighted (i.e., excluding vegetation types not well-represented in this region).

Table 4: Fraction of each land cover class in each product for Yakutian region

The main features are a larger representation of the sparse vegetation class in MODIS (50% and 64% for 5.0 and 5.1 products respectively) compared to 8 and 21% in GLC 2000 and GlobCover 2009, to the detriment of deciduous needleleaf forest (28% in MODIS 5.1 compared to 58% in GlobCover 2009). This disagreement was already pointed out by Frey and Smith (2007) when they compared MODIS product to land cover field-based observations. It can be noted also that the shrub class is represented in the north of the region only in the GLC 2000 product and that regularly flooded areas are more represented in GlobCover and MODIS products compared to GLC 2000, especially in the Lena river delta. The spatial agreements are quantified in Table 5 for the main classes present in this region (i.e., deciduous needleleaf, mixed forests, shrubs, sparse vegetation, herbaceous and bare soils). The statistics were calculated by comparing each product to the GlobCover 2005 one (values above 0.5 are shaded in the table and statistics were not calculated for the classes presenting amounts less than 0.5% like shrubs). As previously, the total agreement percentages do not account for water pixels which could have biased the accuracy assessment. The results confirm the spatial comparisons: the best agreements are obtained for the 2 GlobCover products as expected, even though some classes like mixed forests or herbaceous present some discrepancies, probably linked to the low number of pixels involved since these classes represent about 1% of the GlobCover 2005 map. The comparison with GLC 2000 shows that the lower scores are obtained for the same ambiguous classes which probably present a larger heterogeneity and could suffer from the lower spatial resolution. The evaluation with MODIS maps displays worse statistics especially for shrubs, mixed forests and herbaceous and the total agreement values are all lower than 0.5 except when two products of the same family are compared. A qualitative comparison was further performed with the CAVM product (Walker et al., 2005). The comparison of photo-interpretations presenting continuous fields with pixel based classifications has proven difficult, but on specific regions like in the Lena river delta, the different types of tundra appear better separated in the GLC 2000 and GlobCover 2005 products than in the other ones (not shown in this paper).

Table 5: Agreement percentages on Yakutia for the main classes and comparison of Globcover 2005 with Globcover 2009, GLC 2000, MODIS 5.0 and MODIS 5.1

3.2. Comparison at finer scale on Yakutsk surroundings

In order to check the accuracy of the various products and the impact of the increased resolution of 300m on the spatial representation of the land variability, we focused our comparisons around Yakutsk region taking advantage of the high resolution map provided by A. Fedorov's team (personal communication). This map is shown in Figure 2 (bottom image). It shows clearly a precise delineation of the evergreen needleleaf forest (identified as "Pine" in light green color) and the mixed forest plots (in light blue ("Pine larch") and in pink ("Birch Larch")) within the dominant deciduous needleleaf ecosystem ("Larch" in middle green), particularly on the right bank of the Lena river. On Figure 2, only the corresponding images for the GLC2000 and GlobCover products exhibit this same spatial variability with evergreen needleleaf forests along the river banks in dark green and mixed forests in middle green whereas, the MODIS maps only show the deciduous needleleaf forest in light green. However, the lower spatial resolution of GLC 2000 does not allow representing correctly the river bed and the flooded areas, contrary to the GlobCover 2005 product in which the herbaceous regularly flooded class along the riverbed is well characterized. Therefore, in that region, it seems that GlobCover 2005 better captures the main features of the land coverage.

Figure 2 about here

(Land cover maps with the harmonized legend for Yakutsk region compared to Fedorov's map)

3.3. Comparison in south-west Siberia

The second region where we focused our cross comparison is the south-west part of Siberia where we are interested in analyzing the evolution of agriculture in future works. This region is mostly covered by croplands, deciduous broadleaf and needleleaf forests, herbaceous and sparse vegetation. The grid cell

fractions range between 0.3 and 0.43 for croplands, and are generally lower than 0.2 for the other cover types (Table 6).

Table 6: Fraction of each land cover class in each product for South-West Siberia

Figure 3: Land cover maps with the harmonized legend for South-West Siberia

Table 7: Agreement percentages for South-West Siberia for the main classes and comparison of Globcover 2005 with Globcover 2009, GLC 2000, MODIS 5.0 and MODIS 5.1

The main difference among the LC products stems in the representation of the southern part of the region, covered by sparse vegetation and herbaceous in GlobCover and GLC 2000 maps, and by regularly flooded lands in MODIS. This area at the limit of Kazakhstan is drained by the Irytsh River and presents irrigated croplands which could be identified wrongly as flooded areas. The main cropland area in purple in Figure 3 is well delimited between the forested areas in the north and the sparse vegetated lands southern. We also noted differences in the eastern part of the region which is classified as croplands by MODIS and as sparse vegetation in the other products. This disagreement was analyzed more deeply, by looking at this region with Landsat images. The views show undoubtedly that this region is covered by agricultural fields presenting an unambiguous spatial structure. The classification discrepancies between crop and sparse vegetation highlight the difficulties in separating such ecosystems if several dates per year are not used to assess the intra-seasonal variability, and if a pixel based classification technique is used. An object-based classification methodology should have been better performing in that case in absence of higher spatial resolution imagery. Anyway, MODIS classification appears here, more accurate for crop mapping.

The spatial agreement statistics are presented for the main land cover classes in Table 7. The classes which appear in best agreement are the crops and the mixed forest (except for GLC 2000) with values larger than 0.5. The lowest values are obtained for sparse vegetation which can be mixed with crops or herbaceous ecosystems. Moreover, the deciduous needleleaf and broadleaf forests appear to be better separated in the GlobCover products compared to MODIS and GLC 2000, where all the forested areas are

grouped in the mixed forest class. Finally, the overall agreement percentages are very low, with values never exceeding 0.4 similar to what has been previously shown on Yakutia (except when products of the same family are compared).

3.4. Forest mapping accuracy

Our motivation being the development of PFT maps for land surface modeling, the characterization of biomes and the separation of forested areas from shrublands are particularly important. Indeed, the vertical structure of forests implies different ground shading, aerodynamic and roughness properties, and consequently significant impacts on surface fluxes. Therefore, a product of forest canopy height like the one proposed by Simard et al., 2011, appears interesting for the interpretation of the land cover products legend as well as for their accuracy assessment. This recent product, based on LIDAR measurements, provides at a global scale, the estimation of canopy height at 1km resolution with an error evaluated against ground truth data, less than 6m. For the comparison with land cover products, the forest classes were grouped together. Since the LIDAR product is based on 2005 data, and since the two GlobCover products are very similar, only GlobCover 2005 map was included in the comparison. Figure 4 presents the comparison of the current forest height product with GLC 2000, GlobCover 2005 and the two MODIS products extracted for the year 2005. Qualitatively, the extension of forested areas appears better represented in GlobCover 2005 if a threshold of 10m height is imposed to delineate trees and shrubs. The agreement with the forest class was calculated for the 4 land cover products (MODIS5.0, MODIS5.1, GLC 2000 and GlobCover 2005). The spatial agreements obtained (43.5%, 37.4%, 57.9% and 76.1%) for the 4 products respectively, show clearly that GlobCover 2005 better captures the degree of woodiness at the land surface, which is an essential parameter for vegetation characterization in carbon and water cycle modeling.

3.5. Discussion

The results of the cross comparison of the 5 land cover products studied, on two different regions of Siberia, show differences and similarities which can be explained either by the lack of resolution (for GLC 2000) or by the methodologies used to assess the class separation and interpretation. The agreement among the maps is highest in the zones which present more homogenous landscapes (for example inside

the taiga region) and lower in the transition zones or in sparse ecosystems for which the class definition is determinant. The contribution of higher resolution products is therefore a significant improvement for discriminating vegetation types and for better mapping such regions. For our modeling purposes, since the main objective is to identify the type of ecosystem whatever its density (which will be anyway, provided by the LAI variable used as forcing or prognostically computed by the big-leaf type DGVM), the definition provided by GlobCover or GLC appears more valuable. Furthermore, this class definition allows delineating forested areas more precisely, as was demonstrated in the comparison with the recent forest height product. In addition, GlobCover 2005 provides a more precise legend (compared to GlobCover 2009, even if this level of detail is not ensured globally) and an increased spatial resolution (compared to GLC 2000).

Figure 4 about here (Forest mapping comparison)

For all these reasons, the GlobCover 2005 product was chosen as a basis for the PFT mapping, keeping in mind its class definition, especially the forest classes which can include pixels with spatial coverage as low as 15%. This definition more suitable for land cover type identification will require the merging with other indices to account for vegetation density. Otherwise, it could lead to a likely over representation of forests in transition zones with tundra in the northern latitudes and with herbaceous cover types in the south.

4. PFT mapping

Given the GlobCover 2005 land cover map, our next challenge is to define merging rules to build a PFT map for the ORCHIDEE dynamic global vegetation model (DGVM). For that purpose, we have followed the approach of Poulter et al., 2011 and associated to each GlobCover 2005 land cover class, the corresponding classification in the ORCHIDEE DGVM. In this work, we focused only on the PFTs present in Siberia. Given the previous studies and drawbacks highlighted in the GlobCover 2005 product, the reclassification rules proposed by Poulter et al., 2011, have been slightly modified and adjusted to boreal ecosystems, as described in section 4.2.

4.1. ORCHIDEE model

ORCHIDEE land surface model is a mechanistic dynamic global vegetation model (Krinner et al. 2005), that is part of the IPSL Earth system model (Friedlingstein et al., 2006). It calculates the energy, momentum and hydrological budget of vegetation and soil and all the carbon and nitrogen cycle in the different soil and vegetation pools. Photosynthesis, phenology, allocation of carbon and nitrogen into the different organs, plant growth and mortality, and decomposition of litter and soil organic matter, are derived from primitive equations that depend on vegetation characteristics. ORCHIDEE is built on the concept of PFTs to describe vegetation distributions. Species with similar characteristics are re-grouped together and the model distinguishes 12 PFTs (tropical evergreen and deciduous forests, temperate broadleaf evergreen and deciduous forests, temperate needleleaf forest, boreal needleleaf evergreen and deciduous forests, boreal broadleaf deciduous forest, natural C3 and C4 grassland, and C3 and C4 crops) plus bare soil. In its standard version, the PFTs are defined from two databases, the AVHRR IGBP 1-km global land cover map (Belward et al., 1999) and Olson et al.(1983) biome classification including 96 land types (Vérant et al., 2004). The final map prescribes the fraction of each vegetation type over a resolution cell of 5 km. Therefore, different PFTs can coexist in every grid element, and their fraction can vary when the dynamic vegetation sub-module is activated. Figure 5 presents the standard PFT maps used in ORCHIDEE.

Figure 5 about here (Standard ORCHIDEE PFT maps)

In Siberia, 9 PFTs are present in the standard land cover map. They include four types of forests (temperate and boreal needleleaf evergreen, needleleaf summergreen, broadleaf summergreen forests), C3 grass, C3 crops, unlikely (but very few) C4 crops and bare soils.

4.2. ORCHIDEE PFTs

Table 8 presents the merging rules that have been defined to reclassify the GlobCover 2005 classes present in Siberia, into the ORCHIDEE PFTs. The original crosswalking table from Poulter et al., 2011 was slightly modified to better map the boreal PFTs, based on the ancillary satellite data and considering the LCCS legend for regional land cover types. In particular, the fractions of forested PFTs in the open forests

classes were decreased to account for the lower density of forests in boreal regions, as well as the fractions of bare soil in the sparse vegetation classes to account for mosses in tundra environments. Moreover, because of the absence of the boreal broadleaf evergreen class in the ORCHIDEE PFT classification, this class has been equally distributed between the broadleaf summergreen and the needleleaf evergreen PFT classes. For the same reasons, the lichens were merged with C3 grasses and the shrublands were spread among the C3 grass, the forests and bare soil classes. Further, the percentages of forests, bare soils and grasslands (only C3 in boreal zones), were adjusted with the support of the MODIS VCF products. These data, indeed, permit to assess the land surface heterogeneity and the amount of vegetation inside the pixels. For example, Montesano et al. (2009) in their evaluation of the VCF product on the circumpolar taiga-tundra transition zone, showed the contribution of these continuous fields to capture the forest cover variability and the spatial heterogeneity, especially in land cover transition zones. Therefore, the VCF data for the year 2005 have been upscaled at 1km scale and the fractions of trees, grasslands and bare soil have been extracted and averaged for each Globcover 2005 class. The results permitted a better understanding of the LCCS legend and the PFT reclassification was then performed according to the new merging rules described in Table 8.

Table 8: Merging rules from GlobCover PFTs to ORCHIDEE PFTs

Figure 6: New ORCHIDEE PFT maps

The new PFT maps have been generated from the GlobCover 2005 dataset, keeping the benefit of the high resolution of 300m. The data were aggregated at 1km scale in PFT fractions and the results are presented in Figure 6. The results show the main features characterizing Siberia, i.e., the predominance of needleleaf summergreen forests in the center, broadleaf summergreen and then croplands southern, grasslands and larger fractions of bare soil in the northern latitudes. Compared to the previous maps (see Figure 7 for the differences mapping), the temperate broadleaf PFTs and sparse C4 crops have disappeared, and generally, the fractions are less contrasted. The amount of bare soil and grasslands is larger (up to 15%) in central Siberia, better representing the sparse feature of Siberian forests, and the broadleaf summergreen forests abundance decreased in the northern part of the region, which seems to be more realistic. The

boreal needleleaf summergreen forest covers now a larger area north and south of the previous limited location, where the abundance decreased from 100 to 60%. The water surfaces are also more and better represented.

Figure 7 about here: Difference (New – Standard) PFT maps

The agreement between the original Olson-based PFT map and the new GlobCover-based PFT map was quantified with the commonly used Euclidian distance between the PFT classes (Legendre et al., 2005, Poulter et al., 2011), calculated for each 1km pixel. Eq. 1 presents the expression of the dissimilarity index for a grid cell c , calculated between the 2 classifications (New and Standard) composed of 14 classes (12 PFT + water and ice classes) and their corresponding fractional abundance P in the 2 classifications.

$$D_c = \left[\sum_i^{14} (P_{new_{i,c}} - P_{stan_{i,c}})^2 \right]^{0.5} \quad (1)$$

This index, which is 0 for full agreement and $\sqrt{2}$ for full disagreement is displayed in Figure 8. The agreement is best in the northern latitudes and worse in the center of Siberia where the fractions of grasslands and forested PFTs have been the more modified.

Figure 8 about here: PFT dissimilarity index

5. Discussion and Conclusion

Land cover mapping is crucial for many environmental studies and the re-gathering in PFT classes is necessary for the specific purposes of land surface modeling. In this study focused on Siberia, we compared five medium resolution land cover maps derived from remote sensing and highlighted some discrepancies mostly linked to the legend definition adopted. The strengths and weaknesses of each product were shown and the results led us to choose the GlobCover 2005 product because of its highest spatial resolution and more detailed legend. Therefore, a new PFT map at 1km scale over Siberia has been generated for the ORCHIDEE DGVM. This map shows large differences compared to the standard maps in

the differentiation of broadleaf and needleleaf forests and in the representation of the landscape heterogeneity. The fractions of the various ecosystems are smoothed and seem to better represent the vegetation diversity, thanks to the use of higher resolution datasets for the PFTs mapping.

These differences should significantly impact the DGVM simulations. Indeed, PFTs fractions are used to define the vegetation characteristics in terms of photosynthesis capacity, phenology, roughness, etc. All these properties are determinant for the calculation of the water and carbon fluxes, especially the evapotranspiration and the GPP fluxes. Consequently, such modifications should impact the biosphere-atmosphere exchanges and will be analyzed in further works.

This study also showed the difficulties to link vegetation classes to a limited number of PFT, constrained by global modeling and time computing issues. The absence of a shrub PFT and the solution to distribute the shrubs classes among grasslands, bare soils and forests is not satisfactory. Such vegetation types have so different properties that it appears difficult to well represent the energy and mass transfers with an aggregation of such variability. In the same way, mosses and lichen-dominant ecosystems are not represented in the final PFT map and are assimilated to bare soils, the same for regularly flooded areas and peatlands that have been spread between the grasslands and water classes, which in terms of carbon cycle could lead to significant errors. Therefore, the development of new PFT classes in ORCHIDEE, to better represent these specific northern ecosystems appear to be a priority if one wants to correctly represent boreal ecosystems and their future evolution.

6. Acknowledgements

The authors are grateful to the French ANR CLASSIQUE and EU-FP7 PAGE21 projects for funding this study (internship of J. Lescure). They acknowledge also the MODIS team, Mark Friedl and Damien Sulla-Menasche, the JPL (M. Simard), the Melnikov Permafrost Institute SB RAS (A. Fedorov), ESA and JRC to have made available their datasets for this work. The anonymous reviewers are also acknowledged for their constructive comments which helped to improve the paper.

7. References

- Arino, O., Trebossen, H., Achard, F., Leroy, M., Brockman, C., Defourny, P., 2005: The GLOBCOVER Initiative, Proceedings of the MERIS (A) ATSR Workshop 2005 (ESA SP-597), 26-30 September 2005 ESRIN, Frascati, Italy. Editor: H. Lacoste. Published on CDROM, p.36.1, 2005ESASP.597E..36A
- Arino, O., Ramos Perez, J.J., Kalogirou, V., Bontemps, S., Defourny, P., Van Bogaert, E., 2012: Global Land Cover Map for 2009 (GlobCover 2009). © *European Space Agency (ESA) & Université catholique de Louvain (UCL)*, doi:10.1594/PANGAEA.787668
- Bartalev, S. a., Belward, a. S., Erchov, D. V., & Isaev, a. S., 2003: A new SPOT4-VEGETATION derived land cover map of Northern Eurasia. *International Journal of Remote Sensing*, 24(9), 1977–1982. doi:10.1080/0143116031000066297
- Bartholomé, E. Belward, AS., 2005: GLC 2000: a new approach to global land cover mapping from Earth observation data. *International Journal of Remote Sensing*, 26(9), 1959–1977. doi: 10.1080/01431160412331291297.
- Belward, A., J. Estes, and K. Kline, 1999: The IGBP-DIS Global 1-km Land-Cover Data Set DISCover: A project overview. *Photogramm.Eng. Remote Sens.*, **9**, 1013–1020.
- Bicheron, P., Leroy, M., Brockmann, C., Miras, B., Huc, M., Vancutsem, C., Arino, O., et al., 2006: GLOBCOVER : a 300 m global land cover product for 2005 using ENVISAT MERIS time series, Proceedings of the [28th EARSeL Symposium: Remote Sensing for a Changing Europe](#), December 2004), 1–5.
- Chapin III F.S., M. Sturm, M. C. Serreze, J. P. McFadden, J. R. Key, A. H. Lloyd, A. D. McGuire, T. S. Rupp, A. H. Lynch, J. P. Schimel, J. Beringer, W. L. Chapman, H. E. Epstein, E. S. Euskirchen, L. D. Hinzman, G. Jia, C.-L. Ping, K. D. Tape, C. D. C. Thompson, D. A. Walker, and J. M. Welker, 2005: Role of Land-Surface Changes in Arctic Summer Warming, *Science* 28 October 2005: 310 (5748), 657-660. Published online 22 September 2005 [DOI:10.1126/science.1117368].
- Delbart, N., L. Kergoat, T. Le Toan, J. L'Hermitte and G. Picard, 2005: Determination of phenological dates in boreal regions using Normalised Difference Water Index. *Remote Sensing of Environment*, 97, 26-38.

- Delbart, N. and Picard, G., 2007: Modeling the date of leaf appearance in low-arctic tundra, *Global Change Biology*, vol. 13, 2551-2562.
- Di Gregorio, A., Jansen, L. J. M., & Resources, S., 1998: Land Cover Classification System (LCCS): Classification Concepts and User Manual.
- Dufresne, J.-L., Foujols, M.-A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benshila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J.-P., Ethé, C., Fairhead, L., Fichefet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J.-Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A., Lefebvre, M.-P., Lefevre, F., Levy, C., Li, Z. X., Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J., Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., 2013: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, *Clim. Dynam.*, DOI 10.1007/s00382-012-1636-1
- Euskirchen, E. S.; McGuire, A. D.; Chapin III, F. S.; Yi, S.; Thompson, C. C., 2009, Changes in vegetation in northern Alaska under scenarios of climate change, 2003-2100: Implications for climate feedbacks, *Ecological Applications*, 19: 1022 – 1043.
- Forbes, B.C., Macias Fauria, M. & Zetterberg, P., 2010: [Russian Arctic warming and 'greening' are closely tracked by tundra shrub willows](#). *Global Change Biology*, 16, 1542–1554.
- Frey K.E. and L.C.Smith, 2007: How well do we know northern land cover? Comparison of four global vegetation and wetland products with a new ground-truth database for West Siberia, *Global Biogeochemical Cycles*, 21, GB1016.
- Friedl, M., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., & Huang, X., 2010: MODIS Collection 5 global land cover: Algorithm refinements and characterization of new datasets, *Remote Sensing of Environment*, 114(1), 168–182. doi:10.1016/j.rse.2009.08.016.
- Friedl, M.A., McIver, D.K., Hodges, J.C.F., Zhang, X.Y., Muchoney, D., Strahler, A.H., Woodcock, C.E., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F., Schaaf, C., 2002: [Global land cover mapping from MODIS: algorithms and early results](#). *Remote Sensing of Environment*, 83(2-Jan), 287-302.

Friedlingstein, P., P. Cox, R. Betts, L. Bopp, W. Von Bloh, V. Brovkin, P. Cadule, S. Doney, M. Eby, I. Fung, G. Bala, J. John, C. Jones, F. Joos, T. Kato, M. Kawamiya, W. Knorr, K. Lindsay, H. D. Matthews, T. Raddatz, P. Rayner, C. Reick, E. Roeckner, K.-G. Schnitzler, R. Schnur, K. Strassmann, A. J. Weaver, C. Yoshikawa, and N. Zengq, 2006: Climate–Carbon Cycle Feedback Analysis: Results from the C4MIP Model Intercomparison. *Journal of Climate*, 19, 3337-3353.

Hansen, M., R. DeFries, J.R. Townshend, M. Carroll, C. Dimiceli, and R. Sohlberg, 2003: Global Percent Tree Cover at a Spatial Resolution of 500 Meters: First Results of the MODIS Vegetation Continuous Fields Algorithm. *Earth Interactions*, 7(10), 1–15. doi:10.1175/1087-3562(2003)007<0001:GPTCAA>2.0.CO;2.

Hansen, M., R. DeFries, J.R. Townshend, M. Carroll, C. Dimiceli, and R. Sohlberg (2006). Vegetation Continuous Fields MOD44B, 2001 Percent Tree Cover, Collection 4, University of Maryland, College Park, Maryland, 2001.

Hourdin, F., I. Musat, S. Bony, P. Braconnot, F. Codron, J.-L. Dufresne, L. Fairhead, M.-A. Filiberti, P. Friedlingstein, J.-Y. Grandpeix, G. Krinner, P. LeVan, Z.-X. Li et F. Lott, 2006, The LMDZ4 general circulation model: climate performance and sensitivity to parametrized physics with emphasis on tropical convection, *Climate Dynamics*, 27: 787-813.

Hüttich, C., Herold, M., Schmullius, C., Egorov, V. and Bartalev, S. A., 2007: Indicators of Land Cover Change in Northern Eurasia from SPOT-VEGETATION Time Series Analysis 1998-2005, *International Journal of Remote Sensing* 28 (18) , 4199 - 4206.

Jung, M., K. Henkel, M. Herold and G. Churkina, 2006: Exploiting synergies of global land cover products for carbon cycle modeling, *Remote Sensing of Environment*, 101, 534-553, 2006.

Kaptue Tchente, A. T., Roujean, J.-L., and Faroux, S., 2010: ECOCLIMAP-II: An ecosystem classification and land surface parameters database of Western Africa at 1km resolution for the African Monsoon Multidisciplinary Analysis (AMMA) project. *Remote Sensing of Environment*, 114(5), 961–976. doi:10.1016/j.rse.2009.12.008.

Kaptue Tchente, A. T., Roujean, J.-L., and De Jong S.M., 2011: Comparison and relative assessment of the GLC 2000, GLOBCOVER, MODIS and ECOCLIMAP land cover datasets at the African continental scale, *Int. J. of Applied Earth Observation and Geoinformation*, 13, 207-219.

Krinner, G., Viogy, N., De Noblet-Ducouré, N., 2005 : A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system. *Global Biogeochemical Cycles*, 19(1), doi: 10.1029/2003GB002199.

Legendre, P., Borcard, D., and Pers-Neto, P.R., 2005: Analyzing beta diversity: Partitioning the spatial variation of community composition data, *Eco. Monogr.*, 75, 435-450, 2005.

Lenton, T. M., Held, H., Kriegler, E., Hall, J. W., Lucht, W., Rahmstorf, S., and Schellnhuber, H. J., 2008: Tipping elements in the Earth's climate system, *Proceedings of the National Academy of Science*, 105, 1786-1793, 2008.

Lorant M.M. and S.J. Goetz., 2012: Shrub expansion and climate feedbacks in Arctic tundra. *Environmental Research Letters*, 7 011005 doi:10.1088/1748-9326/7/1/011005.

Lucht, W., Prentice, I. C., Myneni, R. B., Sitch, S., Friedlingstein, P., Cramer, W., Bousquet, P., Buermann, W., and Smith, B., 2002: Climatic control of the high-latitude vegetation greening trend and Pinatubo effect, *Science*, 296, 1687-1689, 2002.

Montesano, P.M., R. Nelson, G. Sun, H. Margolis, A. Kerber, K.J. Ranson, 2009: MODIS tree cover validation for the circumpolar taiga–tundra transition zone, *Remote Sensing of Environment*, 113, 10, 2130-2141, ISSN 0034-4257, <http://dx.doi.org/10.1016/j.rse.2009.05.021>.

Myneni, R.B., Dong, J., Tucker, J.M., Kaufmann, R.K., Kauppi, P.E., Liski, J., Zhou, L., Alexeyev, V. and Hughes, M.K., 2001: A large carbon sink in the woody biomass of Northern forests. *Proceedings of the National Academy of Science*, 98(26): 14784-14789.

Olson, J., J. Watts, and L. Allison, 1983: Carbon in live vegetation of major world ecosystems. *Technical Report W-7405-ENG-26*, Oak Ridge National Laboratory, Oak Ridge, TN, 152 pp.

Pflugmacher, D., Krankina, O. N., Cohen, W. B., Friedl, M. a., Sulla-Menashe, D., Kennedy, R. E., Nelson, T.V. Loboda, T. Kuemmerle, E. Dyukarev, V. Elsakov, V.I. Kharuk, 2011: Comparison and assessment of coarse resolution land cover maps for Northern Eurasia, *Remote Sensing of Environment*, 115(12), 3539–3553. doi:10.1016/j.rse.2011.08.016.

- Poulter, B., P. Ciais, E. Hodson, H. Lischke, F. Maignan, S. Plummer and N.E. Zimmermann, 2011: Plant functional type mapping for earth system models, *Geosci. Model Dev.*, 4, 993-1010, 2011.
- Quaife, T., S. Quegan, M. Disney, P. Lewis, M. Lomas, and F. I. Woodward, 2008: Impact of land cover uncertainties on estimates of biospheric carbon fluxes, *Global Biogeochem. Cycles*, 22,GB4016, doi:10.1029/2007GB003097.
- Simard, M., N. Pinto, J.B. Fisher and A. Baccini, 2011: Mapping forest canopy height globally with spaceborne lidar, *J. Geophys. Res.*, 116, G04021, doi:10.1029/2011JG001708.
- See L.M. and S. Fritz, 2006: A method to compare and improve land cover datasets: Application to the GLC-2000 and MODIS land cover products, *IEEE Transactions on Geoscience and Remote Sensing*, 44(7):1740- 1746 [2006].
- Shepashenko, D., I. McCallum, A. Shvidenko, S. Fritz, F. Kraxner and M. Obersteiner, 2011: A new hybrid land cover dataset for Russia: a methodology for integrating statistics, remote sensing and in situ information, *Journal of Land Use Science*, 6, 4, 245-259.
- Sulla-Menashe, D., M.A. Friedl, O.N. Krankina, A. Baccini, C.E. Woodcock, A. Sibley, G. Sun, V. Kharuk, and V. Elsakov, 2011: Hierarchical mapping of northern Eurasia using MODIS data, *Remote Sensing of Environment*, 115: 392-403.
- Tape, K., M. Sturm and C. Racine, 2006: The evidence for shrub expansion in Northern Alaska and the Pan-Arctic, *Global Change Biology*, [12, 4](#), 686–702.
- Urban, M., S. Hese, M. Herold, S. Pöcking and C. Schmullius, 2010: Pan Arctic Land cover Mapping and Fire Assessment for the ESA Data User Element Permafrost, *PFG Photogrammetrie, Fernerkundung, Geoinformation*, 4, 283-293.
- Verant, S., Laval, K., Polcher, J., and M. De Castro, 2004: Sensitivity of the continental hydrological cycle to the spatial resolution over the Iberian Peninsula, *Journal of Hydrometeorology*, 5, 267-285, 2004.
- Walker, D., Raynolds, M. K., Daniëls, F. J., Einarsson, E., Elvebakk, A., Gould, W., Katenin, A. E., et al., 2005: The Circumpolar Arctic vegetation map. *Journal of Vegetation Science*, 16(3), 267. doi:10.1658/1100-9233(2005)016[0267:TCAVM]2.0.CO;2.

8. Figure captions

Figure 1: Comparison of 5 Land Cover products on Yakutia: GLC2000, GlobCover 2005 and 2009, MODIS 5.0 and 5.1 for 2001, 2005 and 2009, are aggregated at 1km scale to a common legend.

Figure 2: Comparison of 5 Land Cover products on Yakutsk region: GLC2000, GlobCover 2005, MODIS 5.0 and 5.1 for 2001, aggregated at 1km scale to a common legend with the land cover map provided by A. Fedorov (Melnikov Permafrost Institute, Russia).

Figure 3: Comparison of 5 Land Cover products on south-west Siberia: GLC2000, GlobCover 2005 and 2009, MODIS 5.0 and 5.1 for 2001, 2005 and 2009, are aggregated at 1km scale to a common legend.

Figure 4: Comparison of 4 Land Cover products on Yakutia: GLC2000, GlobCover 2005, MODIS 5.0 and 5.1 for 2001, with the forest canopy height product provided by Simard et al., 2011. The forests classes were grouped together at 1km scale.

Figure 5: Standard ORCHIDEE PFT maps. The respective fractions of the following 9 classes: Bare soil (PFT1), Temperate Needleleaf Evergreen (PFT4), Temperate Broadleaf Evergreen (PFT5), Temperate Broadleaf Summergreen (PFT6), Boreal Needleleaf Summergreen (PFT9), C3 Grass (PFT10), C3 Agriculture (PFT12), C4 Agriculture (PFT13), are represented in color scale, from blue (0%) to red (100%).

Figure 6: New ORCHIDEE PFT maps. The respective fractions of the following 8 classes: Bare soil (PFT1), Boreal Needleleaf Evergreen (PFT7), Boreal Broadleaf Summergreen (PFT8), Boreal Needleleaf Summergreen (PFT9), C3 Grass (PFT10), C3 Agriculture (PFT12), Water (PFT14), Snow/Ice (PFT15), are represented in color scale, from blue (0%) to red (100%).

Figure 7: Difference (New-Standard) PFT maps. The percentage differences are represented in color scale, ranging from -1 (blue) to +1 (red).

Figure 8: PFT dissimilarity index ranging from 0 (full agreement) to $\sqrt{2}$ (full disagreement).

9. Table captions

Table 1: List of the Land Cover products used and their characteristics

Table 2: Aligning the legends of global maps: dominant life form type (LFT) and corresponding land cover classes from GLC 2000, GLOBCOVER & MODIS IGBP

Table 3: Harmonized legend used and correspondence with original products classes

Table 4: Fraction of each class in each land cover product for Yakutian region

Table 5: Agreement percentages for Yakutia for the main classes and comparison of Globcover 2005 with Globcover 2009, GLC 2000, MODIS 5.0 and MODIS 5.1

Table 6: Fraction of each class in each land cover product for South-West Siberia

Table 7: Agreement percentages for South-West Siberia for the main land cover classes and comparison of Globcover 2005 with Globcover 2009, GLC 2000, MODIS 5.0 and MODIS 5.1

Table 8: Merging rules from GlobCover 2005 land cover classes to ORCHIDEE PFTs

	Products	Satellite et sensor	Time period	Spatial resolution	Projection	Cover zone	Number of classes
Land Cover	GLC2000 (v1.1)	SPOT-4 (VGT)	2000	1000 m	Plate-Carrée WGS84	Global	22
	MODIS MCD12Q1 (v5.0)	Terra Aqua (MODIS)	2001-2007	500m	Sinusoidal	Global	17
	MODIS MCD12Q1 (v5.1)	Terra Aqua (MODIS)	2001-2010	500 m	Sinusoidal	Global	17
	GlobCover Eastern Eurasia (v2.2) 2005	Envisat (Meris)	2004- 2006	300 m	Plate-Carrée WGS84	Regional map (Eastern Eurasia zone)	40
	GlobCover (v2.2) 2009	Envisat (Meris)	2009	300 m	Plate-Carrée WGS84	Global	22
Auxiliary	Circumpolar Arctic Vegetation Map	NOAA (AVHRR)	1993-1995	1:7.5 M	Lambert Azimuthal	Circumpolar Artic zone	18
	Yakutsk Land Cover (Fedorov et al., personal communication)	Landsat-TM	2002	60m	Lambert Azimuthal	Yakutsk region	12
	Forest height (Simard et al., 2011)	GLAS /ICESAT	2005	1km	Plate-Carrée WGS84	Global	1
	MODIS VCF (V4)	Terra (MODIS)	2000-2010	500 m	Sinusoidal	Global	3

Table 1: List of products used and their characteristics

Dominant LFT	GLC2000	GlobCover 2005	GlobCover 2009	MODIS
Tree	[1] Tree Cover, broadleaved, evergreen [2] Tree Cover, broadleaved, deciduous, closed [3] Tree Cover, broadleaved, deciduous, open [4] Tree Cover, needle-leaved, evergreen [5] Tree Cover, needle-leaved, deciduous [6] Tree Cover, mixed leaf type [7] Tree Cover, regularly flooded, fresh water [8] Tree Cover, regularly flooded, saline water [10] Tree Cover, burnt	[40] Closed to open broadleaved evergreen or semi-deciduous forest [50] Closed broadleaved deciduous forest [60] Open broadleaved deciduous forest/woodland [70] Closed needleleaved evergreen forest [91] Open needleleaved deciduous forest [90] Open needleleaved deciduous or evergreen forest [92] Open (15-40%) needleleaved evergreen forest [100] Closed to open (>15%) mixed broadleaved and needleleaved forest [101] Closed (>40%) mixed broadleaved and needleleaved forest	[40] Closed to open broadleaved evergreen or semi-deciduous forest [50] Closed broadleaved deciduous forest [60] Open broadleaved deciduous forest/woodland [70] Closed needleleaved evergreen forest [90] Open needleleaved deciduous or evergreen forest [100] Closed to open mixed broadleaved and needleleaved forest	[1] Evergreen needleleaf forest [2] Evergreen broadleaf forest [3] Deciduous needleleaf forest [4] Deciduous broadleaf forest [5] Mixed forest 8] Woody savannas [9] Savannas
Shrub	[11] Shrub Cover, closed-open, evergreen [12] Shrub Cover, closed-open, deciduous	[130] Closed to open shrubland [131] Closed to open (>15%) broadleaved or needleleaved evergreen shrubland [134] Closed to open broadleaved deciduous shrubland	[130] Closed to open shrubland	[6] Closed shrublands [7] Open shrublands
Herbaceous	[16] Cultivated and managed areas [13] Herbaceous Cover, closed-open	[11] Post-flooding or irrigated croplands (or aquatic) [12] Post-flooding or irrigated shrub or tree crops [13] Post-flooding or irrigated herbaceous crops [14] Rainfed croplands [15] Rainfed herbaceous crops [16] Rainfed shrub or tree crops [140] Closed to open (>15%) herbaceous vegetation [141] Closed grassland [143] Open grassland	[11] Post-flooding or irrigated croplands (or aquatic) [14] Rainfed croplands [140] Closed to open (>15%) herbaceous vegetation	[12] Croplands [10] Grasslands
Barren	[14] Sparse herbaceous or sparse shrub cover [22] Artificial surfaces and associated areas [19] Bare Areas [21] Snow and Ice	[150] Sparse (<15%) vegetation [151] Sparse (<15%) grassland [152] Sparse (<15%) shrubland [190] Artificial surfaces and associated areas [200] Bare areas [201] Consolidated bare areas (hardpans, gravels, bare rock, stones, boulders) [202] Non-consolidated bare areas (sandy desert) [203] Salt hardpans [220] Permanent snow and ice	[150] Sparse (<15%) vegetation [190] Artificial surfaces and associated area [200] Bare areas [220] Permanent snow and ice	[13] Urban and built-up [16] Barren or sparsely vegetated [15] Snow and ice
Mosaic	[9] Mosaic: Tree Cover / Other natural vegetation [15] Regularly flooded shrub and/or herbaceous cover [17] Mosaic: Cropland / Tree Cover / Other natural vegetation [18] Mosaic: Cropland / Shrub and/or grass cover	[20] Mosaic cropland (50-70%) / vegetation (grassland/shrubland/forest) (20-50%) [21] Mosaic cropland (50-70%) / grassland or shrubland (20-50%) [30] Mosaic vegetation (grassland/shrubland/forest) (50-70%) / cropland (20-50%) [32] Mosaic forest (50-70%) / cropland (20-50%) [110] Mosaic forest or shrubland (50-70%) / grassland (20-50%) [120] Mosaic grassland (50-70%) / forest or shrubland (20-50%) [170] Closed broadleaved forest or shrubland permanently flooded [180] Closed to open grassland or woody vegetation on regularly flooded or waterlogged soil [185] Closed to open grassland on regularly flooded or waterlogged soil	[20] Mosaic cropland (50-70%) / vegetation (grassland/shrubland/forest) (20-50%) [30] Mosaic vegetation (grassland/shrubland/forest) (50-70%) / cropland (20-50%) [110] Mosaic forest or shrubland (50-70%) / grassland (20-50%) [120] Mosaic grassland (50-70%) / forest or shrubland (20-50%) [160] Closed to open (>15%) broadleaved forest regularly flooded [170] Closed (>40%) broadleaved forest or shrubland permanently flooded [180] Closed to open (>15%) grassland or woody vegetation on regularly flooded or waterlogged soil	[11] Permanent wetlands [14] Cropland-natural vegetation mosaic
Water	[20] Water Bodies	[210] Water bodies	[210] Water bodies	[0] Water bodies

Table 2: Aligning legends of global maps: dominant life form type (LFT) and corresponding land cover classes from GLC 2000, GlobCover & MODIS

Classes	GlobCover 2005	GlobCover 2009	MODIS	GLC 2000
Evergreen Needleleaf Forest	92, 70	70	1	4
Evergreen Broadleaf Forest	X	X	2	X
Deciduous Needleleaf Forest	90, 91	90	3, 8, 9	5
Deciduous Broadleaf Forest	50, 60	50, 60	4	2
Mixed Forest	32, 40 ,100	40, 100	5	6
Mixed Forest-Shrubs	110, 12	110, 12	X	9
Shrubs	130, 131, 134	130	6	11, 12
Sparse Vegetation	150, 151, 152	150	7, 16	14
Herbaceous regularly flooded	180, 185	180	10	15
Herbaceous	30, 32, 140	30, 140	X	13
Urban	190	190	13	22
Bare Soil	200, 201, 202, 203	200	X	19
Snow/Ice	220	220	15	21
Water	210	210	0	20
Burnt areas	X	X	X	10
Croplands	11, 12, 13, 14, 15, 16, 20, 21	11, 14, 20	12, 14	16, 17, 18

Table 3: Harmonized legend used and correspondance with original products classes

Classes	GlobCover 2005	GlobCover 2009	MODIS5.0 (2005)	MODIS5.1 (2005)	GLC 2000
Evergreen Needleleaf Forest	0,03	0,00	0,01	0,01	0,04
Evergreen Broadleaf Forest	0,00	0,00	0,00	0,00	0,00
Deciduous Needleleaf Forest	0,56	0,58	0,42	0,28	0,45
Deciduous Broadleaf Forest	0,01	0,01	0,00	0,00	0,01
Mixed Forest	0,01	0,01	0,01	0,03	0,02
Mixed Forest-Shrubs	0,16	0,13	0,00	0,00	0,04
Shrubs	0,00	0,00	0,05	0,00	0,18
Sparse Vegetation	0,17	0,21	0,50	0,64	0,08
Herbaceous regularly flooded	0,01	0,02	0,02	0,04	0,04
Herbaceous	0,00	0,00	0,00	0,00	0,07
Urban	0,00	0,00	0,00	0,00	0,00
Bare Soil	0,05	0,05	0,00	0,00	0,06
Snow/Ice	0,00	0,00	0,00	0,00	0,00
Water	0,00	0,00	0,00	0,00	0,00
Burnt areas	0,00	0,00	0,00	0,00	0,00
Croplands	0,00	0,00	0,00	0,00	0,00

Table 4: Fraction of each class in each product for Yakutia

Classes	GlobCover2005 - GLC2000	GlobCover 2005 - GlobCover 2009	GlobCover 2005 - MODIS 5.0	GlobCover 2005 - MODIS 5.1	GLC 2000 - MODIS 5.0	GLC 2000 - MODIS 5.1	MODIS 5.0 - MODIS 5.1
Deciduous Needleleaf Forest	0,65	0,95	0,64	0,42	0,68	0,70	0,61
Mixed Forest	0,24	0,73	0,25	0,33	0,31	0,23	0,73
Mixed Forest-Shrubs	0,09	0,74	-	-	-	-	-
Shrubs	0,66	-	0,50	-	0,18	0,12	-
Sparse Vegetation	0,36	0,90	0,88	0,85	0,14	0,11	0,86
Herbaceous regularly flooded	0,18	0,97	0,20	0,22	0,04	0,04	0,31
Herbaceous	0,24	0,58	-	-	-	-	-
Bare Soil	0,49	0,83	-	-	-	-	-
Total Agreement	0,60	0,89	0,61	0,52	0,49	0,43	0,76
Total Agreement without Water	0,38	0,68	0,40	0,31	0,28	0,21	0,55

Table 5: Agreement percentages on Yakutia (for the main classes and comparison of Globcover 2005 with Globcover 2009, GLC 2000, MODIS 5.0, MODIS 5.1

Classes	GlobCover 2005	GlobCover 2009	MODIS 5.0 (2005)	MODIS 5.1 (2005)	GLC2000
Evergreen Needleleaf Forest	0,05	0,02	0,06	0,02	0,07
Evergreen Broadleaf Forest	0,00	0,00	0,00	0,00	0,00
Deciduous Needleleaf Forest	0,08	0,12	0,05	0,06	0,04
Deciduous Broadleaf Forest	0,13	0,11	0,00	0,00	0,01
Mixed Forest	0,03	0,06	0,23	0,26	0,24
Mixed Forest-Shrubs	0,05	0,03	0,00	0,00	0,00
Shrubs	0,00	0,00	0,01	0,00	0,02
Sparse Vegetation	0,17	0,13	0,02	0,01	0,03
Herbaceous regularly flooded	0,01	0,01	0,19	0,27	0,17
Herbaceous	0,17	0,10	0,00	0,00	0,00
Urban	0,01	0,00	0,00	0,00	0,00
Bare Soil	0,00	0,01	0,00	0,00	0,00
Snow/Ice	0,00	0,00	0,00	0,00	0,00
Water	0,00	0,00	0,00	0,00	0,00
Burnt areas	0,00	0,00	0,00	0,00	0,00
Croplands	0,30	0,41	0,43	0,37	0,43

Table 6: Fraction of each class in each product for South Siberia

Classes	GlobCover 2005 - GLC2000	GlobCover 2005 - GlobCover 2009	GlobCover 2005 - MODIS 5.0	GlobCover 2005 - MODIS 5.1	GLC 2000 - MODIS 5.0	GLC 2000 - MODIS 5.1	MODIS 5.0 - MODIS 5.1
Evergreen Needleleaf Forest	0,65	0,13	0,54	0,71	0,56	0,71	0,34
Deciduous Needleleaf Forest	0,16	0,68	0,13	0,07	0,10	0,07	0,29
Deciduous Broadleaf Forest	0,55	0,50	-	-	-	-	-
Mixed Forest	0,56	0,50	0,88	0,90	0,36	0,39	0,80
Mixed Forest- Shrubs	-	0,32	-	-	-	-	-
Sparse Vegetation	0,24	0,72	0,03	0,02	0,08	0,01	0,22
Herbaceous regularly flooded	-	-	-	-	-	-	0,93
Herbaceous	-	0,46	-	-	-	-	-
Croplands	0,54	0,96	0,79	0,70	0,42	0,67	0,80
Total Agreement	0,42	0,65	0,31	0,27	0,32	0,39	0,74
Total Agreement without Water	0,40	0,64	0,30	0,26	0,31	0,38	0,72

Table 7: Agreement percentages on South Siberia (for the main classes and comparison of Globcover2005 with Globcover2009, GLC2000, MODIS 5.0, MODIS 5.1)

ID	Globcover description	Bare Soil	Boreal Needleleaf Evergreen	Boreal Broadleaf Summer	Boreal Needleleaf Summer	C3 Grass	C3 Agri	Water	Snow/Ice
11	Post-flooding or irrigated croplands						100,0		
12	Post-flooding or irrigated shrub or tree crops						100,0		
13	Post-flooding or irrigated herbaceous crops						100,0		
14	Rainfed croplands						100,0		
15	Rainfed herbaceous crops						100,0		
16	Rainfed shrub or tree crops						100,0		
20	Mosaic cropland (50-70%)/vegetation (grassland/shrubland/forest) (20-50%)		10,0	15,0		15,0	60,0		
21	Mosaic cropland (50-70%) / grassland or shrubland (20-50%)		10,0		15,0	25,0	50,0		
30	Mosaic vegetation (grassland/shrubland/forest) (50-70%)/cropland (20-50%)		13,8	21,3		25,0	40,0		
32	Mosaic forest (50-70%) / cropland (20-50%)		10,0		50,0		40,0		
40	Closed to open broadleaved evergreen or semi-deciduous forest		47,5	52,5					
50	Closed broadleaved deciduous forest			85,0		15,0			
60	Open broadleaved deciduous forest/woodland	10,0		55,0		35,0			
70	Closed needleleaved evergreen forest		77,5	7,5		15,0			
90	Open needleleaved deciduous or evergreen forest	15,0	20,0	5,0	15,0	45,0			
91	Open needleleaved deciduous forest	10,0			60,0	30,0			
92	Open needleleaved evergreen forest	15,0	47,5	7,5		30,0			
100	Closed to open mixed broadleaved and needleleaved forest	10,0	27,5	37,5	10,0	15,0			
110	Mosaic forest or shrubland (50-70%)/grassland (20-50%)		17,5	32,5	10,0	40,0			
120	Mosaic grassland (50-70%)/forest or shrubland (20-50%)		12,5	22,5	5,0	60,0			
130	Closed to open shrubland	10,0	30,0	30,0		30,0			
131	Closed to open broadleaved or needleleaved evergreen shrubland	10,0	45,0	15,0		30,0			
134	Closed to open broadleaved deciduous shrubland	15,0		40,0		45,0			
140	Closed to open herbaceous vegetation	40,0				60,0			
141	Closed (>40%) grassland	40,0				60,0			
143	Open (15-40%) grassland	60,0				40,0			
150	Sparse (<15%) vegetation	35,0	9,4	9,4	6,3	40,0			
151	Sparse (<15%) grassland	35,0				65,0			
152	Sparse (<15%) shrubland	35,0	7,5	7,5	5,0	45,0			
160	Closed to open (>15%) broadleaved forest regularly flooded		15,0	45,0		20,0		20,0	
170	Closed (>40%) broadleaved forest or shrubland permanently flooded		40,0	40,0				20,0	
180	Closed to open (>15%) grassland or woody vegetation on regularly flooded		15,0	15,0		40,0		30,0	
185	Closed to open (>15%) grassland on regularly flooded or waterlogged soil					70,0		30,0	
190	Artificial surfaces and associated areas	75,0	2,5	2,5		15,0		5,0	
200	Bare areas	100,0							
201	Consolidated bare areas (hardpans, gravels, bare rock, stones, boulders)	100,0							
202	Non-consolidated bare areas (sandy desert)	100,0							
203	Salt hardpans	100,0							
210	Water bodies							100,0	
220	Permanent snow and ice								100,0

Table 8: Merging rules from GlobCover classes to ORCHIDEE PFTs







Figure 1




GLC2000




GlobCover 2005



MODIS 5.0 2001



MODIS 5.1 2001



Vegetation map of the Middle Lena River Basin,
compiled by A.N. Fedorov, Y.I. Torgovkin, A.I. Vasiliev,
M.I. Petrov and A.A. Shestakova

Legend




Figure 2




Figure 3








Figure 4




1 : Bare soil




4 : Temperate Needleleaf Evergreen




5 : Temperate Broadleaf Evergreen




6 : Temperate Broadleaf Summergreen




7 : Boreal Needleleaf Evergreen




9 : Boreal Needleleaf Summergreen



10 : C3 Grass




12 : C3 Agriculture




13 : C4 Agriculture


Figure 5




1 : Bare soil




7 : Boreal Needleleaf Evergreen




8 : Boreal Broadleaf Summergreen




9 : Boreal Needleleaf Summergreen




10 : C3 Grass



12 : C3 Agriculture




14 : Water




15 : Snow/ice


Figure 6




1 : Bare soil




4 : Temperate Needleleaf Evergreen




5 : Temperate Broadleaf Evergreen




6 : Temperate Broadleaf Summergreen




7 : Boreal Needleleaf Evergreen




9 : Boreal Needleleaf Summergreen



10 : C3 Grass



12 : C3 Agriculture
Figure 7



13 : C4 Agriculture




Figure 8