

A new compilation of stomach content data

J. K. Pinnegar et al.

A new compilation of stomach content data for commercially-important pelagic fish species in the Northeast Atlantic

**J. K. Pinnegar¹, N. Goñi², V. M. Trenkel³, H. Arrizabalaga², W. Melle⁴, J. Keating⁵,
and G. Óskarsson⁶**

¹Centre for Environment, Fisheries and Aquaculture Science (Cefas), Pakefield Road, Lowestoft, Suffolk, NR33 0HT, UK

²AZTI Fundazioa, Herrera kaia Portualdea z/g, 20110 Pasaia, Gipuzkoa, Spain

³ Institut français de recherche pour l'exploitation de la mer (IFREMER), BP 21105, 44311 Nantes, France

⁴Institute of Marine Research (IMR), P.O. Box 1870 Nordnes, 5817 Bergen, Norway

⁵Marine and Freshwater Research Centre, Galway-Mayo Institute of Technology (GMIT), Galway, Ireland

⁶Marine Research Institute (MRI), Skulagata 4, 121 Reykjavik, Iceland

Received: 9 January 2014 – Accepted: 1 March 2014 – Published: 14 April 2014

Correspondence to: J. K. Pinnegar (john.pinnegar@cefas.co.uk)

Published by Copernicus Publications.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

[Back](#)

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



Abstract

There is increasing demand for information on predator-prey interactions in the ocean as a result of legislative commitments aimed at achieving sustainable exploitation. However, comprehensive datasets are lacking for many fish species and this has hampered development of multispecies fisheries models and the formulation of effective food-web indicators. This work describes a new compilation of stomach content data for five pelagic fish species (herring, blue whiting, mackerel, albacore and bluefin tuna) sampled across the northeast Atlantic and submitted to the PANGAEA open-access data portal (www.pangaea.de). We provide detailed descriptions of sample origin and of the corresponding database structures. We describe the main results in terms of diet composition and predator-prey relationships. The feeding preferences of small pelagic fish (herring, blue whiting, mackerel) were sampled over a very broad geographic area within the North Atlantic basin, from Greenland in the west, to the Lofoten Islands in the east and from the Bay of Biscay northwards to the Arctic. This analysis revealed significant differences in the prey items selected in different parts of the region at different times of year. Tunas (albacore and bluefin) were sampled in the Bay of Biscay and Celtic Sea. Dominant prey items for these species varied by location, year and season. This data compilation exercise represents one of the largest and most wide-ranging ever attempted for pelagic fish in the north Atlantic. The earliest data included in the database were collected in 1864, whereas the most recent were collected in 2012. Datasets are available at [doi:10.1594/PANGAEA.820041](https://doi.org/10.1594/PANGAEA.820041) and [doi:10.1594/PANGAEA.826992](https://doi.org/10.1594/PANGAEA.826992).

1 Introduction

Food-webs have become a major focus for EU research and maritime policy. The 2008 European Marine Strategy Framework Directive (2008/56/EC) includes a commitment that Member States should work to achieve “Good Environmental Status” (GES) by

ESSDD

7, 197–223, 2014

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



information on pelagic fishes (i.e. those living at or near the surface of the ocean) and especially those with a wide geographic distribution.

The EU Euro-Basin project aims to understand and predict the dynamics of plankton and pelagic fish species in the North Atlantic, and to assess the impacts of climate variability. This project has a particular focus on herring (*Clupea harengus*), mackerel (*Scomber scombrus*) and blue whiting (*Micromesistius poutassou*), which are the most abundant and widespread planktivorous fish species in the region, as well as bluefin tuna (*Thunnus thynnus*) and albacore (*Thunnus alalunga*), top predator species that inhabit the whole North Atlantic basin and carry out trans-oceanic migrations.

In the present paper we provide details of newly digitized information on the diet of these five species, firstly from the UK DAPSTOM database (mackerel, herring and blue whiting), that incorporates information from recent research cruises by the Institute of Marine Research – IMR (Norway), Marine Research Institute – MRI-HAFRO (Iceland), Institut français de recherche pour l'exploitation de la mer – IFREMER (France), – Marine Institute, (Ireland), as well as historic data from the Centre for Environment, Fisheries and Aquaculture Science (Cefas) in the UK. We also examine data from AZTI-Tecnalia (Spain) on bluefin tuna and albacore stomach contents. Datasets derived under the Euro-Basin project have been submitted to, and are available via, the PANGAEA open-access data portal (www.pangaea.de).

2 Data and methods

The two datasets described here provide information on feeding preferences of 27 746 individual herring, 7423 mackerel, 1126 blue whiting, 1526 albacore and 689 bluefin tuna. Samples have been obtained from scientific research surveys (using a variety of different fishing gears), from commercial fishery catches, and from individual fish voluntarily sampled by recreational fishermen. In the following two sub-sections we describe the origins of the data and database structures as well as the geographical and temporal coverage.

ESSDD

7, 197–223, 2014

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.1 The DAPSTOM database

The DAPSTOM database has been in existence for 8 years, having been created in response to a “data-rescue” call from the EU “Network of Excellence” project EU-ROCEANS. The most recent version of the DAPSTOM dataset (Version 4.7, col-
lated in January 2014) includes 226 407 records derived from 449 distinct research
cruises, spanning the period 1837–2012. The database contains information from
254 202 individual predator stomachs and 188 species. As such, this represents one
of the largest and most diverse compilations of food-web information anywhere in the
world. A key component of the DAPSTOM programme has been an online data por-
tal (www.cefas.defra.gov.uk/fisheries-information/fish-stomach-records.aspx) through
which outputs are made freely available to the wider scientific community. In addition,
a subset of the pelagic fish information contained in the full DAPSTOM database has
now been made available via PANGAEA (doi:10.1594/PANGAEA.820041).

As the DAPSTOM initiative has progressed, a relational-database structure has
evolved (in Microsoft Access) that can accommodate all formats of stomach content
information (see Hyslop, 1980), including data collected at the level of individual fish,
pooled samples of multiple fish stomachs, frequency of occurrence data as well as fully
gravimetric information (prey weights or volumes). As a minimum, in Version 4.7 of the
DAPSTOM database, information on the predator species, geographic area and the
number of stomachs examined was required for a dataset to be included. Information
on predator length (or size range) was also widely available.

Central to the relational-database structure is the “DAPSTOM” data table (Fig. 1).
This includes much of the “raw” information about both the predator and prey. The
“DAPSTOM” data table includes 23 information fields (Fig. 1), and a full definition of
each field is provided by Pinnegar (2014). The “HAULS” table contains all information
about the geographic location from which the sample was derived. In most cases this
includes ship name, dates and times, latitudes, longitudes, depth, gear type, ICES area
and any additional information. Each “haul” has been assigned to a predefined “Sea”

ESSDD

7, 197–223, 2014

A new compilation of
stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(e.g. North Sea, Irish Sea, W Ireland, Celtic Sea, Channel, Biscay etc.) and ICES “Division” – a spatial sub-unit used by the International Council for the Exploration of the Sea.

5 A new innovation within version 4.7 of the DAPSTOM database is the inclusion of a “PROVENANCE” look-up table (Fig. 1). The purpose of this is to record the origin of the data, for example whether or not the data has been derived from published sources or based upon “raw” data files from collaborating scientists. Two additional look-up tables have been created to help standardise the taxonomic information that is available to users. The “PREDATOR” look-up table expands on the 3 digit predator names in the
10 “DAPSTOM” table and gives the predator’s full latin name, common name (in English), 10 digit NODC code and TSN identifier. The “PREY” look-up table aims to reduce the enormous number of potential prey names and descriptions to a manageable number of standardized names that can be used for analyses and collation. It corrects historic taxonomy to modern counterparts, and allows aggregation by broad prey groups (e.g.
15 euphausiids, amphipods, copepods, teleosts etc.).

The DAPSTOM dataset has now seen wide usage among ICES Working Groups as well as in a number of theoretical ecology papers (e.g. Rochet et al., 2011; Rossberg et al., 2011; Brose et al., 2006). On the whole, researchers have used the online portal to look at the diet composition of their favoured predator species – however there has
20 also been some interest in making use of historical datasets to determine long-term changes in fish diets at particular localities (Le Quesne and Pinnegar, 2012).

2.2 The AZTI tuna stomach database

25 The feeding ecology of temperate tunas (albacore *Thunnus alalunga* and bluefin tuna *Thunnus thynnus*) in the Bay of Biscay has been investigated through different projects in AZTI-Tecnalia since the mid-2000s. Although there has been little continuity between the projects dealing with the feeding ecology of temperate tunas, a total of 1525 stomachs have been sampled from albacore and 686 stomachs from bluefin tuna between 2004 and 2011.

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The sample origin and methodologies employed have been diverse, samples were collected:

1. During albacore acoustic tracking surveys (summer 2005), albacore dummy archival tagging surveys (summers 2005 and 2006), albacore archival tagging survey (June 2010), bluefin tuna tagging surveys (2009, 2011, 2012). In the case of these samples the location and the hour of each capture was included in the database.
2. Through collaborating recreational fishermen in the Southeastern Bay of Biscay in 2005, 2006 and 2007.
3. Through opportunistic sampling by the canning industry for albacore (2005, 2006, 2010, 2011) and in wholesalers stores for bluefin tuna (2009–2012). In this case, the catch dates and estimated locations of each predator were re-constructed a posteriori using information contained in the logbook of each fishing vessel. No catch hour was recorded in the case of these commercial catches, and the estimated locations correspond to the centers of the ICES statistical rectangles recorded in the logbooks.

The gears used to catch each sampled tuna also differed considerably, these included: (1) rod and reel (RR) primarily used by recreational fishermen; (2) trolling gear (TR) used during albacore tagging surveys; (3) pole and line with live bait (BB), used for most bluefin tuna catches (commercial and surveys) and for part of the commercial albacore catch, (4) pelagic trawl (MWT) used for a further part of the commercial albacore catch.

Due to the lower number of predators sampled in the case of tunas (in comparison with small pelagic fishes such as herring, blue whiting and mackerel), the relatively restricted geographical location of the samples (mostly Bay of Biscay) especially in the case of bluefin tuna, we chose not to include the tuna data into the DAPSTOM data portal but rather to build a simple database in XML format held at AZTI-Tecnalia,

ESSDD

7, 197–223, 2014

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



but also uploaded to PANGAEA (doi:10.1594/PANGAEA.826992). The tuna stomach database is provided as a single table. Each row corresponding to a single predator. For each predator, the information provided included the species, the sample origin (survey or opportunistic sampling), the catch date and location (either measured or assigned according to logbooks information), the size and weight, and the gear used for the catch.

3 Results – data submitted to PANGAEA under the EU project Euro-Basin

3.1 DAPSTOM dataset

Throughout 2013, Euro-Basin partners submitted datasets to the lead author of this paper and these were reformatted into the required DAPSTOM relational tables (Fig. 1). Datasets made available as part of Euro-Basin are summarised in Table 1.

It is clear that the vast majority of the blue whiting data within the DAPSTOM database, were collected in recent years, and explicitly for the purposes of Euro-Basin (1367 records out of 1583), whereas this was not true for herring or mackerel (Table 1). It is also apparent from Tables 1 and 2 that for blue whiting and mackerel the number of database records exceeded the number of stomachs examined, confirming that the data were largely non-pooled records from individual stomachs whereas this was not true for herring where 8508 database records were derived from 27 746 stomachs. The primary explanation for this disparity is the digitisation of “pooled” herring datasets from a historical report by Hardy (1924), but also “pooled” data from Brook and Calderwood (1886) and Scott (1924).

Table 2 shows the number of records and samples by geographic area (Fig. 2), including all larval and juvenile fish. From this table it is apparent that herring, blue whiting and mackerel have been sampled over a huge geographic area, spanning from the Bay of Biscay (~ 43° N), to the high Arctic (~ 73° N) and from Greenland in the west (~ 29° W) to the Lofoten islands in the east (~ 9° E). By contrast the very limited number

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of records for Albacore and Bluefin tuna in the DAPSTOM database were derived from opportunistic sampling in the English Channel and North Sea.

The earliest data included in the DAPSTOM/Euro-Basin dataset is a single record of an albacore tuna stranded on the Channel coast of England in August 1864, whereas the most recent data comes from a single bluefin tuna stranded at Ventnor, Isle of Wight in August 2012. The pelagic fish dataset includes information on the feeding preferences of fish larvae (0.1 to 10 cm in length), as well as adult fish. Specifically, the feeding habits of larval/juvenile herring and mackerel from Plymouth Sound, the Clyde and the North Sea by Lebour (1921, 1924), Marshall et al. (1937, 1939) and Last (1980) respectively.

3.2 AZTI-Tecnalia dataset

The tuna stomach database from AZTI-Tecnalia represents 7 years of sampling from 2004 to 2011. Due to the absence of continuity in the different projects dealing with the feeding ecology of tunas, the sampling could not be performed every year for both species, and no samples were collected in 2008. The lack of temporal continuity is more apparent for bluefin tuna than for albacore. However, a total of 1525 albacore and 686 bluefin tuna stomachs were collected during the study period (Table 3).

4 Discussion

A major limitation of the DAPSTOM dataset is that it comprises a mixture of “pooled” information and data collected from individual fish. Sometimes only information on the number of stomachs containing a particular prey item was available (i.e. “frequency of occurrence”), rather than the actual number of a particular prey item. Hence in any data extraction, outputs should be viewed as providing information on the “minimum number” of prey items consumed. This would have little impact in predator species that consume large prey items one at a time (e.g. fish feeders), but also with regard to most

ESSDD

7, 197–223, 2014

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the newer datasets assembled under Euro-Basin. However, in certain older datasets, the total number of prey items in plankton-eating species such as mackerel, herring and blue whiting would be underestimated. An example would be the historical dataset containing mackerel stomachs off the Cornish coast from Bullen (1908) a component of the “Cefas Historical” records cited in Table 1, as well as the herring datasets digitised from Marshall et al. (1937, 1939).

A further limitation of the DAPSTOM database is paucity of information on prey weights. In many of the constituent datasets no gravimetric information was provided. A result is that it can be difficult to judge the importance of a particular prey item to the overall nourishment of the predator, since a mackerel for example, may draw significantly more nourishment from eating a single fish in comparison with 1000+ copepods. To remedy this situation, we plan to develop an updated “PREY” table that includes average prey weights, and perhaps energy density for each standardised prey type so that numbers consumed can be converted to total weights, however this feature is not yet available.

Several authors have suggested that the preferred prey of blue whiting are euphausiids and hyperiid amphipods, although the relative importance of each of these varies depending on season and locality (Prokopchuk and Sentyabov, 2006; Langøy et al., 2012). The Euro-Basin dataset (Tables 1 and 2) shows similar variability in diet composition depending on sampling location (Fig. 3), with euphausiids dominating in terms of number in Iceland, the Bay of Biscay and the Irish Sea, but hyperiid amphipods dominating in the Norwegian Sea, eastern Greenland/northern Iceland (ICES Sub-Area XIV) and the Celtic Sea. Copepods (mainly *Calanus finmarchicus*) were an additional important prey item in the Norwegian Sea and shrimps (in particular *Pasiphaea sivado*) were commonly observed in stomachs from the Irish Sea. Adult blue whiting migrate in the springtime, to the Porcupine and Rockall areas west of Ireland. During this season they feed very infrequently.

Post larval mackerel feed on a variety of zooplankton and small fish. Published sources suggest that the main zooplankton prey organisms in the North Sea are

ESSDD

7, 197–223, 2014

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



copepods (mainly *Calanus finmarchicus*), and euphausiids (mainly *Meganyctiphanes norvegica*), while fish prey include larval sandeel, herring and sprat (Mehl and Westgård, 1983). In the Norwegian Sea published sources suggest that euphausiids, copepods, pteropod molluscs (*Limacina retroversa*), amphipods, appendicularia and capelin are the main dietary items (Langoy et al., 2012; Prokopchuk and Sentyabov, 2006). The Euro-Basin dataset confirms these broad patterns (Fig. 4), and in every geographic region for which data were available (Table 2), with the exception of the Bay of Biscay, copepods dominated in terms of numerical abundance, especially in Iceland (91 %). However, mysids and hyperiid amphipods contributed a significant additional proportion to the diet of mackerel in the North Sea (16 %). Hyperiids and euphausiids contributed a significant additional proportion to the diet in east Greenland/north Iceland (ICES Sub-Area XIV, 32 %) and phytoplankton, teleosts and chaetognaths contributed a significant proportion in the Celtic Sea (27 %). In the Bay of Biscay, 67 % of the diet composition (by number) was suggested to comprise mackerel eggs (denoted as “teleosts” in Fig. 4), although the vast majority of these data originate from a single research cruise in March 1986 and from a very limited number of haul stations.

For herring, many detailed diet composition studies of have been published, starting with Hardy (1924) and Jespersen (1928). In the Norwegian Sea diet has been shown to vary depending on availability of food and geographic location (Prokopchuk and Sentyabov, 2006; Langøy et al., 2012). *C. finmarchicus* is an important prey in summer (about 77 % by weight), but in certain years appendicularians (*Oikopleura* spp.), amphipods (mainly *Parathemisto abissorum*), and euphausiids are important. Similar variability has been noted for the North Sea, where pteropod molluscs (*Limacina retroversa*), sandeel (*Ammodytes* spp.) larvae, diatoms and the copepod *Temora longicornis* can also be locally important (Hardy, 1924; Savage, 1931). In the Euro-Basin dataset, copepods dominated herring diets in terms of numerical abundance in the Norwegian Sea, Iceland, North Sea and West of Scotland (69, 85, 66, 74 % respectively, Fig. 5). Hyperiid amphipods contributed a significant additional proportion in the Norwegian Sea (24 %), appendicularians contributed a significant but smaller propor-

ESSDD

7, 197–223, 2014

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion in the North Sea (15 %) and barnacle cypris larvae about 21 % in the west of Scotland. In eastern Greenland/northern Iceland (ICES Division XIV), euphausiids were the dominant prey item (63 %), followed by copepods (16 %) and hyperiid amphipods (15 %). In the Irish Sea euphausiids comprised 49 % of the diet and fish eggs (denoted as “teleosts” in Fig. 5, but mostly plaice *Pleuronectes platessa*) contributed a further 31 %. However, these Irish Sea research cruises (in February 2009, 2010 and 2011) were deliberately timed to quantify the seasonal predation mortality imparted by pelagic fish on plaice eggs and larvae, so it is not surprising that this particular prey item featured strongly.

Albacore diet composition in the northeast Atlantic has been reported as being dominated by small, mesopelagic fish, e.g. *Maurolicus muelleri* (pearlside) and *Scomberomorus saurus* (Atlantic saury), as well as also euphausiids and hyperiid amphipods (Pusineri et al., 2005; Goñi et al., 2011). In the AZTI data submitted to PANGAEA similar patterns were observed, with euphausiids being dominant and the most ubiquitous prey in albacore diet. Albacore diet displayed high plasticity with important spatial variability, both latitudinally and in terms of oceanic vs. shelf-break waters. Among small pelagic fish species, *Maurolicus muellerii* was less abundant in our samples than in previous studies, whereas Atlantic saury, blue whiting and anchovy (*Engraulis encrasicolus*) were the major prey in the shelf-break areas of the Bay of Biscay and Celtic Sea. There was also considerable latitudinal variability in Atlantic saury consumption by albacore. This prey represented a larger proportion of albacore diet in northern sampling areas than within the Bay of Biscay. This corroborates previous observations by Aloncle and Delaporte (1974), who related the presence of Atlantic saury to relatively low sea surface temperatures (SST) found out of the Bay of Biscay. This result suggests a potential higher predation impact on Atlantic saury when the summer distribution of albacore shifts westwards, as in 2009–2011.

Blue whiting consumption by albacore appeared to be related to the shelf-break of the Bay of Biscay and Celtic Sea but did not vary significantly with latitude. Consumption patterns suggested very marked inter-annual variation, with particularly low pres-

ence in albacore diet in 2010 and 2011 compared to 2005–2007. Taking into account the decrease in blue whiting biomass in recent years (Payne et al., 2012), this decrease in blue whiting consumption by albacore was likely related to a lower availability of the prey rather than to a shift in feeding preferences. This is corroborated by the observed proportion of blue whiting in the diet of bluefin tuna, which were sampled mostly in shelf-break locations.

Anchovy consumption also displayed seasonal and latitudinal variability, being higher in the late summer and autumn in the southern Bay of Biscay (Fig. 6). Within the Bay of Biscay, we observed a broader spatial distribution of anchovy in albacore diet during 2009–2011 than during 2005–2007. This broader distribution was probably related to the recovery of the anchovy population after a period of collapse between 2005 and 2008. The combined variability of the spatial extension of juvenile anchovies and of albacore distribution in summer months suggests a distinct spatial match/mismatch and predation impact each year.

Bluefin tuna diet in the northeast Atlantic has been reported by Logan et al. (2011). In the Bay of Biscay, euphausiids (*M. norvegica*) and anchovy made up 39 % of prey weight, with relative consumption of each reflecting annual changes in prey abundance. These same data, as well as more recent information have been submitted to PAN-GAEA as part of the Euro-Basin project. Regarding anchovy consumption in particular, the same seasonal variability as in albacore diet was apparent in bluefin tuna diet. This seasonal variability in both predators is likely to be related to anchovy life-cycle and to the ecology of juvenile stages. Adult anchovies usually spawn on the continental shelf during spring. After a planktonic phase, juveniles start forming monospecific schools and leaving the continental shelf to reach oceanic waters (Irigoien et al., 2007) from early August onwards. This shift from the continental shelf to oceanic waters explains the higher consumption of juvenile anchovy by bluefin tunas in the second half of the summer. We also observed inter-annual variability in anchovy consumption, apparently following the interannual variations in anchovy recruitment and distribution.

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Blue whiting is an important prey for bluefin tuna, especially in the first half of the summer when juvenile anchovies are not available. As with albacore, we observed a decrease in blue whiting consumption in the most recent years, following a decrease in recruitment (Payne et al., 2012). Atlantic saury, possibly because of the effect of SST on their distribution, were not as prevalent in bluefin tuna diet as they were in albacore diet, as bluefin tunas were exclusively sampled in the southeastern part of the Bay of Biscay. However, we did notice a higher proportion of saury in bluefin diet in 2011, when it seemed to partly replace anchovy as a high-caloric prey. Bluefin tuna used to be distributed in the North Sea, from where it disappeared in the 1960s, it is thought that in the past they fed primarily on herring and on mackerel in this region (Tiews, 1978). Occasional individuals have been recorded in the North Sea and around the British Isles in recent years. The few recent records available in the Euro-basin/DAPSTOM dataset suggest that they are targeting similar prey animals as was the case in the past.

Herring stocks in the northeast Atlantic support some of the largest fisheries in the World, with total catches in 2011 amounting to 1.5 million tonnes. Mackerel and blue whiting also support important fisheries and together these three species, with very large spawning stock biomasses (11.2, 2.7 and 3.8 million tonnes respectively in 2011), exert significant predation pressure on other commercial fish through feeding on fish eggs and larvae as well as depleting the standing stock of mesozooplankton (Engelhard et al., 2014). Small pelagic fish are themselves important prey items for larger predators, most notably tunas, but also seabirds and marine mammals in the north Atlantic. The total biomass of Albacore in the north Atlantic has been estimated at 76 000 tonnes (in 2012) and bluefin tuna at 285 000 tonnes (ICCAT, 2014). Within Euro-Basin, the datasets described in this paper will be used to calculate overall predation pressure exerted by the various species. The longer-term aspiration (through making these datasets available via PANGAEA and DAPSTOM) is that this action will facilitate the construction of more realistic ecosystem or multispecies fisheries models

ESSDD

7, 197–223, 2014

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that can subsequently be used to provide holistic advice, as now mandated by the EU and international conventions.

Acknowledgements. Research leading to these results received funding from the European Commission FP7 project EURO-BASIN (European Basin-Scale Analysis, Synthesis, and Integration, Grant Agreement 264 933). Development of the DAPSTOM data portal was initially supported through a “data rescue” grant from the EU Network of Excellence programme “EuroOceans”. Further iterations were supported by the UK Department for Environment, Food and Rural Affairs (specifically contracts MF1202 and MF1109). The AZTI tuna stomach sampling was funded through 4 research projects supported by the Basque Government (Agriculture and Fisheries Department).

References

- Aloncle, H. and Delaporte, F.: Données nouvelles sur le germon atlantique *Thunnus alalunga* Bonnaterre 1788 dans le nord-est Atlantique, 2ème Partie, Rev. Trav. Inst. Pêches Marit., 38, 5–102, 1974.
- Brook, G. and Calderwood, W. L.: Report on the food of herring. in: Appendix F, VI – Preliminary Reports on the Food of Fishes, Fourth Annual Report of the Fishery Board for Scotland, being for the year 1885, HM Stationary Office, Edinburgh, 102–128, 1886.
- Brose, U., Jonsson, T., Berlow, E. L., Warren, P., Banasek-Richter, C., Bersier, L. F., Blanchard, J. L., Brey, T., Carpenter, S. R., Blandenier, M. F., Cushing, L., Dawah, H. A., Dell, T., Edwards, F., Harper-Smith, S., Jacob, U., Ledger, M. E., Martinez, N. D., Memmott, J., Mintenbeck, K., Pinnegar, J. K., Rall, B. C., Rayner, T. S., Reuman, D. C., Ruess, L., Ulrich, W., Williams, R. J., Woodward, G., and Cohen, J. E.: Consumer–resource body–size relationships in natural food webs, *Ecology*, 87, 2411–2417, 2006.
- Bullen, G. E.: Plankton studies in relation to the western mackerel fishery, *J. Mar. Biol. Assoc. UK*, 8, 269–302, 1908.
- Dolgov, A. V., Yaragina, N. A., Orlova, E. L., Bogstad, B., Johannesen, E., and Mehl, S.: 20th anniversary of the PINRO–IMR cooperation in the investigations of feeding in the Barents Sea – results and perspectives, in: Proceedings of the 12th Norwegian–Russian Symposium, Tromsø, 21–22 August 2007, IMR/PINRO Report Series, 44–78, 2007.

ESSDD

7, 197–223, 2014

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Engelhard, G. H., Peck, M. A., Rindorf, A., Smout, S. C., van Deurs, M., Raab, K., Andersen, K. H., Garthe, S., Lauerburg, R. A. M., Scott, F., Brunel, T., Aarts, G., van Kooten, T., and Dickey-Collas, M.: Forage fish, their fisheries, and their predators: who drives whom?, ICES J. Mar. Sci., 71, 90–104, 2014.

5 EC: Reform of the Common Fisheries Policy: Communication from the Commission to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions, COM(2011) 417, 2011.

Goñi, N.: Stomach content records of albacore (*Thunnus alalunga*) and bluefin tuna (*Thunnus thynnus*) in the North Atlantic Drift Region focusing on the Bay of Biscay between 2004 and 2011, available at: doi:10.1594/PANGAEA.826992 (last access: March 2014), 2014.

10 Goñi, N., Logan, J., Arrizabalaga, H., Jarry, M., and Lutcavage, M.: Variability of albacore (*Thunnus alalunga*) diet in the Northeast Atlantic and Mediterranean Sea, Mar. Biol., 158, 1057–1073, 2011.

Hardy, A. C.: The herring in relation to its animate environment. Part I. The food and feeding habits of the herring with special reference to the east coast of England, MAFF Fishery Investigations II, 7, 1–39, 1924.

Hyslop, E. J.: Stomach contents analysis – a review of methods and their application, J. Fish Biol., 17, 411–429, 1980.

20 ICCAT: Report for biennial period, 2012–13, Part II (2013) – Vol. 2, SCRS, International Commission for the Conservation of Atlantic Tunas, 343 pp., 2014.

ICES: Report of the study group on multispecies model implementation in the Baltic, International Council for the Exploration of the Sea, ICES CM 1997/J:2, 1997.

ICES: Multispecies considerations for the North Sea stocks, ICES Advice 2013, Book 6, Sect. 6.3.1, June 2013, International Council for the Exploration of the Sea, Copenhagen, Denmark, 2013.

25 Irigoien, X., Fiksen, Ø., Cotano, U., Uriarte, A., Alvarez, P., Arrizabalaga, H., Boyra, G., Santos, M., Sagarminaga, Y., Otheguy, P., Etxebeste, E., Zarauz, L., Artetxe, I., and Motos, L.: Could Biscay Bay anchovy recruit through a spatial loophole?, Prog. Oceanogr., 74, 132–148, 2007.

30 Jespersen, P.: Investigations on the food of the herring in Danish waters, Medd. Komm. f. Havundersogelser. Serie Plankton, 2, 1–149, 1928.

Langøy, H., Nøttestad, L., Skaret, G., Broms, C., and Fernö, A.: Overlap in distribution and diets of Atlantic mackerel (*Scomber scombrus*), Norwegian spring-spawning herring (*Clupea*

harengus) and blue whiting (*Micromesistius poutassou*) in the Norwegian Sea during late summer, Mar. Biol. Res., 8, 442–460, 2012.

Last, J. M.: The food of twenty species of fish larvae in the west-central North Sea, MAFF Fisheries Research Technical Report, 60, 44 pp., 1980.

Lebour, M. V.: The food of young clupeoids, J. Mar. Biol. Assoc. UK, 12, 458–467, 1921.

Lebour, M. V.: The food of young herrings, J. Mar. Biol. Assoc. U. K., 13, 325–330, 1924.

Le Quesne, W. J. F. and Pinnegar, J. K.: The potential impacts of ocean acidification: scaling from physiology to fisheries, Fish. Fisher, 13, 333–344, 2012.

Logan, J. M., Rodríguez-Marín, E., Goñi, N., Barreiro, S., Arrizabalaga, H., Golet, W., and Lutcavage, M.: Diet of young Atlantic bluefin tuna (*Thunnus thynnus*) in eastern and western Atlantic foraging grounds, Mar. Biol., 158, 73–85, 2011.

Mackinson, S., Deas, B., Beveridge, D., and Casey, J.: Mixed-fishery or ecosystem conundrum? Multi-species considerations inform thinking on long-term management of North Sea demersal stocks, Can. J. Fish. Aquat. Sci., 66, 1107–1129, 2009.

Marshall, S. R., Nicholls, A. G., and Orr, A. P.: On the growth and feeding of the larval and post-larval stages of the Clyde herring, J. Mar. Biol. Assoc. UK, 22, 245–267, 1937.

Marshall, S. R., Nicholls, A. G., and Orr, A. P.: On the growth and feeding of young herring in the Clyde, J. Mar. Biol. Assoc. UK, 23, 427–455, 1939.

Mehl, S. and Westgård, T.: The diet and consumption of mackerel in the North Sea, International Council for the Exploration of the Sea, ICES CM 1983/H: 34, 1983.

Payne, M. R., Egan, A., Fässler, S. M. M., Hátún, H. L., Holst, J. C., Jacobsen, J. A., Slotte, A., and Loeng, H.: The rise and fall of the NE Atlantic blue whiting (*Micromesistius poutassou*), Mar. Biol. Res., 8, 475–487, 2012.

Pinnegar, J. K.: DAPSTOM – an Integrated Database and Portal for Fish Stomach Records, Version 4.7, Centre for Environment, Fisheries and Aquaculture Science (Cefas), Lowestoft, UK, February 2014, 39 pp., 2014.

Pinnegar, J. K., Trenkel, V. M., Melle, W., and Óskarsson, J. G.: Stomach content records for pelagic fish (herring, mackerel, blue whiting, albacore and bluefin tuna) in the north-east Atlantic, The Centre for Environment Fisheries and Aquaculture Science, Lowestoft, doi:10.1594/PANGAEA.820041, 2013.

Pusineri, C., Vasseur, Y., Hassani, S., Meynier, L., Spitz, J., and Ridoux, V.: Food and feeding ecology of juvenile albacore, *Thunnus alalunga*, off the Bay of Biscay: a case study, ICES J. Mar. Sci., 62, 116–122, 2005.

ESSDD

7, 197–223, 2014

A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Prokopchuk, I. and Sentyabov, E. Diets of herring, mackerel, and blue whiting in the Norwegian Sea in relation to *Calanus finmarchicus* distribution and temperature conditions, ICES J. Mar. Sci., 63, 117–127, 2006.

Rochet, M.-J., Collie, J. S., Jennings, S., and Hall, S. J.: Does selective fishing conserve community biodiversity? Predictions from a length-based multispecies model, Can. J. Fish. Aquat. Sci., 68, 469–486, 2011.

Rosberg, A. G., Farnsworth, K. D., Satoh, K., and Pinnegar, J. K.: Universal power-law diet partitioning by marine fish and squid, with surprising stability-diversity implications, P. Roy. Soc. B-Biol. Sci., 278, 1617–1625, 2011.

Savage, R. E.: The relation between the feeding of herring off the East coast of England and the plankton of the surrounding waters, MAFF Fishery Investigations II, 12, 1–88, 1931.

Scott, A.: Food of the Irish Sea Herring in 1923, in: Proceedings and Transactions of the Liverpool Biological Society, 38, 115–119, 1924.

Tiews, K.: On the disappearance of bluefin tuna in the North Sea and its ecological implications for herring and mackerel, Rapp. p.-v. Reun., 172, 301–309, 1978.

A new compilation of stomach content data

J. K. Pinnegar et al.

Table 1. Number of stomach content records for pelagic fish species submitted to the DAP-STOM database as part of Euro-Basin. Number of individual stomachs included in parentheses.

Dataset	Period	Herring	Blue whiting	Mackerel	Albacore	Bluefin tuna
IFREMER (France)	2011	0 (0)	133 (117)	0 (0)	0 (0)	0 (0)
IMR (Norway)	2004, 2006	1291 (538)	354 (139)	1772 (635)	0 (0)	0 (0)
MRI (Iceland)	2010, 2011	1610 (823)	274 (158)	3226 (1486)	0 (0)	0 (0)
GMIT (Ireland)	2011	0 (0)	139 (109)	0 (0)	0 (0)	0 (0)
Cefas (UK)	2010, 2011	1101 (961)	467 (366)	6 (3)	0 (0)	0 (0)
Cefas – Historical	1864–2009	4506 (25 424)	216 (237)	5614 (5299)	1 (1)	10 (3)
Total		8508 (27 746)	1583 (1126)	10 618 (7423)	1 (1)	10 (3)

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A new compilation of stomach content data

J. K. Pinnegar et al.

Table 2. Number of records for pelagic fish species submitted to the DAPSTOM stomach content database as part of Euro-Basin, by geographic region. Number of individual stomachs included in parentheses.

ICES Region (Sea)	Herring	Blue whiting	Mackerel	Albacore	Bluefin tuna
VIIIa, b,c (Bay of Biscay)	0 (0)	157 (139)	896 (598)	0 (0)	0 (0)
VIIIf, g,h, j (Celtic Sea)	66 (59)	506 (411)	2804 (2416)	0 (0)	0 (0)
VIIe, d (Channel)	577 (5077)	35 (24)	718 (789)	1 (1)	1 (1)
XIVa, b (E Greenland/N Iceland)	605 (246)	70 (29)	1050 (432)	0 (0)	0 (0)
Va (Iceland)	680 (405)	105 (72)	1356 (672)	0 (0)	0 (0)
VIIa (Irish Sea)	1294 (1285)	183 (166)	29 (19)	0 (0)	0 (0)
XII (North Atlantic)	0 (0)	18 (9)	0 (0)	0 (0)	0 (0)
IVa, b,c (North Sea)	2954 (16 480)	19 (49)	1106 (1225)	0 (0)	9 (2)
IIa (Norwegian Sea)	1616 (710)	435 (187)	2447 (987)	0 (0)	0 (0)
VIIb (West Ireland)	0 (0)	55 (40)	129 (116)	0 (0)	0 (0)
Via (West Scotland)	716 (3484)	0 (0)	83 (169)	0 (0)	0 (0)

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Number of stomach content records for albacore and bluefin tuna, sampled in the Bay of Biscay and submitted to the AZTI tuna stomach database as part of Euro-Basin, by year and fishing gear.

Year	Sample source (gear)	Albacore	Bluefin tuna
2004	Commercial fishery (BB)	0	32
2005	Sonic tracking and archival tagging surveys (TR)	166	12
2005	Recreational fishermen (RR)	162	24
2005	Commercial fishery (MWT)	69	0
2006	Archival tagging survey	49	0
2006	Recreational fishermen	68	3
2006	Commercial fishery (MWT)	79	0
2007	Recreational fishermen (RR)	37	0
2009	Commercial fishery (albacore MWT, bluefin BB)	95	238
2009	Archival tagging survey	0	19
2010	Commercial fishery (TR, BB, MWT)	532	233
2010	Archival tagging survey	34	0
2011	Commercial fishery (TR, BB, MWT)	234	68
2011	Conventional tagging survey (BB)	0	57
Total		1525	686

A new compilation of stomach content data

J. K. Pinnegar et al.




Fig. 1. Structure of the DAPSTOM 4.7 relational database, including a list of the fields contained within each table. For a full description of the field formats and nomenclature see Pinnegar (2014).

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 2. Location of stomach content samples (circles) for the five pelagic species collected in the north-east Atlantic and submitted to the DAPSTOM and AZTI databases.

A new compilation of stomach content data

J. K. Pinnegar et al.




Fig. 3. Diet composition of blue whiting *Micromesistius poutassou* in different parts of the north-east Atlantic (ICES Divisions). Proportions are based on the number of individual prey items.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A new compilation of stomach content data

J. K. Pinnegar et al.




Fig. 4. Diet composition of mackerel *Scomber scombrus* in different parts of the northeast Atlantic (ICES Divisions). Proportions are based on the number of individual prey items.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A new compilation of stomach content data

J. K. Pinnegar et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 5. Diet composition of herring *Clupea harengus* in different parts of the northeast Atlantic (ICES Divisions). Proportions are based on the number of individual prey items.

A new compilation of stomach content data

J. K. Pinnegar et al.




Fig. 6. Diet composition of albacore sampled in distinct areas and periods in the Bay of Biscay during the summer and autumn of 2010.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

