

This discussion paper is/has been under review for the journal Earth System Science Data (ESSD). Please refer to the corresponding final paper in ESSD if available.

# Data compilation on the biological response to ocean acidification: an update

Y. Yang<sup>1</sup>, L. Hansson<sup>2</sup>, and J.-P. Gattuso<sup>3,4,5</sup>

<sup>1</sup>State Key Laboratory of Marine Environmental Science, Xiamen University, 361005 Xiamen, China

<sup>2</sup>International Atomic Energy Agency, Environment Laboratories, 4a Quai Antoine 1er, 98000 Principality of Monaco

<sup>3</sup>CNRS-INSU, Laboratoire d'Océanographie de Villefranche, 181 chemin du Lazaret, 06230 Villefranche-sur-mer, France

<sup>4</sup>Sorbonne Universités, UPMC Univ Paris 06, Observatoire Océanologique, 06230 Villefranche-sur-mer, France

<sup>5</sup>Institute for Sustainable Development and International Relations, Sciences Po, 27 rue Saint Guillaume, 75007 Paris, France

Received: 29 September 2015 – Accepted: 25 October 2015 – Published: 11 November 2015

Correspondence to: Y. Yang (yangyan@xmu.edu.cn)

Published by Copernicus Publications.

Data compilation on  
the biological  
response to ocean  
acidification: an  
update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

The exponential growth of studies on the biological response to ocean acidification over the last few decades has generated a large amount of data. To facilitate data comparison, a data compilation hosted at the data publisher PANGAEA was initiated in 2008 and is updated on a regular basis (doi:10.1594/PANGAEA.149999). By January 2015, a total of 581 data sets (over 4 000 000 data points) from 539 papers had been archived. Here we present the developments of this data compilation five years since its first description by Nisumaa et al. (2010). Most of study sites from which data archived are still in the Northern Hemisphere and the number of archived data from studies from the Southern Hemisphere and polar oceans are still relatively low. Data from 60 studies that investigated the response of a mix of organisms or natural communities were all added after 2010, indicating a welcomed shift from the study of individual organisms to communities and ecosystems. The initial imbalance of considerably more data archived on calcification and primary production than on other processes has improved. There is also a clear tendency towards more data archived from multifactorial studies after 2010. For easier and more effective access to ocean acidification data, the ocean acidification community is strongly encouraged to contribute to the data archiving effort, and help develop standard vocabularies describing the variables and define best practices for archiving ocean acidification data.

## 1 Introduction

The release of carbon dioxide (CO<sub>2</sub>) into the atmosphere by human activities results in an increased flux of CO<sub>2</sub> into a mildly alkaline ocean, resulting in an increase in the concentration of inorganic carbon, and a reduction in pH, carbonate ion concentration, and the capacity of seawater to buffer changes in its chemistry. These changes are collectively known as ocean acidification (Gattuso et al., 2014).

ESSDD

8, 889–912, 2015

### Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Investigations of the effect of ocean acidification on marine organisms and ecosystems have a relatively short history. A wide range of sensitivities to projected rates of ocean acidification exists within and across diverse groups of organisms, with a trend for greater sensitivity in early life stages (Kroeker et al., 2013). Several meta-analyses reveal a pattern of positive and negative impacts (Gattuso et al., 2014) but key uncertainties remain in our understanding of the impacts on organisms, life histories, and ecosystems.

The number of papers addressing biological responses to ocean acidification has increased steeply in the past decade, from 18 papers per year in 2004 to 365 papers in 2014 (Gattuso and Hansson, 2011; OA-ICC bibliographic database, <http://www.tinyurl.com/oaicc-biblio>). It is challenging to compare and synthesize the results of these papers for two reasons. First, data are not easily discoverable and accessible because they are either not archived or archived in different data repositories in varying formats. Second, the carbonate chemistry and ancillary data are often reported in different units and scales, and calculated using different sets of constants, making the carbonate chemistry across studies inconsistent. For example, it is crucial to report the pH scale used, since pH reported on the free scale is about 0.11 to 0.12 units higher than on the total and seawater scales, respectively (Zeebe and Wolf-Gladrow, 2001).

A compilation of ocean acidification biological response data was initiated by the European Network of Excellence for Ocean Ecosystems Analysis (EUR-OCEANS) and the European Project on Ocean Acidification (EPOCA) in 2008 (Nisumaa et al., 2010). This effort ended in 2012 when the EPOCA project came to an end but was resumed in the framework of the Ocean Acidification International Coordination Centre (OA-ICC) in 2013, in close collaboration with Xiamen University and the world data center PAN-GAEA. The goal of the OA-ICC data compilation is to gather data on the biological response to ocean acidification (carbonate chemistry, biogeochemical processes and ancillary data) from published articles and to make them available in a coherent format to the scientific community. As part of the effort, the carbonate system variables are recalculated in a consistent way. Data from papers that report at least two car-

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



bonate chemistry parameters, as well as temperature and salinity, are included in the compilation. The data compilation has been very well received by the ocean acidification community. It has been used in three meta-analyses (Kroeker et al., 2010, 2013; Liu et al., 2010), a modeling study (Muller and Nisbet, 2014) and cited in six other publications (Fiorini et al., 2011; Hendriks and Duarte, 2010; Hoppe et al., 2012; Koeve and Oschlies, 2012; Meyer and Riebesell, 2014; Rokitta, 2012). We report here on the developments of this database five years since its first description by Nisumaa et al. (2010). This update is timely, since 500 data sets from 439 papers (81 % of the total number included in the compilation) have been archived since the publication of Nisumaa et al. (2010).

## 2 Data

The compilation process described in Nisumaa et al. (2010) was followed to maintain consistency. Briefly, papers on the biological response to ocean acidification were identified by searching the OA-ICC news stream (<http://news-oceanacidification-icc.org/>) or through the OA-ICC bibliographic database for older papers. Papers were excluded from the compilation when they only used levels of partial pressure of carbon dioxide ( $p\text{CO}_2$ ) or pH which are not consistent to present-day levels or future scenarios. For example, data collected with a control condition with  $p\text{CO}_2$  values below about  $100\ \mu\text{atm}$  or above  $1700\ \mu\text{atm}$  or  $\text{pH}_T$  (on the total scale) values below about 7.5 or above 8.5 were excluded unless they are environmentally relevant at the study location. Data were requested from the authors by email. Data were either provided by authors, or extracted from tables in the original paper, or downloaded from other data repositories such as the Biological and Chemical Oceanography Data Management Office (BCO-DMO), British Oceanographic Data Centre (BODC), Australian Antarctic Data Centre (AADC). All the data sets archived in the framework of the OA-ICC (since 2013) as well as the projects EPOCA/EUR-OCEANS (2008–2012) have been given the tag “Ocean Acidification International Coordination Centre (OA-ICC)” in PANGAEA.

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Between the beginning of this activity in 2008 and January 2015, 1026 relevant papers were identified. 581 data sets (over 4 000 000 data points) were archived from 539 of these papers. Data from a paper can be archived as several data sets (e.g. <http://doi.pangaea.de/10.1594/PANGAEA.777725>), which explains why out of 539 papers, there were 581 data sets archived. Data of the remaining 487 papers could not be added to the compilation for the following reasons (Fig. 1): (1) less than two carbonate system parameters were measured in 176 papers, preventing the recalculation of the carbonate chemistry, (2) data from 295 papers could not be obtained from the authors; and (3) data from 16 papers were lost by authors.

The first papers having investigated a biological response to decreased pH (e.g. Vernon, 1895) predate the definition of the pH scale by Sørensen (1909) and are obviously not amenable to be included in the data compilation. The earliest data included in the compilation were published in 1967 (Traganza, 1967). No data from the papers published during 1968–1993 could be archived because data were lost or could not be obtained. Data from 41 papers published during 1994 and 2006 were archived. The quantity of data archived for a given year follows the increase in publication rate of biological response papers, with data from 11 papers archived for 2007 to 110 papers for 2014. More than 90 % of the data archived in the compilation come from papers published after 2007.

All the references of papers included in the data compilation have been tagged with the keyword “OAICcdb” in the OA-ICC bibliographic database (<http://www.tinyurl.com/oaicc-biblio>). The OA-ICC bibliographic database was used to retrieve statistical information on the type of papers from which the data archived originated. Keywords describing the geographical region, type of organism and biological process studied, other factors manipulated, as well as the country of affiliation of the first author were extracted from the bibliographic database for statistical analysis. Results are presented as percent of papers from which data were archived. Information on salinity, temperature and carbonate chemistry data archived as part of the data compilation were extracted



## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(11 %) investigated the response of a mix of organisms or natural communities, while no such studies from which data were archived before 2010, indicating a clear shift away from the study of individual organisms to communities and ecosystems, as has been recommended to close existing gaps in ocean acidification science (Riebesell and Gattuso, 2015). This is important as data on competitive and trophic interactions are key to better predict future impacts of ocean acidification. The amount of papers concerning prokaryotes, protists, zooplankton and others (annelids, virus etc.) are relatively low (less than 8 % for each species).

### 2.3 Biological processes

The most studied biological processes are morphology, calcification and growth, representing 33, 30 and 26 % of the papers from which data were archived (Fig. 4). Physiology, photosynthesis, mortality, reproduction and respiration are also well represented (13 to 23 %), followed by primary production, performance, dissolution and nitrogen fixation. Other processes such as community composition, abundance, adaptation, nutrient uptake, toxicity and bleaching are representing 26 % of the papers from which data were archived (see detail description of keywords in the user instructions of the OA-ICC bibliographic database, [www.tinyurl.com/oaicc-biblio](http://www.tinyurl.com/oaicc-biblio)). All processes, except calcification and primary production, are better represented today than they were before 2010, indicating that the initial imbalance of considerably more studies on calcification and primary production than on other processes has improved.

### 2.4 Multiple factors

There is a clear tendency towards more multifactorial studies after 2010. Even though the majority of the compiled papers have only manipulated the carbonate chemistry, their relative contribution in the data compilation decreased from 81 (pre-2010) to 69 %. The main other factors studied in addition to the carbonate system are temperature (14 % of papers), nutrients (6 %) and light (5 %) (Fig. 5). Few papers manipulated oxy-

gen, metal, or toxic compounds. Twenty papers have also reported combined effect of changes in carbonate chemistry with two or three other factors. In general, data from studies on multifactorial impact are still limited in the data compilation. It is one of the major challenges faced by future ocean acidification research (Riebesell and Gattuso, 2015).

## 2.5 Countries of affiliation of first author

Based on affiliation of the first author, most of the data compiled are from articles contributed by European countries (59 %; Fig. 6). Within Europe, 41 % of the papers were published by German scientists and 19 % by UK scientists. The USA (18 %) and Australia (8 %) also contribute a lot to the data compilation. There are still data from many papers which could not be obtained from the authors from these two countries (87 and 49 papers respectively).

## 2.6 Measured carbonate chemistry variables

A complete and consistent set of carbonate system variables was calculated by the R package seacarb (Gattuso et al., 2015) as described by Nisumaa et al. (2010), in order to provide coherent information on the chemistry<sup>1</sup>. The recalculated parameters were archived together with the original ones and were flagged accordingly (“Calculated using seacarb after Nisumaa et al. (2010)” together with a reference to the data curator responsible for the recalculation).

Total alkalinity ( $A_T$ ) is the carbonate chemistry variable that is the most measured (79 % of the data sets; Fig. 7). The other variables measured include pH (70 %), dissolved inorganic carbon ( $C_T$ , 36 %) and the partial pressure of carbon dioxide ( $pCO_2$ , 8 %). Out of the 70 % data sets that measured pH, 38 % reported pH on the National Bureau of Standard (N.B.S.) scale (also referred to National Institute of Standards and

<sup>1</sup>Dissolved inorganic carbon ( $C_T$ ) is referred to DIC in the data compilation which begun before the guidelines shown in the Supplement were defined.

### Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Technology, N.I.S.T., scale), 30 % on the total scale, 1 % on the seawater scale (SWS) and 1 % on the free scale (Fig. 7). Although the number of data sets with pH reported on the NBS scale are still more than on the total scale, the ratio of them has been decreased from 2.2 (pre-2010) to 1.2. The pH value on the total scale is 0.15 units lower than on the N.B.S/N.I.S.T scale and 0.01 units higher than on the SWS scale (Dickson, 2010), which makes the direct comparison of experimental results difficult. All other scales are converted to the total scale in this compilation as recommended in the *Guide to Best Practices in Ocean Acidification Research and Data Reporting* (Dickson, 2010).

## 2.7 Temperature and salinity coverage

The temperature values reported range from  $-3$  to  $34^{\circ}\text{C}$  but most temperature data are between  $18$  and  $21^{\circ}\text{C}$  (Fig. 8a) and few values are below  $0^{\circ}\text{C}$  or above  $30^{\circ}\text{C}$ . Salinity ranges from  $4$  to  $65$  with 60 % of the data points ranging between  $32$  and  $36$  (Fig. 8b). The geographical distribution of the papers considered (Fig. 2) shows that the study sites are distributed over the all oceans of the world, including the high-salinity Red Sea and the low-salinity Baltic Sea. High-salinity lagoons and low-salinity estuarine areas have also been studied, which can explain the large range of salinity in the compilation. There are 36 data points with salinity greater than 50, they were derived from paper on brine algae collected from sea ice in Antarctica (McMinn et al., 2014).

## 2.8 Carbonate chemistry variables coverage

$\text{pH}_T$ ,  $\text{pCO}_2$ ,  $A_T$ ,  $C_T$ , aragonite ( $\Omega_a$ ) and calcite ( $\Omega_c$ ) saturation states and carbonate ion ( $\text{CO}_3^{2-}$ ) data included in this compilation span the range of the average open-ocean surface values from the Last Glacial Maximum (LGM) to 2100 (Fig. 8). The largest number of data points are comprised between  $\text{pH}_T$  7.65 to 7.95,  $\text{pCO}_2$  500 to 1000  $\mu\text{atm}$  and  $C_T$  2100 to 2200  $\mu\text{molkg}^{-1}$  (Fig. 8c, d, and f). This indicates that most studies

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



used carbonate chemistries that are environmentally relevant. Ambient  $p\text{CO}_2$  level is often used as the control treatment, so many  $\text{pH}_T$ ,  $p\text{CO}_2$  and  $C_T$  data points are close to the present-day values (8.07, 384  $\mu\text{atm}$  and 2064  $\mu\text{mol kg}^{-1}$ ). Few data points are close to the value during LGM and 1766, because few studies used preindustrial and “glacial”  $p\text{CO}_2$  level (respectively 267 and 180  $\mu\text{atm}$ ) as treatments. Some data points are indicative of carbonate chemistry that are not environmentally relevant (i.e.  $p\text{CO}_2$  below 150  $\mu\text{atm}$  or above 2000  $\mu\text{atm}$ ,  $\text{pH}_T$  below 7.35 or above 8.45). This is because in some studies they not only used  $p\text{CO}_2$  and  $\text{pH}_T$  relevant to past or for future scenarios as treatments, but also used extreme  $p\text{CO}_2$  and  $\text{pH}_T$  as treatments. Since  $A_T$  is unaffected by the uptake of  $\text{CO}_2$ , it has been recommended to keep it constant in perturbation experiments, for example by bubbling seawater with  $\text{CO}_2$  enriched air (Gattuso and Lavigne, 2009). In the OA-ICC data compilation, most of  $A_T$  data points ranges from 2250 to 2350  $\mu\text{mol kg}^{-1}$ , which is close to the average present-day open-ocean surface values (2325  $\mu\text{mol kg}^{-1}$ ) (Fig. 8e). However, some studies manipulated the carbonate chemistry by adding strong acids and bases without restoring total alkalinity (see Gattuso et al., 2010), which significantly altered  $A_T$ . The number of this kind of study has decreased since the publication of the *Guide to Best Practices in Ocean Acidification Research and Data Reporting* (Cornwall and Hurd, 2015). In addition, some low  $A_T$  data points were archived from studies which performed at low salinity areas, such as the Baltic Sea and estuarine areas. The concentration of  $\text{CO}_3^{2-}$ , as well as  $\Omega_a$  and  $\Omega_c$  exhibit similar distribution patterns, with most number of data points close to their average value of surface seawater for 2100 (Fig. 8g–i).

### 3 Recommendations

As a new scientific field such as ocean acidification develops, the amount of scientific data grows considerably and the need for data archiving, greatly increases to ensure that data are easily accessible for analysis and reuse. Many journals, such as *Proceedings of the Royal Society B-Biological Sciences*, *PLoS ONE* and *Nature*, require

or encourage authors to archive data in public data repositories. Several data journals were launched to help this process. For example, the present journal was launched by the European Geosciences Union in 2008 and *Scientific Data* of the Nature publishing group in 2014. Many funding agencies also have requirements on data archiving. The European Research Council (ERC) mandate funded scientists to deposit primary data in relevant databases as soon as possible after data collection. In the United States, researchers seeking funding from the National Science Foundation (NSF) are required to submit a data management plan as a supplement to the grant application. Research projects such as the European FP7 project EPOCA has provided guidelines on data reporting and discussed the benefits of data archiving (Pesant et al., 2010). Although data archiving is not only of benefit for the scientific community but also a great way to advertise scientific work, get more citations (Piwowar et al., 2007) and initiate new collaborations, it is a difficult task to collect data from authors. It is a slow process and only data from 53% of the relevant papers could be included in the compilation. We therefore encourage the international ocean acidification community to actively participate in the data archiving effort.

Another challenge is the use of consistent variable names as papers sometimes report the same variable with different names, for example “respiration” and “oxygen consumption”. It is recommended to develop standard vocabularies describing the variables. And the standard vocabularies should be used in the metadata template to ensure effective data searching (Jiang et al., 2015). At an expert meeting on the management of ocean acidification biological response data organized by the Ocean Acidification International Coordination Center in 2014, ocean acidification scientists and data managers from major data centers agreed to develop a list of the most common biological parameters from ocean acidification studies (Hansson et al., 2014). Journals are invited to make sure that variables of the carbonate chemistry are reported in full and with the required level of detail (see Supplement). Furthermore, some data have previously been published in PANGAEA by data curators of other projects (e.g. the project “Biological Impacts of Ocean Acidification”, BIOACID), or in

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



other databases (e.g. the Biological and Chemical Oceanography Data Management Office, British Oceanographic Data Centre, Australian Antarctic Data Centre), which induced duplicates when they were archived again for the OA-ICC data compilation with recalculated carbonate chemistry data. It is hoped that all projects and databases can define best practices for archiving ocean acidification data and cooperate to avoid duplication. To this end and to meet the need for easier and more effective access to ocean acidification data, data managers of various institutions are planning to create a joint portal (“one-stop shop”) for access to global ocean acidification data by linking different data repositories (Garcia et al., 2015; Hansson et al., 2014). The Ocean Acidification International Coordination Center will organize a follow-up workshop to the last meeting in 2014 to work on the technical details of the one-stop shop data portal.

**The Supplement related to this article is available online at doi:10.5194/essdd-8-889-2015-supplement.**

*Acknowledgements.* We are grateful to all the authors for the provision of data and to the staff of PANGAEA, especially Stefanie Schumacher, for the considerable help and support it has provided to this data compilation since 2007. The pioneer work of Anne-Marin Nisumaa, data manager of the EPOCA project, is greatly appreciated. This is a contribution of the Ocean Acidification International Coordination Centre (OA-ICC), a project funded through the Peaceful Uses Initiative of the IAEA. The IAEA is grateful for the support provided to its Environment Laboratories by the Government of the Principality of Monaco. The support of the State Key Laboratory of Marine Environmental Science (Xiamen University) is also gratefully acknowledged.

## References

Cornwall, C. E. and Hurd, C. L.: Experimental design in ocean acidification research: problems and solutions, ICES J. Mar. Sci., in press, 2015.

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dickson, A. G.: The carbon dioxide system in seawater: equilibrium chemistry and measurements, in: Guide to Best Practices for Ocean Acidification Research and Data Reporting, edited by: Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J.-P., Publications Office of the European Union, Luxembourg, 17–40, 2010.
- 5 Fiorini, S., Middelburg, J. J., and Gattuso, J.-P.: Testing the effects of elevated  $p\text{CO}_2$  on coccolithophores (Prymnesiophyceae): comparison between haploid and diploid life stages, *J. Phycol.*, 47, 1281–1291, doi:10.1111/j.1529-8817.2011.01080.x, 2011.
- Garcia, H. E., Cosca, C., Kozyr, A., Mayorga, E., Chandler, C., Thomas, R. W., O'Brien, K., Appeltans, W., Hankin, S., Newton, J. A., Gutierrez, A., Gattuso, J.-P., Hansson, L., Zweng, M., and Pfell, B.: Data management strategy to improve global use of ocean acidification data and information, *Oceanography*, 28, 226–228, 2015.
- 10 Gattuso, J.-P. and Hansson, L.: Ocean acidification: background and history, in: Ocean Acidification, edited by: Gattuso, J.-P. and Hansson, L., Oxford University Press, Oxford, 1–27, 2011.
- 15 Gattuso, J.-P. and Lavigne, H.: Technical Note: Approaches and software tools to investigate the impact of ocean acidification, *Biogeosciences*, 6, 2121–2133, doi:10.5194/bg-6-2121-2009, 2009.
- Gattuso, J.-P., Gao, K., Lee, K., Rost, B., and Schulz, K. G.: Approaches and tools to manipulate the carbonate chemistry, in: Guide to Best Practices for Ocean Acidification Research and Data Reporting, edited by: Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J.-P., Publications Office of the European Union, Luxembourg, 41–52, 2010.
- 20 Gattuso, J.-P., Brewer, P. G., Hoegh-Guldberg, O., Kleypas, J. A., Pörtner, H.-O., and Schmidt, D. N.: Ocean acidification, in: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects, Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel of Climate Change, edited by: Field, C. B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., Genova, R. C., Girma, B., Kissel, E. S., Levy, A. N., MacCracken, S., Mastrandrea, P. R., and White, L. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 129–131, 2014.
- 25 Gattuso, J.-P., Epitalon, J.-M., and Lavigne, H.: seacarb: Seawater Carbonate Chemistry with R, R Package Version 3.0.8, available at: <http://CRAN.R-project.org/package=seacarb>, last access: 10 November 2015.
- 30

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Hansson, L., Appeltans, W., and Gattuso, J.-P.: Promoting international collaboration on ocean acidification data management, *EOS T. Am. Geophys. Un.*, 95, 421–422, 2014.
- Hendriks, I. E. and Duarte, C. M.: Ocean acidification: separating evidence from judgment – a reply to Dupont et al., *Estuar. Coast. Shelf Sci.*, 89, 186–190, 2010.
- 5 Hoppe, C. J. M., Langer, G., Rokitta, S. D., Wolf-Gladrow, D. A., and Rost, B.: Implications of observed inconsistencies in carbonate chemistry measurements for ocean acidification studies, *Biogeosciences*, 9, 2401–2405, doi:10.5194/bg-9-2401-2012, 2012.
- Jiang, L.-Q., O'Connor, S. A., Arzayus, K. M., and Parsons, A. R.: A metadata template for ocean acidification data, *Earth Syst. Sci. Data*, 7, 117–125, doi:10.5194/essd-7-117-2015, 2015.
- 10 Koeve, W. and Oschlies, A.: Potential impact of DOM accumulation on  $f\text{CO}_2$  and carbonate ion computations in ocean acidification experiments, *Biogeosciences*, 9, 3787–3798, doi:10.5194/bg-9-3787-2012, 2012.
- Kroeker, K. J., Kordas, R. L., Crim, R. N., and Singh, G. G.: Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms, *Ecol. Lett.*, 13, 1419–1434, 2010.
- 15 Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M., and Gattuso, J.-P.: Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming, *Glob. Chang. Biol.*, 19, 1884–1896, doi:10.1111/gcb.12179, 2013.
- 20 Liu, J., Weinbauer, M. G., Maier, C., Dai, M., and Gattuso, J.-P.: Effect of ocean acidification on microbial diversity, and on microbe-driven biogeochemistry and ecosystem functioning, *Aquat. Microb. Ecol.*, 61, 291–305, doi:10.3354/ame01446, 2010.
- McMinn, A., Müller, M. N., Martin, A., and Ryan, K. G.: The response of Antarctic sea ice algae to changes in pH and  $\text{CO}_2$ , *PLoS One*, 9, e86984, doi:10.1371/journal.pone.0086984, 2014.
- 25 Meyer, J. and Riebesell, U.: Reviews and syntheses: responses of coccolithophores to ocean acidification: a meta-analysis, *Biogeosciences*, 12, 1671–1682, 2014, <http://www.biogeosciences.net/12/1671/2014/>.
- Muller, E. B. and Nisbet, R. M.: Dynamic energy budget modeling reveals the potential of future growth and calcification for the coccolithophore *Emiliania huxleyi* in an acidified ocean, *Glob. Change Biol.*, 20, 2031–2038, doi:10.1111/gcb.12547, 2014.
- 30 Nisumaa, A.-M., Pesant, S., Bellerby, R. G. J., Delille, B., Middelburg, J. J., Orr, J. C., Riebesell, U., Tyrrell, T., Wolf-Gladrow, D., and Gattuso, J.-P.: EPOCA/EUR-OCEANS data com-

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- pilation on the biological and biogeochemical responses to ocean acidification, *Earth Syst. Sci. Data*, 2, 167–175, doi:10.5194/essd-2-167-2010, 2010.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G.-K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M.-F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, doi:10.1038/nature04095, 2005.
- Pesant, S., Alan Hook, L., Lowry, R., Moncoiffé, G., Nisumaa, A.-M., and Pfeil, B.: Safeguarding and sharing ocean acidification data, in: *Guide to Best Practices for Ocean Acidification Research and Data Reporting*, edited by: Riebesell, U., Fabry, V. J., Hansson, L., and Gattuso, J.-P., Publications Office of the European Union, Luxembourg, 243–258, 2010.
- Piwowar, H. A., Day, R. S., and Fridsma, D. B.: Sharing detailed research data is associated with increased citation rate, *PLoS One*, 2, e308, doi:10.1371/journal.pone.0000308, 2007.
- Riebesell, U. and Gattuso, J.-P.: Lessons learned from ocean acidification research, *Nat. Clim. Change*, 5, 12–14, doi:10.1038/nclimate2456, 2015.
- Riebesell, U., Gattuso, J.-P., Thingstad, T. F., and Middelburg, J. J.: Preface “Arctic ocean acidification: pelagic ecosystem and biogeochemical responses during a mesocosm study”, *Biogeosciences*, 10, 5619–5626, 2013, <http://www.biogeosciences.net/10/5619/2013/>.
- Rokitta, S. D.: Characterization of the Life-Cycle Stages of the Coccolithophore *Emiliania huxleyi* and their Responses to Ocean Acidification, PhD thesis, University of Bremen, Bremen, 2012.
- Sørensen, S.: Enzymstudien II: über die Messung und die Bedeutung der Wasserstoffionenkonzentration bei enzymatischen Prozessen, *Biochem. Z.*, 21, 131–200, 1909.
- Steinacher, M., Joos, F., Frölicher, T. L., Plattner, G.-K., and Doney, S. C.: Imminent ocean acidification in the Arctic projected with the NCAR global coupled carbon cycle-climate model, *Biogeosciences*, 6, 515–533, doi:10.5194/bg-6-515-2009, 2009.
- Traganza, E. D.: Dynamics of the carbon dioxide system on the Great Bahama Bank, *B. Mar. Sci.*, 17, 348–366, 1967.
- Vernon, H. M.: The effect of environment on the development of echinoderm larvae: an experimental inquiry into the causes of variation, *Philos. T. Roy. Soc. B*, 186, 577–632, doi:10.1038/515312a, 1895.

Zeebe, R. E. and Wolf-Gladrow, D. A.: CO<sub>2</sub> in Seawater: Equilibrium, Kinetics, Isotopes, Elsevier Oceanography Series, Amsterdam, 2001.

## ESSDD

8, 889–912, 2015

### Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures



Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion







**Figure 1.** Cumulative number of papers for which data have been included in the compilation (“archived”), papers for which data could not be obtained (“not obtained”), papers which reported less than two carbonate system parameters (“incomplete”) and papers for which the data have been lost (“lost”).

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc



Printer-friendly Version

Interactive Discussion



## Data compilation on the biological response to ocean acidification: an update


Y. Yang et al.



**Figure 2.** (a) Location of organisms and natural communities from which data has been archived and (b) distribution of locations according to latitude.

## Data compilation on the biological response to ocean acidification: an update


Y. Yang et al.



**Figure 3.** Taxonomic coverage of the papers from which data have been included in the OA-ICC data compilation, and compared to those archived prior to 2010. Categories marked with \* were not present in Nisumaa et al. (2010).

## Data compilation on the biological response to ocean acidification: an update


Y. Yang et al.



**Figure 4.** Biological processes reported in the papers from which data have been included in the OA-ICC data compilation, compared to those archived prior to 2010. Other processes studied in the papers from which data were archived before 2010 comprise variables such as shell length, width and linear extension of molluscs, bleaching, invasion, orientation, different blood cell concentration, concentration in the tissues of organisms, rate of nitrogen fixation (Nisumaa et al., 2010). The categories marked with \* were not present in Nisumaa et al. (2010) or their definition was different. “Morphology” was included in “Other processes” in Nisumaa et al. (2010) whereas it is considered as a distinct category in the present paper.

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.



**Figure 5.** Papers that have manipulated the carbonate chemistry alone and those that have manipulated the carbonate chemistry as well as other variables. CC, carbonate chemistry. The categories marked with \* were not present in Nisumaa et al. (2010).

[Title Page](#)

[Abstract](#) | [Instruments](#)

[Data Provenance & Structure](#)

[Tables](#) | [Figures](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)


[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





**Figure 6.** Countries of affiliation of first author of papers from which data were archived and papers for which data could not be obtained.

## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.

Title Page

Abstract Instruments

Data Provenance & Structure

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.



**Figure 7.** Variables of the seawater carbonate system reported in the data sets.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Data compilation on the biological response to ocean acidification: an update

Y. Yang et al.



**Figure 8.** (a) Temperature, (b) salinity, (c)  $\text{pH}_T$ , (d)  $p\text{CO}_2$ , (e)  $A_T$ , (f)  $C_T$ , (g) aragonite saturation state, (h) calcite saturation state, and (i) carbonate ion coverage in the compilation. The dashed vertical lines indicate the average open–ocean surface values, during the Last Glacial Maximum (LGM), 1766, 2007, and 2100 (see the values in Table 1.1 of Gattuso and Hansson, 2011).