



An Internally Consistent Dataset of $\delta^{13}\text{C}$ -DIC in the North Atlantic Ocean - NAC13v1

Meike Becker^{1,2}, Nils Andersen³, Helmut Erlenkeuser³, Matthew.P. Humphreys⁴,
Toste Tanhua¹, and Arne Kötzinger^{1,2}

¹GEOMAR, Helmholtz Center for Ocean Research, Kiel, Germany

²Christian Albrechts University Kiel, Kiel, Germany

³Leibniz-Laboratory for Radiometric Dating and Isotope Research, Christian Albrecht University, Kiel, Germany

⁴Ocean and Earth Science, University of Southampton, Southampton, UK

Correspondence to: Meike Becker (mbecker@geomar.de)

Abstract.

The stable carbon isotope composition of dissolved inorganic carbon ($\delta^{13}\text{C}$ -DIC) can be used to quantify fluxes within the carbon system. For example, knowing the $\delta^{13}\text{C}$ signature of the inorganic carbon pool can help to describe the exchange between ocean and atmosphere as well as the amount of anthropogenic carbon in the water column. The measurements can also be used for evaluating modeled carbon fluxes, for making basin wide estimates, studying seasonal and interannual variability or decadal trends in interior ocean biogeochemistry. For all these purposes, it is not only important to have a sufficient amount of data, but these data must also be internally consistent and of high quality.

In this study, we present a $\delta^{13}\text{C}$ -DIC dataset for the North Atlantic, which has undergone secondary quality control. The data originate from oceanographic research cruises between 1981 and 2012. During a primary quality control step based on simple range tests obviously bad data were flagged. In a second quality control step, biases between measurements from different cruises were quantified through a crossover analysis using nearby data of the respective cruises and absolute values of biased cruises were adjusted in the data product. the crossover analysis was possible for 22 of the 29 cruises in our dataset and adjustments were applied to 10 of these. The internal accuracy of this dataset is 0.017‰.

The dataset is available via CDIAC at http://cdiac.ornl.gov/oceans/ndp_096/NAC13v1.html, doi:10.3334/CDIAC/OTG.NAC13v1.

20 1 Introduction

Stable carbon isotope ratios are utilized as a tracer in several applications in marine carbon research. Particularly the stable carbon isotope ratio of dissolved inorganic carbon ($\delta^{13}\text{C}$ -DIC) can be used to enhance the understanding of carbon related processes ranging widely from the estimation of



glacial circulation changes (Curry and Oppo, 2005) to testing the performance of ecosystem models
25 (Schmittner et al., 2013). By observing the temporal development of the lightening of the inorganic carbon pool due to the uptake of CO₂ originating from the burning of ¹³C-depleted fossil fuel carbon, a phenomenon also known as oceanic ¹³C Suess effect, an estimation of the anthropogenic carbon fraction of DIC is possible (Gruber et al., 2002; Körtzinger et al., 2003; Olsen et al., 2006, 2010; Quay et al., 2007; Racapé et al., 2013). Furthermore, $\delta^{13}\text{C}$ can provide information concerning the
30 quantification of biological processes such as net community production (Quay et al., 2009). Using the stable carbon isotope signature facilitates the distinction between anthropogenic, biological and physical drivers of the carbon system.

A sample's stable carbon isotope ratio, $\delta^{13}\text{C}$ -DIC, is expressed as per mil deviation from that of the commonly used standard material Vienna Pee-Dee Belemnite (V-PDB)(Coplen, 1995).

$$35 \quad \delta^{13}\text{C} = \left(\frac{^{13}\text{R}}{^{13}\text{R}_{\text{PDB}}} - 1 \right) \cdot 10^3 \quad (1)$$

with ^{13}R being the ratio of the two stable carbon isotopes ¹³C and ¹²C in the sample.

For basin-wide carbon flux estimates, studies of seasonal variations, or interannual trends it is important to have a dataset of sufficiently high coverage both in space and time. Moreover, the dataset should be free of systematic differences between measurements carried out by different laboratories and on different cruises. However, both criteria are not easily met. Since Isotope Ratio Mass Spectroscopy (IRMS), the common method to analyze $\delta^{13}\text{C}$ -DIC data, is a very time consuming and expensive technique that cannot be performed at sea, data coverage has remained relatively poor. Therefore, several efforts have been made to assemble a dataset containing as many cruises as possible.

45 For oceanic $\delta^{13}\text{C}$ -DIC data this has been done first by Kroopnick (1985) who provided an analysis of the distribution of $\delta^{13}\text{C}$ -DIC in the world's oceans. Over the years more data was accumulated and different data collections emerged (Gruber et al., 1999; Quay et al., 2003, 2007; Schmittner et al., 2013). During recent years, databases like GLODAP (Global Ocean Data Analysis Project) and CARINA (Carbon dioxide in the Atlantic Ocean) were created for carbon-related parameters
50 (Olsen et al., 2016). These projects did not only assemble the data but also conducted a secondary quality control so that systematic biases between individual cruises could be identified and adjusted for (Tanhua et al., 2009; Velo et al., 2009; Tanhua et al., 2010a; Pierrot et al., 2010). Relative to other parameters such as total alkalinity or DIC, however, the dataset for $\delta^{13}\text{C}$ -DIC is still small and disorganized. Therefore, no secondary quality control in which deep water samples from different
55 cruises at the nearby locations, so called crossovers, are compared to each other could be carried out within these collections. Several new cruises have become available for the North Atlantic so that now the present crossover study could be performed for this area. This crossover analysis features




Figure 1. Map of all stations with $\delta^{13}\text{C}$ -DIC data used in this dataset. Data from deeper than 1500 m was available only for the stations in dark red, so only these stations were used for the crossover analysis.

29 cruises of which 22 could be compared quantitatively. Cruises without a quantitatively evaluable crossover were qualitatively related to the corrected dataset.

60 Please note, that for applying the crossover inversion routine we assume that the deep water masses (below 1500 m) are only to a negligible amount influenced by changes due to an increasing amount of anthropogenic carbon. Since the detected differences between some cruises were not consistent with a slowly increasing amount of anthropogenic carbon we think that this consistent dataset is an important step for improving the study of carbon isotope dynamics in the upper 1500 m. In regions
65 for which also the deeper water masses have been shown to contain a high amount of anthropogenic carbon, were neglected crossovers with cruises that took place long before or long after the respective cruise. We believe that no temporal trends have been removed, or created by the 2nd QC procedures employed here. However, care should be exercised for calculating C_{ant} accumulation in water below 1500 m.



Table 1. Information about sample handling and measurements for those cruises where the $\delta^{13}\text{C}$ data have not been published elsewhere.

Cruise ID	Expo code	Laboratory	Analysis period	Sample handling	PI
1	06MT19941012	1	9/2002	-	200 μL $\text{HgCl}_{2\text{sat}}$ / A. Körtzinger / H.
			12/2002	100 mL sample	Erlenkeuser
2	06MT1997-M39	1	1/1998	-	200 μL $\text{HgCl}_{2\text{sat}}$ / A. Körtzinger / H.
			2/2000	100 mL sample	Erlenkeuser
3	06MT1999-M45	1	7/2000	-	50 μL $\text{HgCl}_{2\text{sat}}$ / A. Körtzinger / H.
			6/2002	100 mL sample	Erlenkeuser
4	06MT20010507	1	12/2001	-	50 μL $\text{HgCl}_{2\text{sat}}$ / A. Körtzinger / H.
			9/2002	100 mL sample	Erlenkeuser
5	06MT20030723	1	3/2004	-	100 μL $\text{HgCl}_{2\text{sat}}$ / A. Körtzinger / H.
			10/2004	100 mL sample	Erlenkeuser
6	06MT20040311	1	1/2005	-	200 μL $\text{HgCl}_{2\text{sat}}$ / D.W.R. Wallace / H. Erlenkeuser
			10/2005	100 mL sample	H. Erlenkeuser

¹ Leibniz Laboratory for Radiometric Dating and Isotope Research, Kiel, Germany

2 Data Provenance and Structure

This dataset comprises data and metadata from 29 research cruises/campaigns from several international research groups, in total 6068 samples. Some of these consist of multiple cruises and one is a time series. For the crossover analysis, some consecutive cruises whose data were analyzed together were treated as one cruise. While the focus is on the North Atlantic, four cruises were included that also have stations in the Nordic Seas, and one cruise extends into the South Atlantic. Thereby, consistency with extended quality controlled datasets for these regions is ensured. Since only deep (> 1500 m) samples of each cruise are compared in this study, only cruises with at least one deep station could be included in this analysis.

Figure 1 shows the locations of all stations with $\delta^{13}\text{C}$ -DIC data that are part of this compilation. For cruises that have not been published elsewhere, Table 1 shows a summary of the respective sample handling, the periods during which the samples were analyzed and the responsible PI. Some cruises had $\delta^{13}\text{C}$ -DIC measurements over the entire depth range at every station, whereas others just had one or two stations with deep $\delta^{13}\text{C}$ -DIC data. Most of the cruises were conducted in the subpolar North Atlantic, while the tropical region has relative poor coverage. The temporal and latitudinal distributions of the data are displayed in Figure 2. The data was collected in the North Atlantic between 1981 and 2012, with the majority falling between 1990 and 2005. Considering the




Figure 2. The temporal distribution of the presented dataset ordered by year (left panel) and the sum of all samples at each latitudinal degree (right panel).

seasonal distribution of the data, a bias towards summer time exists, especially towards late summer. The only two cruises which took place between January and March were located south of 42°N. The uncertainty of the $\delta^{13}\text{C}$ -DIC samples analyzed by IRMS is usually reported to be between $\pm 0.12\text{\textperthousand}$ (Gruber et al., 1999) and $\pm 0.03\text{\textperthousand}$ (Quay et al., 2003).

95 The presented dataset consists of 20 columns of which the first 17 are cruise number, station, sampling number, day, month, year, latitude, longitude, maximal depth, maximal sampling depth, bottle number, cast number, temperature, salinity, depth, ctd salinity and pressure. Column 18 contains the adjusted $\delta^{13}\text{C}$ -DIC data, column 19 a quality flag(C13f) and column 20 the QC-flag (C13qc, see Table 2). For bad data the quality flag was set to 'not measured' and therefore column 19 has only
100 two entries (2: good, 9: not measured). Cruises that could be quantitatively compared to each other by the 2nd QC have a '1' in the QC-flag. All others are flagged with '0'.

Additional parameters to most of the cruises can be found in either GLODAPv2 or CARINA. Only the most recent cruises 35TH20060521 and 74DI20120731 are not included in these datasets, but the individual cruise files can be found on CDIAC. The OMEX1NA data are only part of Carina.

105 The respective cruise numbers in GLODAPv2 and CARINA of the cruises shown in the NAC13v1-dataset can be found in the documentation.



Table 2. NAC13v1 data set parameter list, column names used in the data product and the respective units.

parameter	data product parameter name	data product flag name	unit
NAC13v1 cruise number	cruiseno		
Station	station		
Sample number	nosamp		
Day	day		
Month	month		
Year	year		
Latitude	latitude		°N
Longitude	longitude		°E
Bottom depth	maxdepth		m
Pressure of the deepest sample	maxsampdepth		dbar
Bottle number	bottle		
Cast number	cast		
Temperature	temperature		°C
Salinity	salinity		
Depth	depth		m
CTD salinity	ctdsal		
Pressure	pressure		dbar
$\delta^{13}\text{C}$ -DIC	C13	C13f, C13qc	‰

110 3 Computational Analysis

115 In order to derive an internally consistent set of $\delta^{13}\text{C}$ -DIC data in the North Atlantic all publicly available data in this area were assembled and quality controlled (QC) in two steps. At first, a primary QC was performed in order to identify obviously erroneous data, such as wrong positions, time stamps and depths. Also outliers were identified and then flagged by comparing the profiles of each cruise internally. After that, the secondary QC procedure was conducted employing a crossover analysis as described by Tanhua et al. (2010b). This MATLAB based software package compares two cruises at a time, searches for nearby stations, so-called 'crossovers', and calculates differences between all crossovers of the two cruises as additive offsets with the unit ‰. As criterion for identifying crossovers a maximum of 180 nm (3° of latitude) distance between stations was used. From 120 these crossovers, the $\delta^{13}\text{C}$ -DIC data collected deeper than 1500 m was compared on equal potential density. Based on the resulting offsets and standard deviations determined for each of these crossovers a suggestion for a possible adjustment was made. This suggestion was obtained by an



inversion routine using a Weighted Least-Square (WLSQ) and a Weighted Damped Least Square (WDLSQ) model as described by Johnson et al. (2001). The cruise 33MW19930704-1 covers a long
125 distance and is assumed to have high quality data. Therefore this cruise was selected as core cruise and weighted higher than the other cruises. Unfortunately this was the only cruise meeting these two criteria. Several cruises from different years were in good agreement with the core cruise while the other cruises were adjusted towards it. Choosing the appropriate distance criterion for crossover locations is always a compromise between including as many statistically relevant crossovers as possible by selecting a large enough radius on the one hand and trying to have only crossovers between
130 stations that share similar oceanographic characteristics on the other hand. However, reducing the crossover distance to 120 nm reduced the amount of crossovers and the number of cruises that could be quantitatively compared to each other but did not significantly change the suggested magnitude of adjustments of the remaining cruises. Therefore, the $3^\circ \times 3^\circ$ criterion was used instead. For some
135 crossovers in highly variable regions with deep water formation, such as the Labrador Sea and the Nordic Seas, the standard deviation was decreased significantly by restricting the comparison depths to >2000 m. Generally, offsets from crossovers in these highly variable regions, from cruises with a relatively poor data precision or with just a few deep samples were considered in the model with less influence, by weighting the offsets with their uncertainty. In Figure 3 all crossovers between the
140 cruises 06MT20030723 and 33MW19930704-1 are shown as an example, both for the uncorrected as well as for the corrected dataset. All crossovers from the adjusted and the unadjusted dataset can be found at: http://cdiac.ornl.gov/oceans/ndp_096/NAC13v1.html.

Whether an adjustment was applied to the data was decided somewhat subjectively in each case based on a combination of the shape and distribution of individual crossover differences and the
145 suggestions given by the inversion routine with knowledge about the sampling region. After applying the adjustments, the inversion was conducted again and it was checked whether or not the adjustment improved the overall consistency within the entire dataset. Temporal changes of the deep water masses were only considered in this step of the routine when comparing the suggested corrections and the corresponding crossover offsets between cruises in areas where also the deep water $\delta^{13}\text{C}$ -
150 DIC was expected to change over time. In order to get a quantitative description of the internal consistency of the final dataset, a weighted mean using the respective offsets of all crossovers and their standard deviation was calculated (Tanhua et al., 2010a).

$$\text{WM} = \frac{\sum_{i=1}^L D(i)/(\sigma(i))^2}{\sum_{i=1}^L 1/(\sigma(i))^2} \quad (2)$$

L refers to the total number of crossover, D to the respective offset of all crossover and σ is their
155 standard deviation.




Figure 3. Crossovers between the cruises 06MT20030723 and the core cruise 33MW19930704-1. The left hand plot shows the original and the right hand plot the adjusted data. In both cases the distribution of the $\delta^{13}\text{C}$ -DIC on equal density surfaces (left hand side) as well as the mean offset between both cruises (right hand side) is shown. The cruise 06MT20030723 was adjusted by -0.20 ‰ .

4 Adjustments

The data of all cruises as well as locations are shown in Figure 4. The offsets, as well as the corrections suggested by the WDLSQ inversion routine, and the final adjustments are listed in Table 3.

In Figure 5 the results of the WDLSQ inversion are shown before and after the adjustments were applied. Some cruises show quite big deviations from the core cruise. However, we do not know the reason for these biases. Besides the actual sample analysis in the laboratory, also different sampling routines on board the ship, insufficient poisoning and the sample storage time can cause these biases.

A detailed overview of the offset of each crossover in the original as well as the adjusted dataset is given in Table 4 in the supplementary information. Moreover, the evidence for our decision will be presented for each cruise.

4.1 06MT19941012, cruise #1

This cruise on the German R/V Meteor is also known as M30-2 (Körtzinger et al., 1998). The inversion suggested a correction of -0.07 ‰ . The mean offset of all crossovers is 0.11 ‰ too high. Based on the crossover with the core cruise an adjustment of -0.07 ‰ was applied.



Table 3. Overview of all cruises in this dataset. The data of some cruises were combined for the analysis. For more information, please see the detailed description in the 'Adjustments' section. Both, the mean offsets and the corrections suggested by the WDLSQ inversion are shown for the original and the adjusted dataset. In the last column the applied adjustments are displayed. NC indicates that these cruises were not considered in the inversion since they had no statistically significant crossover and the core cruise is marked with C. Cruises with insufficient quality data are denoted 'poor' and not included in the further analysis.

cruise ID	Expo code	Calculated offset		Suggested adjustments		Final adjustments / %
		not adjusted / %	adjusted / %	WDLSQ / %	WDLSQ (adj) / %	
1	06MT19941012	0.11	-0.02	-0.07(±0.10)	-0.01(±0.02)	-0.07
2	06MT1997-M39	-0.02	0.02	0.01(±0.14)	0.00(±0.01)	0
3	06MT1999-M45	0.16	-0.01	-0.14(±0.09)	0.00(±0.01)	-0.15
4	06MT20010507	0.16	0.00	-0.24(±0.10)	0.00(±0.01)	-0.30
5	06MT20030723	0.14	0.03	-0.15(±0.09)	0.00(±0.01)	-0.20
6	06MT20040311	-0.14	-0.02	0.10(±0.09)	0.01(±0.01)	0.10
7	316N19970717	0.17	0.02	-0.06(±0.17)	-0.01(±0.01)	-0.05
8	316N19970815					NC
9	316N20030922					NC
10	316N20031023					NC
11	33RO19980123					NC
12	33MW19910711	-0.02	-0.02	0.00(±0.01)	0.00(±0.01)	0
13	33MW19930704-1	-0.05	0.01	0.00(±0.01)	0.00(±0.01)	C
14	35TH20020611					NC
15	35TH20060521	-0.39	-0.02	0.24(±0.21)	-0.03(±0.05)	0.25
16	58JH19920712					NC
17	58JH19940723					NC
18	64TR19900417					poor
19	74DI20120731	-0.33	-0.13	0.13(±0.28)	0.12(±0.12)	0
20	OMEXINA	-0.14	-0.03	0.03(±0.13)	0.02(±0.02)	0
21	316N19810401	-0.06	0.03	-0.03(±0.10)	-0.01(±0.03)	0




Figure 4. Adjusted $\delta^{13}\text{C}$ -DIC profiles and locations of each cruise. The green profiles represent the data of the specific cruise whereas the gray dots show all profiles in the dataset.

continued on next page




Figure 4.

continued from previous page.

Adjusted $\delta^{13}\text{C}$ -DIC profiles and locations of each cruise. The green profiles represent the data of the specific cruise whereas the gray dots show all profiles in the dataset.




Figure 5. The results of the WDLSQ based inversion routine for the original (blue circles) and the adjusted dataset (red stars). The cruises are plotted at the time the data was collected vs. the suggested correction.

170 4.2 06MT19970515, 06MT19970707 and 6MT19970815, here referred to as
06MT1997-M39, cruise #2

These cruises are also known as M39 cruises with three legs of $\delta^{13}\text{C}$ -DIC sampling (M39-2, M39-4, M39-5) (Körtzinger et al., 1999; Thomas and Ittekkot, 2001). Since each leg of this cruise had only



175 a few stations with $\delta^{13}\text{C}$ -DIC samples, and all these samples were analyzed together, these cruises were summarized for the crossover study. Both, the inversion routine and the single crossover with the adjusted cruises show no evidence for an offset.

4.3 06MT19990711 and 06MT19990813, here referred to as 06MT1999-M45, cruise #3

180 These cruises are also known as M45-2 and M45-3 (Friis et al., 2005). Since both were analyzed together, they were summarized for this crossover study. The inversion suggested a correction of -0.15 ‰ and the mean offset of all crossovers was 0.16 ‰ too high. After applying this adjustment and comparing this cruise to the adjusted dataset, the inversion routine still suggested a small correction. Therefore, an adjustment of -0.20 ‰ was applied.

4.4 06MT20010507, cruise #4

185 This cruise is also known as M50-1 (Friis et al., 2007). The inversion routine suggested a correction of -0.24 ‰, whereas the mean offset was 0.16 ‰ too high. Based on the southern crossover with cruise 06MT20040311 and 316N19970717 an adjustment of -0.30 ‰ was applied.

4.5 06MT20030723, cruise #5

190 This cruise is also known as M59-2 (Friis et al., 2007). The correction suggested by the inversion routine is -0.15 ‰ which matches with the positive offsets of the crossovers, except of those with 33TH20060521. Based on the crossover with the core cruise, an adjustment of -0.15 ‰ was applied.

4.6 06MT20040311, cruise #6

This cruise is also known as M60-5 (Tanhua et al., 2007). The inversion routine indicates that the $\delta^{13}\text{C}$ -DIC data of this cruise is 0.10 ‰ too low. Additionally, the mean offset shows that this data is too low. An adjustment of +0.10 ‰ was applied.

195 4.7 316N19970717, cruise #7 and 316N19970815, cruise #8

These cruises followed the WOCE/GO-Ship standard lines A20 and A22 (Johnson et al., 2003). The inversion suggests a correction of -0.06 ‰ for 316N19970717. It shows one crossover with cruise 06MT20040311 in which a significant positive offset is still visible after cruise 06MT20040311 was corrected. Therefore, an adjustment of -0.05 ‰ was applied for cruise 316N19970717. The cruise 200 316N19970815 does not show a statistically significant crossover.

4.8 316N20030922, cruise #9, and 316N20031023, cruise #10

These cruises, that took place in the tropical western Atlantic, following the A20 and A22 lines, have only one deep station each (Feely et al., 2008). The crossovers of these stations with both, the ad-



justed data of cruise 06MT20040311 and cruise 316N19970717 show a good agreement, suggesting
205 that no adjustment should be applied.

4.9 33RO19980123, cruise #11

This cruise (Lee et al., 2003) has one statistically insignificant crossover with the cruise 06MT20040311 and one with cruise 33MW19930704-1. Both seem to be in good agreement, suggesting that no adjustment should be applied.

210 4.10 33MW19910711, cruise #12, and 33MW19930704-1, cruise #13

The cruise 33MW19930704-1 was considered as core cruise in the present analysis (Forde et al., 1996). The cruise 33MW19910711 extends into the south Atlantic and its crossover with cruise 13 shows no need for an adjustment.

4.11 35TH20020611, cruise #14, and 35TH20060521, cruise #15

215 The latter of these two cruises has a few quantitative crossovers, that show a high offset of $-0.39\text{\textperthousand}$. Furthermore, the inversion suggests a correction of $0.24\text{\textperthousand}$. The high variability of the sampling area south of Iceland, as well as an increasing lightening of the deep water carbon pool over time don't give an adequate explanation for this large deviation and, therefore, an adjustment of $-0.25\text{\textperthousand}$ was applied. The cruise 35TH20020611 shows just a few qualitatively analyzable crossovers, that show
220 a lighter carbon pool compared to earlier cruises and a heavier one compared to the original data of cruise 35TH20060521 (Racapé et al., 2013). After adjusting this cruise, both cruises, which were analyzed in the same laboratory, are not in good agreement anymore which suggests that the earlier cruise also has too low isotope values. However, in the absence of a statistically significant crossover no adjustment was applied to this cruise.

225 4.12 58JH19920712, cruise #16, and 58JH19940723, cruise #17

These two cruises took place in a highly variable area (Gislefoss et al., 1995). No statistically relevant crossover exists but the data are in good agreement with the core cruise and the other adjusted cruises in that area.

4.13 64TR19900417, cruise #18

230 This cruise shows extreme scatter compared to all other cruises and, therefore, was not included into the adjusted product (Rommets et al., 1991). When comparing crossover stations this cruise shows a mean offset to other cruises of about $-1.2\text{\textperthousand}$.



4.14 74DI20120731, cruise #19

Both the inversion and the offset mean of the crossover suggest a correction of +0.13 ‰ for the cruise
235 (Humphreys et al., 2015). This most recent cruise took place near the Scotland-Iceland ridge where the deep water masses cannot be assumed to be constant over time. All crossovers indicate a lower δ¹³C-DIC of this cruise when comparing it with the others which is consistent with an increased amount of anthropogenic carbon. Therefore, no adjustment was applied.

4.15 OMEX1NA, cruise #20

240 During the OMEX1 project in the North Atlantic δ¹³C-DIC samples were taken in January 1994 (Wollast and Chou, 2001). The data is in good agreement with the other cruises in this area and no adjustment was applied.

4.16 316N19810401, cruise #21

245 The cruises 316N19810401, 316N19810416, 316N19810516, 316N19810619, 316N19810721, 316N19810821 and 316N19810923 are combined and usually named Transient Tracers in the Oceans North Atlantic Study (TTO-NAS) (Brewer et al., 1986). The inversion does not suggest any correction for this dataset.

5 Conclusions

250 The finalized, quality controlled dataset of δ¹³C-DIC presented here consists of 22 cruises (some of which consists of multiple legs) that have been quantitatively compared to each other and form an internally consistent dataset. Seven cruises could not be quantitatively compared to the other cruises due to a lack of crossovers and / or deep δ¹³C-DIC data. The internal consistency of the adjusted dataset was calculated to be 0.017‰ based on Equation 2.

255 The database is available at CDIAC via http://cdiac.ornl.gov/oceans/ndp_096/NAC13v1.html, doi:10.3334/CDIAC/OTG.NAC13v1.

6 Acknowledgment

260 We would like to thank all the people, both researchers as well as captains and crews, who spend time at sea and in the lab collecting and measuring the samples and preparing the data that is presented here and the PIs for sharing it. The measurements of the majority of previously not public δ¹³C data was supported by the Deutsche Forschungsgemeinschaft (DFG) through SFB460 and this work was funded by the Future Ocean Excellence Cluster Project CP1140. We also thank Aley Kozyr from CDIAC for preparing the NAC13v1-website on CDIAC.



Table 4. This table shows an overview of all crossovers. The symbol \bullet in each row divides the table into a triangle in the upper right and one in the lower left of the table. In the upper right corner for each crossover the offsets of the original dataset are listed. In the lower left corner the remaining offsets in the adjusted dataset are shown. Not statistically relevant crossovers are displayed with $>$ and $=$ indicating the tendency of not significant crossover. Please note that the offsets shown in the table result from $\delta^{13}\text{C}_{\text{column}} - \delta^{13}\text{C}_{\text{row}}$. All offsets are given in ‰.



References

Brewer, P., Takahashi, T., and Williams, R.: Transient tracers in the oceans (TTO)-Hydrographic data and carbon
265 dioxide systems with revised carbon chemistry data, Tech. rep., Oak Ridge National Lab., TN (United States),
1986.

Coplen, T.: Reporting of stable hydrogen, carbon, and oxygen isotopic abundances, *Geothermics*, 24, 707 –
712, doi:[http://dx.doi.org/10.1016/0375-6505\(95\)00024-0](http://dx.doi.org/10.1016/0375-6505(95)00024-0), 1995.

Curry, W. B. and Oppo, D. W.: Glacial water mass geometry and the distribution of $\delta^{13}\text{C}$ of ΣCO_2 in the
270 western Atlantic Ocean, *Paleoceanography*, 20, doi:[10.1029/2004PA001021](https://doi.org/10.1029/2004PA001021), 2005.

Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., and Hales, B.: Evidence for upwelling of cor-
rosive" acidified" water onto the continental shelf, *Science*, 320, 1490–1492, doi:[10.1126/science.1155676](https://doi.org/10.1126/science.1155676),
2008.

Forde, E. B., Hendee, J. C., and Wanninkhof, R. H.: Hydrograohic, Carbon Dioxide, Nutrient, and Produc-
275 tivity measurements from the South Atlantic during July and August of 1991, Tech. rep., NOAA/Atlantic
Oceanographic and Meteorological Laboratories Miami, Florida 33149, 1996.

Friis, K., Körtzinger, A., Pätsch, J., and Wallace, D. W.: On the temporal increase of anthropogenic CO_2 in the
subpolar North Atlantic, *Deep-Sea Res. Pt I*, 52, 681–698, doi:[10.1016/j.dsr.2004.11.017](https://doi.org/10.1016/j.dsr.2004.11.017), 2005.

Friis, K., Najjar, R., Follows, M., Dutkiewicz, S., Körtzinger, A., and Johnson, K.: Dissolution of calcium
280 carbonate: observations and model results in the subpolar North Atlantic, *Biogeosciences*, 4, 205–213,
doi:[10.5194/bg-4-205-2007](https://doi.org/10.5194/bg-4-205-2007), 2007.

Gislefoss, J., Nydal, R., Skjelvan, I., Nes, A., Østerhus, S., Holmén, K., Jull, T., and Sonninen, E.: Carbon
profiles in the Nordic Seas, Tech. rep., Radiological Dating Laboratory, Trondheim-NTH, 1995.

Gruber, N., Keeling, C. D., Bacastow, R. B., Guenther, P. R., Lueker, T. J., Wahlen, M., Meijer, H. A. J., Mook,
285 W. G., and Stocker, T. F.: Spatiotemporal patterns of carbon-13 in the global surface oceans and the oceanic
suess effect, *Global Biogeochem. Cycles*, 13, 307–335, doi:[10.1029/1999GB900019](https://doi.org/10.1029/1999GB900019), 1999.

Gruber, N., Keeling, C. D., and Bates, N. R.: Interannual Variability in the North Atlantic Ocean Carbon Sink,
Science, 298, 2374–2378, doi:[10.1126/science.1077077](https://doi.org/10.1126/science.1077077), 2002.

Humphreys, M. P., Achterberg, E. P., Griffiths, A. M., McDonald, A., and Boyce, A. J.: Measurements of the
290 stable carbon isotope composition of dissolved inorganic carbon in the northeastern Atlantic and Nordic Seas
during summer 2012, *Earth Sys. Sci. Data*, 7, 127–135, doi:[10.5194/essd-7-127-2015](https://doi.org/10.5194/essd-7-127-2015), 2015.

Johnson, G. C., Robbins, P. E., and Hufford, G. E.: Systematic Adjustments of Hydrographic
Sections for Internal Consistency, *J. Atmos. Ocean. Tech.*, 18, 1234–1244, doi:[10.1175/1520-0426\(2001\)018<1234:SAOHSF>2.0.CO;2](https://doi.org/10.1175/1520-0426(2001)018<1234:SAOHSF>2.0.CO;2), 2001.

295 Johnson, K., Key, R., Millero, F., Sabine, C., Wallace, D., Winn, C., Arlen, L., Erickson, K., Friis, K., Galanter,
M., Goen, J., Rotter, R., Thomas, C., Wilke, R., Takahashi, T., and Sutherland, S.: Carbon Dioxide, Hydro-
graphic, and Chemical Data Obtained During the R/V Knorr Cruises in the North Atlantic Ocean on WOCE
Sections AR24 (November 2 - December 5, 1996) and A24, A20, and A22 (May 30 - September 3, 1997),
Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy,
300 Oak Ridge, Tennessee, doi:[10.3334/CDIAC/otg.ndp082](https://doi.org/10.3334/CDIAC/otg.ndp082), 2003.

Körtzinger, A., Mintrop, L., and Duinker, J. C.: On the penetration of anthropogenic CO_2 into the North Atlantic
Ocean, *J. Geophys. Res.*, 103, 18 681–18 689, doi:[10.1029/98JC01737](https://doi.org/10.1029/98JC01737), 1998.



Körtzinger, A., Rhein, M., and Mintrop, L.: Anthropogenic CO₂ and CFCs in the North Atlantic Ocean-A comparison of man-made tracers, *Geophysical Research Letters*, 26, 2065–2068, doi:10.1029/1999GL900432, 305 1999.

Körtzinger, A., Quay, P. D., and Sonnerup, R. E.: Relationship between anthropogenic CO₂ and the ¹³C Suess effect in the North Atlantic Ocean, *Global Biogeochemical Cycles*, 17, 5–15–20, doi:10.1029/2001GB001427, 1005, 2003.

Kroopnick, P. M.: The distribution of ¹³C of Σ CO₂ in the world oceans, *Deep-Sea Res.*, 32, 57 – 84, 310 doi:10.1016/0198-0149(85)90017-2, 1985.

Lee, K., Choi, S.-D., Park, G.-H., Wanninkhof, R., Peng, T.-H., Key, R., Sabine, C., Feely, R., Bullister, J., Millero, F., et al.: An updated anthropogenic CO₂ inventory in the Atlantic Ocean, *Global Biogeochem. Cycles*, 17, doi:10.1029/2003GB002067, 2003.

Olsen, A., Omar, A. M., Bellerby, R. G., Johannessen, T., Ninnemann, U., Brown, K. R., Olsson, K. A., Olafsson, J., Nondal, G., Kivimäe, C., et al.: Magnitude and origin of the anthropogenic CO₂ increase and ¹³C Suess effect in the Nordic seas since 1981, *Global Biogeochem. Cycles*, 20, doi:10.1029/2005GB002669, 315 2006.

Olsen, A., Omar, A. M., Jeansson, E., Anderson, L. G., and Bellerby, R. G. J.: Nordic seas transit time distributions and anthropogenic CO₂, *J. Geophys. Res. Oceans*, 115, doi:10.1029/2009JC005488, c05005, 2010.

Olsen, A., Key, R. M., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F., and Suzuki, T.: An internally consistent data product for the world ocean: the Global Ocean Data Analysis Project, version 2 (GLODAPv2), *Earth System Science Data Discussions*, 2016, 1–78, doi:10.5194/essd-2015-42, 2016.

Pierrot, D., Brown, P., Van Heuven, S., Tanhua, T., Schuster, U., Wanninkhof, R., and Key, R. M.: CARINA 325 TCO₂ data in the Atlantic Ocean, *Earth Syst. Sci. Data*, 2, 177–187, doi:10.5194/essd-2-177-2010, 2010.

Quay, P. D., Sonnerup, R., Westby, T., Stutsman, J., and McNichol, A.: Changes in the ¹³C/¹²C of dissolved inorganic carbon in the ocean as a tracer of anthropogenic CO₂ uptake, *Global Biogeochem. Cycles*, 17, 1–20, doi:10.1029/2001GB001817, 2003.

Quay, P. D., Sonnerup, R., Stutsman, J., Maurer, J., Körtzinger, A., Padin, X. A., and Robinson, C.: Anthropogenic CO₂ accumulation rates in the North Atlantic Ocean from changes in the ¹³C/¹²C of dissolved inorganic carbon, *Global Biogeochem. Cycles*, 21, 1–15, doi:10.1029/2006GB002761, 2007.

Quay, P. D., Stutsman, J., Feely, R. A., and Juraneck, L. W.: Net community production rates across the subtropical and equatorial Pacific Ocean estimated from air-sea $\delta^{13}\text{C}$ disequilibrium, *Global Biogeochem. Cycles*, 23, 1–15, doi:10.1029/2008GB003193, 2009.

Racapé, V., Pierre, C., Metzl, N., Monaco, C. L., Reverdin, G., Olsen, A., Morin, P., Vázquez-Rodríguez, M., Ríos, A. F., and Pérez, F. F.: Anthropogenic carbon changes in the Irminger Basin (1981–2006): Coupling $\delta^{13}\text{C}$ DIC and DIC observations, *J. Marine Syst.*, 126, 24–32, doi:10.1016/j.jmarsys.2012.12.005, 2013.

Rommets, J., Dapper, R., and Fransz, H.: JGOFS North Atlantic Pilot Project - R.V.Tyro Leg 3, Upper Ocean Processes, NIOZ Data Rpt 1991-3, 1991.

Schmittner, A., Gruber, N., Mix, A., Key, R., Tagliabue, A., and Westberry, T.: Biology and air-sea gas exchange controls on the distribution of carbon isotope ratios ($\delta^{13}\text{C}$) in the ocean, *Biogeosciences*, 10, 5793–5816, doi:10.5194/bg-10-5793-2013, 2013.



345 Tanhua, T., Körtzinger, A., Friis, K., Waugh, D. W., and Wallace, D. W.: An estimate of anthropogenic CO₂ inventory from decadal changes in oceanic carbon content, *P. Natl. Acad. Sci.*, 104, 3037–3042, doi:10.1073/pnas.0606574104, 2007.

Tanhua, T., Brown, P. J., and Key, R. M.: CARINA: nutrient data in the Atlantic Ocean, *Earth Syst. Sci. Data*, 1, 7–24, doi:10.5194/essd-1-7-2009, 2009.

350 Tanhua, T., Steinfeldt, R., Key, R. M., Brown, P., Gruber, N., Wanninkhof, R., Perez, F., Körtzinger, A., Velo, A., Schuster, U., van Heuven, S., Bullister, J. L., Stendardo, I., Hoppema, M., Olsen, A., Kozyr, A., Pierrot, D., Schirnick, C., and Wallace, D. W. R.: Atlantic Ocean CARINA data: overview and salinity adjustments, *Earth Syst. Sci. Data*, 2, 17–34, doi:10.5194/essd-2-17-2010, 2010a.

Tanhua, T., van Heuven, S., Key, R. M., Velo, A., Olsen, A., and Schirnick, C.: Quality control procedures and methods of the CARINA database, *Earth Syst. Sci. Data*, 2, 35–49, doi:10.5194/essd-2-35-2010, 2010b.

355 Thomas, H. and Ittekkot, V.: Determination of anthropogenic CO₂ in the North Atlantic Ocean using water mass ages and CO₂ equilibrium chemistry, *J. Marine Syst.*, 27, 325–336, 2001.

Velo, A., Perez, F. F., Brown, P., Tanhua, T., Schuster, U., and Key, R. M.: CARINA alkalinity data in the Atlantic Ocean, *Earth Syst. Sci. Data*, 1, 45–61, doi:10.5194/essd-1-45-2009, 2009.

Wollast, R. and Chou, L.: The carbon cycle at the ocean margin in the northern Gulf of Biscay, *Deep-Sea Res. Pt II*, 48, 3265 – 3293, doi:10.1016/S0967-0645(01)00040-6, 2001.