Earth Syst. Sci. Data Discuss., https://doi.org/10.5194/essd-2017-126-AC4, 2018 © Author(s) 2018. This work is distributed under the Creative Commons Attribution 4.0 License.



ESSDD

Interactive comment

Interactive comment on "Central-Pacific surface meteorology from the 2016 El Niño Rapid Response (ENRR) field campaign" by Leslie M. Hartten et al.

Leslie M. Hartten et al.

leslie.m.hartten@noaa.gov

Received and published: 25 May 2018

Here is the figure meant to accompany our "Reply to Anonymous Referee #2", which was not included in our reply due to software issues.

Additional information about the data plotted in Figure 1 (originally intended for the caption): Periods during which approximately 25 mm or more rainfall accumulated during a short time are highlighted as "heavy". The ship's latitude and longitude ranges during the legs are as follows: Leg 2, $(2.02^{\circ}-5.24^{\circ}N, 80.50^{\circ}-90.65^{\circ}E)$; Leg 3, $(2.03^{\circ}S-4.83^{\circ}N, 80.49^{\circ}-90.79^{\circ}E)$; Leg 4, $(0.07^{\circ}S-3.53^{\circ}N, 80.5^{\circ}-87.91^{\circ}E)$. Ship data were collected and post-processed into one-minute values by Jim Edson (University of Con-

Printer-friendly version

Discussion paper



necticut), Chris Fairall (NOAA/ESRL/Physical Sciences Division), and Simon deSzoeke (Oregon State University).

Interactive comment on Earth Syst. Sci. Data Discuss., https://doi.org/10.5194/essd-2017-126, 2017.

ESSDD

Interactive comment

Printer-friendly version

Discussion paper



ESSDD

Interactive comment




Fig. 1. Ten-minute values of relative humidity at 10 m and leg-accumulated rainfall from three cruise legs during the DYNAMO campaign. See text for details.

Printer-friendly version

Discussion paper

