Growth characteristics of natural and planted Dahurian larch in northeast China

Bingrui Jia[1*] and Guangsheng Zhou[2*]

[1]*State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China,*

[2]*Chinese Academy of Meteorological Sciences, Beijing 100081, China,*

[*]Corresponding author:

Bingrui Jia, Guangsheng Zhou

Institute of Botany, Chinese Academy of Sciences,

20 Nanxincun, Xiangshan, Haidian District, Beijing 100093, China

E-mail: jiabingrui@ibcas.ac.cn, gszhou@ibcas.ac.cn

Tel: 86-10-62836289, Fax: 86-10-82595962

Abstract. Dahurian larch (*Larix gmelinii* Rupr.) is the dominant species in both natural and planted forests in northeast China, which situated in the southernmost part of the global boreal forest biome and undergoing the greatest climatically induced changes. Published studies (1965–2015) on tree aboveground growth of *Larix gmelinii* forests in northeast China were collected in this study, critically reviewed, and a comprehensive growth data set was developed from 123 sites, which distributed between 40.85 ° N and 53.47 ° N in latitude, between 118.20 ° E and 133.70 ° E in longitude, between 130 m and 1260 m in altitude. The data set was composed of 776 entries, including growth data (mean tree height, mean DBH, mean tree volume and/or stand volume) and the associated information, i.e., geographical location (latitude, longitude, altitude, aspect and slope), climate (mean annual temperature (MAT) and mean annual precipitation (MAP)), stand description (origin, stand age, stand density and canopy density), and sample regime (observing year, plot area and number). It

would provide quantitative references for plantation management practices and boreal forest growth prediction under future climate change. The data set is freely available for noncommercial scientific applications, and the DOI for the data is https://doi.org/10.1594/PANGAEA.880984.

1 Introduction

Boreal forests, the second largest biome in the world, cover about one-third of the Earth's forest area (Achard et al., 2006; Keenan et al., 2015). Dahurian larch (*Larix gmelinii* Rupr.) is a dominant tree species in Chinese boreal forest, which is distributed primarily in northeast China. *Larix gmelinii* forest is also the predominant timber source in China, occupying 55% of Chinese boreal forest area and accounting for 75% of Chinese boreal forest volume (Xu, 1998; Zhou et al., 2002). *Larix gmelinii* forest is situated in the southernmost part of the global boreal forest biome (Shugart et al., 1992) and undergoing the greatest climatically induced changes. Thus understanding the growth characteristics of *Larix gmelinii* forest in China are of critical need for management and prediction under future climate change.

With increased greenhouse effective and climate warming in recent years, forest carbon sink has being paid~~is payed~~ more and more attention by the world (Bastin et al., 2017). Forestation is the main measure to offset the greenhouse gas emission and increase carbon sink. China has the largest area of forest plantations in the world, approximately 79 million ha or one-fourth of world total (FAO, 2015; Payn et al., 2015). The forest cover showed an increasing trend through reforestation in northeast China (Achard et al., 2006). *Larix gmelinii* is an important fast-growing and cold-tolerant tree species used in forestation in northeast China. *Larix gmelinii* is usually planted after fire or logging. The growth rates of *Larix gmelinii* plantations are important indexes in the assessment of forest recovery processes and carbon sequestration potentials, which could supply strategies for post-fire or

post-harvest management. The data set can provide basis for evaluating and predicting the carbon sequestration and its potential of the forestation activities.

Relating the easily measured variables (e.g. diameter at breast height or DBH, tree height~~tree height, diameter~~) to other structural and functional characteristics, is the most common and reliable method for estimating forest biomass, net primary production and biogeochemical budgets (Luo, 1996; Fang et al., 2001). For larch forests in northeast China, synthesis studies mainly focused on biomass and net primary production with ~~a small quantity of~~ increasing samples in recent decade, for example, N=28 (Luo, 1996), N=17 (Wang et al., 2001a), N=18 (Wang et al., 2001b; Zhou et al., 2002), N=36 (Wang et al., 2005), N=83 (Wang et al., 2008), N=50 (Zeng, 2015), N=150 (Zeng et al., 2017). However, large numbers of growth measurements (e.g. age, DBH, tree height~~, DBH~~, volume) have scarcely been studied systematically at the large scale. Therefore, a comprehensive growth data set (N=776) of Dahurian larch in northeast China was developed in this paper.

2 Data and methods

2.1 Research origin descriptors

(1) Identity: Growth data set of natural and planted Dahurian larch in northeast China, version 1.0

(2) Originators:

Bingrui Jia, State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China;

Guangsheng Zhou, Chinese Academy of Meteorological Sciences, Beijing 100081, China.

(3) Period of Study: From January 1965 to December 2015.

(4) Objectives: We conducted a complete literature ~~and book~~ review of published studies on age, DBH, tree height, ~~DBH,~~ and/or volume in *Larix gmelinii* forests including natural and

planted forests in order to develop a growth data set. The data set can be used to analyze growth characteristics of *Larix gmelinii* forests and assess their potential productivity in future climate warming.

2.2 Site description

(1) Site type: Data were derived from 123 study sites in northeastern China. This region includes Heilongjiang, Liaoning, Jilin provinces and the eastern part of Inner Mongolia Autonomous Region (Fig. 1).



Figure 1. Study sites of *Larix gmelinii* across northeastern China, including Heilongjiang, Liaoning, Jilin provinces and the eastern part of Inner Mongolia Autonomous Region. Open circles represent natural forests and dots represent planted forests.

(2) Habitat: *Larix gmelinii* forest is naturally distributed in Great Xing'an Mountains of northeastern China. *Larix gmelinii* was usually planted in barren hills, post-fire or

post-harvest area in northeast China.

(3) Climate: The climate in northeast China is controlled by the high latitude East Asia monsoon, changing from cool temperate to temperate zones from north to south, and from semi-arid to humid zones from west to east. Mean annual temperature (MAT) for these sites ranged from -6.1 to 7.0 ℃, and mean annual precipitation (MAP) from 355 to 926 mm.

2.3 Data sources

Published studies (1965-2015) were collected from available online full-text databases, including China Knowledge Resource Integrated Database (http://www.cnki.net/), China Science and Technology Journal Database (http://www.cqvip.com/), Wanfang Data Knowledge Service Platform (http://www.wanfangdata.com/), ScienceDirect (http://www.sciencedirect.com/), ISI Web of Science (http://isiknowledge.com/) and Springer Link (http://link.springer.com/). The different combination of the terms "Dahurian larch (or *Larix gmelinii*)" with "height", "diameter at breast height (or DBH)", "tree volume" ~~and~~ or "stand volume" were searched in full text. Meanwhile, we also looked up the related books (e.g. Ma, 1992; Wang, 1992; Zhou, 1994; Yang, 2009). We attempted to compile a complete growth data set of natural and planted Dahurian larch in northeastern part of China (between 40.85 °N and 53.47 °N; between 118.20 ℉ and 133.70 °E).

2.4 Data collection criteria

A critical review of the literatures collected from the above-mentioned sources was conducted to obtain reliable growth data using the following criteria:

(1) Scope: The objective of this study was to provide the data for understanding growth characteristics of *Larix gmelinii* natural forests (pure *Larix gmelinii* or its proportion more than 50%) and monoculture plantations. Forest stands included in the data set were restricted to those not recently disturbed by logging, fire, or insect pests. Additionally, the following small numbers of special types were excluded: (i) *Larix gmelinii* afforestation in wetland (Li

et al., 1985; Song & Li, 1990; Huang, 2011; Cui et al., 2013), pastureland (Duan, 2005), or abandoned mine land (Yang et al., 2013); (ii) hybrid test between *Larix gmelinii* and other larch (Deng et al., 2010; Zhang et al., 2005); (iii) low-yield stands in hard environment~~, e.g. igneous rock forest~~ (Wang et al., 1979~~), old man forest (;~~ Wang et al., 1991).

(2) Study design and sampling: DBH and tree height ~~and DBH~~ were averaged from the measurement values of all trees in plots or with random/systematic sampling method. Tree regeneration layer, generally below 5 cm in DBH or 1.3 m in height, was neglected in sampled plots. Stem volume of individual tree was computed from felled-wood samples or local tree volume equations. Stand volume was usually calculated by multiplying mean individual volume with stand density. Besides the growth data (i.e. DBH, height ~~DBH~~ and/or volume), the necessary information should be provided in the original sources, e.g., stand age, stand orign, study site, etc.

(3) Quality control: The data quality has been carefully reviewed by the authors. Data has undergone substantial checking, for example, cross-check for the relevant information from different sources, preliminary correlation analysis among growth variables.

Consequently, 776 records that met the above criteria were selected to develop a comprehensive growth data set of Dahurian larch in China. The data set includes growth characteristics of Dahurian larch (i.e. mean tree height (m), mean DBH (cm), mean tree volume (10^{-3} m^3) and/or stand volume (m^3/ha)). In addition, associate information was included, if available in original sources or ascertainable from other relevant literatures, i.e., geographical location (province location and locality name of study site, latitude (°), longitude (°), altitude (m), aspect and slope (°)), stand description (origin, stand age (years), stand density (trees/ha) and canopy density), climate (mean annual temperature (MAT, ℃) and mean annual precipitation (MAP, mm)), and sample regime (observing year, plot size and number).

2.5 Data structural descriptors

Table 1 Variable information in the data set.

Column code	Definition	Unit	Number	Range
ID	Unique identification number of each record	N/A	776	1—776
Province	Province location of study site	N/A	7764	N/A
Study site	Locality name of study site	N/A	123	N/A
Latitude	Latitude of study site	°	776	40.85—53.47
Longitude	Longitude of study site	°	776	118.20—133.70
Altitude	Altitude of study site	m	776	130—1260
Aspect	Slope direction of study site, including flat slope (FL), sunny slope (SU: South), half-sunny slope (HSU: West, Southwest, Southeast), shady slope (SH: North) and half-shady slope (HSH: East, Northwest, Northeast)	N/A	300	N/A
Slope	Slope degree of study site	°	357	0—60
Origin	Stand origin was classified into natural and planted forests	N/A	776	N/A
MAT	Mean annual temperature, from original study or other related ference	℃	776	-6.1—7.0
MAP	Mean annual precipitation, from original study or other related reference	mm	776	355—926
Age	Stand age, which is generally defined as age since germination in natural forest and since planting in planted forest. Stand age is usually obtained from historical records or tree rings.	years	776	1—280
Height	Mean tree height	m	670	0.24—29.40
DBH	Mean diameter at breast height, base diameter was only given in some young forests and marked with [B]	cm	661697	0.70—34.89
V_{tree}	Mean tree volume, the estimated tree volume data from the two-variable larch equation were marked with [E]	10^{-3} m^3 /tree	696	0.04—935.736
V_{stand}	Stand volume, the estimated stand volume data from the estimated tree volume and stand density were marked with [E]	m^3/ha	590	0.07—975.32
Density	Stand density/Canopy density, planting density was only given in some studies and marked with [P]	trees/ha %	656 150	213—13275 0.2—1.0
Area	Plot area	m^2	397	50—10000
Plot	Plot numbers, i.e. replications	N/A	573	1—25
Year	Investigation year	N/A	533	1954—2014
Reference	Data sources, the sources used to supplement climate information lacking in the original publications were added asterisks.	N/A	226	1965—2015

3 Data estimates and evaluation

3.1 Geographical location

Google Earth (Version: 7.1.8.3036) was used to estimate latitude, longitude, and/or altitude when the geographic coordinates was unavailable in original sources.

3.2 Tree and stand volume

The missing tree and stand volumes were estimated with the available information (mean DBH, mean tree height and stand density). Stem volume of individual tree was calculated with the larch equation in northeast China (V_{tree}= 0.000050168241 $DBH^{1.7582894}$ $H^{1.1496653}$), the best method recommended by the ministry standard of China: Tree volume tables (LY 208-77) (Agriculture and Forestry Ministry of China~~Forestry Administration of China~~, 1978; Liu, 2017). Meanwhile, available 590 pairs of mean tree height and mean DBH in the data set were used to establish the ~~linear~~ H-DBH correlation with power function (R^2=0.~~8377~~8919, see Fig. 2). To calculate tree volume from only one known variable of DBH, tree height was firstly calculated with the power~~linear~~ H-DBH equation in Fig. 2. The estimated stand volume was determined by multiplying the estimated tree volume with stand density.



Figure 2. Relationship between mean tree height and diameter at breast height in the data set.

The accuracy of tree volume was assessed by comparing the extracted data from references with the calculated data from the above-mentioned two-variable tree volume equation (Fig. 3). The coefficient of determination (R^2) was 0.9724 and the slope was 1.~~0737~~0733. Therefore, we were confident in applying the larch volume equation to interpolate tree and stand volume data in this study.



Figure 3. Comparison of available tree volume from references with simulated values from the two-variable tree volume equation (V_{tree}=0.000050168241 $DBH^{1.7582894}$ $H^{1.1496653}$). Open circles (N=317): tree height (H) and diameter at breast height (DBH) were available in the references, solid circles (N=37): DBH was only available in the references and H was estimated with the H-DBH correlation from Fig. 2.

Acknowledgements. We would like to acknowledge ~~the~~ many scientists and researchers whose field measurements made this work possible. This research was supported by the

National Key Research and Development Program of China (2017YFC0503906) and China Special Fund for Meteorological Research in the Public Interest (GYHY201406034).

References

Achard, F., Mollicone, D., Stibig, H., Aksenov, D., Laestadius, L., Li, Z., Popatov, P., and Yaroshenko, A.: Areas of rapid forest-cover change in boreal Eurasia, Forest Ecology and Management, 237, 322-334, https://doi.org/10.1016/j.foreco.2006.09.080, 2006.

Agriculture and Forestry Ministry of China: Tree Volume Tables (LY 208-77), Standard Press of China, Beijing, China, 1978.

Bastin, J.F., Berrahmouni, N., Grainger, A., Maniatis, D., Mollicone, D., Moore, R., Patriarca, C., Picard, N., Sparrow, B., Abraham, E.M., Aloui, K., Atesoglu, A., Attore, F., Bassüllü, Ç., Bey, A., Garzuglia, M., García-Montero, L.G., Groot, N., Guerin, G., Laestadius, L., Lowe, A.J., Mamane, B., Marchi, G., Patterson, P., Rezende, M., Ricci, S., Salcedo, I., Diaz, A.S.P., Stolle, F., Surappaeva, V., and Castro, R.: The extent of forest in dryland biomes, Science, 356, 635-638. https://doi.org/10.1126/science.aam6527, 2017.

Cui, W., Mu, C. C., Lu, H. C., Bao, X., and Wang, B.: Effects of draining for forestation on carbon storage of wetland ecosystem in Daxing'an Mountains of northeastern China, Journal of Beijing Forestry University, 35(5), 28-36, https://doi.org/10.13332/j.1000-1522.2013.05.020, 2013.

Deng, J. F., Zhang, H. G., Zhang, L., Guan, C. Y., and Zhang, L.: Genetic variation of 17-year-old hybrid larch and its superior family selection, Journal of Northeast Forestry University, 38(1), 8-11, https://doi.org/10.13759/j.cnki.dlxb.2010.01.017, 2010.

Duan, W. B.: Suitable types of soils for main tree species of Shelterbelts on pastureland, Chinese Journal of Soil Science, 36(6), 871-876, https://doi.org/10.19336/j.cnki.trtb.2005.06.012, 2005.

Fang, J., Chen, A., Peng, C., Zhao, S., Ci, L.: Changes in forest biomass carbon storage in China between 1949 and 1998, Science, 292(5525), 2320-2322, https://doi.org/10.1126/science.1058629, 2001.

FAO: Global Forest Resources Assessment 2015, UN Food and Agriculture Organization, Rome, Italy, available at: http://www.fao.org/3/a-i4808e.pdf, last access: 8 October 2017, 2015.

Huang, D. Q.: Biomass and carbon storage estimation of *Larix gmelinii*, Thesis, available at: http://d.wanfangdata.com.cn/Thesis/Y2026419, last access: 8 October 2017, 2011.

Keenan, R. J., Reams, G. A., Achard, F., de Freitas, J. V., Grainger, A., and Lindquist, E.: Dynamics of global forest area: Results from the FAO Global Forest Resources Assessment 2015, Forest Ecology and Management, 352, 9-20, https://doi.org/10.1016/j.foreco.2015.06.014, 2015.

Li, J. C., Zhao Y. Y., Gao Y. Q., and Liu G. L.: A preliminary analysis of changes in ecological factors after reforestation in swamp land, Scientia Silvae Sinicae, 21(1), 80-85, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE198501009.htm, last access: 8 October 2017, 1985.

Liu, Q. J.: Tree volume tables of China, China Forestry Publishing House, Beijing, China,

2017.
Luo, T. X.: Patterns of net primary productivity for Chinese major forest types and their mathematical models, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-80061-2006113572.htm, last access: 8 October 2017, 1996.
Ma, C. G.: Selection of optimum species and seed sources for plantations of larch, Beijing Agricultural university press, Beijing, China, 1992.
Payn, T., Carnus, J. M., Freer-Smith, P., Kimberley, M., Kollert, W., Liu, S. R., Orazio, C., Rodriguez, L., Silva, L. N., and Wingfield, M. J.: Changes in planted forests and future global implications, Forest Ecology and Management, 352, 57-67, https://doi.org/10.1016/j.foreco.2015.06.021, 2015.
Shugart, H. H., Leemans, R., and Bonan, G. B.: A system analysis of the global boreal forest, Cambridge University Press, Cambridge, 1-10, 1992.
Song, X. S. and Li, Y. M.: Study on the effect of improvement of fen land and afforestation of *Larix gmelinii*, Forestry Science & Technology, 6, 5-7, https://doi.org/10.13456/j.cnki.lykt.1990.06.003, 1990.
Wang, H. Q., Wu, J. C., Yang, W. B., Wang, S. T., and Song, C. J.: A research on management and utilization of "old man forest" of *Larix gmelinii*, Forestry Science & Technology, 16(3), 12-14, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ199103003.htm, last access: 8 October 2017, 1991.
Wang, W. J., Zu, Y. G., Wang, H. M., Matsuura, Y., Sasa, K., and Koike, T.: Plant biomass and productivity of *Larix gmelinii* forest ecosystems in Northeast China: intra- and inter-species comparison, Eurasian Journal of Forest Research, 8, 21-41, available at: http://hdl.handle.net/2115/22188, last access: 8 October 2017, 2005.
Wang, X. P., Fang, J. Y., and Zhu, B.: Forest biomass and root-shoot allocation in northeast China, Forest Ecology and Management, 255, 4007-4020, https://doi.org/10.1016/j.foreco.2008.03.055, 2008.
Wang, Y. H., Zhou, G. S., and Jiang, Y. L.: Simulation of NPP and evaluation of ecosystem services of *Larix gmelinii* forest, Chinese Journal of Applied Ecology, 12(5), 648-652, https://doi.org/10.13287/j.1001-9332.2001.0157, 2001a.
Wang, Y. H., Zhou, G. S., Jiang, Y. L., and Yang, Z. Y.: Estimating biomass and NPP of Larix forests using forest inventory data(FID), Acta Phytoecologica Sinica, 25(4), 420-425, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-ZWSB200104006.htm, last access: 8 October 2017, 2001b.
Wang, Z.: Larch Forest in China, China Forestry Publishing House, Beijing, China, 1992.
Wang, Z., Xu, Z. B., Tan, Z. X., and Dai, H. C.: Ecological characteristics of *Larix gmelinii* grown in igneous rock in Aershan forest region, Scientia Silvae Sinicae, 4, 300-303, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE197904010.htm, last access: 8 October 2017, 1979.
Xu, H.: Forests in Daxinganling Mountains China, Science Press, Beijing, China, 22-53, 1998.
Yang, C. P.: Study on the provenance of Dahurian larch, Science Press, Beijing, China, 2009.
Yang, Y., Zhang, K., Cai, T. J., and Gu, J. F.: Effect of afforestation species on soil nutrient in dumping site of open coal mine in Daxing'an Mountain, Journal of Northeast Forestry University, 41(9), 54-58, https://doi.org/10.13759/j.cnki.dlxb.2013.09.028, 2013.

Zeng, W. S.: Integrated individual tree biomass simultaneous equations for two larch species in northeastern and northern China, Scandinavian Journal of Forest Research, 30(7), 594-604, https://doi.org/10.1080/02827581.2015.1046481, 2015.

Zeng, W. S., Duo, H. R., Lei, X. D., Chen, X. Y., Wang, X. J., Pu, Y., and Zou, W. T.: Individual tree biomass equations and growth models sensitive to climate variables for Larix spp. in China, European Journal of Forest Research, 136 (2), 233-249, https://doi.org/10.1007/s10342-017-1024-9, 2017.

Zhang, H. G., Zhang, C. L., Lan, S. B., Kang, Y. K., and Pan, B. L.: The analysis and selection for families of hybrid larch, Journal of Nanjing Forestry University: Natural Science Edition, 29(3), 69-72, https://doi.org/10.3969/j.issn.1000-2006.2005.03.017, 2005.

Zhou, G. S., Wang, Y. H., Jiang, Y. L., and Yang, Z. Y.: Estimating biomass and net primary production from forest inventory data: a case study of China's Larix forests, Forest Ecology and Management, 169(1-2), 149-157, https://doi.org/10.1016/s0378-1127(02)00305-5, 2002.

Zhou, X. F.: Long-Term Research on China's Forest Ecosystems, Northeast Forestry University Press, Harbin, China, 1994.

References cited in the dataset

An, J.: Tree species and management effect on soil carbon sequestration and fertilities of plantation forests. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1012445928.htm, last access: 8 October 2017, 2011.

Bi, Y. H.: Study on stand growth models of larch gmelinii with different thinning intensity. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2001005669.htm, last access: 8 October 2017, 2001.

Cao, G. Z.: Analysis of hydrological characteristics and change trend in westen region of Songhua River Basin. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10183-1011098650.htm, last access: 8 October 2017, 2011

Chen, J. D., Pan, B. L., Wu, M. C., Han, C. W., and Yan, W. F.: A summary report on the introduction of *Larix principis-rupprechtii*, Forestry Science & Technology, 4, 2-5, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ198304000.htm, last access: 8 October 2017, 1983.

Chen, L. X. and Xiao, Y.: Evolution and evaluation of soil fertility in forest land in *Larix gmelinii* plantations at different development stages in Daxinganling forest region, Science of Soil and Water Conservation, 4(5), 50-55, https://doi.org/10.16843/j.sswc.2006.05.012, 2006.

Chen, Q., Zhao, Y. S., Xin, Y., Han, X. C., Xia, X. Y., and Han, C. H.: Plant species diversity and correlation with soil factors after vegetation restoration in burned area of Great Xing'an Mountains, Bulletin of Soil and Water Conservation, 32(6), 61-65, 70, https://doi.org/ 10.13961/j.cnki.stbctb.2012.06.040, 2012a.

Chen, Q., Zhao, Y. S., Xin, Y., and Han, X. C.: Litter and soil water-holding capacity of burned area in northern slope of the Great Khingan after vegetation restoration, Chinese Journal of Soil Science, 44(2), 308-313, https://doi.org/10.19336/j.cnki.trtb.2013.02.008, 2013.

Chen, R. M.: A preliminary report on provenance trials of *Larix gmelinii*, Forest Investigation Design, 1, 45-49, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-

LYKC198901014.htm, last access: 8 October 2017, 1989.
Chen, X. W., Wang, C. H., Ding, B. Y., and Zhang, S. Y.: Dynamic thinning system for larch plantation, Journal of Northeast Forestry University, 5(2), 17-21, https://doi.org/10.1007/BF02843226, 1994.
Chen, Y. L., Han, S. J., Zhou, Y. M., Zou, C. J., and Zhang, J. H.: Characteristics of available P in the rhizosphere soil in pure *Juglans mandshurica* and *Larix gmelinii* and their mixed plantation, Chinese Journal of Applied Ecology, 13(7), 790-794, https://doi.org/10.13287/j.1001-9332.2002.0188, 2012b.
Cheng, F. Y.: Spatial variations in non-structural carbonhydrates of *Larix gmelinii*, *Juglans mandshurica* and *Ulmus japonica*. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1015665453.htm, last access: 8 October 2017, 2015.
Cheng, Y. X. and Li, Z. X.: Preliminary study of biomass of three forest types of *Larix gmelinii*, Inner Mongolia Forestry Investigation and Design, 4, 29-39, available at: http://www.cqvip.com/QK/90816X/198904/206948.html, last access: 8 October 2017, 1989.
Chu, X.: Spatial and temporal distribution of fine root biomass of dominant tree species and changes after fire on Great Xing'an Mountains. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1011147000.htm, last access: 8 October 2017, 2011.
Chu, X., Di, X. Y., Zhang, J. L., Cai, H. Y., and Yan, B. Z.: Distribution and seasonal dynamics of fine root biomass of two types of forests in Great Xing'an Mountains, Journal of Northeast Forestry University, 39(5), 36-39, https://doi.org/10.13759/j.cnki.dlxb.2011.05.033, 2011.
Cui, G. F., Cai, T. J., and Yang, W. H.: Soil acidity of *Larix gmelinii* plantation, Journal of Beijing Forestry University, 22(3), 33-36, https://doi.org/10.13332/j.1000-1522.2000.03.008, 2000.
Di, X. Y., Geng, Y. Y., Sun, L., and Hu, H. Q.: Soil microbial biomass dynamics and influence factors in *Larix gmelinii* forest in Daxing'an Mountains, Journal of Northeast Forestry University, 40(5), 1-4, https://doi.org/10.13759/j.cnki.dlxb.2012.05.009, 2012.
Ding, B. Y.: The method and intensity of intermediate cuttings for larch plantation, Journal of Northeast Forestry University, 16(2), 46-57, https://doi.org/10.13759/j.cnki.dlxb.1988.02.007, 1988.
Ding, B. Y., Ju, Y. G., Zhang, S. S., Zhang, S. Y., and Yu, W. J.: Study on the structure of the larch wood community, Journal of North-Eastern Forestry Institute, 4, 11-23, https://doi.org/10.13759/j.cnki.dlxb.1982.04.002, 1982.
Ding, B. Y., Liu, S. R., and Cai, T. J.: Studies on biological productivity of artificial forests of Dahurian larches, Acta Phytoecologica et Geobotanica Sinica, 14(3), 226-236, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-ZWSB199003003.htm, last access: 8 October 2017, 1990.
Dong, X. B.: The impacts of cutting intensity on the growth of larch forest, Journal of Northeast Forestry University, 29(1), 44-47, https://doi.org/10.13759/j.cnki.dlxb.2001.01.011, 2001.
Dong, X. B.: Modes and evaluation of ecological harvesting in Larch plantation areas. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2009132292.htm, last access:

8 October 2017, 2008.
Dong, X. B., Yang, X. C., and Yang, G. X.: Impacts of different cutting modes on soil properties in Larch plantations, Journal of Northeast Forestry University, 35(10), 7-10, https://doi.org/10.13759/j.cnki.dlxb.2007.10.012, 2007.
Dong, Y.: A survey study on the factors influencing the growth of *Larix gmelinii* and *Pinus sylvestris* plantations, Transactions of China Pulp and Paper, Suppl., 134-142, https://doi.org/10.3321/j.issn:1000-6842.2004.z1.021, 2004.
Du, E. Z. and Fang J. Y.: Linking belowground and aboveground phenology in two boreal forests in Northeast China, Oecologia, 176, 883-892, https://doi.org/10.1007/s00442-014-3055-y, 2014.
Du, Y. J., Xu, H. C., and Yu, R. Y.: The effect of understory, litter and small mammals on seedling occurrence of *Larix gmelinii*, Journal of Beijing Forestry University, 15(4), 12-20, https://doi.org/10.13332/j.1000-1522.1993.04.002, 1993.
E, W. F.: Stem respiration spatial-temporal dynamics and characteristics of sapwood/ heartwood of *Larix gmelinii*, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2009133891.htm, last access: 8 October 2017, 2008.
E, W. F., Wang C. K., Yang C. P., Wang, X. C., Zhang, Q. Z., and Zhang, Y. Q.: Effects of provenances on growth characteristics of sapwood and heartwood of *Larix gmelinii* trees, Scientia Silvae Sinicae, 45(6), 109-115, https://doi.org/10.3321/j.issn:1001-7488.2009.06.019, 2009.
Feng, L. and Yang, Y. G.: A study on biomass and production of three types of Dahurian larch virgin forest, Scientia Silvae Sinicae, 21(1), 86-92, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE198501010.htm, last access: 8 October 2017, 1985.
Fu, Y.: Study on *Larix gmelinii* plantation in different planting density in the south of Daxing'anling City, Journal of Inner Mongolia Forestry, 5, 24-26, available at: http://www.cnki.com.cn/Article/CJFDTotal-NMLY198405030.htm, last access: 8 October 2017, 1984.
Gao, M. F., Li, C. H., and He, S. S.: Productivity division of *Larix gmelinii* in Great Xing'an Mountains of Inner Mongolia, Inner Mongolia Forestry Investigation and Design, 28, 75-77, 84, https://doi.org/10.13387/j.cnki.nmld.2005.s1.037, 2005.
Gao, X. W.: Research by the stand density effect on Xing'an larch of natural forest, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10129-1014375561.htm, last access: 8 October 2017, 2014.
Gao, X. W., Zhang, Q. L., Bao, Y., Wang, X. H., Li, D. Y., and Shi, X.: Study on the forest biomass and carbon density under different cutting patterns of Dahurian Larch forest, Forest Resources Management, 3, 91-96, https://doi.org/10.13466/j.cnki.lyzygl.2014.03.020, 2014.
Gu, Y. C.: The necessity of tending and thinning of natural young and middle aged forest, Forest Resources Management, 1, 58-62, https://doi.org/10.13466/j.cnki.lyzygl.1985.01.018, 1985a.
Gu, Y. C.: The succession of forest communities in the forest region of the Greater Xingan Mountains, Acta Phytoecologica et Geobotanica Sinica, 9(1), 64-70, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-ZWSB198501006.htm, last access: 8 October 2017, 1985b

Guan, J. Q., Cai, T. J., and Man, X. L.: Characteristics of snowpack chemistry in different type plantations of Xiaoxing'an Mountains, northeastern China, Journal of Beijing Forestry University, 35(4), 41-46, https://doi.org/10.13332/j.1000-1522.2013.04.020, 2013.

Guo, R. Y., Fu, Y., Gao, S. C., and Jia, Z. S.: Studies on the reasonable planting density of Poplar and larch in Xing-an League, Inner Mongolia Forestry Science and Technology, 2, 20-22, 35, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198402005.htm, last access: 8 October 2017, 1984.

Guo, Y. X.: The variation characteristics of heartwood and sapwood of artificial *Larix gmelinii*, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1015665595.htm, last access: 8 October 2017, 2015.

Guo, Y. X. and Jiang, L. C.: Variation of heartwood and sapwood and growth characteristics of artificial Dahurian Larch, Journal of Northeast Forestry University, 43(6), 93-97, https://doi.org/10.13759/j.cnki.dlxb.20150522.048, 2015.

Han, J. Y., Zhang, H. G., Jia, Q. B., Zhang, Z., and Xu, Y. L.: Genetic variations in growth traits of juvenile stage hybrid larch in southern Xiao Xing'an Mountains and selection of superior families, Journal of Northwest Forestry University, 27(6), 78-82, https://doi.org/10.3969/j.issn.1001-7461.2012.06.16, 2012a.

Han, M. Z.: A research on biomass and production of Larix Dahurica forest communities, Acta Agriculturae Boreali-Sinica, 2(4), 134-138, https://doi.org/10.3321/j.issn:1000-7091.1987.04.022, 1987.

Han, S. L., Zhang, Q. L., AN, H. J., and Yu, B.: Spatial structure characteristics of *Larix gmelinii* natural forest, Journal of Arid Land Resources and Environment, 29(5), 87-92, https://doi.org/10.13448/j.cnki.jalre.2015.157, 2015a.

Han, X. C.: Effect on process of burned areas *Larix gmelinii* plantations restoration in Daxinganling, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1013371072.htm, last access: 8 October 2017, 2013.

Han, X. C., Zhao, Y. S., Xin, Y., and et al.: Soil hydrology characteristics of *Larix gmelinii* plantation in burned forestland in the north of Greater Xing'an Mountains, Journal of Soil and Water Conservation, 26(4), 183-188, https://doi.org/10.13870/j.cnki.stbcxb.2012.04.053, 2012b.

Han, X. C., Zhao, Y. S., and Xin, Y.: Succession process of *Larix gmelinii* forest with artificial restoration after fire in Daxing′anling, Science of Soil and Water Conservation, 13(2), 70-76, https://doi.org/10.16843/j.sswc.2015.02.012, 2015b.

He, H. Y.: Analysis on growth of three species of artifical plantation in Zhangguangcai Mountains-subregion of Xiaoxingan Mountains, Journal of Northeast Forestry University, 16(1), 32-36, https://doi.org/10.13759/j.cnki.dlxb.1988.01.005, 1988.

Hong, Q. L., Cai, Y. X., Wang, Y. H., Zhao, H. X., Hu, C. X., Li, F. R., and Fan, W. Y.: The growth of natural forests of *Larix gmelinii* in Tahe Forestry Bureau Daxingan Mountain District, Journal of Northeast Forestry University, 22(2), 92-97, https://doi.org/10.13759/j.cnki.dlxb.1994.02.017, 1994.

Hu, H. Q., Wei, S. J., and Sun, L.: Estimation of carbon emissions from forest fires in 2010 in Huzhong of Daxing'anling Mountain, Scientia Silvae Sinicae, 48(10), 109-119, https://doi.org/10.11707/j.1001-7488.20121017, 2012.

Huang, C., Meng, F. L., Shan, C. C., and Hua, S.: Techniques for fast growing and high yield

of larch, Journal of Heilongjiang Vocational Institute of Ecological Engineering, 24(1), 28-29, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-HSGX201101014.htm, last access: 8 October 2017, 2011a.

Huang, D. Q.: Biomass and carbon storage estimation of *Larix gmelinii*, Thesis, available at: http://d.wanfangdata.com.cn/Thesis/Y2026419, last access: 8 October 2017, 2011.

Huang, D. Q., Hu, H. Q., Wang, Z. W., and Sun, L.: Dynamic changes in biomass and productivity of *Larix gmelinii* plantation after fire, Journal of Northeast Forestry University, 39(9), 62-64, https://doi.org/10.13759/j.cnki.dlxb.2011.09.025, 2011b.

Huang, S. Z.: The internal cycling of nutrients in fine roots of *Fraxinus Mandshurica* and *Larix gmelinii*, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2006110390.htm, last access: 8 October 2017, 2006.

Huo, H.: The effects of climate warming on photosynthesis and growth of *Larix gmelinii*. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2007187763.htm, last access: 8 October 2017, 2007.

Jia, S. X., Zhao, Y. L., Sun, Y., Chen, L., and Wang, Z. Q.: Effects of nitrogen fertilization on seasonal dynamics of soil microbial biomass carbon and nitrogen in *Larix gmelinii* and *Fraxinus mandshurica* plantations, Chinese Journal of Applied Ecology, 20(9), 2063-2071, https://doi.org/10.13287/j.1001-9332.2009.0380, 2009.

Jiang, H. Y.: Study on hydrological characteristics of forest ecosystem in Daxing'anling, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2009132313.htm, last access: 8 October 2017, 2008.

Jiang, H. Y., Zhao, Y. S., Chen, X. W., Li, W. H., Zhu, W. C., Lv, W. B., and Li, X. P.: Research on soil hydrology characteristics of some main forest type in south part of Daxing'anling, Journal of Soil and Water Conservation, 21(3), 149-153,187, https://doi.org/10.13870/j.cnki.stbcxb.2007.03.033, 2007.

Jiang, H. Y., Zhao, Y. S., Xin, X. J., Ma, W. H., Li, X. P., Sun, C. K., Guo, X. W.: Precipitation distribution laws of different forest stand in Daxing'Anling, Journal of Soil and Water Conservation, 22(6), 197-201, https://doi.org/10.13870/j.cnki.stbcxb.2008.06.044, 2008.

Jiang, L. F.: Study on mechanism of photosynthesis and respiration of *Larix gmelinii* plantations, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2003110988.htm, last access: 8 October 2017, 2003.

Jiang, L. F., Shi, F. C., Li, B., Luo, Y. Q., Chen, J. Q., and Chen, J. K.: Separating rhizosphere respiration from total soil respiration in two larch plantations in northeastern China, Tree Physiology, 25, 1187-1195, https://doi.org/10.1093/treephys/25.9.1187, 2005.

Jiang, L. F., Shi, F. C., Zu, Y. G., Wang, W. J., and Koike, T.: Study on stem respiration of *Larix gmelinii* of different ages and its relationship to environmental factors, Bulletin of Botanical Research, 23(3), 296-301, https://doi.org/10.3969/j.issn.1673-5102.2003.03.012, 2003.

Jiang, R., Sun, L., and Hu, H. Q.: The seasonal dynamics of soil microbial biomass of a *Larix gmelinii* forest after wildfire, African Journal of Microbiology Research, 6(10), 2328-2336, https://doi.org/10.5897/ajmr11.1264, 2012.

Laboratory of Forest Tree and Breeding, Northeast Forestry University(LFTB).: Study on larch heterosis, Forestry Science & Technology, 1, 1-3, https://doi.org/10.13456/j.cnki.lykt.

1979.01.001, 1979.
Li, B.: The impact of different densities on the *Larix gmelinii* plantation. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10129-1013248427.htm, last access: 8 October 2017, 2013a.
Li, B., De, Y. J., Wei, Q., Wang, Y., Wang, Y. B., and Bao, Y. Q.: Effects of three densities on the growth of *Larix gmelinii* plantations, Journal of Inner Mongolia Agricultural University, 34(2), 65-68, https://doi.org/10.16853/j.cnki.1009-3575.2013.02.013, 2013.
Li, J., Luo, Y. Q., Shi, J., Zhou, J., Wang, X. L., Ma L. Y., and Chen C.: Use of phyto-attractant in monitoring and controlling gypsy moth, Journal of Beijing Forestry University, 33(4), 85-90, https://doi.org/10.13332/j.1000-1522.2011.04.010, 2011.
Li, L. N., Zhao, L., Zhang, W. J., and Wang, X. J.: Study on species diverstity in different density of *Larix gmelinii* plantations in Zhalantun Region, Practical Forestry Technology, 8, 9-11, https://doi.org/10.13456/j.cnki.lykt.2014.08.002, 2014.
Li, M., Gu, J. C., Zhang, Z. W., and Wang, Z. Q.: Responses of fine root mass, length, production and turnover to soil nitrogen fertilization in *Larix gmelinii* and *Fraxinus mandshurica* forests in Northeastern China, Journal of Forest Research, 15, 194-201, https://doi.org/10.1007/s10310-009-0176-y, 2010a.
Li, Q., Chen, J., Li, T., and Liu, F. J.: Ecological function of artificial coniferous timber forests in water conservation, Jilin Forestry Science and Technology, 36(3), 14-17, 20, https://doi.org/10.16115/j.cnki.issn.1005-7129.2007.03.004, 2007.
Li, S. L. and Chen, Y. L.: Decomposition and nutrient return of the leaf litter under the pure and mixed plantations of *Juglans mandshrica* and *Larix gmelinii*, Journal of Nanjing Forestry University: Natural Science Edition, 28(5), 59-62, https://doi.org/10.3969/j.issn.1000-2006.2004.05.014, 2004.
Li, X. F.: Comprehensive evaluation of the effects of different thinning intensities on *Larix gmelinii* forest in Great Xing'an Mountains of Inner Mongolia, Inner Mongolia Forestry Investigation and Design, 1, 7-20, available at: http://www.cqvip.com/QK/90816X/199301/1276513.html, last access: 8 October 2017, 1993.
Li, X. F.: Study on genetic diversity of provenances and clones of *Larix gmelinii* (Rupr.) Rupr, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2009134088.htm, last access: 8 October 2017, 2009.
Li, X. F.: Spatio-temporal dynamics and influencing factors of fine roots in five Chinese temperate forest ecosystems, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1013366884.htm, last access: 8 October 2017, 2013b.
Li, X. Y.: The study on dynamic features of several metabolites of larch(*Larix gmelinii*), Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2005136726.htm, last access: 8 October 2017, 2005.
Li, Y. W., Chen, X. H., and Hu, M. Q.: Study on growth and mortality of mature and over-mature *Larix gmelinii* forest in Daxinganling forest region, Forest Resources Management, 1, 38-43, https://doi.org/10.13466/j.cnki.lyzygl.1979.01.007, 1979.
Li, Z. F., Zhao, Y. S., Xin, Y., Kong, X. F., and Wu, C. Y.: Soil water storage and infiltration characteristics under 3 artificial forests in the upper reaches of Ashihe river, Science of Soil and Water Conservation, 8(1), 77- 80, https://doi.org/10.16843/j.sswc.2010.01.014, 2010b.
Li, Z. F., Zhao, Y. S., Xin, Y., Kong, X. F., and Wu, C. Y.: Temporal and spatial variation of

soil water under different forest types in the upper reaches of Ashihe River, Science of Soil and Water Conservation, 8(4), 86-89, https://doi.org/10.16843/j.sswc.2010.04.016, 2010c.
Lin, L.: Afforestation techniques on southward slopes in Greater Khingan Mts. forest district, Journal of Northeast Forestry University, 3, 48-58, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198103005.htm, last access: 8 October 2017, 1981.
Liu, G. L.: Intermediate cutting experiment of larch (Larix sp.) plantation, Journal of North-Eastern Forestry Institute, 12(1), 33-46, https://doi.org/10.13759/j.cnki.dlxb.1984.01.004, 1984.
Liu, G. L. and Xu, M. H.: Study on half-sib breeding plantation of different larchs. Forestry Science and Technology, 2: 19-24. available at: http://www.cnki.com.cn/Article/CJFDTOTAL-JLLK198402003.htm, last access: 8 October 2017, 1984.
Liu, H. L., Cai, T. J., Man, X. L., Chai, R. S., and Lang, Y.: Effects of major forest types of Xiaoxing'an Mountains on the process of snowfall, snow cover and snow melting, Journal of Beijing Forestry University, 34(2), 20-25, https://doi.org/10.13332/j.1000-1522.2012.02.003, 2012.
Liu, J. L., Wang, C. K., and Zhang, Q. Z.: Spatial variations in stem heartwood and sapwood for *Larix gmelinii* trees with various diferentiation classes, Scientia Silvae Sinicae, 50(12), 114-121, https://doi.org/10.11707/j.1001-7488.20141216, 2014.
Liu, S. R., Chai, Y. X., Cai, T. J., and Peng, C. H.: Study on biomass and net primary productivity of Dahurian larch plantation, Journal of Northeast Forestry University, 18(2), 40-46, https://doi.org/10.13759/j.cnki.dlxb.1990.02.006, 1990a.
Liu, S. R., Li, X. M., and Niu, L. M.: The degradation of soil fertility in pure larch plantations in the northeastern part of China, Ecological Engineering, 10, 75-86, https://doi.org/10.1016/s0925-8574(97)10024-6, 1998.
Liu, T. Y.: Carbon storage in Manchurian ash and Dahurian larch plantations with different stand structures. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1012445805.htm, last access: 8 October 2017, 2012.
Liu, W. and Deng, H.: Study on the effect of different afforestation methods on the survival rate of Larix, Forestry Science and Technology Information, 39(4), 4-5, https://doi.org/10.3969/j.issn.1009-3303.2007.04.003, 2007.
Liu, X. T., Du, X. G., Liang, L., Zhang, P., Zhang, B. C., and Zhang, Y. S.: The selection of geographic provenances of *Larix gmelinii* in the eastern slope of Great Xing'an Mountain, Journal of Northeast Forestry University, 18, 9-16, https://doi.org/10.13759/j.cnki.dlxb.1990.s2.003, 1990b.
Liu, X. T., Zhang, H. J., Wang, S. X., Lin, Y. K., and He, X. Y.: Introduction of *Larix olgensis* in Daxing'anling, Journal of Northeast Forestry University, 17(3), 18-25, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198903002.htm, last access: 8 October 2017, 1989.
Liu, Y.: Study on the niche and the production of artificial Korean pine and Larch population. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2003111030.htm, last access: 8 October 2017, 2002.
Liu, Y. C., Wu, M. C., Chu, Y. J., Liu, D. C., and He, Z. J.: Study on the provenance test of *Larix gmelinii*. Forestry Science and Experiment, 35, 1-17, available at: http://www.cqvip.com/QK/92313X/199035/319899.html, last access: 8 October 2017, 1990c.

Lu, X., Hu, H. Q., Sun, L., and Wei, S. J.: Effects of fire disturbance on soil light fraction organic carbon of *Larix gmelinii* forest, Journal of Anhui Agricultural Sciences, 41(29), 11709-11713, https://doi.org/10.13989/j.cnki.0517-6611.2013.29.006, 2013.

Luo, Y. J., Wang, X. K., Zhang, X. Q., and Lu, F.: Biomass and its allocation of forest ecosystems in China, China Forestry Publishing House, Beijing, China, 2013.

Ma, H. B.: Stem sap flow dynamics in relation to environmental factors of Larix gmelinii. Thesis, https://doi.org/10.7666/d.y1473827, 2009.

Ma, J. W.: Study on spectral characteristics of pure stands under different site conditions and phenological stages at Maoershan Area, Journal of Beijing Forestry College, 3, 26-40, https://doi.org/10.13332/j.1000-1522.1983.03.003, 1983.

Ma, J. W., Zhang, H. G., Liu, L., Zhang, Z., and Jia, Q. B.: Genetic variation of hybrid larch at juvenile period and its superior family selection, Journal of Northeast Forestry University, 40(9), 1-5, https://doi.org/10.13759/j.cnki.dlxb.2012.09.019, 2012.

Mao, W. X., Tong, D. H., Zhang, C., Zhao, D. D., Ding, Y. Y., and Jin, S.: Spatial heterogeneity of moisture content of land surface dead fuel in larch stand and effects of sampling methods on moisture estimation, Journal of Northeast Forestry University, 40(5), 29-33, https://doi.org/10.13759/j.cnki.dlxb.2012.05.012, 2012.

Mao, Y. Q., Zhang, J. L., Wang, F. S., Liu, L., Zhao, W., Zhang, H. P., Yuan, L. D., Wang, F. J., and Li, B. C.: Study on the provenance test of *Larix olgensis* and *Larix gmelinii*, Protection Forest Science and Technology, 2, 10-15, https://doi.org/10.13601/j.issn.1005-5215.1993.02.003, 1993.

Mao, Y. Q., Zhang, J. L., Wang, F. S., Liu, L., and Wu, C. J.: A study on introduction of Larch. Scientia Silvae Sinicae 34(4): 48-58. available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE804.006.htm, last access: 8 October 2017, 1998.

Meng, C., Wang, L. H., Wang, J., You, X. F., Liu, Y. Q., and Xing, J.: Diurnal change in soil respiration rate in five types of plantations during non-growing season, Journal of Northeast Forestry University, 37(11), 77-79, 91, https://doi.org/10.13759/j.cnki.dlxb.2009.11.029, 2009.

Meng, X. Q., Li, H. Y., Wang, Y. M., Zhang, Y. W., and Yang, W. Y.: Study on the effects of Coleophora dahurica on tree growth, Journal of Liaoning Forestry Science & Technology, 5, 19-20, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LNLK200005007.htm, last access: 8 October 2017, 2000.

Mu, T. M. and Yan, W.: Preliminary study on the evapotranspiration in Xingan larch forest, Journal of Northeast Forestry University, 2, 21-30, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198202001.htm, last access: 8 October 2017, 1982.

Na, D. C.: A study on genetic diversity of different geographic provenances of *Larix gmelinii*(Rupr.) Rupr and its utilization, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2005136986.htm, last access: 8 October 2017, 2005.

Na, D. C., Yang, C. P., Jiang, J., Xia, D. A., and Wang, H. R.: Provenance division and superior provenance choice of *Larix gmelinii*, Forestry Science & Technology, 30(4), 1-4, https://doi.org/10.3969/j.issn.1001-9499.2005.04.001, 2005.

Piao, C. B., Zhang, W. X., Lin, J., and Ma, A. B.: The preliminary report on the good natural Dahurian larch stand transformation for seed stand, Forestry Science & Technology, 16(3), 17-23, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ199103005.htm,

last access: 8 October 2017, 1991.
Qi, G., Wang, Q. L., Wang, X. C., Qi, L., Wang, Q. W., Ye, Y. J., and Dai, L. M.: Vegetation carbon storage in *Larix gmelinii* plantations in Great Xing'an Mountains, Chinese Journal of Applied Ecology,22(2), 273-279, https://doi.org/10.13287/j.1001-9332.2011.0072, 2011.
Qi, G., Wang, Q. L., Wang, X. C., Yu, D. P., Zhou, L., Zhou, W. M., Peng, S. L., and Dai, L. M.: Soil organic carbon storage in different aged *Larix gmelinii* plantations in Great Xing'an Mountains of Northeast China, Chinese Journal of Applied Ecology, 24(1), 10-16, https://doi.org/10.13287/j.1001-9332.2013.0122, 2013.
Qiu, L.: Temporal dynamic change of soil pool in larch plantations and its contribution to the forest carbon budget in NE China, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1011146228.htm, last access: 8 October 2017, 2011.
Qiu, L., Zu, Y. G., Wang, W. J., Sun, W., Su, D. X., and Zheng, G. Y.: CO_2 flux characteristics and their influence on the carbon budget of a larch plantation in Maoershan region of Northeast China, Chinese Journal of Applied Ecology, 22(1), 1-8, https://doi.org/10.13287/j.1001-9332.2011.0031, 2011.
Quan, X., Wang, C., Zhang, Q., Wang, X., Luo, Y., and Bond-Lamberty, B.: Dynamics of fine roots in five Chinese temperate forests, Journal of Plant Research, 123, 497-507, https://doi.org/10.1007/s10265-010-0322-9, 2010.
Ren, B. P.: Research on growth process and stand density of *Larix gmelinii* plantation, Journal of Inner Mongolia Radio & TV University, 2, 73-75, 85, https://doi.org/10.16162/j.issn.1672-3473.2010.02.022, 2010.
Ren, L. J., Li, G. T., Sun, X. J., Yu, S. P., Zheng, X. W., and Yue, Y. J.: Research on community structure and species diversity of *Larix gmelinii* community in Inner Mongolia Daxing'anling Mountains, Forest Resources Management, 3, 83-87, 118, https://doi.org/10.13466/j.cnki.lyzygl.2013.03.023, 2013.
Shang, Y. L., Zhang, S. H., Chen, Z. C., and Wang, G. Y.: Fine family selection and genetic gain of *Larix gmelinii*, Journal of Northeast Forestry University, 38(7), 123-125, https://doi.org/10.13759/j.cnki.dlxb.2010.07.017, 2010.
Sheng, H. C., Cai, T. J., Zhu, D. G., and Shi, F. C.: Rainfall redistribution and hydrochemical characteristics in the Larch plantation, Journal of Soil and Water Conservation, 23(2), 79-83, 89, https://doi.org/10.13870/j.cnki.stbcxb.2009.02.050, 2009.
Si, H. Z., Yang, S., Li, J., Wang, F. S. and Wen, B. Y.: A study on introduction of *Larix kaempferi*, Forest Science and Technology, 11, 5-7, https://doi.org/10.13456/j.cnki.lykt.1999.11.002, 1999.
Song, C. L., Zhao, P. W., Su, R. N., Hai, L., Zhou, M., and Feng, L.: Dynamic study of photosynthetic characteristics of *Larix gmelinii*, Journal of Inner Mongolia Agricultural University, 29(4), 49-54, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMGM200804010.htm, last access: 8 October 2017, 2008a.
Song, S., Gu, J. C., Quan, X. K., Guo, D. L. and Wang, Z. Q.: Fine-root decomposition of *Fraxinus mandshurica* and *Larix gmelinii* plantations, Journal of Plant Ecology, 32(6), 1227-1237, https://doi.org/10.3773/j.issn.1005-264x.2008.06.003, 2008b.
Song, W. Y.: Carbon density and its distribution of *Pinus koraiensis* and *Larix gmelinii* plantations in different regions of Heilongjiang province. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1014402349.htm, last access: 8 October

2017, 2013.
Song, X. S., Bao, F. L., and Zhang, Y. L.: The analysis of the economic benefit on the fast growing and high yield of *Larix gmelini* plantation, Forestry Science & Technology, 17(2), 16-17, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ199202004.htm, last access: 8 October 2017, 1992.
Song, X. S., Guo, S. Y., and Wang, J. X.: Study on fast growing and high yield of *Larix gmelini* plantation, Forestry Science & Technology, 21(3), 22-24, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ603.009.htm, last access: 8 October 2017, 1996.
Sun, C. Z., Shen, G. F., Li, J. Y., and Jia, L. M.: Study on the present condition and the potentialities of the productivity of main tree species plantation of china II. study on the plantation productivities of Eucalyptus, Larix and Pinus sylvestris var. Mongolica, Forest Research, 14(6), 657-667, https://doi.org/10.13275/j.cnki.lykxyj.2001.06.012, 2001.
Sun, H. Z., Chen, M. Y., Cai, C. J., and Zhu, N.: Effects of different type urban forest plantations on soil fertility, Chinese Journal of Applied Ecology, 20(12), 2871-2876, https://doi.org/10.13287/j.1001-9332.2009.0454, 2009.
Sun, H. Z., Lu, X. J., Chen, M. Y., Cai, C. J., and Zhu, N.: Soil heavy metals concentration in different type plantations in Harbin City, Chinese Journal of Applied Ecology, 22(3), 614-620, https://doi.org/10.13287/j.1001-9332.2011.0126, 2011.
Sun, X. M., Zhang, S. G., Zhou, D. Y., Wang, X. D., Ding, B. and Liu, S. M.: Phenological variation of Larix species and their intra-species and inter-species hybrid families and early selection, Scientia Silvae Sinicae, 44(1), 77-84, https://doi.org/10.3321/j.issn:1001-7488.2008.01.013, 2008.
Sun, Y.: Effects of soil fauna on root biomass, morphology, production and turnover rate in Larch and Ash plantatioins, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2010242477.htm, last access: 8 October 2017, 2010.
Sun, Y., Wang, L. L., Chen, J., and Duan J. P.: Reconstructing mean maximum temperatures of May–August from tree-ring maximum density in North Da Hinggan Mountains, China, Chinese Science Bulletin, 57(16), 2007-2014, https://doi.org/10.1007/s11434-012-5055-9, 2012.
Sun, Z. B. and Sun, X. W.: Selection of species and seed sources for Larix spp. in Chifeng Region, Inner Mongolia, in: Ma, C.G., ed. Selection of optimum species and seed sources for plantations of larch, Beijing Agricultural University Press, Beijing, China, 1992.
Sun, Z. B., Wang, X.W., Ding, M.X., Shi, J.S., Liu, Z., and Bao, G.Z.: Study on selection of optimum species and seed sources for plantations of larch, Forest Science and Technology, 10, 11-13, https://doi.org/10.13456/j.cnki.lykt.1991.10.005, 1991.
Tang, J. Z., Li, Q. Y., Wang, H. X., and Wang, Y. G.: The main raw material forest of larch growth indicators, Forest Investigation Design, 3, 61-63, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKC200903029.htm, last access: 8 October 2017, 2009.
Tian, F., Bi, X. H., Shen, G. T., Zhou, B. and Li, W. B.: Study on the forecast technique and control index of larch caterpillar of a cycle in two years, Forest Research, 11(4), 388-395, https://doi.org/10.13275/j.cnki.lykxyj.1998.04.008, 1998.
Wang, B. S.: The undergrowth succession in larch plantations before and after cutting,

Journal of Northeast Forestry University, 24(5), 82-86, https://doi.org/10.13759/j.cnki.dlxb.1996.05.014, 1996.
Wang, C. K.: Biomass allometric equations for 10 co-occurring tree species in Chinese temperate forests, Forest Ecology and Management, 222, 9-16, https://doi.org/10.1016/j.foreco.2005.10.074, 2006.
Wang, C. K., Gower, S. T., Wang, Y. H., Zhao, H. X., Yan, P., and Bond-Lamberty, B.: The influence of fire on carbon distribution and net primary production of boreal *Larix gmelinii* forests in north-eastern China, Global Change Biology, 7, 719-730, https://doi.org/10.1046/j.1354-1013.2001.00441.x, 2001a.
Wang, C. K. and Yang, J. Y.: Rhizospheric and heterotrophic components of soil respiration in six Chinese temperate forests, Global Change Biology, 13, 123-131, https://doi.org/10.1111/j.1365-2486.2006.01291.x, 2007.
Wang, H. X., Zhang, S. H., Jing, C. Q., Dai, J. G., Liu, Y. T. and Zhou, J. Y.: Cultivation techniques for raw material forest of short rotation Larch, Forestry Science & Technology, 25(5), 7-8, https://doi.org/10.3969/j.issn.1001-9499.2000.05.003, 2000a.
Wang, J. Z.: Study on selection of larch species, Jilin Forestry Science and Technology, 1, 22-26, available at: http://www.cqvip.com/QK/96402X/199501/1795612.html, last access: 8 October 2017, 1995.
Wang, L.: Regeneration rules of the north forest region in western slope of Inner Mongolia Daxing'anling, Inner Mongolia Forestry Investation and Design, S1, 65-69, https://doi.org/10.13387/j.cnki.nmld.2000.s1.022, 2000.
Wang, L., Li, C. H., Yang, J. Y., and Guo, Y. L.: Biomass changes of artificial *Larix gmelinii* from 2002 to 2012 in Helong, Jilin Province, Journal of Green Science and Technology, 2, 11-12, https://doi.org/10.3969/j.issn.1674-9944.2013.02.005, 2013a.
Wang, L. H., Xin, Y., Zou, M. L., Zhao, Y. S., Lei, Y. Y., and Chen, L.: Litter water-holding characteristics in the process of vegetation restoration in severely burned area of Great Xing'an Mountains, Journal of Soil and Water Conservation, 28(6), 262-267, https://doi.org/10.13870/j.cnki.stbcxb.2014.06.048, 2014.
Wang, L. H., Xin, Y., Zou, M. L., Zhao, Y. S., Liu, S. J., and Ren, Q. S.: Plants diversity and biomass distribution of vegetation restoration in burned area of Great Xing'an Mountains, Journal of Beijing Forestry University, 37(12), 41-47, https://doi.org/10.13332/j.1000-1522.20150025, 2015a.
Wang, Q. B., Zhang, H. J., Zhang, S. L., Wang, C. B., and Xue, Y.: Species composition and structure investigation of typical plant community in Daliangzihe National Forest Park, Protection Forest Science and Technology, 7, 14-17, 38, https://doi.org/10.13601/j.issn.1005-5215.2015.07.006, 2015b.
Wang, Q. J., Liu, Y. T., Dai, J. G., and Luo, Y. F.: Effects of different provenances and ABT1 rooting powder on *Larix gmelinii* afforestation, Forestry Science & Technology, 25(4), 4-5, https://doi.org/10.3969/j.issn.1001-9499.2000.04.002, 2000b.
Wang, Q. Y.: Genetic variation of the provenance for *Picea koraiensis*, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2003110985.htm, last access: 8 October 2017, 2003.
Wang, W. Z., Chen, J., Zhang, B. Y., Feng, Y. L. and Wang, Y. T.: Study on the photosynthetic characteristics and the primary productivity of *Larix olgensis*, Journal of Northeast

Forestry University, 22(4), 15-21, https://doi.org/10.13759/j.cnki.dlxb.1994.04.003, 1994.
Wang, X. F., Li, G. T., Wu, H. B., Lv, L. K., and Yue, Y. J.: Research on structure and species diversity of *Larix gmelinii* community in Orqohan, Journal of Southwest Forestry University, 33(5), 69-73, https://doi.org/10.3969/j.issn.2095-1914.2013.05.012, 2013b.
Wang, X. H., Zhang, Q. L., Guo, Y. D., Xing, T. T., Wang, F., Gao, X. W., and Wu, J.: Study on the coarse root biomass and distribution characteristics about natural *Larix gmelinii* forest. Forest Resources Management, 5, 62-66, https://doi.org/10.13466/j.cnki.lyzygl. 2013.05.020, 2013c.
Wang, X. W. and Mao, Z. J.: Temporal and spatial variation in gas exchange in canopy of *Larix gmelinii* plantation, Scientia Silvae Sinicae, 43(11), 43-49, https://doi.org/10.3321/j.issn:1001-7488.2007.11.008, 2007.
Wang, X. W. and Mao, Z. J.: Practicability of 7 light responsive curve models to different plant species, Bulletin of Botanical Research, 29(1), 43-48, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-MBZW200901012.htm, last access: 8 October 2017, 2009.
Wang, Y. H., Zhou, G. S., and Jiang, Y. L.: Simulation of NPP and evaluation of ecosystem services of *Larix gmelinii* forest, Chinese Journal of Applied Ecology, 12(5), 648-652, https://doi.org/10.13287/j.1001-9332.2001.0157, 2001a.
Wang, Y. L.: Biomass, carbon density and distribution characteristics in different age *Larix gmelinii* plantations, Journal of Anhui Agricultural Sciences, 43(15), 144-145,150, https://doi.org/10.13989/j.cnki.0517-6611.2015.15.053, 2015.
Wang, Z.: Larch Forest in China, China Forestry Publishing House, Beijing, China, 73-150, 1992.
Wu, C. Y, Zhao, Y. S., Xin, Y., Kong, X. F. and Li, Z. F.: Effects of structure of the *Larix gmelinii* plantation on soil hydrology characteristic, Science of Soil and Water Conservation, 7(5), 58-62, https://doi.org/10.16843/j.sswc.2009.05.011, 2009.
Wu, X. J., Cai, T. J., Li, H., and Sheng, H. C.: Precipitation hydrochemical characteristic in virgin *Pinus koraiensis* forest and artifical *Larix gmelinii* forest in Liangshui National Nature Reserve, Science of Soil and Water Conservation, 6(6), 37-42, https://doi.org/10.16843/j.sswc.2008.06.007, 2008.
Wu, X. J., Jia, E. L., and Li, J. C.: Preliminary study on the regeneration before clear cutting in *Larix gmelinii* plantation, Forest Science and Technology, 8, 24-26, https://doi.org/10.13456/j.cnki.lykt.1983.08.010, 1983.
Wu, X. L.: Interannual variability of temperature, precipitation and sunshine hours in Wuerqihan Forest Region, Heilongjiang Agricultural Sciences, 10, 113-116, https://doi.org/10.11942/j.issn1002-2767.2015.10.0113, 2015.
Wu, Z. C.: Introduction and afforestation of *Larix olgensis*, Journal of Inner Mongolia Forestry, 12, 20-21, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMLY198612016.htm, last access: 8 October 2017, 1986.
Xia, C. C., Liu, Z. L., Wang, Q. C., and Li, K. L.: Comparison of the growth and biomass production of monoculture and mixed stands of *Larix gmelinii* and *Betula platyphylla*, Journal of Northeast Forestry University, 40(10), 1-3, https://doi.org/10.13759/j.cnki.dlxb.2012.10.021, 2012.
Xiao, S. L. and Yang, J. L.: Individual tree aboveground biomass of *Larix gmelinii* natural

forest in the Northern Greater Khingan Mountains, Scientia Silvae Sinicae, 50(8), 22-29, https://doi.org/10.11707/j.1001-7488.20140804, 2014.

Xiao, Y., Man, X. L., and Fan, J. F.: Soil fertility and water conservation function of main forest types in Gongbiela River Watershed, Journal of Northeast Forestry University, 34(1), 28-30, https://doi.org/10.13759/j.cnki.dlxb.2006.01.010, 2006.

Xin, Y., Zhao, Y. S., and Pan, B. Y.: Influence of *Larix gmelinii* plantations at Eastern Mountainous Region in Heilongjiang Province on water quality, Science of Soil and Water Conservation, 4(2), 29-33, https://doi.org/10.16843/j.sswc.2006.02.006, 2006.

Xin, Y., Zou, M. L., and Zhao, Y.S.: Soil water storage capacity and infiltration properties of *Larix gmelinii* plantation in different growth age after burned in Greater Khingan Mountains, Chinese Journal of Soil Science, 45(4), 823-829, https://doi.org/10.19336/j.cnki.trtb.2014.04.009, 2014.

Xu, H. C. and Du, Y. J.: Amount of falling seed and seedling occurrence dynamics in primitive *Larix gmelinii* forest, Scientia Silvae Sinicae, 29(4), 298-306, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE199304001.htm, last access: 8 October 2017, 1993.

Xu, H. C., Yu, R. Y., Guo, Z. L., Zhou, C. H., and Ma, D. G.: Study on germination conditions of *Larix gmelinii* seedlings in Great Xing'an Mountains, Journal of Beijing Forestry University, 12, 15-25, https://doi.org/10.13332/j.1000-1522.1990.s4.004, 1990.

Xu, H. L., Man, X. L., and Sheng, H. C.: Water conservation function of different types of *Larix gmelinii* forests in Da Hinggan Mountains, Research of Soil and Water Conservation, 18(4), 92-96, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-STBY201104020.htm, last access: 8 October 2017, 2011a.

Xu, H. L., Man, X. L., and Sheng, H. C.: Comprehensive evaluation on water conservation function of the different types of larch natural forests in Da Hinggan Mountains, Chinese Agricultural Science Bulletin, 29(19), 74-78, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-ZNTB201319017.htm, last access: 8 October 2017, 2013.

Xu, J. L.: Study on Larix biomass model and biomass estimation in Dongzhelenghe, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10022-2009161609.htm, last access: 8 October 2017, 2009.

Xu, S. Z. and Han, M. Z.: Regeneration rule and management decision of *Larix gmelinii* in Genhe regions, in Zhou, X. F. ed., Long-Term Research on China's Forest Ecosystems, 709-716, Northeast Forestry University Press, Harbin, China, 1994.

Xu, X. T., Wan, D. Y., Wei, D. F., and Zhou, J. Y.: Growth characteristics and techniques for fast growing and high yield of larch, Inner Mongolia Forestry Investigation and Design, 34(6), 54-55, https://doi.org/10.13387/j.cnki.nmld.2011.06.057, 2011b.

Xu, Y. B., Zhou, Y. Q., Wu, R. X., Liu, Z. S., and Yu, C. H.: Afforestation techniques of *Larix gmelinii* container seedlings, Forestry Science & Technology, 21(3), 5-7, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ603.001.htm, last access: 8 October 2017, 1996.

Xu, Y. Q.: Study on the geographic variations and selection for the optimal provenances of *Larix gmelinii*, Forest Inventory and Planning, 3, 15-23, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-LDGH199303004.htm, last access: 8 October 2017, 1993.

Xue, Z. H.: Investigation of larch plantation in Daxing'anling Mountains, Inner Mongolia Forestry Science and Technology, 2, 18-21, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198702003.htm, last access: 8 October 2017, 1987a.

Xue, Z. H.: Study on tending and thinning of natural young and middle-age *Larix gmelinii* forest, Inner Mongolia Forestry Science and Technology, 4, 1-5, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198704000.htm, last access: 8 October 2017, 1987b.

Xue, Z. H.: Tending and Thinning of *Larix gmelinii* plantation, Inner Mongolia Forestry Science and Technology, 3, 20-23, 30, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK199003004.htm, last access: 8 October 2017, 1990.

Xue, Z. H.: Cultivation and management techniques of fast-growing and high-yielding *Larix gmelinii* Forests, Inner Mongolia Forestry Science & Technology, 1, 11-18, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK199301004.htm, last access: 8 October 2017, 1993.

Yang, B. L., Jia, Q. B., Zhang, H. G., and Zhang, L.: Research on hybrid larch superior families selection, Forestry Science & Technology, 39(2), 1-4, https://doi.org/10.3969/j.issn.1001-9499.2014.02.001, 2014.

Yang, C. P.: Study on the provenance of Dahurian larch, Science Press, Beijing, China, 2009.

Yang, C. P, Jiang, J., Tang, S. S., Li, J. Y. and Wang, H. R.: The provenance test of 21- year old *Larix gmelinii* at Maoershan Area, Journal of Northeast Forestry University, 30(6), 1-5, https://doi.org/10.13759/j.cnki.dlxb.2002.06.001, 2002.

Yang, C. P., Yang, S. W., Xia, D. A., Liu, G. F., Lv, Q. Y. and Zhang, P. G.: Study on the provenance test of Dahurian Larch (II) Selection of the best provenance, Journal of Northeast Forestry University, 19, 68-76, https://doi.org/10.13759/j.cnki.dlxb.1991.s2.011, 1991.

Yang, H. T. and Guo, Y. X.: A primary report on the provenance test of *Larix gmelinii* and *Larix olgensis* in Huanan forest region, Journal of Northeast Forestry University, 18, 65-71, https://doi.org/10.13759/j.cnki.dlxb.1990.s2.010, 1990.

Yang, K., Zhu, J. J., Gu, J. C.,Yu, L. Z., and Wang, Z. Q.: Changes in soil phosphorus fractions after 9 years of continuous nitrogen addition in a *Larix gmelinii* plantation, Annals of Forest Science, 72, 435-442, https://doi.org/10.1007/s13595-014-0444-7, 2015.

Yang, L. J., Zhang, C. J., Li, C. H., Jiang, X. L., and Pan, B. L.: Early selection of superior clone of *Larix gmelinii*, Protection Forest Science and Technology, 4, 35-37, https://doi.org/10.13601/j.issn.1005-5215.2006.04.013, 2006.

Yang, S. W. and Shang, L. T.: Research on selection of larch species, Journal of North-Eastern Forestry Institute, 11(4), 50-56, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198304006.htm, last access: 8 October 2017, 1983.

Yang, S. W., Wang, Q. Y., Liu, G. F., Yang, C. P., Xia, D. A., and Bao, L. F.: Analysis on the effects of genetic improvement and the breeding program of larch, Journal of Northeast Forestry University, 20(1), 1-8, https://doi.org/10.13759/j.cnki.dlxb.1992.01.001, 1992.

Yang, S. W., Yang, C. P., Wang, Q. Y., Liu, G. F., Xia, D. A., and Zhang, P. G.: The provenance selection of the important tree species in the northeast part of China and Inner Mongolia Autonomous Region, Journal of Northeast Forestry University, 21(4), 1-7, https://doi.org/10.13759/j.cnki.dlxb.1993.04.001, 1993.

Yang, S. W., Yang, C. P., Zhang, S. Y., Peng, H. M., Li, J. T., Xia, D. A., Liu, G. F., and Wang, H. R.: A study on the provenance test of *Larix gmelinii* outside of distribution area, Journal of Northeast Forestry University, 18, 1-8, https://doi.org/10.13759/j.cnki.dlxb.1990.s2.002, 1990.

Yang, Y. B. and Liu, Y. Y.: Discussion about planting fast-growing and high-yield plantation in the south of Daxing'anling City, Forestry Science and Technology Information, 37(2), 1-2, https://doi.org/10.3969/j.issn.1009-3303.2005.02.001, 2005.

Yao, C. Y., Wang, Q. J., and Gao, G. Y.: Study on growth status of introduced *Larix olgensis* in Xingan League of Inner Mongolia, Inner Mongolia Forestry Science & Technology, 1, 17-22, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK501.004.htm, last access: 8 October 2017, 1995.

Yao, G. Q., Chi, G. Q., Dong, Z. Q., Hu, W. L., and Jiang, J. B.: A study on the growth and economic effectiveness of four kinds of planted Larch Forest, Journal of Shenyang Agricultural University, 20, 2l7-222, available at: http://www.cqvip.com/QK/93761X/198903/129531.html, last access: 8 October 2017, 1989.

Yao, Y. J., Chen, X. W., and Li, C. S.: Studies on biomass and productivity for Dahuian larch plantations, Journal of Northeast Forestry University, 5(2), 13-16, https://doi.org/10.1007/BF02843225, 1994.

Yu, B. and Wang, L. M.: Analysis on classified stem growing characteristics of *Larix gmelinii* natural forest, Forest Research, 20(4), 452-457, https://doi.org/10.13275/j.cnki.lykxyj.2007.04.014, 2007.

Yu, B., Wu, J. S. G. L.,Wang, B. T., and Wang, L. M.: Analysis on classified stem transformation characteristic of *Larix gmelinii* natural forest at Daxingan Mountains, Acta Ecologica Sinica, 28(11), 5750-5757, https://doi.org/10.3321/j.issn:1000-0933.2008.11.062, 2008.

Yu, B.,Wu, J. S. G. L.,Wang, B. T., Wang, L. M., and Da, B. X. L. T.: Study on the distribution pattern characteristics of different stand structures in *Larix gmelinii* natural forest, Forest Research, 23(1), 83-88, https://doi.org/10.13275/j.cnki.lykxyj.2010.01.013, 2010.

Yu, B., Zang, Q. L.,Wang, L. M., and Wu, J. S. G. L.: Characteristics of biomass and productivity in *Larix gmelinii* natural forests with different stand structures, Journal of Zhejiang A & F University, 28(1), 52-58, https://doi.org/10.3969/j.issn.2095-0756.2011.01.009, 2011.

Yu, B. J. and Yang, C. P.: Study on provenance test of 5 year old *Larix olgensis*, Journal of Northeast Forestry University, 16(3), 27-33, https://doi.org/10.13759/j.cnki.dlxb.1988.03.005, 1988.

Yu, S. L. and Yuan, X. G.: Studies on selection of fine families of Hybrid Larch based on growth traits in seedling stage, Inner Mongolia Forestry Investigation and Design, 32(2), 114-117, https://doi.org/10.13387/j.cnki.nmld.2009.02.025, 2009.

Yuan, G. X., Li, H. X., Chen, G. F., Lou, Y. M., and Mou, Z. J.: Analysis on the introduction of *Pinus banksiana Lamb*, Forestry Science & Technology, 26(5), 9-10, https://doi.org/10.3969/j.issn.1001-9499.2001.05.004, 2001.

Yue, Y. J., Gao, R. H., Li, G. T., Jin, J., and Wang, X. F.: Research on community structure and species diversity of *Larix gmelinii* community in Jinhe Forestry Bureau, Journal of

Inner Mongolia Agricultural University, 35(2), 43-47, https://doi.org/10.16853/j.cnki.1009-3575.2014.02.010, 2014.

Yue, Y. J., Qu, L. C., Song, L. J., Liu, M. D., Zeng, Y. E., Li, G. T., and Wang, F. S.: Distribution patterns and diversity of species of *Larix gmelinii* in Kuduer, Journal of Inner Mongolia Agricultural University, 34(1), 42-47, https://doi.org/10.16853/j.cnki.1009-3575.2013.01.009, 2013.

Zhai, M. P., Yi, W. L., Jia, L. M., and Ma, L. Y.: Study on the characteristics of *Betula platyphylla-Larix gmelinii* Mixed Forest in Xilinji Forest Region, Journal of Beijing Forestry University, 12, 86-91, https://doi.org/10.13332/j.1000-1522.1990.s3.010, 1990a.

Zhai, M. P., Yi, W. L., Jia, L. M., Ma, L. Y., Sun, Y. X., Liu, Z. Y., and Wang, Y. G.: Investigation of *Betula platyphylla-Larix gmelinii* Mixed Forest in Xilinji Forest Region, Journal of Beijing Forestry University, 12, 78-85, https://doi.org/10.13332/j.1000-1522.1990.s3.009, 1990b.

Zhan, J. and Yang, C. P.: Study on the provenance test of Dahurian larch (*Larix gmelinii*), Journal of Northeast Forestry University, 17(6), 22-30, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198906003.htm, last access: 8 October 2017, 1989.

Zhang, G. L., Qu, B. C., Zhang, Y. H., and Wang, H.: Growth analysis and model of *Larix gmelinii* plantation, Protection Forest Science and Technology, 4, 41-43, https://doi.org/10.13601/j.issn.1005-5215.2012.04.056, 2012.

Zhang, H. D. and Zhou, M.: Nutrient characteristics of *Larix gmelinii* forest ecosystem in cold temperate zone, Journal Liaoning Forestry Science & Technology, 3, 5-8, 22, https://doi.org/10.3969/j.issn.1001-1714.2014.03.002, 2014.

Zhang, J.: Study on community structure, biomass and carbon storage of *Larix gmelini* plantation, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10022-2008085203.htm, last access: 8 October 2017, 2008a.

Zhang, J., Sun, Y. J., and Xu, J. L.: Research on growing process of *Larix gmelinii* plantation in Northeast of China, Journal of Northwest Forestry University, 23(6), 179-181, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-XBLX200806045.htm, last access: 8 October 2017, 2008.

Zhang, J. L., Mao, Y. Q., Wang, F. S., Liu, L., Wu, C. J., Zhang, H. P., Yuan, L. D., Zhao, W., and Wang, F. J.: The progeny testing and fine family selection of *Larix gmelinii* and *Larix olgensis*, Protection Forest Science and Technology, 3, 22-26, https://doi.org/10.13601/j.issn.1005-5215.1994.03.003, 1994.

Zhang, L.: Study on stability of families and selection of superior families of hybrid larch. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1011146424.htm, last access: 8 October 2017, 2011.

Zhang, L. Y.: A study on individual tree health assessment of nature *Larix gmelinii* in Daxing'anling, Forestry Science and Technology Information, 45(1), 41-43. https://doi.org/10.3969/j.issn.1009-3303.2013.01.018, 2013.

Zhang, Q. B. and Fang, L.: A study of problems of planting fast-growing and high-yield plantation in secondary forest site on the south slope in Great Xing'an Mountain, Journal of Inner Mongolia Forestry College, 20(3), 37-42, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMGL803.007.htm, last access: 8 October 2017, 1998.

Zhang, S. L., Zhang, H. J., Zhang, W., Guo, Y., Li, X. Z., Zhang, Q., Lu, Y., and Yao, S. P.: Comparison of water-holding characteristics and biomass of litter under different forest types in Xiaoxingan Mountains, Bulletin of Soil and Water Conservation, 35(4), 85-90, https://doi.org/10.13961/j.cnki.stbctb.2015.04.017, 2015a.

Zhang, S. L., Zhang, H. J., Zhang, W., Wang, C. B., Deng, J. H., Song, Y. B., and Zhang, Q.: Soil hydrological properties of main vegetation in Down South of Xiao Xing'an Mountains, Research of Soil and Water Conservation, 22(1), 140-145, https://doi.org/10.13869/j.cnki.rswc.2015.01.027, 2015b.

Zhang, T. W., Yuan, Y. J., Wei, W. S., and Yu, S. L.: The tree-ring chronologies from Moerdaoga in the northeast of Inner Mongolia, Journal of Arid Land Resources and Environment, 23(12), 177-182, https://doi.org/10.13448/j.cnki.jalre.2009.12.008, 2009.

Zhang, X. J.: Effect of close-to-nature transformation of Larch Monoculture stands on regeneration and plant diversity. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2010242257.htm, last access: 8 October 2017, 2010.

Zhang, X. J., Wang, Q. C., Wang, S. L., and Sun, Q.: Effect of the close-to-nature transformation of *Larix gmelinii* pure stands on plant diversity of understory vegetation in Xiaoxing'an Mountains of China, Scientia Silvae Sinicae, 47(1), 6-14, https://doi.org/10.11707/j.1001-7488.20110102, 2011.

Zhang, Y. Q.: Transpiration characteristic of Dahurian Larch trees transplanted from a latitudinal gradient. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2009133900.htm, last access: 8 October 2017, 2008b.

Zhao, H. X., Wang, Y. H., Chai, Y. X., Hu, C. X., Li, F. R., and Fan, W. Y.: On forest growth of northern slope of Daxing'an mountains (Tahe Forest Bureau), Journal of Northeast Forestry University, 24(6), 1-8, https://doi.org/10.13759/j.cnki.dlxb.1996.06.001, 1996.

Zhao, P. W.: Studies on litterfall dynamics and nutrient release regularity of Larix gmelinii in Great Xingan Mountains. Thesis, https://doi.org/10.7666/d.y1473822, 2009.

Zhao, S. J., Yang, J. M., and Bi, X. M.: Study on the fast growing and high yield of larch species and the relationships with their introduction districts, Journal of Hebei Forestry Science and Technology, 2, 2-7, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-HBLK199202001.htm, last access: 8 October 2017, 1992.

Zhao, X. M., Xu, S. T., Ren, J. Z., Wang, M. Z., Yang, S. M., and Jiang, S. W.: A study on provenance test of *Pinus sylvestris L.var mongolica* litvin on Korqin sandy land, Journal of Inner Mongolia Forestry College, 21(3), 44-49, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-NMGL199903008.htm, last access: 8 October 2017, 1999.

Zhou, B. D., Hao, W. K., Zou, Q., Zhao, K. M., Liu, Q. L., and Chen, Y. D.: Study on artificial regeneration in a clear-cut forest of Xiaoxingan Mountains in Liangshui region, Journal of North-Eastern Forestry College, 4, 19-35, available at: http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY196500001.htm, last access: 8 October 2017, 1965.

Zhou, B. Q.: Advices on secondary forest culture on the south slope in Great Xing'an Mountain, Inner Mongolia Forestry Investigation and Design, 18-20, https://doi.org/10.13387/j.cnki.nmld.1998.s1.008, 1998.

Zhou, H. X., Zhang, Y. D., Sun, H. L., and Wu, S. Y.: Soil respiration in temperate secondary forest and *Larix gmelinii* plantation in Northeast China, Chinese Journal of Applied Ecology, 18(12), 2668-2674, https://doi.org/10.13287/j.1001-9332.2007.0477, 2007.

Zhou, J. Q., Liu, X. D., and Guo, H. W.: Surface fuel loading and relevant influencing factors of main forest types in southern Daxing'anling, Journal of Northwest A&F University(Natural Science Edition), 42(6), 131-137, https://doi.org/10.13207/j.cnki.jnwafu.2014.06.010, 2014.

Zhu, H. Y.: Main characters variation among provenances of Fraxinus mandshurica Rup. Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-2009134220.htm, last access: 8 October 2017, 2008.

Zhu, S. Y., Zhou, B., Mao, Z. J., Wang, X. W., and Sun, Y. F.: Space time dynamics of fine root biomass of six forests in Maoershan forest region, Scientia Silvae Sinicae, 42(6), 13-19, https://doi.org/10.3321/j.issn:1001-7488.2006.06.003, 2006.

Zhu, Y.: Forest health assessment of natural Larix gmelinii in Great Xing'an Mountains of China, Thesis, available at: http://cdmd.cnki.com.cn/Article/CDMD-10225-1013371221.htm, last access: 8 October 2017, 2013.

Zhu, Y., Liu, Z. G., and Jin, G. Z.: Health assessment of individual trees in natural *Larix gmelinii* forest in Great Xing'an Mountains of China, Chinese Journal of Applied Ecology, 24(5), 1320-1328, https://doi.org/10.13287/j.1001-9332.2013.0294, 2013.

Zou, L., Tang, Q. M., and Wang, Y.: Ecological distribution of soil microorganism in pure and mixed forests of *Pinus sylvestris var. mongolica* and *Larix gmelini*, Journal of Northeast Forestry University, 38(11), 63-64, 79, https://doi.org/10.13759/j.cnki.dlxb.2010.11.029, 2010.

Zou, M. L., Xin, Y., and Zhao, Y. S.: The influence of vegetation restoration on carbon storage of burned Area in Greater Xing' an Mountains, Journal of Soil and Water Conservation, 27(5), 133-137, 143, https://doi.org/10.13870/j.cnki.stbcxb.2013.05.046, 2013.