Earth Syst. Sci. Data Discuss., https://doi.org/10.5194/essd-2017-99-SC1, 2017 © Author(s) 2017. This work is distributed under the Creative Commons Attribution 4.0 License.



ESSDD

Interactive comment

Interactive comment on "Growth characteristics of natural and planted Dahurian larch in northeast China" by Bingrui Jia and Guangsheng Zhou

F. Achard

frederic.achard@ec.europa.eu

Received and published: 27 October 2017

I would suggest to refer to two papers in the introduction of this interesting paper: Achard et al 2006 and Bastin et al 2017. Achard et al 2006 presents an overview of areas of rapid forest-cover change in Boreal Eurasia, including two active hot spot zones (selective logging) in the Heilongjiang province: Da Hinggan Ling prefecture and Xiao Hinggan Ling prefecture, both including Larix forests. Bastin et al (2017) present a global assessment of forests in dryland zone. The dryland zone used in Bastin et al (2017) covers a large paper of the study zone of this manuscript.

Achard F, et al 2006. Areas of rapid forest cover change in boreal Eurasia. Forest Ecology and Management 237: 322–334 Bastin JF et al, The extent of forest in dryland

Printer-friendly version

Discussion paper



biomes Science 356, 635–638 (2017)

Interactive comment on Earth Syst. Sci. Data Discuss., https://doi.org/10.5194/essd-2017-99, 2017.

ESSDD

Interactive comment

Printer-friendly version

Discussion paper

