



1 **Growth characteristics of natural and planted Dahurian larch in northeast China**

2 Bingrui Jia^{1*} and Guangsheng Zhou^{2*}

3 ¹*State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese*

4 *Academy of Sciences, Beijing 100093, China,*

5 ²*Chinese Academy of Meteorological Sciences, Beijing 100081, China,*

6 ^{*}Corresponding author:

7 Bingrui Jia, Guangsheng Zhou

8 Institute of Botany, Chinese Academy of Sciences,

9 20 Nanxincun, Xiangshan, Haidian District, Beijing 100093, China

10 E-mail: jiabingrui@ibcas.ac.cn, gszhou@ibcas.ac.cn

11 Tel: 86-10-62836289, Fax: 86-10-82595962

12 **Abstract.** Dahurian larch (*Larix gmelinii* Rupr.) is the dominant species in both natural and
13 planted forests in northeast China, which situated in the southernmost part of the global
14 boreal forest biome and undergoing the greatest climatically induced changes. Published
15 studies (1965–2015) on tree aboveground growth of *Larix gmelinii* forests in northeast China
16 were collected in this study, critically reviewed, and a comprehensive growth data set was
17 developed from 123 sites, which distributed between 40.85° N and 53.47° N in latitude,
18 between 118.20° E and 133.70° E in longitude, between 130 m and 1260 m in altitude. The
19 data set was composed of 776 entries, including growth data (mean tree height, mean DBH,
20 mean tree volume and/or stand volume) and the associated information, i.e., geographical
21 location (latitude, longitude, altitude, aspect and slope), climate (mean annual temperature
22 (MAT) and mean annual precipitation (MAP)), stand description (origin, stand age, stand
23 density and canopy density), and sample regime (observing year, plot area and number). It



24 would provide quantitative references for plantation management practices and boreal forest
25 growth prediction under future climate change. The data set is freely available for
26 noncommercial scientific applications, and the DOI for the data is <https://doi.org/10.1594/PANGAEA.880984>.
27
28

29 **1 Introduction**

30 Boreal forests, the second largest biome in the world, cover about one-third of the Earth's
31 forest area (Keenan et al., 2015). Dahurian larch (*Larix gmelinii* Rupr.) is a dominant tree
32 species in Chinese boreal forest, which is distributed primarily in northeast China. *Larix*
33 *gmelinii* forest is also the predominant timber source in China, occupying 55% of Chinese
34 boreal forest area and accounting for 75% of Chinese boreal forest volume (Xu, 1998; Zhou
35 et al., 2002). *Larix gmelinii* forest is situated in the southernmost part of the global boreal
36 forest biome (Shugart et al., 1992) and undergoing the greatest climatically induced changes.
37 Thus understanding the growth characteristics of *Larix gmelinii* forest in China are of critical
38 need for management and prediction under future climate change.

39 With increased greenhouse effective and climate warming in recent years, forest carbon
40 sink is payed more and more attention by the world. Forestation is the main measure to offset
41 the greenhouse gas emission and increase carbon sink. China has the largest area of forest
42 plantations in the world, approximately 79 million ha or one-fourth of world total (FAO,
43 2015; Payn et al., 2015). *Larix gmelinii* is an important fast-growing and cold-tolerant tree
44 species used in forestation in northeast China. *Larix gmelinii* is usually planted after fire or
45 logging. The growth rates of *Larix gmelinii* plantations are important indexes in the
46 assessment of forest recovery processes and carbon sequestration potentials, which could
47 supply strategies for post-fire or post-harvest management. The data set can provide basis for
48 evaluating and predicting the carbon sequestration and its potential of the forestation



49 activities.

50 Relating the easily measured variables (e.g. tree height, diameter) to other structural and
51 functional characteristics, is the most common and reliable method for estimating forest
52 biomass, net primary production and biogeochemical budgets (Luo, 1996; Fang et al., 2001).
53 For larch forests in northeast China, synthesis studies mainly focused on biomass and net
54 primary production with a small quantity of samples, for example, N=28 (Luo, 1996), N=17
55 (Wang et al., 2001a), N=18 (Wang et al., 2001b; Zhou et al., 2002), N=36 (Wang et al., 2005),
56 N=83 (Wang et al., 2008). However, large numbers of growth measurements (e.g. tree height,
57 DBH, volume) have scarcely been studied systematically at the large scale. Therefore, a
58 comprehensive growth data set (N=776) of Dahurian larch in northeast China was developed
59 in this paper.

60

61 2 Data and methods

62 2.1 Research origin descriptors

63 (1) Identity: Growth data set of natural and planted Dahurian larch in northeast China,
64 version 1.0

65 (2) Originators:

66 Bingrui Jia, State Key Laboratory of Vegetation and Environmental Change, Institute of Botany,
67 Chinese Academy of Sciences, Beijing 100093, China;
68 Guangsheng Zhou, Chinese Academy of Meteorological Sciences, Beijing 100081, China.

69 (3) Period of Study: From January 1965 to December 2015


70 (4) Objectives: We conducted a complete literature and book review of published studies on
71 tree height, DBH, and/or volume in *Larix gmelinii* forests including natural and planted
72 forests in order to develop a growth data set. The data set can be used to analyze growth



73 characteristics of *Larix gmelinii* forests and assess their potential productivity in future
74 climate warming.

75 **2.2 Site description**

76 **(1) Site type:** Data were derived from 123 study sites in northeastern China. This region
77 includes Heilongjiang, Liaoning, Jilin provinces and the eastern part of Inner Mongolia
78 Autonomous Region (Fig. 1).



79
80 Figure 1. Study sites of *Larix gmelinii* across northeastern China, including Heilongjiang,
81 Liaoning, Jilin provinces and the eastern part of Inner Mongolia Autonomous Region. Open
82 circles represent natural forests and dots represent planted forests.

83 **(2) Habitat:** *Larix gmelinii* forest is naturally distributed in Great Xing'an Mountains of
84 northeastern China. *Larix gmelinii* was usually planted in barren hills, post-fire or
85 post-harvest area in northeast China.



86 **(3) Climate:** The climate in northeast China is controlled by the high latitude East Asia
87 monsoon, changing from cool temperate to temperate zones from north to south, and from
88 semi-arid to humid zones from west to east. Mean annual temperature (MAT) for these sites
89 ranged from -6.1 to 7.0 °C, and mean annual precipitation (MAP) from 355 to 926 mm.

90 **2.3 Data sources**

91 Published studies (1965-2015) were collected from available online full-text databases,
92 including China Knowledge Resource Integrated Database (<http://www.cnki.net/>), China
93 Science and Technology Journal Database (<http://www.cqvip.com/>), Wanfang Data
94 Knowledge Service Platform (<http://www.wanfangdata.com/>), ScienceDirect (<http://www.sciencedirect.com/>), ISI Web of Science (<http://isiknowledge.com/>) and Springer Link
95 (<http://link.springer.com/>). The different combination of the terms “Dahurian larch (or *Larix*
96 *gmelinii*)” with “height”, “diameter at breast height (or DBH)”, “tree volume” and “stand
97 volume” were searched in full text. Meanwhile, we also looked up the related books (e.g. Ma,
98 1992; Wang, 1992; Zhou, 1994; Yang, 2009). We attempted to compile a complete growth
99 data set of natural and planted Dahurian larch in northeastern part of China (between 40.85 °
100 N and 53.47 °N; between 118.20 °E and 133.70 °E).

102 **2.4 Data collection criteria**

103 A critical review of the literatures collected from the above-mentioned sources was
104 conducted to obtain reliable growth data using the following criteria:

105 **(1) Scope:** The objective of this study was to provide the data for understanding growth
106 characteristics of *Larix gmelinii* natural forests (pure *Larix gmelinii* or its proportion more
107 than 50%) and monoculture plantations. Forest stands included in the data set were restricted
108 to those not recently disturbed by logging, fire, or insect pests. Additionally, the following
109 small numbers of special types were excluded: (i) *Larix gmelinii* afforestation in wetland (Li
110 et al., 1985; Song & Li, 1990; Huang, 2011; Cui et al., 2013), pastureland (Duan, 2005), or



111 abandoned mine land (Yang et al., 2013); (ii) hybrid test between *Larix gmelinii* and other
112 larch (Deng et al., 2010; Zhang et al., 2005); (iii) low-yield stands in hard environment, e.g.
113 igneous rock forest (Wang et al., 1979), old man forest (Wang et al., 1991).

114 **(2) Study design and sampling:** Tree height and DBH were averaged from the measurement
115 values of all trees in plots or with random/systematic sampling method. Tree regeneration
116 layer, generally below 5 cm in DBH or 1.3 m in height, was neglected in sampled plots. Stem
117 volume of individual tree was computed from felled-wood samples or local tree volume
118 equations. Stand volume was usually calculated by multiplying mean individual volume with
119 stand density. Besides the growth data (i.e. height, DBH and/or volume), the necessary
120 information should be provided in the original sources, e.g., study site, stand origin, stand age,
121 etc.

122 **(3) Quality control:** The data quality has been carefully reviewed by the authors. Data has
123 undergone substantial checking, for example, cross-check for the relevant information from
124 different sources, preliminary correlation analysis among growth variables.

125 Consequently, 776 records that met the above criteria were selected to develop a
126 comprehensive growth data set of Dahurian larch in China. The data set includes growth
127 characteristics of Dahurian larch (i.e. mean tree height (m), mean DBH (cm), mean tree
128 volume (10^{-3} m 3) and/or stand volume (m 3 /ha)). In addition, associate information was
129 included, if available in original sources or ascertainable from other relevant literatures, i.e.,
130 geographical location (province location and locality name of study site, latitude (°),
131 longitude (°), altitude (m), aspect and slope (°), stand description (origin, stand age (years)),
132 stand density (trees/ha) and canopy density), climate (mean annual temperature (MAT, °C)
133 and mean annual precipitation (MAP, mm)), and sample regime (observing year, plot size and
134 number).

135 **2.5 Data structural descriptors**



136 Table 1 Variable information in the data set.

Column code	Definition	Unit	Number	Range
ID	Unique identification number of each record	N/A	776	1—776
Province	Province location of study site	N/A	776	N/A
Study site	Locality name of study site	N/A	123	N/A
Latitude	Latitude of study site	°	776	40.85—53.47
Longitude	Longitude of study site	°	776	118.20—133.70
Altitude	Altitude of study site	m	776	130—1260
Aspect	Slope direction of study site, including flat slope (FL), sunny slope (SU: South), half-sunny slope (HSU: West, Southwest, Southeast), shady slope (SH: North) and half-shady slope (HSH: East, Northwest, Northeast)	N/A	300	N/A
Slope	Slope degree of study site	°	357	0—60
Origin	Stand origin was classified into natural and planted forests	N/A	776	N/A
MAT	Mean annual temperature, from original study or other related reference	°C	776	-6.1—7.0
MAP	Mean annual precipitation, from original study or other related reference	mm	776	355—926
Age	Stand age, which is generally defined as age since germination in natural forest and since planting in planted forest. Stand age is usually obtained from historical records or tree rings.	years	776	1—280
Height	Mean tree height	m	670	0.24—29.40
DBH	Mean diameter at breast height, base diameter was only given in some young forests and marked with ^B	cm	661	0.70—34.89
V _{tree}	Mean tree volume, the estimated tree volume data from the two-variable larch equation were marked with ^E	10 ⁻³ m ³ /tree	696	0.04—935.73
V _{stand}	Stand volume, the estimated stand volume data from the estimated tree volume and stand density were marked with ^E	m ³ /ha	590	0.07—975.32
Density	Stand density/Canopy density, planting density was only given in some studies and marked with ^P	trees/ha %	656 150	213—13275 0.2—1.0
Area	Plot area	m ²	397	50—10000
Plot	Plot numbers, i.e. replications	N/A	573	1—25
Year	Investigation year	N/A	533	1954—2014
Reference	Data sources, the sources used to supplement climate information lacking in the original publications were added asterisks.	N/A	226	1965—2015

137

138 3 Data estimates and evaluation

139 3.1 Geographical location

140 Google Earth (Version: 7.1.8.3036) was used to estimate latitude, longitude, and/or altitude
 141 when the geographic coordinates was unavailable in original sources.


142 3.2 Tree and stand volume

143 The missing tree and stand volumes were estimated with the available information (mean
 144 DBH, mean tree height and stand density). Stem volume of individual tree was calculated




145 with the larch equation in northeast China ($V_{tree} = 0.000050168241 \text{ DBH}^{1.7582894} \text{ H}^{1.1496653}$),
146 the best method recommended by the national standard of China: Tree volume tables (LY
147 208-77) (Forestry Administration of China, 1978; Liu, 2017). Meanwhile, available 590 pairs
148 of mean tree height and mean DBH in the data set were used to establish the linear H-DBH
149 correlation (see Fig. 2). To calculate tree volume from only one known variable
150 of DBH, tree height was firstly calculated with the linear H-DBH equation in Fig. 2. The
151 estimated stand volume was determined by multiplying the estimated tree volume with stand
152 density.

153 The accuracy of tree volume was assessed by comparing the extracted data from references
154 with the calculated data from the above-mentioned two-variable tree volume equation (Fig. 3).
155 The coefficient of determination (R^2) was 0.9724 and the slope was 1.0737. Therefore, we
156 were confident in applying the larch volume equation to interpolate tree and stand volume
157 data in this study.



158
159 Figure 2. Relationship between mean tree height and diameter at breast height in the data set.



160
161 Figure 3. Comparison of available tree volume from references with simulated values from
162 the two-variable tree volume equation ($V_{tree} = 0.000050168241 \text{ DBH}^{1.7582894} \text{ H}^{1.1496653}$). Open
163 circles ($N=317$): tree height (H) and diameter at breast height (DBH) were available in the
164 references, solid circles ($N=37$): DBH was only available in the references and H was
165 estimated with the H-DBH correlation from Fig. 2.
166

167 **Acknowledgements.** We would like to acknowledge the many scientists and researchers
168 whose field measurements made this work possible. This research was supported by the
169 National Key Research and Development Program of China (2017YFC0503906) and China
170 Special Fund for Meteorological Research in the Public Interest (GYHY201406034).

171 **References**

- 172 Cui, W., Mu, C. C., Lu, H. C., Bao, X., and Wang, B.: Effects of draining for forestation on
173 carbon storage of wetland ecosystem in Daxing'an Mountains of northeastern China,
174 Journal of Beijing Forestry University, 35(5), 28-36, <https://doi.org/10.13332/j.1000-1522>.



- 175 2013.05.020, 2013.
- 176 Deng, J. F., Zhang, H. G., Zhang, L., Guan, C. Y., and Zhang, L.: Genetic variation of
177 17-year-old hybrid larch and its superior family selection, Journal of Northeast Forestry
178 University, 38(1), 8-11, <https://doi.org/10.13759/j.cnki.dlxz.2010.01.017>, 2010.
- 179 Duan, W. B.: Suitable types of soils for main tree species of Shelterbelts on pastureland,
180 Chinese Journal of Soil Science, 36(6), 871-876, <https://doi.org/10.19336/j.cnki.trtb.2005.06.012>, 2005.
- 181 Fang, J., Chen, A., Peng, C., Zhao, S., Ci, L.: Changes in forest biomass carbon storage in
182 China between 1949 and 1998, Science, 292(5525), 2320-2322, <https://doi.org/10.1126/science.1058629>, 2001.
- 183 FAO: Global Forest Resources Assessment 2015, UN Food and Agriculture Organization,
184 Rome, Italy, available at: <http://www.fao.org/3/a-i4808e.pdf>, last access: 8 October 2017,
185 2015.
- 186 [Forestry Administration of China](#): Tree Volume Tables (National standard # LY 208-77),
187 Standard Press of China, Beijing, China, 1978.
- 188 Huang, D. Q.: Biomass and carbon storage estimation of *Larix gmelinii*, Thesis, available at:
189 <http://d.wanfangdata.com.cn/Thesis/Y2026419>, last access: 8 October 2017, 2011.
- 190 Keenan, R. J., Reams, G. A., Achard, F., de Freitas, J. V., Grainger, A., and Lindquist, E.:
191 Dynamics of global forest area: Results from the FAO Global Forest Resources
192 Assessment 2015, Forest Ecology and Management, 352, 9-20, <https://doi.org/10.1016/j.foreco.2015.06.014>, 2015.
- 193 Li, J. C., Zhao Y. Y., Gao Y. Q., and Liu G. L.: A preliminary analysis of changes in ecological
194 factors after reforestation in swamp land, Scientia Silvae Sinicae, 21(1), 80-85, available at:
195 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE198501009.htm>, last access: 8
196 October 2017, 1985.
- 197 Liu, Q. J.: Tree volume tables of China, China Forestry Publishing House, Beijing, China,
198 2017.
- 199 Luo, T. X.: Patterns of net primary productivity for Chinese major forest types and their
200 mathematical models, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-80061-2006113572.htm>, last access: 8 October 2017, 1996.
- 201 Ma, C. G.: Selection of optimum species and seed sources for plantations of larch, Beijing
202 Agricultural university press, Beijing, China, 1992.
- 203 Payn, T., Carnes, J. M., Freer-Smith, P., Kimberley, M., Kollert, W., Liu, S. R., Orazio, C.,
204 Rodriguez, L., Silva, L. N., and Wingfield, M. J.: Changes in planted forests and future
205 global implications, Forest Ecology and Management, 352, 57-67, <https://doi.org/10.1016/j.foreco.2015.06.021>, 2015.
- 206 Shugart, H. H., Leemans, R., and Bonan, G. B.: A system analysis of the global boreal forest,
207 Cambridge University Press, Cambridge, 1-10, 1992.
- 208 Song, X. S. and Li, Y. M.: Study on the effect of improvement of fen land and afforestation of
209 *Larix gmelinii*, Forestry Science & Technology, 6, 5-7, <https://doi.org/10.13456/j.cnki.lykt.1990.06.003>, 1990.
- 210 Wang, H. Q., Wu, J. C., Yang, W. B., Wang, S. T., and Song, C. J.: A research on management
211 and utilization of "old man forest" of *Larix gmelinii*, Forestry Science & Technology, 16(3),
212 12-14, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ199103003.htm>,



- 219 last access: 8 October 2017, 1991.
- 220 Wang, W. J., Zu, Y. G., Wang, H. M., Matsuura, Y., Sasa, K., and Koike, T.: Plant biomass and
221 productivity of *Larix gmelinii* forest ecosystems in Northeast China: intra- and inter-
222 species comparison, Eurasian Journal of Forest Research, 8, 21-41, available at:
223 <http://hdl.handle.net/2115/22188>, last access: 8 October 2017, 2005.
- 224 Wang, X. P., Fang, J. Y., and Zhu, B.: Forest biomass and root-shoot allocation in northeast
225 China, Forest Ecology and Management, 255, 4007-4020, <https://doi.org/10.1016/j.foreco.2008.03.055>, 2008.
- 226 Wang, Y. H., Zhou, G. S., and Jiang, Y. L.: Simulation of NPP and evaluation of ecosystem
227 services of *Larix gmelinii* forest, Chinese Journal of Applied Ecology, 12(5), 648-652,
228 <https://doi.org/10.13287/j.1001-9332.2001.0157>, 2001a.
- 229 Wang, Y. H., Zhou, G. S., Jiang, Y. L., and Yang, Z. Y.: Estimating biomass and NPP of Larix
230 forests using forest inventory data(FID), Acta Phytocologica Sinica, 25(4), 420-425,
231 available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-ZWSB200104006.htm>, last
232 access: 8 October 2017, 2001b.
- 233 Wang, Z.: Larch Forest in China, China Forestry Publishing House, Beijing, China, 1992.
- 234 Wang, Z., Xu, Z. B., Tan, Z. X., and Dai, H. C.: Ecological characteristics of *Larix gmelinii*
235 grown in igneous rock in Aershan forest region, Scientia Silvae Sinicae, 4, 300-303,
236 available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE197904010.htm>, last
237 access: 8 October 2017, 1979.
- 238 Xu, H.: Forests in Daxinganling Mountains China, Science Press, Beijing, China, 22-53,
239 1998.
- 240 Yang, C. P.: Study on the provenance of Dahurian larch, Science Press, Beijing, China, 2009.
- 241 Yang, Y., Zhang, K., Cai, T. J., and Gu, J. F.: Effect of afforestation species on soil nutrient in
242 dumping site of open coal mine in Daxing'an Mountain, Journal of Northeast Forestry
243 University, 41(9), 54-58, <https://doi.org/10.13759/j.cnki.dlxz.2013.09.028>, 2013.
- 244 Zhang, H. G., Zhang, C. L., Lan, S. B., Kang, Y. K., and Pan, B. L.: The analysis and
245 selection for families of hybrid larch, Journal of Nanjing Forestry University: Natural
246 Science Edition, 29(3), 69-72, <https://doi.org/10.3969/j.issn.1000-2006.2005.03.017>, 2005.
- 247 Zhou, G. S., Wang, Y. H., Jiang, Y. L., and Yang, Z. Y.: Estimating biomass and net primary
248 production from forest inventory data: a case study of China's Larix forests, Forest
249 Ecology and Management, 169(1-2), 149-157, [https://doi.org/10.1016/s0378-1127\(02\)00305-5](https://doi.org/10.1016/s0378-1127(02)00305-5), 2002.
- 250 Zhou, X. F.: Long-Term Research on China's Forest Ecosystems, Northeast Forestry
251 University Press, Harbin, China, 1994.
- 252 **References cited in the dataset**
- 253 An, J.: Tree species and management effect on soil carbon sequestration and fertilities of
254 plantation forests. Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1012445928.htm>, last access: 8 October 2017, 2011.
- 255 Bi, Y. H.: Study on stand growth models of larch gmelinii with different thinning intensity.
256 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2001005669.htm>, last
257 access: 8 October 2017, 2001.
- 258 Cao, G. Z.: Analysis of hydrological characteristics and change trend in western region of
259 Songhua River Basin. Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-10183-](http://cdmd.cnki.com.cn/Article/CDMD-10183-260)



- 263 1011098650.htm, last access: 8 October 2017, 2011
- 264 Chen, J. D., Pan, B. L., Wu, M. C., Han, C. W., and Yan, W. F.: A summary report on the
265 introduction of *Larix principis-rupprechtii*, Forestry Science & Technology, 4, 2-5,
266 available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ198304000.htm>, last
267 access: 8 October 2017, 1983.
- 268 Chen, L. X. and Xiao, Y.: Evolution and evaluation of soil fertility in forest land in *Larix*
269 *gmelinii* plantations at different development stages in Daxinganling forest region, Science
270 of Soil and Water Conservation, 4(5), 50-55, <https://doi.org/10.16843/j.sswc.2006.05.012>,
271 2006.
- 272 Chen, Q., Zhao, Y. S., Xin, Y., Han, X. C., Xia, X. Y., and Han, C. H.: Plant species diversity
273 and correlation with soil factors after vegetation restoration in burned area of Great
274 Xing'an Mountains, Bulletin of Soil and Water Conservation, 32(6), 61-65, 70,
275 <https://doi.org/10.13961/j.cnki.stbctb.2012.06.040>, 2012a.
- 276 Chen, Q., Zhao, Y. S., Xin, Y., and Han, X. C.: Litter and soil water-holding capacity of
277 burned area in northern slope of the Great Khingan after vegetation restoration, Chinese
278 Journal of Soil Science, 44(2), 308-313, <https://doi.org/10.19336/j.cnki.trtb.2013.02.008>,
279 2013.
- 280 Chen, R. M.: A preliminary report on provenance trials of *Larix gmelinii*, Forest Investigation
281 Design, 1, 45-49, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ198901014.htm>, last access: 8 October 2017, 1989.
- 283 Chen, X. W., Wang, C. H., Ding, B. Y., and Zhang, S. Y.: Dynamic thinning system for larch
284 plantation, Journal of Northeast Forestry University, 5(2), 17-21,
285 <https://doi.org/10.1007/BF02843226>, 1994.
- 286 Chen, Y. L., Han, S. J., Zhou, Y. M., Zou, C. J., and Zhang, J. H.: Characteristics of available
287 P in the rhizosphere soil in pure *Juglans mandshurica* and *Larix gmelinii* and their mixed
288 plantation, Chinese Journal of Applied Ecology, 13(7), 790-794, <https://doi.org/10.13287/j.1001-9332.2002.0188>, 2012b.
- 290 Cheng, F. Y.: Spatial variations in non-structural carbohydrates of *Larix gmelinii*, *Juglans*
291 *mandshurica* and *Ulmus japonica*. Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1015665453.htm>, last access: 8 October 2017, 2015.
- 293 Cheng, Y. X. and Li, Z. X.: Preliminary study of biomass of three forest types of *Larix*
294 *gmelinii*, Inner Mongolia Forestry Investigation and Design, 4, 29-39, available at:
295 <http://www.cqvip.com/QK/90816X/198904/206948.html>, last access: 8 October 2017,
296 1989.
- 297 Chu, X.: Spatial and temporal distribution of fine root biomass of dominant tree species and
298 changes after fire on Great Xing'an Mountains. Thesis, available at:
299 <http://cdmd.cnki.com.cn/Article/CDMD-10225-1011147000.htm>, last access: 8 October
300 2017, 2011.
- 301 Chu, X., Di, X. Y., Zhang, J. L., Cai, H. Y., and Yan, B. Z.: Distribution and seasonal
302 dynamics of fine root biomass of two types of forests in Great Xing'an Mountains, Journal
303 of Northeast Forestry University, 39(5), 36-39, <https://doi.org/10.13759/j.cnki.dlxz.2011.05.033>, 2011.
- 305 Cui, G. F., Cai, T. J., and Yang, W. H.: Soil acidity of *Larix gmelinii* plantation, Journal of
306 Beijing Forestry University, 22(3), 33-36, <https://doi.org/10.13332/j.1000-1522.2000.03>.



- 307 008, 2000.
- 308 Di, X. Y., Geng, Y. Y., Sun, L., and Hu, H. Q.: Soil microbial biomass dynamics and influence
309 factors in *Larix gmelinii* forest in Daxing'an Mountains, Journal of Northeast Forestry
310 University, 40(5), 1-4, <https://doi.org/10.13759/j.cnki.dlx.2012.05.009>, 2012.
- 311 Ding, B. Y.: The method and intensity of intermediate cuttings for larch plantation, Journal of
312 Northeast Forestry University, 16(2), 46-57, <https://doi.org/10.13759/j.cnki.dlx.1988.02.007>, 1988.
- 313 Ding, B. Y., Ju, Y. G., Zhang, S. S., Zhang, S. Y., and Yu, W. J.: Study on the structure of the
314 larch wood community, Journal of North-Eastern Forestry Institute, 4, 11-23,
315 <https://doi.org/10.13759/j.cnki.dlx.1982.04.002>, 1982.
- 316 Ding, B. Y., Liu, S. R., and Cai, T. J.: Studies on biological productivity of artificial forests of
317 Dahurian larches, Acta Phytoecologica et Geobotanica Sinica, 14(3), 226-236, available at:
318 <http://www.cnki.com.cn/Article/CJFDTOTAL-ZWSB199003003.htm>, last access: 8
319 October 2017, 1990.
- 320 Dong, X. B.: The impacts of cutting intensity on the growth of larch forest, Journal of
321 Northeast Forestry University, 29(1), 44-47, <https://doi.org/10.13759/j.cnki.dlx.2001.01.011>, 2001.
- 322 Dong, X. B.: Modes and evaluation of ecological harvesting in Larch plantation areas. Thesis,
323 available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009132292.htm>, last access:
324 8 October 2017, 2008.
- 325 Dong, X. B., Yang, X. C., and Yang, G. X.: Impacts of different cutting modes on soil
326 properties in Larch plantations, Journal of Northeast Forestry University, 35(10), 7-10,
327 <https://doi.org/10.13759/j.cnki.dlx.2007.10.012>, 2007.
- 328 Dong, Y.: A survey study on the factors influencing the growth of *Larix gmelinii* and *Pinus*
329 *sylvestris* plantations, Transactions of China Pulp and Paper, Suppl., 134-142,
330 <https://doi.org/10.3321/j.issn:1000-6842.2004.z1.021>, 2004.
- 331 Du, E. Z. and Fang J. Y.: Linking belowground and aboveground phenology in two boreal
332 forests in Northeast China, Oecologia, 176, 883-892, <https://doi.org/10.1007/s00442-014-3055-y>, 2014.
- 333 Du, Y. J., Xu, H. C., and Yu, R. Y.: The effect of understory, litter and small mammals on
334 seedling occurrence of *Larix gmelinii*, Journal of Beijing Forestry University, 15(4), 12-20,
335 <https://doi.org/10.13332/j.1000-1522.1993.04.002>, 1993.
- 336 E, W. F.: Stem respiration spatial-temporal dynamics and characteristics of sapwood/
337 heartwood of *Larix gmelinii*, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009133891.htm>, last access: 8 October 2017, 2008.
- 338 E, W. F., Wang C. K., Yang C. P., Wang, X. C., Zhang, Q. Z., and Zhang, Y. Q.: Effects of
339 provenances on growth characteristics of sapwood and heartwood of *Larix gmelinii* trees,
340 Scientia Silvae Sinicae, 45(6), 109-115, <https://doi.org/10.3321/j.issn:1001-7488.2009.06.019>, 2009.
- 341 Feng, L. and Yang, Y. G.: A study on biomass and production of three types of Dahurian larch
342 virgin forest, Scientia Silvae Sinicae, 21(1), 86-92, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE198501010.htm>, last access: 8 October 2017, 1985.
- 343 Fu, Y.: Study on *Larix gmelinii* plantation in different planting density in the south of
344 Daxing'anling City, Journal of Inner Mongolia Forestry, 5, 24-26, available at:



- 351 <http://www.cnki.com.cn/Article/CJFDTotal-NMLY198405030.htm>, last access: 8 October
352 2017, 1984.
- 353 Gao, M. F., Li, C. H., and He, S. S.: Productivity division of *Larix gmelinii* in Great Xing'an
354 Mountains of Inner Mongolia, Inner Mongolia Forestry Investigation and Design, 28,
355 75-77, 84, <https://doi.org/10.13387/j.cnki.nmld.2005.s1.037>, 2005.
- 356 Gao, X. W.: Research by the stand density effect on Xing'an larch of natural forest, Thesis,
357 available at: <http://cdmd.cnki.com.cn/Article/CDMD-10129-1014375561.htm>, last access:
358 8 October 2017, 2014.
- 359 Gao, X. W., Zhang, Q. L., Bao, Y., Wang, X. H., Li, D. Y., and Shi, X.: Study on the forest
360 biomass and carbon density under different cutting patterns of Dahurian Larch forest,
361 Forest Resources Management, 3, 91-96, <https://doi.org/10.13466/j.cnki.lyzygl.2014.03.020>, 2014.
- 362 Gu, Y. C.: The necessity of tending and thinning of natural young and middle aged forest,
363 Forest Resources Management, 1, 58-62, <https://doi.org/10.13466/j.cnki.lyzygl.1985.01.018>, 1985a.
- 364 Gu, Y. C.: The succession of forest communities in the forest region of the Greater Xingan
365 Mountains, Acta Phytoecologica et Geobotanica Sinica, 9(1), 64-70, available at:
366 <http://www.cnki.com.cn/Article/CJFDTOTAL-ZWSB198501006.htm>, last access: 8
367 October 2017, 1985b
- 368 Guan, J. Q., Cai, T. J., and Man, X. L.: Characteristics of snowpack chemistry in different
369 type plantations of Xiaoxing'an Mountains, northeastern China, Journal of Beijing Forestry
370 University, 35(4), 41-46, <https://doi.org/10.13332/j.1000-1522.2013.04.020>, 2013.
- 371 Guo, R. Y., Fu, Y., Gao, S. C., and Jia, Z. S.: Studies on the reasonable planting density of
372 Poplar and larch in Xing-an League, Inner Mongolia Forestry Science and Technology, 2,
373 20-22, 35, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198402005.htm>,
374 last access: 8 October 2017, 1984.
- 375 Guo, Y. X.: The variation characteristics of heartwood and sapwood of artificial *Larix*
376 *gmelinii*, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1015665595.htm>,
377 last access: 8 October 2017, 2015.
- 378 Guo, Y. X. and Jiang, L. C.: Variation of heartwood and sapwood and growth characteristics
379 of artificial Dahurian Larch, Journal of Northeast Forestry University, 43(6), 93-97,
380 <https://doi.org/10.13759/j.cnki.dlxz.20150522.048>, 2015.
- 381 Han, J. Y., Zhang, H. G., Jia, Q. B., Zhang, Z., and Xu, Y. L.: Genetic variations in growth
382 traits of juvenile stage hybrid larch in southern Xiao Xing'an Mountains and selection of
383 superior families, Journal of Northwest Forestry University, 27(6), 78-82,
384 <https://doi.org/10.3969/j.issn.1001-7461.2012.06.16>, 2012a.
- 385 Han, M. Z.: A research on biomass and production of Larix Dahurica forest communities,
386 Acta Agriculturae Boreali-Sinica, 2(4), 134-138, <https://doi.org/10.3321/j.issn:1000-7091.1987.04.022>, 1987.
- 387 Han, S. L., Zhang, Q. L., AN, H. J., and Yu, B.: Spatial structure characteristics of *Larix*
388 *gmelinii* natural forest, Journal of Arid Land Resources and Environment, 29(5), 87-92,
389 <https://doi.org/10.13448/j.cnki.jalre.2015.157>, 2015a.
- 390 Han, X. C.: Effect on process of burned areas *Larix gmelinii* plantations restoration in
391 Daxinganling, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1015665595.htm>
- 392 393 394



- 395 1013371072.htm, last access: 8 October 2017, 2013.
- 396 Han, X. C., Zhao, Y. S., Xin, Y., and et al.: Soil hydrology characteristics of *Larix gmelinii*
397 plantation in burned forestland in the north of Greater Xing'an Mountains, Journal of Soil
398 and Water Conservation, 26(4), 183-188, <https://doi.org/10.13870/j.cnki.stbcxb.2012.04.053>, 2012b.
- 400 Han, X. C., Zhao, Y. S., and Xin, Y.: Succession process of *Larix gmelinii* forest with
401 artificial restoration after fire in Daxing'anling, Science of Soil and Water Conservation,
402 13(2), 70-76, <https://doi.org/10.16843/j.sswc.2015.02.012>, 2015b.
- 403 He, H. Y.: Analysis on growth of three species of artifical plantation in Zhangguangcai
404 Mountains-subregion of Xiaoxingan Mountains, Journal of Northeast Forestry University,
405 16(1), 32-36, <https://doi.org/10.13759/j.cnki.dlxb.1988.01.005>, 1988.
- 406 Hong, Q. L., Cai, Y. X., Wang, Y. H., Zhao, H. X., Hu, C. X., Li, F. R., and Fan, W. Y.: The
407 growth of natural forests of *Larix gmelinii* in Tahe Forestry Bureau Daxingan Mountain
408 District, Journal of Northeast Forestry University, 22(2), 92-97,
409 <https://doi.org/10.13759/j.cnki.dlxb.1994.02.017>, 1994.
- 410 Hu, H. Q., Wei, S. J., and Sun, L.: Estimation of carbon emissions from forest fires in 2010 in
411 Huzhong of Daxing'anling Mountain, Scientia Silvae Sinicae, 48(10), 109-119,
412 <https://doi.org/10.11707/j.1001-7488.20121017>, 2012.
- 413 Huang, C., Meng, F. L., Shan, C. C., and Hua, S.: Techniques for fast growing and high yield
414 of larch, Journal of Heilongjiang Vocational Institute of Ecological Engineering, 24(1),
415 28-29, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-HSGX201101014.htm>,
416 last access: 8 October 2017, 2011a.
- 417 Huang, D. Q.: Biomass and carbon storage estimation of *Larix gmelinii*, Thesis, available at:
418 <http://d.wanfangdata.com.cn/Thesis/Y2026419>, last access: 8 October 2017, 2011.
- 419 Huang, D. Q., Hu, H. Q., Wang, Z. W., and Sun, L.: Dynamic changes in biomass and
420 productivity of *Larix gmelinii* plantation after fire, Journal of Northeast Forestry University,
421 39(9), 62-64, <https://doi.org/10.13759/j.cnki.dlxb.2011.09.025>, 2011b.
- 422 Huang, S. Z.: The internal cycling of nutrients in fine roots of *Fraxinus Mandshurica* and
423 *Larix gmelinii*, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2006110390.htm>, last access: 8 October 2017, 2006.
- 425 Huo, H.: The effects of climate warming on photosynthesis and growth of *Larix gmelinii*.
426 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2007187763.htm>, last
427 access: 8 October 2017, 2007.
- 428 Jia, S. X., Zhao, Y. L., Sun, Y., Chen, L., and Wang, Z. Q.: Effects of nitrogen fertilization on
429 seasonal dynamics of soil microbial biomass carbon and nitrogen in *Larix gmelinii* and
430 *Fraxinus mandshurica* plantations, Chinese Journal of Applied Ecology, 20(9), 2063-2071,
431 <https://doi.org/10.13287/j.1001-9332.2009.0380>, 2009.
- 432 Jiang, H. Y.: Study on hydrological characteristics of forest ecosystem in Daxing'anling,
433 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009132313.htm>, last
434 access: 8 October 2017, 2008.
- 435 Jiang, H. Y., Zhao, Y. S., Chen, X. W., Li, W. H., Zhu, W. C., Lv, W. B., and Li, X. P.:
436 Research on soil hydrology characteristics of some main forest type in south part of
437 Daxing'anling, Journal of Soil and Water Conservation, 21(3), 149-153, 187,
438 <https://doi.org/10.13870/j.cnki.stbcxb.2007.03.033>, 2007.



- 439 Jiang, H. Y., Zhao, Y. S., Xin, X. J., Ma, W. H., Li, X. P., Sun, C. K., Guo, X. W.:
440 Precipitation distribution laws of different forest stand in Daxing'Anling, Journal of Soil
441 and Water Conservation, 22(6), 197-201, <https://doi.org/10.13870/j.cnki.stcxb.2008.06.044>, 2008.
442
- 443 Jiang, L. F.: Study on mechanism of photosynthesis and respiration of *Larix gmelinii*
444 plantations, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2003110988.htm>, last access: 8 October 2017, 2003.
445
- 446 Jiang, L. F., Shi, F. C., Li, B., Luo, Y. Q., Chen, J. Q., and Chen, J. K.: Separating rhizosphere
447 respiration from total soil respiration in two larch plantations in northeastern China, Tree
448 Physiology, 25, 1187-1195, <https://doi.org/10.1093/treephys/25.9.1187>, 2005.
449
- 450 Jiang, L. F., Shi, F. C., Zu, Y. G., Wang, W. J., and Koike, T.: Study on stem respiration of
451 *Larix gmelinii* of different ages and its relationship to environmental factors, Bulletin of
452 Botanical Research, 23(3), 296-301, <https://doi.org/10.3969/j.issn.1673-5102.2003.03.012>,
453 2003.
454 Jiang, R., Sun, L., and Hu, H. Q.: The seasonal dynamics of soil microbial biomass of
455 a *Larix gmelinii* forest after wildfire, African Journal of Microbiology Research, 6(10),
456 2328-2336, <https://doi.org/10.5897/ajmr11.1264>, 2012.
457 Laboratory of Forest Tree and Breeding, Northeast Forestry University(LFTB).: Study on
458 larch heterosis, Forestry Science & Technology, 1, 1-3, <https://doi.org/10.13456/j.cnki.lykt.1979.01.001>, 1979.
459 Li, B.: The impact of different densities on the *Larix gmelinii* plantation. Thesis, available at:
460 <http://cdmd.cnki.com.cn/Article/CDMD-10129-1013248427.htm>, last access: 8 October
461 2017, 2013a.
462 Li, B., De, Y. J., Wei, Q., Wang, Y., Wang, Y. B., and Bao, Y. Q.: Effects of three densities on
463 the growth of *Larix gmelinii* plantations, Journal of Inner Mongolia Agricultural University,
464 34(2), 65-68, <https://doi.org/10.16853/j.cnki.1009-3575.2013.02.013>, 2013.
465 Li, J., Luo, Y. Q., Shi, J., Zhou, J., Wang, X. L., Ma L. Y., and Chen C.: Use of
466 phyto-attractant in monitoring and controlling gypsy moth, Journal of Beijing Forestry
467 University, 33(4), 85-90, <https://doi.org/10.13332/j.1000-1522.2011.04.010>, 2011.
468 Li, L. N., Zhao, L., Zhang, W. J., and Wang, X. J.: Study on species diversity in different
469 density of *Larix gmelinii* plantations in Zhalantun Region, Practical Forestry Technology, 8,
470 9-11, <https://doi.org/10.13456/j.cnki.lykt.2014.08.002>, 2014.
471 Li, M., Gu, J. C., Zhang, Z. W., and Wang, Z. Q.: Responses of fine root mass, length,
472 production and turnover to soil nitrogen fertilization in *Larix gmelinii* and *Fraxinus
473 mandshurica* forests in Northeastern China, Journal of Forest Research, 15, 194-201,
474 <https://doi.org/10.1007/s10310-009-0176-y>, 2010a.
475 Li, Q., Chen, J., Li, T., and Liu, F. J.: Ecological function of artificial coniferous timber
476 forests in water conservation, Jilin Forestry Science and Technology, 36(3), 14-17, 20,
477 <https://doi.org/10.16115/j.cnki.issn.1005-7129.2007.03.004>, 2007.
478 Li, S. L. and Chen, Y. L.: Decomposition and nutrient return of the leaf litter under the pure
479 and mixed plantations of *Juglans mandshurica* and *Larix gmelinii*, Journal of Nanjing
480 Forestry University: Natural Science Edition, 28(5), 59-62,
481 <https://doi.org/10.3969/j.issn.1000-2006.2004.05.014>, 2004.
482 Li, X. F.: Comprehensive evaluation of the effects of different thinning intensities on *Larix*



- 483 *gmelinii* forest in Great Xing'an Mountains of Inner Mongolia, Inner Mongolia Forestry
484 Investigation and Design, 1, 7-20, available at: <http://www.cqvip.com/QK/90816X/1993011276513.html>, last access: 8 October 2017, 1993.
- 485
- 486 Li, X. F.: Study on genetic diversity of provenances and clones of *Larix gmelinii* (Rupr.) Rupr,
487 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009134088.htm>, last
488 access: 8 October 2017, 2009.
- 489 Li, X. F.: Spatio-temporal dynamics and influencing factors of fine roots in five Chinese
490 temperate forest ecosystems, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1013366884.htm>, last access: 8 October 2017, 2013b.
- 491
- 492 Li, X. Y.: The study on dynamic features of several metabolites of larch(*Larix gmelinii*),
493 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2005136726.htm>, last
494 access: 8 October 2017, 2005.
- 495 Li, Y. W., Chen, X. H., and Hu, M. Q.: Study on growth and mortality of mature and
496 over-mature *Larix gmelinii* forest in Daxinganling forest region, Forest Resources
497 Management, 1, 38-43, <https://doi.org/10.13466/j.cnki.lyzygl.1979.01.007>, 1979.
- 498 Li, Z. F., Zhao, Y. S., Xin, Y., Kong, X. F., and Wu, C. Y.: Soil water storage and infiltration
499 characteristics under 3 artificial forests in the upper reaches of Ashihe river, Science of Soil
500 and Water Conservation, 8(1), 77- 80, <https://doi.org/10.16843/j.sswc.2010.01.014>, 2010b.
- 501 Li, Z. F., Zhao, Y. S., Xin, Y., Kong, X. F., and Wu, C. Y.: Temporal and spatial variation of
502 soil water under different forest types in the upper reaches of Ashihe River, Science of Soil
503 and Water Conservation, 8(4), 86-89, <https://doi.org/10.16843/j.sswc.2010.04.016>, 2010c.
- 504 Lin, L.: Afforestation techniques on southward slopes in Greater Khingan Mts. forest district,
505 Journal of Northeast Forestry University, 3, 48-58, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198103005.htm>, last access: 8 October 2017, 1981.
- 506
- 507 Liu, G. L.: Intermediate cutting experiment of larch (Larix sp.) plantation, Journal of
508 North-Eastern Forestry Institute, 12(1), 33-46, <https://doi.org/10.13759/j.cnki.dlxz.1984.01.004>, 1984.
- 509
- 510 Liu, G. L. and Xu, M. H.: Study on half-sib breeding plantation of different larchs. Forestry
511 Science and Technology, 2: 19-24. available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-JLLK198402003.htm>, last access: 8 October 2017, 1984.
- 512
- 513 Liu, H. L., Cai, T. J., Man, X. L., Chai, R. S., and Lang, Y.: Effects of major forest types of
514 Xiaoxing'an Mountains on the process of snowfall, snow cover and snow melting, Journal
515 of Beijing Forestry University, 34(2), 20-25, <https://doi.org/10.13332/j.1000-1522.2012.02.003>, 2012.
- 516
- 517 Liu, J. L., Wang, C. K., and Zhang, Q. Z.: Spatial variations in stem heartwood and sapwood
518 for *Larix gmelinii* trees with various differentiation classes, Scientia Silvae Sinicae, 50(12),
519 114-121, <https://doi.org/10.11707/j.1001-7488.20141216>, 2014.
- 520
- 521 Liu, S. R., Chai, Y. X., Cai, T. J., and Peng, C. H.: Study on biomass and net primary
522 productivity of Dahurian larch plantation, Journal of Northeast Forestry University, 18(2),
523 40-46, <https://doi.org/10.13759/j.cnki.dlxz.1990.02.006>, 1990a.
- 524
- 525 Liu, S. R., Li, X. M., and Niu, L. M.: The degradation of soil fertility in pure larch plantations
526 in the northeastern part of China, Ecological Engineering, 10, 75-86,
527 [https://doi.org/10.1016/s0925-8574\(97\)10024-6](https://doi.org/10.1016/s0925-8574(97)10024-6), 1998.
- 528 Liu, T. Y.: Carbon storage in Manchurian ash and Dahurian larch plantations with different



- 527 stand structures. Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1012445805.htm>, last access: 8 October 2017, 2012.
- 528 Liu, W. and Deng, H.: Study on the effect of different afforestation methods on the survival
529 rate of Larix, Forestry Science and Technology Information, 39(4), 4-5,
530 <https://doi.org/10.3969/j.issn.1009-3303.2007.04.003>, 2007.
- 531 Liu, X. T., Du, X. G., Liang, L., Zhang, P., Zhang, B. C., and Zhang, Y. S.: The selection of
532 geographic provenances of *Larix gmelinii* in the eastern slope of Great Xing'an Mountain,
533 Journal of Northeast Forestry University, 18, 9-16, <https://doi.org/10.13759/j.cnki.dlxb.1990.s2.003>, 1990b.
- 534 Liu, X. T., Zhang, H. J., Wang, S. X., Lin, Y. K., and He, X. Y.: Introduction of *Larix olgensis*
535 in Daxing'anling, Journal of Northeast Forestry University, 17(3), 18-25, available at:
536 <http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198903002.htm>, last access: 8
537 October 2017, 1989.
- 538 Liu, Y.: Study on the niche and the production of artificial Korean pine and Larch population.
539 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2003111030.htm>, last
540 access: 8 October 2017, 2002.
- 541 Liu, Y. C., Wu, M. C., Chu, Y. J., Liu, D. C., and He, Z. J.: Study on the provenance test of
542 *Larix gmelinii*. Forestry Science and Experiment, 35, 1-17, available at: <http://www.cqvip.com/QK/92313X/199035/319899.html>, last access: 8 October 2017, 1990c.
- 543 Lu, X., Hu, H. Q., Sun, L., and Wei, S. J.: Effects of fire disturbance on soil light fraction
544 organic carbon of *Larix gmelinii* forest, Journal of Anhui Agricultural Sciences, 41(29),
545 11709-11713, <https://doi.org/10.13989/j.cnki.0517-6611.2013.29.006>, 2013.
- 546 Luo, Y. J., Wang, X. K., Zhang, X. Q., and Lu, F.: Biomass and its allocation of forest
547 ecosystems in China, China Forestry Publishing House, Beijing, China, 2013.
- 548 Ma, H. B.: Stem sap flow dynamics in relation to environmental factors of *Larix gmelinii*.
549 Thesis, <https://doi.org/10.7666/d.y1473827>, 2009.
- 550 Ma, J. W.: Study on spectral characteristics of pure stands under different site conditions and
551 phenological stages at Maoershan Area, Journal of Beijing Forestry College, 3, 26-40,
552 <https://doi.org/10.13332/j.1000-1522.1983.03.003>, 1983.
- 553 Ma, J. W., Zhang, H. G., Liu, L., Zhang, Z., and Jia, Q. B.: Genetic variation of hybrid larch
554 at juvenile period and its superior family selection, Journal of Northeast Forestry
555 University, 40(9), 1-5, <https://doi.org/10.13759/j.cnki.dlxb.2012.09.019>, 2012.
- 556 Mao, W. X., Tong, D. H., Zhang, C., Zhao, D. D., Ding, Y. Y., and Jin, S.: Spatial
557 heterogeneity of moisture content of land surface dead fuel in larch stand and effects of
558 sampling methods on moisture estimation, Journal of Northeast Forestry University, 40(5),
559 29-33, <https://doi.org/10.13759/j.cnki.dlxb.2012.05.012>, 2012.
- 560 Mao, Y. Q., Zhang, J. L., Wang, F. S., Liu, L., Zhao, W., Zhang, H. P., Yuan, L. D., Wang, F. J.,
561 and Li, B. C.: Study on the provenance test of *Larix olgensis* and *Larix gmelinii*, Protection
562 Forest Science and Technology, 2, 10-15, <https://doi.org/10.13601/j.issn.1005-5215.1993.02.003>, 1993.
- 563 Mao, Y. Q., Zhang, J. L., Wang, F. S., Liu, L., and Wu, C. J.: A study on introduction of Larch.
564 Scientia Silvae Sinicae 34(4): 48-58. available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE804.006.htm>, last access: 8 October 2017, 1998.
- 565 Meng, C., Wang, L. H., Wang, J., You, X. F., Liu, Y. Q., and Xing, J.: Diurnal change in soil
566



- 571 respiration rate in five types of plantations during non-growing season, Journal of
572 Northeast Forestry University, 37(11), 77-79, 91, <https://doi.org/10.13759/j.cnki.dlxb.2009.11.029>, 2009.
- 573 Meng, X. Q., Li, H. Y., Wang, Y. M., Zhang, Y. W., and Yang, W. Y.: Study on the effects of
574 Coleophora dahurica on tree growth, Journal of Liaoning Forestry Science & Technology,
575 5, 19-20, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LNLK200005007.htm>, last access: 8 October 2017, 2000.
- 576 Mu, T. M. and Yan, W.: Preliminary study on the evapotranspiration in Xingan larch forest,
577 Journal of Northeast Forestry University, 2, 21-30, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198202001.htm>, last access: 8 October 2017, 1982.
- 578 Na, D. C.: A study on genetic diversity of different geographic provenances of *Larix gmelinii*(Rupr.) Rupr and its utilization, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2005136986.htm>, last access: 8 October 2017, 2005.
- 579 Na, D. C., Yang, C. P., Jiang, J., Xia, D. A., and Wang, H. R.: Provenance division and
580 superior provenance choice of *Larix gmelinii*, Forestry Science & Technology, 30(4), 1-4,
581 available at: <https://doi.org/10.3969/j.issn.1001-9499.2005.04.001>, 2005.
- 582 Piao, C. B., Zhang, W. X., Lin, J., and Ma, A. B.: The preliminary report on the good natural
583 Dahurian larch stand transformation for seed stand, Forestry Science & Technology, 16(3),
584 17-23, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ199103005.htm>, last access: 8 October 2017, 1991.
- 585 Qi, G., Wang, Q. L., Wang, X. C., Qi, L., Wang, Q. W., Ye, Y. J., and Dai, L. M.: Vegetation
586 carbon storage in *Larix gmelinii* plantations in Great Xing'an Mountains, Chinese Journal
587 of Applied Ecology, 22(2), 273-279, <https://doi.org/10.13287/j.1001-9332.2011.0072>, 2011.
- 588 Qi, G., Wang, Q. L., Wang, X. C., Yu, D. P., Zhou, L., Zhou, W. M., Peng, S. L., and Dai, L.
589 M.: Soil organic carbon storage in different aged *Larix gmelinii* plantations in Great
590 Xing'an Mountains of Northeast China, Chinese Journal of Applied Ecology, 24(1), 10-16,
591 available at: <https://doi.org/10.13287/j.1001-9332.2013.0122>, 2013.
- 592 Qiu, L.: Temporal dynamic change of soil pool in larch plantations and its contribution to the
593 forest carbon budget in NE China, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1011146228.htm>, last access: 8 October 2017, 2011.
- 594 Qiu, L., Zu, Y. G., Wang, W. J., Sun, W., Su, D. X., and Zheng, G. Y.: CO₂ flux characteristics
595 and their influence on the carbon budget of a larch plantation in Maoershan region of
596 Northeast China, Chinese Journal of Applied Ecology, 22(1), 1-8,
597 available at: <https://doi.org/10.13287/j.1001-9332.2011.0031>, 2011.
- 598 Quan, X., Wang, C., Zhang, Q., Wang, X., Luo, Y., and Bond-Lamberty, B.: Dynamics of fine
599 roots in five Chinese temperate forests, Journal of Plant Research, 123, 497-507,
600 available at: <https://doi.org/10.1007/s10265-010-0322-9>, 2010.
- 601 Ren, B. P.: Research on growth process and stand density of *Larix gmelinii* plantation,
602 Journal of Inner Mongolia Radio & TV University, 2, 73-75, 85,
603 available at: <https://doi.org/10.16162/j.issn.1672-3473.2010.02.022>, 2010.
- 604 Ren, L. J., Li, G. T., Sun, X. J., Yu, S. P., Zheng, X. W., and Yue, Y. J.: Research on
605 community structure and species diversity of *Larix gmelinii* community in Inner Mongolia
606 Daxing'anling Mountains, Forest Resources Management, 3, 83-87, 118,
607 available at: <https://doi.org/10.13466/j.cnki.lyzygl.2013.03.023>, 2013.



- 615 Shang, Y. L., Zhang, S. H., Chen, Z. C., and Wang, G. Y.: Fine family selection and genetic
616 gain of *Larix gmelinii*, Journal of Northeast Forestry University, 38(7), 123-125,
617 <https://doi.org/10.13759/j.cnki.dlxz.2010.07.017>, 2010.
- 618 Sheng, H. C., Cai, T. J., Zhu, D. G., and Shi, F. C.: Rainfall redistribution and hydrochemical
619 characteristics in the Larch plantation, Journal of Soil and Water Conservation, 23(2),
620 79-83, 89, <https://doi.org/10.13870/j.cnki.stbcxb.2009.02.050>, 2009.
- 621 Si, H. Z., Yang, S., Li, J., Wang, F. S. and Wen, B. Y.: A study on introduction of *Larix*
622 *kaempferi*, Forest Science and Technology, 11, 5-7, <https://doi.org/10.13456/j.cnki.lykt.1999.11.002>, 1999.
- 623 Song, C. L., Zhao, P. W., Su, R. N., Hai, L., Zhou, M., and Feng, L.: Dynamic study of
624 photosynthetic characteristics of *Larix gmelinii*, Journal of Inner Mongolia Agricultural
625 University, 29(4), 49-54, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-NMGM200804010.htm>, last access: 8 October 2017, 2008a.
- 626 Song, S., Gu, J. C., Quan, X. K., Guo, D. L. and Wang, Z. Q.: Fine-root decomposition of
627 *Fraxinus mandshurica* and *Larix gmelinii* plantations, Journal of Plant Ecology, 32(6),
628 1227-1237, <https://doi.org/10.3773/j.issn.1005-264x.2008.06.003>, 2008b.
- 629 Song, W. Y.: Carbon density and its distribution of *Pinus koraiensis* and *Larix gmelinii*
630 plantations in different regions of Heilongjiang province. Thesis, available at:
631 <http://cdmd.cnki.com.cn/Article/CDMD-10225-1014402349.htm>, last access: 8 October
632 2017, 2013.
- 633 Song, X. S., Bao, F. L., and Zhang, Y. L.: The analysis of the economic benefit on the fast
634 growing and high yield of *Larix gmelini* plantation, Forestry Science & Technology, 17(2),
635 16-17, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ199202004.htm>, last access: 8 October
636 2017, 1992.
- 637 Song, X. S., Guo, S. Y., and Wang, J. X.: Study on fast growing and high yield of *Larix*
638 *gmelini* plantation, Forestry Science & Technology, 21(3), 22-24, available at:
639 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ603.009.htm>, last access: 8 October
640 2017, 1996.
- 641 Sun, C. Z., Shen, G. F., Li, J. Y., and Jia, L. M.: Study on the present condition and the
642 potentialities of the productivity of main tree species plantation of china II. study on the
643 plantation productivities of Eucalyptus, Larix and Pinus sylvestris var. Mongolica, Forest
644 Research, 14(6), 657-667, <https://doi.org/10.13275/j.cnki.lykxyj.2001.06.012>, 2001.
- 645 Sun, H. Z., Chen, M. Y., Cai, C. J., and Zhu, N.: Effects of different type urban forest
646 plantations on soil fertility, Chinese Journal of Applied Ecology, 20(12), 2871-2876,
647 <https://doi.org/10.13287/j.1001-9332.2009.0454>, 2009.
- 648 Sun, H. Z., Lu, X. J., Chen, M. Y., Cai, C. J., and Zhu, N.: Soil heavy metals concentration in
649 different type plantations in Harbin City, Chinese Journal of Applied Ecology, 22(3),
650 614-620, <https://doi.org/10.13287/j.1001-9332.2011.0126>, 2011.
- 651 Sun, X. M., Zhang, S. G., Zhou, D. Y., Wang, X. D., Ding, B. and Liu, S. M.: Phenological
652 variation of Larix species and their intra-species and inter-species hybrid families and early
653 selection, Scientia Silvae Sinicae, 44(1), 77-84, <https://doi.org/10.3321/j.issn:1001-7488.2008.01.013>, 2008.
- 654 Sun, Y.: Effects of soil fauna on root biomass, morphology, production and turnover rate in
655 Larch and Ash plantatioins, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD->



- 659 10225-2010242477.htm, last access: 8 October 2017, 2010.
660 Sun, Y., Wang, L. L., Chen, J., and Duan J. P.: Reconstructing mean maximum temperatures
661 of May–August from tree-ring maximum density in North Da Hinggan Mountains, China,
662 Chinese Science Bulletin, 57(16), 2007-2014, <https://doi.org/10.1007/s11434-012-5055-9>,
663 2012.
664 Sun, Z. B. and Sun, X. W.: Selection of species and seed sources for *Larix* spp. in Chifeng
665 Region, Inner Mongolia, in: Ma, C.G., ed. Selection of optimum species and seed sources
666 for plantations of larch, Beijing Agricultural University Press, Beijing, China, 1992.
667 Sun, Z. B., Wang, X.W., Ding, M.X., Shi, J.S., Liu, Z., and Bao, G.Z.: Study on selection of
668 optimum species and seed sources for plantations of larch, Forest Science and Technology,
669 10, 11-13, <https://doi.org/10.13456/j.cnki.lykt.1991.10.005>, 1991.
670 Tang, J. Z., Li, Q. Y., Wang, H. X., and Wang, Y. G.: The main raw material forest of larch
671 growth indicators, Forest Investigation Design, 3, 61-63, available at:
672 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKC200903029.htm>, last access: 8
673 October 2017, 2009.
674 Tian, F., Bi, X. H., Shen, G. T., Zhou, B. and Li, W. B.: Study on the forecast technique and
675 control index of larch caterpillar of a cycle in two years, Forest Research, 11(4), 388-395,
676 <https://doi.org/10.13275/j.cnki.lykxyj.1998.04.008>, 1998.
677 Wang, B. S.: The undergrowth succession in larch plantations before and after cutting,
678 Journal of Northeast Forestry University, 24(5), 82-86, <https://doi.org/10.13759/j.cnki.dlxz.1996.05.014>, 1996.
679 Wang, C. K.: Biomass allometric equations for 10 co-occurring tree species in Chinese
680 temperate forests, Forest Ecology and Management, 222, 9-16,
681 <https://doi.org/10.1016/j.foreco.2005.10.074>, 2006.
682 Wang, C. K., Gower, S. T., Wang, Y. H., Zhao, H. X., Yan, P., and Bond-Lamberty, B.: The
683 influence of fire on carbon distribution and net primary production of boreal *Larix gmelinii*
684 forests in north-eastern China, Global Change Biology, 7, 719-730,
685 <https://doi.org/10.1046/j.1354-1013.2001.00441.x>, 2001a.
686 Wang, C. K. and Yang, J. Y.: Rhizospheric and heterotrophic components of soil respiration
687 in six Chinese temperate forests, Global Change Biology, 13, 123-131,
688 <https://doi.org/10.1111/j.1365-2486.2006.01291.x>, 2007.
689 Wang, H. X., Zhang, S. H., Jing, C. Q., Dai, J. G., Liu, Y. T. and Zhou, J. Y.: Cultivation
690 techniques for raw material forest of short rotation Larch, Forestry Science & Technology,
691 25(5), 7-8, <https://doi.org/10.3969/j.issn.1001-9499.2000.05.003>, 2000a.
692 Wang, J. Z.: Study on selection of larch species, Jilin Forestry Science and Technology, 1,
693 22-26, available at: <http://www.cqvip.com/QK/96402X/199501/1795612.html>, last access:
694 8 October 2017, 1995.
695 Wang, L.: Regeneration rules of the north forest region in western slope of Inner Mongolia
696 Daxing'anling, Inner Mongolia Forestry Investation and Design, S1, 65-69,
697 <https://doi.org/10.13387/j.cnki.nmld.2000.s1.022>, 2000.
698 Wang, L., Li, C. H., Yang, J. Y., and Guo, Y. L.: Biomass changes of artificial *Larix gmelinii*
699 from 2002 to 2012 in Helong, Jilin Province, Journal of Green Science and Technology, 2,
700 11-12, <https://doi.org/10.3969/j.issn.1674-9944.2013.02.005>, 2013a.
701 Wang, L. H., Xin, Y., Zou, M. L., Zhao, Y. S., Lei, Y. Y., and Chen, L.: Litter water-holding



- 703 characteristics in the process of vegetation restoration in severely burned area of Great
704 Xing'an Mountains, Journal of Soil and Water Conservation, 28(6), 262-267,
705 <https://doi.org/10.13870/j.cnki.stbcxb.2014.06.048>, 2014.
- 706 Wang, L. H., Xin, Y., Zou, M. L., Zhao, Y. S., Liu, S. J., and Ren, Q. S.: Plants diversity and
707 biomass distribution of vegetation restoration in burned area of Great Xing'an Mountains,
708 Journal of Beijing Forestry University, 37(12), 41-47, <https://doi.org/10.13332/j.1000-1522.20150025>, 2015a.
- 709 Wang, Q. B., Zhang, H. J., Zhang, S. L., Wang, C. B., and Xue, Y.: Species composition and
710 structure investigation of typical plant community in Daliangzihe National Forest Park,
711 Protection Forest Science and Technology, 7, 14-17, 38, <https://doi.org/10.13601/j.issn.1005-5215.2015.07.006>, 2015b.
- 712 Wang, Q. J., Liu, Y. T., Dai, J. G., and Luo, Y. F.: Effects of different provenances and ABT1
713 rooting powder on *Larix gmelinii* afforestation, Forestry Science & Technology, 25(4), 4-5,
714 <https://doi.org/10.3969/j.issn.1001-9499.2000.04.002>, 2000b.
- 715 Wang, Q. Y.: Genetic variation of the provenance for *Picea koraiensis*, Thesis, available at:
716 <http://cdmd.cnki.com.cn/Article/CDMD-10225-2003110985.htm>, last access: 8 October
717 2017, 2003.
- 718 Wang, W. Z., Chen, J., Zhang, B. Y., Feng, Y. L. and Wang, Y. T.: Study on the photosynthetic
719 characteristics and the primary productivity of *Larix olgensis*, Journal of Northeast
720 Forestry University, 22(4), 15-21, <https://doi.org/10.13759/j.cnki.dlx.1994.04.003>, 1994.
- 721 Wang, X. F., Li, G. T., Wu, H. B., Lv, L. K., and Yue, Y. J.: Research on structure and species
722 diversity of *Larix gmelinii* community in Orqohan, Journal of Southwest Forestry
723 University, 33(5), 69-73, <https://doi.org/10.3969/j.issn.2095-1914.2013.05.012>, 2013b.
- 724 Wang, X. H., Zhang, Q. L., Guo, Y. D., Xing, T. T., Wang, F., Gao, X. W., and Wu, J.: Study
725 on the coarse root biomass and distribution characteristics about natural *Larix gmelinii*
726 forest. Forest Resources Management, 5, 62-66, <https://doi.org/10.13466/j.cnki.lyzygl.2013.05.020>, 2013c.
- 727 Wang, X. W. and Mao, Z. J.: Temporal and spatial variation in gas exchange in canopy of
728 *Larix gmelinii* plantation, Scientia Silvae Sinicae, 43(11), 43-49,
729 <https://doi.org/10.3321/j.issn:1001-7488.2007.11.008>, 2007.
- 730 Wang, X. W. and Mao, Z. J.: Practicability of 7 light responsive curve models to different
731 plant species, Bulletin of Botanical Research, 29(1), 43-48, available at:
732 <http://www.cnki.com.cn/Article/CJFDTOTAL-MBZW200901012.htm>, last access: 8
733 October 2017, 2009.
- 734 Wang, Y. H., Zhou, G. S., and Jiang, Y. L.: Simulation of NPP and evaluation of ecosystem
735 services of *Larix gmelinii* forest, Chinese Journal of Applied Ecology, 12(5), 648-652,
736 <https://doi.org/10.13287/j.1001-9332.2001.0157>, 2001a.
- 737 Wang, Y. L.: Biomass, carbon density and distribution characteristics in different age *Larix*
738 *gmelinii* plantations, Journal of Anhui Agricultural Sciences, 43(15), 144-145,150,
739 <https://doi.org/10.13989/j.cnki.0517-6611.2015.15.053>, 2015.
- 740 Wang, Z.: Larch Forest in China, China Forestry Publishing House, Beijing, China, 73-150,
741 1992.
- 742 Wu, C. Y., Zhao, Y. S., Xin, Y., Kong, X. F. and Li, Z. F.: Effects of structure of the *Larix*
743 *gmelinii* plantation on soil hydrology characteristic, Science of Soil and Water



- 747 Conservation, 7(5), 58-62, <https://doi.org/10.16843/j.sswc.2009.05.011>, 2009.
- 748 Wu, X. J., Cai, T. J., Li, H., and Sheng, H. C.: Precipitation hydrochemical characteristic in
749 virgin *Pinus koraiensis* forest and artifical *Larix gmelinii* forest in Liangshui National
750 Nature Reserve, Science of Soil and Water Conservation, 6(6), 37-42,
751 <https://doi.org/10.16843/j.sswc.2008.06.007>, 2008.
- 752 Wu, X. J., Jia, E. L., and Li, J. C.: Preliminary study on the regeneration before clear cutting
753 in *Larix gmelinii* plantation, Forest Science and Technology, 8, 24-26,
754 <https://doi.org/10.13456/j.cnki.lykt.1983.08.010>, 1983.
- 755 Wu, X. L.: Interannual variability of temperature, precipitation and sunshine hours in
756 Wuerqihan Forest Region, Heilongjiang Agricultural Sciences, 10, 113-116,
757 <https://doi.org/10.11942/j.issn1002-2767.2015.10.0113>, 2015.
- 758 Wu, Z. C.: Introduction and afforestation of *Larix olgensis*, Journal of Inner Mongolia
759 Forestry, 12, 20-21, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLY198612016.htm>, last access: 8 October 2017, 1986.
- 760 Xia, C. C., Liu, Z. L., Wang, Q. C., and Li, K. L.: Comparison of the growth and biomass
761 production of monoculture and mixed stands of *Larix gmelinii* and *Betula platyphylla*,
762 Journal of Northeast Forestry University, 40(10), 1-3,
763 <https://doi.org/10.13759/j.cnki.dlxz.2012.10.021>, 2012.
- 764 Xiao, S. L. and Yang, J. L.: Individual tree aboveground biomass of *Larix gmelinii* natural
765 forest in the Northern Greater Khingan Mountains, Scientia Silvae Sinicae, 50(8), 22-29,
766 <https://doi.org/10.11707/j.1001-7488.20140804>, 2014.
- 767 Xiao, Y., Man, X. L., and Fan, J. F.: Soil fertility and water conservation function of main
768 forest types in Gongbiela River Watershed, Journal of Northeast Forestry University, 34(1),
769 28-30, <https://doi.org/10.13759/j.cnki.dlxz.2006.01.010>, 2006.
- 770 Xin, Y., Zhao, Y. S., and Pan, B. Y.: Influence of *Larix gmelinii* plantations at Eastern
771 Mountainous Region in Heilongjiang Province on water quality, Science of Soil and Water
772 Conservation, 4(2), 29-33, <https://doi.org/10.16843/j.sswc.2006.02.006>, 2006.
- 773 Xin, Y., Zou, M. L., and Zhao, Y.S.: Soil water storage capacity and infiltration properties of
774 *Larix gmelinii* plantation in different growth age after burned in Greater Khingan
775 Mountains, Chinese Journal of Soil Science, 45(4), 823-829,
776 <https://doi.org/10.19336/j.cnki.trtb.2014.04.009>, 2014.
- 777 Xu, H. C. and Du, Y. J.: Amount of falling seed and seedling occurrence dynamics in
778 primitive *Larix gmelinii* forest, Scientia Silvae Sinicae, 29(4), 298-306, available at:
779 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE199304001.htm>, last access: 8
780 October 2017, 1993.
- 781 Xu, H. C., Yu, R. Y., Guo, Z. L., Zhou, C. H., and Ma, D. G.: Study on germination conditions
782 of *Larix gmelinii* seedlings in Great Xing'an Mountains, Journal of Beijing Forestry
783 University, 12, 15-25, <https://doi.org/10.13332/j.1000-1522.1990.s4.004>, 1990.
- 784 Xu, H. L., Man, X. L., and Sheng, H. C.: Water conservation function of different types of
785 *Larix gmelinii* forests in Da Hinggan Mountains, Research of Soil and Water Conservation,
786 18(4), 92-96, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-STBY201104020.htm>, last access: 8 October 2017, 2011a.
- 787 Xu, H. L., Man, X. L., and Sheng, H. C.: Comprehensive evaluation on water conservation
788 function of the different types of larch natural forests in Da Hinggan Mountains, Chinese
789
790



- 791 Agricultural Science Bulletin, 29(19), 74-78, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-ZNTB201319017.htm>, last access: 8 October 2017, 2013.
- 792 Xu, J. L.: Study on Larix biomass model and biomass estimation in Dongzhelenghe, Thesis,
793 available at: <http://cdmd.cnki.com.cn/Article/CDMD-10022-2009161609.htm>, last access:
794 8 October 2017, 2009.
- 795 Xu, S. Z. and Han, M. Z.: Regeneration rule and management decision of *Larix gmelinii* in
796 Genhe regions, in Zhou, X. F. ed., Long-Term Research on China's Forest Ecosystems,
797 709-716, Northeast Forestry University Press, Harbin, China, 1994.
- 798 Xu, X. T., Wan, D. Y., Wei, D. F., and Zhou, J. Y.: Growth characteristics and techniques for
799 fast growing and high yield of larch, Inner Mongolia Forestry Investigation and Design,
800 34(6), 54-55, <https://doi.org/10.13387/j.cnki.nmld.2011.06.057>, 2011b.
- 801 Xu, Y. B., Zhou, Y. Q., Wu, R. X., Liu, Z. S., and Yu, C. H.: Afforestation techniques of *Larix*
802 *gmelinii* container seedlings, Forestry Science & Technology, 21(3), 5-7, available at:
803 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ603.001.htm>, last access: 8 October
804 2017, 1996.
- 805 Xu, Y. Q.: Study on the geographic variations and selection for the optimal provenances of
806 *Larix gmelinii*, Forest Inventory and Planning, 3, 15-23, available at:
807 <http://www.cnki.com.cn/Article/CJFDTOTAL-LDGH199303004.htm>, last access: 8
808 October 2017, 1993.
- 809 Xue, Z. H.: Investigation of larch plantation in Daxing'anling Mountains, Inner Mongolia
810 Forestry Science and Technology, 2, 18-21, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198702003.htm>, last access: 8 October 2017, 1987a.
- 811 Xue, Z. H.: Study on tending and thinning of natural young and middle-age *Larix gmelinii*
812 forest, Inner Mongolia Forestry Science and Technology, 4, 1-5, available at:
813 <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198704000.htm>, last access: 8
814 October 2017, 1987b.
- 815 Xue, Z. H.: Tending and Thinning of *Larix gmelinii* plantation, Inner Mongolia Forestry
816 Science and Technology, 3, 20-23, 30, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK199003004.htm>, last access: 8 October 2017, 1990.
- 817 Xue, Z. H.: Cultivation and management techniques of fast-growing and high-yielding *Larix*
818 *gmelinii* Forests, Inner Mongolia Forestry Science & Technology, 1, 11-18, available at:
819 <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK199301004.htm>, last access: 8
820 October 2017, 1993.
- 821 Yang, B. L., Jia, Q. B., Zhang, H. G., and Zhang, L.: Research on hybrid larch superior
822 families selection, Forestry Science & Technology, 39(2), 1-4,
823 <https://doi.org/10.3969/j.issn.1001-9499.2014.02.001>, 2014.
- 824 Yang, C. P.: Study on the provenance of Dahurian larch, Science Press, Beijing, China, 2009.
- 825 Yang, C. P., Jiang, J., Tang, S. S., Li, J. Y. and Wang, H. R.: The provenance test of 21-year
826 old *Larix gmelinii* at Maoershan Area, Journal of Northeast Forestry University, 30(6), 1-5,
827 <https://doi.org/10.13759/j.cnki.dlxb.2002.06.001>, 2002.
- 828 Yang, C. P., Yang, S. W., Xia, D. A., Liu, G. F., Lv, Q. Y. and Zhang, P. G.: Study on the
829 provenance test of Dahurian Larch (II) Selection of the best provenance, Journal of
830 Northeast Forestry University, 19, 68-76, <https://doi.org/10.13759/j.cnki.dlxb.1991.s2.011>,
831 1991.
- 832
- 833
- 834



- 835 Yang, H. T. and Guo, Y. X.: A primary report on the provenance test of *Larix gmelinii* and
836 *Larix olgensis* in Huanan forest region, Journal of Northeast Forestry University, 18, 65-71,
837 <https://doi.org/10.13759/j.cnki.dlx.1990.s2.010>, 1990.
- 838 Yang, K., Zhu, J. J., Gu, J. C., Yu, L. Z., and Wang, Z. Q.: Changes in soil phosphorus
839 fractions after 9 years of continuous nitrogen addition in a *Larix gmelinii* plantation,
840 Annals of Forest Science, 72, 435-442, <https://doi.org/10.1007/s13595-014-0444-7>, 2015.
- 841 Yang, L. J., Zhang, C. J., Li, C. H., Jiang, X. L., and Pan, B. L.: Early selection of superior
842 clone of *Larix gmelinii*, Protection Forest Science and Technology, 4, 35-37,
843 <https://doi.org/10.13601/j.issn.1005-5215.2006.04.013>, 2006.
- 844 Yang, S. W. and Shang, L. T.: Research on selection of larch species, Journal of
845 North-Eastern Forestry Institute, 11(4), 50-56, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198304006.htm>, last access: 8 October 2017, 1983.
- 846 Yang, S. W., Wang, Q. Y., Liu, G. F., Yang, C. P., Xia, D. A., and Bao, L. F.: Analysis on the
847 effects of genetic improvement and the breeding program of larch, Journal of Northeast
848 Forestry University, 20(1), 1-8, <https://doi.org/10.13759/j.cnki.dlx.1992.01.001>, 1992.
- 849 Yang, S. W., Yang, C. P., Wang, Q. Y., Liu, G. F., Xia, D. A., and Zhang, P. G.: The
850 provenance selection of the important tree species in the northeast part of China and Inner
851 Mongolia Autonomous Region, Journal of Northeast Forestry University, 21(4), 1-7,
852 <https://doi.org/10.13759/j.cnki.dlx.1993.04.001>, 1993.
- 853 Yang, S. W., Yang, C. P., Zhang, S. Y., Peng, H. M., Li, J. T., Xia, D. A., Liu, G. F., and Wang,
854 H. R.: A study on the provenance test of *Larix gmelinii* outside of distribution area, Journal
855 of Northeast Forestry University, 18, 1-8, <https://doi.org/10.13759/j.cnki.dlx.1990.s2.002>,
856 1990.
- 857 Yang, Y. B. and Liu, Y. Y.: Discussion about planting fast-growing and high-yield plantation
858 in the south of Daxing'anling City, Forestry Science and Technology Information, 37(2),
859 1-2, <https://doi.org/10.3969/j.issn.1009-3303.2005.02.001>, 2005.
- 860 Yao, C. Y., Wang, Q. J., and Gao, G. Y.: Study on growth status of introduced *Larix olgensis*
861 in Xingan League of Inner Mongolia, Inner Mongolia Forestry Science & Technology, 1,
862 17-22, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK501.004.htm>,
863 last access: 8 October 2017, 1995.
- 864 Yao, G. Q., Chi, G. Q., Dong, Z. Q., Hu, W. L., and Jiang, J. B.: A study on the growth and
865 economic effectiveness of four kinds of planted Larch Forest, Journal of Shenyang
866 Agricultural University, 20, 217-222, available at: <http://www.cnki.com.cn/Article/QK/93761X/198903/129531.html>, last access: 8 October 2017, 1989.
- 867 Yao, Y. J., Chen, X. W., and Li, C. S.: Studies on biomass and productivity for Dahui'an larch
868 plantations, Journal of Northeast Forestry University, 5(2), 13-16,
869 <https://doi.org/10.1007/BF02843225>, 1994.
- 870 Yu, B. and Wang, L. M.: Analysis on classified stem growing characteristics of *Larix gmelinii*
871 natural forest, Forest Research, 20(4), 452-457,
872 <https://doi.org/10.13275/j.cnki.lykxyj.2007.04.014>, 2007.
- 873 Yu, B., Wu, J. S. G. L., Wang, B. T., and Wang, L. M.: Analysis on classified stem
874 transformation characteristic of *Larix gmelinii* natural forest at Daxingan Mountains, Acta
875 Ecologica Sinica, 28(11), 5750-5757, <https://doi.org/10.3321/j.issn:1000-0933.2008.11.062>,
876 2008.
- 877



- 879 Yu, B., Wu, J. S. G. L., Wang, B. T., Wang, L. M., and Da, B. X. L. T.: Study on the
880 distribution pattern characteristics of different stand structures in *Larix gmelinii* natural
881 forest, Forest Research, 23(1), 83-88, <https://doi.org/10.13275/j.cnki.lykxyj.2010.01.013>,
882 2010.
- 883 Yu, B., Zang, Q. L., Wang, L. M., and Wu, J. S. G. L.: Characteristics of biomass and
884 productivity in *Larix gmelinii* natural forests with different stand structures, Journal of
885 Zhejiang A & F University, 28(1), 52-58, <https://doi.org/10.3969/j.issn.2095-0756>.
886 2011.01.009, 2011.
- 887 Yu, B. J. and Yang, C. P.: Study on provenance test of 5 year old *Larix olgensis*, Journal of
888 Northeast Forestry University, 16(3), 27-33, <https://doi.org/10.13759/j.cnki.dlxz.1988.03.005>, 1988.
- 889 Yu, S. L. and Yuan, X. G.: Studies on selection of fine families of Hybrid Larch based on
890 growth traits in seedling stage, Inner Mongolia Forestry Investigation and Design, 32(2),
891 114-117, <https://doi.org/10.13387/j.cnki.nmld.2009.02.025>, 2009.
- 892 Yuan, G. X., Li, H. X., Chen, G. F., Lou, Y. M., and Mou, Z. J.: Analysis on the introduction
893 of *Pinus banksiana* Lamb, Forestry Science & Technology, 26(5), 9-10,
894 <https://doi.org/10.3969/j.issn.1001-9499.2001.05.004>, 2001.
- 895 Yue, Y. J., Gao, R. H., Li, G. T., Jin, J., and Wang, X. F.: Research on community structure
896 and species diversity of *Larix gmelinii* community in Jinhe Forestry Bureau, Journal of
897 Inner Mongolia Agricultural University, 35(2), 43-47,
898 <https://doi.org/10.16853/j.cnki.1009-3575.2014.02.010>, 2014.
- 899 Yue, Y. J., Qu, L. C., Song, L. J., Liu, M. D., Zeng, Y. E., Li, G. T., and Wang, F. S.:
900 Distribution patterns and diversity of species of *Larix gmelinii* in Kuduer, Journal of Inner
901 Mongolia Agricultural University, 34(1), 42-47, <https://doi.org/10.16853/j.cnki.1009-3575.2013.01.009>, 2013.
- 902 Zhai, M. P., Yi, W. L., Jia, L. M., and Ma, L. Y.: Study on the characteristics of *Betula*
903 *platyphylla-Larix gmelinii* Mixed Forest in Xilinji Forest Region, Journal of Beijing
904 Forestry University, 12, 86-91, <https://doi.org/10.13332/j.1000-1522.1990.s3.010>, 1990a.
- 905 Zhai, M. P., Yi, W. L., Jia, L. M., Ma, L. Y., Sun, Y. X., Liu, Z. Y., and Wang, Y. G.:
906 Investigation of *Betula platyphylla-Larix gmelinii* Mixed Forest in Xilinji Forest Region,
907 Journal of Beijing Forestry University, 12, 78-85,
908 <https://doi.org/10.13332/j.1000-1522.1990.s3.009>, 1990b.
- 909 Zhan, J. and Yang, C. P.: Study on the provenance test of Dahurian larch (*Larix gmelinii*),
910 Journal of Northeast Forestry University, 17(6), 22-30, available at:
911 <http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198906003.htm>, last access: 8
912 October 2017, 1989.
- 913 Zhang, G. L., Qu, B. C., Zhang, Y. H., and Wang, H.: Growth analysis and model of *Larix*
914 *gmelinii* plantation, Protection Forest Science and Technology, 4, 41-43,
915 <https://doi.org/10.13601/j.issn.1005-5215.2012.04.056>, 2012.
- 916 Zhang, H. D. and Zhou, M.: Nutrient characteristics of *Larix gmelinii* forest ecosystem in
917 cold temperate zone, Journal Liaoning Forestry Science & Technology, 3, 5-8, 22,
918 <https://doi.org/10.3969/j.issn.1001-1714.2014.03.002>, 2014.
- 919 Zhang, J.: Study on community structure, biomass and carbon storage of *Larix gmelini*
920 plantation, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10022>



- 923 -2008085203.htm, last access: 8 October 2017, 2008a.
- 924 Zhang, J., Sun, Y. J., and Xu, J. L.: Research on growing process of *Larix gmelinii* plantation
925 in Northeast of China, Journal of Northwest Forestry University, 23(6), 179-181, available
926 at: <http://www.cnki.com.cn/Article/CJFDTOTAL-XBLX200806045.htm>, last access: 8
927 October 2017, 2008.
- 928 Zhang, J. L., Mao, Y. Q., Wang, F. S., Liu, L., Wu, C. J., Zhang, H. P., Yuan, L. D., Zhao, W.,
929 and Wang, F. J.: The progeny testing and fine family selection of *Larix gmelinii* and *Larix*
930 *olgensis*, Protection Forest Science and Technology, 3, 22-26,
931 <https://doi.org/10.13601/j.issn.1005-5215.1994.03.003>, 1994.
- 932 Zhang, L.: Study on stability of families and selection of superior families of hybrid larch.
933 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1011146424.htm>,
934 last access: 8 October 2017, 2011.
- 935 Zhang, L. Y.: A study on individual tree health assessment of nature *Larix gmelinii* in
936 Daxing'anling, Forestry Science and Technology Information, 45(1), 41-43.
937 <https://doi.org/10.3969/j.issn.1009-3303.2013.01.018>, 2013.
- 938 Zhang, Q. B. and Fang, L.: A study of problems of planting fast-growing and high-yield
939 plantation in secondary forest site on the south slope in Great Xing'an Mountain, Journal
940 of Inner Mongolia Forestry College, 20(3), 37-42, available at: <http://www.cnki.com.cn/>
941 Article/CJFDTOTAL-NMGL803.007.htm, last access: 8 October 2017, 1998.
- 942 Zhang, S. L., Zhang, H. J., Zhang, W., Guo, Y., Li, X. Z., Zhang, Q., Lu, Y., and Yao, S. P.:
943 Comparison of water-holding characteristics and biomass of litter under different forest
944 types in Xiaoxingan Mountains, Bulletin of Soil and Water Conservation, 35(4), 85-90,
945 <https://doi.org/10.13961/j.cnki.stbctb.2015.04.017>, 2015a.
- 946 Zhang, S. L., Zhang, H. J., Zhang, W., Wang, C. B., Deng, J. H., Song, Y. B., and Zhang, Q.:
947 Soil hydrological properties of main vegetation in Down South of Xiao Xing'an Mountains,
948 Research of Soil and Water Conservation, 22(1), 140-145,
949 <https://doi.org/10.13869/j.cnki.rswc.2015.01.027>, 2015b.
- 950 Zhang, T. W., Yuan, Y. J., Wei, W. S., and Yu, S. L.: The tree-ring chronologies from
951 Moerdaoga in the northeast of Inner Mongolia, Journal of Arid Land Resources and
952 Environment, 23(12), 177-182, <https://doi.org/10.13448/j.cnki.jalre.2009.12.008>, 2009.
- 953 Zhang, X. J.: Effect of close-to-nature transformation of Larch Monoculture stands on
954 regeneration and plant diversity. Thesis, available at: <http://cdmd.cnki.com.cn/Article/>
955 CDMD-10225-2010242257.htm, last access: 8 October 2017, 2010.
- 956 Zhang, X. J., Wang, Q. C., Wang, S. L., and Sun, Q.: Effect of the close-to-nature
957 transformation of *Larix gmelinii* pure stands on plant diversity of understory vegetation in
958 Xiaoxing'an Mountains of China, Scientia Silvae Sinicae, 47(1), 6-14,
959 <https://doi.org/10.11707/j.1001-7488.20110102>, 2011.
- 960 Zhang, Y. Q.: Transpiration characteristic of Daurian Larch trees transplanted from a
961 latitudinal gradient. Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009133900.htm>, last access: 8 October 2017, 2008b.
- 963 Zhao, H. X., Wang, Y. H., Chai, Y. X., Hu, C. X., Li, F. R., and Fan, W. Y.: On forest growth
964 of northern slope of Daxing'an mountains (Tahe Forest Bureau), Journal of Northeast
965 Forestry University, 24(6), 1-8, <https://doi.org/10.13759/j.cnki.dlxb.1996.06.001>, 1996.
- 966 Zhao, P. W.: Studies on litterfall dynamics and nutrient release regularity of *Larix gmelinii* in



- 967 Great Xingan Mountains. Thesis, <https://doi.org/10.7666/d.y1473822>, 2009.
- 968 Zhao, S. J., Yang, J. M., and Bi, X. M.: Study on the fast growing and high yield of larch
969 species and the relationships with their introduction districts, Journal of Hebei Forestry
970 Science and Technology, 2, 2-7, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-HBLK199202001.htm>, last access: 8 October 2017, 1992.
- 971 Zhao, X. M., Xu, S. T., Ren, J. Z., Wang, M. Z., Yang, S. M., and Jiang, S. W.: A study on
972 provenance test of *Pinus sylvestris L.var mongolica* litvin on Korqin sandy land, Journal of
973 Inner Mongolia Forestry College, 21(3), 44-49, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-NMGL199903008.htm>, last access: 8 October 2017, 1999.
- 974 Zhou, B. D., Hao, W. K., Zou, Q., Zhao, K. M., Liu, Q. L., and Chen, Y. D.: Study on
975 artificial regeneration in a clear-cut forest of Xiaoxingan Mountains in Liangshui region,
976 Journal of North-Eastern Forestry College, 4, 19-35, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY196500001.htm>, last access: 8 October 2017, 1965.
- 977 Zhou, B. Q.: Advices on secondary forest culture on the south slope in Great Xing'an
978 Mountain, Inner Mongolia Forestry Investigation and Design, 18-20,
979 <https://doi.org/10.13387/j.cnki.nmld.1998.s1.008>, 1998.
- 980 Zhou, H. X., Zhang, Y. D., Sun, H. L., and Wu, S. Y.: Soil respiration in temperate secondary
981 forest and *Larix gmelinii* plantation in Northeast China, Chinese Journal of Applied
982 Ecology, 18(12), 2668-2674, <https://doi.org/10.13287/j.1001-9332.2007.0477>, 2007.
- 983 Zhou, J. Q., Liu, X. D., and Guo, H. W.: Surface fuel loading and relevant influencing factors
984 of main forest types in southern Daxing'anling, Journal of Northwest A&F
985 University(Natural Science Edition), 42(6), 131-137, <https://doi.org/10.13207/j.cnki.jnwafu.2014.06.010>, 2014.
- 986 Zhou, H. Y.: Main characters variation among provenances of *Fraxinus mandshurica* Rup.
987 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009134220.htm>, last
988 access: 8 October 2017, 2008.
- 989 Zhu, S. Y., Zhou, B., Mao, Z. J., Wang, X. W., and Sun, Y. F.: Space time dynamics of fine
990 root biomass of six forests in Maoershan forest region, *Scientia Silvae Sinicae*, 42(6),
991 13-19, <https://doi.org/10.3321/j.issn:1001-7488.2006.06.003>, 2006.
- 992 Zhu, Y.: Forest health assessment of natural *Larix gmelinii* in Great Xing'an Mountains of
993 China, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1013371221.htm>, last access: 8 October 2017, 2013.
- 994 Zhu, Y., Liu, Z. G., and Jin, G. Z.: Health assessment of individual trees in natural *Larix*
995 *gmelinii* forest in Great Xing'an Mountains of China, Chinese Journal of Applied Ecology,
996 24(5), 1320-1328, <https://doi.org/10.13287/j.1001-9332.2013.0294>, 2013.
- 997 Zou, L., Tang, Q. M., and Wang, Y.: Ecological distribution of soil microorganism in pure and
998 mixed forests of *Pinus sylvestris var. mongolica* and *Larix gmelini*, Journal of Northeast
999 Forestry University, 38(11), 63-64, 79, <https://doi.org/10.13759/j.cnki.dlxz.2010.11.029>,
1000 2010.
- 1001 Zou, M. L., Xin, Y., and Zhao, Y. S.: The influence of vegetation restoration on carbon
1002 storage of burned Area in Greater Xing'an Mountains, Journal of Soil and Water
1003 Conservation, 27(5), 133-137, 143, <https://doi.org/10.13870/j.cnki.stbcxb.2013.05.046>,
1004 2013.
- 1005
- 1006
- 1007
- 1008
- 1009