



1 **Growth characteristics of natural and planted Dahurian larch in northeast China**

2 Bingrui Jia<sup>1\*</sup> and Guangsheng Zhou<sup>2\*</sup>

3 <sup>1</sup>*State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese*

4 *Academy of Sciences, Beijing 100093, China,*

5 <sup>2</sup>*Chinese Academy of Meteorological Sciences, Beijing 100081, China,*

6 \*Corresponding author:

7 Bingrui Jia, Guangsheng Zhou

8 Institute of Botany, Chinese Academy of Sciences,

9 20 Nanxincun, Xiangshan, Haidian District, Beijing 100093, China

10 E-mail: jiabingrui@ibcas.ac.cn, gszhou@ibcas.ac.cn

11 Tel: 86-10-62836289, Fax: 86-10-82595962

12 **Abstract.** Dahurian larch (*Larix gmelinii* Rupr.) is the dominant species in both natural and  
13 planted forests in northeast China, which situated in the southernmost part of the global  
14 boreal forest biome and undergoing the greatest climatically induced changes. Published  
15 studies (1965–2015) on tree aboveground growth of *Larix gmelinii* forests in northeast China  
16 were collected in this study, critically reviewed, and a comprehensive growth data set was  
17 developed from 123 sites, which distributed between 40.85° N and 53.47° N in latitude,  
18 between 118.20° E and 133.70° E in longitude, between 130 m and 1260 m in altitude. The  
19 data set was composed of 776 entries, including growth data (mean tree height, mean DBH,  
20 mean tree volume and/or stand volume) and the associated information, i.e., geographical  
21 location (latitude, longitude, altitude, aspect and slope), climate (mean annual temperature  
22 (MAT) and mean annual precipitation (MAP)), stand description (origin, stand age, stand  
23 density and canopy density), and sample regime (observing year, plot area and number). It



24 would provide quantitative references for plantation management practices and boreal forest  
25 growth prediction under future climate change. The data set is freely available for  
26 noncommercial scientific applications, and the DOI for the data is [https://doi.org/10.1594/](https://doi.org/10.1594/PANGAEA.880984)  
27 [PANGAEA.880984](https://doi.org/10.1594/PANGAEA.880984).

28

## 29 **1 Introduction**

30 Boreal forests, the second largest biome in the world, cover about one-third of the Earth's  
31 forest area (Keenan et al., 2015). Dahurian larch (*Larix gmelinii* Rupr.) is a dominant tree  
32 species in Chinese boreal forest, which is distributed primarily in northeast China. *Larix*  
33 *gmelinii* forest is also the predominant timber source in China, occupying 55% of Chinese  
34 boreal forest area and accounting for 75% of Chinese boreal forest volume (Xu, 1998; Zhou  
35 et al., 2002). *Larix gmelinii* forest is situated in the southernmost part of the global boreal  
36 forest biome (Shugart et al., 1992) and undergoing the greatest climatically induced changes.  
37 Thus understanding the growth characteristics of *Larix gmelinii* forest in China are of critical  
38 need for management and prediction under future climate change.

39 With increased greenhouse effective and climate warming in recent years, forest carbon  
40 sink is paid more and more attention by the world. Forestation is the main measure to offset  
41 the greenhouse gas emission and increase carbon sink. China has the largest area of forest  
42 plantations in the world, approximately 79 million ha or one-fourth of world total (FAO,  
43 2015; Payn et al., 2015). *Larix gmelinii* is an important fast-growing and cold-tolerant tree  
44 species used in forestation in northeast China. *Larix gmelinii* is usually planted after fire or  
45 logging. The growth rates of *Larix gmelinii* plantations are important indexes in the  
46 assessment of forest recovery processes and carbon sequestration potentials, which could  
47 supply strategies for post-fire or post-harvest management. The data set can provide basis for  
48 evaluating and predicting the carbon sequestration and its potential of the forestation



49 activities.

50 Relating the easily measured variables (e.g. tree height, diameter) to other structural and  
51 functional characteristics, is the most common and reliable method for estimating forest  
52 biomass, net primary production and biogeochemical budgets (Luo, 1996; Fang et al., 2001).  
53 For larch forests in northeast China, synthesis studies mainly focused on biomass and net  
54 primary production with a small quantity of samples, for example, N=28 (Luo, 1996), N=17  
55 (Wang et al., 2001a), N=18 (Wang et al., 2001b; Zhou et al., 2002), N=36 (Wang et al., 2005),  
56 N=83 (Wang et al., 2008). However, large numbers of growth measurements (e.g. tree height,  
57 DBH, volume) have scarcely been studied systematically at the large scale. Therefore, a  
58 comprehensive growth data set (N=776) of Dahurian larch in northeast China was developed  
59 in this paper.

60

## 61 **2 Data and methods**

### 62 **2.1 Research origin descriptors**

63 **(1) Identity:** Growth data set of natural and planted Dahurian larch in northeast China,  
64 version 1.0

#### 65 **(2) Originators:**

66 Bingrui Jia, State Key Laboratory of Vegetation and Environmental Change, Institute of Botany,  
67 Chinese Academy of Sciences, Beijing 100093, China;

68 Guangsheng Zhou, Chinese Academy of Meteorological Sciences, Beijing 100081, China.

69 **(3) Period of Study:** From January 1965 to December 2015


70 **(4) Objectives:** We conducted a complete literature and book review of published studies on  
71 tree height, DBH, and/or volume in *Larix gmelinii* forests including natural and planted  
72 forests in order to develop a growth data set. The data set can be used to analyze growth



73 characteristics of *Larix gmelinii* forests and assess their potential productivity in future  
74 climate warming.

## 75 2.2 Site description

76 (1) **Site type:** Data were derived from 123 study sites in northeastern China. This region  
77 includes Heilongjiang, Liaoning, Jilin provinces and the eastern part of Inner Mongolia  
78 Autonomous Region (Fig. 1).



79

80 Figure 1. Study sites of *Larix gmelinii* across northeastern China, including Heilongjiang,  
81 Liaoning, Jilin provinces and the eastern part of Inner Mongolia Autonomous Region. Open  
82 circles represent natural forests and dots represent planted forests.

83 (2) **Habitat:** *Larix gmelinii* forest is naturally distributed in Great Xing'an Mountains of  
84 northeastern China. *Larix gmelinii* was usually planted in barren hills, post-fire or  
85 post-harvest area in northeast China.



86 **(3) Climate:** The climate in northeast China is controlled by the high latitude East Asia  
87 monsoon, changing from cool temperate to temperate zones from north to south, and from  
88 semi-arid to humid zones from west to east. Mean annual temperature (MAT) for these sites  
89 ranged from -6.1 to 7.0 °C, and mean annual precipitation (MAP) from 355 to 926 mm.

### 90 **2.3 Data sources**

91 Published studies (1965-2015) were collected from available online full-text databases,  
92 including China Knowledge Resource Integrated Database (<http://www.cnki.net/>), China  
93 Science and Technology Journal Database (<http://www.cqvip.com/>), Wanfang Data  
94 Knowledge Service Platform (<http://www.wanfangdata.com/>), ScienceDirect (<http://www.sciencedirect.com/>), ISI Web of Science (<http://isiknowledge.com/>) and Springer Link  
96 (<http://link.springer.com/>). The different combination of the terms “Dahurian larch (or *Larix  
97 gmelinii*)” with “height”, “diameter at breast height (or DBH)”, “tree volume” and “stand  
98 volume” were searched in full text. Meanwhile, we also looked up the related books (e.g. Ma,  
99 1992; Wang, 1992; Zhou, 1994; Yang, 2009). We attempted to compile a complete growth  
100 data set of natural and planted Dahurian larch in northeastern part of China (between 40.85 °  
101 N and 53.47 °N; between 118.20 °E and 133.70 °E).

### 102 **2.4 Data collection criteria**

103 A critical review of the literatures collected from the above-mentioned sources was  
104 conducted to obtain reliable growth data using the following criteria:

105 **(1) Scope:** The objective of this study was to provide the data for understanding growth  
106 characteristics of *Larix gmelinii* natural forests (pure *Larix gmelinii* or its proportion more  
107 than 50%) and monoculture plantations. Forest stands included in the data set were restricted  
108 to those not recently disturbed by logging, fire, or insect pests. Additionally, the following  
109 small numbers of special types were excluded: (i) *Larix gmelinii* afforestation in wetland (Li  
110 et al., 1985; Song & Li, 1990; Huang, 2011; Cui et al., 2013), pastureland (Duan, 2005), or



111 abandoned mine land (Yang et al., 2013); (ii) hybrid test between *Larix gmelinii* and other  
112 larch (Deng et al., 2010; Zhang et al., 2005); (iii) low-yield stands in hard environment, e.g.  
113 igneous rock forest (Wang et al., 1979), old man forest (Wang et al., 1991).

114 **(2) Study design and sampling:** Tree height and DBH were averaged from the measurement  
115 values of all trees in plots or with random/systematic sampling method. Tree regeneration  
116 layer, generally below 5 cm in DBH or 1.3 m in height, was neglected in sampled plots. Stem  
117 volume of individual tree was computed from felled-wood samples or local tree volume  
118 equations. Stand volume was usually calculated by multiplying mean individual volume with  
119 stand density. Besides the growth data (i.e. height, DBH and/or volume), the necessary  
120 information should be provided in the original sources, e.g., study site, stand origin, stand age,  
121 etc.

122 **(3) Quality control:** The data quality has been carefully reviewed by the authors. Data has  
123 undergone substantial checking, for example, cross-check for the relevant information from  
124 different sources, preliminary correlation analysis among growth variables.

125 Consequently, 776 records that met the above criteria were selected to develop a  
126 comprehensive growth data set of Dahurian larch in China. The data set includes growth  
127 characteristics of Dahurian larch (i.e. mean tree height (m), mean DBH (cm), mean tree  
128 volume ( $10^{-3} \text{ m}^3$ ) and/or stand volume ( $\text{m}^3/\text{ha}$ )). In addition, associate information was  
129 included, if available in original sources or ascertainable from other relevant literatures, i.e.,  
130 geographical location (province location and locality name of study site, latitude (°),  
131 longitude (°), altitude (m), aspect and slope (°), stand description (origin, stand age (years),  
132 stand density (trees/ha) and canopy density), climate (mean annual temperature (MAT, °C)  
133 and mean annual precipitation (MAP, mm)), and sample regime (observing year, plot size and  
134 number).

## 135 **2.5 Data structural descriptors**



136 Table 1 Variable information in the data set.

Column code	Definition	Unit	Number	Range
ID	Unique identification number of each record	N/A	776	1—776
Province	Province location of study site	N/A	776	N/A
Study site	Locality name of study site	N/A	123	N/A
Latitude	Latitude of study site	°	776	40.85—53.47
Longitude	Longitude of study site	°	776	118.20—133.70
Altitude	Altitude of study site	m	776	130—1260
Aspect	Slope direction of study site, including flat slope (FL), sunny slope (SU: South), half-sunny slope (HSU: West, Southwest, Southeast), shady slope (SH: North) and half-shady slope (HSH: East, Northwest, Northeast)	N/A	300	N/A
Slope	Slope degree of study site	°	357	0—60
Origin	Stand origin was classified into natural and planted forests	N/A	776	N/A
MAT	Mean annual temperature, from original study or other related reference	°C	776	-6.1—7.0
MAP	Mean annual precipitation, from original study or other related reference	mm	776	355—926
Age	Stand age, which is generally defined as age since germination in natural forest and since planting in planted forest. Stand age is usually obtained from historical records or tree rings.	years	776	1—280
Height	Mean tree height	m	670	0.24—29.40
DBH	Mean diameter at breast height, base diameter was only given in some young forests and marked with <sup>B</sup>	cm	661	0.70—34.89
V <sub>tree</sub>	Mean tree volume, the estimated tree volume data from the two-variable larch equation were marked with <sup>E</sup>	10 <sup>-3</sup> m <sup>3</sup> /tree	696	0.04—935.73
V <sub>stand</sub>	Stand volume, the estimated stand volume data from the estimated tree volume and stand density were marked with <sup>E</sup>	m <sup>3</sup> /ha	590	0.07—975.32
Density	Stand density/Canopy density, planting density was only given in some studies and marked with <sup>P</sup>	trees/ha %	656 150	213—13275 0.2—1.0
Area	Plot area	m <sup>2</sup>	397	50—10000
Plot	Plot numbers, i.e. replications	N/A	573	1—25
Year	Investigation year	N/A	533	1954—2014
Reference	Data sources, the sources used to supplement climate information lacking in the original publications were added asterisks.	N/A	226	1965—2015

137

138 **3 Data estimates and evaluation**139 **3.1 Geographical location**

140 Google Earth (Version: 7.1.8.3036) was used to estimate latitude, longitude, and/or altitude

141 when the geographic coordinates was unavailable in original sources.

142 **3.2 Tree and stand volume**


143 The missing tree and stand volumes were estimated with the available information (mean

144 DBH, mean tree height and stand density). Stem volume of individual tree was calculated



145 with the larch equation in northeast China ( $V_{\text{tree}} = 0.000050168241 \text{ DBH}^{1.7582894} \text{ H}^{1.1496653}$ ),  
146 the best method recommended by the national standard of China: Tree volume tables (LY  
147 208-77) (Forestry Administration of China, 1978; Liu, 2017). Meanwhile, available 590 pairs  
148 of mean tree height and mean DBH in the data set were used to establish the linear H-DBH  
149 correlation ( $R^2=0.8377$ , see Fig. 2). To calculate tree volume from only one known variable  
150 of DBH, tree height was firstly calculated with the linear H-DBH equation in Fig. 2. The  
151 estimated stand volume was determined by multiplying the estimated tree volume with stand  
152 density.


153 The accuracy of tree volume was assessed by comparing the extracted data from references  
154 with the calculated data from the above-mentioned two-variable tree volume equation (Fig. 3).  
155 The coefficient of determination ( $R^2$ ) was 0.9724 and the slope was 1.0737. Therefore, we  
156 were confident in applying the larch volume equation to interpolate tree and stand volume  
157 data in this study.



158

159 Figure 2. Relationship between mean tree height and diameter at breast height in the data set.





160

161 Figure 3. Comparison of available tree volume from references with simulated values from  
162 the two-variable tree volume equation ( $V_{\text{tree}}=0.000050168241 \text{ DBH}^{1.7582894} \text{ H}^{1.1496653}$ ). Open  
163 circles (N=317): tree height (H) and diameter at breast height (DBH) were available in the  
164 references, solid circles (N=37): DBH was only available in the references and H was  
165 estimated with the H-DBH correlation from Fig. 2.

166

167 **Acknowledgements.** We would like to acknowledge the many scientists and researchers  
168 whose field measurements made this work possible. This research was supported by the  
169 National Key Research and Development Program of China (2017YFC0503906) and China  
170 Special Fund for Meteorological Research in the Public Interest (GYHY201406034).

#### 171 **References**

172 Cui, W., Mu, C. C., Lu, H. C., Bao, X., and Wang, B.: Effects of draining for forestation on  
173 carbon storage of wetland ecosystem in Daxing'an Mountains of northeastern China,  
174 Journal of Beijing Forestry University, 35(5), 28-36, <https://doi.org/10.13332/j.1000-1522>.



- 175 2013.05.020, 2013.
- 176 Deng, J. F., Zhang, H. G., Zhang, L., Guan, C. Y., and Zhang, L.: Genetic variation of  
177 17-year-old hybrid larch and its superior family selection, *Journal of Northeast Forestry*  
178 *University*, 38(1), 8-11, <https://doi.org/10.13759/j.cnki.dlxb.2010.01.017>, 2010.
- 179 Duan, W. B.: Suitable types of soils for main tree species of Shelterbelts on pastureland,  
180 *Chinese Journal of Soil Science*, 36(6), 871-876, [https://doi.org/10.19336/j.cnki.trtb.](https://doi.org/10.19336/j.cnki.trtb.2005.06.012)  
181 2005.06.012, 2005.
- 182 Fang, J., Chen, A., Peng, C., Zhao, S., Ci, L.: Changes in forest biomass carbon storage in  
183 China between 1949 and 1998, *Science*, 292(5525), 2320-2322, [https://doi.org/10.1126/](https://doi.org/10.1126/science.1058629)  
184 [science.1058629](https://doi.org/10.1126/science.1058629), 2001.
- 185 FAO: Global Forest Resources Assessment 2015, UN Food and Agriculture Organization,  
186 Rome, Italy, available at: <http://www.fao.org/3/a-i4808e.pdf>, last access: 8 October 2017,  
187 2015.
- 188 Forestry Administration of China: Tree Volume Tables (National standard # LY 208-77),  
189 Standard Press of China, Beijing, China, 1978.
- 190 Huang, D. Q.: Biomass and carbon storage estimation of *Larix gmelinii*, Thesis, available at:  
191 <http://d.wanfangdata.com.cn/Thesis/Y2026419>, last access: 8 October 2017, 2011.
- 192 Keenan, R. J., Reams, G. A., Achard, F., de Freitas, J. V., Grainger, A., and Lindquist, E.:  
193 Dynamics of global forest area: Results from the FAO Global Forest Resources  
194 Assessment 2015, *Forest Ecology and Management*, 352, 9-20, [https://doi.org/10.1016/](https://doi.org/10.1016/j.foreco.2015.06.014)  
195 [j.foreco.2015.06.014](https://doi.org/10.1016/j.foreco.2015.06.014), 2015.
- 196 Li, J. C., Zhao Y. Y., Gao Y. Q., and Liu G. L.: A preliminary analysis of changes in ecological  
197 factors after reforestation in swamp land, *Scientia Silvae Sinicae*, 21(1), 80-85, available at:  
198 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE198501009.htm>, last access: 8  
199 October 2017, 1985.
- 200 Liu, Q. J.: Tree volume tables of China, China Forestry Publishing House, Beijing, China,  
201 2017.
- 202 Luo, T. X.: Patterns of net primary productivity for Chinese major forest types and their  
203 mathematical models, Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-](http://cdmd.cnki.com.cn/Article/CDMD-80061-2006113572.htm)  
204 [80061-2006113572.htm](http://cdmd.cnki.com.cn/Article/CDMD-80061-2006113572.htm), last access: 8 October 2017, 1996.
- 205 Ma, C. G.: Selection of optimum species and seed sources for plantations of larch, Beijing  
206 Agricultural university press, Beijing, China, 1992.
- 207 Payn, T., Carnus, J. M., Freer-Smith, P., Kimberley, M., Kollert, W., Liu, S. R., Orazio, C.,  
208 Rodriguez, L., Silva, L. N., and Wingfield, M. J.: Changes in planted forests and future  
209 global implications, *Forest Ecology and Management*, 352, 57-67, [https://doi.org/10.1016/](https://doi.org/10.1016/j.foreco.2015.06.021)  
210 [j.foreco.2015.06.021](https://doi.org/10.1016/j.foreco.2015.06.021), 2015.
- 211 Shugart, H. H., Leemans, R., and Bonan, G. B.: A system analysis of the global boreal forest,  
212 Cambridge University Press, Cambridge, 1-10, 1992.
- 213 Song, X. S. and Li, Y. M.: Study on the effect of improvement of fen land and afforestation of  
214 *Larix gmelinii*, *Forestry Science & Technology*, 6, 5-7, [https://doi.org/10.13456/j.cnki.lykt.](https://doi.org/10.13456/j.cnki.lykt.1990.06.003)  
215 [1990.06.003](https://doi.org/10.13456/j.cnki.lykt.1990.06.003), 1990.
- 216 Wang, H. Q., Wu, J. C., Yang, W. B., Wang, S. T., and Song, C. J.: A research on management  
217 and utilization of "old man forest" of *Larix gmelinii*, *Forestry Science & Technology*, 16(3),  
218 12-14, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ199103003.htm>,



- 219 last access: 8 October 2017, 1991.
- 220 Wang, W. J., Zu, Y. G., Wang, H. M., Matsuura, Y., Sasa, K., and Koike, T.: Plant biomass and  
221 productivity of *Larix gmelinii* forest ecosystems in Northeast China: intra- and inter-  
222 species comparison, Eurasian Journal of Forest Research, 8, 21-41, available at:  
223 <http://hdl.handle.net/2115/22188>, last access: 8 October 2017, 2005.
- 224 Wang, X. P., Fang, J. Y., and Zhu, B.: Forest biomass and root-shoot allocation in northeast  
225 China, Forest Ecology and Management, 255, 4007-4020, [https://doi.org/10.1016/j.foreco.](https://doi.org/10.1016/j.foreco.2008.03.055)  
226 2008.03.055, 2008.
- 227 Wang, Y. H., Zhou, G. S., and Jiang, Y. L.: Simulation of NPP and evaluation of ecosystem  
228 services of *Larix gmelinii* forest, Chinese Journal of Applied Ecology, 12(5), 648-652,  
229 <https://doi.org/10.13287/j.1001-9332.2001.0157>, 2001a.
- 230 Wang, Y. H., Zhou, G. S., Jiang, Y. L., and Yang, Z. Y.: Estimating biomass and NPP of Larix  
231 forests using forest inventory data(FID), Acta Phytocologica Sinica, 25(4), 420-425,  
232 available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-ZWSB200104006.htm>, last  
233 access: 8 October 2017, 2001b.
- 234 Wang, Z.: Larch Forest in China, China Forestry Publishing House, Beijing, China, 1992.
- 235 Wang, Z., Xu, Z. B., Tan, Z. X., and Dai, H. C.: Ecological characteristics of *Larix gmelinii*  
236 grown in igneous rock in Aershan forest region, Scientia Silvae Sinicae, 4, 300-303,  
237 available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE197904010.htm>, last  
238 access: 8 October 2017, 1979.
- 239 Xu, H.: Forests in Daxinganling Mountains China, Science Press, Beijing, China, 22-53,  
240 1998.
- 241 Yang, C. P.: Study on the provenance of Dahurian larch, Science Press, Beijing, China, 2009.
- 242 Yang, Y., Zhang, K., Cai, T. J., and Gu, J. F.: Effect of afforestation species on soil nutrient in  
243 dumping site of open coal mine in Daxing'an Mountain, Journal of Northeast Forestry  
244 University, 41(9), 54-58, <https://doi.org/10.13759/j.cnki.dlxb.2013.09.028>, 2013.
- 245 Zhang, H. G., Zhang, C. L., Lan, S. B., Kang, Y. K., and Pan, B. L.: The analysis and  
246 selection for families of hybrid larch, Journal of Nanjing Forestry University: Natural  
247 Science Edition, 29(3), 69-72, <https://doi.org/10.3969/j.issn.1000-2006.2005.03.017>, 2005.
- 248 Zhou, G. S., Wang, Y. H., Jiang, Y. L., and Yang, Z. Y.: Estimating biomass and net primary  
249 production from forest inventory data: a case study of China's Larix forests, Forest  
250 Ecology and Management, 169(1-2), 149-157, [https://doi.org/10.1016/s0378-1127\(02\)](https://doi.org/10.1016/s0378-1127(02)00305-5)  
251 00305-5, 2002.
- 252 Zhou, X. F.: Long-Term Research on China's Forest Ecosystems, Northeast Forestry  
253 University Press, Harbin, China, 1994.
- 254 **References cited in the dataset**
- 255 An, J.: Tree species and management effect on soil carbon sequestration and fertilities of  
256 plantation forests. Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-10225-](http://cdmd.cnki.com.cn/Article/CDMD-10225-1012445928.htm)  
257 [1012445928.htm](http://cdmd.cnki.com.cn/Article/CDMD-10225-1012445928.htm), last access: 8 October 2017, 2011.
- 258 Bi, Y. H.: Study on stand growth models of larch *gmelinii* with different thinning intensity.  
259 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2001005669.htm>, last  
260 access: 8 October 2017, 2001.
- 261 Cao, G. Z.: Analysis of hydrological characteristics and change trend in western region of  
262 Songhua River Basin. Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10183->



- 263 1011098650.htm, last access: 8 October 2017, 2011
- 264 Chen, J. D., Pan, B. L., Wu, M. C., Han, C. W., and Yan, W. F.: A summary report on the  
265 introduction of *Larix principis-rupprechtii*, Forestry Science & Technology, 4, 2-5,  
266 available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ198304000.htm>, last  
267 access: 8 October 2017, 1983.
- 268 Chen, L. X. and Xiao, Y.: Evolution and evaluation of soil fertility in forest land in *Larix*  
269 *gmelinii* plantations at different development stages in Daxinganling forest region, Science  
270 of Soil and Water Conservation, 4(5), 50-55, <https://doi.org/10.16843/j.sswc.2006.05.012>,  
271 2006.
- 272 Chen, Q., Zhao, Y. S., Xin, Y., Han, X. C., Xia, X. Y., and Han, C. H.: Plant species diversity  
273 and correlation with soil factors after vegetation restoration in burned area of Great  
274 Xing'an Mountains, Bulletin of Soil and Water Conservation, 32(6), 61-65, 70,  
275 <https://doi.org/10.13961/j.cnki.stbctb.2012.06.040>, 2012a.
- 276 Chen, Q., Zhao, Y. S., Xin, Y., and Han, X. C.: Litter and soil water-holding capacity of  
277 burned area in northern slope of the Great Khingan after vegetation restoration, Chinese  
278 Journal of Soil Science, 44(2), 308-313, <https://doi.org/10.19336/j.cnki.trtb.2013.02.008>,  
279 2013.
- 280 Chen, R. M.: A preliminary report on provenance trials of *Larix gmelinii*, Forest Investigation  
281 Design, 1, 45-49, available at: [http://www.cnki.com.cn/Article/CJFDTOTAL-](http://www.cnki.com.cn/Article/CJFDTOTAL-LYKC198901014.htm)  
282 [LYKC198901014.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-LYKC198901014.htm), last access: 8 October 2017, 1989.
- 283 Chen, X. W., Wang, C. H., Ding, B. Y., and Zhang, S. Y.: Dynamic thinning system for larch  
284 plantation, Journal of Northeast Forestry University, 5(2), 17-21,  
285 <https://doi.org/10.1007/BF02843226>, 1994.
- 286 Chen, Y. L., Han, S. J., Zhou, Y. M., Zou, C. J., and Zhang, J. H.: Characteristics of available  
287 P in the rhizosphere soil in pure *Juglans mandshurica* and *Larix gmelinii* and their mixed  
288 plantation, Chinese Journal of Applied Ecology, 13(7), 790-794, [https://doi.org/10.13287/](https://doi.org/10.13287/j.1001-9332.2002.0188)  
289 [j.1001-9332.2002.0188](https://doi.org/10.13287/j.1001-9332.2002.0188), 2012b.
- 290 Cheng, F. Y.: Spatial variations in non-structural carbohydrates of *Larix gmelinii*, *Juglans*  
291 *mandshurica* and *Ulmus japonica*. Thesis, available at: [http://cdmd.cnki.com.cn/Article/](http://cdmd.cnki.com.cn/Article/CDMD-10225-1015665453.htm)  
292 [CDMD-10225-1015665453.htm](http://cdmd.cnki.com.cn/Article/CDMD-10225-1015665453.htm), last access: 8 October 2017, 2015.
- 293 Cheng, Y. X. and Li, Z. X.: Preliminary study of biomass of three forest types of *Larix*  
294 *gmelinii*, Inner Mongolia Forestry Investigation and Design, 4, 29-39, available at:  
295 <http://www.cqvip.com/QK/90816X/198904/206948.html>, last access: 8 October 2017,  
296 1989.
- 297 Chu, X.: Spatial and temporal distribution of fine root biomass of dominant tree species and  
298 changes after fire on Great Xing'an Mountains. Thesis, available at:  
299 <http://cdmd.cnki.com.cn/Article/CDMD-10225-1011147000.htm>, last access: 8 October  
300 2017, 2011.
- 301 Chu, X., Di, X. Y., Zhang, J. L., Cai, H. Y., and Yan, B. Z.: Distribution and seasonal  
302 dynamics of fine root biomass of two types of forests in Great Xing'an Mountains, Journal  
303 of Northeast Forestry University, 39(5), 36-39, [https://doi.org/10.13759/j.cnki.dlxb.](https://doi.org/10.13759/j.cnki.dlxb.2011.05.033)  
304 [2011.05.033](https://doi.org/10.13759/j.cnki.dlxb.2011.05.033), 2011.
- 305 Cui, G. F., Cai, T. J., and Yang, W. H.: Soil acidity of *Larix gmelinii* plantation, Journal of  
306 Beijing Forestry University, 22(3), 33-36, <https://doi.org/10.13332/j.1000-1522.2000.03>.



- 307 008, 2000.
- 308 Di, X. Y., Geng, Y. Y., Sun, L., and Hu, H. Q.: Soil microbial biomass dynamics and influence  
309 factors in *Larix gmelinii* forest in Daxing'an Mountains, Journal of Northeast Forestry  
310 University, 40(5), 1-4, <https://doi.org/10.13759/j.cnki.dlxb.2012.05.009>, 2012.
- 311 Ding, B. Y.: The method and intensity of intermediate cuttings for larch plantation, Journal of  
312 Northeast Forestry University, 16(2), 46-57, [https://doi.org/10.13759/j.cnki.dlxb.1988.02.](https://doi.org/10.13759/j.cnki.dlxb.1988.02.007)  
313 007, 1988.
- 314 Ding, B. Y., Ju, Y. G., Zhang, S. S., Zhang, S. Y., and Yu, W. J.: Study on the structure of the  
315 larch wood community, Journal of North-Eastern Forestry Institute, 4, 11-23,  
316 <https://doi.org/10.13759/j.cnki.dlxb.1982.04.002>, 1982.
- 317 Ding, B. Y., Liu, S. R., and Cai, T. J.: Studies on biological productivity of artificial forests of  
318 Dahurian larches, Acta Phytocologica et Geobotanica Sinica, 14(3), 226-236, available at:  
319 <http://www.cnki.com.cn/Article/CJFDTOTAL-ZWSB199003003.htm>, last access: 8  
320 October 2017, 1990.
- 321 Dong, X. B.: The impacts of cutting intensity on the growth of larch forest, Journal of  
322 Northeast Forestry University, 29(1), 44-47, [https://doi.org/10.13759/j.cnki.dlxb.2001.01.](https://doi.org/10.13759/j.cnki.dlxb.2001.01.011)  
323 011, 2001.
- 324 Dong, X. B.: Modes and evaluation of ecological harvesting in Larch plantation areas. Thesis,  
325 available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009132292.htm>, last access:  
326 8 October 2017, 2008.
- 327 Dong, X. B., Yang, X. C., and Yang, G. X.: Impacts of different cutting modes on soil  
328 properties in Larch plantations, Journal of Northeast Forestry University, 35(10), 7-10,  
329 <https://doi.org/10.13759/j.cnki.dlxb.2007.10.012>, 2007.
- 330 Dong, Y.: A survey study on the factors influencing the growth of *Larix gmelinii* and *Pinus*  
331 *sylvestris* plantations, Transactions of China Pulp and Paper, Suppl., 134-142,  
332 <https://doi.org/10.3321/j.issn:1000-6842.2004.z1.021>, 2004.
- 333 Du, E. Z. and Fang J. Y.: Linking belowground and aboveground phenology in two boreal  
334 forests in Northeast China, Oecologia, 176, 883-892, [https://doi.org/10.1007/s00442-014-](https://doi.org/10.1007/s00442-014-3055-y)  
335 3055-y, 2014.
- 336 Du, Y. J., Xu, H. C., and Yu, R. Y.: The effect of understory, litter and small mammals on  
337 seedling occurrence of *Larix gmelinii*, Journal of Beijing Forestry University, 15(4), 12-20,  
338 <https://doi.org/10.13332/j.1000-1522.1993.04.002>, 1993.
- 339 E, W. F.: Stem respiration spatial-temporal dynamics and characteristics of sapwood/  
340 heartwood of *Larix gmelinii*, Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-](http://cdmd.cnki.com.cn/Article/CDMD-10225-2009133891.htm)  
341 [10225-2009133891.htm](http://cdmd.cnki.com.cn/Article/CDMD-10225-2009133891.htm), last access: 8 October 2017, 2008.
- 342 E, W. F., Wang C. K., Yang C. P., Wang, X. C., Zhang, Q. Z., and Zhang, Y. Q.: Effects of  
343 provenances on growth characteristics of sapwood and heartwood of *Larix gmelinii* trees,  
344 Scientia Silvae Sinicae, 45(6), 109-115, [https://doi.org/10.3321/j.issn:1001-7488.](https://doi.org/10.3321/j.issn:1001-7488.2009.06.019)  
345 2009.06.019, 2009.
- 346 Feng, L. and Yang, Y. G.: A study on biomass and production of three types of Dahurian larch  
347 virgin forest, Scientia Silvae Sinicae, 21(1), 86-92, available at: [http://www.cnki.com.cn/](http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE198501010.htm)  
348 [Article/CJFDTOTAL-LYKE198501010.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE198501010.htm), last access: 8 October 2017, 1985.
- 349 Fu, Y.: Study on *Larix gmelinii* plantation in different planting density in the south of  
350 Daxing'anling City, Journal of Inner Mongolia Forestry, 5, 24-26, available at:



- 351 <http://www.cnki.com.cn/Article/CJFDTotal-NMLY198405030.htm>, last access: 8 October  
352 2017, 1984.
- 353 Gao, M. F., Li, C. H., and He, S. S.: Productivity division of *Larix gmelinii* in Great Xing'an  
354 Mountains of Inner Mongolia, Inner Mongolia Forestry Investigation and Design, 28,  
355 75-77, 84, <https://doi.org/10.13387/j.cnki.nmld.2005.s1.037>, 2005.
- 356 Gao, X. W.: Research by the stand density effect on Xing'an larch of natural forest, Thesis,  
357 available at: <http://cdmd.cnki.com.cn/Article/CDMD-10129-1014375561.htm>, last access:  
358 8 October 2017, 2014.
- 359 Gao, X. W., Zhang, Q. L., Bao, Y., Wang, X. H., Li, D. Y., and Shi, X.: Study on the forest  
360 biomass and carbon density under different cutting patterns of Dahurian Larch forest,  
361 Forest Resources Management, 3, 91-96, [https://doi.org/10.13466/j.cnki.lyzygl.2014.03.](https://doi.org/10.13466/j.cnki.lyzygl.2014.03.020)  
362 020, 2014.
- 363 Gu, Y. C.: The necessity of tending and thinning of natural young and middle aged forest,  
364 Forest Resources Management, 1, 58-62, [https://doi.org/10.13466/j.cnki.lyzygl.1985.01.](https://doi.org/10.13466/j.cnki.lyzygl.1985.01.018)  
365 018, 1985a.
- 366 Gu, Y. C.: The succession of forest communities in the forest region of the Greater Xingan  
367 Mountains, Acta Phytoecologica et Geobotanica Sinica, 9(1), 64-70, available at:  
368 <http://www.cnki.com.cn/Article/CJFDTOTAL-ZWSB198501006.htm>, last access: 8  
369 October 2017, 1985b
- 370 Guan, J. Q., Cai, T. J., and Man, X. L.: Characteristics of snowpack chemistry in different  
371 type plantations of Xiaoxing'an Mountains, northeastern China, Journal of Beijing Forestry  
372 University, 35(4), 41-46, <https://doi.org/10.13332/j.1000-1522.2013.04.020>, 2013.
- 373 Guo, R. Y., Fu, Y., Gao, S. C., and Jia, Z. S.: Studies on the reasonable planting density of  
374 Poplar and larch in Xing-an League, Inner Mongolia Forestry Science and Technology, 2,  
375 20-22, 35, available at: [http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198402005.](http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198402005.htm)  
376 htm, last access: 8 October 2017, 1984.
- 377 Guo, Y. X.: The variation characteristics of heartwood and sapwood of artificial *Larix*  
378 *gmelinii*, Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-10225-1015665595.](http://cdmd.cnki.com.cn/Article/CDMD-10225-1015665595.htm)  
379 htm, last access: 8 October 2017, 2015.
- 380 Guo, Y. X. and Jiang, L. C.: Variation of heartwood and sapwood and growth characteristics  
381 of artificial Dahurian Larch, Journal of Northeast Forestry University, 43(6), 93-97,  
382 <https://doi.org/10.13759/j.cnki.dlxb.20150522.048>, 2015.
- 383 Han, J. Y., Zhang, H. G., Jia, Q. B., Zhang, Z., and Xu, Y. L.: Genetic variations in growth  
384 traits of juvenile stage hybrid larch in southern Xiao Xing'an Mountains and selection of  
385 superior families, Journal of Northwest Forestry University, 27(6), 78-82,  
386 <https://doi.org/10.3969/j.issn.1001-7461.2012.06.16>, 2012a.
- 387 Han, M. Z.: A research on biomass and production of *Larix Dahurica* forest communities,  
388 Acta Agriculturae Boreali-Sinica, 2(4), 134-138, [https://doi.org/10.3321/j.issn:1000-7091.](https://doi.org/10.3321/j.issn:1000-7091.1987.04.022)  
389 1987.04.022, 1987.
- 390 Han, S. L., Zhang, Q. L., AN, H. J., and Yu, B.: Spatial structure characteristics of *Larix*  
391 *gmelinii* natural forest, Journal of Arid Land Resources and Environment, 29(5), 87-92,  
392 <https://doi.org/10.13448/j.cnki.jalre.2015.157>, 2015a.
- 393 Han, X. C.: Effect on process of burned areas *Larix gmelinii* plantations restoration in  
394 Daxinganling, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225->



- 395 1013371072.htm, last access: 8 October 2017, 2013.
- 396 Han, X. C., Zhao, Y. S., Xin, Y., and et al.: Soil hydrology characteristics of *Larix gmelinii*  
397 plantation in burned forestland in the north of Greater Xing'an Mountains, Journal of Soil  
398 and Water Conservation, 26(4), 183-188, [https://doi.org/10.13870/j.cnki.stbcxb.2012.04.](https://doi.org/10.13870/j.cnki.stbcxb.2012.04.053)  
399 053, 2012b.
- 400 Han, X. C., Zhao, Y. S., and Xin, Y.: Succession process of *Larix gmelinii* forest with  
401 artificial restoration after fire in Daxing'anling, Science of Soil and Water Conservation,  
402 13(2), 70-76, <https://doi.org/10.16843/j.sswc.2015.02.012>, 2015b.
- 403 He, H. Y.: Analysis on growth of three species of artificial plantation in Zhangguangcai  
404 Mountains-subregion of Xiaoxingan Mountains, Journal of Northeast Forestry University,  
405 16(1), 32-36, <https://doi.org/10.13759/j.cnki.dlxb.1988.01.005>, 1988.
- 406 Hong, Q. L., Cai, Y. X., Wang, Y. H., Zhao, H. X., Hu, C. X., Li, F. R., and Fan, W. Y.: The  
407 growth of natural forests of *Larix gmelinii* in Tahe Forestry Bureau Daxingan Mountain  
408 District, Journal of Northeast Forestry University, 22(2), 92-97,  
409 <https://doi.org/10.13759/j.cnki.dlxb.1994.02.017>, 1994.
- 410 Hu, H. Q., Wei, S. J., and Sun, L.: Estimation of carbon emissions from forest fires in 2010 in  
411 Huzhong of Daxing'anling Mountain, Scientia Silvae Sinicae, 48(10), 109-119,  
412 <https://doi.org/10.11707/j.1001-7488.20121017>, 2012.
- 413 Huang, C., Meng, F. L., Shan, C. C., and Hua, S.: Techniques for fast growing and high yield  
414 of larch, Journal of Heilongjiang Vocational Institute of Ecological Engineering, 24(1),  
415 28-29, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-HSGX201101014.htm>,  
416 last access: 8 October 2017, 2011a.
- 417 Huang, D. Q.: Biomass and carbon storage estimation of *Larix gmelinii*, Thesis, available at:  
418 <http://d.wanfangdata.com.cn/Thesis/Y2026419>, last access: 8 October 2017, 2011.
- 419 Huang, D. Q., Hu, H. Q., Wang, Z. W., and Sun, L.: Dynamic changes in biomass and  
420 productivity of *Larix gmelinii* plantation after fire, Journal of Northeast Forestry University,  
421 39(9), 62-64, <https://doi.org/10.13759/j.cnki.dlxb.2011.09.025>, 2011b.
- 422 Huang, S. Z.: The internal cycling of nutrients in fine roots of *Fraxinus Mandshurica* and  
423 *Larix gmelinii*, Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-10225-](http://cdmd.cnki.com.cn/Article/CDMD-10225-2006110390.htm)  
424 2006110390.htm, last access: 8 October 2017, 2006.
- 425 Huo, H.: The effects of climate warming on photosynthesis and growth of *Larix gmelinii*.  
426 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2007187763.htm>, last  
427 access: 8 October 2017, 2007.
- 428 Jia, S. X., Zhao, Y. L., Sun, Y., Chen, L., and Wang, Z. Q.: Effects of nitrogen fertilization on  
429 seasonal dynamics of soil microbial biomass carbon and nitrogen in *Larix gmelinii* and  
430 *Fraxinus mandshurica* plantations, Chinese Journal of Applied Ecology, 20(9), 2063-2071,  
431 <https://doi.org/10.13287/j.1001-9332.2009.0380>, 2009.
- 432 Jiang, H. Y.: Study on hydrological characteristics of forest ecosystem in Daxing'anling,  
433 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009132313.htm>, last  
434 access: 8 October 2017, 2008.
- 435 Jiang, H. Y., Zhao, Y. S., Chen, X. W., Li, W. H., Zhu, W. C., Lv, W. B., and Li, X. P.:  
436 Research on soil hydrology characteristics of some main forest type in south part of  
437 Daxing'anling, Journal of Soil and Water Conservation, 21(3), 149-153,187,  
438 <https://doi.org/10.13870/j.cnki.stbcxb.2007.03.033>, 2007.



- 439 Jiang, H. Y., Zhao, Y. S., Xin, X. J., Ma, W. H., Li, X. P., Sun, C. K., Guo, X. W.:  
440 Precipitation distribution laws of different forest stand in Daxing'Anling, Journal of Soil  
441 and Water Conservation, 22(6), 197-201, [https://doi.org/10.13870/j.cnki.stbcxb.2008.](https://doi.org/10.13870/j.cnki.stbcxb.2008.06.044)  
442 06.044, 2008.
- 443 Jiang, L. F.: Study on mechanism of photosynthesis and respiration of *Larix gmelinii*  
444 plantations, Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-10225-](http://cdmd.cnki.com.cn/Article/CDMD-10225-2003110988.htm)  
445 2003110988.htm, last access: 8 October 2017, 2003.
- 446 Jiang, L. F., Shi, F. C., Li, B., Luo, Y. Q., Chen, J. Q., and Chen, J. K.: Separating rhizosphere  
447 respiration from total soil respiration in two larch plantations in northeastern China, Tree  
448 Physiology, 25, 1187-1195, <https://doi.org/10.1093/treephys/25.9.1187>, 2005.
- 449 Jiang, L. F., Shi, F. C., Zu, Y. G., Wang, W. J., and Koike, T.: Study on stem respiration of  
450 *Larix gmelinii* of different ages and its relationship to environmental factors, Bulletin of  
451 Botanical Research, 23(3), 296-301, <https://doi.org/10.3969/j.issn.1673-5102.2003.03.012>,  
452 2003.
- 453 Jiang, R., Sun, L., and Hu, H. Q.: The seasonal dynamics of soil microbial biomass of  
454 a *Larix gmelinii* forest after wildfire, African Journal of Microbiology Research, 6(10),  
455 2328-2336, <https://doi.org/10.5897/ajmr11.1264>, 2012.
- 456 Laboratory of Forest Tree and Breeding, Northeast Forestry University(LFTB): Study on  
457 larch heterosis, Forestry Science & Technology, 1, 1-3, [https://doi.org/10.13456/j.cnki.lykt.](https://doi.org/10.13456/j.cnki.lykt.1979.01.001)  
458 1979.01.001, 1979.
- 459 Li, B.: The impact of different densities on the *Larix gmelinii* plantation. Thesis, available at:  
460 <http://cdmd.cnki.com.cn/Article/CDMD-10129-1013248427.htm>, last access: 8 October  
461 2017, 2013a.
- 462 Li, B., De, Y. J., Wei, Q., Wang, Y., Wang, Y. B., and Bao, Y. Q.: Effects of three densities on  
463 the growth of *Larix gmelinii* plantations, Journal of Inner Mongolia Agricultural University,  
464 34(2), 65-68, <https://doi.org/10.16853/j.cnki.1009-3575.2013.02.013>, 2013.
- 465 Li, J., Luo, Y. Q., Shi, J., Zhou, J., Wang, X. L., Ma L. Y., and Chen C.: Use of  
466 phyto-attractant in monitoring and controlling gypsy moth, Journal of Beijing Forestry  
467 University, 33(4), 85-90, <https://doi.org/10.13332/j.1000-1522.2011.04.010>, 2011.
- 468 Li, L. N., Zhao, L., Zhang, W. J., and Wang, X. J.: Study on species diversity in different  
469 density of *Larix gmelinii* plantations in Zhalantun Region, Practical Forestry Technology, 8,  
470 9-11, <https://doi.org/10.13456/j.cnki.lykt.2014.08.002>, 2014.
- 471 Li, M., Gu, J. C., Zhang, Z. W., and Wang, Z. Q.: Responses of fine root mass, length,  
472 production and turnover to soil nitrogen fertilization in *Larix gmelinii* and *Fraxinus*  
473 *mandshurica* forests in Northeastern China, Journal of Forest Research, 15, 194-201,  
474 <https://doi.org/10.1007/s10310-009-0176-y>, 2010a.
- 475 Li, Q., Chen, J., Li, T., and Liu, F. J.: Ecological function of artificial coniferous timber  
476 forests in water conservation, Jilin Forestry Science and Technology, 36(3), 14-17, 20,  
477 <https://doi.org/10.16115/j.cnki.issn.1005-7129.2007.03.004>, 2007.
- 478 Li, S. L. and Chen, Y. L.: Decomposition and nutrient return of the leaf litter under the pure  
479 and mixed plantations of *Juglans mandshurica* and *Larix gmelinii*, Journal of Nanjing  
480 Forestry University: Natural Science Edition, 28(5), 59-62,  
481 <https://doi.org/10.3969/j.issn.1000-2006.2004.05.014>, 2004.
- 482 Li, X. F.: Comprehensive evaluation of the effects of different thinning intensities on *Larix*





- 483 *gmelinii* forest in Great Xing'an Mountains of Inner Mongolia, Inner Mongolia Forestry  
484 Investigation and Design, 1, 7-20, available at: [http://www.cqvip.com/QK/90816X/](http://www.cqvip.com/QK/90816X/199301/1276513.html)  
485 199301/1276513.html, last access: 8 October 2017, 1993.
- 486 Li, X. F.: Study on genetic diversity of provenances and clones of *Larix gmelinii* (Rupr.) Rupr.,  
487 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009134088.htm>, last  
488 access: 8 October 2017, 2009.
- 489 Li, X. F.: Spatio-temporal dynamics and influencing factors of fine roots in five Chinese  
490 temperate forest ecosystems, Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-](http://cdmd.cnki.com.cn/Article/CDMD-10225-1013366884.htm)  
491 10225-1013366884.htm, last access: 8 October 2017, 2013b.
- 492 Li, X. Y.: The study on dynamic features of several metabolites of larch (*Larix gmelinii*),  
493 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2005136726.htm>, last  
494 access: 8 October 2017, 2005.
- 495 Li, Y. W., Chen, X. H., and Hu, M. Q.: Study on growth and mortality of mature and  
496 over-mature *Larix gmelinii* forest in Daxinganling forest region, Forest Resources  
497 Management, 1, 38-43, <https://doi.org/10.13466/j.cnki.lyzygl.1979.01.007>, 1979.
- 498 Li, Z. F., Zhao, Y. S., Xin, Y., Kong, X. F., and Wu, C. Y.: Soil water storage and infiltration  
499 characteristics under 3 artificial forests in the upper reaches of Ashihe river, Science of Soil  
500 and Water Conservation, 8(1), 77- 80, <https://doi.org/10.16843/j.sswc.2010.01.014>, 2010b.
- 501 Li, Z. F., Zhao, Y. S., Xin, Y., Kong, X. F., and Wu, C. Y.: Temporal and spatial variation of  
502 soil water under different forest types in the upper reaches of Ashihe River, Science of Soil  
503 and Water Conservation, 8(4), 86-89, <https://doi.org/10.16843/j.sswc.2010.04.016>, 2010c.
- 504 Lin, L.: Afforestation techniques on southward slopes in Greater Khingan Mts. forest district,  
505 Journal of Northeast Forestry University, 3, 48-58, available at: [http://www.cnki.com.cn/](http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198103005.htm)  
506 Article/CJFDTOTAL-DBLY198103005.htm, last access: 8 October 2017, 1981.
- 507 Liu, G. L.: Intermediate cutting experiment of larch (*Larix* sp.) plantation, Journal of  
508 North-Eastern Forestry Institute, 12(1), 33-46, [https://doi.org/10.13759/j.cnki.dlxb.](https://doi.org/10.13759/j.cnki.dlxb.1984.01.004)  
509 1984.01.004, 1984.
- 510 Liu, G. L. and Xu, M. H.: Study on half-sib breeding plantation of different larchs. Forestry  
511 Science and Technology, 2: 19-24. available at: [http://www.cnki.com.cn/Article/](http://www.cnki.com.cn/Article/CJFDTOTAL-JLLK198402003.htm)  
512 CJFDTOTAL-JLLK198402003.htm, last access: 8 October 2017, 1984.
- 513 Liu, H. L., Cai, T. J., Man, X. L., Chai, R. S., and Lang, Y.: Effects of major forest types of  
514 Xiaoxing'an Mountains on the process of snowfall, snow cover and snow melting, Journal  
515 of Beijing Forestry University, 34(2), 20-25, [https://doi.org/10.13332/j.1000-1522.](https://doi.org/10.13332/j.1000-1522.2012.02.003)  
516 2012.02.003, 2012.
- 517 Liu, J. L., Wang, C. K., and Zhang, Q. Z.: Spatial variations in stem heartwood and sapwood  
518 for *Larix gmelinii* trees with various differentiation classes, Scientia Silvae Sinicae, 50(12),  
519 114-121, <https://doi.org/10.11707/j.1001-7488.20141216>, 2014.
- 520 Liu, S. R., Chai, Y. X., Cai, T. J., and Peng, C. H.: Study on biomass and net primary  
521 productivity of Dahurian larch plantation, Journal of Northeast Forestry University, 18(2),  
522 40-46, <https://doi.org/10.13759/j.cnki.dlxb.1990.02.006>, 1990a.
- 523 Liu, S. R., Li, X. M., and Niu, L. M.: The degradation of soil fertility in pure larch plantations  
524 in the northeastern part of China, Ecological Engineering, 10, 75-86,  
525 [https://doi.org/10.1016/s0925-8574\(97\)10024-6](https://doi.org/10.1016/s0925-8574(97)10024-6), 1998.
- 526 Liu, T. Y.: Carbon storage in Manchurian ash and Dahurian larch plantations with different



- 527 stand structures. Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-](http://cdmd.cnki.com.cn/Article/CDMD-10225-1012445805.htm)  
528 10225-1012445805.htm, last access: 8 October 2017, 2012.
- 529 Liu, W. and Deng, H.: Study on the effect of different afforestation methods on the survival  
530 rate of *Larix*, Forestry Science and Technology Information, 39(4), 4-5,  
531 <https://doi.org/10.3969/j.issn.1009-3303.2007.04.003>, 2007.
- 532 Liu, X. T., Du, X. G., Liang, L., Zhang, P., Zhang, B. C., and Zhang, Y. S.: The selection of  
533 geographic provenances of *Larix gmelinii* in the eastern slope of Great Xing'an Mountain,  
534 Journal of Northeast Forestry University, 18, 9-16, [https://doi.org/10.13759/j.cnki.dlxb.](https://doi.org/10.13759/j.cnki.dlxb.1990.s2.003)  
535 1990.s2.003, 1990b.
- 536 Liu, X. T., Zhang, H. J., Wang, S. X., Lin, Y. K., and He, X. Y.: Introduction of *Larix olgensis*  
537 in Daxing'anling, Journal of Northeast Forestry University, 17(3), 18-25, available at:  
538 <http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198903002.htm>, last access: 8  
539 October 2017, 1989.
- 540 Liu, Y.: Study on the niche and the production of artificial Korean pine and Larch population.  
541 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2003111030.htm>, last  
542 access: 8 October 2017, 2002.
- 543 Liu, Y. C., Wu, M. C., Chu, Y. J., Liu, D. C., and He, Z. J.: Study on the provenance test of  
544 *Larix gmelinii*. Forestry Science and Experiment, 35, 1-17, available at: [http://www.cqvip.](http://www.cqvip.com/QK/92313X/199035/319899.html)  
545 [com/QK/92313X/199035/319899.html](http://www.cqvip.com/QK/92313X/199035/319899.html), last access: 8 October 2017, 1990c.
- 546 Lu, X., Hu, H. Q., Sun, L., and Wei, S. J.: Effects of fire disturbance on soil light fraction  
547 organic carbon of *Larix gmelinii* forest, Journal of Anhui Agricultural Sciences, 41(29),  
548 11709-11713, <https://doi.org/10.13989/j.cnki.0517-6611.2013.29.006>, 2013.
- 549 Luo, Y. J., Wang, X. K., Zhang, X. Q., and Lu, F.: Biomass and its allocation of forest  
550 ecosystems in China, China Forestry Publishing House, Beijing, China, 2013.
- 551 Ma, H. B.: Stem sap flow dynamics in relation to environmental factors of *Larix gmelinii*.  
552 Thesis, <https://doi.org/10.7666/d.y1473827>, 2009.
- 553 Ma, J. W.: Study on spectral characteristics of pure stands under different site conditions and  
554 phenological stages at Maershan Area, Journal of Beijing Forestry College, 3, 26-40,  
555 <https://doi.org/10.13332/j.1000-1522.1983.03.003>, 1983.
- 556 Ma, J. W., Zhang, H. G., Liu, L., Zhang, Z., and Jia, Q. B.: Genetic variation of hybrid larch  
557 at juvenile period and its superior family selection, Journal of Northeast Forestry  
558 University, 40(9), 1-5, <https://doi.org/10.13759/j.cnki.dlxb.2012.09.019>, 2012.
- 559 Mao, W. X., Tong, D. H., Zhang, C., Zhao, D. D., Ding, Y. Y., and Jin, S.: Spatial  
560 heterogeneity of moisture content of land surface dead fuel in larch stand and effects of  
561 sampling methods on moisture estimation, Journal of Northeast Forestry University, 40(5),  
562 29-33, <https://doi.org/10.13759/j.cnki.dlxb.2012.05.012>, 2012.
- 563 Mao, Y. Q., Zhang, J. L., Wang, F. S., Liu, L., Zhao, W., Zhang, H. P., Yuan, L. D., Wang, F. J.,  
564 and Li, B. C.: Study on the provenance test of *Larix olgensis* and *Larix gmelinii*, Protection  
565 Forest Science and Technology, 2, 10-15, [https://doi.org/10.13601/j.issn.1005-5215.](https://doi.org/10.13601/j.issn.1005-5215.1993.02.003)  
566 1993.02.003, 1993.
- 567 Mao, Y. Q., Zhang, J. L., Wang, F. S., Liu, L., and Wu, C. J.: A study on introduction of Larch.  
568 Scientia Silvae Sinicae 34(4): 48-58. available at: [http://www.cnki.com.cn/Article/](http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE804.006.htm)  
569 [CJFDTOTAL-LYKE804.006.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE804.006.htm), last access: 8 October 2017, 1998.
- 570 Meng, C., Wang, L. H., Wang, J., You, X. F., Liu, Y. Q., and Xing, J.: Diurnal change in soil



- 571 respiration rate in five types of plantations during non-growing season, Journal of  
572 Northeast Forestry University, 37(11), 77-79, 91, [https://doi.org/10.13759/j.cnki.dlxb.](https://doi.org/10.13759/j.cnki.dlxb.2009.11.029)  
573 2009.11.029, 2009.
- 574 Meng, X. Q., Li, H. Y., Wang, Y. M., Zhang, Y. W., and Yang, W. Y.: Study on the effects of  
575 *Coleophora dahurica* on tree growth, Journal of Liaoning Forestry Science & Technology,  
576 5, 19-20, available at: [http://www.cnki.com.cn/Article/CJFDTOTAL-LNLK200005007.](http://www.cnki.com.cn/Article/CJFDTOTAL-LNLK200005007.htm)  
577 [htm](http://www.cnki.com.cn/Article/CJFDTOTAL-LNLK200005007.htm), last access: 8 October 2017, 2000.
- 578 Mu, T. M. and Yan, W.: Preliminary study on the evapotranspiration in Xingan larch forest,  
579 Journal of Northeast Forestry University, 2, 21-30, available at: [http://www.cnki.com.cn/](http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198202001.htm)  
580 [Article/CJFDTOTAL-DBLY198202001.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198202001.htm), last access: 8 October 2017, 1982.
- 581 Na, D. C.: A study on genetic diversity of different geographic provenances of *Larix*  
582 *gmelinii*(Rupr.) Rupr and its utilization, Thesis, available at: [http://cdmd.cnki.com.cn/](http://cdmd.cnki.com.cn/Article/CDMD-10225-2005136986.htm)  
583 [Article/CDMD-10225-2005136986.htm](http://cdmd.cnki.com.cn/Article/CDMD-10225-2005136986.htm), last access: 8 October 2017, 2005.
- 584 Na, D. C., Yang, C. P., Jiang, J., Xia, D. A., and Wang, H. R.: Provenance division and  
585 superior provenance choice of *Larix gmelinii*, Forestry Science & Technology, 30(4), 1-4,  
586 <https://doi.org/10.3969/j.issn.1001-9499.2005.04.001>, 2005.
- 587 Piao, C. B., Zhang, W. X., Lin, J., and Ma, A. B.: The preliminary report on the good natural  
588 Dahurian larch stand transformation for seed stand, Forestry Science & Technology, 16(3),  
589 17-23, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ199103005.htm>,  
590 last access: 8 October 2017, 1991.
- 591 Qi, G., Wang, Q. L., Wang, X. C., Qi, L., Wang, Q. W., Ye, Y. J., and Dai, L. M.: Vegetation  
592 carbon storage in *Larix gmelinii* plantations in Great Xing'an Mountains, Chinese Journal  
593 of Applied Ecology, 22(2), 273-279, <https://doi.org/10.13287/j.1001-9332.2011.0072>, 2011.
- 594 Qi, G., Wang, Q. L., Wang, X. C., Yu, D. P., Zhou, L., Zhou, W. M., Peng, S. L., and Dai, L.  
595 M.: Soil organic carbon storage in different aged *Larix gmelinii* plantations in Great  
596 Xing'an Mountains of Northeast China, Chinese Journal of Applied Ecology, 24(1), 10-16,  
597 <https://doi.org/10.13287/j.1001-9332.2013.0122>, 2013.
- 598 Qiu, L.: Temporal dynamic change of soil pool in larch plantations and its contribution to the  
599 forest carbon budget in NE China, Thesis, available at: [http://cdmd.cnki.com.cn/Article/](http://cdmd.cnki.com.cn/Article/CDMD-10225-1011146228.htm)  
600 [CDMD-10225-1011146228.htm](http://cdmd.cnki.com.cn/Article/CDMD-10225-1011146228.htm), last access: 8 October 2017, 2011.
- 601 Qiu, L., Zu, Y. G., Wang, W. J., Sun, W., Su, D. X., and Zheng, G. Y.: CO<sub>2</sub> flux characteristics  
602 and their influence on the carbon budget of a larch plantation in Maershan region of  
603 Northeast China, Chinese Journal of Applied Ecology, 22(1), 1-8,  
604 <https://doi.org/10.13287/j.1001-9332.2011.0031>, 2011.
- 605 Quan, X., Wang, C., Zhang, Q., Wang, X., Luo, Y., and Bond-Lamberty, B.: Dynamics of fine  
606 roots in five Chinese temperate forests, Journal of Plant Research, 123, 497-507,  
607 <https://doi.org/10.1007/s10265-010-0322-9>, 2010.
- 608 Ren, B. P.: Research on growth process and stand density of *Larix gmelinii* plantation,  
609 Journal of Inner Mongolia Radio & TV University, 2, 73-75, 85,  
610 <https://doi.org/10.16162/j.issn.1672-3473.2010.02.022>, 2010.
- 611 Ren, L. J., Li, G. T., Sun, X. J., Yu, S. P., Zheng, X. W., and Yue, Y. J.: Research on  
612 community structure and species diversity of *Larix gmelinii* community in Inner Mongolia  
613 Daxing'anling Mountains, Forest Resources Management, 3, 83-87, 118,  
614 <https://doi.org/10.13466/j.cnki.lyzygl.2013.03.023>, 2013.



- 615 Shang, Y. L., Zhang, S. H., Chen, Z. C., and Wang, G. Y.: Fine family selection and genetic  
616 gain of *Larix gmelinii*, Journal of Northeast Forestry University, 38(7), 123-125,  
617 <https://doi.org/10.13759/j.cnki.dlxb.2010.07.017>, 2010.
- 618 Sheng, H. C., Cai, T. J., Zhu, D. G., and Shi, F. C.: Rainfall redistribution and hydrochemical  
619 characteristics in the Larch plantation, Journal of Soil and Water Conservation, 23(2),  
620 79-83, 89, <https://doi.org/10.13870/j.cnki.stbcb.2009.02.050>, 2009.
- 621 Si, H. Z., Yang, S., Li, J., Wang, F. S. and Wen, B. Y.: A study on introduction of *Larix*  
622 *kaempferi*, Forest Science and Technology, 11, 5-7, <https://doi.org/10.13456/j.cnki.lykt.1999.11.002>, 1999.
- 623
- 624 Song, C. L., Zhao, P. W., Su, R. N., Hai, L., Zhou, M., and Feng, L.: Dynamic study of  
625 photosynthetic characteristics of *Larix gmelinii*, Journal of Inner Mongolia Agricultural  
626 University, 29(4), 49-54, available at: [http://www.cnki.com.cn/Article/CJFDTOTAL-](http://www.cnki.com.cn/Article/CJFDTOTAL-NMGM200804010.htm)  
627 [NMGM200804010.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-NMGM200804010.htm), last access: 8 October 2017, 2008a.
- 628 Song, S., Gu, J. C., Quan, X. K., Guo, D. L. and Wang, Z. Q.: Fine-root decomposition of  
629 *Fraxinus mandshurica* and *Larix gmelinii* plantations, Journal of Plant Ecology, 32(6),  
630 1227-1237, <https://doi.org/10.3773/j.issn.1005-264x.2008.06.003>, 2008b.
- 631 Song, W. Y.: Carbon density and its distribution of *Pinus koraiensis* and *Larix gmelinii*  
632 plantations in different regions of Heilongjiang province. Thesis, available at:  
633 <http://cdmd.cnki.com.cn/Article/CDMD-10225-1014402349.htm>, last access: 8 October  
634 2017, 2013.
- 635 Song, X. S., Bao, F. L., and Zhang, Y. L.: The analysis of the economic benefit on the fast  
636 growing and high yield of *Larix gmelini* plantation, Forestry Science & Technology, 17(2),  
637 16-17, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ199202004.htm>,  
638 last access: 8 October 2017, 1992.
- 639 Song, X. S., Guo, S. Y., and Wang, J. X.: Study on fast growing and high yield of *Larix*  
640 *gmelini* plantation, Forestry Science & Technology, 21(3), 22-24, available at:  
641 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ603.009.htm>, last access: 8 October  
642 2017, 1996.
- 643 Sun, C. Z., Shen, G. F., Li, J. Y., and Jia, L. M.: Study on the present condition and the  
644 potentialities of the productivity of main tree species plantation of china II. study on the  
645 plantation productivities of Eucalyptus, Larix and Pinus sylvestris var. Mongolica, Forest  
646 Research, 14(6), 657-667, <https://doi.org/10.13275/j.cnki.lykxyj.2001.06.012>, 2001.
- 647 Sun, H. Z., Chen, M. Y., Cai, C. J., and Zhu, N.: Effects of different type urban forest  
648 plantations on soil fertility, Chinese Journal of Applied Ecology, 20(12), 2871-2876,  
649 <https://doi.org/10.13287/j.1001-9332.2009.0454>, 2009.
- 650 Sun, H. Z., Lu, X. J., Chen, M. Y., Cai, C. J., and Zhu, N.: Soil heavy metals concentration in  
651 different type plantations in Harbin City, Chinese Journal of Applied Ecology, 22(3),  
652 614-620, <https://doi.org/10.13287/j.1001-9332.2011.0126>, 2011.
- 653 Sun, X. M., Zhang, S. G., Zhou, D. Y., Wang, X. D., Ding, B. and Liu, S. M.: Phenological  
654 variation of Larix species and their intra-species and inter-species hybrid families and early  
655 selection, Scientia Silvae Sinicae, 44(1), 77-84, [https://doi.org/10.3321/j.issn:1001-7488.](https://doi.org/10.3321/j.issn:1001-7488.2008.01.013)  
656 2008.01.013, 2008.
- 657 Sun, Y.: Effects of soil fauna on root biomass, morphology, production and turnover rate in  
658 Larch and Ash plantations, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD->



- 659 10225-2010242477.htm, last access: 8 October 2017, 2010.
- 660 Sun, Y., Wang, L. L., Chen, J., and Duan J. P.: Reconstructing mean maximum temperatures  
661 of May–August from tree-ring maximum density in North Da Hinggan Mountains, China,  
662 Chinese Science Bulletin, 57(16), 2007-2014, <https://doi.org/10.1007/s11434-012-5055-9>,  
663 2012.
- 664 Sun, Z. B. and Sun, X. W.: Selection of species and seed sources for *Larix* spp. in Chifeng  
665 Region, Inner Mongolia, in: Ma, C.G., ed. Selection of optimum species and seed sources  
666 for plantations of larch, Beijing Agricultural University Press, Beijing, China, 1992.
- 667 Sun, Z. B., Wang, X.W., Ding, M.X., Shi, J.S., Liu, Z., and Bao, G.Z.: Study on selection of  
668 optimum species and seed sources for plantations of larch, Forest Science and Technology,  
669 10, 11-13, <https://doi.org/10.13456/j.cnki.lykt.1991.10.005>, 1991.
- 670 Tang, J. Z., Li, Q. Y., Wang, H. X., and Wang, Y. G.: The main raw material forest of larch  
671 growth indicators, Forest Investigation Design, 3, 61-63, available at:  
672 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKC200903029.htm>, last access: 8  
673 October 2017, 2009.
- 674 Tian, F., Bi, X. H., Shen, G. T., Zhou, B. and Li, W. B.: Study on the forecast technique and  
675 control index of larch caterpillar of a cycle in two years, Forest Research, 11(4), 388-395,  
676 <https://doi.org/10.13275/j.cnki.lykxyj.1998.04.008>, 1998.
- 677 Wang, B. S.: The undergrowth succession in larch plantations before and after cutting,  
678 Journal of Northeast Forestry University, 24(5), 82-86, [https://doi.org/10.13759/  
679 j.cnki.dlxb.1996.05.014](https://doi.org/10.13759/j.cnki.dlxb.1996.05.014), 1996.
- 680 Wang, C. K.: Biomass allometric equations for 10 co-occurring tree species in Chinese  
681 temperate forests, Forest Ecology and Management, 222, 9-16,  
682 <https://doi.org/10.1016/j.foreco.2005.10.074>, 2006.
- 683 Wang, C. K., Gower, S. T., Wang, Y. H., Zhao, H. X., Yan, P., and Bond-Lamberty, B.: The  
684 influence of fire on carbon distribution and net primary production of boreal *Larix gmelinii*  
685 forests in north-eastern China, Global Change Biology, 7, 719-730,  
686 <https://doi.org/10.1046/j.1354-1013.2001.00441.x>, 2001a.
- 687 Wang, C. K. and Yang, J. Y.: Rhizospheric and heterotrophic components of soil respiration  
688 in six Chinese temperate forests, Global Change Biology, 13, 123-131,  
689 <https://doi.org/10.1111/j.1365-2486.2006.01291.x>, 2007.
- 690 Wang, H. X., Zhang, S. H., Jing, C. Q., Dai, J. G., Liu, Y. T. and Zhou, J. Y.: Cultivation  
691 techniques for raw material forest of short rotation Larch, Forestry Science & Technology,  
692 25(5), 7-8, <https://doi.org/10.3969/j.issn.1001-9499.2000.05.003>, 2000a.
- 693 Wang, J. Z.: Study on selection of larch species, Jilin Forestry Science and Technology, 1,  
694 22-26, available at: <http://www.cqvip.com/QK/96402X/199501/1795612.html>, last access:  
695 8 October 2017, 1995.
- 696 Wang, L.: Regeneration rules of the north forest region in western slope of Inner Mongolia  
697 Daxing'anling, Inner Mongolia Forestry Investation and Design, S1, 65-69,  
698 <https://doi.org/10.13387/j.cnki.nmld.2000.s1.022>, 2000.
- 699 Wang, L., Li, C. H., Yang, J. Y., and Guo, Y. L.: Biomass changes of artificial *Larix gmelinii*  
700 from 2002 to 2012 in Helong, Jilin Province, Journal of Green Science and Technology, 2,  
701 11-12, <https://doi.org/10.3969/j.issn.1674-9944.2013.02.005>, 2013a.
- 702 Wang, L. H., Xin, Y., Zou, M. L., Zhao, Y. S., Lei, Y. Y., and Chen, L.: Litter water-holding



- 703 characteristics in the process of vegetation restoration in severely burned area of Great  
704 Xing'an Mountains, *Journal of Soil and Water Conservation*, 28(6), 262-267,  
705 <https://doi.org/10.13870/j.cnki.stbcxb.2014.06.048>, 2014.
- 706 Wang, L. H., Xin, Y., Zou, M. L., Zhao, Y. S., Liu, S. J., and Ren, Q. S.: Plants diversity and  
707 biomass distribution of vegetation restoration in burned area of Great Xing'an Mountains,  
708 *Journal of Beijing Forestry University*, 37(12), 41-47, [https://doi.org/10.13332/j.1000-1522.](https://doi.org/10.13332/j.1000-1522.20150025)  
709 20150025, 2015a.
- 710 Wang, Q. B., Zhang, H. J., Zhang, S. L., Wang, C. B., and Xue, Y.: Species composition and  
711 structure investigation of typical plant community in Daliangzihe National Forest Park,  
712 *Protection Forest Science and Technology*, 7, 14-17, 38, [https://doi.org/10.13601/](https://doi.org/10.13601/j.issn.1005-5215.2015.07.006)  
713 [j.issn.1005-5215.2015.07.006](https://doi.org/10.13601/j.issn.1005-5215.2015.07.006), 2015b.
- 714 Wang, Q. J., Liu, Y. T., Dai, J. G., and Luo, Y. F.: Effects of different provenances and ABT1  
715 rooting powder on *Larix gmelinii* afforestation, *Forestry Science & Technology*, 25(4), 4-5,  
716 <https://doi.org/10.3969/j.issn.1001-9499.2000.04.002>, 2000b.
- 717 Wang, Q. Y.: Genetic variation of the provenance for *Picea koraiensis*, Thesis, available at:  
718 <http://cdmd.cnki.com.cn/Article/CDMD-10225-2003110985.htm>, last access: 8 October  
719 2017, 2003.
- 720 Wang, W. Z., Chen, J., Zhang, B. Y., Feng, Y. L. and Wang, Y. T.: Study on the photosynthetic  
721 characteristics and the primary productivity of *Larix olgensis*, *Journal of Northeast*  
722 *Forestry University*, 22(4), 15-21, <https://doi.org/10.13759/j.cnki.dlxb.1994.04.003>, 1994.
- 723 Wang, X. F., Li, G. T., Wu, H. B., Lv, L. K., and Yue, Y. J.: Research on structure and species  
724 diversity of *Larix gmelinii* community in Orqohan, *Journal of Southwest Forestry*  
725 *University*, 33(5), 69-73, <https://doi.org/10.3969/j.issn.2095-1914.2013.05.012>, 2013b.
- 726 Wang, X. H., Zhang, Q. L., Guo, Y. D., Xing, T. T., Wang, F., Gao, X. W., and Wu, J.: Study  
727 on the coarse root biomass and distribution characteristics about natural *Larix gmelinii*  
728 forest. *Forest Resources Management*, 5, 62-66, [https://doi.org/10.13466/j.cnki.lyzygl.](https://doi.org/10.13466/j.cnki.lyzygl.2013.05.020)  
729 2013.05.020, 2013c.
- 730 Wang, X. W. and Mao, Z. J.: Temporal and spatial variation in gas exchange in canopy of  
731 *Larix gmelinii* plantation, *Scientia Silvae Sinicae*, 43(11), 43-49,  
732 <https://doi.org/10.3321/j.issn:1001-7488.2007.11.008>, 2007.
- 733 Wang, X. W. and Mao, Z. J.: Practicability of 7 light responsive curve models to different  
734 plant species, *Bulletin of Botanical Research*, 29(1), 43-48, available at:  
735 <http://www.cnki.com.cn/Article/CJFDTOTAL-MBZW200901012.htm>, last access: 8  
736 October 2017, 2009.
- 737 Wang, Y. H., Zhou, G. S., and Jiang, Y. L.: Simulation of NPP and evaluation of ecosystem  
738 services of *Larix gmelinii* forest, *Chinese Journal of Applied Ecology*, 12(5), 648-652,  
739 <https://doi.org/10.13287/j.1001-9332.2001.0157>, 2001a.
- 740 Wang, Y. L.: Biomass, carbon density and distribution characteristics in different age *Larix*  
741 *gmelinii* plantations, *Journal of Anhui Agricultural Sciences*, 43(15), 144-145,150,  
742 <https://doi.org/10.13989/j.cnki.0517-6611.2015.15.053>, 2015.
- 743 Wang, Z.: Larch Forest in China, China Forestry Publishing House, Beijing, China, 73-150,  
744 1992.
- 745 Wu, C. Y., Zhao, Y. S., Xin, Y., Kong, X. F. and Li, Z. F.: Effects of structure of the *Larix*  
746 *gmelinii* plantation on soil hydrology characteristic, *Science of Soil and Water*



- 747 Conservation, 7(5), 58-62, <https://doi.org/10.16843/j.sswc.2009.05.011>, 2009.
- 748 Wu, X. J., Cai, T. J., Li, H., and Sheng, H. C.: Precipitation hydrochemical characteristic in  
749 virgin *Pinus koraiensis* forest and artificial *Larix gmelinii* forest in Liangshui National  
750 Nature Reserve, Science of Soil and Water Conservation, 6(6), 37-42,  
751 <https://doi.org/10.16843/j.sswc.2008.06.007>, 2008.
- 752 Wu, X. J., Jia, E. L., and Li, J. C.: Preliminary study on the regeneration before clear cutting  
753 in *Larix gmelinii* plantation, Forest Science and Technology, 8, 24-26,  
754 <https://doi.org/10.13456/j.cnki.lykt.1983.08.010>, 1983.
- 755 Wu, X. L.: Interannual variability of temperature, precipitation and sunshine hours in  
756 Wuerqihan Forest Region, Heilongjiang Agricultural Sciences, 10, 113-116,  
757 <https://doi.org/10.11942/j.issn1002-2767.2015.10.0113>, 2015.
- 758 Wu, Z. C.: Introduction and afforestation of *Larix olgensis*, Journal of Inner Mongolia  
759 Forestry, 12, 20-21, available at: [http://www.cnki.com.cn/Article/CJFDTOTAL-](http://www.cnki.com.cn/Article/CJFDTOTAL-NMLY198612016.htm)  
760 [NMLY198612016.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-NMLY198612016.htm), last access: 8 October 2017, 1986.
- 761 Xia, C. C., Liu, Z. L., Wang, Q. C., and Li, K. L.: Comparison of the growth and biomass  
762 production of monoculture and mixed stands of *Larix gmelinii* and *Betula platyphylla*,  
763 Journal of Northeast Forestry University, 40(10), 1-3,  
764 <https://doi.org/10.13759/j.cnki.dlxb.2012.10.021>, 2012.
- 765 Xiao, S. L. and Yang, J. L.: Individual tree aboveground biomass of *Larix gmelinii* natural  
766 forest in the Northern Greater Khingan Mountains, Scientia Silvae Sinicae, 50(8), 22-29,  
767 <https://doi.org/10.11707/j.1001-7488.20140804>, 2014.
- 768 Xiao, Y., Man, X. L., and Fan, J. F.: Soil fertility and water conservation function of main  
769 forest types in Gongbiela River Watershed, Journal of Northeast Forestry University, 34(1),  
770 28-30, <https://doi.org/10.13759/j.cnki.dlxb.2006.01.010>, 2006.
- 771 Xin, Y., Zhao, Y. S., and Pan, B. Y.: Influence of *Larix gmelinii* plantations at Eastern  
772 Mountainous Region in Heilongjiang Province on water quality, Science of Soil and Water  
773 Conservation, 4(2), 29-33, <https://doi.org/10.16843/j.sswc.2006.02.006>, 2006.
- 774 Xin, Y., Zou, M. L., and Zhao, Y.S.: Soil water storage capacity and infiltration properties of  
775 *Larix gmelinii* plantation in different growth age after burned in Greater Khingan  
776 Mountains, Chinese Journal of Soil Science, 45(4), 823-829,  
777 <https://doi.org/10.19336/j.cnki.trtb.2014.04.009>, 2014.
- 778 Xu, H. C. and Du, Y. J.: Amount of falling seed and seedling occurrence dynamics in  
779 primitive *Larix gmelinii* forest, Scientia Silvae Sinicae, 29(4), 298-306, available at:  
780 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKE199304001.htm>, last access: 8  
781 October 2017, 1993.
- 782 Xu, H. C., Yu, R. Y., Guo, Z. L., Zhou, C. H., and Ma, D. G.: Study on germination conditions  
783 of *Larix gmelinii* seedlings in Great Xing'an Mountains, Journal of Beijing Forestry  
784 University, 12, 15-25, <https://doi.org/10.13332/j.1000-1522.1990.s4.004>, 1990.
- 785 Xu, H. L., Man, X. L., and Sheng, H. C.: Water conservation function of different types of  
786 *Larix gmelinii* forests in Da Hinggan Mountains, Research of Soil and Water Conservation,  
787 18(4), 92-96, available at: [http://www.cnki.com.cn/Article/CJFDTOTAL-](http://www.cnki.com.cn/Article/CJFDTOTAL-STBY201104020.htm)  
788 [STBY201104020.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-STBY201104020.htm), last access: 8 October 2017, 2011a.
- 789 Xu, H. L., Man, X. L., and Sheng, H. C.: Comprehensive evaluation on water conservation  
790 function of the different types of larch natural forests in Da Hinggan Mountains, Chinese



- 791 Agricultural Science Bulletin, 29(19), 74-78, available at: [http://www.cnki.com.cn/Article/](http://www.cnki.com.cn/Article/CJFDTOTAL-ZNTB201319017.htm)  
792 [CJFDTOTAL-ZNTB201319017.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-ZNTB201319017.htm), last access: 8 October 2017, 2013.
- 793 Xu, J. L.: Study on *Larix* biomass model and biomass estimation in Dongzhelenghe, Thesis,  
794 available at: <http://cdmd.cnki.com.cn/Article/CDMD-10022-2009161609.htm>, last access:  
795 8 October 2017, 2009.
- 796 Xu, S. Z. and Han, M. Z.: Regeneration rule and management decision of *Larix gmelinii* in  
797 Genhe regions, in Zhou, X. F. ed., Long-Term Research on China's Forest Ecosystems,  
798 709-716, Northeast Forestry University Press, Harbin, China, 1994.
- 799 Xu, X. T., Wan, D. Y., Wei, D. F., and Zhou, J. Y.: Growth characteristics and techniques for  
800 fast growing and high yield of larch, Inner Mongolia Forestry Investigation and Design,  
801 34(6), 54-55, <https://doi.org/10.13387/j.cnki.nmld.2011.06.057>, 2011b.
- 802 Xu, Y. B., Zhou, Y. Q., Wu, R. X., Liu, Z. S., and Yu, C. H.: Afforestation techniques of *Larix*  
803 *gmelinii* container seedlings, Forestry Science & Technology, 21(3), 5-7, available at:  
804 <http://www.cnki.com.cn/Article/CJFDTOTAL-LYKJ603.001.htm>, last access: 8 October  
805 2017, 1996.
- 806 Xu, Y. Q.: Study on the geographic variations and selection for the optimal provenances of  
807 *Larix gmelinii*, Forest Inventory and Planning, 3, 15-23, available at:  
808 <http://www.cnki.com.cn/Article/CJFDTOTAL-LDGH199303004.htm>, last access: 8  
809 October 2017, 1993.
- 810 Xue, Z. H.: Investigation of larch plantation in Daxing'anling Mountains, Inner Mongolia  
811 Forestry Science and Technology, 2, 18-21, available at: [http://www.cnki.com.cn/Article/](http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198702003.htm)  
812 [CJFDTOTAL-NMLK198702003.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198702003.htm), last access: 8 October 2017, 1987a.
- 813 Xue, Z. H.: Study on tending and thinning of natural young and middle-age *Larix gmelinii*  
814 forest, Inner Mongolia Forestry Science and Technology, 4, 1-5, available at:  
815 <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK198704000.htm>, last access: 8  
816 October 2017, 1987b.
- 817 Xue, Z. H.: Tending and Thinning of *Larix gmelinii* plantation, Inner Mongolia Forestry  
818 Science and Technology, 3, 20-23, 30, available at: [http://www.cnki.com.cn/Article/](http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK199003004.htm)  
819 [CJFDTOTAL-NMLK199003004.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK199003004.htm), last access: 8 October 2017, 1990.
- 820 Xue, Z. H.: Cultivation and management techniques of fast-growing and high-yielding *Larix*  
821 *gmelinii* Forests, Inner Mongolia Forestry Science & Technology, 1, 11-18, available at:  
822 <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK199301004.htm>, last access: 8  
823 October 2017, 1993.
- 824 Yang, B. L., Jia, Q. B., Zhang, H. G., and Zhang, L.: Research on hybrid larch superior  
825 families selection, Forestry Science & Technology, 39(2), 1-4,  
826 <https://doi.org/10.3969/j.issn.1001-9499.2014.02.001>, 2014.
- 827 Yang, C. P.: Study on the provenance of Dahurian larch, Science Press, Beijing, China, 2009.
- 828 Yang, C. P., Jiang, J., Tang, S. S., Li, J. Y. and Wang, H. R.: The provenance test of 21- year  
829 old *Larix gmelinii* at Maershan Area, Journal of Northeast Forestry University, 30(6), 1-5,  
830 <https://doi.org/10.13759/j.cnki.dlxb.2002.06.001>, 2002.
- 831 Yang, C. P., Yang, S. W., Xia, D. A., Liu, G. F., Lv, Q. Y. and Zhang, P. G.: Study on the  
832 provenance test of Dahurian Larch (II) Selection of the best provenance, Journal of  
833 Northeast Forestry University, 19, 68-76, <https://doi.org/10.13759/j.cnki.dlxb.1991.s2.011>,  
834 1991.





- 835 Yang, H. T. and Guo, Y. X.: A primary report on the provenance test of *Larix gmelinii* and  
836 *Larix olgensis* in Huanan forest region, Journal of Northeast Forestry University, 18, 65-71,  
837 <https://doi.org/10.13759/j.cnki.dlxb.1990.s2.010>, 1990.
- 838 Yang, K., Zhu, J. J., Gu, J. C., Yu, L. Z., and Wang, Z. Q.: Changes in soil phosphorus  
839 fractions after 9 years of continuous nitrogen addition in a *Larix gmelinii* plantation,  
840 Annals of Forest Science, 72, 435-442, <https://doi.org/10.1007/s13595-014-0444-7>, 2015.
- 841 Yang, L. J., Zhang, C. J., Li, C. H., Jiang, X. L., and Pan, B. L.: Early selection of superior  
842 clone of *Larix gmelinii*, Protection Forest Science and Technology, 4, 35-37,  
843 <https://doi.org/10.13601/j.issn.1005-5215.2006.04.013>, 2006.
- 844 Yang, S. W. and Shang, L. T.: Research on selection of larch species, Journal of  
845 North-Eastern Forestry Institute, 11(4), 50-56, available at: [http://www.cnki.com.cn/](http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198304006.htm)  
846 [Article/CJFDTOTAL-DBLY198304006.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198304006.htm), last access: 8 October 2017, 1983.
- 847 Yang, S. W., Wang, Q. Y., Liu, G. F., Yang, C. P., Xia, D. A., and Bao, L. F.: Analysis on the  
848 effects of genetic improvement and the breeding program of larch, Journal of Northeast  
849 Forestry University, 20(1), 1-8, <https://doi.org/10.13759/j.cnki.dlxb.1992.01.001>, 1992.
- 850 Yang, S. W., Yang, C. P., Wang, Q. Y., Liu, G. F., Xia, D. A., and Zhang, P. G.: The  
851 provenance selection of the important tree species in the northeast part of China and Inner  
852 Mongolia Autonomous Region, Journal of Northeast Forestry University, 21(4), 1-7,  
853 <https://doi.org/10.13759/j.cnki.dlxb.1993.04.001>, 1993.
- 854 Yang, S. W., Yang, C. P., Zhang, S. Y., Peng, H. M., Li, J. T., Xia, D. A., Liu, G. F., and Wang,  
855 H. R.: A study on the provenance test of *Larix gmelinii* outside of distribution area, Journal  
856 of Northeast Forestry University, 18, 1-8, <https://doi.org/10.13759/j.cnki.dlxb.1990.s2.002>,  
857 1990.
- 858 Yang, Y. B. and Liu, Y. Y.: Discussion about planting fast-growing and high-yield plantation  
859 in the south of Daxing'anling City, Forestry Science and Technology Information, 37(2),  
860 1-2, <https://doi.org/10.3969/j.issn.1009-3303.2005.02.001>, 2005.
- 861 Yao, C. Y., Wang, Q. J., and Gao, G. Y.: Study on growth status of introduced *Larix olgensis*  
862 in Xingan League of Inner Mongolia, Inner Mongolia Forestry Science & Technology, 1,  
863 17-22, available at: <http://www.cnki.com.cn/Article/CJFDTOTAL-NMLK501.004.htm>,  
864 last access: 8 October 2017, 1995.
- 865 Yao, G. Q., Chi, G. Q., Dong, Z. Q., Hu, W. L., and Jiang, J. B.: A study on the growth and  
866 economic effectiveness of four kinds of planted Larch Forest, Journal of Shenyang  
867 Agricultural University, 20, 217-222, available at: [http://www.cqvip.com/QK/93761X/](http://www.cqvip.com/QK/93761X/198903/129531.html)  
868 [198903/129531.html](http://www.cqvip.com/QK/93761X/198903/129531.html), last access: 8 October 2017, 1989.
- 869 Yao, Y. J., Chen, X. W., and Li, C. S.: Studies on biomass and productivity for Dahurian larch  
870 plantations, Journal of Northeast Forestry University, 5(2), 13-16,  
871 <https://doi.org/10.1007/BF02843225>, 1994.
- 872 Yu, B. and Wang, L. M.: Analysis on classified stem growing characteristics of *Larix gmelinii*  
873 natural forest, Forest Research, 20(4), 452-457,  
874 <https://doi.org/10.13275/j.cnki.lykxyj.2007.04.014>, 2007.
- 875 Yu, B., Wu, J. S. G. L., Wang, B. T., and Wang, L. M.: Analysis on classified stem  
876 transformation characteristic of *Larix gmelinii* natural forest at Daxingan Mountains, Acta  
877 Ecologica Sinica, 28(11), 5750-5757, <https://doi.org/10.3321/j.issn:1000-0933.2008.11.062>,  
878 2008.



- 879 Yu, B., Wu, J. S. G. L., Wang, B. T., Wang, L. M., and Da, B. X. L. T.: Study on the  
880 distribution pattern characteristics of different stand structures in *Larix gmelinii* natural  
881 forest, *Forest Research*, 23(1), 83-88, <https://doi.org/10.13275/j.cnki.lykxyj.2010.01.013>,  
882 2010.
- 883 Yu, B., Zang, Q. L., Wang, L. M., and Wu, J. S. G. L.: Characteristics of biomass and  
884 productivity in *Larix gmelinii* natural forests with different stand structures, *Journal of*  
885 *Zhejiang A & F University*, 28(1), 52-58, [https://doi.org/10.3969/j.issn.2095-0756](https://doi.org/10.3969/j.issn.2095-0756.2011.01.009).  
886 2011.01.009, 2011.
- 887 Yu, B. J. and Yang, C. P.: Study on provenance test of 5 year old *Larix olgensis*, *Journal of*  
888 *Northeast Forestry University*, 16(3), 27-33, [https://doi.org/10.13759/j.cnki.dlxb](https://doi.org/10.13759/j.cnki.dlxb.1988.03.005).  
889 1988.03.005, 1988.
- 890 Yu, S. L. and Yuan, X. G.: Studies on selection of fine families of Hybrid Larch based on  
891 growth traits in seedling stage, *Inner Mongolia Forestry Investigation and Design*, 32(2),  
892 114-117, <https://doi.org/10.13387/j.cnki.nmld.2009.02.025>, 2009.
- 893 Yuan, G. X., Li, H. X., Chen, G. F., Lou, Y. M., and Mou, Z. J.: Analysis on the introduction  
894 of *Pinus banksiana* Lamb, *Forestry Science & Technology*, 26(5), 9-10,  
895 <https://doi.org/10.3969/j.issn.1001-9499.2001.05.004>, 2001.
- 896 Yue, Y. J., Gao, R. H., Li, G. T., Jin, J., and Wang, X. F.: Research on community structure  
897 and species diversity of *Larix gmelinii* community in Jinhe Forestry Bureau, *Journal of*  
898 *Inner Mongolia Agricultural University*, 35(2), 43-47,  
899 <https://doi.org/10.16853/j.cnki.1009-3575.2014.02.010>, 2014.
- 900 Yue, Y. J., Qu, L. C., Song, L. J., Liu, M. D., Zeng, Y. E., Li, G. T., and Wang, F. S.:  
901 Distribution patterns and diversity of species of *Larix gmelinii* in Kuduer, *Journal of Inner*  
902 *Mongolia Agricultural University*, 34(1), 42-47, [https://doi.org/10.16853/j.cnki.1009-](https://doi.org/10.16853/j.cnki.1009-3575.2013.01.009)  
903 [3575.2013.01.009](https://doi.org/10.16853/j.cnki.1009-3575.2013.01.009), 2013.
- 904 Zhai, M. P., Yi, W. L., Jia, L. M., and Ma, L. Y.: Study on the characteristics of *Betula*  
905 *platyphylla-Larix gmelinii* Mixed Forest in Xilinji Forest Region, *Journal of Beijing*  
906 *Forestry University*, 12, 86-91, <https://doi.org/10.13332/j.1000-1522.1990.s3.010>, 1990a.
- 907 Zhai, M. P., Yi, W. L., Jia, L. M., Ma, L. Y., Sun, Y. X., Liu, Z. Y., and Wang, Y. G.:  
908 Investigation of *Betula platyphylla-Larix gmelinii* Mixed Forest in Xilinji Forest Region,  
909 *Journal of Beijing Forestry University*, 12, 78-85,  
910 <https://doi.org/10.13332/j.1000-1522.1990.s3.009>, 1990b.
- 911 Zhan, J. and Yang, C. P.: Study on the provenance test of Dahurian larch (*Larix gmelinii*),  
912 *Journal of Northeast Forestry University*, 17(6), 22-30, available at:  
913 <http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY198906003.htm>, last access: 8  
914 October 2017, 1989.
- 915 Zhang, G. L., Qu, B. C., Zhang, Y. H., and Wang, H.: Growth analysis and model of *Larix*  
916 *gmelinii* plantation, *Protection Forest Science and Technology*, 4, 41-43,  
917 <https://doi.org/10.13601/j.issn.1005-5215.2012.04.056>, 2012.
- 918 Zhang, H. D. and Zhou, M.: Nutrient characteristics of *Larix gmelinii* forest ecosystem in  
919 cold temperate zone, *Journal Liaoning Forestry Science & Technology*, 3, 5-8, 22,  
920 <https://doi.org/10.3969/j.issn.1001-1714.2014.03.002>, 2014.
- 921 Zhang, J.: Study on community structure, biomass and carbon storage of *Larix gmelini*  
922 plantation, Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10022>



- 923 -2008085203.htm, last access: 8 October 2017, 2008a.
- 924 Zhang, J., Sun, Y. J., and Xu, J. L.: Research on growing process of *Larix gmelinii* plantation  
925 in Northeast of China, Journal of Northwest Forestry University, 23(6), 179-181, available  
926 at: <http://www.cnki.com.cn/Article/CJFDTOTAL-XBLX200806045.htm>, last access: 8  
927 October 2017, 2008.
- 928 Zhang, J. L., Mao, Y. Q., Wang, F. S., Liu, L., Wu, C. J., Zhang, H. P., Yuan, L. D., Zhao, W.,  
929 and Wang, F. J.: The progeny testing and fine family selection of *Larix gmelinii* and *Larix*  
930 *olgensis*, Protection Forest Science and Technology, 3, 22-26,  
931 <https://doi.org/10.13601/j.issn.1005-5215.1994.03.003>, 1994.
- 932 Zhang, L.: Study on stability of families and selection of superior families of hybrid larch.  
933 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-1011146424.htm>,  
934 last access: 8 October 2017, 2011.
- 935 Zhang, L. Y.: A study on individual tree health assessment of nature *Larix gmelinii* in  
936 Daxing'anling, Forestry Science and Technology Information, 45(1), 41-43.  
937 <https://doi.org/10.3969/j.issn.1009-3303.2013.01.018>, 2013.
- 938 Zhang, Q. B. and Fang, L.: A study of problems of planting fast-growing and high-yield  
939 plantation in secondary forest site on the south slope in Great Xing'an Mountain, Journal  
940 of Inner Mongolia Forestry College, 20(3), 37-42, available at: [http://www.cnki.com.cn/  
941 Article/CJFDTOTAL-NMGL803.007.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-NMGL803.007.htm), last access: 8 October 2017, 1998.
- 942 Zhang, S. L., Zhang, H. J., Zhang, W., Guo, Y., Li, X. Z., Zhang, Q., Lu, Y., and Yao, S. P.:  
943 Comparison of water-holding characteristics and biomass of litter under different forest  
944 types in Xiaoxingan Mountains, Bulletin of Soil and Water Conservation, 35(4), 85-90,  
945 <https://doi.org/10.13961/j.cnki.stbctb.2015.04.017>, 2015a.
- 946 Zhang, S. L., Zhang, H. J., Zhang, W., Wang, C. B., Deng, J. H., Song, Y. B., and Zhang, Q.:  
947 Soil hydrological properties of main vegetation in Down South of Xiao Xing'an Mountains,  
948 Research of Soil and Water Conservation, 22(1), 140-145,  
949 <https://doi.org/10.13869/j.cnki.rswc.2015.01.027>, 2015b.
- 950 Zhang, T. W., Yuan, Y. J., Wei, W. S., and Yu, S. L.: The tree-ring chronologies from  
951 Moerdaoga in the northeast of Inner Mongolia, Journal of Arid Land Resources and  
952 Environment, 23(12), 177-182, <https://doi.org/10.13448/j.cnki.jalre.2009.12.008>, 2009.
- 953 Zhang, X. J.: Effect of close-to-nature transformation of Larch Monoculture stands on  
954 regeneration and plant diversity. Thesis, available at: [http://cdmd.cnki.com.cn/Article/  
955 CDMD-10225-2010242257.htm](http://cdmd.cnki.com.cn/Article/CDMD-10225-2010242257.htm), last access: 8 October 2017, 2010.
- 956 Zhang, X. J., Wang, Q. C., Wang, S. L., and Sun, Q.: Effect of the close-to-nature  
957 transformation of *Larix gmelinii* pure stands on plant diversity of understory vegetation in  
958 Xiaoxing'an Mountains of China, Scientia Silvae Sinicae, 47(1), 6-14,  
959 <https://doi.org/10.11707/j.1001-7488.20110102>, 2011.
- 960 Zhang, Y. Q.: Transpiration characteristic of Dahurian Larch trees transplanted from a  
961 latitudinal gradient. Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-  
962 10225-2009133900.htm](http://cdmd.cnki.com.cn/Article/CDMD-10225-2009133900.htm), last access: 8 October 2017, 2008b.
- 963 Zhao, H. X., Wang, Y. H., Chai, Y. X., Hu, C. X., Li, F. R., and Fan, W. Y.: On forest growth  
964 of northern slope of Daxing'an mountains (Tahe Forest Bureau), Journal of Northeast  
965 Forestry University, 24(6), 1-8, <https://doi.org/10.13759/j.cnki.dlxb.1996.06.001>, 1996.
- 966 Zhao, P. W.: Studies on litterfall dynamics and nutrient release regularity of *Larix gmelinii* in



- 967 Great Xingan Mountains. Thesis, <https://doi.org/10.7666/d.y1473822>, 2009.
- 968 Zhao, S. J., Yang, J. M., and Bi, X. M.: Study on the fast growing and high yield of larch  
969 species and the relationships with their introduction districts, Journal of Hebei Forestry  
970 Science and Technology, 2, 2-7, available at: [http://www.cnki.com.cn/Article/  
971 CJFDTOTAL-HBLK199202001.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-HBLK199202001.htm), last access: 8 October 2017, 1992.
- 972 Zhao, X. M., Xu, S. T., Ren, J. Z., Wang, M. Z., Yang, S. M., and Jiang, S. W.: A study on  
973 provenance test of *Pinus sylvestris L. var mongolica* litvin on Korqin sandy land, Journal of  
974 Inner Mongolia Forestry College, 21(3), 44-49, available at: [http://www.cnki.com.cn/  
975 Article/CJFDTOTAL-NMGL199903008.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-NMGL199903008.htm), last access: 8 October 2017, 1999.
- 976 Zhou, B. D., Hao, W. K., Zou, Q., Zhao, K. M., Liu, Q. L., and Chen, Y. D.: Study on  
977 artificial regeneration in a clear-cut forest of Xiaoxingan Mountains in Liangshui region,  
978 Journal of North-Eastern Forestry College, 4, 19-35, available at: [http://www.cnki.com.cn/  
979 Article/CJFDTOTAL-DBLY196500001.htm](http://www.cnki.com.cn/Article/CJFDTOTAL-DBLY196500001.htm), last access: 8 October 2017, 1965.
- 980 Zhou, B. Q.: Advices on secondary forest culture on the south slope in Great Xing'an  
981 Mountain, Inner Mongolia Forestry Investigation and Design, 18-20,  
982 <https://doi.org/10.13387/j.cnki.nmld.1998.s1.008>, 1998.
- 983 Zhou, H. X., Zhang, Y. D., Sun, H. L., and Wu, S. Y.: Soil respiration in temperate secondary  
984 forest and *Larix gmelinii* plantation in Northeast China, Chinese Journal of Applied  
985 Ecology, 18(12), 2668-2674, <https://doi.org/10.13287/j.1001-9332.2007.0477>, 2007.
- 986 Zhou, J. Q., Liu, X. D., and Guo, H. W.: Surface fuel loading and relevant influencing factors  
987 of main forest types in southern Daxing'anling, Journal of Northwest A&F  
988 University(Natural Science Edition), 42(6), 131-137, [https://doi.org/10.13207/j.cnki.  
989 jnwafu.2014.06.010](https://doi.org/10.13207/j.cnki.jnwafu.2014.06.010), 2014.
- 990 Zhu, H. Y.: Main characters variation among provenances of *Fraxinus mandshurica* Rup.  
991 Thesis, available at: <http://cdmd.cnki.com.cn/Article/CDMD-10225-2009134220.htm>, last  
992 access: 8 October 2017, 2008.
- 993 Zhu, S. Y., Zhou, B., Mao, Z. J., Wang, X. W., and Sun, Y. F.: Space time dynamics of fine  
994 root biomass of six forests in Maoershan forest region, Scientia Silvae Sinicae, 42(6),  
995 13-19, <https://doi.org/10.3321/j.issn:1001-7488.2006.06.003>, 2006.
- 996 Zhu, Y.: Forest health assessment of natural *Larix gmelinii* in Great Xing'an Mountains of  
997 China, Thesis, available at: [http://cdmd.cnki.com.cn/Article/CDMD-10225-  
998 1013371221.htm](http://cdmd.cnki.com.cn/Article/CDMD-10225-1013371221.htm), last access: 8 October 2017, 2013.
- 999 Zhu, Y., Liu, Z. G., and Jin, G. Z.: Health assessment of individual trees in natural *Larix*  
1000 *gmelinii* forest in Great Xing'an Mountains of China, Chinese Journal of Applied Ecology,  
1001 24(5), 1320-1328, <https://doi.org/10.13287/j.1001-9332.2013.0294>, 2013.
- 1002 Zou, L., Tang, Q. M., and Wang, Y.: Ecological distribution of soil microorganism in pure and  
1003 mixed forests of *Pinus sylvestris var. mongolica* and *Larix gmelini*, Journal of Northeast  
1004 Forestry University, 38(11), 63-64, 79, <https://doi.org/10.13759/j.cnki.dlxb.2010.11.029>,  
1005 2010.
- 1006 Zou, M. L., Xin, Y., and Zhao, Y. S.: The influence of vegetation restoration on carbon  
1007 storage of burned Area in Greater Xing' an Mountains, Journal of Soil and Water  
1008 Conservation, 27(5), 133-137, 143, <https://doi.org/10.13870/j.cnki.stbcxb.2013.05.046>,  
1009 2013.