

Interactive comment on “A synthetic satellite dataset of *E. huxleyi* spatio-temporal distributions and their impacts on Arctic and Subarctic marine environments (1998–2016)” by Dmitry Kondrik et al.

NEUKERMANS (Referee)

griet.neukermans@obs-vlfr.fr

Received and published: 29 November 2018

Kondrik and collaborators present a 19-year satellite time series of *Emiliana huxleyi* bloom area, calcite content, and associated increase in in-water pCO₂ in four selected areas of the high-latitude northern hemisphere. The dataset is only partly unique, in the sense that a 19-year global remote sensing dataset of *E. huxleyi* bloom extent, coccolith concentration, and PIC content can also be easily obtained elsewhere. Therefore uniqueness only applies to pCO₂. This dataset could be useful, but I request a few substantial modifications that I believe are necessary to improve understanding and

[Printer-friendly version](#)

[Discussion paper](#)



quality of the dataset: (1) some flaws in the dataset (pointed out below, 1a and 1b) will need to be fixed, (2) error estimates for remotely sensed quantities must be provided, and (3) in its present form, the study/data is not correctly positioned within the state-of-the-art literature and other available datasets.

(1a) It appears from Fig. 4 that the *E. huxleyi* bloom dataset includes false positives, a problem that is particularly evident in the Bering Sea (1998-2001) where the authors have detected blooms initiating in winter and lasting about 10 months as previously reported from ocean colour remote sensing data (Iida et al., 2002). However, ship-borne measurements have identified resuspended diatom frustules as the cause of these bright waters in winter-spring instead of *E. huxleyi* blooms (Broerse et al., 2003). This invalidates the authorial *E. huxleyi* bloom detection algorithm and all derived products in the Bering Sea from late fall to spring. I further fail to see how the algorithms used by the authors (Kondrik et al. 2017; Kondrik et al. 2018) to detect *E. huxleyi* blooms present an advance to NASA's standard method of *E. huxleyi* bloom classification (Brown and Yoder, 1994), and many other subsequent bloom detection methods (Iglesias-Rodriguez et al., 2002; Iida et al., 2002; Iida et al., 2012; Moore et al., 2012). (1b) The remote sensing algorithm for pCO₂ estimation is a simple linear regression between observations of Delta_pCO₂ and remote sensing reflectance R_{rs} in a blue waveband. This relationship is strictly empirical and does not appear to have theoretical grounds; I believe the user should be aware of this. Not surprisingly, there is an enormous spread along this regression line such that for a given reflectance value the estimated Delta_pCO₂ has a confidence interval with a width of 50 ppm and even wider for denser blooms. Furthermore, the residuals of the regression are clearly unevenly distributed, with a strong tendency to underestimate Delta_pCO₂ at higher reflectances. This relationship should be explicitly stated, which is not presently the case, including all relevant regression statistics, and especially a figure showing the observations and the fitted line so that the user can better grasp the errors of the algorithm. (2) Whereas the statistics of the validation of the retrieved coccolith concentration are given in section 2.2, the accompanying figure is missing.

No uncertainty assessment is given for pCO₂ (see previous comment). (3) A 19-year global remote sensing dataset of PIC concentration merging all ocean colour satellite missions can be obtained here: <http://www.globcolour.info/> in temporal resolutions ranging from daily to monthly, spatial resolution ranging from 4km to 100km, and various geographical projections. From PIC concentration, coccolith concentration can be derived using a fixed mass per coccolith (as you do too), and PIC content can also be easily derived by combining with a climatology for Mixed layer depth available here http://www.ifremer.fr/cerweb/deboyer/mld/Surface_Mixed_Layer_Depth.php. I therefore suggest you remove all statements of uniqueness of your PIC dataset (e.g., page 2, lines 24-26). The statements on page 2 lines 11-16, "Prior to the publication of Kondrik et al. (2018), no attempts have been undertaken to retrieve from space... No concatenated time series data are available to date on the associated bloom intensity..." are thus simply incorrect. I also suggest you appropriately reference the work of (Shutler et al., 2013) entitled "Coccolithophore surface distributions in the North Atlantic and their modulation of the air-sea flux of CO₂ from 10 years of satellite Earth observation data", which is very similar to your work on remote sensing of pCO₂ in E_{hux} blooms, but is mentioned nowhere. Page 2 Line 8-10: "Until recently, only few satellite studies were published on the typical locations of E. huxleyi blooms and associated concentrations of PIC in surface waters within the bloom area". It appears to me you missed a vast body of literature: (Balch et al., 1991; Balch et al., 1996; Gordon et al., 2001; Smyth et al., 2004; Signorini and McClain, 2009; Moore et al., 2012; Hopkins et al., 2015; Balch et al., 2016; Neukermans et al., 2018) etc.

Further comments : Title : add "blooms" after "E. huxleyi" Abstract : delete "detailed information on E. huxleyi impacts within the bloom area on marine environments", as this suggests that you are detailing ecological impacts

P1, L16 : "Ongoing climate change is a background of numerous emerging hot topics." is a rather meaningless opening sentence. P1 L25 : Rivero-Calle is not the right reference for poleward expansion of coccolithophores, instead use (Winter et al., 2014;

[Printer-friendly version](#)[Discussion paper](#)

Neukermans et al., 2018). "gradually propagating in the poleward direction" ; the poleward expansion is not gradual, as expansion rates exhibit stark jumps as demonstrated in (Neukermans et al., 2018). P2, L1-4 : a lot of statements for only one reference. P2, L23 : replace 1918-2016 by 1998-2016 P2, L20 : remove "original" P3 L1 : spell out OC CCI P6 L1 : "in the cause of satellite processing" ?, rephrase P7 L10-15 and L24-28 : same paragraph appears twice. P7 L31 : "1,105,6800 km²" commas are in the wrong place

References : Balch WM, Bates NR, Lam PJ, Twining BS, Rosengard SZ, Bowler BC, Drapeau DT, Garley R, Lubelczyk LC, Mitchell C, et al. 2016. Factors regulating the Great Calcite Belt in the Southern Ocean and its biogeochemical significance. *Global Biogeochem Cycles* 30(8): 1124–1144. Wiley-Blackwell. doi: 10.1002/2016GB005414
Balch WM, Holligan PM, Ackleson SG, Voss KJ. 1991. Biological and optical properties of mesoscale coccolithophore blooms in the Gulf of Maine. *Limnol Oceanogr* 36(4): 629–643. doi: 10.4319/lo.1991.36.4.0629
Balch WM, Kilpatrick KA, Holligan P, Harbour D, Fernandez E. 1996. The 1991 coccolithophore bloom in the central North Atlantic. 2. Relating optics to coccolith concentration. *Limnol Oceanogr* 41(8): 1684–1696. doi: 10.4319/lo.1996.41.8.1684
Broerse AT., Tyrrell T, Young J., Poulton A., Merico A, Balch W., Miller P. 2003. The cause of bright waters in the Bering Sea in winter. *Cont Shelf Res* 23(16): 1579–1596. doi: 10.1016/j.csr.2003.07.001
Brown CW, Yoder JA. 1994. Coccolithophorid blooms in the global ocean. *J Geophys Res* 99(C4): 7467. doi: 10.1029/93JC02156
Gordon HR, Boynton GC, Balch WM, Groom SB, Harbour DS, Smyth TJ. 2001. Retrieval of coccolithophore calcite concentration from SeaWiFS Imagery. *Geophys Res Lett* 28(8): 1587–1590. doi: 10.1029/2000GL012025
Hopkins J, Henson SA, Painter SC, Tyrrell T, Poulton AJ. 2015. Phenological characteristics of global coccolithophore blooms. *Global Biogeochem Cycles* 29(2): 239–253. doi: 10.1002/2014GB004919
Iglesias-Rodriguez MD, Brown CW, Doney SC, Kleypas JA, Kolber D, Kolber Z, Hayes PK, Falkowski PG. 2002. Representing key phytoplankton functional groups in ocean carbon cycle models: Coccolithophorids. *Global Biogeochem Cycles* 16(4): 47-1-47–20. doi: 10.1029/2001GB001454
Iida T,

Mizobata K, Saitoh S-I. 2012. Interannual variability of coccolithophore *Emiliana huxleyi* blooms in response to changes in water column stability in the eastern Bering Sea. *Cont Shelf Res* 34: 7–17. Pergamon. doi: 10.1016/J.CSR.2011.11.007

lida T, Saitoh SI, Miyamura T, Toratani M, Fukushima H, Shiga N. 2002. Temporal and spatial variability of coccolithophore blooms in the eastern Bering Sea, 1998-2001. *Prog Oceanogr* 55(1–2): 165–175. Pergamon. doi: 10.1016/S0079-6611(02)00076-9

Moore TS, Dowell MD, Franz BA. 2012. Detection of coccolithophore blooms in ocean color satellite imagery: A generalized approach for use with multiple sensors. *Remote Sens Environ* 117: 249–263. doi: 10.1016/j.rse.2011.10.001

Neukermans G, Oziel L, Babin M. 2018. Increased intrusion of warming Atlantic water leads to rapid expansion of temperate phytoplankton in the Arctic. *Glob Chang Biol* 24(6): 2545–2553. Wiley/Blackwell (10.1111). doi: 10.1111/gcb.14075

Shutler JD, Land PE, Brown CW, Findlay HS, Donlon CJ, Medland M, Snooke R, Blackford JC. 2013. Coccolithophore surface distributions in the North Atlantic and their modulation of the air-sea flux of CO₂ from 10 years of satellite Earth observation data. *Biogeosciences* 10(4): 2699–2709. doi: 10.5194/bg-10-2699-2013

Signorini SR, McClain CR. 2009. Environmental factors controlling the Barents Sea spring-summer phytoplankton blooms. *Geophys Res Lett* 36(10): L10604. doi: 10.1029/2009GL037695

Smyth TJ, Tyrell T, Tarrant B. 2004. Time series of coccolithophore activity in the Barents Sea, from twenty years of satellite imagery. *Geophys Res Lett* 31(11): L11302. doi: 10.1029/2004GL019735

Winter A, Henderiks J, Beaufort L, Rickaby REM, Brown CW. 2014. Poleward expansion of the coccolithophore *Emiliana huxleyi*. *J Plankton Res* 36(2): 316–325. Oxford University Press. doi: 10.1093/plankt/fbt110

Interactive comment on Earth Syst. Sci. Data Discuss., <https://doi.org/10.5194/essd-2018-101>, 2018.

Printer-friendly version

Discussion paper

