

A comprehensive global oceanic dataset of helium isotope and tritium measurements

William J. Jenkins¹, Scott C. Doney^{1,2}, Michaela Fendrock¹, Rana Fine³, Toshitaka Gamo⁴, Philippe Jean-Baptiste⁵, Robert Key⁶, Birgit Klein⁷, John E. Lupton⁸, Robert Newton⁹, Monika Rhein¹⁰, Wolfgang Roether¹⁰, Yuji Sano⁴, Reiner Schlitzer¹¹, Peter Schlosser¹², Jim Swift¹³

Correspondence to: William J. Jenkins (wjenkins@whoi.edu)

Abstract. Tritium and helium isotope data provide key information on ocean circulation, ventilation, and mixing, as well as the rates of biogeochemical processes, and deep-ocean hydrothermal processes. We present here global oceanic datasets of tritium and helium isotope measurements made by numerous researchers and laboratories over a period exceeding 60 years. The dataset has a DOI:10.25921/c1sn-9631 and is available at <https://www.nodc.noaa.gov/ocads/data/0176626.xml>, or alternately <http://odv.awi.de/data/ocean/jenkins-tritium-helium-data-compilation/> and includes approximately 60,000 valid tritium measurements, 63,000 valid helium isotope determinations, 57,000 dissolved helium concentrations, and 34,000 dissolved neon concentrations. Some quality control has been applied in that questionable data have been flagged and clearly compromised data excluded entirely. Appropriate metadata has been included, including geographic location, date, and sample depth. When available, we include water temperature, salinity, and dissolved oxygen. Data quality flags and data originator information (including methodology) are also included. This paper provides an introduction to the dataset along with some discussion of its broader qualities and graphics.

¹ Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

² Now at University of Virginia, Charlottesville, VA 22904, USA

³ RSMAS, University of Miami, Miami, FL 33149, USA

⁴ Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Chiba 277-8564, Japan

⁵ LSCE, CEA-CNRS-UVSQ, CEA/Saclay, 91191 Gif-sur-Yvette cedex, France

⁶ Princeton University, Princeton, New Jersey, USA

⁷ BSH, 20359 Hamburg, Germany

⁸ NOAA Pacific Marine Environmental Laboratory, Newport, Oregon, USA.

⁹ LDEO, Columbia University, Palisades, N.Y.

¹⁰ IUP, University Bremen, D28359 Bremen, Germany

¹¹ Alfred Wegener Institute, 27568 Bremerhaven, Germany

¹² LDEO, Columbia University, now at Arizona State University, Arizona, U.S.A.

¹³ SIO, University of California San Diego, La Jolla, CA, USA

1 Introduction

The global oceanic distributions of tritium (^3H , a radioactive isotope of hydrogen with a [half-life of](#) 12.3 years), its daughter product ^3He , and helium isotopes in general arise from the complicated interplay of ocean ventilation, circulation, and mixing, with the hydrologic cycle, air-sea exchange, and geological volatile input. Observations of the delivery of tritium to the ocean and its redistribution are a useful tool for diagnosing gyre- and basin-scale ventilation and circulation (Doney et al., 1992; Doney and Jenkins, 1994; Dorsey and Peterson, 1976; Dreisigacker and Roether, 1978; Fine and Ostlund, 1977; Fine et al., 1987; Fine et al., 1981; Jenkins et al., 1983; Jenkins and Rhines, 1980; Michel and Suess, 1975; Miyake et al., 1975; Ostlund, 1982; Sarmiento, 1983; Weiss and Roether, 1980; Weiss et al., 1979).

In shallow waters, away from sea-floor hydrothermal vents, the combination of tritium and ^3He may be used to determine the time elapsed since a water parcel was at the sea surface, making it a useful tool for diagnosing ventilation and circulation on seasonal through decade time-scales (Jenkins, 1987, 1998, 1977). The ingrowth and evasion of tritiogenic ^3He from the thermocline is also useful as a flux gauge for constraining the rate of nutrient return to the ocean surface (Jenkins, 1988; Jenkins and Doney, 2003; Stanley et al., 2015) as well as upwelling in coastal regions (Rhein et al., 2010). Finally, the distribution of helium isotopes in the deep sea provides important quantitative constraints on the impact of submarine hydrothermal venting on many elements because the global hydrothermal helium flux is well known (Bianchi et al., 2010; Holzer et al., 2017; Schlitzer, 2016). This makes ^3He useful as a flux gauge (German et al., 2016; Jenkins et al., 1978; Jenkins et al., 2018; Lupton and Jenkins, 2017; Resing et al., 2015; Roshan et al., 2016). Consequently, there have been numerous measurements of these properties over the years, particularly under the aegis of major observational programs like GEOSECS (Bainbridge et al., 1987), TTO (Jenkins and Smethie, 1996), WOCE (www.nodc.noaa.gov/woce/wdiu/diu_summaries/whp/index.htm), CLIVAR (www.clivar.org), GO-SHIP (www.go-ship.org), and GEOTRACES (www.geotraces.org). It seems valuable to assemble all existing data, including those measured prior to and outside of these programs, along with appropriate metadata, in one place to facilitate further use and analysis. This is a report of these efforts.

2 Methods

2.1 Tritium measurement methodology

There are at present three distinct methods for the determination of tritium in water samples: direct measurement of tritium abundance by accelerator mass spectrometry (AMS), radioactive counting of tritium decay rate, and daughter product (^3He) ingrowth method. The first method (AMS) has not been used for the measurement of environmental tritium levels, but is better suited to measuring high tritium concentrations in small samples, largely for biomedical tracer research (Brown et al., 2005; Chiarappa-Zucca et al., 2002; Giagola et al., 1984; Roberts et al., 2000). The second method usually involves isotopic enrichment of hydrogen from the water samples either by electrolysis (e.g., Ostlund and Werner, 1962) or thermal diffusion (Israel, 1962) followed by low level radioactive counting; either by liquid scintillation (Momoshima et al., 1983) or gas

proportional counting (Bainbridge et al., 1961). Measurements are made relative to prepared standards (Unterweger et al., 1980) and accuracy appears to be limited by the reproducibility of the enrichment process to 3-10% (Cameron, 1967).

The third method, ^3He ingrowth, is a three step method. First, it involves degassing of a quantity (~1 to 1000 ml) of water to remove all dissolved helium. Second, the degassed water is stored in a helium leak-tight container (usually low He-permeability aluminosilicate glass or metal) for a period of several weeks to a year or more. Experience indicates that it is necessary to shelter the stored samples from cosmic rays since there is a latitude-dependent cosmogenic ^3He production rate that masquerades as “tritium signal” (Lott and Jenkins, 1998). Finally, the ingrown ^3He is extracted from the water sample and mass spectrometrically analysed (Clarke et al., 1976; Ludin et al., 1997). The last method, although it involves a potentially lengthy incubation period, is chemically simpler and does not involve isotopic enrichment steps. As such, it offers intrinsically greater accuracy (limited by standardization of the mass spectrometer, typically better than 1%) and lower ultimate detection limit (Jenkins et al., 1983; Lott and Jenkins, 1998).

2.2 Helium isotope measurement methodology

Water samples are usually drawn from Niskin bottles into a helium leak-tight container either for ship-board (Lott and Jenkins, 1998; Roether et al., 2013) or shore-based gas extraction. The latter involves either clamped (Weiss, 1968) or crimped copper tubing (Young and Lupton, 1983). The extracted gases are subsequently purified and concentrated, usually cryogenically (Lott, 2001; Lott and Jenkins, 1984; Ludin et al., 1997) and expanded into a mass spectrometer for isotopic analysis. While time-of-flight mass spectrometry has been used (Mamyryn, 2001; Mamyryn et al., 1970) most oceanic helium isotope measurements have been made using ~~specialty-designed~~ multi-collector magnetic sector instruments (where ions are electrostatically accelerated and deflected by a magnetic field, e.g., Bayer et al., 1989; Clarke et al., 1969; Lott and Jenkins, 1998, 1984; Ludin et al., 1997). Measurements are typically standardized to marine air and corrected for any sample-size-dependent ratio effects determined by measurement of different-sized air aliquots. Depending on the amount of tritium in the water and the length of time a water sample is stored prior to gas extraction, it is generally necessary to correct the $^3\text{He}/^4\text{He}$ results for decay of tritium during storage. For very deep and old samples, e.g., at 3000 m in the North Pacific, where tritium concentrations are very low, this correction may be inconsequential.

2.3 Helium and neon concentration measurements

To make full use of the helium isotopic ratio measurements, one needs to know at least the concentration of helium in the samples as well. This allows investigators to calculate the actual concentration of both ^3He and ^4He in samples, and with some assumptions, to estimate the amount of non-atmospheric ^3He – whether it be from tritium decay or hydrothermal input – and ^4He . The estimate can be further improved with knowledge of neon concentrations. Since neon is similar to helium in terms of its solubility in seawater and there are no known significant non-atmospheric sources of neon in the ocean, its concentration is a direct tracer of processes like air bubble injection at the sea surface. As an example, we show in Figure 1 the relationship

between the helium and neon saturation anomalies¹⁴ for approximately 2000 near-surface samples. The saturation anomalies range from significantly below zero to as much as 10-15%. The negative values may arise from lower barometric pressures or air-sea disequilibrium during warming. However, there may be systematic offsets due to laboratory standardization as well. The higher values may also reflect such biases, but also may be due to air bubble entrapment during sampling at sea. The solid line in Figure 1 represents a type-II linear regression of the data and has a slope of near unity, which is expected given the similarity of the two gases. However, the slope is greater than the precise ratio expected (0.8 to 0.9 depending on temperature) based on their solubilities and atmospheric abundances, possibly due to differing air-sea exchange rates. Consideration must be given to these and other factors when using this data (e.g., Fuchs et al., 1987; Roether et al., 1998).

Helium and neon concentration measurements are typically made by mass spectrometric peak height manometry; that is, by comparison of major isotope ion currents (⁴He⁺ and ²⁰Ne⁺) between the unknown ~~to~~and helium ~~and~~ /neon derived from an aliquot of marine air. The air aliquot size is determined from a knowledge of the barometric pressure, relative humidity, and temperature at which the previously evacuated air standard reservoir was filled, and the volume of the aliquot. Generally the ion current is assumed to be a linear function of the sample size (number of atoms) over some narrow range, but also can be corrected by construction of a standard curve using ~~different sized multiple~~ aliquots. Some measurements (notably the GEOSECS expedition) were made by splitting the ~~gas~~ extracted ~~from the~~ water sample and measuring the helium and neon contents ~~separately~~ using isotope dilution (with ³He and ²²Ne spikes).

2.4 Data organization

We have compiled a comprehensive dataset consisting of helium isotope and tritium measurements in oceanic waters made by numerous laboratories over the past 6 decades. The dataset includes ~60,000 tritium and ~63,000 helium isotope measurements, ~57,000 dissolved helium concentrations, and ~34,000 dissolved neon concentrations in ocean water taken from 1952 to 2015 (for tritium) and from 1967 to 2015 for helium. In addition~~al~~ to “spot sampling”, there are ~ 380 cruises, with sampling from >5,400 locations for tritium and ~5,600 locations for helium. The helium data are from 8 different laboratories, and the tritium data from 15 laboratories world-wide. In addition to including measurement uncertainties, a data quality flag, and data source, each data point is accompanied by location (latitude, longitude, depth) and time (decimal year) of sampling. When available, water temperature, salinity, and dissolved oxygen measurements are included.

A number of the earliest measurements were obtained from publications. In those cases, the publication source is given. If the data were transcribed from tables, the table number and page is also given. In the event that the data were only available graphically, a computer program to digitize the data from plots was used, and in the rare cases where graph quality was

¹⁴ The saturation anomaly of a gas is the percent deviation of the measured concentration relative to the expected concentration of the sample in equilibrium with air at one atmosphere, for example as defined by $\Delta He = 100 (C/C^* - 1)$ where C^* is the solubility equilibrium concentration at the temperature and salinity of the sample.

sufficiently poor to degrade the precision of the data, the uncertainties were commensurately increased to reflect it. Where data had been assigned a Digital Object Identifier (DOI), this is also included.

The dataset ([provided in several digital formats described below](#)) consists of three tables. The REFERENCES table is a list of the data sources keyed by the text variable “Reference_Code” found in the main data table. ~~This should in principle provide~~ [This provides](#) attribution ~~and/or~~ more information regarding the data origin. The METHODS table provides a more complete description of the ~~methods~~ fields “Tritium_Method” and “Helium_Method” in the main data table. This is intended to provide useful interpretive information regarding how the sampling and/or measurements were accomplished. The main data table fields are described in ~~the~~ Table 1. Most data fields have an associated quality flag field whose meaning is summarized in Table 2. Following WOCE ocean data convention, normal acceptable data are associated with a quality flag of 2, whereas questionable data have a flag of 3. Results obtained by averaging 2 or more replicates are signified with a flag of 6. When fields are missing for a given record, the data is entered as -999 and the corresponding quality flag is 9. The tritium, helium, helium isotope, and neon data also have an associated uncertainty field (e.g., “Tritium_Error”) which is the estimated uncertainty in the data points. This is either provided by the data measurer or an estimate based on described procedures, and can vary greatly between methods and laboratories so the user is advised to be aware of this value.

In the spirit of the WOCE/CLIVAR/GO-SHIP¹⁵ convention, the combination of ExpoCode, Station, CastNo, and Bottle should uniquely define a sample. That is, no two data records should have the same combination of these values. This has been followed with most of the information here: when a sample’s station, cast, or bottle number were not provided (in the case of literature data), arbitrary but unique numbers were assigned. In order to supplement this identification we added a unique integer record ID number.

2.5 Data formats and availability

The data set’s DOI is 10.25921/c1sn-9631. The data is available for download at the U.S. National Oceanic and Atmospheric Administration’s National Centers for Environmental Information web site <https://www.nodc.noaa.gov/ocads/data/0176626.xml> and ~~alternatly~~ <http://odv.awi.de/data/ocean/jenkins-tritium-helium-data-compilation/> in a number of formats ~~for maximum flexibility~~. For maximum flexibility, we suggest one of the following three database formats: Microsoft Access®, PostgreSQL, or ODV (Ocean Data View). In addition, the three tables are available as 4 files (the main data table is split in two to avoid spreadsheet row-number limitations) in Microsoft Excel®

¹⁵ WOCE is the World Ocean Circulation Experiment (e.g., see <https://www.nodc.noaa.gov/woce/>), CLIVAR is the Climate and Ocean – Variability, Predictability, and Change (e.g., see <http://www.clivar.org/about>), and GO-SHIP is the Global Ocean Ship-Based Hydrographic Investigations Program (see <http://www.go-ship.org/>)

or as a comma separated plain text files. [The data is also available in NetCDF format.](#) Finally, the data table is available for download as a MATLAB® binary data file.

3 Graphics and examples **Scope and nature of the dataset**

We provide some [example](#) graphics to indicate the scope and nature of the data holdings. These include time histories of analyses per year for both types of measurements (Figure [42](#)) and maps of sampling locations (Figure [23](#)). The intent is to provide a broad overview of the character of the data sets while not over-interpreting its details and features.

The temporal distribution of oceanic tritium measurements begins in the early 1950s with the development of enrichment and counting capabilities suitable for environmental levels, and the recognition of the existence of cosmogenic tritium production (Comog and Libby, 1941; Currie et al., 1956; Grosse et al., 1951; Kaufman and Libby, 1954; Libby, 1946) and the desire to measure its distribution in the hydrologic cycle. The advent of atmospheric thermonuclear tests in the 1950s and early 1960s dwarfed the natural global inventory (Weiss and Roether, 1980), which motivated an increase in oceanic measurements in the 1960s. The initiation of global ocean chemistry, hydrographic, and tracer survey efforts (especially GEOSECS¹⁶) further increased this activity. A final boost to tritium measurement rates occurred with the development of the ³He regrowth method (Clarke et al., 1976) coupled with even more ambitious global surveys (such as WOCE, CLIVAR and GO-SHIP).

The helium sampling time history was basically initiated and motivated by the discovery of primordial ³He injection into the deep waters (Clarke et al., 1969) which drove the inclusion of helium isotope measurements in the GEOSECS program. It was quickly realized that the existence of *tritiumogenic* ³He (that produced by the *in situ* decay of tritium) offered the potential for a dating tool as well (Jenkins et al., 1972; Jenkins and Clarke, 1976), which spurred continued helium isotope measurements in the global surveys. These were enabled by a number of laboratories coming “on-line” in the 1970s and 1980s.

The tritium and helium sampling locations shown in Figure [2-3](#) are dominated by the global survey programs cruise tracks, but also include a number of ocean island monitoring sites (especially for tritium). One difference in the two maps is the extra tritium sampling in the Arctic, perhaps largely driven by H.G. Ostlund’s motivation to exploit this isotope’s potential in studying the Arctic fresh water system (Ostlund, 1982).

It is well known that the delivery of bomb tritium to the ocean was a reflection of the distribution of the atmospheric tests, and occurred in two principle modes: a dominant pulse-like injection in the northern hemisphere and a much smaller more “diffuse” input into the southern hemisphere (Doney et al., 1992). [The former mode wasere dominated by the early 1960s American and USSR tests that occurred just prior to the 1963 Partial Test Ban Treaty, while the southern hemispheric tests were carried out predominantly by the U.K. and France. Weiss and Roether \(1980\) estimated the delivery of approximately 1500 GCi to the northern hemisphere and 480 GCi to the southern hemisphere by the end of 1972. This-The latitudinal trends](#) can be seen in

Figure [34](#), which is plot of near surface (<50 m depth) water tritium concentrations vs. time for number of latitude bands. Most

¹⁶ GEOSECS is the Geochemical Ocean Sections Survey

striking is the northward increase in the concentration (y-axis) ranges. The time-latitude trends in Figure 3-4 reflect this globally asymmetric delivery, but much of the structure caused by regional variations in atmospheric input and ocean circulation may be masked due to conflating major ocean basins in the groupings.

As a transient tracer, tritium offers an opportunity to visualize the ventilation of deep waters on decade to century time scales.

5 ~~BA~~ benchmark observations of water column tritium concentrations during the GEOSECS Atlantic Expedition reveals North Atlantic deep water formation in a graphic manner (Ostlund et al., 1974). Valuable also is the evolution of the tritium distributions as they penetrate the subtropical thermocline and also intermediate and deep waters. Figure 5 shows four “snapshots” of tritium distributions ~~along-on~~ a section along approximately 52°W in the North Atlantic between the South American and North American coasts (see map inset). Tritium concentrations are decay-corrected to a common mid-point in
10 time (1997) for comparison. While the last three occupations are conveniently along the same cruise track (courtesy WOCE/CLIVAR/GO-SHIP), the first is a composite from several cruises of opportunity taken over an approximately 1 year period at roughly the same longitude. One can readily see the downward propagation and ultimate dispersion of the bomb tritium pulse within the main thermocline (the upper 1000 m) and the progressive ingrowth of tritium at intermediate (1500-2500m depth) and in the bottom layers (~4000 m) is striking. ~~One can also see upper-water-the repartitioning of tritium inventories in the upper waters, both due to vertical exchange and ventilation, but also due to the continued accumulation of this bomb fallout isotope at subtropical latitudes~~ (similar to bomb radiocarbon, as observed by Broecker et al., 1995) ~~from more tritium-rich waters to the north.~~

Equally important ~~is the bottom contour-hugging nature of the deep and intermediate levels, with a delayed arrival in the south~~ ~~corresponding to-is evidence of~~ the southward propagation of the transient tracer along the deep western boundary current system from Nordic and Labrador Seas (Doney and Jenkins, 1994). ~~This first appears in 1982 at the northern end of the section between 3000 and 4000 m depth near the bottom of the continental slope. As time progresses, it grows in at intermediate (~1500 m) and deeper (3500 – 4000 m) depths at the southern end. This clearly marks the net advective time-scales for the deep- and intermediate-depth western boundary currents.~~

25 -
Figure 4-6 shows the corresponding helium isotope anomaly distribution for ~~these-the tritium~~ sections-. Interpretation ~~of the helium isotope ratio anomaly~~ is a little more complicated ~~than the invasion of bomb tritium~~, but the build-up of tritiogenic ³He within the main thermocline is an important diagnostic of vertical transport for the subtropical main thermocline. Its retention and back-flux to the ocean surface is a uniquely valuable transient tracer observation, one that parallels the build-up and reflux
30 of inorganic nutrients in the thermocline (e.g., nitrate and phosphate) but in a quantifiably defined manner. Observations of surface water ³He excesses (not shown here) have been used as flux gauges to quantify/constrain regional-scale new production rates (Jenkins, 1988; Jenkins and Doney, 2003; Stanley et al., 2015) as well as upwelling rates (Rhein et al., 2010).

We also include a time series (plotting only the upper 2000 dbar [in Figure 7](#)) for stations within 200km of Bermuda ([32.3 N, 64.7 W](#)) in the subtropical North Atlantic, which highlights some high temporal resolution features in the penetration of tritium into, and ingrowth of ^3He within the main thermocline and intermediate waters at one location.

- 5 Perhaps the most notable features of the oceanic distribution of [excess- \$^3\text{He}\$ the helium isotope ratio anomaly](#) are the large tongues emanating from mid-ocean ridges and other volcanic edifices on the sea floor (see [the map of helium isotope ratio anomaly at 2500 m depth in Figure 87](#)), driven by the roughly order-of-magnitude higher $^3\text{He}/^4\text{He}$ ratio in the earth's mantle compared to the atmosphere (Clarke et al., 1969; Kurz and Jenkins, 1981; Kurz et al., 1982; Lupton and Craig, 1975). It was in fact the initial discovery (Clarke et al., 1969) that sparked continued interest in the oceanic distribution of helium isotopes.
- 10 [In contrast, one sees a different pattern deeper down \(4000 m, in Figure 9\), where one sees the relatively \$^3\text{He}\$ -impoverished bottom waters flowing northward into the abyssal Pacific.](#) The ongoing survey of deep helium features on both [large-basin](#) and [small-regional](#) scales continues today, particularly in support of “flux gauge” studies of other trace elements and metals influenced by seafloor hydrothermal processes (e.g., Jenkins et al., 2018; Resing et al., 2015; Roshan et al., 2016). These are based on recent model-based estimates of the global flux of hydrothermal ^3He , which center around 550 mol/y (Bianchi et al.,
- 15 2010; Holzer et al., 2017; Schlitzer, 2016). This estimate can be usefully compared to the expected global flux of *tritogenic* ^3He . The global tritium production by atmospheric nuclear weapons tests has been estimated to be of order 3 GCi (Weiss and Roether, 1980, corrected from 1972 to 1963) to 5 GCi (Michel, 1976). Given that the bulk of the tritium “impulse” entered the hydrologic system, and subsequently the oceans within a decade or so, one can argue that the production rate of tritogenic ^3He was of order 3000 mol/y in the mid 1970s. By the mid 1990s, this would be of order 1000 mol/y. Separating the two “types”
- 20 of ^3He (tritogenic vs. primordial) in the northern hemisphere, where the bulk of the tritium delivery occurred (Doney et al., 1992; Weiss and Roether, 1980), is relatively simple in that the former appears largely at the sea surface and the latter is concentrated in older, deeper waters. The separation in the southern hemisphere is not so simple.

4 Acknowledgments

This dataset represents the hard work over many decades of numerous individuals that are not included in the authorship list of this paper. We list their names and affiliations at the time of their contributions in Table 3. The list focuses on those who made the measurements rather than those who may have used the data. We apologize if there are others that we may have missed in this list.

We also would like to recognize that the ability to make the measurements presented in this dataset was a consequence of the pioneering work of more than a few inventive and talented individuals. While space does not permit mentioning them all here,

30 we felt it appropriate to highlight a pair of pioneering scientists who conducted landmark studies on ocean tritium and ^3He measurements.

4.1 W. Brian Clarke (1937-2002)

Although not the first to measure $^3\text{He}/^4\text{He}$ in the environment (that was done by Aldrich and Nier, 1948), Brian Clarke made the first reported helium isotope measurements in seawater (Clarke et al., 1969). He made his first measurements using a modified single stage magnetic sector, single collector mass spectrometer to a precision of about 2%. Brian developed the first compact all-metal branch tube mass spectrometer specifically designed to make $^3\text{He}/^4\text{He}$ measurements ultimately to a precision of 0.1 to 0.2%. At the time, conventional wisdom dictated that such measurements (let alone *precision* measurements) were not possible with a single stage magnetic sector instrument for such high (10^6) abundance ratios, but Clarke forged ahead anyway. He initially constructed two instruments in the early 1970s, using one at McMaster University in Hamilton, Ontario, Canada, and setting the other machine up at the Scripps Institute of Oceanography in La Jolla, CA, USA for H. Craig and J. Lupton. These instruments were used in support of the GEOSECS program, and subsequently for a wide variety of other research projects. One of his students (Jenkins) moved to the Woods Hole Oceanographic Institution in Woods Hole, MA, USA, where he extended Clarke's design to construct three other instruments. A post-doctoral investigator from his laboratory (Z. Top) moved to RSMAS at the University of Miami and constructed a similar machine. Brian also collaborated with a U.K. mass spectrometer company to make a commercially produced mass spectrometer available to the global scientific community. In his early career, Brian contributed to the study of meteorites and nuclear physics using isotope mass spectrometry. Beginning in 1969 Clarke published a series of ground-breaking papers on $^3\text{He}/^4\text{He}$ measurements in seawater and lakes (Clarke et al., 1969, 1970; Clarke and Kugler, 1973; Craig and Clarke, 1970; Craig et al., 1975; Jenkins and Clarke, 1976; Top and Clarke, 1983; Torgersen and Clarke, 1985). He contributed research to geology, hydrology, limnology, nuclear physics, medicine, and other disciplines, and was active until 2002 publishing on ^3He -related evidence for/against cold fusion (Clarke, 2001; Clarke and Oliver, 2003; Clarke et al., 2001).

In addition to developing a mass spectrometer capable of measuring $^3\text{He}/^4\text{He}$ ratios to order 0.1% on sub-nanomolar gas samples, Brian also created a method to measure environmental levels of tritium (^3H) in water samples by the ^3He regrowth technique (Clarke et al., 1976), which has become the *de facto* state of the art in low level tritium measurements. In addition to being intrinsically simpler than the traditional low level method (which involved electrolytic enrichment combined with low-level proportional gas counting), this method has proved to be more precise (by more than a factor of 4) and extended the detection limit to lower levels (by as much as an order of magnitude) (Bayer et al., 1989; Jenkins et al., 1983; Lott and Jenkins, 1998).

Brian was inventive and ingenious in the laboratory with a remarkable ability to recognize opportunities where no one else could, and to pursue them to ultimate success. He was an accomplished glass blower, and constructed vacuum lines and research apparatus from scratch using many different kinds of glass (see Figure 810). Brian kept unusual work hours when not teaching; generally arriving at the laboratory after lunch-time and toiling into the night. Working with him was a delight due to his good nature and whimsical sense of Irish humor, but sometimes challenging because he was an inveterate prankster.

4.2 H. Göte Östlund (1923-2016)

H. Göte Östlund was a pioneer in U.S. ocean tracer measurements, establishing a world class, low-level counting laboratory dedicated to the measurement of tritium and radiocarbon at the University of Miami. He made distinguished contributions to Ocean, Atmosphere, and Groundwater sciences, in particular to understanding of the time scales of the transport of fluids through these systems. Östlund had a life-long devotion to the high-quality measurement of radioactive species. He received a B.S. in chemistry in 1949 and a Ph.D. in chemistry in 1958. Both were from the University of Stockholm. Between 1959 and 1961 Göte developed the electrolytic enrichment of tritium and deuterium for low-level environmental tritium measurements by gas proportional counting at the Radioactive Dating Laboratory of the Swedish Geological Survey. In the early 1960s, he came to the Rosenstiel School of the University of Miami. At the Rosenstiel School, he built a world-class tritium and radiocarbon counting laboratory that set new standards for low level counting. His laboratory also processed the samples rapidly, and he generously shared his data with colleagues. As a result, relatively routine collection and analysis of large quantities of samples in a timely manner enabled oceanographers to use the tracer data to gain new insights into the times scales of oceanographic processes. His work paved the way for the acceptance of the next generation of tracer oceanographers, those measuring tritium and helium-3 by mass spectrometry, and those measuring the chlorofluorocarbons.

He played a key role in the creation and execution of early global ocean survey programs. ~~Although his early interest focused on atmospheric transport, it quickly extended to the hydrologic cycle and the oceans.~~ He developed electrolytic enrichment techniques for low-level environmental tritium measurements by gas proportional counting (see Figure 119).

Östlund was a member of the scientific steering committee for the Geochemical Ocean Sections Study (GEOSECS), which was the first global scale survey of chemical, isotopic and radiochemical tracers in the ocean (1972-1978). He produced the first large-scale, high quality mapping of the distribution of tritium in the oceans, which opened oceanographer's eyes to the dynamic and rapid penetration of bomb-produced tracers into the deep ocean.

Göte Östlund published over 100 papers in peer-reviewed journals on a wide range of subjects. Although his early interests focused on many areas including atmospheric transport, they quickly extended to the hydrologic cycle and the ocean where he applied radioactive tracers to a spectrum of scientific problems. As a student, he participated in the discovery of the anaesthetic Xylocain. Soon after coming to Miami, he used tritium to show that evaporation from the ocean is the major ~~energyfuel~~ source for hurricanes. To collect samples he even flew into the eye of a hurricane. He and a colleague were the first to use tritium data to show that vertical mixing in the upper layers of the open ocean was an order of magnitude smaller than predicted by mass balance and theory. This was corroborated twenty years later by other investigators using new techniques. A major interest of his was the Arctic Ocean. There he quantified the contributions from ice melt, runoff and precipitation to the freshwater budget.

This budget plays a critical role in the global overturning circulation, which is the leading candidate for modulating decadal to centennial climate.

Göte was involved in the planning and implementation and served on the scientific steering committees of early global change programs: Geochemical Ocean Sections (GEOSECS), which was the first global scale survey of chemical, isotopic and

radiochemical tracers in the ocean (1972-1978), followed in the 1980s by Transient Tracers in the Oceans (TTO). The data his laboratory produced, collected under the auspices of these programs, have furthered our understanding of the time scales of ocean processes. For example, the data have been used to estimate the flux of anthropogenic carbon dioxide into the ocean, the rate of exchange between the atmosphere and ocean, and rates of deep-water formation. He produced the first large-scale, high quality mapping of the distribution of tritium in the ocean, which opened a new vista on the dynamic and rapid penetration of bomb-produced tracers into the deep western North Atlantic Ocean. Göte's leadership as a member and coordinator of the scientific advisory committee for GEOSECS and TTO and his vision and credibility in seeing that an accelerator mass spectrometry facility for ^{14}C analysis was established in the United States - had a large influence on ocean science. The big oceanographic programs of the past 50 years (GEOSECS through WOCE, CLIVAR and GO-SHIP) have provided platforms for obtaining large quantities of high quality tracer data.

Göte was soft-spoken and gentle in demeanor, and generous with his time, advice, and data. He set an example and benchmark for subsequent generations of tracer geochemists for responsibility, honesty, and fairness.

4.3 Financial Support

This synthesis work was funded under the auspices of a U.S. National Science Foundation grant number OCE-1434000. Financial support for the actual measurements came from a wide variety of different research grants from many agencies in many countries, far too numerous to list here. The first author (Jenkins) is grateful to a number of U.S. funding sources, most notably the National Science Foundation, NOAA, DOE, and ONR.

5 References

- Aldrich, L. T. and Nier, A. O.: The occurrence of ^3He in natural sources of helium, *Physical Review*, 74, 1590-1594, 1948.
- Bainbridge, A. E., Ostlund, H. G., Craig, H., Broecker, W. S., and Spencer, D. W.: GEOSECS Atlantic, Pacific, and Indian Ocean Expeditions: Shore-based data and graphics, GEOSECS ATLAS, 1987.
- Bainbridge, A. E., Sandoval, P., and Suess, H. E.: Natural tritium measurements by ethane counting, *Science*, 134, 552-553, 1961.
- Bayer, R., Schlosser, P., Bonisch, G., Rupp, H., Zaucker, F., and Zimmek, G.: Performance and blank components of a mass spectrometric system for routine measurement of helium isotopes and tritium by ^3He ingrowth method, *Sitzungsberichte der Heidelberger Akademie der Wissenschaften Mathematisch-naturwissenschaftliche Klasse*, 5, 241-279, 1989.
- Bianchi, D., Sarmiento, J. L., Gnanadesikan, A., Key, R. M., Schlosser, P., and Newton, R.: Low helium flux from the mantle inferred from simulations of oceanic helium isotope data, *Earth and Planetary Science Letters*, 297, 379-386, 2010.

- Broecker, W. S., Sutherland, S. C., and Smethie, W. M.: Oceanic radiocarbon: separation of the natural and bomb components, *Global Biogeochemical Cycles*, 9, 263-288, 1995.
- Brown, K., Dingley, K. H., and Turteltaub, K. W.: Accelerator mass spectrometry for biomedical research, *Methods in Enzymology*, 402, 423-443, 2005.
- 5 Cameron, J. F.: Radioactive dating and methods of low-level counting, Monaco, 2-10 March 1967 1967, 543-574, 1967.
- Chiarappa-Zucca, M. L., Dingley, K. H., Roberts, M. L., Velsko, C. A., and Love, A. H.: Sample preparation for quantification of tritium by acclerator mass spectrometry, *Analytical Chemistry*, 74, 6285-6290, 2002.
- Clarke, W. B.: Search for ^3He and ^4He in Arata-Style Palladium Cathods I: a negative result, *Fusion Science and Technology*, 40, 147, 2001.
- 10 Clarke, W. B., Beg, M. A., and Craig, H.: Excess ^3He in the sea: evidence for terrestrial primordial helium, *Earth and Planetary Science Letters*, 6, 213-220, 1969.
- Clarke, W. B., Beg, M. A., and Craig, H.: Excess Helium 3 at the North Pacific GEOSECS station, *Journal of Geophysical Research*, 75, 7676-7678, 1970.
- 15 Clarke, W. B., Jenkins, W. J., and Top, Z.: Determination of tritium by spectrometric measurement of ^3He ., *International Journal of Applied Radiation and Isotopes*, 27, 515-525, 1976.
- Clarke, W. B. and Kugler, G.: Dissolved helium in groundwater: a possible method for uranium and thorium prospecting, *Economic Geology*, 68, 243-251, 1973.
- Clarke, W. B. and Oliver, B. M.: Response to "Comments on 'Search for ^3He and ^4He in Arata-style palladium cathodes I: a negative result" and "Search for ^3He and ^4He in Arata-style palladium cathodes II: evidence for tritium production", *Fusion Science and Technology*, 43, 135-136, 2003.
- 20 Clarke, W. B., Oliver, B. M., McKubre, M. C. H., Tanzella, F. L., and Tripodi, P.: Search for ^3He and ^4He in Arata-Style Palladium Cathodes II: Evidence of tritium production, *Fusion Science and Technology*, 40, 152, 2001.
- Cornog, R. and Libby, W. F.: Production of radioactive hydrogen by neutron bombardment of boron and nitrogen, *Physical Review*, 59, 1046, 1941.
- 25 Craig, H. and Clarke, W. B.: Oceanic ^3He : contribution from cosmogenic tritium, *Earth and Planetary Science Letters*, 9, 45-48, 1970.
- Craig, H., Clarke, W. B., and Beg, M. A.: Excess ^3He in deep water on the East Pacific Rise, *Earth and Planetary Science Letters*, 26, 125-132, 1975.
- 30 Currie, L. A., Libby, W. F., and Wolfgang, R. L.: Tritium production by high energy protons, *Physical Review*, 101, 1557-1563, 1956.
- Doney, S. C., Glover, D. M., and Jenkins, W. J.: A model function of the global bomb-tritium distribution in precipitation, 1960-1986, *Journal of Geophysical Research*, 97, 5481-5492, 1992.
- Doney, S. C. and Jenkins, W. J.: Ventilation of the deep western boundary current and abyssal Western North Atlantic: Estimates from tritium and ^3He distributions, *Journal of Physical Oceanography*, 24, 638-659, 1994.
- 35 Dorsey, H. G. and Peterson, W. H.: Tritium in the Arctic Ocean and East Greenland Current, *Earth and Planetary Science Letters*, 32, 342-350, 1976.
- Dreisigacker, E. and Roether, W.: Tritium and ^{90}Sr in North Atlantic surface water, *Earth and Planetary Science Letters*, 38, 301-312, 1978.
- 40 Fine, R. A. and Ostlund, H. G.: Source function for tritium transport models in the Pacific, *Geophysical Research Letters*, 4, 461-464, 1977.
- Fine, R. A., Peterson, W. H., and Ostlund, H. G.: The penetration of tritium into the tropical Pacific, *Journal of Physical Oceanography*, 17, 553-564, 1987.

- Fine, R. A., Reid, J. L., and Ostlund, H. G.: Circulation of tritium in the Pacific Ocean, *Journal of Physical Oceanography*, 11, 3-14, 1981.
- Fuchs, G., Roether, W., and Schlosser, P.: Excess ^3He in the ocean surface layer, *Journal of Geophysical Research*, 92, 6559-6568, 1987.
- 5 German, C. R., Casciotti, K. L., Dutay, J.-C., Heimburger, L. E., Jenkins, W. J., Measures, C., Mills, R. A., Obata, H., Schlitzer, R., Tagliabue, A., Turner, D. R., and Whitby, H.: Hydrothermal impacts on trace element and isotope ocean biogeochemistry, *Philos T Roy Soc A*, 374, 20130035, 2016.
- Glagola, B. G., Phillips, G. W., Marlow, K. W., Myers, L. T., and Omohundro, R. J.: Low level tritium detection using accelerator mass spectrometry, *Nuclear Instruments and Methods in Physics Research B*, 5, 221-225, 10 1984.
- Grosse, A. V., Johnston, W. M., Wolfgang, R. L., and Libby, W. F.: Tritium in nature, *Science*, 113, 1-2, 1951.
- Holzer, M., DeVries, T., Bianchi, D., Newton, R., Schlosser, P., and Winckler, G.: Objective estimates of mantle ^3He in the ocean and implications for constraining the deep ocean circulation, *Earth and Planetary Science Letters*, 458, 305-314, 2017.
- 15 Israel, G. W.: Messung des Tritium-Jahresganges im Regen 1960-1961 nach Isotopenanreicherung im Trnnrohr, *Z. Naturforsch.*, 17a, 925-929, 1962.
- Jenkins, W. J.: ^3H and ^3He in the Beta Triangle: Observations of gyre ventilation and oxygen utilization rates, *Journal of Physical Oceanography*, 17, 763-783, 1987.
- Jenkins, W. J.: Nitrate flux into the euphotic zone near Bermuda, *Nature*, 331, 521-523, 1988.
- 20 Jenkins, W. J.: Studying Thermocline Ventilation and Circulation Using Tritium and ^3He , *Journal of Geophysical Research*, 103, 15817-15831, 1998.
- Jenkins, W. J.: Tritium-helium dating in the Sargasso Sea: a measurement of oxygen utilization rates., *Science*, 196, 291-292, 1977.
- Jenkins, W. J., Beg, M. A., Clarke, W. B., Wangersky, P. J., and Craig, H.: Excess ^3He in the Atlantic Ocean., *Earth and Planetary Science Letters*, 16, 122-130, 1972.
- 25 Jenkins, W. J. and Clarke, W. B.: The distribution of ^3He in the western Atlantic Ocean, *Deep-Sea Research*, 23, 481-494, 1976.
- Jenkins, W. J. and Doney, S. C.: The Subtropical Nutrient Spiral, *Global Biogeochemical Cycles*, 17, 1110 doi:1110.1029/2003GB002085, 2003.
- 30 Jenkins, W. J., Edmond, J. M., and Corliss, J. B.: Excess ^3He and ^4He in Galapagos submarine hydrothermal waters, *Nature*, 272, 156-158, 1978.
- Jenkins, W. J., Lott, D. E., Pratt, M. W., and Boudreau, R. D.: Anthropogenic tritium in South Atlantic bottom water, *Nature*, 305, 45-46, 1983.
- Jenkins, W. J., Lott, D. E. I., German, C. R., Cahill, K. L., Goudreau, J., and Longworth, B. E.: The deep distributions of helium isotopes, radiocarbon, and noble gases along the U.S. GEOTRACES East Pacific zonal transect (GP16), *Marine Chemistry*, 201, 167-182, 2018.
- 35 Jenkins, W. J. and Rhines, P. B.: Tritium in the deep North Atlantic Ocean, *Nature*, 286, 877-880, 1980.
- Jenkins, W. J. and Smethie, W. M.: Transient tracers track ocean climate signals, *Oceanus*, 39, 29-32, 1996.
- Kaufman, S. and Libby, W. F.: The natural distribution of tritium, *Physical Review*, 93, 1337-1344, 1954.
- 40 Kurz, M. D. and Jenkins, W. J.: The distribution of helium in oceanic basalt glasses, *Earth and Planetary Science Letters*, 53, 41-54, 1981.
- Kurz, M. D., Jenkins, W. J., and Hart, S. R.: Helium isotopic systematics of oceanic islands and mantle heterogeneity, *Nature*, 297, 43-47, 1982.

- Libby, W. F.: Atmospheric helium three and radiocarbon from cosmic radiation, *Physical Review*, 59, 671-672, 1946.
- Lott, D. E.: Improvements in noble gas separation methodology: a nude cryogenic trap, *Geochemistry, Geophysics, Geosystems*, 2, 10.129/2001GC000202, 2001.
- 5 Lott, D. E. and Jenkins, W. J.: Advances in the analysis and shipboard processing of tritium and helium samples, *International WOCE Newsletter*, 30, 27-30, 1998.
- Lott, D. E. and Jenkins, W. J.: An automated cryogenic charcoal trap system for helium isotope mass spectrometry, *Review of Scientific Instruments*, 55, 1982-1988, 1984.
- 10 Ludin, A., Weppernig, R., Bonisch, G., and Schlosser, P.: Mass spectrometric measurement of helium isotopes and tritium in water samples, 1997. 1997.
- Lupton, J. E. and Craig, H.: Excess He-3 in oceanic basalts: evidence for terrestrial primordial helium, *Earth and Planetary Science Letters*, 26, 133-139, 1975.
- Lupton, J. E. and Jenkins, W. J.: Evolution of the South Pacific helium plume over the past 3 decades, *Geochemistry Geophysics Geosystems*, 18, 1810-1823, 2017.
- 15 Mamyrin, B. A.: Time-of-flight mass spectrometry (concepts, achievements and prospects), *International Journal of Mass Spectrometry*, 206, 251-266, 2001.
- Mamyrin, B. A., Anufriyev, S. G., Kamenskiy, I. L., and Tolstikhin, I. I.: Determination of the isotopic composition of atmospheric helium, *Geochemistry International*, 6, 498-505, 1970.
- Michel, R. L.: Tritium inventories of the world oceans and their implications, *Nature*, 263, 103-106, 1976.
- 20 Michel, R. L. and Suess, H. E.: Bomb tritium in the Pacific Ocean, *Journal of Geophysical Research*, 80, 4139-4152, 1975.
- Miyake, Y., Shimada, T., Sugimura, Y., Shigehara, K., and Saruhashi, K.: Distribution of tritium in the Pacific Ocean, *Records of Oceanographic Works in Japan*, 13, 17-32, 1975.
- Momoshima, N., Nakamura, Y., and Takashima, Y.: VIAL EFFECT AND BACKGROUND SUBTRACTION METHOD IN
25 LOW-LEVEL TRITIUM MEASUREMENT BY LIQUID SCINTILLATION-COUNTER, *International Journal of Applied Radiation and Isotopes*, 34, 1623-1626, 1983.
- Ostlund, H. G.: The residence time of the freshwater component in the Arctic Ocean, *Journal of Geophysical Research*, 87, 2035-2043, 1982.
- 30 Ostlund, H. G., Dorsey, H. G., and Rooth, C. G.: GEOSECS North Atlantic Radiocarbon and Tritium Results, *Earth and Planetary Science Letters*, 23, 69-86, 1974.
- Ostlund, H. G. and Werner, E.: The electrolytic enrichment of tritium and deuterium for natural tritium measurements., *Vienna*, 95 pp., 1962.
- Resing, J. A., Sedwick, P. N., German, C. R., Jenkins, W. J., Moffett, J. W., Sohst, B. M., and Tagliabue, A.: Basin-Scale transport of hydrothermal dissolved metals across the South Pacific Ocean, *Nature*, 523, 203-206, 2015.
- 35 Rhein, M., Dengler, M., Sultenfuss, J., Hummels, R., Huttler-Kabus, S., and Bourles, B.: Upwelling and associated heat flux in the equatorial Atlantic inferred from helium isotope disequilibrium, *J. Geophys. Res.-Oceans*, 115, doi:10.1029/2009JC005772, 2010.
- Roberts, M. L., Hamme, R. W., Dingley, K. H., Chiarappa-Zucca, M. L., and Love, A. H.: A compact tritium AMS system, *Nuclear Instruments and Methods in Physics Research B*, 172, 262-267, 2000.
- 40 Roether, W., Vogt, M., Vogel, S., and Sültenfuß, J.: Combined sample collection and gas extraction for the measurement of helium isotopes and neon in natural waters, *Deep Sea Research Part I: Oceanographic Research Papers*, 76, 27-34, 2013.

- Roether, W., Well, R., Putzka, A., and Ruth, C.: Component separation of oceanic helium, *Journal of Geophysical Research*, 103, 27931-27946, 1998.
- Roshan, S., Wu, J., and Jenkins, W. J.: Long-range transport of hydrothermal dissolved Zn in the tropical South Pacific, *Marine Chemistry*, 183, 25-32, 2016.
- 5 Sarmiento, J. L.: A simulation of bomb tritium entry into the Atlantic Ocean, *Journal of Physical Oceanography*, 13, 1924-1939, 1983.
- Schlitzer, R.: Quantifying He fluxes from the mantle using multi-tracer data assimilation, *Philos T Roy Soc A*, 374, DOI: 10.1098/rsta.2015.0288, 2016.
- 10 Stanley, R. H. R., Jenkins, W. J., Doney, S. C., and Lott, D., III: The ^3He flux gauge in the Sargasso Sea: a determination of physical nutrient fluxes to the euphotic zone at the Bermuda Atlantic time series site, *Biogeosciences* 12, 5199-5210, 2015.
- Top, Z. and Clarke, W. B.: Helium, neon, and tritium in the Black Sea, *Journal of Marine Research*, 41, 1-17, 1983.
- Torgersen, T. and Clarke, W. B.: Helium accumulation in groundwater, 1: an evaluation of sources and the continental flux of crustal He-4 in the Great Artesian Basin, Australia, *Geochimica et Cosmochimica Acta*, 49, 15 1211-1218, 1985.
- Unterweger, M. P., Coursey, B. M., Schima, F. J., and Mann, W. B.: Preparation and calibration of the 1978 National Bureau of Standards tritiated-water standards, *International Journal of Applied Radiation and Isotopes*, 31, 611-614, 1980.
- 20 Weiss, R. F.: Piggyback sampler for dissolved gas studies on sealed water samples, *Deep-Sea Research*, 15, 695-699, 1968.
- Weiss, W. M. and Roether, W.: The rates of tritium input to the world oceans, *Earth and Planetary Science Letters*, 49, 435-446, 1980.
- Weiss, W. M., Roether, W., and Dreisigacker, E.: Tritium in the North Atlantic: inventory, input and transfer to the deep water. In: *The Behavior of tritium in the Environment*, International Atomic Energy Agency, Vienna, 25 1979.
- Young, C. and Lupton, J. E.: An ultratight fluid sampling system using cold-welded copper tubing., *EOS Transactions AGU*, 64, 735, 1983.

6 Tables

- 30 **Table 1:** Fields (columns) in the main data table

Field Name	Field Type	Field Description
ExpoCode	Short Text	Unique string identifying cruise/expedition
Sect_ID	Short Text	String identifying Ocean Section (WOCE/CLIVAR/GEOTRACES)
Station	Short Text	Station name or number
CastNo	Short Text	Cast name or number at that Station
Bottle	Short Text	Bottle name or number on that cast
StaDate	Number	Decimal year of sampling

Latitude	Number	North latitude in decimal degrees (from -90 to +90)
Longitude	Number	East longitude in decimal degrees (from -180 to +180)
StaDepth	Number	Bottom depth at station location in meters
Pressure	Number	Bottle depth (actually pressure) measured in dbar
Temperature	Number	In Situ temperature in degrees centigrade
Temperature_Flag	Integer	Temperature quality flag (see QF table)
Salinity	Number	Sample salinity in PSU
Salinity_Flag	Integer	Salinity quality flag (see QF table)
Oxygen	Number	Dissolved oxygen in umol/kg
Oxygen_Flag	Integer	Dissolved oxygen flag (see QF table)
Tritium	Number	Tritium in TU at time of sampling
Tritium_Error	Number	Uncertainty in TU at time of sampling
Tritium_Flag	Integer	Tritium quality flag (see QF table)
Tritium_PI	Short Text	Principle investigator or measurer of tritium
Tritium_PI_Inst	Short Text	Institution or laboratory where tritium was measured
Tritium_Method	Short Text	Short descriptor of tritium sampling/analysis method
DelHe3	Number	Helium isotope ratio anomaly relative to atmosphere in percent
DelHe3_Error	Number	Uncertainty in helium isotope ratio anomaly
DelHe3_Flag	Integer	Helium isotope ratio anomaly quality flag (see QF table)
Helium	Number	Dissolved helium concentration in nmol/kg
Helium_Error	Number	Uncertainty in dissolved helium concentration in nmol/kg
Helium_Flag	Integer	Dissolved Helium quality flag (see QF table)
Neon	Number	Dissolved neon concentration in nmol/kg
Neon_Error	Number	Uncertainty in dissolved neon concentration in nmol/kg
Neon_Flag	Integer	Dissolved neon quality flag (see QF table)
Helium_PI	Short Text	Principle investigator or measurer of helium (and neon)
Helium_PI_Inst	Short Text	Institution or laboratory where tritium was measured
Helium_Method	Short Text	Short descriptor of helium sampling/analysis method
Reference_Code	Short Text	Data origin or link to paper discussing data
Reference_Source	Short Text	Data Source within reference (e.g., table, figure) if relevant
DOI	Short Text	Digital Object Identifier of original data set (if existing)
Comment	Short text	Additional information or comments

Record_ID	Long Integer	Unique record identifier number
-----------	--------------	---------------------------------

Table 2: Quality flag meaning

Quality Flag Number	Meaning
2	Normal data, no problems reported
3	Questionable data: may not fit profile or some other doubt
6	Average of 2 or more measurements
9	Missing (null) data

5 **Table 3:** Contributing analysts that are not authors on this paper

A.E. Bainbridge	UCSD, La Jolla, CA, USA
R. Bayer	U. Heidelberg, Heidelberg, Germany
F. Begemann	U. Chicago, Chicago, IL, USA
U. Beyerle	ETH, Zurich, Switzerland
W.S. Broecker	LDEO, Pallsades, NY, USA
M. Butzin	University of Bremen, Bremen, Germany
W.B. Clarke*	McMaster University, Hamilton, ON, Canada
K.O. Dockins	UCSD, La Jolla, CA, USA
H.G. Dorsey	RSMAS, Miami, FL, USA
E. Eriksson	IMS, Stockholm, Sweden
E. Fourré	CEA-Saclay, France
B.J. Giletti	LDEO, Pallsades, NY, USA
A.V. Grosse	RITU, Philadelphia, PA, USA
J.R. Harries	Australian AEC, Sutherland, NSW, Australia
T. Kaji	Kyushu University, Fukuoka, Japan
S. Kaufman	U. Chicago, Chicago, IL, USA
J.L. Kulp	LDEO, Pallsades, NY, USA
W.F. Libby	U. Chicago, Chicago, IL, USA
D.E. Lott III	WHOI, Woods Hole, MA, USA
A.Luden	LDEO, Pallsades, NY, USA
L. Merlivat	Sorbonne University, Paris, France

R. Michel	UCSD, La Jolla, CA, USA
Y. Miyake	GRA, Tokyo, Japan
K.O. Munnich	U. Heidelberg, Heidelberg, Germany
A.O. Nier	U. Minnesota, Minneapolis, MN, USA
M. Nonaka	IPRC & SOES, Tokyo, Japan
H.G. Ostlund*	RSMAS, Miami, FL, USA
C. Postlethwaite	NOC-SOES, Southampton, U.K.
P.D. Quay	University of Washington, Seattle, WA, USA
R.S.H.R. Stanley	WHOI, Woods Hole, MA, USA
S. Stark	NOC-SOES, Southampton, U.K.
R. Steinfeldt	IUP, University Bremen, Germany
H.E. Suess	UCSD, La Jolla, CA, USA
J. Sültenfuss	IUP, University Bremen, Germany
N. Takahata	ORI, University of Tokyo, Tokyo, Japan
A. Tamuly	University of Quebec, Rimouski, PQ, Canada
C.B. Taylor	INS, Lower Hutt, New Zealand
Z. Top	RSMAS, Miami, FL, USA
T. Torgersen	WHOI, Woods Hole, MA, USA
K.A. Van Scoy	RSMAS, Miami, FL, USA
C. Walker	WHOI, Woods Hole, MA, USA
W. Weiss	U. Heidelberg, Heidelberg, Germany
P.M. Williams	UCSD, La Jolla, CA, USA

7 Figures




Figure 1: A scatter plot of helium and neon saturation anomalies for nearly 2000 near-surface (< 20 m depth) samples from the data set. The solid line is a type-II linear regression of the data.

Commented [SCD1]: Bill, it looks like you deleted the figure; I added it back





Figure 21: Time distributions of annual tritium (upper panel) and helium (lower panel) measurements.

Tritium Sampling Locations



Helium Sampling Locations




Figure 33: Tritium and helium sample locations




Figure 43: Global ocean surface water (depth < 50 m) tritium concentrations (in TU) for selected latitude bands.




Figure 54: Four meridional tritium sections along roughly 52°W in the North Atlantic taken in 1982, 1997, 2003, and 2012. The tritium concentrations have been decay corrected to a common time (January 1, 1997) for comparison. Contour intervals are 0.2 TU₉₇ and measurement uncertainties are of order 0.01 TU₉₇ or better. Due to differences in cruise tracks, the topography for the 1982 occupation differs from the others.




Figure 65: Four meridional $\delta^3\text{He}$ sections along roughly 52°W in the North Atlantic taken in 1982, 1997, 2003, and 2012. Contour intervals are 1% and measurement uncertainties are 0.15%. Due to differences in cruise tracks, the topography for the 1982 occupation differs from the others.




Figure 76: A time series of tritium (upper) and helium isotope measurements in the vicinity of Bermuda (North Atlantic). Tritium values have been decay-corrected to a common time (January 1, 1997). White dots indicate sampling depths and times.




Figure 87: A map of $\delta^3\text{He}$ values at approximately 2500 m depth. The values plotted are simply an average of all measurements within a 1° square between 2250 and 2750 dbar. Depths shallower than 2500 m are masked in gray and sampling locations indicated by light gray dots. (Perhaps Reiner or I may do something better here, such as interpolate onto a neutral density surface.)

5

Formatted: Normal




Figure 9: A map of $\delta^3\text{He}$ values at approximately 4000 m depth. The values plotted are simply an average of all measurements within a 1° square between 3750 and 4250 dbar. Depths shallower than 4000 m are masked in gray and sampling locations indicated by light gray dots.




Figure 108: W. Brian Clarke, working on a high vacuum helium extraction apparatus (early 1970s)

Formatted: Normal



Figure [119](#): H. Göte Östlund preparing a gas sample for low level counting analysis (mid 1960s).

Response to reviewers for “A comprehensive global oceanic dataset of helium isotope and tritium measurements” by William J. Jenkins, Scott C. Doney , Michaela Fendrock, Rana Fine, Toshitaka Gamo, Philippe Jean-Baptiste, Robert Key, Birgit Klein, John E. Lupton1, Robert Newton, Monika Rhein, Wolfgang Roether, Yuji Sano, Reiner Schlitzer, Peter Schlosser, Jim Swift

Anonymous Referee #1

The authors have organized all existing oceanic measurements of helium-isotope and tritium data into a single comprehensive data set, complete with uncertainty estimates and valuable metadata on data quality and measurement methods. This tracer data is invaluable to oceanography and collecting it all in a single place is a significant service to the scientific community. This short paper accompanying the data set is appropriate and useful. I found the descriptions of the analytical methods useful and I enjoyed the tributes to pioneers Clarke and Ostlund. Subject to minor revisions, I recommend publication.

Thank you for the kind comments.

My main comment is that with 22% of the data being neon measurements, there should be at least a paragraph discussing the use of the neon data with some references, and perhaps a figure on neon. I assume neon, having only stable natural isotopes, is used to constrain air-sea gas exchange, but I don't know of all its uses in oceanography and suspect I'm not alone in this. Some discussion on neon would be useful. Perhaps a graphic on observed neon surface saturation could be shown.

The Reviewer makes a very good point, we had overlooked spelling out the relative importance of the He and Ne concentration measurements. We added a paragraph in section 2.3 explaining the value of both measurements, and presented a composite scatter plot of their respective supersaturations in global surface waters.

Minor comments just concern occasional clarity issues and typographical/grammatical errors:

P1 L19 typo: : : : with a half-life of 12.3 years : : : “half-life of” is missing

Thank you for catching that. This has been corrected.

P2 L24 typo: either measurement “of” or “measurement” should be “measuring”

Thank you for catching that. This has been corrected.

P2 L27 water samples -> gas samples ? (electrolysis enriches the gas, not the water)

Thank you for catching that. This has been corrected.

P3 L13 “magnetic sector instrument” please explain what this means on first usage (e.g., is it different from a mass spectrometer deflecting ions as they traverse a magnetic field?)

That is correct. We have added a brief phrase to make it clearer.

P3 L22: could the ionization state on 4He and 20Ne in the iron currents be indicated here?

5 This has now been done.

P3 L22 “helium/neon” -> “helium or neon” otherwise this is suggesting that the helium/neon ratio is somehow involved, which presumably is not the intent.

10 Good point. Now corrected.

P4 L2 typo: additional -> addition

Now corrected.

15 P4 L13: “The dataset : : :.” On a first reading, I immediately wanted to know what the file format was – it might be nice to add a parenthetical “(digital formats are specified below)”.

Phrase added.

20 P4 L14 “in principle” seems inappropriate here. Perhaps you mean: “Where available, this provides : : :”

Rephrased.

25 P4 L15 “methods fields” – delete “methods”

Deleted

P4 L18 typo: “in the Table 1” -> “in Table 1”

30 Corrected.

Section 2.5 (Data formats and availability) Why is data in netcdf format not included? Arguably this is the standard for portable data in the geosciences.

35 We now provide a netcdf version of the data.

Section 3 “Graphics and examples” seems like a very poor heading for a section because it is so generic and non-descriptive. How about “Scope and nature of the dataset” instead?

40 This is a good suggestion. We have changed the section heading accordingly.

P5 L4 “: : : for maximum flexibility. For maximum flexibility, we : : :” Delete the first “for maximum flexibility”

45 Phrase deleted.

P5 L10 “example graphics” delete “example”

Deleted.

50

P5 L22 delete “basically” or make precise qualifying statement

Word deleted.

5 P6 L5 There were also a number atmospheric, as well as submarine, nuclear tests in the Southern Hemisphere by the British and French militaries. Please add a few sentences of discussion on this and perhaps point out if this is discernible in the available oceanographic data (some of the spikes in the southern bands?) What was the estimated tritium production by nuclear testing in each hemisphere?

10 We added some additional discussion, but please understand that we have tried to strike a reasonable balance by describing the general features of the data set without digressing into too much detail regarding the causal forces behind it. The latter would be the subject of a much larger study. We also urge some caution regarding the interpretation of “spikes” in the latitude bands, particularly early in the records, because they may well reflect local (anomalous) fallout extremes rather than excursions in regional inventories. They are, after all, individual
15 measurements.

L11 P6 there is a grammar issue: “A benchmark observations” should this be singular? – please check sentence and fix.

20 This has been rectified.

L13 P6 Figure 5 should be Figure 4.

Figure numbering has been updated.

25

L14 P6 “along a section along” -> “along a section at”

Redundancy fixed.

30 L20 P6 “Equally important is the bottom-contour-hugging : :” This sentence is very unclear. What does it mean for a “level” to hug a “bottom contour”? It is unclear what in the figure is being described – please reword for clarity.

This has been re-worded to be clearer.

35 L23 P6 Figure 4 should be Figure 5.

Figure numbers corrected.

Also “those” sections – obscure – same as in Figure 4 – please restate.

40

Done.

L30 P6 “We also include a : :” You need to explicitly reference “Figure 6”

45 Done.

L1 P7 “excess 3He” what does “excess” mean here? Do you mean elevated relative to the atmospheric 3He/4He ratio or excess over global mean or something else? Please clarify and perhaps just refer to delta3He as that is precisely defined.

50

We agree. We now revert to the helium isotope ratio anomaly, for which there is a clear definition.

P9 L9-10 and L16-17 "Although his early : : and the ocean" This is a repeat of the exact same sentence – should probably have been deleted from L9-10

5

Redundant sentence deleted.

P9 L19 "fuel" seems a bit like the wrong word here. "Energy source" ?

10

Thank you, "energy source" is a better term.

Figure 4: why is the decay-corrected tritium in the upper ocean higher in 2012 than in 2003? This seems like a prominent feature of this figure that should be briefly discussed in the text.

15

We have added a brief explanation.

Figure 6: "vicinity" what is the lat and lon?

20

Added.

Figure 7: "Perhaps Reiner and I : : ." Please remove internal commentary among coauthors (i.e., please proof-read your paper and captions : :.)

25

Sorry. Comment removed.

Anonymous Referee #2

Jenkins et al. (ESSD-2018-136) introduces a dataset of oceanic helium and tritium measurements, which is open to the world via www with DOI. The dataset is definitely valuable for any earth scientists, especially oceanographers. Analytical methods used are valid. Various formats of the dataset provided are helpful. I have nothing to say without some tiny/minor points before its publication on ESSD.

P1L15: Add "." between depth and When

Period added.

P1L19: with a half-life of 12.3y?

Corrected.

P2L2-5: Numerous refs are nice but their generations seem biased in 20th century. Please introduce nice papers in 21st century if possible.

These references are largely weighted to the 20th century because the work involved was largely published there. We don't want to create an extensive bibliography of all work done in this area as it would detract from the main purpose of the paper.

P2L12-14: In turn, generations of refs here seem biased in 21st century. Please introduce classic/pioneering papers if possible.

We included only the most recent determinations of the global 3He flux because they are by far the most accurate. Those papers refer to the earlier, foundational, but less accurate determinations.

P2L16: Please add refs for each of these programs.

We have added the references.

P3L17: How deep and old? Please introduce approximate criteria.

The second part of the sentence actually provides the criteria. Rather than specify a specific depth, which would not apply in some areas than others, we give an example.

P3L23: 4He and 20Ne are NOT ion currents.

We have corrected this (see Reviewer 1 comments).

P4L6: Please add density into the dataset because of its importance in oceanography.

Density is a derived and calculable property given the sample's pressure, temperature, and salinity. In addition, it requires a reference pressure (i.e., does one refer to the sea surface, some fixed level – e.g., 3000 dbar – or do you use neutral density?).

P5L2: I request the dataset distributed by another (non-US) web server for more robust accessibility if possible.

We have now supplied an EU-based mirror site (referred to in the manuscript) at <http://odv.awi.de/data/ocean/jenkins-tritium-helium-data-compilation/>

P6L13: figure number?

Corrected.

P6L23: figure number?

Corrected.

P6L30: Figure 6?

Corrected.

P7L5: How large and small?

Descriptors added.

P7L18: I cannot understand why numerous hard workers in Table 3 are not included in the authorship list. I know analyses of helium and tritium in seawater (also sampling of oceanic water) are definitely hard. I think the list should include the hard workers.

The reviewers request a formidable task: we are well aware that the hard work and efforts made by many people in making these measurements. The authorship of this article reflects the contributions of the individuals assembling this dataset, not the original generation of the data. We have made considerable effort to provide within the dataset the correct attribution for the data originators and we hope this adequate recognition of their contributions.

Figure 7: Another map for 3000m (or 3500m) seems nice to figure out the impact from hydrothermal activity on the ridges represented in the current 2500m map.

We have included a contrasting map at the 4000m horizon as well, which we think better emphasizes the differences.