

# Authors' Response to the Reviewers' Comments (PAPER: The ISC Bulletin as a comprehensive source of earthquake source mechanisms, <https://doi.org/10.5194/essd-2018-143>)

K. Lentas, D. Di Giacomo, J. Harris, D. A. Storchak

April 3, 2019

We have taken the Reviewers' comments into account in the preparation of this revised version of the paper, which contains some text changes and additions, to fully address their requests. Moreover, we extended the data time period covered in this manuscript until the end of the data year 2018, as opposed to October 2018 when the manuscript was first submitted. The data added until the end of 2018 does not change the content and results discussed. Here follows our response to the Reviewers' comments and an explanation of the points that we have changed. Attached to our response is the revised version of manuscript with changes tracked in blue for the text addition/changes and in red for the text deleted.

Throughout this document the Reviewers' Comments (RC) are reproduced in bold.

## **1 Reviewer 1**

### **1.1 "ISC computed source mechanisms"**

Reviewer 1 expressed the query on why the ISC computed focal mechanism solutions are not discussed in the paper even though they are briefly mentioned in the abstract. The Reviewer claimed that it does not make sense to omit a part of the source mechanism solutions that can be found in the ISC Bulletin since the scope of the paper is to discuss the ISC as a comprehensive source of information on earthquake source mechanisms.

We appreciated the Reviewer's comment and have now included the ISC computed solutions in our analysis throughout the paper. Text has been added in Section 2 (page 3, lines 3-9) and in Section 5 (page 8, lines 27-29) and all the figures in the revised manuscript have been rebuilt in order to include the ISC computed focal mechanism solutions. The additional data did not change the main points of this paper.

## 1.2 “ISC and other global sources”

Reviewer 1 suggested to present more information on the reported mechanism solutions by each agency in the ISC Bulletin in terms of geographical distribution, magnitude range, number of reported mechanisms, time distribution and provided information. The Reviewer proposed to do so at least for the agencies that report solutions more systematically, covering a long period of time in order to point out differences and complementarity.

Information regarding the number of reported solutions, type of methodology being applied (where available) and time distribution has already been presented in Figure 2. In order to address the rest features in the Reviewer’s comment we added Table 1 where we present the magnitude range covered by each agency and the type of parametric data that is being reported by each agency. Moreover, we added Figure 4 which shows the geographical and magnitude distributions covered by the most systematically reporting agencies hoping to help the users of the ISC Bulletin to identify areas where local/regional agencies offer complementary information to global agencies. Text is also added in Section 2 (page 3, lines 11-25) to give some examples in specific areas on how local agencies cover gaps in magnitude range left by global agencies. Details on the methodologies being applied by different agencies are also presented in Section 3 (page 4, lines 25-35, and page 5, lines 1-15). Implications of combining solutions determined by different agencies using different techniques are discussed in Section 5 (see for example page 8, lines 30-36, and page 9, lines 4-10).

## 1.3 “Section 3 restructuring”

Reviewer 1 had objections regarding the structure of Section 3 entitled “Source mechanism variability” and proposed to restructure it and further discuss some parts of it. Specifically, the Reviewer suggested to start by discussing the point source approximation as a concept, discuss the different types of point source models (focal mechanisms, moment tensors) and finally the data being used in different methodologies (polarities, amplitude ratios, waveform modelling) and differences among moment tensors and centroid moment tensors, and centroid locations and hypocentre locations.

Moreover, the Reviewer suggested to further discuss a couple of examples briefly mentioned in the originally submitted manuscript (page 4, lines 8-12 of the original manuscript) and suggest if possible how to select a reference source mechanism for earthquakes with multiple solutions. Then, to discuss further the reasons why some earthquakes show inconsistency between the reported mechanisms and finally, to give more information on how the fault styles in Figure 5 (in the original manuscript, Figure 6 in the revised manuscript) are obtained.

Following the Reviewer’s comments, we have rewritten (almost entirely) Section 3 in order to comply with the suggestions mentioned above. Specifically, we added two new paragraphs in the beginning of Section 3 (page 4, lines 1-18) to address the Reviewer’s comments on the point source approximation and point source models. The text has been reformed in page 4 (lines 25-35) and page 5 (lines 1-15) in order to discuss differences in moment tensors and centroid moment tensors, and

centroid and hypocentre locations and make it easier for the reader to follow.

Text has been added in page 6 (lines 3-16) in order to discuss more thoroughly two case studies of large variability in reported source models as suggested by the Reviewer. New text in page 6 (lines 23-35) addresses the Reviewer's comments on identifying cases with large intra-event variability in source models and potential reasons that may contribute in these cases (see also page 6, lines 19-22).

Text in Section 5 (page 8, lines 25-36, and page 9, lines 4-18) explains further the scope of this paper, discusses complexities in different source models and gives some general hints on data quality characteristics which can help user of the ISC Bulletin to select a source model that may be more robust or more appropriate for their research.

Finally, we expanded the text at page 7 (lines 1-25) to better explain how the annual counts of Figure 6 (Figure 5 in the original manuscript) are obtained following the fault styles of Zoback (1992). Examples to events in the ISC Bulletin have also been added with links to specific events.

## 1.4 “Figures”

Reviewer 1 suggested a few improvements in the existing figures, namely:

1. Figure 1: to show the temporal evolution of target magnitudes or and histograms discussing the magnitude distribution. - We split the data period covered in this paper in six parts and added a subplot for each time period (decade) in order to show the time evolution in magnitude range coverage in the entire ISC Bulletin.
2. Figure 2: institute acronyms should be listed in an additional table. - We added Table 1 to address this comment.
3. Figure 3: add titles for the upper and bottom panels including information regarding agencies and minimum magnitude. - We addressed this comment accordingly.
4. Figure 5: add a symbol legend. - We added a legend in this figure (Figure 6 in the revised manuscript).
5. Figure 5: what are undefined earthquakes? - We explain this in the revised text (page 7, line 6)

## 1.5 Other minor comments

We addressed all the other points raised by Reviewer 1 as follows (following the same order as in the Reviewer's text):

1. page 1, line 8: clarify the meaning of 90% or reformulate the text. - We addressed this comment by adding more information (page 1, lines 8-9).

2. page4, line 5: maximum intra-event rotation (it needs better explanation, introduce the concept clearly). - We added new text in page 5 (lines 28-31) to address this comment.
3. page 4, line 14: "...can be as high as 100°". - We added new text in some earlier stage (page 5, lines 24-28) to address this comment and give an a physical explanation for the rotation angle.
4. page 5, line 8: "be aware of the techniques being used." Is the bulletin reporting details in techniques, data, azimuthal coverage? - The ISC Bulletin reports the number of stations and errors in the collected source models where available. This varies from agency to agency. This is partly a reason for the existnce of this paper, thus, to make users aware of this information. Text was added in pages 6 (lines 20-22) and 8 (lines 24-29).
5. page 5, line 12: "centroid or not" should be "centroid or hypocentre". - We corrected this (page 8, line 22).
6. page 5, line 10: "substantial mislocations" should be "substantial differences". - We corrected this (page 8, line 19).
7. page 5, line 21: should be 80% (not 90%). - This is right, we corrected it (page 9, line 23).
8. bottom of page 6: the similarity of both focal mechanisms and moment tensors is discussed only using a Kagan angle and the discussion is done on DC components. - We added some text to clarify this (page 9, lines 24-27).

## 2 Reviewer 2

### 2.1 "Which mechanism to use?"

Reviewer 2 mentions that the manuscript does not offer an answer in the question which mechanism to use in the case of multiple solutions for the same earthquake. In fact this question is mentioned in two out of four comments. The Reviewer claims that the current paper has tried to answer this question but the answer is not given. The Reviewer also proposed to develop the manuscript towards the direction of presenting the ISC computed focal mechanisms as more reliable, similar to the ISC event locations.

Let us first clarify that the purpose of this manuscript is an attempt to draw the Researchers' attention on the availability of source models in the ISC Bulletin. It is true that for decades the ISC put all of the effort in analysing and relocating seismic events with the use of reported parametric data, whilst not much, if any, has been done to promote the availability of collected source models. In fact the usage of the dedicated online web search on source mechanisms in 2018 corresponds to less than 2% of the total use of the ISC web site!

In this paper we aim to acknowledge the agencies reporting source models in the ISC Bulletin, discuss the differences in techniques being used by the major contributors, and highlight the complexities in combining source models following

different concepts and being determined by different techniques. For this reason we added new text in Section 3 (page 4, lines 1-18) and pages 8 (lines 24-36) and 9 (lines 4-19), also combining some of the comments posed by Reviewer 1. Moreover, in order to highlight how different agencies can act complementary to each other and provide a more comprehensive coverage in space, time and magnitude we added Figure 4 and new text in page 3 (lines 11-25).

Regarding the ISC computed focal mechanisms, we added these mechanism solutions in the analysis of the revised manuscript (also suggested by Reviewer 1) but we cannot claim that the ISC focal mechanism solutions are more reliable than other solutions provided by other agencies, since they can be fundamentally different in comparison with them. Take as an example a centroid based moment tensor solution (see for example additional text in page 4, lines 1-18, and page 6, lines 3-22). Finally, given all the above explained complexities, we gave only general recommendations described in page 8 (lines 24-36) and page 9 (lines 4-19). Yet again, we have made clear in the manuscript that it is not our purpose to recommend the use of one data provider over another, but we provide all the information we can collect and let the Researchers to choose this with respect to the needs of their research.

## **2.2 “Figures”**

Reviewer 2 commented that the Figures only cover statistical information and do not serve the purpose of the manuscript. We believe that enhancing the manuscript with new text, it is clearer and more obvious how the Figures presented in this paper can help readers to better understand the content of the ISC Bulletin regarding the availability of earthquake source models. The newly added Figure 4 should also help further towards this direction.

## **2.3 “Figure 5”**

In conjunction with Reviewer’s 2 first comment, more information was requested on how to choose a proper or target mechanism solution in Figure 5 for earthquakes with multiple solutions.

To address this point we have expanded the text first by recalling the fault styles according to Zoback (1992) and secondly by summarizing how we proceeded to characterize an event when multiple solutions are available. Examples for different cases have been given and links to real cases in the ISC Bulletin added. We believe that the revised text clarifies the scope of original Figure 5 (now Figure 6 in the revised manuscript). The revised text regards the entire page 7.

## **3 Additional changes**

We added a few new references to the manuscript, updated the figures and corresponding captions and at times streamlined the text.

# The ISC Bulletin as a comprehensive source of earthquake source mechanisms

Konstantinos Lentas <sup>1</sup>, Domenico Di Giacomo <sup>1</sup>, James Harris <sup>1</sup>, and Dmitry A. Storchak <sup>1</sup>

<sup>1</sup>International Seismological Centre, Pipers Lane, Thatcham, Berkshire, RG19 4NS, UK

**Correspondence:** Konstantinos Lentas (kostas@isc.ac.uk)

**Abstract.** In this article we summarize the availability of earthquake source mechanisms in the Bulletin of the International Seismological Centre (ISC). The bulletin in its current status contains ~81,000 seismic events with only one associated mechanism solution, and ~~~2225~~225,000 events with at least two associated source mechanisms. The main sources of earthquake mechanisms in the ISC Bulletin are reported solutions provided by data contributors, and ISC computed focal mechanisms based on first motion polarities. Given the importance of using pre-determined fault plane solutions in different types of studies, here we focus only on the reported mechanisms and we briefly discuss the methodologies adopted by major data providers to the ISC and investigate the intra-event variability of the source mechanisms. We conclude that the overall agreement among different earthquake ~~focal~~ mechanisms for the same event as reported by different sources can ~~be show a similarity coefficient~~ as high as ~~9080~~90.80%, based on the rotation angles of their best fitting double couple solutions, for the majority of the cases. The earthquake source mechanisms discussed in this work are freely available within the ISC Bulletin websearch at <http://doi.org/10.31905/D808B830>.

## 1 Introduction

The International Seismological Centre (ISC, [www.isc.ac.uk](http://www.isc.ac.uk)) currently collects station readings, hypocentre solutions and other earthquake bulletin data from approximately 150 agencies around the world. The ISC Bulletin contains over ~~7.7.6~~7.76 million seismic events (mostly earthquakes, as well as chemical and nuclear explosions, mine blasts and mining induced events, and other types of seismic events), and approximately ~~237 million individual~~256 million associated seismic station readings of arrival times, amplitudes, periods and first motion polarities (~~International Seismological Centre, 2018, database last accessed in October 2018~~)([International Seismological Centre, 2018, database last accessed in October 2018](#)).

Considerable effort is put into making sure that the station readings reported by different agencies belong to the correct seismic event. In the first instance, all parametric data sent to the ISC is collected and grouped automatically in unique seismic events. As soon as an event is created it is made openly available via the online ISC Bulletin ([www.isc.ac.uk/iscbulletin](http://www.isc.ac.uk/iscbulletin)). Secondly, the ISC analysts manually review (two/three years behind real-time) the collected station readings and hypocentre solutions for seismic events larger than approximately 3.5. If all conditions are met (details at <http://www.isc.ac.uk/iscbulletin/review/>), the ISC also recomputes location and magnitude (currently only MS and mb) by combining all the available phase arrival times and amplitude measurements, respectively. ISC location and magnitude procedures have recently been improved (Bondár and Storchak, 2011).

The ISC aims to increase the number of collected bulletins from national data centres or other sources (Willemann and Storchak, 2001) and improve its procedures in earthquake location and magnitude determinations (e.g., Bondár and Storchak, 2011; Di Giacomo and Storchak, 2015; Weston et al., 2018). As a result, the ISC Bulletin has proved to be a very useful resource for seismologists and geoscientists in general, as demonstrated by the vast use of ISC datasets in many research papers, including works on new tomographic models and global tectonics (e.g., Kennett et al., 1995; Rezapour and Pearce, 1998; Bormann et al., 2009; Hayes et al., 2012; Adam and Romanowicz, 2015; Zhan and Kanamori, 2016; Euler and Wyession, 2017; Lay et al., 2017). Recently the ISC has started to compute its own focal mechanisms (freely available in the reviewed bulletin) by using first motion polarities both from reported bulletins and picked automatically from waveform data (Lentas, 2017). In addition, the ISC Bulletin contains a substantial amount of source mechanisms (Fig. 1) calculated using different data and techniques as reported from various agencies working at local/regional and/or global scales, predominantly covering the period from mid 1970s till present.

In this paper, we aim to emphasize the availability of ~~reported (i.e., not computed by the ISC)~~ source mechanisms in the ISC Bulletin and discuss the different features of those solutions, aiming at helping ISC data users to decide how best to use the database according to the needs of their research.

## 2 Source mechanism contributions to the ISC Bulletin

There are currently ~~64~~65 agencies in the ISC Bulletin which have reported in the past or continue to report source mechanism solutions to the ISC (Fig. 2). By using the term source mechanisms we refer to both moment tensor solutions (and their associated best fitting double couple mechanisms) and pure double couple mechanisms of a point source. Table 1 shows details of the type of reported source mechanism solutions by each agency.

Major contributors of global source mechanisms include the Global Centroid Moment Tensor Project (GCMT, [www.globalcmt.org](http://www.globalcmt.org), Dziewonski et al., 1981; Ekström et al., 2012), the US National Earthquake Information Center (NEIC, or NEIS prior to 1984), and for regional earthquakes, the National Research Institute for Earth Science and Disaster Resilience (NIED) in Japan and the Pacific Northwest Seismic Network (PNSN).

Note that prior to data year 2006 the agency code HRVD (Harvard University) was used throughout the ISC Bulletin for GCMT solutions. Here we use a unique agency code for these source mechanisms and replace the HRVD agency code with the GCMT code throughout the ISC Bulletin. This is already done for the time period 1976-1979 covered by the first part of the ISC rebuild project (1964-1979, Storchak et al., 2017). After completion of the ISC rebuild project, all remaining HRVD source mechanism solutions will be available under the GCMT code. Moreover, moment tensor solutions for 76 intermediate depth earthquakes and 104 deep earthquakes from 1962 to 1976 have been added under the GCMT agency code (Chen et al., 2001; Huang et al., 1997).

Since the mid 1990s numerous other agencies, mainly national data centres, started reporting source mechanism solutions to the ISC. This has resulted in a steep increase of available mechanism solutions in the ISC Bulletin (Fig. 1). Nevertheless, the coverage and completeness of seismic events with associated source mechanisms is not uniform and primarily depends on

the tectonics and the associated seismicity in different regions, the station coverage and the practices of the reporting agencies (Fig. 3).

In May 2018 the ISC published in the online ISC Bulletin its first automatic focal mechanism solutions obtained from reported first motion polarities and automatic picks of waveform data. These are focal mechanisms for reviewed and relocated earthquakes in the Reviewed ISC Bulletin with  $m_b^{ISC} \geq 4.5$ , starting from data month January 2011. Since then new focal mechanism are routinely added for every data month added in the Reviewed ISC Bulletin. Moreover, we have published focal mechanism solutions obtained from reported polarities for the ISC relocated earthquakes ( $m_b^{ISC} > 3.5$ ) covering the time period 1964 - 1984 as part of the rebuild project (Storchak et al., 2017) and focal mechanism solutions obtained from reported polarities in the ISS (1938-1963) Bulletins (see <http://www.isc.ac.uk/projects/focalmechs/>). The gap currently shown in Figure 2 between 1985 and 2010 is expected to be bridged gradually until the completion of the rebuild project.

Figure 4 shows the geographical and magnitude distribution of earthquakes with source mechanisms reported by the agencies which systematically send their mechanism solutions to the ISC (see also Fig. 2 for numbers of reported source mechanisms by agency and distribution in time). Local agencies are important to complement the results of global agencies as they cover events with lower magnitudes. For example, GCMT computes source mechanisms for global earthquakes with magnitude 5.0 and above, but also slightly lower ( $\sim 4.5$ ) depending on the area and station coverage. Similarly, NEIC covers earthquakes with a minimum magnitude of  $M_W \sim 4.5$  on a global scale. IGP reports earthquakes with magnitude  $M_W$  5.5 and above, whilst at the ISC we attempt to determine the focal mechanisms of earthquakes with  $m_b \geq 4.5$ . On the other hand, agencies like JMA and NIED cover a much wider magnitude range together, in comparison to global agencies, offering a more complete coverage in earthquake source mechanisms for Japan. Similar observations can be drawn for agencies BRK, ECX and PNSN covering the seismicity along the western coast of the United States, several European agencies (e.g., MED-RCMT and ZUR-RMT) covering the seismicity of Europe, and agencies RSNC and WEL for Colombia and New Zealand, respectively. This obviously introduces some heterogeneity in the available solutions for different areas and for different magnitude ranges, as a result of the different techniques applied by different agencies for the determination of their source models. Moreover, different agencies report different types of source mechanism and associated parametric data (nodal planes, moment tensor components, principal axes - see also Table 1 for details). More details will be given in Section 3.

All the available source mechanisms are included in the ISC Bulletin. However, users particularly interested in focal mechanisms can search using either a dedicated tool at <http://www.isc.ac.uk/iscbulletin/search/fmechanisms/> or webservice at <http://www.isc.ac.uk/iscbulletin/search/webservices/fmechanisms/>. Search parameters include date, area, magnitude, depth and agency code. Search outputs are available either in a comma separated CSV-like format with one line per mechanism solution, or in QuakeML format. Included in the output are the ISC event identifier, scalar moments, moment tensor components, nodal planes, principal axes and the hypocentre/centroid parameters for each mechanism solution, where applicable. The format is explained in detail at <http://www.isc.ac.uk/iscbulletin/search/fmechanisms/csvoutput/>.

### 3 Source mechanism variability



The majority of the observed global seismicity is characterized by crustal (shallow) earthquakes which occur as the result of a sudden release of accumulated strain across a seismic fault of finite dimensions. A seismic source whose energy is recorded at stations located at distances of several wavelengths from the source, can be approximated as a point source. The point source model provides a simple and convenient approach in order to simulate the seismic radiation. Nonetheless, for larger earthquakes (for example mega-thrust earthquakes, Tsai et al., 2005; Lentas et al., 2014; Ye et al., 2016), and/or earthquakes observed at distances close to the source (a few kilometers), the point source approximation is not sufficient, and ideally, the rupture propagation history and finite fault characteristics should be taken into account when attempting to model the seismic source. For deep earthquakes, on the other hand, non conventional models have been proposed (e.g., Okal, 2001; Meng et al., 2014), but the point source approximation is also being used for the sake of simplicity and processing consistency with crustal (shallow) earthquakes.

A seismic point source is described in principal by a double couple system of equivalent body forces which are represented by two unit vectors, the normal and slip vectors. These vectors are defined by the orientation of the fault and the direction of slip in terms of the strike, dip and rake angles (e.g., Aki and Richards, 2002). Different techniques follow different concepts for determining the source model of a point source. Some algorithms solve directly for the geometry of a planar fault, meaning the strike, dip and rake of the fault and auxiliary planes assuming a pure double couple mechanism. Other techniques determine the six components of the moment tensor. This is a mathematical representation of the equivalent body forces acting on a seismic point source, and can be decomposed into an isotropic component, a compensated linear vector dipole (CLVD) and a best fitting double couple mechanism describing the geometry of a planar fault.

Source mechanism solutions can be determined by using two main data types: (i) parametric data such as first motion *P*-wave polarities and amplitude ratios, and (ii) waveform data modelling. A vast variety of techniques and algorithms have been developed over the last few decades using different concepts and data. The most robust results are obtained by waveform modelling methods. Even though techniques based on polarities depend strongly on the network geometry and the station azimuthal coverage, they can still be very useful in determining the focal mechanisms of small earthquakes and aftershock sequences using local networks (Shearer, 1998). ~~Focal mechanisms-~~

~~Focal mechanisms expressed in terms of strike, dip and rake angles,~~ based on first motion polarities ~~are reported for example by JMA (Nakamura and Mochizuki, 1988), PNSN (FPFIT code by Reasenber and Oppenheimer, 1985) and ISC (Lentas, 2017). They depend on pre-determined locations and~~ reflect the geometry of the seismic fault at the initial breaking of the rupture, ~~whereas waveform modelling techniques usually consider.~~ Waveform modelling techniques on the other hand usually consider a group of phases (body waves and/or surface waves) and provide source mechanisms closer to the dominant component of the entire rupture geometry.

~~Moreover, different techniques follow different concepts on determining the source model of a point source. Some algorithms solve directly for the geometry of a planar fault, meaning the~~ Their source models can be expressed as strike, dip and rake of the fault and auxiliary planes assuming a pure double couple mechanism. This is more common in first motion polarity-based techniques (i.e., Reasenber and Oppenheimer, 1985, which is used by PNSN), but it can also be the case in waveform angles or moment tensor components and they can either be based on a pre-determined location or can be centroid solutions. Centroid

based techniques determine the six elements of the moment tensor, the centroid location (latitude, longitude and depth of a point source) and the origin time simultaneously. ~~Waveform~~ modelling methods such as the SCARDEC technique (Vallée et al., 2010) ~~reported~~, whose source models are reported to the ISC by agency IGP. ~~These techniques usually depend~~, make use of body-wave phases and the NEIC location. The obtained source models are expressed as strike, dip and rake angles of a pure double couple source. Source models of other waveform modelling techniques based on a pre-determined location. ~~Other techniques determine the six components~~ hypocentre location are also routinely reported in the ISC Bulletin, such as the NIED solutions for the Japanese area (Fukuyama and Kawai, 1998). In this case the reported source models are expressed in terms of the six elements of the moment tensor ~~which is a mathematical representation of the equivalent body forces acting on a seismic point source, and can be decomposed into an isotropic component, a compensated linear vector dipole (CLVD) and a~~ and the best fitting double couple ~~mechanism which describes the geometry of a planar fault~~ solution. Centroid based techniques like the GCMT ~~, the MedNet Regional Centroid Moment Tensors ((Dziewonski et al., 1981; Ekström et al., 2012), MED-RCMT ); the Zurich Moment Tensors ((Pondrelli et al., 2011), ZUR-RMT )~~ (Braunmiller et al., 2002) and others applied by the NEIC (Dziewonski et al., 1981; Ekström et al., 2012; Braunmiller et al., 2002; Pondrelli et al., 2011; ?; ?; Benz and Herrmann, 2014) use this concept and simultaneously determine the centroid location. (Hayes et al., 2009; Benz and Herrmann, 2014) provide centroid locations, moment tensor components and best fitting double couple solutions.

Source models for all the above mentioned techniques can be found in the ISC Bulletin (see Fig. 2). ~~Other automated moment tensors which depend on a pre-determined hypocentre location are also routinely reported in the ISC Bulletin, for example, by NIED in Japan (Fukuyama and Kawai, 1998).~~ Moreover, taking into consideration additional differences in velocity models, station distribution and observations in different waveform frequency bands that are being used by different techniques, some variation among source mechanisms reported by different agencies for the same seismic event ~~has is~~ to be expected.

Figure 4-5 shows the frequency distribution of available source mechanism solutions per event in the ISC Bulletin, and the frequency distribution of maximum intra-event rotation angle for the events having at least two reported mechanism solutions. The rotation angle describes the transformation of a double couple mechanism into another arbitrary mechanism through 3-D rotations (Kagan, 1991). ~~The vast majority peak between~~ To further visualise this it is worth noting that the rotation angle between two double couple mechanism solutions can vary between  $0^\circ$  and  $120^\circ$ , where  $0^\circ$  corresponds to perfect match and  $120^\circ$  describes absolute mismatch which physically translates into mechanism solutions showing for example perpendicular strike orientations and/or conflicting fault types (e.g., normal compared to thrust, or normal compared to pure strike slip, and so on). Consequently, in order to determine the maximum intra-event rotation angle we list all the available best fitting double couple source mechanism solutions for each event and we calculate rotation angles for all possible pairs. We then pick the maximum value as describing the greatest mismatch between the available mechanisms, and hence, the extent of expected differences in strike, dip and rake (see for example Fig. 1 in Cesca et al., 2013).

A substantial number of earthquakes in the ISC Bulletin ( $\sim 40\%$ ) show intra-event rotation angles between  $10^\circ$  and  $2025^\circ$  (see Figure 5). Cases showing large differences, above  $40^\circ$ , ~~are not rare occasionally occur~~ and can be partly explained by earthquakes showing complex rupture, such as the 2002 November 3,  $M_W$  7.9, Denali Central Alaska earthquake (Ozacar et al., 2003) (e.g., Ozacar et al., 2003) and the doublet 2012 December 7,  $M_W$  7.2, east coast of Honshu earthquake (Lay et al., 2013) (e.g., Lay et al., 2013) which show

intra-event rotation angles up to  $60^\circ$  and  $90^\circ$  respectively, or comparing automatic source mechanism solutions such as the 2015 May 25,  $M_W$  5.2, eastern Honshu earthquake which shows intra-event rotation angles up to  $\sim 90^\circ$ .

In the case of the Denali event ([http://www.isc.ac.uk/cgi-bin/web-db-v4?event\\_id=6123395&out\\_format=IMS1.0&request=COMPREHENSIVE](http://www.isc.ac.uk/cgi-bin/web-db-v4?event_id=6123395&out_format=IMS1.0&request=COMPREHENSIVE)), the earthquake started as a thrust event and then ruptured along the curved strike-slip Denali fault (Ozacar et al., 2003). Similarly, Lay et al. (2013) suggested the case of a doublet event that began with a  $M_W$  7.2, thrust earthquake ([http://www.isc.ac.uk/cgi-bin/web-db-v4?event\\_id=607215270&out\\_format=IMS1.0&request=COMPREHENSIVE](http://www.isc.ac.uk/cgi-bin/web-db-v4?event_id=607215270&out_format=IMS1.0&request=COMPREHENSIVE)) and followed by a  $M_W$  7.1-7.2, normal-faulting earthquake ([http://www.isc.ac.uk/cgi-bin/web-db-v4?event\\_id=602005586&out\\_format=IMS1.0&request=COMPREHENSIVE](http://www.isc.ac.uk/cgi-bin/web-db-v4?event_id=602005586&out_format=IMS1.0&request=COMPREHENSIVE)) for the case of the 2012 east coast of Honshu earthquake. Since the routinely reported mechanism solutions for these complex events are based on the point source approximation, it is no surprise that different episodes of rupture are captured by different methods using different data. For example, the Global CMT, by mainly using long period surface waves (and in some cases long period body waves), has detected the strike-slip nature of the Denali earthquake. In contrast, NEIC source mechanism captured the initial stage of the rupture and shows evidence of thrust faulting. Note that large intra-event rotation angles can occur also for moderate earthquake, such as for the 2015 May 25,  $M_W$  5.2, eastern Honshu earthquake. Indeed, for this event the intra-event rotation angles are up to  $\sim 90^\circ$ . It is likely that such variability is due to differences in the methods and data applied (first-motion by JMA and ISC versus waveform modelling by NIED and GCMT) rather than to rupture complexity. Substantial intra-event differences are also very common as a result of multiple solutions reported by PNSN for small earthquakes with poorly constrained source mechanisms, such as the 1981 February 11,  $M_d$  2.5, Washington earthquake where the maximum intra-event rotation angle can be as high as  $100^\circ$ . This is very common in the case of PNSN reported mechanisms due to the first motion technique that is being used which provides multiple solutions if the data is not adequate for the determination of a single well constrained solution. Quality characterization found in the comments in the online ISC Bulletin can help users to identify the most robust mechanism solution among multiple PNSN provided mechanisms for the same event.

~~As source mechanisms are~~ Intra-event rotation angles are not currently calculated and published in a systematic way in the ISC Bulletin, and hence, the identification of cases of substantial differences in reported source models is not part of the ISC standard procedures. Researchers who are interested in earthquake source model validations and assessment are encouraged to make use of the ISC Bulletin for this purpose and apply their own schemes. As already mentioned, the source models in the ISC Bulletin are not reviewed by the analysts. However, we frequently carry out health checks of the bulletin, namely, removing duplicates and checking for consistency between moment tensors and best fitting double couple solutions, or moment tensors and principal axes, or double couple solutions and principal axes and so on. This has resulted in correcting a few cases (less than 1% of the entire bulletin) where either typos from the data providers or bugs in the database parsing algorithms have been identified and fixed. As part of this process we did not detect systematic inconsistencies between source models provided by different agencies for the same seismic event that has to do with the type of source mechanism, or depth, or a specific region. The most common case in this respect is source mechanisms provided by PNSN for small magnitude events along the western coast of United States due to the type of methodology that is being applied which determines multiple solutions when the data quality is poor.

Source mechanisms are also used in a variety of studies (e.g., tectonics, stress patterns, cluster analysis) ~~in Figure 5 we show~~ aiming at characterizing an event or a set of events by the fault style. Several classifications of fault styles are available in the literature (see, e.g., C  l  rier, 2010, for an overview). For this work we adopt the classification proposed by Zoback (1992) for the World Stress Map Project (<http://www.world-stress-map.org/>). With such classification an event is assigned one of the following fault styles: thrust, normal, strike-slip, normal with strike-slip component, thrust with strike-slip component and undefined for events not fitting in any of the previous categories (similarly to the “odd” group in Frohlich, 1992). Figure 6 shows the annual number of earthquakes grouped ~~by the faulting styles proposed by Zoback (1992) with~~ according to Zoback (1992) and the sole intent of ~~showcasing the figure is to showcase~~ one of the possible uses for the source mechanisms in the ISC Bulletin.

Considering the source mechanisms in the ISC Bulletin, as discussed so far, it is obvious that for events with only one mechanism available the fault style is easily assigned. The same also applies to events with multiple solutions all of the same fault style (e.g., [http://www.isc.ac.uk/cgi-bin/web-db-v4?event\\_id=3021752&out\\_format=IMS1.0&request=COMPREHENSIVE](http://www.isc.ac.uk/cgi-bin/web-db-v4?event_id=3021752&out_format=IMS1.0&request=COMPREHENSIVE)). However, for events with more than one solution it is not always possible to assign a fault style with the solutions at hand. This happens, for example, when an event has two source mechanisms and one being thrust and the other normal (e.g., [http://www.isc.ac.uk/cgi-bin/web-db-v4?event\\_id=602214316&out\\_format=IMS1.0&request=COMPREHENSIVE](http://www.isc.ac.uk/cgi-bin/web-db-v4?event_id=602214316&out_format=IMS1.0&request=COMPREHENSIVE)). Similarly, if an event has multiple solutions, we may have source mechanisms falling into more than two fault styles (e.g., [http://www.isc.ac.uk/cgi-bin/web-db-v4?event\\_id=602431903&out\\_format=IMS1.0&request=COMPREHENSIVE](http://www.isc.ac.uk/cgi-bin/web-db-v4?event_id=602431903&out_format=IMS1.0&request=COMPREHENSIVE)) or without a unique maximum in the number of source mechanisms belonging to a fault style (e.g., [http://www.isc.ac.uk/cgi-bin/web-db-v4?event\\_id=602945524&out\\_format=IMS1.0&request=COMPREHENSIVE](http://www.isc.ac.uk/cgi-bin/web-db-v4?event_id=602945524&out_format=IMS1.0&request=COMPREHENSIVE)). In such cases we do not assign a fault style to an event. If, instead, out of the fault style distribution within an event there is a more recurrent fault style (e.g., [http://www.isc.ac.uk/cgi-bin/web-db-v4?event\\_id=2944860&out\\_format=IMS1.0&request=COMPREHENSIVE](http://www.isc.ac.uk/cgi-bin/web-db-v4?event_id=2944860&out_format=IMS1.0&request=COMPREHENSIVE)), we still assign a fault style to the event. Therefore, in Figure 6 we also show the “Discrepant” category for those events where we could not assign a specific fault style. Note that also complex earthquakes, such as the ones previously mentioned, may fall into this category. The annual percentage of events falling into the “Discrepant” category is usually between 0 and 5%, with a maximum of 8% in 2000. The occurrence of such “Discrepant” events should not discourage the use of source mechanisms from the ISC Bulletin. Indeed, this category of events can highlight complex events or events for which more studies are needed.

Not surprisingly, Figure 6 shows how the thrust earthquakes dominate the annual occurrences, with the only exceptions for year 2000 (significant strike-slip aftershock sequence following the 2000 Tottori, Japan, earthquake) and 2011 (aftershocks of the 2011 Tohoku earthquake, due to the stress field change, were characterized by many normal-fault earthquakes, as shown, e.g., by Hasegawa et al. (2012)). ~~Figure 5 also shows the annual number of earthquakes for which we could not assign a fault style. This is due to large intra-event variability of Hasegawa et al., 2012). Such use of the source mechanisms due to complexities, as mentioned earlier, or to conflicting reports. Note that classifications different from Zoback (1992) have been proposed (see, e.g., C  l  rier, 2010, for an overview), and users can apply their preferred classification from the source mechanisms within the ISC Bulletin~~ in the ISC Bulletin can be used for event characterizations and applied, e.g. to studies concerning regional stress patterns (e.g., Balfour et al., 2011), magnitude (e.g., Lomax and Michelini, 2009), tsunami/tsunamigenic earthquakes (e.g., Okal and Newman, 2001), and other seismological fields.

#### 4 Data availability

- 5 The earthquake source mechanisms summarized in this work are freely available within the ISC Bulletin websearch at <http://doi.org/10.31905/D808B830>. All data used in this paper are maintained at the ISC ([www.isc.ac.uk](http://www.isc.ac.uk), last accessed ~~20 November~~ ~~2018~~ 15 March 2019).

#### 5 Summary and conclusions

The ISC offers the most comprehensive bulletin of global seismicity in terms of hypocentre solutions, phase arrivals, magnitudes and amplitude measurements. In this paper we ~~presented~~ present an additional aspect of the ISC Bulletin, namely its source mechanism content and the opportunity for ISC users to complement the source mechanism information with all other data included in the ISC Bulletin.

The ISC has a mandate to collect as much parametric data as possible from various sources around the world and make it freely available to the seismological as well as to a broad geoscience community. As a result, the magnitude range of earthquakes covered by available mechanism solutions is larger than individual global catalogues such as GCMT or NEIC. However, this feature inevitably leads to a higher heterogeneity in the solutions due to different methods adopted by each provider. Thus, users are advised to be aware of the techniques being used in the computations of the various source models in the ISC Bulletin. For example, centroid based mechanism solutions should be used together with centroid locations, since: (i) both the centroid mechanism and centroid location are parts of the same output, and (ii) substantial ~~mislocations~~ differences may exist among centroid locations and standard hypocentre locations fitting the observed phase arrival times of body waves. To facilitate this the CSV format provided by the online ISC Bulletin indicates whether a mechanism solution is centroid or ~~no~~ hypocentre. Source mechanisms obtained from pre-determined standard hypocentre locations can be used together with the provider's hypocentre solution or the prime hypocentre solution in the ISC Bulletin. However, large differences in depth may be present in some cases among the prime hypocentre solution and the solution provided by the mechanism's agency. Moreover, information regarding the quality of the obtained mechanism solutions such as the number of stations being used and/or errors in the obtained source models is also provided in the ISC Bulletin where available from the reporting agency. The latter is more common in moment tensor solutions. Detailed quality information is routinely provided for the ISC focal mechanism solutions both in the comments section in the online ISC Bulletin as well as by clicking on the "ISC Focal Mechanism" logo on the top of the online bulletin.

30 In the case of multiple available source mechanism solutions for the same event the question "which source model should I pick for my research from the ISC Bulletin" arises from a researcher's practical point of view. Since the source model determination problem is not fully resolved and the point source approximation is still the standard in routine earthquake mechanism representation, unfortunately there is no easy and straightforward answer to this question. The purpose of this paper is to give insights to the content of the ISC Bulletin regarding the availability of earthquake source models for seismologists, and further highlight the complexity of the earthquake source and the associated point source models for the broad geoscience community. Thus, only general suggestions can be given without attempting or willing to discriminate the data in the ISC

5 Bulletin in right or wrong. Users of the bulletin should keep in mind that different techniques are based on different data  
(first motion polarities, body wave amplitude ratios, body/surface wave modelling) and as a result they weight differently  
various episodes in the rupture history which ultimately can have a strong effect on the final solution. For example, slip  
takes place across seismic faults that are not necessarily planar, but their orientation can vary with length. Moreover, the  
use of local/regional or teleseismic data can be another component of source model variations, and in conjunction with the  
uncertainties in velocity models that are being used to simulate the wave propagation in the Earth's interior it is advisable to  
10 take into consideration the details of the methodologies being used by different source mechanism providers.

In all cases, the ISC will not deprecate solutions without being revised by the data provider. The ISC cannot encourage the  
use of source models of one data provider over another. We try to include as much information as possible in the comments  
section of the online ISC Bulletin regarding each mechanism solution. Researchers are free to make use of any source models  
they might think that fit their research and to facilitate this we advise users to pay attention to quality information provided  
15 in the comments section, namely, the number of stations and components being used, the azimuthal gap and so on. Another  
tool for selecting the most appropriate source model is the association of the mechanism solution with the type of location  
being used (hypocentre/centroid) as stated above. In this paper we briefly summarized the methods being used by major model  
reporters. However, users still need to identify the most appropriate source models for their research.

Similar to the variation of hypocentre solutions in the ISC Bulletin, multiple source mechanisms for the same seismic event,  
20 when available, can provide a measure of the posterior uncertainties with respect to data errors and modelling techniques.  
Despite variations in methods and data used to compute the solutions, we showed that in most cases there is a good agreement  
among multiple solutions provided by different agencies. The intra-event variability in the ISC Bulletin was quantified by  
the maximum rotation angle and is well constrained up to  $20^\circ$ , which corresponds to a similarity coefficient of  $\sim 90\%$   
(Cesca et al., 2013). As already mentioned above both moment tensors and focal mechanism solutions are included in the  
25 ISC Bulletin. Since we needed to compare results obtained from these fundamentally different concepts, obtained by different  
techniques, we decided to focus on pure double couple and moment tensor best fitting double couple mechanisms, which  
describe the geometry of the seismic source. For this reason the use of the rotation angle (Kagan, 1991) was selected. Different  
classification techniques and metrics could be applied (i.e., Helffrich, 1997; Frohlich and Davis, 1999) but for the purpose  
of the current article the rotation angle is considered to be an adequate metric of source mechanism variability. Similarly, by  
30 applying Zoback (1992) fault styles classification, we observe a large intra-event variability for up to 8% of the earthquakes  
per year ~~a large intra-event variability~~.

Source mechanisms are currently not reviewed by ISC analysts, and the user should pick, if required, the preferred one  
when multiple solutions are available for an event. The ISC values all agencies reporting source mechanism solutions and  
encourages new ones to submit theirs. In parallel, we recommend researchers ~~to~~ make a more systematic use of the earthquake  
source mechanisms in the ISC Bulletin in future studies.

5 *Author contributions.* KL was the leading author of the paper and compiled most of the figures. DDG vetted the data for the magnitude and fault styles plots, JH maintains the database and webservice and DAS obtained the funding for the work and established and maintained connections with many data providers. All authors contributed to the manuscript and approved the final version.

*Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* The [authors wish to thank two anonymous Reviewers for their comments and suggestions which hepled to improve this manuscript.](#) The work done at the ISC is possible thanks to the support of its members ([www.isc.ac.uk/members/](http://www.isc.ac.uk/members/)) and sponsors ([www.isc.ac.uk/sponsors/](http://www.isc.ac.uk/sponsors/)). Work partially funded by NSF Grants 1417970 and 1811737, and a USGS Award G18AP00035. ~~Some figures~~ [Figures](#) were drawn using the Generic Mapping Tools (Wessel et al., 2013) [and the Matplotlib python library \(Hunter, 2007\).](#)

## 5 References


- Adam, J.-C. and Romanowicz, B.: Global scale observations of scattered energy near the inner-core boundary: Seismic constraints on the base of the outer-core, *Physics of the Earth and Planetary Interiors*, 245, 103–116, <https://doi.org/10.1016/j.pepi.2015.06.005>, <https://doi.org/10.1016/j.pepi.2015.06.005>, 2015.
- Aki, K. and Richards, P. G.: *Quantitative Seismology: Theory and Methods*, University Science Books, 2 edn., 2002.
- 10 Balfour, N. J., Cassidy, J. F., Dosso, S. E., and Mazzotti, S.: Mapping crustal stress and strain in southwest British Columbia, *Journal of Geophysical Research*, 116, <https://doi.org/10.1029/2010jb008003>, <https://doi.org/10.1029/2010jb008003>, 2011.
- Benz, H. M. and Herrmann, R. B.: Rapid Estimates of the Source Time Function and Mw using Empirical Green’s Function Deconvolution—Rapid Estimates of the Source Time Function and Mw using EGF Deconvolution, *Bulletin of the Seismological Society of America*, 104, 1812, <https://doi.org/10.1785/0120130325>, <http://dx.doi.org/10.1785/0120130325>, 2014.
- 15 Bondár, I. and Storchak, D. A.: Improved location procedures at the International Seismological Centre, *Geophysical Journal International*, 186, 1220–1244, <https://doi.org/10.1111/j.1365-246x.2011.05107.x>, <https://doi.org/10.1111/j.1365-246x.2011.05107.x>, 2011.
- Bormann, P., Liu, R., Xu, Z., Ren, K., Zhang, L., and Wendt, S.: First Application of the New IASPEI Teleseismic Magnitude Standards to Data of the China National Seismographic Network, *Bulletin of the Seismological Society of America*, 99, 1868–1891, <https://doi.org/10.1785/0120080010>, <https://doi.org/10.1785/0120080010>, 2009.
- 20 Braunmiller, J., Kradolfer, U., Baer, M., and Giardini, D.: Regional moment tensor determination in the European–Mediterranean area — initial results, *Tectonophysics*, 356, 5–22, [https://doi.org/10.1016/s0040-1951\(02\)00374-8](https://doi.org/10.1016/s0040-1951(02)00374-8), [https://doi.org/10.1016/s0040-1951\(02\)00374-8](https://doi.org/10.1016/s0040-1951(02)00374-8), 2002.
- Célérier, B.: Remarks on the relationship between the tectonic regime, the rake of the slip vectors, the dip of the nodal planes, and the plunges of the P, B, and T axes of earthquake focal mechanisms, *Tectonophysics*, 482, 42–49, <https://doi.org/10.1016/j.tecto.2009.03.006>, <https://doi.org/10.1016/j.tecto.2009.03.006>, 2010.
- 25 Cesca, S., Şen, A. T., and Dahm, T.: Seismicity monitoring by cluster analysis of moment tensors, *Geophysical Journal International*, 196, 1813–1826, <https://doi.org/10.1093/gji/ggt492>, <https://doi.org/10.1093/gji/ggt492>, 2013.
- Chen, P. F., Nettles, M., Okal, E. A., and Ekström, G.: Centroid moment tensor solutions for intermediate-depth earthquakes of the WWSSN–HGLP era (1962–1975), *Physics of the Earth and Planetary Interiors*, 124, 1–7, [https://doi.org/10.1016/s0031-9201\(00\)00220-x](https://doi.org/10.1016/s0031-9201(00)00220-x), [https://doi.org/10.1016/s0031-9201\(00\)00220-x](https://doi.org/10.1016/s0031-9201(00)00220-x), 2001.
- 30 Di Giacomo, D. and Storchak, D. A.: A scheme to set preferred magnitudes in the ISC Bulletin, *Journal of Seismology*, 20, 555–567, <https://doi.org/10.1007/s10950-015-9543-7>, <https://doi.org/10.1007/s10950-015-9543-7>, 2015.
- Dziewonski, A. M., Chou, T.-A., and Woodhouse, J. H.: Determination of earthquake source parameters from waveform data for studies of global and regional seismicity, *Journal of Geophysical Research: Solid Earth*, 86, 2825–2852, <https://doi.org/10.1029/jb086ib04p02825>, <https://doi.org/10.1029/jb086ib04p02825>, 1981.
- 35 Ekström, G., Nettles, M., and Dziewoński, A.: The global CMT project 2004–2010: Centroid-moment tensors for 13, 017 earthquakes, *Physics of the Earth and Planetary Interiors*, 200-201, 1–9, <https://doi.org/10.1016/j.pepi.2012.04.002>, <https://doi.org/10.1016/j.pepi.2012.04.002>, 2012.
- Euler, G. G. and Wyssession, M. E.: Geographic variations in lowermost mantle structure from the ray parameters and decay constants of core-diffracted waves, *Journal of Geophysical Research: Solid Earth*, 122, 5369–5394, <https://doi.org/10.1002/2017jb013930>, <https://doi.org/10.1002/2017jb013930>, 2017.




- 5 Frohlich, C.: Triangle diagrams: ternary graphs to display similarity and diversity of earthquake focal mechanisms, *Physics of the Earth and Planetary Interiors*, 75, 193–198, [https://doi.org/10.1016/0031-9201\(92\)90130-n](https://doi.org/10.1016/0031-9201(92)90130-n), [https://doi.org/10.1016/0031-9201\(92\)90130-n](https://doi.org/10.1016/0031-9201(92)90130-n), 1992.
- Frohlich, C. and Davis, S. D.: How well constrained are well-constrained T, B, and P axes in moment tensor catalogs?, *Journal of Geophysical Research: Solid Earth*, 104, 4901–4910, <https://doi.org/10.1029/1998jb900071>, <https://doi.org/10.1029/1998jb900071>, 1999.
- Fukuyama, E., S. I. D. S. D. and Kawai, H.: Automated seismic moment tensor determination by using on-line broadband seismic waveforms, *Journal of the Seismological Society of Japan*, 2, 149–156, 1998.
- 10 Hasegawa, A., Yoshida, K., Asano, Y., Okada, T., Iinuma, T., and Ito, Y.: Change in stress field after the 2011 great Tohoku-Oki earthquake, *Earth and Planetary Science Letters*, 355-356, 231–243, <https://doi.org/10.1016/j.epsl.2012.08.042>, <https://doi.org/10.1016/j.epsl.2012.08.042>, 2012.
- Hayes, G. P., Rivera, L., and Kanamori, H.: Source Inversion of the W-Phase: Real-time Implementation and Extension to Low Magnitudes, *Seismological Research Letters*, 80, 817–822, <https://doi.org/10.1785/gssrl.80.5.817>, <https://doi.org/10.1785/gssrl.80.5.817>, 2009.
- 15 Hayes, G. P., Wald, D. J., and Johnson, R. L.: Slab1.0: A three-dimensional model of global subduction zone geometries, *Journal of Geophysical Research: Solid Earth*, 117, <https://doi.org/10.1029/2011jb008524>, <https://doi.org/10.1029/2011jb008524>, 2012.
- Helffrich, G. R.: How good are routinely determined focal mechanisms? Empirical statistics based on a comparison of Harvard, USGS and ERI moment tensors, *Geophysical Journal International*, 131, 741–750, <https://doi.org/10.1111/j.1365-246x.1997.tb06609.x>, <https://doi.org/10.1111/j.1365-246x.1997.tb06609.x>, 1997.
- 20 //doi.org/10.1111/j.1365-246x.1997.tb06609.x, 1997.
- Huang, W.-c., Okal, E. A., Ekström, G., and Salganik, M. P.: Centroid moment tensor solutions for deep earthquakes predating the digital era: the World-Wide Standardized Seismograph Network dataset (1962–1976), *Physics of the Earth and Planetary Interiors*, 99, 121–129, [https://doi.org/10.1016/s0031-9201\(96\)03177-9](https://doi.org/10.1016/s0031-9201(96)03177-9), [https://doi.org/10.1016/s0031-9201\(96\)03177-9](https://doi.org/10.1016/s0031-9201(96)03177-9), 1997.
- Hunter, J. D.: Matplotlib: A 2D graphics environment, *Computing In Science & Engineering*, 9, 90–95, 2007.
- 25 International Seismological Centre: On-line Bulletin, Internatl. Seismol. Cent., Thatcham, United Kingdom, <http://www.isc.ac.uk>, 2018.
- ISS: International Seismological Summary, annual volumes, 1938-1963.
- Kagan, Y. Y.: 3-D rotation of double-couple earthquake sources, *Geophysical Journal International*, 106, 709–716, <https://doi.org/10.1111/j.1365-246x.1991.tb06343.x>, <https://doi.org/10.1111/j.1365-246x.1991.tb06343.x>, 1991.
- Kennett, B. L. N., Engdahl, E. R., and Buland, R.: Constraints on seismic velocities in the Earth from traveltimes, *Geophysical Journal International*, 122, 108–124, <https://doi.org/10.1111/j.1365-246x.1995.tb03540.x>, <https://doi.org/10.1111/j.1365-246x.1995.tb03540.x>, 1995.
- 30 Lay, T., Duputel, Z., Ye, L., and Kanamori, H.: The December 7, 2012 Japan Trench intraplate doublet ( $M_w$  7.2, 7.1) and interactions between near-trench intraplate thrust and normal faulting, *Physics of the Earth and Planetary Interiors*, 220, 73–78, <https://doi.org/10.1016/j.pepi.2013.04.009>, <https://doi.org/10.1016/j.pepi.2013.04.009>, 2013.
- Lay, T., Ye, L., Koper, K. D., and Kanamori, H.: Assessment of teleseismically-determined source parameters for the April 25, 2015  $M_w$  7.9 Gorkha, Nepal earthquake and the May 12, 2015  $M_w$  7.2 aftershock, *Tectonophysics*, 714-715, 4–20, <https://doi.org/10.1016/j.tecto.2016.05.023>, <https://doi.org/10.1016/j.tecto.2016.05.023>, 2017.
- 35 Lentas, K.: Towards routine determination of focal mechanisms obtained from first motion P-wave arrivals, *Geophysical Journal International*, 212, 1665–1686, <https://doi.org/10.1093/gji/ggx503>, <https://doi.org/10.1093/gji/ggx503>, 2017.
- Lentas, K., Ferreira, A., Clévéde, E., and Roch, J.: Source models of great earthquakes from ultra low-frequency normal mode data, *Physics of the Earth and Planetary Interiors*, 233, 41–67, <https://doi.org/10.1016/j.pepi.2014.05.011>, <https://doi.org/10.1016/j.pepi.2014.05.011>, 2014.

- 5 Lomax, A. and Michelini, A.: Mwpd: a duration-amplitude procedure for rapid determination of earthquake magnitude and tsunamigenic potential from P waveforms, *Geophysical Journal International*, 176, 200–214, <https://doi.org/10.1111/j.1365-246x.2008.03974.x>, <https://doi.org/10.1111/j.1365-246x.2008.03974.x>, 2009.
- Meng, L., Ampuero, J.-P., and Bürgmann, R.: The 2013 Okhotsk deep-focus earthquake: Rupture beyond the metastable olivine wedge and thermally controlled rise time near the edge of a slab, *Geophysical Research Letters*, 41, 3779–3785, <https://doi.org/10.1002/2014gl059968>, <https://doi.org/10.1002/2014gl059968>, 2014.
- 10 Nakamura, M. and Mochizuki, E.: Focal mechanism solutions and their reliability determined by P-wave first motions (in Japanese), *Quart. J. Seis.*, 67, 11–20, 1988.
- Okal, E. A.: “Detached” deep earthquakes: are they really?, *Physics of the Earth and Planetary Interiors*, 127, 109–143, [https://doi.org/10.1016/s0031-9201\(01\)00224-2](https://doi.org/10.1016/s0031-9201(01)00224-2), [https://doi.org/10.1016/s0031-9201\(01\)00224-2](https://doi.org/10.1016/s0031-9201(01)00224-2), 2001.
- 15 Okal, E. A. and Newman, A. V.: Tsunami earthquakes: the quest for a regional signal, *Physics of the Earth and Planetary Interiors*, 124, 45–70, [https://doi.org/10.1016/s0031-9201\(01\)00187-x](https://doi.org/10.1016/s0031-9201(01)00187-x), [https://doi.org/10.1016/s0031-9201\(01\)00187-x](https://doi.org/10.1016/s0031-9201(01)00187-x), 2001.
- Ozacar, A. A., Beck, S. L., and Christensen, D. H.: Source process of the 3 November 2002 Denali fault earthquake (central Alaska) from teleseismic observations, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2003gl017272>, <https://doi.org/10.1029/2003gl017272>, 2003.
- 20 Pondrelli, S., Salimbeni, S., Morelli, A., Ekström, G., Postpischl, L., Vannucci, G., and Boschi, E.: European–Mediterranean Regional Centroid Moment Tensor catalog: Solutions for 2005–2008, *Physics of the Earth and Planetary Interiors*, 185, 74–81, <https://doi.org/10.1016/j.pepi.2011.01.007>, <https://doi.org/10.1016/j.pepi.2011.01.007>, 2011.
- Reasenber, P. and Oppenheimer, D.: FPFIT, FPLOT and FPPAGE; Fortran computer programs for calculating and displaying earthquake fault-plane solutions, <https://doi.org/10.3133/ofr85739>, <https://doi.org/10.3133/ofr85739>, 1985.
- 25 Rezapour, M. and Pearce, R. G.: Bias in surface-wave magnitude  $M_s$  due to inadequate distance corrections, *Bulletin of the Seismological Society of America*, 88, 43–61, 1998.
- Shearer, P. M.: Evidence from a cluster of small earthquakes for a fault at 18 km depth beneath Oak Ridge, southern California, *Bulletin of the Seismological Society of America*, 88, 1327–1336, 1998.
- Storchak, D. A., Harris, J., Brown, L., Kathrin, L., Shumba, B., Verney, R., Di Giacomo, D., and Korger, E. I. M.: Rebuild of the Bulletin of the International Seismological Centre (ISC), part 1: 1964–1979, *Geoscience Letters*, 4, <https://doi.org/10.1186/s40562-017-0098-z>, <https://doi.org/10.1186/s40562-017-0098-z>, 2017.
- 30 Tsai, V. C., Nettles, M., Ekström, G., and Dziewonski, A. M.: Multiple CMT source analysis of the 2004 Sumatra earthquake, *Geophysical Research Letters*, 32, <https://doi.org/10.1029/2005gl023813>, <https://doi.org/10.1029/2005gl023813>, 2005.
- Vallée, M., Charléty, J., Ferreira, A. M. G., Delouis, B., and Vergoz, J.: SCARDEC: a new technique for the rapid determination of seismic moment magnitude, focal mechanism and source time functions for large earthquakes using body-wave deconvolution, *Geophysical Journal International*, 184, 338–358, <https://doi.org/10.1111/j.1365-246x.2010.04836.x>, <https://doi.org/10.1111/j.1365-246x.2010.04836.x>, 2010.
- Wessel, P., Smith, W. H. F., Scharroo, R., Luis, J., and Wobbe, F.: Generic Mapping Tools: Improved Version Released, *Eos, Transactions American Geophysical Union*, 94, 409–410, <https://doi.org/10.1002/2013eo450001>, 2013.
- Weston, J., Engdahl, E., Harris, J., Di Giacomo, D., and Storchak, D.: ISC-EHB: reconstruction of a robust earthquake data set, *Geophysical Journal International*, 214, 474–484, <https://doi.org/10.1093/gji/ggy155>, <https://doi.org/10.1093/gji/ggy155>, 2018.


- 5 Willemann, R. J. and Storchak, D. A.: Data Collection at the International Seismological Centre, *Seismological Research Letters*, 72, 440–453, <https://doi.org/10.1785/gssrl.72.4.440>, <https://doi.org/10.1785/gssrl.72.4.440>, 2001.
- Ye, L., Lay, T., Kanamori, H., and Rivera, L.: Rupture characteristics of major and great ( $M_w \geq 7.0$ ) megathrust earthquakes from 1990 to 2015: 1. Source parameter scaling relationships, *Journal of Geophysical Research: Solid Earth*, 121, 826–844, <https://doi.org/10.1002/2015jb012426>, <https://doi.org/10.1002/2015jb012426>, 2016.
- Zhan, Z. and Kanamori, H.: Recurring large deep earthquakes in Hindu Kush driven by a sinking slab, *Geophysical Research Letters*, 43, 7433–7441, <https://doi.org/10.1002/2016gl069603>, <https://doi.org/10.1002/2016gl069603>, 2016.
- Zoback, M. L.: First- and second-order patterns of stress in the lithosphere: The World Stress Map Project, *Journal of Geophysical Research*, 97, 11 703–11 728, <https://doi.org/10.1029/92jb00132>, <https://doi.org/10.1029/92jb00132>, 1992.




**Figure 1.** Number of source mechanisms in the ISC Bulletin from January 1964 to ~~October 2018~~ December 2018 and normalized frequency - magnitude distributions with respect to different time periods. A magnitude value for each event with an associated source mechanism is selected following the scheme described in Di Giacomo and Storchak (2015). Note the high peak in the top subplot in 2011 which is associated with the 2011 Tohoku earthquake aftershock sequence. ~~The~~ A slight dip for year 2018 is only apparent as the data collection is not ~~complete~~ yet complete.





**Figure 2.** Time distribution of source mechanisms in the ISC Bulletin (January 1964 – ~~October~~ December 2018) reported by different agencies. The numbers in brackets next to each agency code shows the total number of reported source mechanism solutions. Red lines indicate waveform inversion techniques, blue lines indicate first motion polarity techniques and grey colour shows cases where there is no information available on the techniques being used or we could not verify them. A detailed list of the reporting agencies ~~can be found at~~ is shown in Table 1.



**Figure 3.** Global maps showing the number of events with at least one source mechanism reported in the ISC Bulletin for the time period from January 1964 to ~~October-December~~ 2018, in a  $1^\circ$  by  $1^\circ$  grid, for mechanism solutions reported by local and regional agencies (top), and global agencies (bottom, [[GCMT](#), [HRVD](#), [GCMFIPGP](#), [NEISISC](#), [MOS](#), [NEIC](#), [IPGPNEIS](#)]). Minimum magnitude ( $M_{\min.}$ ) following the scheme described in Di Giacomo and Storchak (2015) is shown on top of each map.




**Figure 4.** [Geographical and frequency - magnitude distribution \(following the scheme described in Di Giacomo and Storchak \(2015\)\) of earthquakes with source mechanisms reported by the agencies which systematically send their mechanism solutions to the ISC. Locations on the maps are colour coded by depth. The agency codes are shown on top of each map and their details can be found in Table 1.](#)



**Figure 5.** Frequency distribution of available source mechanisms per event in the ISC Bulletin for the time period from January 1964 to ~~October~~ December 2018 (top), and intra-event maximum rotation angle frequency distribution for the seismic events having at least two mechanism solutions available (bottom).



Annual number of source mechanisms grouped by Zoback (1992) fault styles. Dark blue circles, red pentagons, green squares, orange triangles, light blue inverted triangles and olive diamonds represent, respectively, thrust, normal, strike-slip, normal with strike-slip component, thrust with strike-slip component and undefined earthquakes. Black octagons are earthquakes for which the intra-variability of the source mechanisms does not allow us to assign an earthquake to a fault style. These are earthquakes with large rotation angles (see Figure 4). The percentage of such earthquakes goes up to 8% of earthquakes per



**Figure 6.** Annual number of source mechanisms grouped by Zoback (1992) fault styles. Dark blue circles, red pentagons, green squares, orange triangles, light blue inverted triangles and olive diamonds represent, respectively, thrust, normal, strike-slip, normal with strike-slip component, thrust with strike-slip component and undefined earthquakes. Black octagons are earthquakes for which the intra-variability of the source mechanisms does not allow us to assign an earthquake to a fault style. These are earthquakes with large rotation angles (see Figure 5). The percentage of such earthquakes goes up to 8% of earthquakes per year. The grey area for year 2018 denotes that the data collection is not yet complete. See main text for details.

**Table 1.** Agencies reporting source mechanism solutions at the ISC for the time period from January 1964 to December 2018. The cross symbol (x) denotes the type of parametric data that is reported from each agency to the ISC.

Agency code	Name	Country	Mag. range	Nodal planes	Moment tensor	Principal axes
AEIC	Alaska Earthquake Information Center	U.S.A.	3.6-5.0		X	
AFAR	The Afar Depression: Interpretation of the 1960-2000 Earthquakes	Israel	3.8-6.6	X		
ATA	The Earthquake Research Center Ataturk University	Turkey	3.9-4.5	X		
ATH	National Observatory of Athens	Greece	3.7-5.5	X		
BER	University of Bergen	Norway	1.1-4.0	X		
BGS	British Geological Survey	United Kingdom	1.2-4.6	X		
BRK	Berkeley Seismological Laboratory	U.S.A.	2.9-7.2	X		
BUD	Geodetic and Geophysical Research Institute	Hungary	2.0-4.7	X	X	X
CASC	Central American Seismic Center	Costa Rica	3.1-6.5	X		
DNK	Geological Survey of Denmark and Greenland	Denmark	2.5-4.6	X		
ECX	Centro de Investigación Científica y de Educación Superior de Ensenada	Mexico	0.9-5.8	X		
FUNV	Fundación Venezolana de Investigaciones Sismológicas	Venezuela	1.9-4.2	X		
GBZT	Marmara Research Center	Turkey	3.7-5.4	X		
GCMT	The Global CMT Project	U.S.A.	4.0-9.1	X	X	X
GRAL	National Council for Scientific Research	Lebanon	1.4-3.3	X		
GTFE	German Task Force for Earthquakes	Germany	2.2-4.6	X		
GUC	Centro Sismológico Nacional, Universidad de Chile	Chile	2.9-4.3	X		
HLW	National Research Institute of Astronomy and Geophysics	Egypt	2.5-4.9	X		X
HYB	National Geophysical Research Institute	India	3.0-3.8	X		
IAG	Instituto Andaluz de Geofísica	Spain	3.2-6.8	X	X	X
IEC	Institute of the Earth Crust, SB RAS	Russia	3.3-6.3	X		X
INET	Instituto Nicaragüense de Estudios Territoriales - INETER	Nicaragua	3.5-6.0	X		
IPGP	Institut de Physique du Globe de Paris	France	5.2-8.2	X		
ISC	International Seismological Centre	United Kingdom	3.6-9.1	X		X
ISN	Iraqi Meteorological and Seismology Organisation	Iraq	5.0	X		
JMA	Japan Meteorological Agency	Japan	1.6-8.3	X		X
JSN	Jamaica Seismic Network	Jamaica	2.5-5.2	X		
LDO	Lamont-Doherty Earth Observatory	U.S.A.	3.5-4.6	X	X	
MED RCMT	MedNet Regional Centroid - Moment Tensors	Italy	3.6-6.9	X	X	X
MEX	Instituto de Geofísica de la UNAM	Mexico	2.9-7.0	X		
MOS	Geophysical Survey of Russian Academy of Sciences	Russia	5.1-9.1	X		X
NAO	Stifelsen NORISAR	Norway	2.5	X		
NCEDC	Northern California Earthquake Data Center	U.S.A.	3.0-5.2		X	
NDI	National Centre for Seismology of the Ministry of Earth Sciences of India	India	3.5-7.7	X		
NEIC	National Earthquake Information Center	U.S.A.	2.4-9.1	X	X	X
NEIS	National Earthquake Information Service	U.S.A.	5.3-7.7	X	X	X
NIED	National Research Institute for Earth Science and Disaster Prevention	Japan	3.1-9.1	X	X	
NSSC	National Syrian Seismological Center	Syria	1.6-4.6	X		
OSPL	Observatorio Sismológico Politécnico Loyola	Dominican Republic	2.2-5.8	X		
OTT	Canadian Hazards Information Service, Natural Resources Canada	Canada	3.6-7.8	X		
PAS	California Institute of Technology	U.S.A.	3.3-7.2	X	X	X
PGC	Pacific Geoscience Centre	Canada	3.5-6.6	X	X	X
PNSN	Pacific Northwest Seismic Network	U.S.A.	0.0-6.8	X		X
PRE	Council for Geoscience	South Africa	2.2-6.5	X		
REN	MacKay School of Mines	U.S.A.	3.3-4.2		X	
ROM	Istituto Nazionale di Geofisica e Vulcanologia	Italy	3.1-6.6	X	X	
RSNC	Red Sismológica Nacional de Colombia	Colombia	1.6-6.3	X		
SEA	Geophysics Program AK-50	U.S.A.	3.2-3.3		X	
SJA	Instituto Nacional de Prevención Sísmica	Argentina	2.6-5.9	X		
SLM	Saint Louis University	U.S.A.	2.2-6.0	X	X	X
SNET	Servicio Nacional de Estudios Territoriales	El Salvador	1.3-5.6	X		
SSS	Centro de Estudios y Investigaciones Geotécnicas del San Salvador	El Salvador	3.6-6.0	X		
THE	Department of Geophysics, Aristotle University of Thessaloniki	Greece	4.2-6.1	X		
THR	International Institute of Earthquake Engineering and Seismology (IIEES)	Iran	4.5-6.1	X		
TUL	Oklahoma Geological Survey	U.S.A.	2.9-3.7		X	
UAF	Department of Geosciences	U.S.A.	0.5-5.2	X	X	
UCDES	Department of Earth Sciences	United Kingdom	5.5-6.7	X		
UCR	Sección de Sismología, Vulcanología y Exploración Geofísica	Costa Rica	2.6-6.0	X		
UESG	School of Geosciences	United Kingdom	5.7-6.2	X		
UPA	Universidad de Panama	Panama	1.8-6.6	X		
UPIES	Institute of Earth and Environmental Science	Germany	4.3-6.4	X	X	
UPSL	University of Patras, Department of Geology	Greece	3.3-6.8	X		
USGS	United States Geological Survey	U.S.A.	3.2-7.8	X	X	X
UOSS	The University of Utah Seismograph Stations	U.S.A.	3.3-3.8		X	