



Present-day surface deformation of the Alpine Region inferred from geodetic techniques

Laura Sánchez¹, Christof Völksen², Alexandr Sokolov^{1,2}, Herbert Arenz¹, Florian Seitz¹

5 ¹Technische Universität München, Deutsches Geodätisches Forschungsinstitut (DGFI-TUM), Arcisstr. 21, 80333 München, Germany

²Bayerische Akademie der Wissenschaften, Erdmessung und Glaziologie, Alfons-Goppel-Str. 11, 80539 München, Germany

Correspondence to: Laura Sánchez (lm.sanchez@tum.de)

10 **Abstract.** We provide a present-day surface-kinematics model for the Alpine region and surroundings based on a high-level data analysis of about 300 geodetic stations continuously operating over more than 12 years. This model includes a deformation model, a continuous surface-kinematic (velocity) field, and a strain field consistently assessed for the entire Alpine mountain belt. Special care is given to the use of the newest GNSS processing standards to determine high-precise 3D station coordinates. The coordinate solution refers to the reference frame

15 IGB08, epoch 2010.0. The mean precision of the station positions at the reference epoch is ± 1.1 mm in N and E and ± 2.3 mm in height. The mean precision of the station velocities is ± 0.2 mm/a in N and E and ± 0.4 mm/a in the height. The deformation model is derived from the pointwise station velocities using a geodetic least-squares collocation approach with empirically determined covariance functions. According to our results, no significant horizontal deformation is detected in the Western Alps, while across the Southern and Eastern Alps the deformation vectors describe a progressive eastward rotation toward Pannonia. This kinematic pattern makes also evident an increasing magnitude of the deformation from 0.1 mm/a in the western part of Switzerland up to about

20 1.5 mm/a in the Austrian Alps. The largest shortenings are observed along the southern front of the Eastern Alps (in the northern area of the Venetian-Friuli Basin) and in the northern part of the Apennine Peninsula, where they reach 2 mm/a and 3 mm/a, respectively. The averaged accuracy of the horizontal deformation model is ± 0.2 mm/a.

25 Regarding the vertical kinematics, our results clearly show an still on-going averaged uplift of 1.8 mm/a of the entire mountain chain, with exception of the southern part of the Western Alps, where no significant uplift (less than 0.5 mm/a) is detected. The fastest uplift rates (more than 2 mm/a) occur in the central area of the Western Alps, in the Swiss Alps and in the Southern Alps in the boundary region between Switzerland, Austria and Italy. The general uplift observed across the Alpine mountain chain decreases toward the outer regions to stable values

30 between 0.0 and 0.5 mm/a and, in some cases, to subsidence like in the Liguro-Provençal and Vienna Basins, where vertical rates of -0.8 mm/a and -0.3 mm/a are observed respectively. In the surroundings, three regional subsidence regimes are identified: the Rhone-Bresse Graben with -0.8 mm/a, the Rhine Graben with -1.3 mm/a, and the Venetian-Friuli Basin with -1.5 mm/a. The estimated uncertainty of our vertical motion model across the Alpine mountain belt is about ± 0.3 mm/a. The strain field inferred from the deformation model shows two main

35 contrasting strain regimes: shortening across the south-eastern front of the Alps and the northern part of the Dinarides, and extension in the Apennines. The pattern of the strain principal axes indicates that the compression directions are more or less perpendicular to the thrust belt fronts, reaching maximum values of $20 \times 10^{-9} \text{ a}^{-1}$ in the Venetian-Friuli and Po Basins. Across the Alpine mountain belt, we observe a slight dilatation regime in the Western Alps, which smoothly changes to a contraction regime in West Austria and South Germany, reaching



maximum shortening values of $6 \times 10^{-9} \text{ a}^{-1}$ in the north-eastern Austria. The numerical results of this study are available at <https://doi.pangaea.de/10.1594/PANGAEA.886889>.

Keywords. Surface kinematics, deformation, strain field, vertical motion, GNSS velocities, Alpine region.

1. Introduction

- 5 The Alps orogeny is the result of the continental collision of the African and European plates, more precisely of the push of the African plate through the Adriatic Promontory (also called Adria or Apulian microplate; Channell et al., 1979) against Central Europe. A wide-range of multidisciplinary studies based on geologic, geophysical and geodetic data have demonstrated that the Alpine orogeny is active from the Jurassic until the Present (e.g.; Dewey et al., 1973, 1989; Channell and Horváth, 1976; Dercourt et al., 1986; Le Pichon et al., 1988; Mueller and Kahle, 1993; Ustaszewski et al., 2008; Handy et al., 2010, 2015). The tectonic evolution and present geodynamic framework have been primarily deduced from seismotectonic syntheses, and it is clear that the Alpine area and, in a more extended sense, the Mediterranean represent a variety of lithosphere blocks of different age, thickness and rheology leading to complex kinematic processes, including subduction, back-arc spreading, rifting and thrust, reverse and strike-slip faulting. These processes are superimposed on those phenomena associated to any orogeny; i.e., uplift, deformation, erosion, metamorphism, foredeep basins changes, etc. During the last three decades, the Global Navigation Satellite Systems (GNSS), like GPS (Global Positioning System) and GLONASS (GLObalnaja NAWigazionnaja Sputnikowaja Sistema) became a fundamental tool to observe, model and understand present-day kinematic processes, as they provide precise geodetic constraints with a high spatial resolution allowing the improvement of geodynamic models, especially at diffuse plate boundaries like the African-Eurasian boundary.
- 10 15 20 25
- The use of GPS to detect tectonic surface deformations in Europe started already in the 1990s (e.g., Fejes et al., 1993; Kahle et al. 1994; Chéry et al., 1995; Kaniuth et al., 1995, 1999; van Mierlo et al., 1996; Calais et al., 1998, 2000; Ferhat et al., 1998; Calais 1999; Grenerczy et al., 2000; Sue et al. 2000). Given the complexity of the deformation zone in the southern part of the continent, scientists concentrate on selected areas to deploy observing stations and to develop local deformation models. Recent studies can be classified following the geographic subdivision of the Alps into Western, Central, Eastern and Southern Alps (Fig. 1):
- the Western Alps in south-eastern France, the boundary region France-Italy and south-western Switzerland; e.g., Tesauro et al. (2006), Delacou et al. (2008), Larroque et al. (2009), Nguyen et al. (2016), Nocquet et al. (2016);
 - the Central/Eastern Alps in Austria and the adjacent regions of Switzerland, Liechtenstein, Germany, Italy and Slovenia; e.g., Haslinger et al. (2006), Weber et al., (2006), Gosar et al. (2007), Sue et al. (2007), Brückl et al. (2010), Brockmann et al. (2012);
 - the junction zone of the Eastern Alps, the Dinarides and the Pannonian Basin: Bada et al. (2007), Bus et al. (2009), Caporali et al. (2009), Möller et al. (2011);
 - the Southern Alps in northern Italy (also considered as a boundary zone for the computation of deformation models in the Adriatic region): Grenerczy et al. (2005), Grenerczy and Kenyeres (2006), D'Agostino et al. (2005; 2011), Devoti et al. (2008; 2011), Serpelloni et al. (2005; 2013). Cuffaro et al. (2010), Métois et al. (2015).
- 30 35



These studies describe in general deformation models of regional scope and cover isolated segments of the Alpine mountain chain and its forelands. The objective of this work is the consistent determination of a continuous horizontal and vertical surface-kinematic field of the Alpine area that provides an integral picture of the on-going deformation processes in the entire region. A network of about 300 continuously operating GNSS (CO-GNSS) stations with observations collected over 12.4 years is used for the precise determination of station positions and velocities. Based on these results, a continuous kinematic field is derived using a geodetic least-squares collocation (LSC) approach with empirically determined covariance functions. Main results are a deformation model, a continuous surface-kinematic (velocity) field, and a strain field consistently assessed for the entire Alpine mountain belt. The core contribution of this work is the homogeneous analysis of a large number of freely available data from CO-GNSS stations located in the Alpine region, using unified processing standards and a common reference frame for the complete time-span covered by the observations. Special care is given to the application of the newest GNSS processing standards like absolute corrections to the GNSS antenna phase centre variations, troposphere delay estimation based on numerical weather models, atmospheric and oceanic tide loading effects, etc. To ensure consistency, we also use reprocessed GNSS satellite orbits referring to the IGS08/IGb08 reference frame (Rebischung et al. 2012) and including the most recent models adopted by the International GNSS Service (IGS) for the orbit determination, such as the albedo model of Rodríguez-Solano et al. (2012), the satellite antenna thrust effect, and the a-priori solar radiation pressure model switch-off (Steigenberger et al. 2014). The IGS08/IGb08 reference frame corresponds to a subset of GNSS stations of the ITRF2008 (International Terrestrial Reference Frame 2008, Altamimi et al., 2011), which are used by the IGS as fiducial points for the computation of the satellite ephemeris and the Earth orientation parameters (EOP).

In this context, after a brief description of the geological and tectonic framework of the Alpine region, the third section of this paper provides details about the geographical distribution of the GNSS stations, the availability of the GNSS observations, and the data sources. Afterwards, we present in Section 4 the GNSS data processing strategy, the geodetic datum realisation, the analysis of the position time series, and the computation of a cumulative (multi-year) solution for the estimation of precise station positions and velocities. Section 5 concentrates on the computation of the kinematic model: correlations between the station velocities and the tectonic settings are analysed, the modelling methodology is presented, and the vertical and horizontal deformation models are briefly described. Finally, the strain field computation is summarised in Section 6. The reliability of the results obtained in this study is validated by contrasting them with the deformation patterns and the displacement rates published in the existing literature. With this work, we are providing a present-day surface kinematic model for the Alpine region based on a high-level GNSS data processing.

2. Geological and tectonic framework

We focus on the estimation of the present-day kinematics in the Alps and surrounding areas based on GNSS measurements gained during the last decade. Previous processes associated to the tectonic evolution and their effects are usually deduced from geophysical modelling, but they are beyond the scope of this work. The main tectonic structures underlying the Alps and its forelands are the Eurasian plate in the North and the West, the Adriatic and Ligurian microplates in the South, and the Pannonian fragment in the East (Fig. 1). The boundaries between these units are mainly inferred from (natural and controlled) seismic data and tomographic inversion; see details in Pfiffner et al. (1990), Waldhauser et al. (1998), Doglioni and Carminati (2002), Lippitsch et al. (2003),



Kissling et al. (2006), Brückl et al. (2007). The push of the African plate against the Eurasian plate causes an under-thrusting motion beneath the Southern Alps as indicated, for instance, by the Friuli seismicity. In addition, the Adriatic Promontory rotates counter-clockwise, indenting the Eastern Alps and Dinarides in the North (e.g., Nocquet and Calais, 2004; Handy et al., 2015) and subducting beneath the Hellenides in the South (Greneczy et al., 2005; Bennett et al. 2008). The polarity of subduction changes from one area to the other: while the European lithosphere is subducted beneath the Alps, the African lithosphere is subducted beneath the Apennines and the Dinarides-Hellenides (Argnani, 2009). In this way, Alps, Apennines, and Dinarides-Hellenides are still active orogens, although at different rates (D'Agostino et al., 2005; Devoti et al. 2008; Cuffaro et al., 2010; Cheloni et al., 2014). Each subduction is associated with the usual vertical motions; i.e., subsidence in the foreland basin and uplift in the mountain belt. The continuous curved orogenic belt built by the Alps together with the Carpathians and the Dinarides-Hellenides bifurcates and encircles the Pannonian Basin system (Doglioni, 1987; Royden, 1993, Doglioni and Carminate, 2002; Horváth et al., 2006). South of the Alps are the Po and the Venetian-Friulian Basins. As mentioned, these three basins: Pannonian, Po and Venetian-Friulian Basins, are still foreland active basins; i.e., they continue subsiding and receiving sediments (e.g., Cuffaro et al., 2010; Serpelloni et al., 2006; 2013). The Po Basin is also part of the northern foreland basin of the Apennines (e.g., Laubscher, 1988; Argnani, 2009). In the west of the Apennines and south of the Western Alps, we find the Liguro-Provençal and the Tyrrhenian Basins. They are separated by the Corso-Sardinia block (Serpelloni et al., 2006; 2013). The northern foreland basin of the Alps is the Molasse Basin (Ziegler, 1990; Kuhlemann and Kempf, 2002). It runs between the Jura Mountains and the Alps in the north-western part of Switzerland and continues along the Bavarian and Austrian Alps up to Vienna. In contrast to the Pannonian and Po Basins, it is not active as foreland basin; on the contrary, large parts of it are being uplifted and eroded (Becker, 1999). On the West, the Alps are boarded by the Rhone-Bresse Graben. In general, it can be stated that the deformation in the Alpine region is dominated by the counter-clockwise motion of the Adria microplate, causing compression in the Eastern Alps, dextral shear in the Central Alps and a very slow deformation in the Western Alps.





Figure 1: Simplified tectonic map of the Alps and surrounding areas showing the major tectonic features and faults, modified from Schmid et al. (2004) and Bousquet et al. (2012). Geographic subdivision of the Alps: WA – Western Alps, CA – Central Alps, EA – Eastern Alps, SA – Southern Alps. Tectonic plates/blocks: EU – Eurasia, ADR – Adria, LIG – Liguria,



PAN – Pannonia, CS – Corso-Sardinia. Basins: LP – Liguro-Provençal, TS – Tyrrhenian, PO – Po, VF – Venetian-Friuli, MB – Molasse, PB – Pannonian, SB – Styrian, VB – Vienna. Graben: RBG – Rhone-Bresse, RHG – Rhine. Mountains and mountain-belts: AP – Apennines, PY – Pyrenees, DI – Dinarides-Hellenides, JM – Jura, MC – Massif Central. Topography and bathymetry after the ETOPO1 model (Amante and Eakins, 2009).

5 3. Distribution of the CO-GNSS stations

The determination of the surface kinematics in the Alpine region is based on the observations collected at 306 CO-GNSS stations (Fig. 2). Stations located at distances up to some thousand kilometres from the Alps are included to improve the geometry of the network and to ensure a reliable geodetic datum realisation. Several sites are part of the global IGS network (Dow et al., 2009), the EUREF Permanent Network (EPN) (Bruyninx et al., 2012, 2013) and the *Integriertes Geodätisches Referenznetz Deutschlands* (GREF). EUREF stands for the Reference Frame Sub-Commission for Europe of the International Association of Geodesy (IAG). The CO-GNSS station distribution along the Alpine mountain belt is densified with more than 30 GAIN (Geodetic Alpine Integrated Network) stations. Five of these stations were installed and are maintained by DGFI-TUM (Seitz et al., 2014). The GAIN network was established in the frame of the ALPS-GPSQUAKENET project, which was funded by the European Union under the INTERREG IIIB Alpine Space program from 2004 to 2007 (Aoudia et al., 2007). Additional data from CO-GNSS sites in Austria were provided by the Space Research Institute (*Institut für Weltraumforschung*, Graz) of the Austrian Academy of Sciences (Haslinger and Strangle, 2006). Another valuable source of data for this study is the Friuli Deformation Network (FreDNet) established in 2002 (Zuliani et al., 2002, Rossi et al. 2013). While FreDNet densifies the station distribution in the Eastern Alps, the Western Alps are very well covered by CO-GNSS sites of the RENAG (*Réseau National GPS*, France, Nocquet et al., 2016) and Orpheon networks. Orpheon is a geodetic network providing real time and augmentation services in France. The CO-GNSS network processed in this study does not include stations in Switzerland. However, to get a homogeneous data coverage in the Central Alps, we use IGB08-based station velocities of the Automated GNSS Network for Switzerland (AGNES) provided by Brockmann et al. (2012). Figure 2 shows the station distribution according to the data sources.



5 **Figure 2: CO-GNSS stations processed for the estimation of the surface-kinematics in the Alps.** The stations of the contributing CO-GNSS arrays are color-coded. Topography and bathymetry after the ETOPO1 model (Amante and Eakins, 2009).

The GNSS data cover the time-span from 2004-01-01 to 2016-05-30. However, the amount of the available GNSS observations varies over the years (Fig. 3). The period with the maximum number of active stations within the network is between 2012 and 2015, reaching nearly 180 CO-GNSS stations every day. The total number of the daily observation files processed in this study is about 0.57 million.




Figure 3: Amount of daily observation files processed in this study.

4. Analysis of GNSS data and determination of precise station positions and velocities

15 Effects of on-going geodynamic processes may be inferred from crustal movements detected by the analysis of repeated (or continuous) geodetic observations. The basic idea is to determine (and compare) the geometry of the network formed by the geodetic stations at certain epochs. The changes of the network geometry over large time-spans (e.g. years) are interpreted as a deformation of the network caused by geodynamic processes underlying the studied area. To ensure a high reliability in the determination of the network geometry, the geodetic measurements



5 must be processed over the entire time-span following strict standards and procedures. In this study, we apply therefore the same analysis strategy usually applied for the establishment of GNSS-based regional geodetic reference frames like SIRGAS (*Sistema de Referencia Geocéntrico para las Américas*, e.g. Sánchez et al., 2013; Sánchez and Drewes 2016) or the EUREF Permanent Network (EPN) as a realisation of the European Terrestrial Reference System (ETRS); e.g., Völksen (2009a; 2009b). The data analysis is based on the conventions outlined by the International Earth Rotation and Reference Systems Service (IERS) for the determination of the ITRF (Petit and Luzum 2010) and the GNSS-specific guidelines defined by the IGS for the precise processing of the global GNSS network (IGS, 2014).

10 In a first processing step, free weekly normal equations (NEQ) are computed. *Free* means in this context that the geometry of the GNSS network is fully consistent with the GNSS orbits, but the network's origin and orientation are not constrained yet. They will be constrained in a posterior processing step, when the geodetic datum for the GNSS network is introduced. The free weekly NEQ covering the complete time-span from January 2004 to May 2016 are combined to a multi-year solution with constant velocity for each station. This includes an iterative time-series analysis to detect outliers and discontinuities in the weekly station positions. In the multi-year solution, the geodetic datum is realised with respect to the coordinates (positions and velocities) of selected reference stations. 15 The station velocities represent the mean displacement per year of the CO-GNSS stations and they are the primary input data to model the surface kinematics. The following sections describe the key aspects of the GNSS data analysis performed in this study.

4.1 GNSS network weekly solutions

20 The processing of the daily GNSS observations is accomplished using the double-difference baseline approach and the least-squares adjustment implemented in the Bernese GNSS Software V.5.2 (Dach et al. 2015). The main processing characteristics are:

- GPS and GLONASS observations at a 30-s sampling rate are considered.
- The basic observable is the L3 ionosphere-free linear combination.
- 25 - The satellite orbits, satellite clock offsets, and EOPs are fixed to the values generated by the IGS processing centre CODE (Centre for Orbit Determination in Europe, Dach et al., 2017; Steigenberger et al., 2014; <ftp://ftp.aiub.unibe.ch/aiub/CODE/>). To ensure consistency, reprocessed (2004-2013) and final CODE (2014-2016) products based on the reference frame IGS08/IGb08 are used for the analysis.
- The antenna phase centre offsets and direction-dependent phase centre corrections for both satellite 30 transmitting and ground receiving antennas are taken from the model `igs08.atx` (Schmid et al., 2016; http://igsceb.jpl.nasa.gov/igsceb/station/general/pcv_archive/).
- The tropospheric zenith delay is modelled using the Global Mapping Function (GMF, Böhm et al., 2006). Here, the a-priori zenith hydrostatic delay (ZPD) values are derived from gridded coefficients based on the Global Pressure Temperature (GPT) model (Böhm et al., 2006) and are refined by computing zenith-wet delays (ZWD) in a 1-hour interval (also using GMF). In addition, to model azimuthal asymmetries, 35 horizontal gradient parameters are estimated using the model described by Chen and Herring (1997).
- Corrections for the solid Earth tide, permanent tide, and solid Earth pole tide are applied as described in Petit and Luzum (2010). The ocean tide loading is estimated with the FES2004 model (Letellier, 2004) and the atmospheric tide loading caused by the semidiurnal constituents S1 and S2 is estimated following 40 the model of van Dam and Ray (2010). The coefficients for the ocean tide loading are provided by M.S.



Bos and H.-G. Scherneck at <http://holt.oso.chalmers.se/loading/>. The coefficients for the atmospheric tide loading are provided by T. van Dam at <http://geophy.uni.lu/ggfc-atmosphere/tide-loading-calculator.html>.

- 5 - Non-tidal loadings such as atmospheric pressure, ocean bottom pressure, or surface hydrology are not reduced.
- The seven daily free normal equations corresponding to a week are combined to generate a free normal equation per week. In this combination, stations with large residuals in any daily normal equation (more than ± 10 mm in the horizontal or more than ± 15 mm in the vertical) are removed from the corresponding daily equation and the weekly combination is recomputed.


10 4.2 Determination of precise station positions and velocities

The determination of the station coordinates is performed within a multi-year solution. In a first step, all free weekly NEQ available from January 2004 to May 2016 are combined and solved using the program ADDNEQ2 of the Bernese GNSS Software V.5.2 (Dach et al., 2015). Based on this preliminary solution, in a second step, residual time-series for each station are generated to identify outliers or discontinuities that may mislead the
15 computation of the station velocities. In this study, we use as thresholds ± 10 mm for the northeast component and ± 20 mm for height (about fourfold the mean RMS). Sporadically occurring outliers are removed from the respective weekly NEQ. If outliers reflect a discontinuity (i.e. they appear in successive NEQ), a new position is estimated for this station. Most of these discontinuities are caused by the replacement of antennas and receivers, either because of defects or to modernise the equipment. In some cases, the discontinuities are caused by seismic
20 effects and they may be followed by changes in the linear movement of the station; thus new position and velocity values must be estimated for the affected stations, see an example in Fig. 4. Once outliers are removed and discontinuities are identified, a new multi-year solution is computed, new residual time-series are generated and the time-series analysis is repeated to identify remaining outliers or discontinuities. This process is iteratively performed until no more outliers or discontinuities remain. The geodetic datum is realised by imposing two
25 minimum constraint conditions with respect to the IGB08 coordinates of the reference stations. The no-net translation (NNT) condition ensures that the origin [$X=Y=Z=0$] of our network coincides with the origin of the network built by the reference stations. The no-net-rotation (NNR) condition aligns the orientation of our network to the orientation defined by the network built by the reference stations. In this study, we selected six IGB08 sites (WTZR, ZIMM, GRAS, GRAZ, MEDI, LROC; Fig. 7) as reference stations (see Rebeschung et al. 2012; <ftp://igs-rf.ign.fr/pub/IGb08/IGb08.snz>). The main selection criteria is based on the homogenous distribution of the
30 reference stations across the CO-GNSS network, rare incidents of equipment changes, and a nearly complete coverage of the time-series from January 2004 to May 2016.


In general, the station position time-series describe clear seasonal motions (Fig. 5), which are assumed to be a consequence of non-modelled geophysical loading or local effects superimposed with GNSS inherent systematic
35 errors (e.g., Blewitt et al., 2001; Collilieux et al., 2010, 2012; Drewes et al., 2013; King et al., 2008; Ray et al., 2008). Therefore, to increase the reliability of the constant velocities required for modelling the surface kinematics, stations with time-series shorter than two years are excluded from further computations. The station position time-series of the entire network present an averaged RMS value of about ± 1 mm in the horizontal component, and ± 2.5 mm in the height (Fig. 6). The final multi-year solution for the CO-GNSS network covering the Alpine region
40 refers to the IGB08, epoch 2010.0. While the averaged RMS precision for the station positions at the reference



epoch is ± 1.1 mm in N and E and ± 2.3 mm in height, the averaged RMS precision for the station velocities is ± 0.2 mm/a in N and E and ± 0.4 mm/a in height. Figures 7 and 8 show the horizontal station velocities (after removing the Eurasian plate motion) and the vertical station velocities, respectively.



5 **Figure 4: Position time-series of the station BOLG located in Bologna, Italy.** Discontinuities (indicated by blue vertical lines) are caused by seismic events. Piecewise sinusoidal lines in light green represent a functional model approximating the seasonal motions detected at the station.



10 **Figure 5: Position time-series of the station KARL located in Karlsruhe, Germany.** Annual signals are well observed in the North and East components. Equipment change (indicated by the blue vertical lines) seems to be not affecting the time-series.




Figure 6: Mean RMS values of the station position time-series for the period January 2004 to May 2016.

4.3 Velocity solution

- 5 According to the results presented in Fig. 7, we observe in the Alps-Pannonian-Dinarides junction zone horizontal velocities in the range 1 - 4 mm/a relative to the stable part of the Eurasian plate. These large relative velocities are controlled by the north-eastward motion of the Adria plate towards the Southern and Central Alps and the Dinarides. Based on our few CO-GNSS sites in the Apennines, we see apparently an extension of 2 - 4 mm/a across these mountains and a 0.5 - 1 mm/a shortening across the southern front of the Eastern Alps. A progressive
- 10 eastward rotation of the velocities toward the Pannonian Basin is also detected. The relative velocities of the stations located in the Western Alps and surrounding areas like the Po Basin, the Rhone-Bresse Graben and southern France are very small (0.1 - 0.4 mm/a), allowing to infer a very slow horizontal deformation in this region. Nocquet et al. (2016) call this a *virtually zero horizontal velocity*. As expected, stations situated in the stable part of the Eurasian plate show velocities around zero; some exceptions reflect mostly local effects.
- 15 While the motion of the Eurasian plate dominates the horizontal velocities, the vertical velocities (Fig. 8) mirror in general the orogenic processes in the Alpine region: uplift of the mountain chains and subsidence of the active foreland basins. A wide area of the Western Alps presents an uplift rate of approximately 2.2 mm/a decreasing westwards. In fact, stations located west of the Alpine foreland show zero vertical rates. The southern part of the Eastern Alps shows a slower vertical uplift rate of about 1.0 - 1.5 mm/a, which decreases towards the North. The
- 20 CO-GNSS stations located in the northern margin of the Alps in Germany and Austria present uplift rates of less than 0.6 mm/a decreasing eastwards. Subsidence rates of about 1 mm/a are observed in the Rhine and Rhone-Bresse Grabens as well as in the Venetian-Friuli Basin. They increase southwards and reach up to 2 mm/a at the coasts in Marseille and Venice, respectively.




Figure 7: Horizontal station velocities relative to the Eurasian plate with 95% error ellipses. Labels identify the reference stations used for the geodetic datum definition: GRAS (Caussols, France), GRAZ (Graz, Austria), LROC (La Rochelle, France), MEDI (Medicina, Italy), WTZR (Bad Koetzing, Germany), ZIMM (Zimmerwald, Switzerland). Topography and bathymetry after the ETOPO1 model (Amante and Eakins, 2009).

5

In general, it can be stated that the vertical velocities shown in Fig. 8 are a result of the combination of three main mechanisms: (i) uplift caused by the ongoing convergence of Eurasia and Africa; (ii) orogenic gravitational movement due to erosion, collapse and associated isostatic compensation; and (iii) isostatic rebound caused by the melting of the Alpine ice shields since the last glacial maximum 22,000 years ago; i.e., Glacial Isostatic Adjustment (GIA). The orogenic gravitational movement is much slower than the GIA; however, it is clear that the eroded rock masses are redistributed to the surroundings with lower elevation (delta valleys and sedimentary basins). This may cause a bedrock uplift and surface height lowering in the mountains as well as a bedrock subsidence and a regional surface height increase in the foreland basins. According to Persaud and Pfiffner (2004), Barletta et al. (2006), and Stocchi et al. (2005), the crustal uplift caused by GIA in the Alpine region varies from 0.06 to 0.20 mm/a. This assessment suggests that the first two mechanisms (plate convergence and orogenic gravitational movement) are the main contributors to the vertical rates obtained in this study. According to this, and recognizing that the GIA effects are smaller than the estimated accuracy of the vertical velocities, we assume GIA effects so far negligible.

10

15




Figure 8: Vertical station velocities: Red arrows indicate uplift, blue arrows represent subsidence, green bars show 95% confidence. Labels identify the reference stations used for the geodetic datum definition: GRAS (Caussois, France), GRAZ (Graz, Austria), LROC (La Rochelle, France), MEDI (Medicina, Italy), WTZR (Bad Koetzing, Germany), ZIMM (Zimmerwald, Switzerland). Topography and bathymetry after the ETOPO1 model (Amante and Eakins, 2009).

5

5. Surface-kinematic modelling based on CO-GNSS network solutions

Based on the precise station velocities estimated previously, the objective of this section is to describe the surface-kinematic modelling performed within this study. We distinguish three main components: a deformation model, a continuous velocity field, and a strain field. A common trend motion is removed from the CO-GNSS station velocities to obtain the so-called residual velocities. The common trend motion represents a large-scale variation, while the residual velocities represent short-scale variations. The pointwise residual velocities are correlated with the existing tectonic structures (Fig. 1) and then, they are interpolated to a regular grid. The interpolation of the residual velocities provides us with the deformation model (sections 5.2 and 5.3), which is the basis for the computation of the strain field (section 6). The common trend motion removed from the initial station velocities is restored to the deformation model to get a continuous velocity field (section 5.4). Various geodetic and geophysical methods based on mathematical interpolation approaches or physical crust models may be applied to represent the continuous surface deformation from pointwise velocities. A frequently used interpolation method is the least-squares collocation (e.g. Moritz 1973, Drewes 1978); to include the physical properties of the deforming body, the finite element method is an appropriate tool (e.g. Meissl 1981, Heidbach and Drewes 2003). It has been demonstrated that for the sole representation of the Earth surface deformation, the results of both methods are nearly identical (e.g. Drewes and Heidbach 2005). Therefore, we use here the least-squares collocation method.

10

15

20

5.1 Least-squares collocation (LSC) approach



The primary principle of LSC is the correlation of physical parameters (in this case, station velocities) between adjacent points. The observations are divided into a systematic linearized part (or trend) and two independent random parts: the signal and the observation error (or noise). The parameters describing the systematic component and the stochastically correlated signals are estimated by minimising the noise. The estimated parameters and signals together with a given spatial correlation allow the prediction (interpolation) of *fictional measurements* at regions where no observations exist. In spatial applications, the correlation function is usually considered as a function dependent on the distance. The so-called correlation distance d should be in agreement with the spacing of the stations and the wavelength of the expected signal. If d is smaller than the mean station spacing, artefacts may appear in the interpolation. If d is too large, a smoothing similar to a filtering effect may occur. According to the station distribution in our network (Fig. 2), we use 100 km as the maximum distance and a grid size of 25 km x 25 km for the interpolation. This grid size is chosen as appropriate since it corresponds to the mean station spacing in the most densely covered region within the network (i.e., the French and Swiss Alps). A larger grid size would average out the results and this filtering should be avoided. In areas with poor station coverage, the nearest four stations are used for the LSC prediction.


According to the tectonic plate boundary model PB2002 (Bird, 2003), we assume that the Adria, Pannonia, and Liguria blocks constitute a deformation zone along the southern Eurasian plate boundary. We do not consider small lithospheric blocks rotating with independent Euler vectors, but a continuous and viscous lithosphere deforming under certain kinematic boundary conditions as suggested, for instance, by Flesch et al. (2000); Vergnolle et al. (2007); or Copley (2008). However, as horizontal and vertical station motions are dominated by different mechanisms (see Section 4.3), the LSC procedure is applied separately: as a vector prediction for the horizontal component and as a scalar prediction for the height component. In the empirical procedure, the geocentric Cartesian station positions (X, Y, Z) and velocities (v_x, v_y, v_z) obtained within the multi-year solution (Section 4.2) are transformed together with their precision values into ellipsoidal coordinates (φ, λ, h) and $(v_\varphi, v_\lambda, v_h)$. The station velocities are reduced by means of the Eurasian plate motion. The corresponding rotation vector Ω ($260.74^\circ \pm 0.53^\circ\text{E}$, $55.14^\circ \pm 0.27^\circ\text{N}$, 0.2598 ± 0.0011 °/Ma) is inferred from our CO-GNSS stations located on the stable part of the plate and it is nearly identical with the rotation vector derived from the ITRF2008 (Drewes, 1982; 2009). Finally, the empirical covariance functions are determined using the approach described by Drewes and Heidbach (2005) and further applied in Sánchez and Drewes (2016). In this procedure, outliers with discrepancy larger than 0.5 mm/a with respect to the velocities of the surrounding stations are considered to be affected by local effects and they are not included in the collocation.

5.2 Horizontal deformation model

The deformation model presented in Fig. 9 describes the surface kinematics of the Alpine region with respect to the Eurasian plate. Two distinct kinematic patterns can clearly be seen: the Western Alps as well as the western and northern surrounding areas (Liguro-Provençal Basin, Massif Central Mountains, Rhone-Bresse Graben, Jura Mountains and Molasse Basin) show a small and statistically insignificant motion, indicating that these regions move north-eastward with the Eurasian plate. This stationary pattern contrasts with the pronounced N- and NE-oriented movement predicted in the Southern and Eastern Alps as well as in the Apennine Peninsula, the Venetian-Friuli Basin and the northern part of the Dinarides. It is also evident that the magnitude of deformation is increasing



5 from 0.1 mm/a in the western part of Switzerland to the South and to the East, reaching 3.0 mm/a in the northern part of the Apennine Peninsula and about 1.5 mm/a in the Austrian Alps, respectively. The two kinematic patterns are apparently separated by the so-called Periadriatic line, continued by the Tonale line to the West and the Brenner fault to the East. They confirm a fault system in a WE-direction along the boundary between the Southern Alps and the Adriatic region (around latitude 46°N, see Fig. 1). The magnitude of the deformation vectors across the westernmost part of the Po Basin and the south of the Western Alps is less than 0.2 mm/a. The averaged residual motion with respect to the Western Alps is not larger than 0.1 mm/a, indicating that presently there is no shortening in this region. These findings are confirmed by Nocquet et al. (2016), who suggest an upper bound of 0.3 mm/a (with 95% confidence) for possible right-lateral strike slip motion in the western Alpine area.



10 **Figure 9: GNSS-inferred horizontal deformation model of the Alpine region.** Red arrows represent the predicted surface deformation. Blue arrows represent the input residual velocities. Green dots show the CO-GNSS station distribution. Uncertainty ellipses are separately shown in Fig. 10. Topography and bathymetry after the ETOPO1 model (Amante and Eakins, 2009).

15 Unlike the Western Alps, where the deformation vectors indicate a very small (insignificant) internal surface deformation, the Alpine area between longitudes 6°E and 16°E presents an interesting kinematic pattern. The deformation vectors describe a progressive eastward rotation toward the Pannonian segment, starting with a north orientation near longitude 6°E and reaching N20°E close longitude 16°E. These vectors also make evident a shortening of about 2 mm/a across the southern front of the Eastern Alps, in the northern area of the Venetian-Friuli Basin. This is in agreement with the results published by Métois et al. (2015) and Cheloni et al. (2014). Although our CO-GNSS network presents a sparse station distribution in the northern Apennines and the Dinarides, our prediction supports the conclusions of D'Agostino et al. (2014) and Nocquet (2012) in that regions. In the former case, we identify a deformation roughly oriented to the northeast with the largest magnitudes (~ 3



mm/a) across the highest topographic features of the Apennines toward the Adriatic Sea. In the opposite direction, toward the Tyrrhenian Sea, the deformation magnitudes decrease up to 1.5 mm/a. According to Mantovani et al. (2015), this may be an evidence of a faster motion of the outer Apennine belt with respect to the inner one. In the case of the Dinarides, a shortening of about 3.0 mm/a across the Adriatic coastline is found. This magnitude decreases toward the Pannonian segment up to 0.5 mm/a. In the southern front of the Alps, the orientation of the deformation vectors presents a slightly progressive westward rotation from the Venetian surroundings toward the Po Basin. These findings are quantitatively equivalent to the results presented by Möller et al. (2011) and Devoti et al. (2011). Figure 10 shows the uncertainty (95% of confidence) of the horizontal surface deformation model. The largest values (from ± 0.5 to ± 1.5 mm/a) occur at those regions with poor station coverage (mainly Apennines, Dinarides and easternmost part of the Austrian Alps). In general, the mean accuracy of the model is ± 0.2 mm/a.




Figure 10: Estimated uncertainty (95% confidence) of the GNSS-based horizontal deformation model of the Alpine region. Green dots show the CO-GNSS station distribution. Topography and bathymetry after the ETOPO1 model (Amante and Eakins, 2009).

15 5.3 Vertical deformation model

The present-day vertical motion of the Alps and their surrounding areas is displayed in Fig. 11. Our results clearly show a still on-going averaged uplift of 1.8 mm/a of the entire mountain chain, with exception of the southern part of the Western Alps (6.5°E, 44°N), where no significant uplift (less than 0.5 mm/a) is detected. The largest uplift rates (more than 2 mm/a) occur in the central area of the Western Alps (7°E, 45°N), in the Swiss Alps (8°E, 46.5°N), and in the Southern Alps in the boundary region between Switzerland, Austria and Italy (11°E, 46.6°N). The uplift kinematics we present here for the Western Alps are consistent with the vertical motion patterns identified by Nguyen et al. (2016) and Nocquet et al. (2016). Our vertical rates in the Swiss Alps also agree



qualitatively with similar results inferred by Brockmann et al. (2012) using not only GNSS observations, but also levelling data.

In the northwest area, across the Rhone-Bresse and Rhine Grabens, a well-defined regional subsidence pattern with varying rates is observed. Apparently, the largest subsidence of around -1.3 mm/a happens in the Rhine Graben (7°E , 48.5°N), while in the Rhone-Bresse Graben (5°E , 46°N – 48°N) the largest rate reaches -0.8 mm/a. In the complete studied area, the largest subsidence rates are close to -1.5 mm/a and are located west of the Venetian-Friuli Basin. We do not detect a significant subsidence in the western part of the Po Basin; this may be a consequence of the poor distribution of our CO-GNSS stations in that region. Results of previous studies like Cuffaro et al. (2010) or Serpelloni et al. (2013) suggest subsidence rates up to -11 mm/a near Modena, Italy (11°E , 44.6°N). However, we do not detect these extreme values. This may be explained by the fact that we eliminate stations with strong local effects before the computation of the vertical-kinematics model, while the two studies mentioned base their conclusions on punctual station velocities. It is well-known that GNSS systematic errors and non-modelled local effects accumulate mainly in the estimation of the vertical coordinates. Therefore, we gave special care to identify and to remove those stations reflecting local effects (like water extraction, tunnel construction, landslides, etc.) that may mislead the estimation of vertical motions mirroring mountain building or subsidence processes in the Alpine region.





Figure 11: GNSS-inferred vertical deformation model of the Alpine region. Green dots show the CO-GNSS station distribution. Topography after the ETOPO1 model (Amante and Eakins, 2009).

20

The general uplift observed across the Alpine mountain chain decreases toward the outer regions to stable values between 0.0 and 0.5 mm/a and in some cases to subsidence motions like the Liguro-Provençal (5°E , 43.6°N) and Vienna (15.5°E , 48.5°N) Basins, where vertical rates of -0.8 mm/a and -0.3 mm/a are respectively detected. Figure



12 displays the estimated uncertainty of the predicted uplift and subsidence rates. The largest values (from ± 0.6 to ± 0.8 mm/a) occur in those areas with poor station coverage, mainly in the Apennine Peninsula and the Po Basin. The mean precision of the vertical motion model computed in this study for the Alpine mountain belt is about ± 0.3 mm/a.



5


Figure 12: Estimated uncertainty (95% confidence) of the GNSS-based vertical deformation model of the Alpine region. Dark blue dots show the CO-GNSS station distribution. Topography after the ETOPO1 model (Amante and Eakins, 2009).

5.4 Continuous velocity field

GNSS techniques are currently the primary tool for precise positioning. In applications of high-reliability
10 (accuracy in 1 cm level or better), the GNSS data processing requires the reference station coordinates in the same
reference frame and at the same observation epoch as the GNSS satellite orbits. In active seismic regions, strong
earthquakes cause large changes in the station positions and velocities of the geodetic reference stations, disabling
the use of previous coordinates (see for instance co-seismic and post-seismic effects of the El Maule earthquake
15 in the Latin American reference frame presented by Sánchez and Drewes (2016)). To ensure the long-term stability
of the geodetic reference frames, the transformation of station positions between different epochs requires the
computation of reliable continuous surface deformation (or velocity) models. Based on this kind of models,
geodesists monitor the kinematics of reference frames; infer appropriate transformation parameters between pre-
seismic and post-seismic (deformed) coordinates (specially in official matters like legal borders, cadastre, land
20 management, etc.), and interpolate surface motions arising from plate tectonics or crustal deformations in areas
where no geodetic stations are established. Consequently, after computing the deformation model (section 5.2),
the Eurasian plate motion removed from the CO-GNSS station velocities is restored to the deformation vectors to



get a continuous velocity field referring to the ITRF. As shown in Fig. 13, the velocity field mirrors the present-day surface kinematics as a combination of deformation patterns and the Eurasian plate tectonic motion, which is the dominant kinematic regime in Europe.



5

Figure 13: Continuous velocity field for the Alpine region referring to the ITRF2008 (above: horizontal velocity field, below: vertical velocity field). Topography and bathymetry after the ETOPO1 model (Amante and Eakins, 2009).

6. Strain field

The strain field represents the zones with compression, extension (or dilatation) and shear strain on the Earth's surface. In this study, the strain rates are computed from the deformation model (Fig. 9) as it provides a better



(smoothed) surface coverage than the residual station velocities (Fig. 7). This strategy allows to reduce random errors occurring at individual stations and enhances the major trends on strain patterns. Our computations are based on the infinitesimal strain theory, assuming that the deformation is much smaller (indeed, infinitesimally smaller) than any relevant dimension of the body, in this case the Alpine region. It means, very small deformation changes (at the mm/a level) over large distances (at the km level) between grid points are presumed. In the empirical procedure, the deformation vectors computed at the grid points are used for the estimation of deformation gradients, and these gradients are used for the computation of the strain components following the equations described by Lambeck (1988), pages 194-196. Even though the deformation field is available in three components, we concentrate only on the horizontal component. Thus, the obtained strain field is a horizontal projection of the surface deformation. The core contribution of a strain field is the possibility of distinguishing large-scale features (e.g. surface deformation, mountain building, etc.) from areas with large strain rates, which are often seismically active.




Figure 14: Strain distribution in the Alpine Region. Blue shades represent dilatation; red shades represent compression. Topography after the ETOPO1 model (Amante and Eakins, 2009).

Figure 14 shows the principal strain components (compression and dilatation) and the first strain invariant (sum of strain components) obtained in this study. Two main contrasting strain regimes are noticed – shortening across the south-eastern front of the Alps and the northern part of the Dinarides, and extension in the Apennines. The pattern of the strain principal axes indicates that the compression directions are more or less perpendicular to the thrust belt fronts (see Fig. 1). This compression presents maximum strain-rate values of about $20 \times 10^{-9} \text{ a}^{-1}$ in the Venetian-Friuli and Po Basins. These findings are in agreement with previous works published by Serpelloni et al. (2005, 2006), Caporali et al. (2009), Bennett et al. (2012) and Cheloni et al. (2014). Nevertheless, the lack of data in the eastern margin of the Adriatic Sea does not permit to conclude if this shortening continues with similar



magnitudes to the South Dinarides. As already stated by D'Agostino et al. (2014), dilatation is dominant in the north-western part of the Apennine Peninsula. Apparently, this dilatation continues to the West across the Po Basing, although at minor rates. However, poor station coverage in this area prevents from accurate conclusions from these observations.

5 Across the Alpine mountain belt, we observe a slight dilatation regime in the Western Alps, which smoothly changes to a contraction regime in West Austria and South Germany, reaching maximum shortening values of $6 \times 10^{-9} \text{ a}^{-1}$ in the north-eastern part of Austria (eastern margin of the Molasse Basin). The strain-rates obtained for the French and Swiss Alps are in the same order of uncertainty of our results; however, they are in agreement with results presented in previous studies like Calais et al. (2002) and Delacou et al. (2008). The magnitude and orientation of the strain principal axes we observe in the boundary region between Switzerland, Austria and Germany are very similar to the strain-rate patterns described by Tesauro et al. (2006).

10 The largest shear strain of about $20 \times 10^{-9} \text{ a}^{-1}$ is detected in the south-eastern part of the Alps (Fig. 15). The sharp boundaries between the left-lateral and right-lateral shear zones and a quiet area reveal the triple junction between the Eurasian, Adriatic and Pannonian plates (Möller et al., 2011). According to Brückl and Hammer (2014), the approximate location of the triple junction is 13.7°E , 46.6°N , which is well correlated with the strain model computed in this study.

15 The Western Alps and surroundings do not reveal any specific pattern of the shear strain while the Dinarides show a compressional strain with a right-lateral shear of about 10 to $20 \times 10^{-9} \text{ a}^{-1}$. The resulting principal and shear strain fields show that the main tectonic activity is ongoing on the plate boundaries.




Figure 15: Shear strain distribution in the Alpine Region. Blue and red shades represent the left-lateral and right-lateral shear, respectively. For illustration purposes, the left-lateral shear is considered as negative and the right-lateral as positive. Topography after the ETOPO1 model (Amante and Eakins, 2009).



Conclusions

In this study, we provide a present-day surface-kinematics model homogeneously assessed for the whole Alpine mountain chain. It is based on a high-level GNSS data analysis covering 12.4 years of continuous geodetic observations. Together with a coordinate solution for more than 300 CO-GNSS stations, our results comprise a deformation model, a continuous velocity field, and a strain field covering the area between 4°E-16°E and 43°N-50°N at a spatial interval of 25 km x 25 km. The averaged uncertainty of our model is ± 0.2 mm/a in the horizontal component and ± 0.3 mm/a in the height component. Although the station coverage is heterogeneous, our model describes coherent patterns of on-going uplift processes and horizontal deformations with respect to the Eurasian Plate. The determination of the constant velocities, which form the basis for the deformation model and strain field, was particularly complex. Since each little discontinuity had to be revealed, analysed and taken into account in determining the constant velocity of each site. Just because the geodetic network consists of many permanently operating GNSS stations, which have to fulfil different purposes, the analysis was difficult and required a lot of manual intervention. Some sites fulfil geodetic services requiring frequent updates in the set-up, which in turn cause inconsistencies in the time series that had to be recovered. Not all of these changes were well documented. The GNSS data processed in this study are unique as they cover the entire Alpine range and the associated foreland. The comparison of our derived deformation patterns with the results of other studies, whose research areas were mostly limited to smaller regions of the Alps, has proven the high reliability of our model. With this data set, a deformation model of the Alps derived from a uniform analysis is now available.

Supplementary data

- Station positions and velocities as well as velocity and deformation fields computed in the frame of this study are available through the PANGAEA (Data Publisher for Earth and Environmental Science) platform at <https://doi.pangaea.de/10.1594/PANGAEA.886889>. The dataset collection is composed of
- ALPS2017.SNX: Multi-year solution for geocentric Cartesian station positions and velocities referring to the IGB08, epoch 2010.0 in SINEX, the Solution (Software/technique) INdependent EXchange format.
 - ALPS2017_XYZ.CRD: Geocentric Cartesian station positions.
 - ALPS2017_XYZ.VEL: Geocentric Cartesian station velocities.
 - ALPS2017_NEH.CRD: Ellipsoidal station positions.
 - ALPS2017_NEH.VEL: Ellipsoidal station velocities.
 - ALPS2017_EPR.VEL: Station velocities relative to the Eurasian Plate.
 - ALPS2017_DEF_HZ.GRD: Surface deformation model of the Alpine Region.
 - ALPS2017_DEF_VT.GRD: Vertical deformation model of the Alpine Region.
 - ALPS2017_STR.GRD: Strain field of the Alpine Region.
 - ALPS2017_VEL.GRD: Continuous velocity field of the Alpine Region.

Acknowledgements

This study has been possible thanks to the support of different organizations providing us with the observational data of their CO-GNSS stations. We thank the International GNSS Service (IGS), the Reference Frame Subcommittee for Europe of the International Association of Geodesy (EUREF), the Federal Agency for Cartography and Geodesy of Germany (BKG), the Space Research Institute of the Austrian Academy of Sciences,



the Geodetic Alpine Integrated Network (GAIN), the Centro Ricerche Sismologiche (CRS) of the Istituto Nazionale di Oceanografia e di Geofisica Sperimentale (OGS), the Réseau National GPS France, and the Orpheon network. We are also grateful to our colleague E. Brockmann from swisstopo for providing us with the coordinate solution of the AGNES (Automated GNSS Network for Switzerland) network. The maps in this paper were
5 compiled using the GMT (Generic Mapping Tools) software package (Wessel et al., 2013).

References

- Altamimi Z., Collilieux X., and Métivier L.: ITRF2008: an improved solution of the international terrestrial reference frame, *J. Geod.*, 85(8): 457-473, doi:10.1007/s00190-011-0444-4, 2011.
- Amante C. and Eakins B.W.: ETOPO1 Global Relief Model converted to PanMap layer format. NOAA-National
10 Geophysical Data Center, PANGAEA, doi:10.1594/PANGAEA.769615, 2009.
- Aoudia A., Amodio A., Troisi C., Manzino A., Luchetta A., Völksen C., Drewes H., Zivcic M., Van der Voerd J., Fosson J.P., Carraro C., Zampedri G., Walpersdorf A., Laffi R., Fiononi M., Barzaghi R., Sabadini R., Pasetti L. and Sguero D.: ALPS GPSQUAKENET. Alpine Integrated GPS Network: Real-Time Monitoring and Master Model for Continental Deformation and Earthquake Hazard. Intergared III B Project –Alpine Space,
15 2007.
- Argnani A.: Plate Tectonics and the Boundary between Alps and Apennines, *Italian Journal of Geoscience*, 128(2): 317-330, doi:10.3301/IJG.2009.128.2.317, 2009.
- Bada G., Horváth F., Dövényi P., Szafián P., Windhoffer G. and Cloetingh S.: Present-day stress field and tectonic inversion in the Pannonian basin, *Global and Planetary Change* 58:165-180,
20 doi:10.1016/j.gloplacha.2007.01.007, 2007.
- Barletta V.R., Ferrari C., Diolaiuti G., Carnielli T., Sabadini R. and Smiraglia C.: Glacier shrinkage and modeled uplift of the Alps, *Geophys. Res. Lett.*, 33: L14307, doi:10.1029/2006GL026490, 2006.
- Becker A.: In situ stress data from the Jura Mountains - new results and interpretation. *Terra Nova*, 11: 9-15. doi:10.1046/j.1365-3121.1999.00215.x, 1999.
- 25 Bennett R.A., Hreinsdóttir S., Buble G., Bašić T., Bašić Ž, Marjanović M., Casale G., Gendaszek A. and Cowan D.: Eocene to present subduction of southern Adria mantle lithosphere beneath the Dinarides, *Geology* 36: 3–6, doi:10.1130/G24136A.1, 2008.
- Bennett R.A., Serpelloni E., Hreinsdóttir S., Brandon M.T., Buble G., Basic T., Casale G., Cavaliere A., Anzidei M., Marjonovic M., Minelli G., Molli G. and Montanari A.: Syn-convergent extension observed using the RETREAT GPS network, northern Apennines, Italy, *J. Geophys. Res.*, 117: B04408,
30 doi:10.1029/2011JB008744, 2012.
- Bird P.: An updated digital model for plate boundaries, *Geochem. Geophys. Geosyst.*, 4: 1027, doi:10.1029/2001GC000252, 2003.
- Blewitt G., Lavallée D., Clarke P. and Nurudinov K.: A new global model of Earth deformation: Seasonal cycle detected, *Science*, 294(5550): 2342–2345, doi:10.1126/science.1065328, 2001.
- 35 Böhlm J., Niell A., Tregoning P. and Schuh H.: Global mapping function (GMF): a new empirical mapping function based on numerical weather model data, *Geophys. Res. Lett.* 33: L07304, doi:10.1029/2005GL025546, 2006.
- Bousquet R., Schmid S.M., Zeilinger G, Oberhänsli R., Rosenberg C., Molli G., Robert C., Wiederkehr M. and Rossi Ph.: Tectonic framework of the Alps, CCGM/CGMW, retrieved from <http://www.geodynalps.org>, 2012.



- Brockmann E., Ineichen D., Marti U., Schaer S., Schlatter A. and Villiger A.: Determination of Tectonic Movements in the Swiss Alps using GNSS and Levelling, In: Kenyon S., Pacino M., Marti U. (eds) Geodesy for Planet Earth, IAG Symposia, 136: 689–695, doi:10.1007/978-3-642-20338-1_85, 2012.
- Brückl E., Bleibinhaus F., Gosar A., Grad M., Guterch A., Hrubcová P., Keller R., Majdański M., Šumanovac F.,
5 Tiira T., Yliniemi J., Hegedüs E. and Thybo, H.: Crustal structure due to collisional and escape tectonics in the Eastern Alps region based on profiles Alp01 and Alp02 from the ALP 2002 seismic experiment, *J. Geophys. Res.* 112: B06308, doi:10.1029/2006JB004687, 2007.
- Brückl E., Behm M., Decker K., Grad M., Guterch A., Keller G. and Thybo H.: Crustal structure and active tectonics in the eastern Alps, *Tectonics* 29, doi:10.1029/2009TC002491, 2010.
- 10 Brückl E. and Hammer C.: Eduard Suess' conception of the Alpine orogeny related to geophysical data and models, *Austrian J. of Earth Sciences*, 107(1), Vienna, 2014.
- Bruyninx C., Habrich H., Söhne W., Kenyeres A., Stangl, G. and Völksen C.: Enhancement of the EUREF permanent network services and products, IAG Symposia 138: 27-34, doi:10.1007/978-3-642-20338-1_4, 2012.
- 15 Bruyninx C., Habrich H., Kenyeres A., Söhne W., Stangl G. and Völksen C.: EUREF Permanent Network. International GNSS Service 2012 Technical Report, R. Dach and Y. Jean (eds), Astronomical Institute University of Bern, pp. 101-107, doi:10.7892/boris.80303, 2013.
- Bus Z., Grenerczy G., Tóth L. and Mónus P.: Active crustal deformation in two seismogenic zones of the Pannonian region – GPS versus seismological observations, *Tectonophysics* 474: 343-352, doi:10.1016/j.tecto.2009.02.045, 2009.
- 20 Calais E.: Continuous GPS measurements across the Western Alps, 1998-1998, *Geophys. J. Int.*, 38: 221-230, doi:10.1046/j.1365-246x.1999.00862.x, 1999.
- Calais E., Barlier F., Bayer R., Boucher C., Chéry J., Cotton F., Gariel J.C., Jouanne F., Martinod J. and Vigny C.: A permanent GPS network for monitoring crustal deformation in the Western Alps, *Annales Geophys* 6, Suppl. 1: C393, 1998.
- 25 Calais E., Galisson L., Stéphan J.-F., Delteil J., Deverchère J., Larroque C., Mercier de Lépinay B., Popoff M. and Sosson M.: Crustal strain in the Southern Alps, 1948-1998, *Tectonophysics*, 319: 1-17, doi:10.1016/S0040-1951(00)00029-9, 2000.
- Calais E., Nocquet J.M., Jouanne F., and Tardy M.: Current strain regime in the Western Alps from continuous
30 Global Positioning System measurements, 1996–2001, *Geology*, 30 (7): 651–654. doi:10.1130/0091-7613(2002)030<0651:CSRITW>2.0.CO;2, 2002.
- Caporali A., Aichhorn C., Barlik M., Becker M., Fejes I., Gerhatova L., Ghitau D., Grenerczy G., Hefty J., Krauss S., Medak D., Milev G., Mojzes M., Mulic M., Nardo A., Pesec P., Rus T., Simek J., Sledzinski J., Solaric M., Stangl G., Stopar B., Vespe F. and Virag G.: Surface kinematics in the Alpine-Carpathian-Dinaric and Balkan
35 region inferred from a new multi-network GPS combination solution, *Tectonophysics*, 474(1), 295–321, doi:10.1016/j.tecto.2009.04.035, 2009.
- Channell J.E.T. and Horváth F.: The African/Adriatic promontory as a paleogeographical premise for Alpine orogeny and plate movements in the Carpathó-Balkan region. *Tectonophysics* 35: 71–101, doi:10.1016/0040-1951(76)90030-5, 1976.
- 40 Channell J.E.T., D'Argenio B. and Horváth F.: Adria, the African promontory, in mesozoic Mediterranean palaeogeography, *Earth-Science Reviews*, 15(3): 213-292, doi:10.1016/0012-8252(79)90083-7, 1979.



- Cheloni D., D'Agostino N. and Selvaggi G.: Interseismic coupling, seismic potential and earthquake recurrence on the southern front of the Eastern Alps (NE Italy), *J. Geophys. Res. Solid Earth*, 119: 4448-4468, doi:10.1002/2014JB010954, 2014.
- Chen G. and Herring T.A.: Effects of atmospheric azimuthal asymmetry on the analysis of space geodetic data, *J. Geophys. Res.*, 102(B9): 20489-20502, doi:10.1029/97JB01739, 1997.
- 5 Chéry J., Vigny C., Meyer B., Ferhat G., Anzidei M., Bayer R., Boloh L., Briole P., Deschamps A., Feigl K., Gamond J.F., Geiger A., Jouanne F., Kasser M., Le Pape M., Martinod J., Ménard G., Ruegg J.C., Scheubel J.M. and Walsh J.J.: Global Positioning System network monitors the Western Alps. *EOS Trans Am Geophys Union* August 27, doi:10.1029/95EO00299, 1995.
- 10 Collilieux X., Altamimi Z., Coulot D., van Dam T. and Ray J.: Impact of loading effects on determination of the International Terrestrial Reference Frame, *Adv. Space Res.*, 45: 144–154, doi:10.1016/j.asr.2009.08.024, 2010.
- Collilieux X., van Dam T., Ray J., Coulot D., Métivier L. and Altamimi Z.: Strategies to mitigate aliasing of loading signals while estimating GPS frame parameters, *J. Geod.*, 86: 1–14, doi:10.1007/s00190-011-0487-6, 2012.
- 15 Copley A.: Kinematics and dynamics of the southeastern margin of the Tibetan Plateau, *Geophys. J. Int.*, 174(3): 1081-1100, doi:10.1111/j.1365-246X.2008.03853.x, 2008.
- Cuffaro M., Riguzzi F., Scrocca D., Antonioli F., Carminati E., Livani M. and Doglioni C.: On the geodynamics of the northern Adriatic plate, *Rend Fis Acc Lincei*, 21 (Sopl 1): S253-S279, doi: 10.1007/s12210-010-0098-9, 2010.
- 20 D'Agostino N., Cheloni D., Mantenuto S., Selvaggi G., Michelini A. and Zuliani D.: Interseismic strain accumulation in the eastern Southern Alps (NE Italy) and deformation at the eastern boundary of the Adria block observed by CGPS measurements, *Geophys. Res. Lett.*, 32, doi:10.1029/2006GL024266, 2005.
- D'Agostino N., Mantenuto S., D'Anastasio E., Giuliani R., Mattone M., Calcaterra S., Gambino P. and Bonci L.: Evidence for localized active extension in the central Apennines (Italy) from global positioning system observations, *Geology*, 39(4): 291–294, doi:10.1130/G31796.1, 2011.
- 25 D'Agostino N., England P., Hunstad I. and Selvaggi G.: Gravitational potential energy and active deformation in the Apennines, *Earth Planet. Sci. Lett.*, 397: 121-132, doi:10.1016/j.epsl.2014.04.013, 2014.
- Dach R., Lutz S., Walser P. and Fridez P. Eds.: *Bernese GNSS Software Version 5.2*. University of Bern, 2015.
- Dach R., Schaer S., Arnold D., Prange L., Sidorov D., Sušnik A., Villiger A. and Jäggi A.: CODE final product series for the IGS. Published by Astronomical Institute, University of Bern. URL: <http://www.aiub.unibe.ch/download/CODE>; doi:10.7892/boris.75876.2, 2017.
- 30 Delacou B., Sue Ch., Nocquet J.M., Champagnac J.D., Allanic C. and Burkhard M.: Quantification of strain rate in the Western Alps using geodesy: comparisons with seismotectonics, *Swiss J Geosci* 101: 377-385, doi:10.1007/s00015-008-1271-3, 2008.
- 35 Dercourt J., Zonenshain L.P., Ricou L.-E., Kazmin V.G., Le Pichon X., Knipper A.L., Grandjacquet C., Sbertshikov I.M., Geysant J., Lepvrier C., Pechersky D.H., Boulin J., Sibuet J.-C., Savostin L.A., Sorokhtin O., Westphal M., Bazhenov M.L., Lauer J.P. and Biju-Duval B.: Geological evolution of the tethys belt from the atlantic to the pamirs since the LIAS, *Tectonophysics*, 123 (1-4): 241-315, doi:10.1016/0040-1951(86)90199-X, 1986.



- Devoti R., Esposito A., Pietrantonio G., Pisani A.R. and Riguzzi F.: Evidence of large scale deformation patterns from GPS data in the Italian subduction boundary, *Earth Planet. Sci. Lett.* 311: 230-241, doi:10.1016/j.epsl.2011.09.034, 2011.
- 5 Devoti R., Riguzzi F., Cuffaro M. and Doglioni C.: New GPS constraints on the kinematics of the Apennines subduction, *Earth Planet. Sci. Lett.*, 273:163-174, doi:10.1016/j.epsl.2008.06.031, 2008.
- Dewey J.F., Pitman W.C.III, Ryan W.B.F. and Bonnin J.: Plate Tectonics and the Evolution of the Alpine System, *Geological Society of America Bulletin* 84(10): 3137–3180, doi:10.1130/0016-7606(1973)84<3137:PTATEO>2.0.CO;2, 1973.
- 10 Dewey J. F., Helman M. L., Knott S. D., Turco E. and Hutton D. H. W.: Kinematics of the western Mediterranean, *Geological Society, London, Special Publications*, 45: 265-283, doi:10.1144/GSL.SP.1989.045.01.15, 1989.
- Doglioni C.: Tectonics of the Dolomites (southern alps, northern Italy), *Journal of Structural Geology*, 9(2): 181-193, doi:10.1016/0191-8141(87)90024-1, 1987.
- Doglioni C. and Carminati E.: The effects of four subductions in NE-Italy, *Mem. Sci. Geol.* 54:1-4, 2002.
- 15 Dow J.M., Neilan R.E. and Rizos C.: The International GNSS Service in a hanging landscape of Global Navigation Satellite Systems, *J. Geod.*, 83: 191–198, doi:10.1007/s00190-008-0300-3, 2009.
- Drewes H.: Experiences with least squares collocation as applied to interpolation of geodetic and geophysical quantities, *Proceedings 12th Symposium on Mathematical Geophysics, Caracas, Venezuela*, 1978.
- Drewes H.: A geodetic approach for the recovery of global kinematic plate parameters, *Bull. Geod.* 56: 70-79, doi:10.1007/BF02525609, 1982.
- 20 Drewes H.: The Actual Plate Kinematic and Crustal Deformation Model APKIM2005 as Basis for a Non-Rotating ITRF. In: Drewes H. (eds) *Geodetic Reference Frames, IAG Symposia*, 134: 95-99, doi:10.1007/978-3-642-00860-3_15, 2009.
- Drewes H. and Heidbach O.: Deformation of the South American crust estimated from finite element and collocation methods. In: Sansò F (ed.) *A Window on the Future of Geodesy, IAG Symposia* 128: 544-549, doi:10.1007/3-540-27432-4_92, 2005.
- 25 Drewes H., Angermann D. and Seitz M.: Alternative definitions of the terrestrial reference system and its realization in reference frames. In: Altamimi Z., Collilieux X. (Eds.) *Reference Frames for Applications in Geosciences, IAG Symposia* 138: 39-44, doi:10.1007/978-3-642-32998-2_7, 2013.
- 30 Fejes I., Barlik M., Busics I., Packelski W., Rogowsky J., Sledzinsky J. and Zielinsky J.B.: The Central Europe Regional Geodynamics Project, paper presented at 2nd International Seminar on “GPS in Central Europe”, *Hun Acad Sci Penc, Hungary, Apr.* 27-29, 1993.
- Ferhat G., Feigl K.L., Ritz J.F. and Souriau A.: Geodetic measurement of tectonic deformation in the Southern Alps and Provence, France, 1947–1994. *Earth Planet. Sci. Lett.* 159: 35-46, doi:10.1016/S0012-821X(98)00065-X, 1998.
- 35 Flesch L.M., Holt W.E., Haines A.J. and Bingming Shen-Tu B.: Dynamics of the Pacific-North American Plate Boundary in the Western United States, *Science*, 287(5454): 834-836, doi:10.1126/science.287.5454.834, 2000.
- Gosar A., Šebela S., Košťák B. and Stemberk J.: Micro-deformation monitoring of active tectonic structures in W Slovenia, *Acta Geodyn. Geomater.* 4(1-145): 87-98, 2007.



- Grenerczy G., Kenyeres A. and Fejes I.: Present crustal movements and strain distribution in Central Europe inferred from GPS measurements, *J. Geophys. Res.*, 105(B9): 21835-21846, doi:10.1029/2000JB900127, 2000.
- 5 Grenerczy G., Sella G., Stein S. and Kenyeres, A.: Tectonic implications of the GPS velocity field in the northern Adriatic region, *Geophys. Res. Lett.*, 32: L16311, doi:10.1029/2005GL022947, 2005.
- Grenerczy G. and Kenyeres A.: Crustal deformation between Adria and the European platform from space geodesy, in the Adria microplate: GPS geodesy, tectonics and hazards, edited by Pinter N., Grenerczy G., Weber J., Stein S. and Medak D., *NATO Science Series IV: Earth and Environmental Sciences*, 61: 321-334, 2006.
- 10 Handy M.R., Schmid S.M., Bousquet R., Kissling E. and Bernoulli D.: Reconciling plate-tectonic reconstructions with the geological-geophysical record of spreading and subduction in the Alps, *Earth. Sci. Rev.*, 102: 121-158, 2010.
- Handy M.R., Ustaszewski K. and Kissling E.: Reconstructing the Alps–Carpathians–Dinarides as a key to understanding switches in subduction polarity, slab gaps and surface motion, *Int. J. Earth Sci. (Geol. Rundsch)* 104: 1-26, doi:10.1007/s00531-014-1060-3, 2015.
- 15 Haslinger C., Kraus S. and Stangl G.: The Intra-Plate Velocities of GPS Permanent Stations of the Eastern Alps, *Vermessung & Geoinformation*, 2: 66-72, ISSN 0029-9650, 2006.
- Haslinger C. and Stangl G.: The First Austrian Velocity Field derived from GPS. In: Report on the Symposium of the IAG Sub-commission for Europe (EUREF) held in Riga, Latvia, 14 - 17 June 2006. Retrieved from <http://www.euref.eu/symposia/2006Riga/Symposium2006-Riga.html>, 2006.
- 20 Heidbach O. and Drewes H.: 3-D finite element model of major tectonic processes in the Eastern Mediterranean, *Geol. Soc. Spec. Publ.*, 212: 261-274, doi:10.1144/GSL.SP.2003.212.01.17, 2003.
- Horváth F., Bada G., Szafián P., Tari G., Ádám A. and Cloetingh S.: Formation and deformation of the Pannonian Basin: Constraints from observational data, *Geological Society Memoir*, 32: 191-206 doi:10.1144/GSL.MEM.2006.032.01.11, 2006.
- 25 IGS: Specifications for the 2nd data reprocessing campaign of the International GNSS Service, retrieved from <http://acc.igs.org/reprocess2.html>, 2014.
- Kahle H.G., Müller M.V., Mueller St., Veis G., Billiris H., Paradissis D., Drewes H., Kaniuth K., Stuber K., Tremel H., Zerbini S., Pezzoli L. and Corrado G.: The convergent African/Eurasian plate boundary and associated crustal movements: GPS investigations in the Calabrian and West Hellenic arcs. *Proc. 8th Int. Symp. on Recent Crustal Movements, Kobe/Japan* 333-342, 1994.
- 30 Kaniuth K., Drewes H., Tremel H., Stuber K., Zerbini S., Pezzoli L. and Corrado G.: Crustal deformations in the Calabrian Arc area from five years of GPS measurements. *Proc. of ISTANBUL-94, 1st Turkish Internat. Symp. on Deformations, Istanbul*, 781-790, 1995.
- 35 Kaniuth K., Drewes H., Stuber K., Tremel H., Kahle H.G., Peter Y., Zerbini S., Tonti G. and Fagard H.: Crustal deformations in the central Mediterranean derived from the WHAT A CAT GPS project. In: *Proceedings of the 13th Working Meeting on European VLBI for Geodesy and Astrometry, Viechtach, Febr. 12-13*, edited by W. Schlüter and H. Hase, 192-197, 1999.
- 40 King M.A., Watson C.S., Penna N.T. and Clarke P.J.: Subdaily signals in GPS observations and their effect at semiannual and annual periods, *Geophys. Res. Lett.*, 35, L03302; doi:10.1029/2007GL032252, 2008.



- Kissling E., Schmid S. M., Lippitsch R., Ansoerge J. and Fügenschuh B.: Lithosphere structure and tectonic evolution of the Alpine arc: new evidence from high-resolution teleseismic tomography, *Geological Society, London, Memoirs*, 32: 129-145, doi:10.1144/GSL.MEM.2006.032.01.08, 2006.
- Kuhlemann J. and Kempf O.: Post-Eocene evolution of the North Alpine Foreland Basin and its response to Alpine tectonics, *Sedimentary Geology*, 152 (1-2): 45-78, doi:10.1016/S0037-0738(01)00285-8, 2002.
- 5 Lambeck K.: *Geophysical Geodesy: The slow deformations of the Earth*. Oxford University Press, New York, 1988.
- Larroque C., Delouis B., Godel B. and Nocquet J.M.: Active deformation at the southwestern Alps-Ligurian basin junction (France-Italy boundary): Evidence for recent change from compression to extension in the Argentera massif, *Tectonophysics* 467: 22–34, doi:10.1016/j.tecto.2008.12.013 , 2009.
- 10 Laubscher H.P.: The arcs of the Western Alps and the Northern Apennines: an updated view, *Tectonophysics*, 146(1-4): 67-78, doi:10.1016/0040-1951(88)90082-0, 1988.
- Le Pichon X., Bergerat F. and Roulet M.J.: Plate kinematics and tectonics leading to the Alpine belt formation; A new analysis. In: Clark S.P. Jr., Clark Burchfiel B. and Suppe J. (eds.) *Processes in Continental Lithospheric Deformation*, Geological Society of America, special paper 2018: 111-131, 1988.
- 15 Letellier T.: *Etude des ondes de marée sur les plateaux continentaux*. Thèse doctorale, Université de Toulouse III, Ecole Doctorale des Sciences de l'Univers, de l'Environnement et de l'Espace, 237 p, 2004.
- Lippitsch R., Kissling E. and Ansoerge J.: Upper mantle structure beneath the Alpine orogen from high-resolution teleseismic tomography, *J. Geophys. Res.*, 108 (B8): 2376, doi:10.1029/2002JB002016, 2003.
- 20 Mantovani E., Viti M., Cenni N., Babbucci D. and Tamburelli C.: Present Velocity Field in the Italian Region by GPS Data: Geodynamic/Tectonic Implications, *Int. J. Geosciences*, 6: 1285-1316. doi:10.4236/ijg.2015.612103, 2015.
- Meissl P.: *The use of finite elements in physical geodesy*, Rep. Dep. Geod. Sci. and Surv., The Ohio State Univ., Columbus, No.313, 1981.
- 25 Métois M., D'Agostino N., Avallone A., Chamot-Rooke N., Rabaute A., Duni L., Kuka N., Koci R. and Georgiev I.: Insights on continental collisional processes from GPS data: Dynamics of the peri-Adriatic belts, *J. Geophys. Res., Solid Earth*, 120: 8701–8719, doi:10.1002/2015JB012023, 2015.
- Möller G., Brückl E. and Weber R.: Active tectonic deformation at the transition from the European and Pannonian domain monitored by a local GNSS network: *Vermessung & Geoinformation* 2: 138-148, 2011.
- 30 Moritz H.: Least squares collocation, *Dt. Geod. Komm.*, Nr. A 75, 1973.
- Mueller S. and Kahle H.-G.: *Crust-Mantle Evolution, Structure and Dynamics of the Mediterranean-Alpine Region*. In Smith D.E. and Turcotte D.L. (eds.) *Contributions of Space Geodesy to Geodynamics: Crustal Dynamics*, American Geophysical Union, Washington, D.C., doi:10.1029/GD023p0249, 1993.
- Nguyen H.N., Vernant Ph., Mazzotti S., Khazaradze G. and Asensio E.: 3D GPS velocity field and its implications on the present-day postorogenic deformation of the Western Alps and Pyrenees, *Solid Earth Discuss.*, doi:10.5194/se-2016-78, 2016.
- 35 Nocquet J.M. and Calais E.: Geodetic measurements of crustal deformation in the Western Mediterranean and Europe, *Pure appl Geophys* 161: 661–681, doi:10.1007/s00024-003-2468-z, 2004.
- Nocquet J.M., Sue C., Walpersdorf A., Tran T., Lenôtre N., Vernant P., Cushing M., Jouanne F., Masson F., Baize S., Chéry J. and van der Beek V.A.: Present-day uplift of the western Alps, *Nature, Scientific Reports* 6: 28404, doi:10.1038/srep28404, 2016.
- 40



- Nocquet J.M.: Present-day kinematics of the Mediterranean: A comprehensive overview of GPS results, *Tectonophysics*, 579: 220-242, doi:10.1016/j.tecto.2012.03.037, 2012.
- Persaud M. and Pfiffner O.A.: Active Deformation in the Eastern Swiss Alps: Post-Glacial Faults, Seismicity and Surface Uplift. *Tectonophysics*, 385:59-84, doi:10.1016/j.tecto.2004.04.020, 2004.
- 5 Petit G. and Luzum B. (eds.): IERS Conventions 2010. IERS Technical Note 36. Verlag des Bundesamtes für Kartographie und Geodäsie, Frankfurt a.M., 2010.
- Pfiffner O.A., Frei W., Valasek P., Stäubli M., Levato L., DuBois L., Schmid S.M. and Smithson S.B.: Crustal shortening in the Alpine Orogen: Results from deep seismic reflection profiling in the eastern Swiss Alps, Line NFP 20-east, *Tectonics*, 9(6): 1327–1355, doi:10.1029/TC009i006p01327, 1990.
- 10 Ray J., Altamimi Z., Collilieux X. and van Dam T.M.: Anomalous harmonics in the spectra of GPS position estimates, *GPS Solutions*, 12(1): 55–64; doi:10.1007/s10291-007-0067-7, 2008.
- Rebischung P., Griffiths J., Ray J., Schmid R., Collilieux X. and Garayt B.: IGS08: the IGS realization of ITRF2008. *GPS Solutions*, 16 (4): 483-494, doi:10.1007/s10291-011-0248-2, 2012.
- Rodríguez-Solano C.J., Hugentobler U., Steigenberger, P. and Lutz S.: Impact of Earth radiation pressure on GPS position estimates, *J. Geod.*, 86: 309, doi:10.1007/s00190-011-0517-4, 2012.
- 15 Rossi G., Fabris P. and Zuliani D.: Slow transients recorded by the cGPS network FreDNet at the northern Adria microplate boundary (NE-Italy). In: EGU General Assembly Conference Abstracts, 15: 8417, 2013.
- Royden L.H.: Evolution of retreating subduction boundaries formed during continental collision, *Tectonics*, 12(3): 629–638, doi:10.1029/92TC02641, 1993.
- 20 Sánchez L. and Drewes H.: Crustal deformation and surface kinematics after the 2010 earthquakes in Latin America, *J. Geodynamics* 102: 1-23, doi:10.1016/j.jog.2016.06.005, 2016.
- Sánchez L., Seemüller W., Drewes H., Mateo L., González G., Silva A., Pampillón J., Martínez W., Cioce V., Cisneros D. and Cimbaro S.: Long-term stability of the SIRGAS reference frame and episodic station movements caused by the seismic activity in the SIRGAS region, *IAG Symposia* 138: 153-161, doi:10.1007/978-3-642-32998-2_24, 2013.
- 25 Schmid R., Dach R., Collilieux X., Jäggi A., Schmitz M. and Dilssner F.: Absolute IGS antenna phase center model igs08.atx: status and potential improvements, *J. Geod.* 90(4): 343-364, doi:10.1007/s00190-015-0876-3, 2016.
- Schmid S.M., Fügenschuh B., Kissling E. and Schuster R.: Tectonic map and overall architecture of the Alpine orogeny, *Eclogae geol. Helv.* 97: 93-117, doi:10.1007/s00015-004-1113-x, 2004.
- 30 Seitz F., Arenz H. and Leismüller F.: Position measurements at five permanent GPS stations in the Bavarian Alps as part of the Geodetic Alpine Integrated Network (GAIN) of the ALPS-GPSQUAKENET project. Deutsches Geodätisches Forschungsinstitut, Munich, doi:10.1594/PANGAEA.834193, 2014.
- Serpelloni E., Anzidei M., Baldi P., Casula G. and Galvani A.: Crustal velocity and strain-rate fields in Italy and surrounding regions: New results from the analysis of permanent and non-permanent GPS networks, *Geophys. J. Int.*, 161(3): 861–880, doi:10.1111/j.1365-246X.2005.02618.x, 2005.
- 35 Serpelloni E., Adzic M., Baldi P., Casula G. and Galvani A.: GPS measurement of active strains across the Apennines, *Annals of Geophysics, Supplement to Vol. 49* (1), doi:10.4401/ag-5756, 2006.
- Serpelloni E., Faccenna C., Spada G., Dong D. and Williams S.D.: Vertical GPS ground motion rates in the Euro-Mediterranean region: New evidence of velocity gradients at different spatial scales along the Nubia-Eurasia plate boundary, *J. Geophys. Res. Solid Earth*, 118(11): 6003–6024, doi:10.1002/2013JB010102, 2013.
- 40



- Steigenberger P., Lutz S., Dach R., Schaer S. and Jäggi A.: CODE repro2 product series for the IGS. Published by Astronomical Institute, University of Bern. URL: http://www.aiub.unibe.ch/download/REPRO_2013, doi:10.7892/boris.75680, 2014.
- 5 Stocchi P., Spada G. and Cianetti S.: Isostatic rebound following the Alpine deglaciation: impact on the sea level variations and vertical movements in the Mediterranean region, *Geophysical Journal International*, 162(1): 137-147, doi:10.1111/j.1365-246X.2005.02653.x, 2005.
- Sue C., Martinod J., Tricart P., Thouvenot F., Gamond J.F., Frechet J., Marinier D., Glot J.P. and Grasso J.R.: Active deformation in the inner western Alps inferred from comparison between 1972-classical and 1996-GPS geodetic surveys, *Tectonophysics*, 320: 17-29, doi:10.1016/S0040-1951(00)00024-X, 2000.
- 10 Sue Ch., Delacou B., Champagnac J.D., Allanic C. and Burkhard M.: Aseismic deformation in the Alps: GPS vs. seismic strain quantification, *Terra Nova* 19 (3): 182-188, 2007.
- Tesauro M., Hollenstein Ch., Egli R., Geiger A. and Kahle H.-G.: Analysis of central western Europe deformation using GPS and seismic data, *J. Geodyn.* 42: 194-209, doi:10.1016/j.jog.2006.08.001, 2006.
- Ustaszewski K., Schmid S.M., Fügenschuh B. Tischler M., Kissling E. and Spakman W.: A map-view restoration of the Alpine-Carpathian-Dinaridic system for the Early Miocene, *Swiss J. Geosci.* 101(Suppl 1): 273-294, doi:10.1007/s00015-008-1288-7, 2008.
- 15 van Dam T. and Ray R.: S1 and S2 atmospheric tide loading effects for geodetic applications, data retrieved from <http://geophy.uni.lu/ggfc-atmosphere/tide-loading-calculator.html>, 2010.
- van Mierlo J., Oppen S. and Vogel M.: Monitoring of recent crustal movements in the Eastern Alps with the Global Positioning System (GPS), *Tectonophysics*, 275: 273-283, doi:10.1016/S0040-1951(97)00032-2, 1996.
- Vergnolle M., Calais E. and Dong L.: Dynamics of continental deformation in Asia, *J. Geophys. Res.*, 112: B11403, doi:10.1029/2006JB004807, 2007.
- Völksen C., Walpersdorf A., Aoudia A., Barzaghi R., Borghi A. and Cannizzaro L.: The ALPS GPSQUAKENET project – A permanent GPS network in the Alps. In: *Bollettino di Geodesia e Science Affini*, Journal of Istituto Geografico Militare. Vol. LXVIII, 1: 1-17, 2009a.
- 25 Völksen C., Árnadóttir T., Geirsson H., and Valsson G.: Present day geodynamics in Iceland monitored by a permanent network of continuous GPS stations, *J Geody*, 48(3): 279-283, doi:j.jog.2009.09.033, 2009b.
- Waldhauser F., Kissling E., Ansorge J. and Mueller St.: Three dimensional interface modelling with two-dimensional seismic data: the Alpine crust-mantle boundary, *Geophysical Journal International*, 135 (1): 264–278, doi:10.1046/j.1365-246X.1998.00647.x, 1998.
- 30 Weber J., Vrabec M., Stopar B., Pavlovčič Prešeren P. and Dixon T.: The PIVO-2003 experiment: a GPS study of Istria peninsula and Adria microplate motion, and active tectonics in Slovenia. In: Pinter, N. et al. (eds.): *The Adria Microplate: GPS Geodesy, Tectonics, and Hazards*, 305-320, Springer, 2006.
- Wessel, P., Smith W. H. F., Scharroo R., Luis J. F. and Wobbe F.: *Generic Mapping Tools: Improved version released*, *EOS Trans. AGU*, 94, 409-410, 2013.
- 35 Ziegler P.A.: *Geological Atlas of Western and Central Europe* Shell Int. Petrol. Maatschap. Geol. Soc., London, 239 pp., 1990.
- Zuliani D., Battaglia M., Murray M.H., Michellini A., Burgmann R. and Marson I.: FREDNET: A continuous GPS geodetic network monitoring crustal deformation in NE Italy. *AGU Fall Meeting*, Poster presentation, 40 2002.