

57 years (1960-2017) of snow and meteorological observations from a mid-altitude mountain site (Col de Porte, France, 1325 m alt.)

Yves Lejeune¹, Marie Dumont¹, Jean-Michel Panel¹, Matthieu Lafaysse¹, Philippe Lapalus¹, Erwan Le Gac¹, Bernard Lesaffre¹, and Samuel Morin¹

¹Univ. Grenoble Alpes, Université de Toulouse, Météo-France, CNRS, CNRM, Centre d'Etudes de la Neige, Grenoble, France

Abstract. In this paper, we introduce and provide access to a daily (1960-2017) and hourly (1993-2017) ~~dataset~~ datasets of snow and meteorological data measured at the Col de Porte site, 1325 m a.s.l, ~~Chartreuse~~ Chartreuse, France. Site metadata and ancillary measurements such as soil properties and masks of the incident solar radiation are also provided. Weekly snow profiles are made available from September 1993 to ~~April 2015-~~ March 2018. A detailed study of the uncertainties originating from both ~~measurements~~ measurement errors and spatial variability within the measurement site is provided for several variables. We show that the estimates of the ratio of diffuse to total shortwave broadband irradiance is affected by an uncertainty of ± 0.21 (no unit). The estimated root mean squared deviation, ~~that can be mainly attributed to~~ which mainly represents spatial variability, is ± 10 cm for snow depth, ± 25 kg m⁻² for ~~snow-water~~ water equivalent of snow cover and ± 1 K for soil temperature (± 0.4 K during the snow season). The daily dataset can be used to quantify the effect of climate change at this site with a ~~reduction~~ decrease of the mean snow depth (Dec. 1st to April 30th) of 39 cm from 1960-1990 to 1990-2017 (40 % of the mean snow depth for 1960-1990) and an increase in temperature of + 0.90 K for the same periods. Finally, we show that the daily and hourly datasets are useful and appropriate for driving and evaluating a snowpack model over such a long period. The data are placed on the repository of the Observatoire des Sciences de l'Univers de Grenoble (OSUG) datacenter : <http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018>.

15 1 Introduction

The Col de Porte (CDP) site is a ~~meadow~~ mid-elevation meadow site located at 1325 m altitude (45.30° N, 5.77° E) in the Chartreuse mountain range. This observation site has been operated since 1959 in collaboration with several academic and non-academic partners (<https://www.umn-cnrm.fr/spip.php?rubrique218>). Daily measurements of snow depth, air temperature and precipitation amount have been performed since 1960. Hourly measurement of meteorological and snow variables required ~~for~~ running and evaluating to run and evaluate detailed snowpack model such as Crocus (Brun et al., 1992; Vionnet et al., 2012) started in 1987 and have been almost continuous during the snow season since snow season 1993-1994. Measured data are manually and automatically checked and corrected using the measurements of several sensors and ~~if required~~ meteorological analyses (SAFRAN, Durand et al., 1999) if required, thus ensuring the quality and continuity of the dataset.

Such a dataset provides a unique framework to drive and evaluate snowpack models over a long period. Indeed Essery et al. (2013) demonstrated that the evaluation of snowpack models can be misleading if performed over only a few snow seasons ~~only~~.

~~In the~~. In recent years, such datasets with ~~a varying level of details~~ varying levels of detail have been made public for several snow sites (e.g. Essery et al., 2016) and have motivated the publication of a special issue in Earth System Science Data to gather openly available detailed meteorological and hydrological observational archives from long-term research catchments in well-instrumented mountain regions around the world. This initiative arises from a GEWEX Hydroclimatology Panel cross-cut project, INARCH, the International Network for Alpine Research Catchment Hydrology.

CDP is part of several ~~observations~~ observation networks at the local level (Observatoire des Sciences de l'Univers de Grenoble, OSUG), at the national scale (Observation pour l'Experimentation et la Recherche en Environnement CryObsClim and Systemes d'Observation et d'Experimentation au long terme pour la Recherche en Environnement des glaciers GlacioClim) and contributes to OZCAR (Observatoires de la Zone Critique Applications et Recherches), one of the French components of the eLTER European Research Infrastructure (International Long-term Ecological Research Networks, Gaillardet et al., in review). It is also ~~labeled as a member~~ a reference station of the World Meteorological Observation (WMO) Global Cryospheric Watch Cryonet network and of the INARCH network. CDP snow and meteorological observations have been selected as an indicator of climate change ~~effect~~ effects at medium elevation by the national climate change observatory (ONERC). ~~CDP~~ The CDP dataset has been used as driving and evaluation data in several snow model intercomparison projects : SnowMIP (Etchevers et al., 2004) and ESM-SnowMIP (Krinner et al., 2018). CDP is also an ideal place for specific snow related measurements campaigns, e.g. ~~intercomparison of measurement methods for solid precipitation~~ the WMO Solid Precipitation Intercomparison Experiment (SPICE), measurement of the spectral reflectance of snow (Dumont et al., 2017; Tuzet et al., 2017), snow surface roughness (Picard et al., 2016), snow under forest (Sicart et al., 2017).

The objectives of the present paper are (i) ~~extending to extend~~ the hourly dataset published in Morin et al. (2012) from 1993-2011 to 1993-2017, (ii) ~~providing to provide~~ a daily dataset over the 1960-2017 period and (iii) ~~providing to provide~~ estimates of the uncertainties of several variables due to both spatial variability within the observation site and measurements uncertainties. The paper first describes the site and the dataset. The second section is dedicated to ~~provide estimates of measurements~~ providing estimates of measurement uncertainties and spatial variability within the site and the last section describes some examples of the use of this dataset. ~~Text similar to Morin et al. (2012) is underlined.~~

25 **2 Data description**

The Col de Porte site (Fig. 1) is a grassy meadow surrounded by mainly coniferous (spruces) and some lobed-leave trees. All the instruments are located within an area of $40 \times 50 \text{ m}^2$ (Fig. 2, ~~Tab.~~ Tables 2, 3, 4). The height of the trees ranges from 10 to 40 m. Note that all datasets are provided in universal time coordinate (UTC).

~~Picture of the site taken on 2014-03-10 from the South barrier, looking toward North.~~



Figure 1. [Picture of the site taken on 2014-03-10 from the South barrier, looking toward North.](#)




Figure 2. Schematic view of the experimental sites with sensor locations. The sensors indicated in yellow are for meteorological variables. The sensors indicated in red are not used anymore as of 2018, and those in blue correspond to snow measurements. Areas 23 and 24 correspond to soil temperature and humidity measurements. The correspondence between numbering and sensors is indicated in Tables 2, 3 and 4. For the sake of clarity, when a location is cited in the text, the reference to Fig. 2 is omitted and the location is directly linked to the figure or the corresponding table. The three dark blue asterisks correspond to the three hemispherical Webcam locations. The dedicated experimental area has been used for specific experiments, e.g. Dumont et al. (2017) or Bouilloud and Martin (2006).

Schematic view of the experimental sites with sensor locations. The sensors indicated in yellow are for meteorological variables. The sensors indicated in red are not used anymore as of 2018, those in blue corresponds to snow measurements. Areas 23 and 24 corresponds to soil temperature and humidity measurements. The correspondance between numbering and sensors are indicated in Tabs. 2, 3 and 4. The 3 emoticons correspond to the 3 Webcam locations.

5 2.1 Radiation masks




Figure 3. Masks measured at location 31 on July 1998 (left panel) and on June 2018 (right panel). Upper and lower mask elevations are represented by the coloured areas. Elevations are given in degrees, the center is 60 degrees elevation.

Surrounding trees and topography ~~are masking mask~~ part of the shortwave radiation. Masks were measured at ~~location 31~~ location 31 (corresponding to the measurements of the incoming shortwave radiation, see Fig. 2 and ~~Tab. Table 2~~) with 5° resolution in azimuth for two dates: July 1998 ~~and June 2018. (using a theodolite)~~ and June 2018 (using a compass and a clinometer). Masks are provided as a .csv file (doi:10.17187/CRYOBSCLIM.CDP.2018.SolarMask), they contain 3 values for each azimuth that ~~corresponds correspond~~ to: lower elevation, upper elevation and occultation percentage (p_{occ} , visually estimated) defined as follows (Fig. 3). Below the lower mask elevation, there is no direct radiation. Above the upper mask elevation, 100 percent of the direct radiation is available and between the two, only $100 - p_{occ}$ percent of the direct radiation is available. These masks are applied for the calculation of the direct/diffuse shortwave incoming radiation as explained in ~~section Sect.~~ Sect. 2.3.1. The discrepancies between the two masks are most likely due to changes of the vegetation (growing and ~~trees cut tree cutting~~ in 1999, see Morin et al., 2012).

~~Masks measured at location 31 on July 1998 (left panel) and on June 2018 (right panel). Upper and lower mask elevations are represented by the solid lines. Elevations are given in degrees, the center is 60 degrees elevation.~~

2.2 Soil and vegetation properties

Soil properties were measured close to ~~location-33 (see Fig-2)~~ location 33 on 29 September 2008, close to ~~location-24 on October-2nd~~ location 24 on 2 October 2012 and close to location 30 on 18 October 2017.

On 29 September 2008, the soil properties were measured over the first meter as illustrated by Fig. 4. The layering of the soil was estimated visually and is provided in ~~Tab-~~Table 1. The soil properties (particles size analysis, organic matter, nitrogen and carbon total content) were also analyzed down to 87 cm depth. The dataset is provided as a .csv file (soil_properties_2008.csv). On ~~October-2nd~~ 2 October 2012, the same analysis was conducted over the first 30 cm of soil at ~~location-24~~ location 24 along with measurements of the soil dry density. The dataset is provided as a .csv file (soil_properties_2012.csv). The two .csv files are available as doi:10.17178/CRYOBSCLIM.CDP.2018.Soil.

Table 1. Visual characterization of the soil layers corresponding to Fig. 4 on 29 September 2008.

Top depth (cm)	Bottom depth (cm)	Visual texture
0	5	organic soil with grass roots
5	18	organic soil without roots
18	47	clay and sand
47	70	grey clay and sand
70	87	grey clay
87	100	pebbles and grey clay, no sampling

On 18 ~~oetober~~ October 2017, the soil densities were ~~analyzed~~ analysed for the first 30 cm. At that time, the soil dry density was $1100 \pm 67 \text{ kg m}^{-3}$ without considering the vegetation. The soil wet density was $1475 \pm 59 \text{ kg m}^{-3}$. These values are the mean and standard deviation of 2 measurements over 0-10 cm depth and 2 at 20-30 cm depth close to ~~location-30-~~location 30. No significant differences between the two sampling depths were ~~measured~~ observed. On the same day, the vegetation (grass) dry and wet mass were measured on a 50 by 50 cm surface at the same location. The measurements result in a value of 1.92 kg m^{-2} for wet mass and 1.54 kg m^{-2} for dry mass. The height of the grass (roughly 5 cm) during the time of measurements can be considered as typical for late fall. Note that the grass is frequently cut during summer. These measured soil and vegetation properties can be useful ~~to constrain~~ for constraining soil and vegetation schemes which are often coupled with snowpack models (Decharme et al., 2013).



Figure 4. Soil profile of 1 meter depth performed close to **location 33**-location 33 on 29 September 2008. [The visual characterization provided in Table 1 can be seen on this picture.](#)

2.3 Meteorological hourly data, 1993-2017

The meteorological hourly dataset over 1993-2017 is an extension of the meteorological dataset provided in Morin et al. (2012) ~~An in which an~~ extensive description of the dataset is available ~~in this study. Below are reported.~~ ~~Below~~ only changes that happened after 2011 and additional details not provided in Morin et al. (2012) ~~are reported.~~

5 The dataset is provided as a continuous hourly dataset since 1993, so that it can be easily used to drive snowpack models. The partitioning of the dataset between *in-situ* data and the output of the meteorological analysis and downscaling tool SAFRAN (~~Durand et al., 1999~~) (~~Durand et al., 1999, 2009b~~) is the same as in ~~figure Fig.~~ 4 of Morin et al. (2012). For years 2011 to 2015, *in-situ* data are restricted to the period ~~20th 20~~ October of one year to ~~10th 10~~ June of the next year. ~~Starting on 10th Summer~~ ~~in situ data are thus missing (calibration of the sensors during summer) from 1993 to 2015. Starting on 10~~ June 2015, all
10 data are *in-situ* year-round except for very short periods with observation issues. An *in situ* flag is provided together with the meteorological data (value = 1 for *in situ* data).

Table 2 provides an ~~update update~~ of the type of sensors used for meteorological measurements with respect to ~~Tab. Table~~ 1 in Morin et al. (2012). The dataset is provided in netCDF format (doi:10.17178/CRYOBSCLIM.CDP.2018.MetInsitu) in the standard format for SURFEX surface model meteorological inputs (Vionnet et al., 2012; Masson et al., 2013). The ~~atmospherie~~
15 ~~atmospheric~~ pressure value corresponds to the mean climatological value at CDP.

2.3.1 Shortwave incoming radiation

The ~~meterological meteorological~~ dataset provides both total and diffuse incoming broadband ~~radiations at location 31. radiation~~ ~~at~~ location 31. The diffuse shortwave radiation is not measured but calculated from total shortwave and longwave incident radiation and air temperature as described in the following.

20 The first step of the procedure is to compute a cloudiness value, η (no unit, between 0 for clear sky and 1 for fully overcast) from measured air temperature T_{air} (K), longwave radiation LW_{down} (W m^{-2}) and specific humidity using ~~Equations 1 and 2~~ ~~from Berliand (1952) Eqs. (1) and (2) from Berliand (1952); Etchevers (2000).~~

$$LW_{down} = 1.05 \varepsilon \sigma T_{air}^4 \quad (1)$$

$$\varepsilon = 0.58 + 0.9 k(0.09 + 0.2\eta)\eta^2 + 0.06 e_{air} - 0.05 * \sqrt{e_{air}} (1 - 1 - k(0.09 + 0.2\eta)\eta^2) \quad (2)$$

25 ~~$k(\eta) = (0.09 + 0.2\eta)\eta^2$~~ $k(\eta) = (0.09 + 0.2\eta)\eta^2 \quad (3)$

where σ is the Stefan-Boltzman constant, and e_{air} is the water vapour partial pressure calculated from measured T_{air} and relative humidity, expressed in hPa. ~~The correction factor 1.05 in Eq. (1) accounts for the additional longwave radiation that~~

Table 2. Overview of the sensors used to gather the hourly meteorological data, between 1993 and 2017 at Col de Porte, France. The locations refer to Fig. 2.

Variable	Location	Sensor	Period of operation	Height	Unit	Integration method
Air temperature	12	PT 100/ 3 wires	... → 1996/1997	1.5 m*	K	Instantaneous
	12	PT 100/ 4 wires	1997/1998 → ...	1.5 m*	K	Instantaneous
	mast	PT 100/ 4 wires	1997/1998 → ...	3.1 m	K	Instantaneous
Relative humidity	13	SPSI MU-C.1/MUTA.2	... → 1994/1995	1.5 m*	%RH	Instantaneous
	13	Vaisala HMP 35DE	1995/1996 → 2005/2006	1.5 m*	%RH	Instantaneous
	13	Vaisala HMP 45D	2006/2007 → ...	1.5 m*	%RH	Instantaneous
Windspeed	2	Laumonier – heated	1997/1998 → ...	10 m	m s^{-1}	Integrated (60 min)
	7	Chauvin Arnoux Tavid 87 – non-heated	whole record	10 m	m s^{-1}	Integrated (60 min)
	15	Laumonier – heated	2000/2001 → 2014-2015	3.3 m	m s^{-1}	Integrated (60 min)
	3	Thies Ultrasonic anemometer – heated	March 2012 → ...	10 m	m s^{-1}	Integrated (60 min)
	18	Thies Ultrasonic anemometer – heated	Dec. 2013 → ...	3,3 m	m s^{-1}	Integrated (60 min)
Inc. shortwave radiation	31	Kipp & Zonen CM7	... → 15 March 1996	1.2 m*	W m^{-2}	Integrated (50 min)
	31	Kipp & Zonen CM14	15 March 1996 → Oct. 31st 2015	1.2 m*	W m^{-2}	Integrated (50 min)
	31	Kipp & Zonen CMP10	Nov. 2015 → ...	1.2 m*	W m^{-2}	Integrated (50 min)
Inc. longwave radiation	30	Eppley PIR	... → 2010/2011	1.2 m*	W m^{-2}	Integrated (50 min)
	30	Kipp & Zonen CG4	2010/2011 → Oct. 2015	1.2 m*	W m^{-2}	Integrated (50 min)
	30	Kipp & Zonen CGR4	Oct. 2015 → ...	1.2 m*	W m^{-2}	Integrated (50 min)
Precipitation	9	PG2000 heated (2000 cm ²), tipping bucket	whole record	2.75 m	$\text{kg m}^{-2} \text{s}^{-1}$	Difference
	1	PG2000 non-heated (2000 cm ²), tipping bucket	whole record	2.75 m	$\text{kg m}^{-2} \text{s}^{-1}$	Difference
	20	GEONOR (200 cm ²) with windshield, weighing gauge	whole record	3 m	$\text{kg m}^{-2} \text{s}^{-1}$	Difference
	17°	GEONOR T-200B-3 (200 cm ²), weighing gauge	Dec. 2013 → ...	3.1 m	$\text{kg m}^{-2} \text{s}^{-1}$	Difference
	19°	GEONOR T-200B-3 (200 cm ²) with windshield, weighing gauge	Dec. 2013 → ...	3.1 m	$\text{kg m}^{-2} \text{s}^{-1}$	Difference
	34°	OTT Pluvio 2 OTT (400 cm ²) with windshield, weighing gauge	Dec. 2013 → ...	3.1 m	$\text{kg m}^{-2} \text{s}^{-1}$	Difference*

* Height adjusted manually above snow surface (\approx weekly).

◊ The sensors were installed for the WMO SPICE project and are used in this study only to complement the dataset if a problem exists for the reference sensor.

* Amount processed in non-real-time (filtered values).

is reaching the sensor due to the presence of surrounding trees. Eq. (2) solution does not necessarily range between 0 and 1, η must be bounded between 0 and 1 when solving the equation.

The calculated value of η is then used to partition the total measured shortwave radiation into direct and diffuse fraction using the radiative transfer model from Vauge (1983) and the measured mask described in SeeSect. 2.

5 An additional shortwave radiation sensor (Delta-T SPN1 -heated) ~~has been installed at location 5 (Fig. 2)~~ was installed at location 5 in September 2016 (9.5 m above ground) and measures both diffuse and total shortwave radiation over the 400-2700 nm range.

A comparison between these measured and calculated direct/diffuse ~~distribution distributions~~ is provided in SeeSect. 3.1.

2.3.2 Longwave incident radiation

10 The sensor for incident longwave radiation was replaced in October 2015 by a Kipp&Zonen CGR4 sensor (~~location 30 in Fig. 2~~location 30). Figure 5 displays the comparison of the measured incident longwave radiation with simulated longwave radiation

from SAFRAN based on monthly average averages. It shows that the bias deviation between SAFRAN and the measurements displays two large breaks of -10 W m^{-2} in October 2015 and in autumn 2010 (corresponding to another sensor replacement, Tab. Table 2). Based on the hypothesis that the newest sensor is the reference since can be used as a reference because it was fully calibrated outside at the Physikalisch-Meteorologisches Observatorium (Davos, Switzerland) outside and inside with a blackbody, the dataset was corrected as follow : $+10 -10 \text{ W m}^{-2}$ from 1993 to November 2010 and $-10 +10 \text{ W m}^{-2}$ from November 2010 to November 2015. Since SAFRAN is the only available reference and does not account for local conditions, e.g. cloudiness, due to its coarse spatial resolution, it is unfortunately not possible currently to investigate with more temporal refinement this instrumental bias. This correction, although ranging in spanning the uncertainty values provided by sensor manufacturer the manufacturer, is of large significance for snowpack modelling considering the high sensitivity of the snowpack to processes governed by this variable (e.g. Raleigh et al., 2015; Sauter and Obleitner, 2015; Quéno et al., 2017). Using the Crocus snowpack model with or without the corrections leads to a shift in the melt-out date ranging between 5 and 10 days.




Figure 5. Monthly average of the difference between measured downward longwave and SAFRAN estimates. The two vertical black lines indicate the sensor changes (cf Tab. Table 2). The blue lines correspond to the raw time series and the green one to the corrected time series.

2.3.3 Precipitation

Precipitation data are handled according to Morin et al. (2012). Precipitation data are manually partitioned between liquid and solid phase using all relevant sources of data at the site, namely snow depth, surface albedo, surface and air temperatures and differences between heated and non-heated rain gauges (locations 1 and 9). The precipitation values provided in the dataset are based on the reference gauge, GEONOR, at ~~location 20~~-location 20. Other OTT and GEONOR gauges are used ~~in complement to~~ to complement the reference sensor measurements. Hourly solid precipitation measurements are corrected for undercatch depending on temperature and wind speed, as described in Morin et al. (2012). From 2013 to 2017, the wind measurement used for the correction ~~is placed at location~~ was the one placed at location 18 instead of location 15, since the ultrasonic sensor at location 18 is more accurate than the wind sensor at location 15. Note that locations 15 and 18 (Fig. 2), which better represents the wind at the level of the different precipitation gauges are very close, i.e. a few meters, so that the wind speed values are not significantly different between the two locations.

~~Overview of the sensors used to gather the hourly meteorological data, between 1993 and 2017 at Col de Porte, France. The index refers to the location of the sensor represented in Fig. 2. Text similar to Morin et al. (2012) is underlined.~~

2.4 Snow and soil data, 1993-2017

The ~~evaluation~~hourly~~hourly evaluation~~ dataset over 1993-2017 is an extension of the evaluation dataset provided in Morin et al. (2012). An extensive description of the dataset is available in the latter study. Below ~~are reported~~ only changes that happened after 2011 and additional details not provided in Morin et al. (2012) are reported. The hourly dataset is provided as a netCDF file (doi:10.17178/CRYOBSCLIM.CDP.2018.HourlySnow). Within this dataset, the soil temperature, soil humidity and settling disk temperature are raw measurements (uncorrected).

Table 3 provides an update of the type of sensors used for evaluation measurements with respect to ~~Tab.~~Table 2 in Morin et al. (2012).

Starting ~~on~~in October 2010, the snow depth at ~~location-32~~ location 32 has been measured with a Dimetix Laser ranger. The field-of-view is a few mm diameter spot and the accuracy provided by the manufacturer is ± 1.5 mm. Since October 2010, the snow depth ~~measurements~~measurement provided in the dataset (reference snow depth) is the ~~measurements~~measurement of the Dimetix Laser ranger. Data from the other ~~sensors~~snow depth sensors and precipitation amount are used to ~~evaluate and~~ correct the Laser data from small artefacts.

The surface temperature reference values contained in the dataset mainly originates from the Kipp&Zonen ~~upward~~pyrgeometer (location-25upward pyrgeometer (location 25), same sensor as ~~location-30 in Tab.~~ location 30 in Table 2). Since september 2010, these data are complemented by the other surface temperature sensors with a conical field of view shown in ~~Tab.~~Table 3. The reference surface temperature is bounded to 273.15 K when snow is present on the ground.

New sensors for soil temperature and humidity have been installed in October 2012 at several depths (-0.05, -0.1, -0.2, -0.3 m) at ~~location-23~~ location 23 close (roughly 2 m) to ~~location-24 (Fig. 2)~~ location 24 where the older soil temperature sensors were located. In total, for location 23, 3 probes are placed at 10 cm depth roughly 10 cm away from each other. In the following they are referred as s1_loc23_10, s2_loc23_10 and s3_loc23_10. At 20 cm depth, there is only two probes roughly 10 cm away from each other that are referred as s1_loc23_20, s2_loc23_20.

The differences between the measurements at these two locations are discussed in ~~See Sect.~~ 3.4. It must be underlined that the soil humidity measurements show that the soil is almost always saturated by liquid water when snow is present. This characteristic may not be typical for mountain slopes (e.g. Williams et al., 2009) and may be difficult to reproduce with usual soil models.

The measurements of the vertical profile of snowpack properties as described in Fierz et al. (2009) are also provided in caaml format (version 56) according to the International Association for Cryospheric Science (IACS) standard. They can be visualized using the niViz software ~~-(niviz.org)~~. An example is displayed in Fig. 6 for 13 January 2001. These profiles are available on a weekly basis from September 1993 to ~~April-2015-March 2018~~ (doi:10.17178/CRYOBSCLIM.CDP.2018.SnowProfile).

Table 3. Overview of the sensors used to gather the hourly and daily snow and soil data, between 1993 and 2017 at Col de Porte, France. Note that outgoing shortwave and longwave radiation is measured using instruments similar to the corresponding incoming radiation, described in Table 2. Note also that snow surface temperature can be derived from the outgoing longwave radiation sensor, in addition to the sensors presented here. The ~~index refers~~ locations refer to ~~the location of the sensor represented in~~ Fig. 2. Text similar to Morin et al. (2012) is underlined.

Variable	Location	Sensor	Period of operation	Height	Unit	Time resolution	Integration method
Snow depth	33	BEN ultrasonic depth gauge	... → 1999/2000	3 m	m	hourly	Instantaneous
	33	FNX ultrasonic depth gauge	2000/2001 → 2008/2009	3 m	m	hourly	Instantaneous
	33	Campbell Ultra-sound depth gauge SR50A	2009/2010 → ...	3.5 m	m	hourly	Instantaneous
	32	Dimetix Laser ranger	2010/2011 → ...	3.1 m	m	hourly	Instantaneous
	6°	Campbell Ultra-sound depth gauge SR50	Jan. 2014 → ...	4.1 m	m	hourly	Instantaneous
	6°	Campbell Ultra-sound depth gauge SR50ATH	Jan. 2014 → ...	4.1 m	m	hourly	Instantaneous
	6°	Jenoptik Laser ranger	Jan. 2014 → ...	4.1 m	m	hourly	Instantaneous
	6°	Dimetix Laser ranger	Jan. 2014 → ...	4.1 m	m	hourly	Instantaneous
	hatched	Snowpit (up to three values)	whole record	N.A.	m	≈ weekly	N.A.
Water equivalent of snow cover	16	Cosmic-Ray Neutron sensor	2001/2002 → ...	0 m	kg m ⁻²	daily	24h integration
	16	Cosmic-Ray Neutron sensor ^a	2008/2009 → ...	0 m	kg m ⁻²	daily	24h integration
	hatched	Snowpit (up to three values)	whole record	N.A.	kg m ⁻²	≈ weekly	N.A.
Runoff	11	5 m ² lysimeter – tipping gauge	... → March 1994	0 m	kg m ⁻² s ⁻¹	hourly	Difference
	11	5 m ² lysimeter – scale	March 1994 → ...	0 m	kg m ⁻² s ⁻¹	hourly	Difference
	14	1 m ² lysimeter – tipping gauge	... → Dec. 1996	0 m	kg m ⁻² s ⁻¹	hourly	Difference
	14	1 m ² lysimeter – scale	Dec. 1996 → ...	0 m	kg m ⁻² s ⁻¹	hourly	Difference
Surface temperature	22	Testo term Pyroterm	... → 2016/10	1.2 m ^b	K	hourly	Instantaneous
	21	Campbell IR120	Nov. 2015 → ...	0.8 m ^b	K	hourly	Instantaneous
	28	Heitronics KT15	2010/2011 → ...	3.2 m	K	hourly	Instantaneous
	4°	Campbell IR120	Jan. 2014 → ...	4.1 m	K	hourly	Instantaneous
Soil temperature	24	PT 100/3 wires	... → 1996/1997	-0.1 m	K	hourly	Instantaneous
	24	PT 100/4 wires	1997/1998 → ...				
	24	PT 100/3 wires	... → 1996/1997	-0.2 m	K	hourly	Instantaneous
	24	PT 100/4 wires	1997/1998 → ...				
	24	PT 100/3 wires	... → 1996/1997	-0.5 m	K	hourly	Instantaneous
	24	PT 100/4 wires	1997/1998 → ...				
	23	PT 100/4 wires	Oct. 2012 → ...	-0.05 m	K	hourly	Instantaneous
	23	PT 100/4 wires	Oct. 2012 → ...	-0.10 m	K	hourly	Instantaneous
	23	PT 100/4 wires	Oct. 2012 → ...	-0.20 m	K	hourly	Instantaneous
	23	PT 100/4 wires	Oct. 2012 → ...	-0.30 m	K	hourly	Instantaneous
Soil moisture	23	Delta-T ML2x ThetaProbe Moisture sensor	2012/10 → ...	-0.05 m	m ³ m ⁻³	hourly	Instantaneous
	23	Delta-T ML2x ThetaProbe Moisture sensor	Oct. 2012 → ...	-0.10 m	m ³ m ⁻³	hourly	Instantaneous
	23	Delta-T ML2x ThetaProbe Moisture sensor	Oct. 2012 → ...	-0.20 m	m ³ m ⁻³	hourly	Instantaneous
	23	Delta-T ML2x ThetaProbe Moisture sensor	Oct. 2012 → ...	-0.30 m	m ³ m ⁻³	hourly	Instantaneous
Settling disks temp.	27 and 29	PT 100/3 wires	... → 1996/1997	variable	K	hourly	Instantaneous
	27 and 29	PT 100/4 wires	1997/1998 → ...				
Settling disks height	27 and 29	In-house positioning system	whole record ^c	variable	m	hourly	Instantaneous
Ground flux	24	Hukseflux HFP01	since 2010/2011	0	W m ⁻²	hourly	Instantaneous

^a Sensor including shielding for ground-originating neutrons (reduced data scatter).

^b Height adjusted manually above snow surface (≈ weekly).

^c Progressive migration from mercury to solid state electric contact.

◊ The sensors have been installed for the WMO SPICE project and are used in this study only to complement the dataset if a problem exists for the reference sensor.




Figure 6. Example of snow profile measured on 13 January 2001 vizualized using niViz software.

2.5 1960-2017 Data

Table 4 describes the daily dataset that combines snow and meteorological measurements. The dataset is provided in netCDF format (doi:10.17178/CRYOBSCLIM.CDP.2018.MetSnowDaily). Variable names correspond to the names listed in ~~Tab-~~[Table 4](#). Within this daily dataset, the total precipitation dataset is not corrected for undercatch, ~~in~~ contrary to rain and snow datasets (starting in Sept. 1993). The total precipitation dataset is also not measured by the same sensor ~~as-used for~~ the rain and snow datasets (cf ~~Tab-~~[Table 4](#)). The total precipitation dataset is measured with the PG2000 sensor~~thus mitigating the impact of the undercatch correction, for which the undercatch plays a minor role compared to the GEONOR due to the 10 times larger collecting surface area (Table 2)~~. In addition, the total precipitation time ~~serie-series~~ may be qualified as inhomogeneous in time due to the various changes in precipitation gauges. The daily SWE automatic measurements (~~loc-16~~location 16, [Table 3](#)) are discarded for snow season 2015/2016 due to a disfunction of the sensor. Note also that the daily albedo data are uncorrected for local snow surface slope.

The hourly meteorological dataset that contains the whole SAFRAN reanalysis (Durand et al., 2009a) at Col de Porte for the period 1960-2017 is provided in order to drive snowpack simulation over the whole period. The dataset is provided in netCDF format (doi:10.17178/CRYOBSCLIM.CDP.2018.MetSafran) in the standard format for SURFEX meteorological inputs (Vionnet et al., 2012; Masson et al., 2013). The solar mask measured in 1998 (Fig. 3) is accounted for in this dataset.

Table 4. Description of the daily dataset between 1960 and 2017 at Col de Porte, France. The [index refers](#) [locations refer](#) to the location of the sensor represented in Fig. 2.

Variable	Location	Sensor	Period of operation	Unit	Description
T_{min}	12	PT100	... → 1993	K	Min. temp. between 00:00 (day D) and 24:00 (day D)
	12	cf Table 2	1993 → ...	K	Min. temp between 06:00 (day D-1) and 06:00 (day D)
T_{max}	12	PT100	... → 1993	K	Max. temp. between 00:00 (day D) and 24:00 (day D)
	12	cf Table 2	1993 → ...	K	Max. temp between 06:00 (day D) and 06:00 (day D+1)
snow_depth_auto	close to 33	automatic sensor	... → 1977/1978	m	Snow depth 06:00 (day D)
	33	Ultra-sound depth gauge BEN	1978/1979 → 1999/2000	m	Snow depth 06:00 (day D)
	33-6	cf Table 3	1993 → ...	m	Snow depth 06:00 (day D)
snow_depth_pit	hatched	manual	1963/1964 → 7 Feb. 1996	m	Irregular frequency
	hatched	manual	8 Feb. 1996 → ...	m	Weekly
snow_depth_pit_north	hatched	manual	2001/2002 → ...	m	Weekly
snow_depth_pit_south	hatched	manual	2001/2002 → ...	m	Weekly
swe_auto	16	cf Table 3	2001/2002 → ...	kg m ⁻²	Daily (not available for 2015-2016)
swe_pit	hatched	manual	1963/1964 → 7 Feb. 1996	kg m ⁻²	Irregular frequency, SWE core 38.5 and 25. cm ²
	hatched	manual	8 Feb. 1996 → ...	kg m ⁻²	Weekly, SWE core 100 cm ²
swe_pit_north	hatched	manual	2001/2002 → ...	kg m ⁻²	Weekly, SWE core 100 cm ²
swe_pit_south	hatched	manual	2001/2002 → ...	kg m ⁻²	Weekly, SWE core 100 cm ²
total_precipitation	9	cf Table 2	1960/1961 → 2004/2005	kg m ⁻²	Daily sum of precipitation not corrected for undercatch 06:00 (day D) to 06:00 (day D+1)
rain [◊]	20	cf Table 2	1993/1994 → ...	kg m ⁻²	Daily sum of corrected liquid precipitation, 06:00 (day D) to 06:00 (day D+1)
snow [◊]	20	cf Table 2	1993/1994 → ...	kg m ⁻²	Daily sum of corrected solid precipitation, 06:00 (day D) to 06:00 (day D+1)
height of new snow	33,27	calculated from snow depth measurement and settlement disks	whole record	cm	Daily sum of new snow, 06:00 (day D) to 06:00 (day D+1)
albedo_daily	26 and 31	cf Table 2	2005/2006 → ...	NA	Ratio of the daily sums of reflected and incident shortwave radiations
albedo_daily_flag	26 and 31	NA	2005/2006 → ...	NA	Number of hourly measurements used to calculate daily albedo

[◊] Note that rain and snow variables are provided only when *in situ* measurements are available (i.e *in situ* flag of [Tab. Table 2](#) - see also [figure Fig. 4](#) in Morin et al., 2012).

3 Spatial variability and measurements uncertainties

The dataset presented in this study is, like any observation dataset, affected by different sources of uncertainties. Regardless of whether these data ~~should be~~ are used for model evaluation or process ~~study~~ studies, characterizing their associated uncertainties is essential for a proper use of the data. The uncertainties of the dataset may come from ~~measurements~~ measurement uncertainties (including instrumental and environmental uncertainties) but also from the spatial variability of the variables within the measurement plot.

A lower bound of the uncertainty for each variable can be estimated from the information provided by the sensor manufacturer. Some variables are measured at different locations within the field sites and by different sensors. This provides a better insight of the uncertainty associated ~~to~~ with both sources for each variable. Lafaysse et al. (2017) already provided a first estimate of the ~~uncertainty associated to~~ uncertainties associated with snow depth, ~~snow water equivalent~~ water equivalent of snow cover, bulk density, broadband albedo, soil temperature and snow surface temperature. In this section, we extend the period and the number of points used for the ~~uncertainties evaluation~~ uncertainty evaluations for snow depth, ~~snow water equivalent~~ water equivalent of snow cover and soil temperature for which several measurements are available over a sufficiently long period. We also provide ~~uncertainties assessment~~ uncertainty assessments of the direct/diffuse incident shortwave radiation estimates (cf ~~see 2.3.1 Sect. 2.3.1 for the calculation of the estimates~~). Note that an update on the uncertainties for snow surface temperature and broadband albedo is not provided in this study (lack of a sufficient number of sensors) though their uncertainty estimates are crucial for snow model evaluation. In this respect, we recommend the use of uncertainty values provided in Lafaysse et al. (2017) for these two variables.

3.1 Direct/diffuse shortwave incoming radiation

A first source of uncertainties in the calculation of the distribution of the measured broadband shortwave radiation into diffuse and direct radiation originates from the uncertainties of the mask used for the calculation (cf ~~see Sect. 2, Fig. 3~~). Using the methodology explained in ~~See Sect. 2.3.1~~, we estimate the direct and diffuse shortwave incoming radiation based on the mask from 1998 and the mask from 2018 for two snow seasons (~~Sept-1st September~~ to 30 June) : 2015-2016 and 2016-2017. The mean difference (mask measured in ~~2017-2018~~ minus mask measured in 1998) and root mean square deviation (RMSD) computed between diffuse components (over non zero values only) are -1.30 W m^{-2} and 10.1 W m^{-2} . The mean difference and RMSD for the diffuse to total ratio are -0.02 and 0.10 , respectively. The histogram of differences ~~are is~~ provided in Fig. 7a.

The accuracy of the methodology described in ~~See Sect. 2.3.1~~ has also been evaluated using the measurements of total and diffuse ~~radiations from location 2~~ radiation from location 5 (at 10 m above ground) and the mask measured in October 2017 at the same location. The comparison is done from ~~September-1st September~~ 2016 to 30 June 2017 during daylight (i.e., if total measured shortwave ~~is~~ larger than 4 W m^{-2}). The mean difference between the estimated and simulated diffuse component is -15.26 W m^{-2} (RMSD: 53 W m^{-2}). The mean difference and RMSD computed for the diffuse to total ratio are -0.08 and 0.21 . ~~The histograms of differences are~~, respectively. ~~The histogram of differences is~~ provided in Fig. 7b. This shows that the




Figure 7. Comparison of different broadband diffuse to total shortwave radiation ratio, r . (a) Difference in ratio estimated with the mask measured in June 2018 and in June 1998 at ~~location 25~~-location 25. Statistics are calculated during daylight from ~~Sep-1st~~-September 2015 to 30 June 2017 excluding July and August for each year. (b) Difference in ratio estimated with the 2017 mask (measured at ~~location 5~~-location 5, ~~21~~ October 21st-2017) and the measured ratio at ~~location 5~~-location 5. Statistics are calculated during daylight from ~~September~~ 1st-September 2016 to 30 June 2017.

estimation of the diffuse radiation ~~is slightly biased low~~ has a slightly negative bias and that this uncertainty has to be taken into account for applications such as radiative balance calculation for which the direct/diffuse distribution has a significant impact. It also shows that the methodology applied to partition the direct and diffuse components has a larger impact on the uncertainty than the change in solar masks shown in Fig. 3.

5 3.2 Snow depth

Table 5. Statistics of the comparisons between the different snow depth measurements represented in Fig. 8.

Sensors	Number of times	Deviation (m)	RMSD (m)	Period
Nivose 1 - h_{ref}	22498	-0.007	0.039	Sept. 2009 to June 2016
Mast - h_{ref}	22225	0.013	0.036	Sept. 2009 to June 2016
Pit - h_{ref}	874	0.053	0.077	Sept. 1960- June 2017
North Pit - h_{ref}	261	0.124	0.128	Sept. 2001 to June 2017
South Pit - h_{ref}	261	0.107	0.108	Sept. 2001 to June 2017




Figure 8. Comparison of snow depth measurements at different locations. h_{ref} corresponds to ~~location 33~~-location 33. (a) Difference in measured snow depth between the ultrasound sensor placed ~~on at~~ the Nivose 1 ~~location (Fig. 2)~~-location and the reference snow depth (~~locations 32-33 in Fig. 2~~locations 32-33) in blue. In red, differences between the measured snow depth at ~~location 6 (Fig. 2)~~location 6 and the reference snow depth. The ~~difference-differences~~ are calculated from snow season 2009/2010 to snow season 2015/2016 using only data from 20 ~~Sept-September~~ to 10 June. Data where both locations indicate 0 snow depth are excluded from the statistics. (b) Difference in measured snow depth between the manual snow depth measurements at ~~snowpit field location (Fig. 2)~~-snowpit field location and reference automatic snow depth (~~location 33~~location 33) in blue, between manual snow depth measurement in the snow pit south field and reference in grey and snow pit north field and reference in red. Difference values are calculated over ~~the~~ 1960-2017 period for the pit value and 2001-2017 for north and south pits. Data where both locations indicate 0 snow depth are excluded from the statistics. Corresponding statistics are provided in ~~Tab-Table~~ Table 5.

Figure 8 compares the snow depth reference value mostly measured at ~~location 32-33~~location 32-33, h_{ref} (~~Fig. 2~~) with several other measurements of snow depth : in panel (a) with respect to automatic snowdepth measurements at ~~locations "Nivose 1" and 6 (see Fig. 2) and~~locations "Nivose 1" and 6 ~~and~~ in panel (b) with respect to manual snow depth measurement in snow pit fields (main, north and south, blue hatched areas in Fig. 2). For panel (a), the comparison is done over the 2009-2016 period and
 5 ~~the automated measurement have been manually corrected for any blank period and measurements inconsistency- any blank or inconsistent measurement period in the "Nivose 1" (resp. mast) sensor was discarded from the comparison.~~ For panel (b), the comparison with the main snow pit field is done over 1960-2017 and for the ~~pits~~-south and north ~~pits~~ over 2001-2017. For each sensor, the number of points used to calculate the statistics are in ~~Tab-Table~~ Table 5.

Figure 8a and Table 5 show that the three automatic measurements exhibit deviations lower than 1.3 cm and that the RMSD is
 10 lower than 4 cm. Higher discrepancies are found between the reference automatic measurements and the manual measurements (~~Figure Fig. 8b~~) with mean deviation reaching almost ~~14-13~~ cm and RMSD ~~14-13~~ cm. These higher difference values might be attributed to the local slope, aspect, and small topographic features within the three snow pit ~~fields-area~~ ~~field areas~~ and to

the higher ~~measurements uncertainties associated to~~ measurement uncertainty associated with manual measurements. Extreme difference values corresponds to the end of the snow season when the snow cover is patchy. Picard et al. (2016) installed during the 2014-2015 snow season an automatic scanning laser meter close to ~~location 6~~ location 6 that scanned an area of 100-200 m². During this snow season, the laser measurements indicated a spatial variability of the snow depth within the footprint that can reach 7-10 cm (RMSD). We thus recommend the use of ± 10 cm uncertainty value for snow depth in any evaluation to represent the spatial variability within the site, ~~that is~~ comparable to the values used in Lafaysse et al. (2017).

3.3 ~~Snow water~~ Water equivalent of snow cover




Figure 9. Comparison of SWE measurements at different locations. (a) Difference in measured SWE between the manual measurement in the snow pit field (Fig. 2) and the reference automatic SWE (SWE_{ref} , ~~location 16~~ location 16 in Fig. 2) in blue. The difference are calculated over the period 2001-2017 (no reference data for 2015/2016 snow season). Data where both locations indicate 0 SWE are excluded from the statistics. Note that the manual measurements from snow pit south and north are used for the SWE sensor (~~location 16~~ location 16) calibration. (b) Difference in manually measured SWE between the snowpit field south and the snow pit field location in blue, between the snow pit field north and south locations in green and snow pit north field and snow pit field in red. Difference are calculated over the 2001-2017 period. Data where both locations indicate 0 SWE are excluded from the statistics. Numerical values are provided in Tab. Table 6.

Figure 9 and ~~Tab. 6 compares~~ Table 6 compare the SWE automatic measurements at ~~location 16~~ (Fig. 2) location 16 with the manual ~~measured in~~ measurements from the main snow pit field (panel a) and the three locations for manual SWE measurements (panel b). The statistics are calculated over the ~~2001-2016~~ 2001-2017 period. It must be underlined that the **SWE**

Table 6. Statistics of the comparisons between the different SWE measurements represented in Fig. 9.

Sensors	Number of dates	Deviation (kg m^{-2})	RMSD (kg m^{-2})	Period
Pit - SWE _{ref}	244	-16.83	24.44	Sept. 2001 to June 2017
South Pit - Pit	239	17.37	25.09	Sept. 2001 to June 2017
North Pit - South Pit	260	-6.69	17.66	Sept. 2001 to June 2017
South Pit - Pit	239	11.84	20.01	Sept. 2001 to June 2017

~~automatic-automatic~~ SWE sensor is calibrated using the manual measurements at snow pit fields south and north. The average of the annual maximum value of SWE_{ref} during this period is $389 \pm 104 \text{ kg m}^{-2}$.

Figure 9 and ~~Tab.~~ Table 6 show that the mean difference between the automatic and ~~the~~ manual measurements in the main snow pit field reaches -17 kg m^{-2} with RMSD of almost 25 kg m^{-2} . The comparison between the three locations of manual measurements displays RMSD reaching 25 kg m^{-2} , i.e. 8.6 % of average peak SWE values. This value is consistent with the spatial variability of snow depth and can probably be used as an estimate of the uncertainty associated ~~to~~ with the SWE dataset both due to ~~measurements-measurement~~ errors and spatial variability.

3.4 Soil temperature

Table 7. Statistics of the comparisons between the different soil temperature measurements represented in Fig. 10.

Sensors	Depth (cm)	Number of dates	Deviation (K)	RMSD (K)	Period
s2_loc23_10 - s1_loc23_10	10	15084	0.034	0.110	Dec. 2015 to June 2017
s3_loc23_10 - s1_loc23_10	10	15084	-0.094	0.244	Dec. 2015 to June 2017
s3_loc23_10 - s2_loc23_10	10	15084	0.128	0.182	Dec. 2015 to June 2017
loc_23 - loc_24	10	11396	-0.108	0.415	Dec. 2015 to June 2017 (snow season)
loc_23 - loc_24	10	3688	-1.059	1.100	Dec. 2015 to June 2017 (summer)
s2_loc23_20 - s1_loc23_20	20	15084	0.093	0.118	Dec. 2015 to June 2017
loc_23 - loc_24	20	11396	-0.224	0.390	Dec. 2015 to June 2017 (snow season)
loc_23 - loc_24	20	3688	-0.943	0.961	Dec. 2015 to June 2017 (summer)

Figure 10 and ~~Tab.~~ Table 7 compare the different soil temperature measurements at 10 and 20 cm depths for ~~locations 23 and 24 (Fig. 2).~~ The first column locations 23 and 24. The left panels in Fig. 10 display the statistics of the different temperature probes ~~located index 23 and spaces at~~ location 23 and spaced by roughly 10 cm -(s1_loc23_10, s2_loc23_10 and s3_loc23_10 and s1_loc23_20, s2_loc23_20, resp.). It indicates that the RMSD between the 3 probes is lower than 0.25 K .-The second column compares location 24 (Table 7). The right panels in Fig. 10 compare locations 24 (old sensors) and 23 (new sensors mean) for two periods : summer (20 June to 10 October) and snow season (11 October to 19 June). During the snow season, the two locations show a small mean deviation of -0.11 K and an RMSD of 0.42 K , while during summer the mean deviation is roughly -1.06 K leading to RMSD of 1.10 K (Table 7). Note that these two locations are spaced by only a few meters only (see Fig. 2). The temperature difference between the two sensors may be attributed to differences in soil properties, local topography




Figure 10. Comparison between the different soil temperature measurements at 10 cm (panels a and b) and 20 cm (panels c and d) depths. Panels a and c compare the new sensors (3 probes roughly 10 cm away of each other) at location 23 to the old ones (2 probes roughly 10 cm away of each other at -20 cm). Panels b and d compare the average values of the new sensors (location 23) to the old ones (location 24). Statistics are calculated from December 2015 to July 2017. Summer (panels b and d, in red) corresponds to the period between 20 June 2016 and 10 October 2016 and 20 June 2017 to 31 July 2017. The rest of the dates corresponds to snow season (panels b and d, in blue). Numerical values are provided in [Table 7](#).

and shading. The larger differences in summer may be due to (i) larger heterogeneity in soil wetness and (ii) the absence of the snow cover that spatially tempers the surface temperature signal in winter.

From these observations, a lower bound of the uncertainty of the soil temperature measurements (spatial variability and measurements errors) is roughly 1.10 K during summer, roughly 0.42 K during the snow season and a little higher than 0.5 K averaged over the whole year.

4 Data use

4.1 Temperature, snow depth and precipitation since 1960

Fig. 11 displays the evolution of mean snow depth, air temperature and total precipitation from ~~Dec.-1st to April~~ December to 30th April of each snow season for the whole period of the dataset (Dec. 1960 - April 2017). This figure shows an example of a direct use of the dataset to study the past evolution of winter conditions at Col de Porte. It demonstrates that the decrease in mean snow depth reduction between 1960-1990 and 1990-2017 is 39 cm (40 % of the mean snow depth for 1960-1990), while the air temperature has increased ~~of by~~ 0.90 °C over the same period ~~and while~~ the total precipitation does not exhibit a significant trend. This indicates that at this site, the reduction of the snow cover is mainly due to the increase in temperature and its consequences (e.g. higher snow/rain limit ~~when during~~ precipitation and higher melt rates). ~~This long time serie contributes~~ These long time series contribute to placing long term climate change impact studies on mountain snow conditions in the context of past changes (Verfaillie et al., 2018).

4.2 Snow model evaluation

This dataset has been widely used to drive and evaluate snow models (e.g. ~~Essery et al., 2013; Magnusson et al., 2015; Decharme et al., 2016; Lafaysse et al., 2017; Piazzini et al., 2018~~ Essery et al., 2013; Wever et al.) A list of the studies using CDP dataset is available at <http://www.umr-cnrm.fr/spip.php?article533>. ~~It has also been included in the Earth System Model – Snow Model Intercomparison Project (-)~~

5 Data availability

The database (doi:10.17178/CRYOBSCLIM.CDP.2018) presented and described in this article is available for download at <http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018>. ~~Tab-~~ Table 8 provides the links to the different datasets.

25 6 Conclusions

This paper describes and provides access to the daily snow and meteorological dataset measured at the Col de Porte site, 1325 m a.s.l, Chartreuse, France for the period 1960-2017. The hourly dataset of snow and meteorological observations for the period 1993-2017 is made available along with weekly snow profiles from September 1993 to March 2018, soil properties and




Figure 11. Evolution of mean snow depth, air temperature and total precipitation over 1960-2017. The mean and total values are calculated over the period Dec-1st December to April-30th April of each snow season. The black lines are the 15-year moving means.

Table 8. Link to the dataset repository

Dataset	Period	Format	Repository
Solar Mask	July 1998 and June 2018	csv	http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018.SolarMask
Soil properties	29 September 2008 and October 2nd 2012	csv	http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018.Soil
Hourly in situ meteorological data	August 1st 1993 to 31 July 2017	netCDF	http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018.MetInsitu
Hourly SAFRAN meteorological data	August 1st 1960 to 31 July 2017	netCDF	http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018.MetSafran
Daily snow and meteorological data	August 1st 1960 to 31 July 2017	netCDF	http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018.MetSnowDaily
Hourly snow data	August 1st 1960 to 31 July 2017	netCDF	http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018.HourlySnow
Snow profiles	September 1993 to March 2018	caaml	http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018.SnowProfile

[solar radiation masks. Based on measurements at several locations within the measurement field, we estimated the uncertainties and spatial variability of : the ratio between solar diffuse and total irradiance, snow depth, water equivalent of snow cover and soil temperature. The data are placed on the repository of the Observatoire des Sciences de l'Univers de Grenoble \(OSUG\) datacenter : <http://dx.doi.org/10.17178/CRYOBSCLIM.CDP.2018>.](#)

- 5 *Author contributions.* Y.L. and J.-M. P. endorse the responsibility of the experimental site and of the instruments. M.D. led the consolidation of the data set and wrote this manuscript together with all co-authors. E. L. and J.-M. P. ensure the proper working of the instruments.

Acknowledgements. Many thanks are expressed to people of CNRM/CEN who have assisted in the collection, collation and archive of this unique dataset. We thank in particular É. Pougatch, Y. Danielou, J.-L. Dumas for their crucial work on the database. Generating and distributing this dataset directly benefitted from the LabEX OSUG@2020 (ANR10 LABX56). The authors are also thankful to EDF/DTG,

- 10 ONF, SOERE CryObsClim, GCW and the IDEX Univ. Grenoble Alpes Cross Disciplinary Project Trajectories. The authors also ~~thanks~~ [thank](#) Arnaud Foulquier (LECA) for his help with soil and vegetation properties measurements and Laurent Bourges (OSUG) for the establishment of the data repository and the allocation of dedicated dois. [The authors are grateful to R. Essery, C. Fierz and one anonymous referee for their useful comments on the manuscript, and to M. Bavay and C. Fierz for their help with niViz and caaml formats.](#)

References

- Berliand, M.: Determining the net long-wave radiation of the Earth with consideration of the effect of cloudiness, *Izv. Akad. Nauk. SSSR Ser. Geofiz.*, 1, 64–78, 1952.
- Bouilloud, L. and Martin, E.: A coupled model to simulate snow behavior on roads, *J. Appl. Meteorol.*, 45, 500–516, 2006.
- 5 Brun, E., David, P., Sudul, M., and Brunot, G.: A numerical model to simulate snow-cover stratigraphy for operational avalanche forecasting, *J. Glaciol.*, 38, 13 – 22, [http://refhub.elsevier.com/S0165-232X\(14\)00138-4/ff0155](http://refhub.elsevier.com/S0165-232X(14)00138-4/ff0155), 1992.
- Decharme, B., Martin, E., and Faroux, S.: Reconciling soil thermal and hydrological lower boundary conditions in land surface models, *Journal of Geophysical*, 118, 7819–7834, doi:10.1002/jgrd.50631, 2013.
- Decharme, B., Brun, E., Boone, A., Delire, C., Le Moigne, P., and Morin, S.: Impacts of snow and organic soils parameterization on northern
10 Eurasian soil temperature profiles simulated by the ISBA land surface model, *The Cryosphere*, 10, 853–877, doi:10.5194/tc-10-853-2016, <http://www.the-cryosphere.net/10/853/2016/>, 2016.
- Dumont, M., Arnaud, L., Picard, G., Libois, Q., Lejeune, Y., Nabat, P., Voisin, D., and Morin, S.: In situ continuous visible and near-infrared spectroscopy of an alpine snowpack, *The Cryosphere*, 11, 1091–1110, doi:10.5194/tc-11-1091-2017, <https://www.the-cryosphere.net/11/1091/2017/>, 2017.
- 15 Durand, Y., Giraud, G., Brun, E., Mérindol, L., and Martin, E.: A computer-based system simulating snowpack structures as a tool for regional avalanche forecasting, *J. Glaciol.*, 45, 469–484, 1999.
- Durand, Y., Giraud, G., Laternser, M., Etchevers, P., Mérindol, L., and Lesaffre, B.: Reanalysis of 47 Years of Climate in the French Alps (1958–2005): Climatology and Trends for Snow Cover, *J. Appl. Meteor. Climat.*, 48, 2487–2512, doi:10.1175/2009JAMC1810.1, 2009a.
- Durand, Y., Giraud, G., Laternser, M., Etchevers, P., Mérindol, L., and Lesaffre, B.: Reanalysis of 44 Yr of Climate in the French Alps
20 (1958–2002): Methodology, Model Validation, Climatology, and Trends for Air Temperature and Precipitation., *J. Appl. Meteor. Climat.*, 48, 429–449, doi:10.1175/2008JAMC1808.1, 2009b.
- Essery, R., Morin, S., Lejeune, Y., and Menard, C. B.: A comparison of 1701 snow models using observations from an alpine site, *Advances in water resources*, 55, 131–148, doi:10.1016/j.advwatres.2012.07.013, 2013.
- Essery, R., Kontu, A., Lemmetyinen, J., Dumont, M., and Ménard, C. B.: A 7-year dataset for driving and evaluating snow models at
25 an Arctic site (Sodankylä, Finland), *Geoscientific Instrumentation, Methods and Data Systems*, 5, 219–227, doi:10.5194/gi-5-219-2016, <http://www.geosci-instrum-method-data-syst.net/5/219/2016/>, 2016.
- Etchevers, P.: Modélisation de la phase continentale du cycle de l’eau à l’échelle régionale. Impact de la modélisation de la neige sur l’hydrologie du Rhône, Ph.D. thesis, PhD thesis, Université Paul Sabatier, Toulouse, 2000.
- Etchevers, P., Martin, E., Brown, R., Fierz, C., Lejeune, Y., Bazile, E., Boone, A., Dai, Y.-J., Essery, R., Fernandez, A., Gusev, Y., Jordan, R., Koren, V., Kowalczyk, E., Nasonova, N. O., Pyles, R. D., Schlosser, A., Shmakin, A. B., Smirnova, T. G., Strasser, U., Verseghy, D., Yamazaki, T., and Yang, Z.-L.: Intercomparison of the surface energy budget simulated by several snow models (SNOWMIP project),
30 *Ann. Glaciol.*, 38, 150 – 158, doi:10.3189/172756404781814825, 2004.
- Fierz, C., Armstrong, R. L., Durand, Y., Etchevers, P., Greene, E., McClung, D. M., Nishimura, K., Satyawali, P. K., and Sokratov, S. A.: The international classification for seasonal snow on the ground, IHP-VII Technical Documents in Hydrology n 83, IACS Contribution n
35 1, <http://unesdoc.unesco.org/images/0018/001864/186462e.pdf>, 2009.
- Gaillardet, J., Braud, I., Hankard, F., et al.: OZCAR, the French network of critical zones observatories, *Vadoze Zone Journal*, in review.

- Krinner, G., Derksen, C., Essery, R., Flanner, M., Hagemann, S., Clark, M., Hall, A., Rott, H., Brutel-Vuilmet, C., Kim, H., Ménard, C. B., Mudryk, L., Thackeray, C., Wang, L., Arduini, G., Balsamo, G., Bartlett, P., Boike, J., Boone, A., Chéruy, F., Colin, J., Cuntz, M., Dai, Y., Decharme, B., Derry, J., Ducharne, A., Dutra, E., Fang, X., Fierz, C., Ghattas, J., Gusev, Y., Haverd, V., Kontu, A., Lafaysse, M., Law, R., Lawrence, D., Li, W., Marke, T., Marks, D., Nasonova, O., Nitta, T., Niwano, M., Pomeroy, J., Raleigh, M. S., Schaedler, G., Semenov, V., Smirnova, T., Stacke, T., Strasser, U., Svenson, S., Turkov, D., Wang, T., Wever, N., Yuan, H., and Zhou, W.: ESM-SnowMIP: Assessing models and quantifying snow-related climate feedbacks, *Geoscientific Model Development Discussions*, 2018, 1–32, doi:10.5194/gmd-2018-153, <https://www.geosci-model-dev-discuss.net/gmd-2018-153/>, 2018.
- Lafaysse, M., Cluzet, B., Dumont, M., Lejeune, Y., Vionnet, V., and Morin, S.: A multiphysical ensemble system of numerical snow modelling, *The Cryosphere*, 11, 1173–1198, doi:10.5194/tc-11-1173-2017, <http://www.the-cryosphere-discuss.net/tc-2016-287/>, 2017.
- 10 Magnusson, J., Wever, N., Essery, R., Helbig, N., Winstral, A., and Jonas, T.: Evaluating snow models with varying process representations for hydrological applications, *Water Resources Research*, 51, 2707–2723, 2015.
- Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouysse, F., Brousseau, P., Brun, E., Calvet, J.-C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essauoui, K., Gibelin, A.-L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Lebeau-pin Brossier, C., Lemonsu, A., Mahfouf, J.-F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy, G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of Earth surface variables and fluxes, *Geoscientific Model Development*, 6, 929–960, doi:10.5194/gmd-6-929-2013, 2013.
- 15 Morin, S., Lejeune, Y., Lesaffre, B., Panel, J.-M., Poncet, D., David, P., and Sudul, M.: A 18-years long (1993 - 2011) snow and meteorological dataset from a mid-altitude mountain site (Col de Porte, France, 1325 m alt.) for driving and evaluating snowpack models, *Earth Syst. Sci. Data*, 4, 13–21, doi:10.5194/essd-4-13-2012, 2012.
- Piazzì, G., Thirel, G., Campo, L., and Gabellani, S.: A particle filter scheme for multivariate data assimilation into a point-scale snowpack model in an Alpine environment, *The Cryosphere*, 12, 2287–2306, doi:10.5194/tc-12-2287-2018, <https://www.the-cryosphere.net/12/2287/2018/>, 2018.
- Picard, G., Arnaud, L., Panel, J.-M., and Morin, S.: Design of a scanning laser meter for monitoring the spatio-temporal evolution of snow depth and its application in the Alps and in Antarctica, *The Cryosphere*, 10, 1495–1511, doi:10.5194/tc-10-1495-2016, <https://www.the-cryosphere.net/10/1495/2016/>, 2016.
- 25 Quéno, L., Karbou, F., Vionnet, V., and Dombrowski-Etchevers, I.: Satellite products of incoming solar and longwave radiations used for snowpack modelling in mountainous terrain, *Hydrology and Earth System Sciences Discussions*, 2017, 1–33, doi:10.5194/hess-2017-563, <https://www.hydrol-earth-syst-sci-discuss.net/hess-2017-563/>, 2017.
- 30 Raleigh, M., Lundquist, J., and Clark, M.: Exploring the impact of forcing error characteristics on physically based snow simulations within a global sensitivity analysis framework, *Hydrology and Earth System Sciences*, 19, 3153–3179, 2015.
- Sauter, T. and Obleitner, F.: Assessing the uncertainty of glacier mass-balance simulations in the European Arctic based on variance decomposition, *Geoscientific Model Development*, 8, 3911–3928, doi:10.5194/gmd-8-3911-2015, <https://www.geosci-model-dev.net/8/3911/2015/>, 2015.
- 35 Sicart, J. E., Ramseyer, V., Lejeune, Y., Essery, R., Webster, C., and Rutter, N.: Spatial and Temporal Variabilities of Solar and Longwave Radiation Fluxes below a Coniferous Forest in the French Alps, in: *AGU Fall Meeting Abstracts*, 2017.

- Tuzet, F., Dumont, M., Lafaysse, M., Picard, G., Arnaud, L., Voisin, D., Lejeune, Y., Charrois, L., Nabat, P., and Morin, S.: A multilayer physically based snowpack model simulating direct and indirect radiative impacts of light-absorbing impurities in snow, *The Cryosphere*, 11, 2633–2653, doi:10.5194/tc-11-2633-2017, <https://www.the-cryosphere.net/11/2633/2017/>, 2017.
- Vauge, P. d. B.: Le gisement solaire-Evaluation de la ressource énergétique, *Ciel et Terre*, 99, 62, 1983.
- 5 Verfaillie, D., Lafaysse, M., Déqué, M., Eckert, N., Lejeune, Y., and Morin, S.: Multi-component ensembles of future meteorological and natural snow conditions for 1500 m altitude in the Chartreuse mountain range, Northern French Alps, *The Cryosphere*, 12, 1249–1271, doi:10.5194/tc-12-1249-2018, <https://www.the-cryosphere.net/12/1249/2018/>, 2018.
- Vionnet, V., Brun, E., Morin, S., Boone, A., Martin, E., Faroux, S., Moigne, P. L., and Willemet, J.-M.: The detailed snowpack scheme Crocus and its implementation in SURFEX v7.2, *Geosci. Model. Dev.*, 5, 773–791, doi:10.5194/gmd-5-773-2012, 2012.
- 10 Wever, N., Fierz, C., Mitterer, C., Hirashima, H., and Lehning, M.: Solving Richards Equation for snow improves snowpack meltwater runoff estimations in detailed multi-layer snowpack model, *The Cryosphere*, 8, 257–274, doi:10.5194/tc-8-257-2014, 2014.
- Williams, C., McNamara, J., and Chandler, D.: Controls on the temporal and spatial variability of soil moisture in a mountainous landscape: the signature of snow and complex terrain, *Hydrology and Earth System Sciences*, 13, 1325–1336, 2009.