

# The Global Fire Atlas of individual fire size, duration, speed, and direction

5 Niels Andela<sup>1,2</sup>, Douglas C. Morton<sup>1</sup>, Louis Giglio<sup>3</sup>, Ronan Paugam<sup>4</sup>, Yang Chen<sup>2</sup>, Stijn Hantson<sup>2</sup>, Guido R. van der Werf<sup>5</sup>, and James T. Randerson<sup>2</sup>

<sup>1</sup>Biospheric Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

<sup>2</sup>Department of Earth System Science, University of California, Irvine, CA 92697, USA

10 <sup>3</sup>Department of Geographical Sciences, University of Maryland, College Park, MD 20742, USA

<sup>4</sup>Centre Europe´en de Recherche et de Formation Avanc´ee en Calcul Scientifique, URA1875, CNRS, Toulouse, France.

<sup>5</sup>Faculty of Science, Vrije Universiteit Amsterdam, Amsterdam, Netherlands

15 *Correspondence to:* Niels Andela (niels.andela@nasa.gov)

15

**Abstract.** Natural and human-ignited fires affect all major biomes, altering ecosystem structure, biogeochemical cycles, and atmospheric composition. Satellite observations provide global data on spatiotemporal patterns of biomass burning and evidence for rapid changes in global fire activity in response to land management and climate. Satellite imagery also provides detailed information on the daily or sub-daily position of fires that can be used to understand the dynamics of individual fires. The Global Fire Atlas is a new global dataset that tracks the dynamics of individual fires to determine the timing and location of ignitions and fire size, duration, daily expansion, fire line length, speed, and direction of spread. Here we present the underlying methodology and Global Fire Atlas results for 2003-2016 derived from daily moderate resolution (500 m) Collection 6 MCD64A1 burned area data. The algorithm identified 13.3 million individual fires over the study period, and estimated fire perimeters were in good agreement with independent data for the continental United States. A small number of large fires dominated sparsely populated arid and boreal ecosystems, while burned area in agricultural and other human-dominated landscapes was driven by high ignition densities that resulted in numerous smaller fires. Long-duration fires in the boreal regions and natural landscapes in the humid tropics suggest that fire-season length exerts a strong control on fire size and total burned area in these areas. In arid ecosystems with low fuel densities, high fire spread rates resulted in large, short-duration fires that quickly consumed available fuels. Importantly, multi-day fires contributed the majority of burned area in all biomass burning regions. A first analysis of the largest, longest, and fastest fires that occurred around the world revealed coherent regional patterns of extreme fires driven by large-scale climate forcing. Global Fire Atlas data are publicly available through [www.globalfiredata.org](http://www.globalfiredata.org), and individual fire information and summary data products provide new information for benchmarking fire models within ecosystem and Earth system models, understanding vegetation-fire feedbacks, improving global emissions estimates, and characterizing the changing role of fire in the Earth system.

40

## 1 Introduction

Worldwide, fires burn an area larger than the size of the European Union every year (Randerson et al., 2012; Giglio et al., 2013). The majority of burned area occurs in grasslands and savannas, fire-adapted ecosystems where fires maintain open landscapes by reducing shrub and tree cover (Scholes and Archer, 1997; Abreu et al., 2017). However, all major biomes burn. Climate controls global patterns of fire activity by driving vegetation productivity and fuel build up as well as fuel conditions (Bowman et al., 2009). Humans are the dominant source of ignitions in most flammable ecosystems, but human activities also reduce fire sizes through landscape fragmentation and fire suppression (Archibald et al., 2012; Taylor et al., 2016; Balch et al., 2017).

Over the past ~~two decades~~ 18 years, socio-economic development and corresponding changes in human land use have considerably reduced fire activity in fire-dependent grasslands and savannas worldwide (Andela et al., 2017). At the same time, warming climate has dried fuels and has increased the length of fire seasons across the globe (Jolly et al., 2015), which is particularly important in forested ecosystems with abundant fuels (e.g., Kasischke and Turetsky, 2006; Aragão et al., 2018). Fire activity increases non-linearly in response to drought conditions in populated areas of the humid tropics (Alencar et al., 2011; Field et al., 2016), resulting in large scale degradation of tropical ecosystems (van der Werf et al., 2008; Morton et al., 2013b; Brando et al., 2014), and extensive periods of poor air quality (Johnston et al., 2012; Lelieveld et al., 2015; Koplitz et al., 2016). Moreover, increasing population densities in highly flammable biomes also amplify the socio-economic impacts of wildfires related to air quality or damage to houses and infrastructure (Moritz et al., 2014; Knorr et al., 2016). Despite the importance of understanding changing global fire regimes for ecosystem services, human well being, climate, and conservation, our current understanding of changing global fire regimes is limited because existing satellite data products detect actively burning pixels or burned area, but not individual fires and their behavior.

Frequent observations from moderate-resolution, polar-orbiting satellites may provide information on individual fire behavior in addition to estimates of total burned area. Several recent studies have shown that fire-affected pixels can be separated into clusters based on spatial and temporal proximity. This information can be used to study the number and size distributions of individual fires (Archibald and Roy, 2009; Hantson et al., 2015; Oom et al., 2016), fire shapes (Nogueira et al., 2017; Laurent et al., 2018), and the location of ignition points (Benali et al., 2016; Fusco et al., 2016). One limitation of fire clustering algorithms that rely on spatial and temporal proximity of fire pixels is the inability to separate individual fires within large burn patches that contain multiple ignition points, a frequent phenomenon in grassland biomes. To address the possibility of multiple ignition points, other algorithms have specifically tracked the spread of individual fires in time and space, with demonstrated improvements for isolating ignition points and constraining final fire perimeters (Frantz et al., 2016; Andela et al., 2017). In addition to the size and ignition points of individual fires, other studies used daily or sub-daily detections of fire activity to track growth dynamics of fires (Loboda and Csiszar, 2007; Coen and Schroeder, 2013; Veraverbeke et al., 2014; Sá et al., 2017). Together, these studies highlight the strengths and limitations of using daily or sub-daily satellite imagery to derive information on individual fires and their behavior over time.

Here we present the Global Fire Atlas of individual fires based on a new methodology to identify the location and timing of fire ignitions and estimate fire size, duration, daily expansion, fire line, speed, and direction of spread. The Global Fire Atlas is derived from the Moderate Resolution Imaging Spectroradiometer (MODIS) collection 6 burned area dataset (Giglio et al., 2018) and estimated day of burn information at 500 m resolution. Individual fire data were generated starting in 2003, when combined data from the Terra and Aqua satellites provide greater burn date certainty. The algorithm for the Global Fire Atlas tracks the daily progression of individual fires at 500 m resolution to produce a set

of metrics on individual fire behavior in standard raster and vector data formats. Together, these Global Fire Atlas data layers provide an unprecedented look at global fire behavior and changes in fire dynamics during 2003-2016. The data are freely available at <http://www.globalfiredata.org>, and new years will be added to the dataset following the availability of global burned area data.

95

## 2 Data and Methods


Here we developed a method to isolate individual fires from daily moderate resolution burned area data. The approach used two filters to account for uncertainties in the day of burn in order to map the location and timing of fire ignitions and the extent and duration of individual fires (Fig. 1). Subsequently, we tracked the growth dynamics of each individual fire to estimate the daily expansion, daily fire line, speed and direction of spread. Based on the Global Fire Atlas algorithm, burned area was broken down into seven fire characteristics in three steps (Fig. 1b). First, burned area was described as the product of ignitions and individual fire sizes. Second, fire size was further separated into fire duration and a daily expansion component. Third, the daily fire expansion was subdivided into fire speed, the length of the fire line, and the direction of spread. The Global Fire Atlas algorithm can be applied to any moderate resolution daily global burned area product, and the quality of the resulting dataset depends both on the Fire Atlas algorithm as well as the underlying burned area estimates. ~~MCD64A1 collection 6 burned area dataset (Giglio et al., 2018); for example, Here we applied the algorithm to the MCD64A1 collection 6 burned area dataset (Giglio et al., 2018) and~~ the minimum detected fire size is therefore one MODIS pixel (21 ha). Several studies have shown that the MCD64A1 col. 6 burned area product provides a considerable improvement compared to previous generation of moderate resolution global burned area products (Padilla et al., 2015; Giglio et al., 2018; Humber et al., 2018). We also present an initial effort to validate the higher order Global Fire Atlas products using independent fire perimeter data for the continental US and active fire detections to assess estimated fire duration and the temporal accuracy of individual fire dynamics.

100

105

110

115



**Figure 1: Flow chart showing the data-processing steps and resulting products.** (a) The Global Fire Atlas algorithm tracks individual fires and their day-to-day behavior based on the MCD64A1 collection 6 500 m daily burned area product starting in 2003. (b) Decomposition of burned area into seven different

120

components of the fire regime in the Global Fire Atlas. The output includes two annual shape file layers (.shp) of ignition location and individual fire perimeters with corresponding database files (.dbf) providing summary information for each individual fire, including the seven key characteristics (b). In addition, four global raster maps on the 500 m sinusoidal MODIS grid (.tif) provide details on the day-to-day fire behavior. Finally, data are summarized in a monthly 0.25° gridded product based on average values of individual fires. Global Fire Atlas data-layers are described in more detail in Table A1.

## 2.1 Individual fires: ignitions, size, perimeter and duration


Large burn patches are often made up of multiple individual fires that may burn simultaneously or at different points in time during the burning season, particularly in frequently burning grasslands and savannas with a high density of ignitions from human activity. Separating large clusters of burned area into individual fires is therefore critical to understand the fire regime in human-dominated landscapes. To isolate individual fires, clusters of adjacent burned area for a given fire season (12 months centered on the month of maximum burned area) were subdivided into individual fires based on the spatial structure of estimated burn dates in the MCD64A1 burned area product. Although we allow individual fires to burn from one fire season into the next, we processed the data on a per-fire-season basis in each 10° x 10° MODIS tile. In the rare case a pixel burned twice during a single fire season (<1%), we retained only the earliest burn date. This approach results in a small reduction of total burned area, but allows us to produce user friendly global annual layers in both gridded and shapefile format. This format allowed us to create global annual 500 m data layers with minimal loss of information. To locate candidate ignition points within each burned area cluster, we mapped the “local minima,” defined as a single grid cell or group of adjacent grid cells with the same burn date surrounded by grid cells with later burn dates. However, because of orbital coverage and cloud cover, burn date estimates are somewhat uncertain (Giglio et al., 2013), which results in many local minima that may not correspond to actual ignition points. We applied a three-step procedure to address burn date uncertainty and distinguish individual fires. First, we developed a filter to adjust the burn date of local minima that do not correspond to ignition points. Second, we set a “fire persistence” threshold that determines how long a fire may take to spread from one 500 m grid cell into the next, to distinguish individual fires that are adjacent but occurred at different times in the burning season. Third, we developed a second filter to correct for outliers in the burn date that occurred along the edges of large fires. Each of these steps is described in detail below.

The ignition point filter is based on the assumption that the fires progress in a logical manner through space and time. First, all local minima were mapped within the original field of burn dates (Fig. 2a and b). Next, each local minimum was replaced by the nearest later burn date in time of the surrounding grid cells, and a new map of local minima was created. If the original local minimum remained as a part of a new, larger local minimum with a later burn date, the fire followed a logical progression in time and space and the original local minimum was retained. If the local minimum disappeared, the original local minimum was likely the product of an inconsistency within the field of burn dates rather than a true ignition point and the burn date was adjusted forward in time to remove the original local minimum. This step can be repeated several times, with each new iteration further reducing the number of local minima and increasing the confidence in ignition points, but, each iteration may also result in greater adjustment of the original burn date (Fig. A1). Here we implemented three iterations of the ignition point filter to remove most local minima that did not spread forward in time while limiting the scope of burn date adjustments (e.g. Figs. 2c and d ~~and~~ A1 and A2). For short duration fires, the ignition points were retained associated with largest possible number of iterations. In addition In all cases, if several local minima were all connected through a single cluster of grid cells with the same burn date, only the local minimum with the earliest burn date or largest number of grid cells was retained, unless the required adjustment of the burn date was larger than the specified burn date uncertainty in the MCD64A1 product. By design, the ignition point filter cannot adjust the earliest burn date of a fire, and thus has no influence on estimated fire duration.

175 To establish the location and date of ignition points, as well as to track the daily growth and extent of  
individual fires, we used a “fire persistence” threshold that determines how long a fire may take to spread  
| A2A3). For example, if an ignition point was adjacent to a fire that burned earlier in the season, this  
threshold allowed the ignition point to be mapped as separate local minima despite the presence of  
180 adjacent burned grid cells with earlier burn dates. On the other hand, when an active fire is covered by  
dense clouds or smoke, multiple days can pass before a new observation can be made, resulting in a break  
in fire continuity and increasing the risk of artificially splitting single fires into multiple parts. Using such  
a threshold is particularly important to distinguish individual fires in frequently burning savannas and  
highly fragmented agricultural landscapes, where many individual small fires may occur within a  
relatively short time span. Because there are no reference datasets on global fire persistence, we used a  
185 spatially-varying fire persistence threshold that depends on fire frequency (Andela et al., 2017). We  
assumed that frequently-burning landscapes are generally characterized by faster fires and higher ignition  
densities, increasing the likelihood of having multiple ignition points within large burn patches, while  
infrequently burning landscapes will generally be characterized by slower fire spread rates and/or fewer  
ignitions. In addition, frequently burning landscapes often face a pronounced dry season characterized by  
190 low cloud cover, while infrequently burning landscapes may experience a shorter dry season with greater  
obscuration by clouds. Therefore, we used a 4-day fire persistence threshold for 500 m grid cells that  
burned more than 3 times during the study period (2003 - 2016), and a 6, 8 and 10-day fire persistence  
| period for grid cells that burned 3 times, 2 times~~2-3, 1-2~~, or 1 time, respectively. These threshold values  
broadly correspond to biomes, with shorter persistence values for tropical regions and human-dominated  
landscapes, and longer threshold values for temperate and boreal ecosystems with high fuel loads (Fig.  
195 A2).

Based on the location and date of the established ignition points and the fire persistence thresholds, we  
tracked the growth of each individual fire through time to determine its size, perimeter, and duration (Fig  
2f). For each day of year, we allowed individual fires to grow into the areas that burned on that specific  
200 day, as long as the difference in burn dates between two pixels was equal to or smaller than the fire  
persistence threshold of the pixel of origin. When two actively burning fires met each other, as on day 255  
for the example fires shown in Fig. 2, grid cells that burned on the day of the merger were divided based  
on nearest distance to the fire perimeter on the previous day.

205 Burn date uncertainty may also lead to multiple “extinction points,” outliers in the estimated day of burn  
along the edges of a fire. Environmental conditions such as cloud cover complicate the precise estimation  
of the date of fire extinction, as rainfall events extinguish many fires, and pixels at the edge of the fire  
may be partially burned and therefore harder to detect. In addition, the contextual relabeling phase of the  
MCD64A1 algorithm increases burn date uncertainty for extinction points based on a longer consistency  
210 threshold (Giglio et al., 2009). We used a second filtering step to adjust the burn date for extinction  
points, if required. Outliers were adjusted to the nearest burn date back in time, if (1) they represented a  
cluster no more than 1 to 4 grid cells ( $0.21 - 0.9 \text{ km}^2$ ) along the edge of a fire that was at least 10 times  
larger and (2) the difference in burn dates was larger than the fire persistence threshold of the adjacent  
grid cells and thus mapped as a new fire along the edge of the larger fire. If these criteria were met, the  
215 outliers were adjusted to the nearest burn date back in time, and incorporated within the larger  
neighboring fire. However, if these criteria were not met (e.g., for burned areas larger than 4 grid cells),  
the original burn dates and ignition points were left unadjusted, resulting in separate fires. For the  
example fires shown in Fig. 2, the adjustment of these outliers affected four grid cells (Fig. 2e) and  
effectively reduced the number of ignition points (and resulting individual fires) from five (Fig. 2d) to  
220 two (Fig. 2f). After adjusting these outliers (extinction points), and including them within the larger fires,  
we estimated the size ( $\text{km}^2$ ), duration (days) and perimeter (km) of each individual fire based on the  
adjusted burn dates.




225 **Figure 2: Example of the algorithm to account for uncertainty in the “day of burn” and identify**  
**individual fires within large clusters of adjacent burned pixels.** (a) The original MCD64A1 collection  
 6 day of burn for one burnt patch in the Brazilian Cerrado (2015), and (b) local minima or “ignition  
 230 points” identified within the original day of burn field. (c) Burn date adjustment based on the filter that  
 removes local minima that do not progress continuously through time and space (positive adjustment),  
 235 and (d) the corresponding estimate of ignition points based on the adjusted day of burn field. (e) Further  
 burn date adjustment based on the removal of outliers along the edge of the fire (negative adjustment of  
 extinction points), and (f) the final estimate of ignition locations and date by the Global Fire Atlas based  
 on the combined adjustments shown in (e). In (f), the colored lines indicate the final estimates of fire  
 perimeters.

## 2.2 Daily fire expansion: fire line, speed, and direction of spread

The revised day of burn estimates were used to track the daily expansion ( $\text{km}^2 \text{day}^{-1}$ ) and length of the fire  
 line (km) for each individual fire. The daily estimates of fire line length were based on the daily perimeter  
 240 of the fire, where we assumed that once the fire reaches the edge of the burn scar, this part of the  
 perimeter stops burning after one day (Fig. 3a). The expansion of the fire ( $\text{km}^2 \text{day}^{-1}$ ) is the area burned by  
 a fire each day. The average speed of the fire line ( $\text{km} \text{day}^{-1}$ ) can now be calculated as the expansion ( $\text{km}^2$   
 245  $\text{day}^{-1}$ ) divided by the length of the fire line (km) on the same day. However, this estimate of fire line  
 includes the head, flank and backfire, while it is typically the head-fire that moves fastest and may be  
 responsible for most of the burned area. Moreover, fire dynamics tend to be highly variable in space and  
 time. To understand the spatial variability and distribution of fire speeds, we therefore used an alternative  
 method to estimate the speed and direction of fire spread for each individual 500 m grid cell.

To estimate the speed and direction of spread (Fig. 3), we calculated the “most likely” path of the fire to  
 250 reach each individual 500 m grid cell based on shortest distance. More specifically, for each grid cell we  
 estimated the shortest route to connect the grid cell between two points: 1) the nearest point on the fire  
 line with the same day of burn and 2) the nearest point on the previous day’s fire line. This route was  
 forced to follow areas burned on the specific day. For each point on this route, or “fire path,” the speed of  
 the fire ( $\text{km} \text{day}^{-1}$ ) was estimated as the length of the path (km) divided by one day ( $\text{day}^{-1}$ ) and the  
 255 direction as the direction of the next grid cell on the fire path. Since each grid cell is surrounded by 8  
 other grid cells, this resulted in eight possible spread directions: north, northeast, east, southeast, south,

260 southwest, west, and northwest. For ignition points that represented a cluster of 500 m grid cells with the same burn date, we assumed that the fire originated in the center point of the cluster (pixel with largest distance to the final fire perimeter by the end of day 1) and spreads towards the perimeter of the fire by the end of day 1 over the course of one day. For single pixel fires, we assumed the fire burned across 463 m (1 pixel) during a single day and we did not assign a direction of spread. Similarly, fires of all sizes that burned on a single day were not assigned a direction of spread. We corrected estimates of both speed and direction for the orientation between 500 m grid cells on the MODIS sinusoidal projection that varies with location. When a particular grid cell formed part of multiple “fire paths,” the earliest time of arrival or the highest fire speed and corresponding direction of spread were retained. This assures a logical progression of the fire in time and space and corresponds to fires typically moving fastest in a principal direction and then spreading more slowly along the flank.



270 **Figure 3: Sub-daily estimates of fire progression can be used to estimate spatiotemporal variation in fire speed and direction of spread.** (a) daily progression of the fire line, (b) interpolated estimates of sub-daily time of arrival, (c) fire speed ( $\text{km day}^{-1}$ ), and (d) direction of spread. The light gray areas in (a) are burned areas between fire lines and correspond to areas of relatively high fire speed. White areas were not burned.

275

### 2.3 Validation

280 Few large-scale datasets are available on daily or sub-daily fire dynamics, highlighting the novelty of the Global Fire Atlas dataset but also posing challenges for validation. Here we used two alternative datasets for this purpose. First, we used active fire detections to assess the temporal accuracy of the Global Fire Atlas burn date. Second, we compared fire perimeters to independent fire perimeter data for the continental US. Finally Third, we combined the independent data on fire perimeters with active fire detections to evaluate the Global Fire Atlas fire duration estimates. Finally, we compared Global Fire Atlas data to a small (manually compiled) dataset of daily fire perimeters from the US Forest Service.

285 We used the 375 m resolution active fire detections (VNP14IMGML C1) derived from the Visible  
Infrared Imaging Radiometer Suite (VIIRS) instrument aboard the Suomi National Polar-orbiting  
Partnership (Suomi-NPP) satellite (Schroeder et al., 2014). Active fire detections provide accurate  
290 information on the burn date, particularly in ecosystems with low fuel loads where fires will typically be  
active during only a single day in each particular grid cell. We compared the date of active fire detections  
from VIIRS within each larger 500 m MODIS grid cell (based on VIIRS center point) to the adjusted  
MCD64A1 day of burn to understand the temporal precision of the derived Global Fire Atlas products. If  
several active fire detections were available for a single 500 m MODIS grid cell we used the date closest  
295 to the mean. We compared all 500 m MODIS grid cells with corresponding active fire detection during  
the overlapping data period (2012 – 2016) for four different ecosystems globally: (1) forests (including all  
forests), (2) shrublands (including open and closed shrublands), (3) woody savannas and (4) savannas and  
grasslands, with land cover type derived from MODIS MCD12Q1 collection 5.1 data for 2012 using the  
University of Maryland (UMD) classification (Friedl et al., 2002).

300 We compared fire perimeters from the Global Fire Atlas to fire perimeter estimates from the Monitoring  
Trends in Burn Severity (MTBS) project during their overlapping period (2003 – 2015). The MTBS  
project provides semi-automated estimates of fire perimeters based on 30 m Landsat data for fires with a  
minimum size of 1000 acres (405 ha) in the western US and 500 acres (202 ha) in the eastern US  
(Eidenshink et al., 2007; Sparks et al., 2015). In order to determine overlap between MTBS and Fire Atlas  
305 perimeter estimates, we rasterized the MTBS perimeters onto the 500 m MODIS sinusoidal grid including  
all 500 m grid cells with their center point within the higher resolution (30 m) MTBS fire perimeter. For  
all overlapping fire perimeters, we compared the original MTBS fire perimeter information with the Fire  
Atlas estimates of fire perimeters. In cases with multiple overlapping perimeters, fires with the largest  
overlapping surface area were compared.

310 We also combined MTBS fire perimeters with VIIRS active fire detections to derive an alternative  
estimate of fire duration (2012 – 2015). In order to determine the fire duration, we first determined the  
median burn date of each fire according to the MCD64A1 burned area data. Subsequently, we included all  
VIIRS active fire detections before and after the median or ‘center’ burn date until a period of three fire-  
315 free days was reached. Any active fire detections that occurred outside this timeframe were excluded to  
avoid overestimation of the fire duration due to smoldering or possible false detections before or after the  
fire. Two thresholds were used to select a subset of MTBS and Fire Atlas perimeters for validation of  
estimated fire duration. Fires were first matched based on perimeters, with maximum of a threefold  
difference between perimeters. Second, we further selected MTBS perimeters with VIIRS active fire  
320 detections for at least 25% of the 500 m Fire Atlas grid cells. These thresholds excluded 51% of the  
overlapping fire perimeters, but reduced errors originating from cloud cover or differences in the  
underlying burned area estimates (e.g., resolution, methodology) to evaluate estimated fire duration.  
Similar to the burn date validation, comparisons of fire perimeters and fire duration with MTBS data over  
the continental US were grouped into four land cover types: (1) forests, (2) shrublands, (3) woody  
325 savannas and (4) savannas and grasslands.


330 For specific large wildfires across the western USA, the US Forest Service National Infrared Operations  
(NIROPS; <https://fsapps.nwcg.gov/nirops/>) derives estimates of daily fire perimeters for fire management  
purposes by collecting night-time high resolution infrared imagery. This imagery is manually analyzed by  
trained specialists to extract the active fire front. Although these data provide a wealth of information,  
only few fires were completely and precisely documented. From their database we were able to extract 15  
large fires for which daily perimeter information was available. Although insufficient for full scale  
validation, results provide valuable insights into the strengths and shortcomings of the Global Fire Atlas  
estimates of individual fire size, duration and expansion rates. In addition, we compared day-to-day  
335 expansion rates ( $\text{km}^2 \text{ day}^{-1}$ ) of individual large fires across both datasets. If multiple Global Fire Atlas



perimeters overlapped with a single US Forest Service fire perimeter, we compared the fires with the largest overlapping surface area.

### 3 Results

#### 340 3.1 Validation




345 **Figure 4: Per pixel global Comparison-comparison of burn dates derived from the MCD64A1 burned area product, adjusted burn dates of the Global Fire Atlas, and VIIRS active fire detections (2012 – 2016).** (a) Forests, (b) shrublands, (c) woody savannas, and (d) savannas and grasslands. Negative values indicate pixels with a burned area day of burn earlier than the first corresponding VIIRS active fire detection, zero indicates no difference in day of burn between both datasets, and positive numbers indicate a delayed detection of burned area compared to active fire detections.

350 At the pixel scale, estimated burn dates from burned area and active fire products were comparable (Fig. 4), with greater variability across biomes than from minor burn date adjustments in the Global Fire Atlas algorithm. Burn dates estimated from MODIS burned area and VIIRS active fire detections were least comparable in high-biomass ecosystems with lower fire spread rates. In forests and woody savannas 24% and 35% of burned pixels were detected on the same day and 54% and 67% within  $\pm 1$  day, respectively

355 (Fig. 4a and c). With decreasing biomass, the direct correspondence between burn dates from burned area  
 and active fire detections increased to 41% (same day) and 80% ( $\pm 1$  day) in shrublands (Fig. 4b) and  
 40% (same day) and 75% ( $\pm 1$  day) in savannas and grasslands (Fig. 4d). These differences likely stem  
 360 from the combined increase in uncertainty of burn date in higher-biomass ecosystems and influence of  
 fire persistence (multiple active fire days in a single 500 m grid cell) on the ability to reconcile the timing  
 of burned area and active fire detections in these ecosystems. Several factors may account for the positive  
 bias in the 500 m day of burn from burned area compared to active fire detections, including orbital  
 coverage, cloud and smoke obscuration, and different thresholds between burned area and active fire  
 algorithms regarding the burnt fraction of a 500 m grid cell. The adjustments made to the burn date here,  
 365 required to effectively determine the extent and duration of individual fires, had a relatively small effect  
 on the overall accuracy but tended to reduce the negative bias in burn dates and increase the positive bias  
 (i.e. delayed burn date compared to active fire detection, see red and black lines in Fig 4). In line with  
 these findings, we found good agreement between a 3-day running average of Global Fire Atlas and US  
 Forest service estimates of daily fire expansion, but reduced correspondence for daily estimates of fire  
 growth rates due to uncertainty in the day-of-burn of the burned area product (Fig. B1).

370




**Figure 5: Comparison of fire perimeter estimates based on the Global Fire Atlas and MTBS for the continental US (2003 – 2015).** (a) Forests, (b) shrublands, (c) woody savannas, and (d) savannas and grasslands. Red lines indicate the slope between both datasets based on ordinary least squares (OLS) with corresponding  $r^2$  values, while blue lines are based on orthogonal distance regression (ODR). For the scatter plots, darker gray or black indicates a greater density of points.

375

380 For fire perimeters, the best agreement between the Global Fire Atlas and MTBS was found in forests and  
 385 shrublands, where the Global Fire Atlas reproduced 65% and 61% of the observed variance in MTBS fire  
 390 perimeters, respectively (Fig. 5). Less agreement was found for woody savannas (38%) and savannas and  
 grasslands (41%). However, uncertainty exists in both datasets. Orthogonal distance regression (ODR)  
 accommodates uncertainties in both datasets and generally resulted in slopes closer to the 1:1 line,  
 indicating closer correspondence, on average, in absolute perimeter estimates for the two datasets. An in-  
 depth comparison of the performance of the Global Fire Atlas and the MTBS datasets for several  
 grassland fires in Kansas (USA) suggested that differences originated both from the underlying burned  
 area datasets and the methodologies (Fig. B+B2). For this particular grassland in Kansas, the MCD64A1  
 product estimated less burned area compared to the Landsat-based MTBS dataset, resulting in  
 fragmentation of larger burn scars into disconnected patches. However, the daily temporal resolution of  
 the MCD64A1 burned area product allowed for recognition of individual ignition points within larger  
 burn patches of fast moving grassland fires that cannot be separated using infrequent Landsat imagery  
 (Fig. B2). In addition, the 30 m spatial resolution of the MTBS perimeters may result in more irregularity  
 and therefore in longer fire perimeter estimates compared to the 500 m Fire Atlas perimeters (Fig. B1).  
 Combined, these tradeoffs in spatial and temporal resolution resulted in less agreement between fire  
 perimeters in woody savannas (Fig. 5c) and savannas and grasslands (Fig. 5d).

395



400 **Figure 6: Comparison of fire duration estimates from the Global Fire Atlas and the combination of VIIRS active fire detections within MTBS fire perimeters for the continental US (2012 – 2015).** (a) Forests, (b) shrublands, (c) woody savannas, and (d) savannas and grasslands. Red lines indicate the slope between both datasets based on ordinary least squares (OLS) with corresponding  $r^2$  values, while blue


lines are based on orthogonal distance regression (ODR). For the scatter plots, darker gray or black indicates a greater density of points. This comparison used a subset of MTBS and Fire Atlas perimeters based on selection criteria for perimeter overlap and VIIRS active fire detections (see Section 2.3).

405 Initial validation of fire duration estimates from the Global Fire Atlas highlighted the differences in the sensitivity of satellite-based burned area and active fire products to fire lifetime (Fig. 6). Similar to fire perimeters, the best agreement in fire duration estimates was found for forests, where the Global Fire Atlas reproduced 51% of the observed variance of the fire duration estimates based on combining MTBS fire perimeters with active fire detections. Shrublands, woody savannas, and savannas and grasslands had lower correlations, with 27%, 30% and 33% of the variance explained, respectively. The orthogonal distance regression resulted in slopes close to the one-to-one line for shrublands and savannas and grasslands, indicating reasonable agreement. Fire duration was clearly underestimated for forested ecosystems with high fuel loads, as fires may continue to smolder for days (resulting in active fire detections) after the fire has stopped expanding.

415 The comparison of Global Fire Atlas data to a small dataset (n = 15) of daily perimeters of large wildfires in primarily forested cover types mapped by the US Forest Service yielded good correspondence between estimates of fire size, duration, and expansion rates (Fig. 7). The improved comparison of fire size (cf. Fig. 5a and 7a) could be related to the US Forest Service data being more accurate than MTBS, but likely also represents the good performance of the Global Fire Atlas (e.g. compare Figs. 7a, b and c to Figs. 7d, e and f) and underlying burned area products (Fusco et al., 2019) for relatively large fires. In contrast to the suggested underestimate of fire duration shown in Fig. 6a, these data suggest the Global Fire Atlas may slightly overestimate fire duration. This difference may reflect the fact that active fire detections may be triggered by smoldering while the burned area product will only register the initial changes in surface reflectance from fire. Based on a small underestimate of overall burned area and overestimate of fire duration by the Global Fire Atlas, the average daily fire expansion rates based on US Forest Service data were higher than estimates based on Global Fire Atlas data (Fig. 7c and f).

420

425




430 **Figure 7: Comparison of Global Fire Atlas (GFA) and US Forest Service (FS) data for a selected**  
**number of large wildfires in the US. Comparison of (a) fire size, (b) duration, and (c) average daily**  
**expansion rate for all fires (N=15). (d, e and f) are like (a, b and c) but for fires smaller than 250 km<sup>2</sup>**  
**(N=12). Correlation coefficients are provided based on linear regression with (yellow) and without**  
**(green) intercept, assuming a non-zero intercept could indicate a structural offset between both datasets.**


435

### 3.2 Characterizing global fire regimes

Over the 14-year study period we identified 13,250,145 individual fires with an average size of 4.4 km<sup>2</sup> (Table 1) and minimum size of one MODIS pixel (21 ha or 0.21 km<sup>2</sup>). On average, largest fires were  
 440 found in Australia (17.9 km<sup>2</sup>), boreal North America (6.0 km<sup>2</sup>), and northern hemisphere Africa (5.1 km<sup>2</sup>), while central America (1.7 km<sup>2</sup>), equatorial Asia (1.8 km<sup>2</sup>), and Europe (2.0 km<sup>2</sup>) had the smallest average fire sizes (Table 1). Spatial patterns of number of ignitions and fire sizes were markedly different and often inversely related (Fig. 78). Burned area in agricultural regions and parts of the humid tropics, particularly in Africa, resulted from high densities of fire ignitions and relatively small fires, consistent  
 445 with widespread use of fire for land management. Large fires accounted for most of the burned area in arid regions, high latitudes, and other natural areas with low population densities and a sufficiently long season of favorable fire weather (Fig. 78).



450 | **Figure 78:** Average global burned area (MCD64A1), ignition density, and fire size over the study  
 period 2003 – 2016. For any given area (a) Burned-burned area in km<sup>2</sup> per year is-would be the product  
 of (b) ignitions per year and (c) fire size in km<sup>2</sup>. However, because the size of a 0.25° grid cell varies with  
 latitude we have converted the units of burned area to fraction (%) per year and of ignitions to number per  
 455 | km<sup>2</sup> per year for spatial consistency.




**Figure 89:** Average fire duration (a), fire line length (b), and daily expansion (c) over the study period 2003 – 2016. Fire size (see Fig. 7c) is the product of fire duration (a) and daily fire expansion (c).

460 Global patterns of fire duration and expansion rates provide new insights in the occurrence of large fires, as the size of each fire ( $\text{km}^2$ ) is the product of fire duration (days) and daily fire expansion rate ( $\text{km}^2 \text{day}^{-1}$ ). Individual fires that burned for a week or more occurred frequently across the productive tropical grasslands and in boreal regions (Fig. 8a9a, Table 2). In these regions, fire duration exerted a strong control on fire size and total burned area. On average, human-dominated landscapes such as deforestation frontiers or agricultural regions experienced smaller and shorter fires compared to natural landscapes (Table 2). Fire duration was also relatively short in semiarid grasslands and shrublands characterized by high daily fire expansion rates, based on the development of long fire lines (Fig. 8b-9b and c) and high velocity. In these regions, fire duration and size were likely limited by fuel connectivity. In line with these findings, largest average daily expansion rates were found in Australia ( $1.7 \text{ km}^2 \text{day}^{-1}$ ), northern hemisphere Africa ( $0.9 \text{ km}^2 \text{day}^{-1}$ ) and southern hemisphere Africa ( $0.9 \text{ km}^2 \text{day}^{-1}$ ), and smallest expansion rates in central America ( $0.3 \text{ km}^2 \text{day}^{-1}$ ), equatorial Asia ( $0.3 \text{ km}^2 \text{day}^{-1}$ ), and southeast Asia ( $0.4 \text{ km}^2 \text{day}^{-1}$ ; Table 1).

465  
470





475 | **Figure 910:** Average speed of the fire (a) and the dominant direction of fire spread (b) over the  
 study period 2003 – 2016. For each 0.25° grid cell the direction was estimated as the dominant fire  
 spread direction of fires larger than 10 km<sup>2</sup> within the grid cell. We focused on larger fires (≥ 10 km<sup>2</sup>) to  
 determine the dominant spread direction, because large fires will generally express a clearer  
 spatiotemporal structure of fire spread at 500 m daily resolution. Pie charts show the fraction of individual  
 480 larger fires (≥ 10 km<sup>2</sup>) by dominant spread direction for each continent.

| The fastest fires occurred in arid grasslands and shrublands (Fig. 910a), where fuel structure, climate  
 conditions, and emergent properties of large wildfires contribute to high fire spread rates. Relatively high  
 fire speeds were also observed in some parts of the boreal zone, particularly in central and western  
 485 Canada. Lowest fire velocities were observed in infrequently burning humid tropical regions where fire  
 spread was influenced by higher fuel loads and humidity (Table 1). At all scales, estimated fire direction  
 exhibited considerable complexity (Fig. 910b). With some regional exceptions, no clear dominant spread  
 direction was found in South America or Africa. Based on the underlying 500 m data layers, landscape  
 structure and drainage patterns played an important role in controlling individual fire spread direction in  
 490 the humid tropics. Fire spread direction also varied considerably within individual fires, and the dominant  
 direction typically represented less than half of the pixels. Fire spread direction was more consistent in the  
 arid tropics, as demonstrated by the northwest and southeast orientation of fire spread in Australia,  
 consistent with the dominant wind directions. At mid-latitudes, we found evidence for more east and  
 westward fire progression in Europe and Asia and northwest and southeast spread direction in North  
 495 America, broadly consistent with the orientation of mountain ranges and other topographic features within  
 the key biomass burning regions.

500

**Table 1: Fire attributes for each Global Fire Emissions Database (GFED) region during 2003 – 2016.** Ignitions are the summed ignitions over the study period (2003 – 2016). For size, duration, expansion, and speed the mean values are shown for individual fires and weighted by fire size (between parenthesis). For ignitions, regions with over one million ignitions are shown in red and lower values in blue, for other fire aspects values equal to or above the global average are shown in red and below the global average in blue. A map of the GFED regions is shown in the annex material (Fig. [B2aB3a](#)).

GFED Region	Ignitions (2003-2016)	Size (km <sup>2</sup> )	Duration (days)	Expansion (km <sup>2</sup> day <sup>-1</sup> )	Speed (km day <sup>-1</sup> )
World	13250145	4.4 (395.9)	4.5 (14.7)	0.6 (14.5)	0.9 (3.2)
BONA	57613	6.0 (202.8)	5.4 (23.3)	0.5 (6.8)	1.0 (4.3)
TENA	137900	2.9 (136.7)	4.7 (13.4)	0.5 (8.8)	0.8 (3.7)
CEAM	229245	1.7 (28.3)	4.3 (12.2)	0.3 (1.5)	0.7 (1.4)
NHSA	242359	3.1 (50.1)	5.1 (12.4)	0.5 (3.3)	0.8 (2.1)
SHSA	1320177	3.0 (90.6)	4.7 (13.8)	0.5 (4.8)	0.7 (2.3)
EURO	71233	2.0 (30.7)	4.6 (10.3)	0.4 (2.7)	0.7 (2.0)
MIDE	86783	2.3 (22.0)	4.0 (9.8)	0.5 (2.1)	0.8 (1.9)
NHAF	3517808	5.1 (186.2)	4.4 (14.7)	0.7 (8.6)	0.9 (3.0)
SHAF	5000436	4.3 (232.5)	4.5 (13.5)	0.7 (9.6)	0.9 (2.6)
BOAS	363279	3.7 (116.8)	4.5 (15.6)	0.5 (6.8)	1.0 (4.1)
CEAS	807739	3.2 (339.7)	4.2 (11.5)	0.5 (22.7)	0.8 (5.6)
SEAS	937810	2.2 (27.8)	4.1 (13.2)	0.4 (1.8)	0.7 (1.8)
EQAS	117870	1.8 (13.5)	5.5 (16.4)	0.3 (0.8)	0.7 (1.3)
AUST	358807	17.9 (2030.6)	5.0 (20.5)	1.7 (59.5)	1.2 (6.1)

505

**Table 2: Fire attributes by GFED fire type during 2003 – 2016.** Ignitions are the summed ignitions over the study period (2003 – 2016). For size, duration, expansion, and speed, the mean values are shown for individual fires and weighted by fire size (between parenthesis). For agriculture, we only included fires with >90% of burned area classified as cropland. For ignitions, fire types with over one million ignitions are shown in red and lower values in blue, for other fire aspects values equal to or above the global average are shown in red and below the global average in blue. A map of the GFED fire types is shown in the annex material (Fig. [B2bB3b](#)).

510


GFED fire type	Ignitions (2003-2016)	Size (km <sup>2</sup> )	Duration (days)	Expansion (km <sup>2</sup> day <sup>-1</sup> )	Speed (km day <sup>-1</sup> )
All	13250145	4.4 (395.9)	4.5 (14.7)	0.6 (14.5)	0.9 (3.2)
Boreal forest	197124	5.2 (149.2)	5.4 (20.1)	0.6 (6.5)	1.0 (4.2)
Temporal forest	178909	2.5 (84.1)	4.1 (14.0)	0.4 (4.2)	0.8 (2.8)
Deforestation	909826	1.4 (28.7)	3.8 (13.7)	0.3 (1.4)	0.6 (1.4)
Savanna	9809719	5.1 (447.5)	4.6 (14.9)	0.7 (16.2)	0.9 (3.4)
Agriculture	1631918	1.4 (26.4)	3.4 (10.3)	0.3 (2.0)	0.7 (1.9)

### 3.3 Fire extremes

515

The world's largest individual fires were mostly found in sparsely populated arid and semiarid grasslands and shrublands of interior Australia, Africa, and central Asia (Fig. [10a11a](#)). Strikingly, fires of these proportions were nearly absent in North and South America, possibly due to higher landscape fragmentation and different management practices, including active fire suppression. In arid regions of

520 Southern Africa and Australia, large fires typically followed La Niña periods (e.g., 2011 and 2012), when  
 increased rainfall and productivity increase fuel connectivity (Chen et al., 2017). The largest fire in the  
 Global Fire Atlas occurred in northern Australia, burning across 40,026 km<sup>2</sup> (about the size of  
 Switzerland or the Netherlands) over a period of 72 days with an average speed of 19 km day<sup>-1</sup>, following  
 the 2007 La Niña. The longest fires burned for over 2 months in seasonal regions of the humid tropics and  
 high-latitude forests (Fig. 40b11b). Drought conditions in 2007 and 2010 caused multiple fires to burn  
 525 | synchronously for over two months across tropical forests and savannas in South America. Highest fire  
 velocities typically occurred in areas of low fuel loads. While fires larger than 2500 km<sup>2</sup> were nearly  
 absent from arid grass and shrublands in the North and South America, patterns of extremely fast-moving  
 fires in arid grass and shrublands were similar to other continents. Fast-moving fires also show evidence  
 of synchronization, for example with several extremely fast fires burning across the steppe of eastern  
 530 | Kazakhstan during 2003 (Fig. 40e11c).



535 | **Figure 4011:** Location and year of the largest, longest, and fastest fires over the study period 2003 –  
 2016. (a) fires larger than 2500 km<sup>2</sup>, (b) fires longer than 60 days, and (c) fires with an average velocity  
 larger than 25 km day<sup>-1</sup>. The background image depicts mean MODIS normalized difference vegetation  
 index (NDVI, 2003 – 2016), an indicator for large scale vegetation patterns and available fuels.

## 4 Discussion

540 The Global Fire Atlas is the first freely available global dataset to provide daily information on seven key  
fire characteristics: ignition timing and location, fire size, duration, daily expansion, daily fire line, speed  
and direction of spread based on moderate resolution burned area data. Over the 2003 – 2016 study  
period, we identified nearly one million individual fires ( $\geq 21$  ha) each year (Table 1). Characteristics of  
545 | these fires varied widely across ecosystems and land use types. In arid regions and other fire-prone  
natural landscapes, most of the burned area resulted from a small number of large fires (Fig. 78). Fire  
sizes declined along gradients of increasing rainfall and human activity, with larger numbers of small fires  
in the humid tropics or other human-dominated landscapes. Multiday fires were the norm across nearly all  
landscapes, with some large fires in productive tropical grasslands and boreal regions burning for over  
550 | two months during drought periods (Fig. 1011). The dominant control on fire sizes also varied across  
ecosystems; fire duration was the principal control on fire sizes in boreal forests, whereas fuels limited the  
size of fast-moving fires in arid grasslands and shrublands (Figs. 8-9 and 109). Characterizing fire  
behavior across large scales is key for understanding fire-vegetation feedbacks, emissions estimates, fire  
prediction, effective fire management, and modeling of fires within ecosystem models. Satellite remote  
555 sensing has been widely used to characterize global pyrogeography (Archibald et al., 2013) and fire-  
climate interactions (Westerling et al., 2006; Alencar et al., 2011; Morton et al., 2013a; Field et al., 2016;  
Young et al., 2017). Nonetheless, large-scale understanding of individual fire behavior has remained  
elusive without consistent global data products such as the Global Fire Atlas.

560 Both climate and human activity exert a strong control on global burned area (Bowman et al., 2009) and  
contribute to rapidly changing fire regimes worldwide (Jolly et al., 2015; Andela et al., 2017; Earl and  
Simmonds, 2018). Moreover, increasing human presence in fire prone ecosystems requires increased  
efforts to actively manage fires for ecosystem conservation and human wellbeing (Moritz et al., 2014;  
Knorr et al., 2016). The ignition location, spread, and duration of individual fires can be used to address  
565 | new questions of fire-climate interactions and changing influence of human activity on fire behavior, as  
each of these aspects may respond differently to variability or change. For example, recent studies have  
suggested that climate warming and drying may increase fire size and burned area in the tropics (Hantson  
et al., 2017) and at higher latitudes (Yang et al., 2015). Our findings suggest that an increase in the length  
of the fire season may be the dominant driver for increases in fire activity in these ecosystems, as fire  
570 | duration was a strong control on eventual fire sizes and burned area (Figs. 87, 98 and 110). Investigating  
fire-climate interactions and human controls on burned area using the Fire Atlas data layers will benefit  
management efforts and science investigations, as fires alter vegetation structure (Bond et al., 2005;  
Staver et al., 2011), biogeochemical cycles (Bauters et al., 2018; Pellegrini et al., 2018) and climate  
(Randerson et al., 2006; Ward et al., 2012).

575 The Global Fire Atlas provides several new constraints that could improve the representation of fires in  
ecosystem and Earth system models. Fire models embedded in dynamic vegetation models are important  
tools for understanding the changing role of fires in the Earth system and the ecosystem impacts of fires  
(Hantson et al., 2016; Rabin et al., 2017). Most global models of fire activity are calibrated using satellite-  
580 | derived estimates of total burned area or active fires (Hantson et al., 2016), rather than individual fire  
characteristics. As a result, these fire models capture the spatial distribution of global fire activity but not  
burned area trends (Andela et al., 2017) or interannual variability that may increase fire spread rates or  
duration. Models range from simple empirical schemes to complex, process-based representations of  
individual fires (Hantson et al., 2016; Rabin et al., 2017). Process-based models estimate burned area as  
585 | the product of fire ignitions and size, while many models include a dynamic rate of spread to determine  
eventual fire sizes (e.g. SPITFIRE; Thonicke et al., 2010) but use arbitrary threshold values for key  
parameters such as fire duration (Hantson et al., 2016). We found that global patterns of fire duration,  
ignitions, size, and rate of spread (i.e. speed) varied widely across ecosystems and human land

management types, and thus these Global Fire Atlas data products provide additional pathways to benchmark models of various levels of complexity. While only a few models include multiday fires (e.g.,  
590 Pfeiffer et al., 2013; Le Page et al., 2015; Ward et al., 2018), we found that multiday fires were the norm across most biomes, and fire duration forms an important control on eventual fire sizes and burned area in many natural ecosystems with abundant fuels. In a similar fashion, many models assume relatively homogeneous fuel beds, while our results suggest that landscape features and vegetation patterns result in highly heterogeneous fuel beds that form a strong control on fire spread (speed and direction). Large differences in fire behavior across ecosystems and management strategies may improve fire emissions estimates and emissions forecasting, particularly when combined with active fire detections to better characterize different fire stages including the smoldering phase (Kaiser et al., 2012). Recent studies have shown that fire emissions factors may vary widely depending on fire-behavior (Van Leeuwen and Van Der Werf, 2011; Parker et al., 2016; Reisen et al., 2018), while improved knowledge of fire-climate interactions are crucial for emissions forecasting (Di Giuseppe et al., 2018).

The Global Fire Atlas methodology builds on a range of previous studies that have used remote sensing  
605 daily moderate resolution satellite imagery to estimate individual fire sizes (Archibald and Roy, 2009; Hantson et al., 2015; Frantz et al., 2016; Andela et al., 2017), shape (Nogueira et al., 2017; Laurent et al., 2018), duration (Frantz et al., 2016) and spread dynamics (Loboda and Csiszar, 2007; Coen and Schroeder, 2013; Sá et al., 2017). We provide the first fire progression-based algorithm to map individual fires across all biomes, including the first global estimates of ignition locations and timing, duration, daily expansion, fire line, speed and direction of spread. Several previous studies have estimated fire size distributions based on a flood-fill algorithm, where all neighboring pixels within a certain time threshold  
610 are classified as the same fire (Archibald and Roy, 2009; Hantson et al., 2015). Interestingly, we found similar spatial patterns of fire size (cf. Fig. 8 and Archibald et al., 2013; Hantson et al., 2015), although absolute estimates may show large differences based on the “cut off” value used within the flood-fill approach (Oom et al., 2016), and to a lesser extent by the fire persistence threshold used here. Spatial patterns of fire size and duration also compared favorably with estimates of Frantz et al. (2016) for southern Africa (Fig. 8a9a) and estimates of fire speed by Loboda et al. (2007) for central Asia (Fig. 9a10a). Here we compared our results to fire perimeter estimates from the MTBS (Eidenshink et al., 2007; Sparks et al., 2015) for validation purposes. Good agreement was found for forested ecosystems and shrublands, but results differed more in grassland biomes (Fig. 5). Interestingly, we found that the poor agreement in grasslands stemmed from differences in the spatial and temporal resolution of the burned area estimates (Fig. B2). In line with previous studies, we found that ~~While~~ the coarser resolution (500 m) of the MODIS burned area data used to develop the Global Fire Atlas sometimes underestimated overall burned area (e.g. Randerson et al., 2012; Roteta et al., 2019), fragmented-fragmenting individual large fires. However, the Landsat-based MTBS data at 30 m resolution were unable to distinguish individual fires within large burn patches of fast-moving grassland fires based on infrequent Landsat  
620 satellite overpasses (Fig. B1B2).

Validation of Global Fire Atlas fire perimeter estimates for the continental US revealed several important limitations and opportunities for further development of individual fire characterization using satellite burned area data. In addition to the validation of fire perimeters, we also investigated the temporal accuracy of the Global Fire Atlas (Fig. 4) as well as the fire duration estimates (Fig. 6) based on active fire detections. Reasonable correlations ( $r^2$  ranging from 0.3 to 0.5) were found between Global Fire Atlas and fire duration estimates based on a combination of MTBS fire perimeters and VIIRS active fire detections. Disagreement partly originated from differences in fire perimeter estimates as well as differences between the day-of-burn estimates derived from the MCD64A1 burned area data and VIIRS active fire detections. Moreover, the uncertainty in the burn date of the underlying burned area product is typically at least one day, resulting in a large uncertainty in the fire duration estimates of shorter fires.  
630 Global Fire Atlas data therefore performed best for large fires (Figs. 6 and 7). Particular care should be taken when using the Global Fire Atlas for cropland regions for two main reasons. First, mapping burned

640 area in croplands is notoriously difficult using moderate resolution satellite data, as typical crop residue  
burning is often too small to detect (Randerson et al., 2012; Giglio et al., 2013). Second, we allow a fire 4  
– 10 days to spread from one grid cell into the next (fire persistence threshold), which may be more  
representative for natural landscapes than croplands with synchronized small fire activity at specific  
645 points in the crop cycle. The temporal accuracy of the Global Fire Atlas adjusted burned area compared to  
VIIRS active fire detections ranged from 41% on the same day and 80% within  $\pm 1$  day in shrublands to  
and 24% (same day) and 54% ( $\pm 1$  day) in forests. However, in forested ecosystems the use of active fire  
detections for validation purposes is not ideal, as fires may smolder for days resulting in active fire  
detections a long time after the fire front has passed. Understanding the temporal accuracy of the Global  
Fire Atlas products is important for linking individual fire dynamics to fire weather, and we found good  
650 agreement between Global Fire Atlas and US Forest Service fire expansion using a 3-day running  
average, but less good agreement for individual days based on burn date uncertainty (Fig. B1). Other  
parameters, including fire speed and direction of spread, were not validated during this stage. However,  
our comparison to daily fire perimeter estimates from the US Forest Service show good agreement in  
terms of average expansion rates, suggesting reasonable overall estimates of speed (Fig. 7). Overall, there  
is a need to develop additional validation methodologies and data products to advance our understanding  
655 of satellite-derived estimates of individual fire behavior, building on the long-standing efforts for burned  
area (Boschetti et al., 2009) and active fires (Schroeder et al., 2008).

The Global Fire Atlas provides the first consistent, global assessment of individual fire behavior. Further  
development of the Fire Atlas product suite is possible based on improvements in the underlying burned  
660 area data, including new products at higher spatial resolution (e.g., VIIRS), and additional constraints  
from active fire detections. The Global Fire Atlas algorithm provides a flexible framework that can be  
easily adjusted to work at different spatial and/or temporal resolutions. In particular, daily burned area  
products do not resolve the diurnal cycle of fire activity, that may vary widely across fire regimes  
(Freeborn et al., 2011; Andela et al., 2015). A better understanding of the drivers of fire persistence and  
665 fuel loads across biomes and ecosystem gradients is also important.

## 5 Data availability

670 The data are freely available at <http://www.globalfiredata.org> in standard data product formats and  
updates for subsequent years will be distributed pending availability of MCD64A1 burned area data and  
associated research funding. Global per-fire-year shapefiles of the ignition locations (point) and  
individual fire perimeters (polygon) contain attribute tables with a unique fire ID, ignition location, start  
and end dates, size, duration, and average values of the daily expansion, daily fire line, speed, and  
675 direction of spread (Fig. 1, Table A1). In addition, gridded 500 m global maps of the Global Fire Atlas  
adjusted burn dates, daily fire line, speed and direction of spread are available in GeoTIFF format. A  
monthly gridded product is also available at 0.25° resolution. Global Fire Atlas data products can also be  
visualized and evaluated using an online tool at [globalfiredata.org](http://globalfiredata.org) to explore individual fire characteristics  
for a selected region of interest.

## 680 6 Conclusions

The Global Fire Atlas is a new publicly available global dataset on seven key fire characteristics: ignition  
location and timing, fire size, duration, daily expansion, daily fire line, speed, and direction of spread.  
Over the 2003 – 2016 study period, we identified 13,250,145 individual fires ( $\geq 21$  ha) based on the  
685 moderate resolution MCD64A1 collection 6 burned area data. Striking differences were observed among  
global fire regimes along gradients of ecosystem productivity and human land use. In general, in  
ecosystems of abundant fuel and low human influence, large fires of long duration dominated total burned


690

area, with large numbers of small fires contributing most to overall burned area in human-dominated regions or areas too wet for frequent fires. Fires moved quickly through arid ecosystems with low fuel densities but fire sizes were eventually limited by fuels from natural or human landscape fragmentation. The dataset enables new lines of investigation for understanding vegetation-fire feedbacks, climatic and human controls on global burned area, fire forecasting, emissions modeling, and benchmarking of global fire models.

700 **Table A1: Overview of the Global Fire Atlas data-layers.** The shapefiles of ignition locations (point) and fire perimeters (polygon) contain attribute tables with summary information for each individual fire, while the underlying 500 m gridded layers reflect the day-to-day behavior of the individual fires. In addition, we provide aggregate monthly layers at 0.25° resolution for regional and global analyses.


	Shapefile attributes*	500 m daily gridded	0.25° monthly gridded
<b>Ignitions</b>	location and timing	-	sum
<b>Perimeter (km)</b>	per fire	-	-
<b>Size (km<sup>2</sup>)</b>	per fire	-	average
<b>Duration (days)</b>	per fire	-	average
<b>Daily fire line (km)</b>	average per fire	yes	average
<b>Daily fire expansion (km<sup>2</sup> day<sup>-1</sup>)</b>	average per fire	-	average
<b>Speed (km day<sup>-1</sup>)</b>	average per fire	yes	average
<b>Direction of spread (-)</b>	dominant per fire	yes	dominant
<b>Day of burn</b>	-	yes	-

\* vector data are derived from the underlying 500 m MODIS data.



705 **Figure A1: Burn date adjustment to remove local minima that are not associated with ignition points.** (a) MCD64A1 burn date estimate for the 2015 example fires in the Cerrado, (b) local minima within (a). (c) Burn date adjustment after the first iteration, and (d) resulting local minima. (e) Burn date adjustment after the second iteration, and (f) resulting local minima. (g) Burn date adjustment after the third iteration, and (h) resulting local minima. Note that for these particular fires there was no difference between (e and f) and (g and h), and the final iteration has no added value here. We found that multiple  
710 iterations were particularly beneficial for slow moving fires in forested ecosystems.






715

**Figure A2: Tradeoffs between reducing local minima not associated with ignition locations and adjustments made to the global burned area product.** (a) Local minima (ignitions) detected within the daily 500 m global burned area data for 2015 after different number of iterations of the ignition point filter, (b) corresponding fraction of burned area pixels with adjusted burn date, and (c) corresponding number of burned area pixels adjusted divided by the reduction in ignition count. In this study, we used three iterations of the ignition point filter (indicated with the intermittent lines in figures a, b and c), and “0 iterations” refers to the original MCD64A1 col. 6 burned area data.

720




725

**Figure A2A3: Average fire persistence threshold at 0.25° resolution.** The fire persistence threshold determines how long a fire may take to spread from one 500 m grid cell into the next. We used a 4-day fire persistence threshold for 500 m grid cells that burned more than 3 times during the study period (2003 - 2016), and a 6, 8 and 10-day fire persistence period for grid cells that burned 2-3, 1-2, or 1 time, respectively.

Appendix B: Supporting material for the results and discussion


730




**Figure B1: Comparison of daily Global Fire Atlas and US Forest Service data for a selected number of well characterized wildfires in the US.** (a) The accumulated daily fire size (for all fires,  $N=15$ ) illustrates the ability of the Global Fire Atlas to reproduce individual large fire sizes at any specific day over the fire lifetime (each blue dot indicates the size of a specific fire on a specific day). (b) A 3-day running average of the daily growth or “expansion” of each fire ( $\text{km}^2 \text{ day}^{-1}$ ) and (c) the daily expansion on each day of each fire. Figures (d), (e), and (f) are like (a), (b), and (c), but for US Forest Service fire sizes smaller than  $500 \text{ km}^2$  or expansion rates lower than  $250 \text{ km}^2 \text{ day}^{-1}$  and corresponding Global Fire Atlas estimates (see intermittent boxes on top-figures).

735

740



745 **Figure B2: Comparison of Global Fire Atlas ~~and MTBS~~ perimeters and ignition locations to**  
**estimates based on MTBS and VIIRS for frequently-burning grasslands in Kansas, USA. (a) Global**  
**Fire Atlas adjusted burn dates from MCD64A1, (b) per-pixel comparison of adjusted burn dates used**  
**within the Global Fire Atlas (GFA) to the day of the (first) active fire detection from VIIRS, (c) ignition**  
**points as estimated by the Global Fire Atlas, (d) manually interpreted ignition locations (red circles) based**  
**on VIIRS active fire detections on top of MTBS fire perimeters, (e) individual fires as estimated by the**  
**Global Fire Atlas, and (ef) the MTBS burned area and individual fires—and (d) individual fires as**  
**estimated by the Global Fire Atlas.** Here, MCD64A1 data underestimates the total burned area compared  
750 to the visual interpretation of Landsat data within the MTBS project, resulting in fragmentation of  
individual large fires. However, the daily temporal resolution of MODIS imagery allows the Global Fire  
Atlas to distinguish individual fires and ignition points within larger burn scars that cannot be resolved  
from infrequent Landsat observations used to delineate fire perimeters within the MTBS project. Broad  
patterns of ignition locations identified by the Global Fire Atlas are confirmed by manual interpretation of  
755 patterns inferred from VIIRS active fire detections (d).



760 **Figure B2B3:** Global Fire Emissions Database (GFED) regions and dominant GFED fire types used for Tables 1 and 2. (a) GFED regions used in Table 1, and (b) GFED dominant fire type as used in  
 765 Table 2. Abbreviations of the GFED regions shown in (a) are: boreal North America (BONA), temperate North America (TENA), Central America (CEAM), northern hemisphere South America (NHSA), southern hemisphere South America (SHSA), Europe (EURO), Middle East (MIDE), northern hemisphere Africa (NHAF), southern hemisphere Africa (SHAF), boreal Asia (BOAS), Central Asia (CEAS) southeast Asia (SEAS), equatorial Asia (EQAS), and Australia and New Zealand (AUST). Abbreviations of the GFED fire types shown in (b) are: boreal forests (BOAF), Temperate forests (TMPF), Tropical forest deforestation (DEFO), savanna (SAVA) and agriculture (AGRI).

770 **Author contributions.** NA, DCM, and JTR designed the study. NA carried out the data processing and analysis. All authors contributed to the interpretation of the results and writing of the manuscript.

**Competing interests.** The authors declare that they have no conflict of interest.

775 **Acknowledgements.** This work was supported by NASA's Carbon Monitoring System program (grant 80NSSC18K0179) and the Gordon and Betty Moore Foundation (grant GBMF3269). We thank Thomas Mellin of the USFS for granting access to the daily fire perimeter collected over the western USA.



## References

- 780 Abreu, R. C. R., Hoffmann, W. A., Vasconcelos, H. L., Pilon, N. A., Rossatto, D. R. and Durigan, G.: The biodiversity cost of carbon sequestration in tropical savanna, *Sci. Adv.*, 3, e1701284, doi:10.1126/sciadv.1701284, 2017.
- 785 Alencar, A., Asner, G. P., Knapp, D. and Zarin, D.: Temporal variability of forest fires in eastern Amazonia., *Ecol. Appl.*, 21, 2397–2412, doi:10.1890/10-1168.1, 2011.
- Andela, N., Kaiser, J. W., van der Werf, G. R. and Wooster, M. J.: New fire diurnal cycle characterizations to improve fire radiative energy assessments made from MODIS observations, *Atmos. Chem. Phys.*, 15, 8831–8846, doi:10.5194/acp-15-8831-2015, 2015.
- 790 Andela, N., Morton, D. C., Giglio, L., Chen, Y., Van Der Werf, G. R., Kasibhatla, P. S., Defries, R. S., Collatz, G. J., Hantson, S., Kloster, S., Bachelet, D., Forrest, M., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Yue, C. and Randerson, J. T.: A human-driven decline in global burned area, *Science*, 356, 1356–1362, doi:10.1126/science.aal4108, 2017.
- 795 Aragão, L. E. O. C., Anderson, L. O., Fonseca, M. G., Rosan, T. M., Vedovato, L. B., Wagner, F. H., Silva, C. V. J., Silva Junior, C. H. L., Arai, E., Aguiar, A. P., Barlow, J., Berenguer, E., Deeter, M. N., Domingues, L. G., Gatti, L., Gloor, M., Malhi, Y., Marengo, J. A., Miller, J. B., Phillips, O. L. and Saatchi, S.: 21st Century drought-related fires counteract the decline of Amazon deforestation carbon emissions, *Nat. Commun.*, 9, 536, doi:10.1038/s41467-017-02771-y, 2018.
- 800 Archibald, S., Lehmann, C. E. R., Gómez-Dans, J. L. and Bradstock, R. A.: Defining pyromes and global syndromes of fire regimes, *Proc. Natl. Acad. Sci. U. S. A.*, 110, 6442–6447, doi:10.1073/pnas.1211466110, 2013.
- Archibald, S. and Roy, D. P.: Identifying individual fires from satellite-derived burned area data, *IEEE Int. Geosci. Remote Sens. Symp. Proc.*, 9, 160–163, doi:10.1109/IGARSS.2009.5417974, 2009.
- Archibald, S., Staver, A. C. and Levin, S. A.: Evolution of human-driven fire regimes in Africa, *Proc. Natl. Acad. Sci.*, 109, 847–852, doi:10.1073/pnas.1118648109, 2012.
- 805 Balch, J. K., Bradley, B. A., Abatzoglou, J. T., Nagy, R. C., Fusco, E. J. and Mahood, A. L.: Human-started wildfires expand the fire niche across the United States, *Proc. Natl. Acad. Sci. U. S. A.*, 114, 2946–2951, doi:10.1073/pnas.1617394114, 2017.
- 810 Bauters, M., Drake, T. W., Verbeeck, H., Bodé, S., Hervé-Fernández, P., Zito, P., Podgorski, D. C., Boyemba, F., Makelele, I., Cizungu Ntaboba, L., Spencer, R. G. M. and Boeckx, P.: High fire-derived nitrogen deposition on central African forests, *Proc. Natl. Acad. Sci. U. S. A.*, 115, 549–554, doi:10.1073/pnas.1714597115, 2018.
- Benali, A., Russo, A., Sá, A. C. L., Pinto, R. M. S., Price, O., Koutsias, N. and Pereira, J. M. C.: Determining fire dates and locating ignition points with satellite data, *Remote Sens.*, 8, 326, doi:10.3390/rs8040326, 2016.
- 815 Bond, W. J., Woodward, F. I. and Midgley, G. F.: The global distribution of ecosystems in a world without fire, *New Phytol.*, 165, 525–537, doi:10.1111/j.1469-8137.2004.01252.x, 2005.
- Boschetti, L., Roy, D. P. and Justice, C. O.: International Global Burned Area Satellite Product Validation Protocol. In: CEOS-CalVal (Ed.), Part I–production and standardization of validation reference data, Committee Earth Obs. Satell. USA, pp. 1-11, 2009.
- 820 Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D’Antonio, C. M., Defries, R. S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Marston, J. B., Moritz, M. A., Prentice, I. C., Roos, C. I., Scott, A. C., Swetnam, T. W., van der Werf, G. R. and Pyne, S. J.: Fire in the Earth system, *Science*, 324, 481–484,

- doi:10.1126/science.1163886, 2009.
- 825 Brando, P. M., Balch, J. K., Nepstad, D. C., Morton, D. C., Putz, F. E., Coe, M. T., Silvério, D., Macedo, M. N., Davidson, E. A., Nóbrega, C. C., Alencar, A. and Soares-Filho, B. S.: Abrupt increases in Amazonian tree mortality due to drought-fire interactions., *Proc. Natl. Acad. Sci. U. S. A.*, 111, 6347–6352, doi:10.1073/pnas.1305499111, 2014.
- 830 Chen, Y., Morton, D. C., Andela, N., Van Der Werf, G. R., Giglio, L. and Randerson, J. T.: A pan-tropical cascade of fire driven by El Niño/Southern Oscillation, *Nat. Clim. Chang.*, 7, 906–911, doi:10.1038/s41558-017-0014-8, 2017.
- Coen, J. L. and Schroeder, W.: Use of spatially refined satellite remote sensing fire detection data to initialize and evaluate coupled weather-wildfire growth model simulations, *Geophys. Res. Lett.*, 40, 5536–5541, doi:10.1002/2013GL057868, 2013.
- 835 Earl, N. and Simmonds, I.: Spatial and Temporal Variability and Trends in 2001–2016 Global Fire Activity, *J. Geophys. Res. Atmos.*, 123, 2524–2536, doi:10.1002/2017JD027749, 2018.
- Eidenshink, J., Schwind, B., Brewer, K., Zhu, Z.-L., Quayle, B. and Howard, S.: A Project for Monitoring Trends in Burn Severity, *Fire Ecol.*, 3, 3–21, doi:10.4996/fireecology.0301003, 2007.
- 840 Field, R. D., Werf, G. R. Van Der, Fanin, T., Fetzer, E. J., Fuller, R., Jethva, H., Levy, R., van der Werf, G. R., Fanin, T., Fetzer, E. J., Fuller, R., Jethva, H., Levy, R., Livesey, N. J., Luo, M., Torres, O. and Worden, H. M.: Indonesian fire activity and smoke pollution in 2015 show persistent nonlinear sensitivity to El Niño-induced drought, *Proc. Natl. Acad. Sci. U. S. A.*, 113, 9204–9209, doi:10.1073/pnas.1524888113, 2016.
- 845 Frantz, D., Stellmes, M., Röder, A. and Hill, J.: Fire spread from MODIS burned area data: Obtaining fire dynamics information for every single fire, *Int. J. Wildl. Fire*, 25, 1228–1237, doi:10.1071/WF16003, 2016.
- Freeborn, P. H., Wooster, M. J. and Roberts, G.: Addressing the spatiotemporal sampling design of MODIS to provide estimates of the fire radiative energy emitted from Africa, *Remote Sens. Environ.*, 115, 475–489, doi:10.1016/j.rse.2010.09.017, 2011.
- 850 Friedl, M. A., McIver, D. K., Hodges, J. C. F., Zhang, X. Y., Muchoney, D., Strahler, A. H., Woodcock, C. E., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F. and Schaaf, C.: Global land cover mapping from MODIS: algorithms and early results, *Remote Sens. Environ.*, 83, 287–302, doi:10.1016/S0034-4257(02)00078-0, 2002.
- 855 Fusco, E. J., Abatzoglou, J. T., Balch, J. K., Finn, J. T. and Bradley, B. A.: Quantifying the human influence on fire ignition across the western USA, *Ecol. Appl.*, 26, 2388–2399, doi:10.1002/eap.1395, 2016.
- Fusco, E. J., Finn, J. T., Abatzoglou, J. T., Balch, J. K., Dadashi, S. and Bradley, B. A.: Detection rates and biases of fire observations from MODIS and agency reports in the conterminous United States, *Remote Sens. Environ.*, 220, 30–40, doi:10.1016/j.rse.2018.10.028, 2019.
- 860 Giglio, L., Boschetti, L., Roy, D. P., Humber, M. L. and Justice, C. O.: The Collection 6 MODIS burned area mapping algorithm and product, *Remote Sens. Environ.*, 217(July), 72–85, doi:10.1016/j.rse.2018.08.005, 2018.
- 865 Giglio, L., Loboda, T., Roy, D. P., Quayle, B. and Justice, C. O.: An active-fire based burned area mapping algorithm for the MODIS sensor, *Remote Sens. Environ.*, 113, 408–420, doi:10.1016/j.rse.2008.10.006, 2009.
- Giglio, L., Randerson, J. T. and van der Werf, G. R.: Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4), *J. Geophys. Res. Biogeosciences*,

- 118, 317–328, doi:10.1002/jgrg.20042, 2013.
- 870 Di Giuseppe, F., Rémy, S., Pappenberger, F. and Wetterhall, F.: Using the Fire Weather Index (FWI) to improve the estimation of fire emissions from fire radiative power (FRP) observations, *Atmos. Chem. Phys.*, 18, 5359–5370, doi:10.5194/acp-18-5359-2018, 2018.
- 875 Hantson, S., Arneth, A., Harrison, S. P., Kelley, D. I., Prentice, I. C., Rabin, S. S., Archibald, S., Mouillot, F., Arnold, S. R., Artaxo, P., Bachelet, D., Ciais, P., Forrest, M., Friedlingstein, P., Hickler, T., Kaplan, J. O., Kloster, S., Knorr, W., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Meyn, A., Sitch, S., Spessa, A., van der Werf, G. R., Voulgarakis, A. and Yue, C.: The status and challenge of global fire modelling, *Biogeosciences*, 13, 3359–3375, doi:10.5194/bg-2016-17, 2016.
- Hantson, S., Pueyo, S. and Chuvieco, E.: Global fire size distribution is driven by human impact and climate, *Glob. Ecol. Biogeogr.*, 24, 77–86, doi:10.1111/geb.12246, 2015.
- 880 Hantson, S., Scheffer, M., Pueyo, S., Xu, C., Lasslop, G., Van Nes, E. H., Holmgren, M. and Mendelsohn, J.: Rare, Intense, Big fires dominate the global tropics under drier conditions, *Sci. Rep.*, 7, 14374, doi:10.1038/s41598-017-14654-9, 2017.
- Humber, M. L., Boschetti, L., Giglio, L. and Justice, C. O.: Spatial and temporal intercomparison of four global burned area products, *Int. J. Digit. Earth*, 8947, 1–25, doi:10.1080/17538947.2018.1433727, 2018.
- 885 Johnston, F. H., Henderson, S. B., Chen, Y., Randerson, J. T., Marlier, M., Defries, R. S., Kinney, P., Bowman, D. M. J. S. and Brauer, M.: Estimated global mortality attributable to smoke from landscape fires, *Environ. Health Perspect.*, 120, 695–701, doi:10.1289/ehp.1104422, 2012.
- Jolly, W. M., Cochrane, M. A., Freeborn, P. H., Holden, Z. A., Brown, T. J., Williamson, G. J. and Bowman, D. M. J. S.: Climate-induced variations in global wildfire danger from 1979 to 2013, *Nat. Commun.*, 6, 7537, doi:10.1038/ncomms8537, 2015.
- 890 Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razinger, M., Schultz, M. G., Suttie, M. and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9(1), 527–554, doi:10.5194/bg-9-527-2012, 2012.
- 895 Kasischke, E. S. and Turetsky, M. R.: Recent changes in the fire regime across the North American boreal region - Spatial and temporal patterns of burning across Canada and Alaska, *Geophys. Res. Lett.*, 33, L09703, doi:10.1029/2006GL025677, 2006.
- Knorr, W., Arneth, A. and Jiang, L.: Demographic controls of future global fire risk, *Nat. Clim. Chang.*, 6, 781–785, doi:10.1038/nclimate2999, 2016.
- 900 Koplitz, S. N., Mickley, L. J., Marlier, M. E., Buonocore, J. J., Kim, P. S., Liu, T., Sulprizio, M. P., DeFries, R. S., Jacob, D. J., Schwartz, J., Pongsiri, M. and Myers, S. S.: Public health impacts of the severe haze in Equatorial Asia in September-October 2015: Demonstration of a new framework for informing fire management strategies to reduce downwind smoke exposure, *Environ. Res. Lett.*, 11, 094023, doi:10.1088/1748-9326/11/9/094023, 2016.
- 905 Laurent, P., Mouillot, F., Yue, C., Ciais, P., Moreno, M. V. and Nogueira, J. M. P.: FRY, a global database of fire patch functional traits derived from space-borne burned area products, *Sci. Data*, 5, 180132, doi:10.1038/sdata.2018.132, 2018.
- Van Leeuwen, T. T. and Van Der Werf, G. R.: Spatial and temporal variability in the ratio of trace gases emitted from biomass burning, *Atmos. Chem. Phys.*, 11, 3611–3629, doi:10.5194/acp-11-3611-2011, 2011.
- 910 Lelieveld, J., Evans, J. S., Fnais, M., Giannadaki, D. and Pozzer, A.: The contribution of outdoor air pollution sources to premature mortality on a global scale, *Nature*, 525, 367–371,



- doi:10.1038/nature15371, 2015.
- Loboda, T. V. and Csiszar, I. A.: Reconstruction of fire spread within wildland fire events in Northern Eurasia from the MODIS active fire product, *Glob. Planet. Change*, 56, 258–273, doi:10.1016/j.gloplacha.2006.07.015, 2007.
- 915 Moritz, M. A., Batllori, E., Bradstock, R. A., Gill, A. M., Handmer, J., Hessburg, P. F., Leonard, J., McCaffrey, S., Odion, D. C., Schoennagel, T. and Syphard, A. D.: Learning to coexist with wildfire, *Nature*, 515, 58–66, doi:10.1038/nature13946, 2014.
- Morton, D. C., Collatz, G. J., Wang, D., Randerson, J. T., Giglio, L. and Chen, Y.: Satellite-based assessment of climate controls on US burned area, *Biogeosciences*, 10, 247–260, doi:10.5194/bg-10-247-2013, 2013a.
- 920 Morton, D. C., Page, Y. Le, Defries, R., Collatz, G. J. and Hurtt, G. C.: Understorey fire frequency and the fate of burned forests in southern Amazonia, *Phil. Trans. R. Soc. B*, 368, 20120163, doi:10.1098/rstb.2012.0163, 2013b.
- 925 Nogueira, J. M. P., Ruffault, J., Chuvieco, E. and Mouillot, F.: Can we go beyond burned area in the assessment of global remote sensing products with fire patch metrics?, *Remote Sens.*, 9, 7, doi:10.3390/rs9010007, 2017.
- Oom, D., Silva, P. C., Bistinas, I. and Pereira, J. M. C.: Highlighting biome-specific sensitivity of fire size distributions to time-gap parameter using a new algorithm for fire event individuation, *Remote Sens.*, 8, 663, doi:10.3390/rs8080663, 2016.
- 930 Padilla, M., Stehman, S. V., Ramo, R., Corti, D., Hantson, S., Oliva, P., Alonso-Canas, I., Bradley, A. V., Tansey, K., Mota, B., Pereira, J. M. and Chuvieco, E.: Comparing the accuracies of remote sensing global burned area products using stratified random sampling and estimation, *Remote Sens. Environ.*, 160, 114–121, doi:10.1016/j.rse.2015.01.005, 2015.
- 935 Le Page, Y., Morton, D., Bond-Lamberty, B., Pereira, J. M. C. and Hurtt, G.: HESFIRE: A global fire model to explore the role of anthropogenic and weather drivers, *Biogeosciences*, 12, 887–903, doi:10.5194/bg-12-887-2015, 2015.
- Parker, R. J., Boesch, H., Wooster, M. J., Moore, D. P., Webb, A. J., Gaveau, D. and Murdiyarso, D.: Atmospheric CH<sub>4</sub> and CO<sub>2</sub> enhancements and biomass burning emission ratios derived from satellite observations of the 2015 Indonesian fire plumes, *Atmos. Chem. Phys.*, 16, 10111–10131, doi:10.5194/acp-16-10111-2016, 2016.
- 940 Pellegrini, A. F. A., Ahlström, A., Hobbie, S. E., Reich, P. B., Nieradzik, L. P., Staver, A. C., Scharenbroch, B. C., Jumpponen, A., Anderegg, W. R. L., Randerson, J. T. and Jackson, R. B.: Fire frequency drives decadal changes in soil carbon and nitrogen and ecosystem productivity, *Nature*, 53, 194–198, doi:10.1038/nature24668, 2018.
- 945 Pfeiffer, M., Spessa, A. and Kaplan, J. O.: A model for global biomass burning in preindustrial time: LPJ-LMfire (v1.0), *Geosci. Model Dev.*, 6, 643–685, doi:10.5194/gmd-6-643-2013, 2013.
- Rabin, S. S., Melton, J. R., Lasslop, G., Bachelet, D., Forrest, M., Hantson, S., Kaplan, J. O., Li, F., Mangeon, S., Ward, D. S., Yue, C., Arora, V. K., Hickler, T., Kloster, S., Knorr, W., Nieradzik, L., Spessa, A., Folberth, G. A., Sheehan, T., Voulgarakis, A., Kelley, D. I., Colin Prentice, I., Sitch, S., Harrison, S. and Arneeth, A.: The Fire Modeling Intercomparison Project (FireMIP), phase 1: Experimental and analytical protocols with detailed model descriptions, *Geosci. Model Dev.*, 10, 1175–1197, doi:10.5194/gmd-10-1175-2017, 2017.
- 950 Randerson, J. T., Chen, Y., van der Werf, G. R., Rogers, B. M. and Morton, D. C.: Global burned area and biomass burning emissions from small fires, *J. Geophys. Res.*, 117, G04012, doi:10.1029/2012JG002128, 2012.
- 955

- 960 Randerson, J. T., Liu, H., Flanner, M. G., Chambers, S. D., Jin, Y., Hess, P. G., Pfister, G., Mack, M. C., Treseder, K. K., Welp, L. R., Chapin, F. S., Harden, J. W., Goulden, M. L., Lyons, E., Neff, J. C., Schuur, E. A. G. and Zender, C. S.: The impact of boreal forest fire on climate warming, *Science*, 314, 1130–1132, doi:10.1126/science.1132075, 2006.
- Reisen, F., Meyer, C. P., Weston, C. J. and Volkova, L.: Ground-based Field Measurements of PM<sub>2.5</sub> Emission Factors from Flaming and Smouldering Combustion in Eucalypt Forests, *J. Geophys. Res. Atmos.*, doi:10.1029/2018JD028488, 2018.
- 965 Roteta, E., Bastarrika, A., Padilla, M., Storm, T. and Chuvieco, E.: Development of a Sentinel-2 burned area algorithm: Generation of a small fire database for sub-Saharan Africa, *Remote Sens. Environ.*, 222(September 2017), 1–17, doi:10.1016/j.rse.2018.12.011, 2019.
- Sá, A. C. L., Benali, A., Fernandes, P. M., Pinto, R. M. S., Trigo, R. M., Salis, M., Russo, A., Jerez, S., Soares, P. M. M., Schroeder, W. and Pereira, J. M. C.: Evaluating fire growth simulations using satellite active fire data, *Remote Sens. Environ.*, 190, 302–317, doi:10.1016/j.rse.2016.12.023, 2017.
- 970 Scholes, R. J. and Archer, S. R.: Tree-grass interactions in savannas, *Annu. Rev. Ecol. Syst.*, 28, 517–544, doi:10.1146/annurev.ecolsys.28.1.517, 1997.
- Schroeder, W., Oliva, P., Giglio, L. and Csiszar, I. A.: The New VIIRS 375m active fire detection data product: Algorithm description and initial assessment, *Remote Sens. Environ.*, 143, 85–96, doi:10.1016/j.rse.2013.12.008, 2014.
- 975 Schroeder, W., Prins, E., Giglio, L., Csiszar, I., Schmidt, C., Morisette, J. and Morton, D.: Validation of GOES and MODIS active fire detection products using ASTER and ETM+ data, *Remote Sens. Environ.*, 112, 2711–2726, doi:10.1016/j.rse.2008.01.005, 2008.
- Sparks, A. M., Boschetti, L., Smith, A. M. S., Tinkham, W. T., Lannom, K. O. and Newingham, B. A.: An accuracy assessment of the MTBS burned area product for shrub-steppe fires in the northern Great Basin, United States, *Int. J. Wildl. Fire*, 24, 70–78, doi:10.1071/WF14131, 2015.
- 980 Staver, A. C., Archibald, S. and Levin, S. A.: The Global Extent and Determinants of Savanna and Forest as Alternative Biome States, *Science*, 334, 230–232, doi:10.1126/science.1210465, 2011.
- Taylor, A. H., Trouet, V., Skinner, C. N. and Stephens, S.: Socioecological transitions trigger fire regime shifts and modulate fire – climate interactions in the Sierra, *Proc. Natl. Acad. Sci. U. S. A.*, 113, 13684–13689, doi:10.1073/pnas.1609775113, 2016.
- 985 Thonicke, K., Spessa, A., Prentice, I. C., Harrison, S. P., Dong, L. and Carmona-Moreno, C.: The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas emissions: results from a process-based model, *Biogeosciences*, 7, 1991–2011, doi:10.5194/bg-7-1991-2010, 2010.
- 990 Veraverbeke, S., Sedano, F., Hook, S. J., Randerson, J. T., Jin, Y. and Rogers, B. M.: Mapping the daily progression of large wildland fires using MODIS active fire data, *Int. J. Wildl. Fire*, 23, 655–667, doi:10.1071/WF13015, 2014.
- Ward, D. S., Kloster, S., Mahowald, N. M., Rogers, B. M., Randerson, J. T. and Hess, P. G.: The changing radiative forcing of fires: global model estimates for past, present and future, *Atmos. Chem. Phys.*, 12, 10857–10886, doi:10.5194/acp-12-10857-2012, 2012.
- 995 Ward, D. S., Shevliakova, E., Malyshev, S. and Rabin, S.: Trends and Variability of Global Fire Emissions Due To Historical Anthropogenic Activities, *Global Biogeochem. Cycles*, 32, 122–142, doi:10.1002/2017GB005787, 2018.
- 1000 van der Werf, G. R., Dempewolf, J., Trigg, S. N., Randerson, J. T., Kasibhatla, P. S., Giglio, L., Murdiyarso, D., Peters, W., Morton, D. C., Collatz, G. J., Dolman, A. J. and DeFries, R. S.: Climate regulation of fire emissions and deforestation in equatorial Asia., *Proc. Natl. Acad. Sci. U. S. A.*, 105,

20350–20355, doi:10.1073/pnas.0803375105, 2008.

Westerling, A. L., Hidalgo, H. G., Cayan, D. R. and Swetnam, T. W.: Warming and earlier spring increase western US forest wildfire activity, *Science*, 313, 940–943, doi:10.1126/science.1128834, 2006.

1005 Yang, J., Tian, H., Tao, B., Ren, W., Pan, S., Liu, Y. and Wang, Y.: A growing importance of large fires in conterminous United States during 1984-2012, *J. Geophys. Res. Biogeosciences*, 120, 2625–2640, doi:10.1002/2015JG002965, 2015.

Young, A. M., Higuera, P. E., Duffy, P. A. and Hu, F. S.: Climatic thresholds shape northern high-latitude fire regimes and imply vulnerability to future climate change, *Ecography*, 40, 606–617, doi:10.1111/ecog.02205, 2017.

1010