

Dear Editor,

Below you will find our point-by-point responses to the reviewer comments along with the changes we made to the manuscript. At the end you will find the revised manuscript with all changes shown in red.

Thank you for processing our manuscript.

Martin Stengel and co-authors.

Anonymous Referee #1

General comments:

This article describes the third version of the Cloud Climate Change Initiative AVHRR-PM dataset; a 35 year climatology based on measurements in 5 spectral bands from the Advanced Very High Resolution Radiometer (AVHRR) instruments on board several polar orbiting satellites. This dataset includes both cloud properties retrieved from the AVHRR measurements and surface and top of atmosphere irradiances calculated using these retrieved cloud properties. The article describes changes since the previous version of the dataset and presents some evaluation of the dataset and comparisons with the previous version.

This article is generally very well written. The description of the cloud retrieval algorithm is quite brief, but this is appropriate as the previous version of the algorithm is described in detail in a previous publication (Stengel et al. 2017). The description of the radiative transfer calculations is also quite brief and in my opinion more detail is required here (see comments below for suggestions). The dataset was accessed through the given identifier and appeared to be complete and consistent with the description in the article.

Recommendation:

Accept pending minor revisions.

Minor Comments:

Referee comment:

1. Why does this article focus specifically on AVHRR-pm, as opposed to describing datasets for multiple instruments as in Stengel et al (2017)? At the very least, it seems odd not to include the AVHRR-am dataset in the scope of this article. On similar lines, would it be possible to produce a product combining AVHRR-am and AVHRR-pm measurements? Presumably the additional sampling of the diurnal cycle would lead to smaller errors in the radiation diurnal cycle corrections.

Author's response:

The Stengel et al. (2017) paper aimed at introducing all available version 2 datasets generated in the framework of the Cloud_cci project. Only a subset of those (AVHRR-AM, AVHRR-PM, ATSR2-AATSR) were reprocessed building on new developments leading to corresponding version 3 datasets. ATSR2-AATSR version 3 data will be introduced in a separate paper, which is soon to be submitted. For AVHRR we decided to put the focus more or less entirely on AVHRR-PM as this dataset is longer, more stable and of higher quality than AVHRR-AM. On the other hand, for the period covered by the AM satellites NOAA-17 and METOP-A (NOAA-12 and NOAA-15 are very difficult to handle due to their twilight orbit), the AVHRR-AM datasets is of similar quality as AVHRR-PM and does indeed provided the possibility for combining AVHRR-AM and AVHRR-PM to reduced sampling problems, although only for the years 2002 and beyond. Another difficulty is the availability of the 1.6mic channel instead of the 3.7mic channel as available on nearly all PM satellites. In the data availability section, the existence of the AVHRR-AMv3 is reflected. We will add that there is a potential to combine with AVHRR-PMv3, but also mentioning the difficulties for NOAA-12 and NOAA-15.

Author's changes to the manuscript:

We added the following sentence to the data availability section: "The AVHRR-AMv3 dataset provides the feasibility to be combined with AVHRR-PMv3 to increase sampling frequency. However, for the period of NOAA-12 and NOAA-15 the AVHRR-AMv3 dataset is of reduced quality due to the difficult twilight orbits of NOAA-12 and NOAA-15."

Referee comment:

2. It's very difficult to see any differences in most of Figs. 1,2,5,6,7,8. I would consider including difference plots, either instead of the v2/CERES images, or as an additional row/column.

Author's response:

We have included differences plots in Figures 1,2,5,6,7,8.

Author's changes to the manuscript:

We updated Figures 1,2,5,6,7,8 , modified their captions and added/modified a few sentences in the text linking to the difference plots.

Referee comment:

3. Is there any attempt to account for changes in the surface albedo with the angle of incident light in the SW radiative transfer calculations (e.g. Wang et al 2007)? Perhaps this could explain some of the differences between the CERES and AVHRR-pm surface SW upwelling irradiances?

Author's response:

Yes, for ocean surfaces we have built in an empirical method to adjust the surface albedo as a quadratic function of the angle of incident light. For land surfaces no adjustment is made. We will include a statement on this in the text.

Author's changes to the manuscript:

On page 13 line 2 we will change the sentence to "The diurnal cycle of SZA is then used to rescale the incoming and reflected solar radiation, adjust the surface albedo (using an empirical quadratic function of SZA) and the atmospheric path length for a given set of time stamps throughout the local day."

Referee comment:

4. With regards to the LW diurnal cycle correction factors, are separate factors derived for clear and cloudy scenes?

Author's response:

No, this is not the case, but probably something to consider for the future. Thanks. In this context we noticed that the LW diurnal cycle correction is only applied for land surfaces, which is not reflected in the manuscript yet. We will add this now.

Author's changes to the manuscript:

On page 13 line 7 we will modify the sentence to "For longwave radiation, a diurnal cycle correction is applied over land based on a cosine fit to an observed mean...."

Referee comment:

5. If I understand the radiative transfer model correctly, it requires the cloud to be split into layers. If this is the case, how do you determine how many layers to include cloud in (i.e. where is the cloud base?). I would expect this to have a reasonably large impact on the calculated surface LW downwelling irradiance.

Author's response:

We assume the radiative transfer in BUGSrad is meant which is employed to derive the broadband fluxes. Here we assume only one cloud layer with its top being place at the derived cloud top height. Using derived optical thickness and effective radius the geometrical thickness, thus the cloud base height is estimated. This is actually described in more detail in an ATBD, which we will include a reference to.

Author's changes to the manuscript:

On page 12 line 22 we will add the sentence "The reader is referred to ATBD (2019) for more details on the calculation of the broadband fluxes." Along with including the following reference :

ATBD – Algorithm Theoretical Baseline Document (ATBD) of CC4CL Broadband Radiative Flux Retrieval - ESA Cloud_cci, 2019, Issue 1, Rev. 1; 14/10/2019, available from <http://www.esa-cloud-cci.org/?q=documentation>, 2019.

Referee comment:

6. Page 15, Line 11 -13. I don't follow the argument that "the larger standard deviations...is primarily related to variances in surface albedo and cloud cover which tend to have significant annual cycles". Relating the larger s.d. to the surface albedo variance makes sense, but I don't understand why the cloud cover variance will lead to a larger s.d. as it also affects the downwelling SW irradiance.

Author's response:

Right. We will remove cloud cover.

Author's changes to the manuscript:

The revised version of that sentence will read "The larger standard deviations retrieved from the solar reflected radiation is primarily related to variances in surface albedo which tend to have significant annual cycles"

Referee comment:

7. For TOA radiation, clear-sky differences between CERES and AVHRR-pm are attributed to sampling differences. Presumably this is relatively easy to test by calculating a CERES-like value from the AVHRR-pm product?

Author's response:

Thank you for this suggestion. We performed a little experiment (covering 3 months) in which we emulated the CERES-like clear-sky sampling. And yes, when doing so the global mean TOA LW flux was increased by approx. 3W/m^2 , thus the deviation to CERES was reduced. This emphasises that at least parts of the deviation can be explained by sampling. We will include the results of this experiment in the text.

Author's changes to the manuscript:

The following sentence was added: "This could be confirmed by a 3-months covering test run in which Cloud_cci LWF^{up_toa} was only averaged over clear-sky cases, which led to an increase by about $3\sim\text{W m}^{-2}$ for the global mean value."

Referee comment:

8. To further demonstrate the usefulness of the radiation products, it would be good to see some further comparisons with other datasets, such as the ERA-Interim reanalysis, or the GEWEX radiation budget data. Perhaps you could add a couple of extra lines in table 8 to show mean values for other products?

Author's response:

Good suggestion. We will add value for ERA-Interim to table 8, and addition to tables 9 and 10. We prefer not to add the GEWEX SRB dataset as it does not fully cover the period 2003-2016 chosen for corresponding comparisons.

Author's changes to the manuscript:

We will add ERA-Interim values to Tables 8, 9 and 10.

Referee comment:

9. I really appreciate the effort undertaken to provide useful and accurate uncertainty estimates for the cloud variables. It would be very helpful to have some estimate of uncertainty in the computed radiation variables too. This could be based on further radiative transfer calculations using different cloud inputs to represent the uncertainty in the input cloud profiles, though this may be time consuming. Alternatively, a simple quality variable to indicate when the radiation calculation is uncertain due to larger uncertainty in the input cloud profiles could potentially be quite helpful.

Author's response:

Thank you for this suggestion. We have in indeed planned to provide uncertainty estimates for the radiation variables too. For the presented dataset version this was however not feasible due to time constraints, but we certainly have this on the to-do list for next versions. For the time being the radiation validation results presented do provide some guidance wrt. to certainty/uncertainty of the radiation products, although not on pixels/grid-cell level. As the determination of the radiation product heavily depend on the derived cloud properties and their uncertainty it is indeed wise to inform the

users of the data that the provided cloud property uncertainties give hints on the certainty/uncertainty in the radiation product already in the current dataset version. We will include a comment on this in the text.

Author's changes to the manuscript:

We added the following sentence at the end of Section 3.1 “In contrast to the cloud properties, the radiative fluxes in the presented dataset version are not accompanied by uncertainty estimates on pixel level. While the validation results presented below provide a general guidance to the quality of the radiative fluxes, user of the data are also encouraged to inspect the pixel-level uncertainties of the cloud properties as these are dominant input to the calculation of the fluxes.”

Referee comment:

10. Looking at the daily data, there appear to be some artifacts in the retrieved cloud water path at the edges of the swaths for the descending overpasses (e.g. for the 1 June 2016 data). These do seem to correspond with very large uncertainty estimates. In such cases, where the uncertainty is much larger than the retrieved value I wonder whether it would be better to replace the retrieved value with a missing data value? In particular, I have concerns about these retrieved values undergoing further processing (e.g. passed to radiation calculations, or used in monthly mean/histogram products) and the information about the large uncertainty associated with the particular retrieval being lost.

Author's response:

Thank you for this observation and feedback. The descending cloud water path is based on the night-time retrievals of CER and COT, which are experimental products as listed in the manuscript. To emphasise this we will add a corresponding comment below Table 1. For monthly aggregations these night-time products are not considered. A quick inspection of the LW fluxes confirmed that this issue does not seemed to have a significant impact on the LW fluxes. SW fluxes are zero anyway during night time.

Author's changes to the manuscript:

Added a comment below Table 1.

Anonymous Referee #2

General comments:

Summary of paper: The authors describe a data set of global cloud properties based on AVHRR observations, available since 1982. The data set is an update of version 2, with the main changes the use of artificial neural networks for cloud mask and cloud phase detection, and additional cloud radiative properties. Both versions 2 and 3 are evaluated against the best available retrievals from other satellite and ground-based products. Standard verification metrics indicate overall improvement in most cloud properties, with some deterioration in ice cloud top height. The cloud radiative properties compare well against CERES observations.

Review: This paper is generally very well written. It is mostly complete and useful information for anyone wishing to use this data set. Comments mostly concern some further clarification sought and perhaps slightly different presentation of the differences between version 2 and 3. The DOI links to a suitably presented web page describing the data. Overall, the recommendation is to accept this paper with minor corrections.

Minor comment:

Referee comment:

Statistical significance. It is not immediately clear whether any of the differences in skill between v2 and v3 reported in the tables are statistically significant, although the large sample suggests these are. However, it should be possible and it would help the reader if the maps in figures 1, 2, 5, 6, 7, and 8 could include (i) difference/bias maps and (ii) stippling/hashing/shading for statistically significant differences. Most of these maps are visually similar and might hide key differences due to the colour scale used. A different way of presenting the various data sets, including additional maps of bias and statistically significant differences, would help inform the reader how the new data set compares against existing data sets.

Author's response:

Thank you for this suggestion. We updated Figures 1,2,5,6,7 and 8 to include difference maps as suggested. Unfortunately, we have not always all information available to calculate and provide the statistical significance of the differences, which are for this reason not included.

Author's changes to the manuscript:

We updated Figures 1,2,5,6,7,8 , modified their captions and added/modified a few sentences in the text linking to the difference plots.

Line-by-line comments:

Referee comment:

p2. l28-30. This sentence is difficult to read, especially the first part.

Author's response:

Thanks. We will rephrase that sentence.

Author's changes to the manuscript:

The rephrased sentence will read "For the MODIS cloud record however there is the potential to be combined with high quality TOA radiation measurements made by the Clouds and the Earth's Radiant Energy System (CERES) sensors mounted on board the same platforms (Terra and Aqua)."

Referee comment:

p2. l33. "limitations". Limitations to do what?

Author's response:

We mean the limitation to resolve cloud and their radiative effect at smaller scales than the CERES footprint size. We will rephrase that sentence.

Author's changes to the manuscript:

The rephrased sentence will read "Limitations to resolve small scale clouds and their radiative effect might arise from the coarse spatial resolution of CERES (footprint size of approximately 30 km) and from the fact that the clear-sky fluxes are exclusively based on clear-sky pixels (and interpolation of clear-sky fluxes for gap filling on monthly scales), for which the spatio-temporal sampling is reduced and the meteorological conditions are likely to be biased."

Referee comment:

p3. l3-7. Please provide references for the WCRP GEWEX data and the ISCCP DX data.

Author's response:

Thank you. We will include the reference Stackhouse (2011) and Rossow and Schiffer (1990).

Author's changes to the manuscript:

Included reference Stackhouse (2011) and Rossow and Schiffer (1990).

Referee comment:

p3. l10-14. "based on the rationale above". It is clear why these data are required, compared to the MODIS/CERES and GEWEX data sets. However, what are the other data sets based on AVHRR lacking (PATMOS-x, CLARA-A2, Cloud-cci) that this paper will address with Cloud cci 3? A sentence on p2, line 19-21 would help clarify the shortcomings of those existing data sets.

Author's response:

Thanks for this suggestion. We will add the suggested sentence; however we find that sentence better placed on p3 l 13, as we need the provision of a full suite of cloud and radiation properties to make the point.

Author's changes to the manuscript:

We will add the following sentence near line 13 on page 3: "The availability of the full suites of cloud and radiative flux properties will also make these data superior to the already existing AVHRR-based datasets mentioned above."

Referee comment:

p3. l20-31. This paragraph pre-empts the findings ("superior") and methodology. The relevant information is better placed in section 2.1.

Author's response:

Thank you for this suggestion. We kind of see the point. However, algorithm developments (content of section 2.1) are only one reason for this finding. The longer time period covered and in particular the inclusion of the radiative broadband fluxes are strong contributors to this finding/conclusion as well. As these characteristics are motivated in the introduction we would like to leave that paragraph where it is right now.

Author's changes to the manuscript:

None.

Referee comment:

p4. l2. Please add that table 1 contains all abbreviations used throughout the text. CER had not been introduced in the main text prior to p6 and it took a moment to figure out its meaning.

Author's response:

Thank you. We'll do that.

Author's changes to the manuscript:

We will modify that sentence: "The set of cloud properties included in v3 is identical to v2 and is outlined in the upper part of Table 1, which also gives all cloud property abbreviations used throughout the paper."

Referee comment:

p4. l16. "much larger set". How do the two sets of training data compare? Did both v2 and v3 use CALIOP, but v2 just used fewer overpasses?

Author's response:

Yes. Both used CALIOP and for v2 we used much fewer overpasses compared to v3 (about a factor of 10). We will clarify in the text.

Author's changes to the manuscript:

The corresponding sentence will be changed to "The ANN for cloud detection (ANNmask) has been retrained using a much larger set of training data (approx. 10 times more collocation data used for v3 than for v2), which is composed of...."

Referee comment:

p6. l16. A "lower" CTP mean is not explained by more very low-level clouds, which have higher CTP. It appears that over the West Pacific and Maritime Continent, mean CTP has generally increased, which could be due to detection of more low-level clouds. Please re-consider this statement.

Author's response:

Thank you. This is indeed a type. A "higher" CTP it is. We will revise that sentence.

Author's changes to the manuscript:

That sentence will be changed to "Mean CTP is higher in v3 than in v2 in the Tropics...."

Referee comment:

p7. l1. Regarding the validation, did the authors consider performing the validation separately for daytime and nighttime observations? The algorithms use different channels and the authors consider nighttime COT and CER "experimental". It would be useful to understand the algorithm performance for different times of the day.

Author's response:

It is indeed useful. For this manuscript however we would prefer presenting just the general figures as the discussion would get to extended otherwise. Important to consider in this respect that there is a Project Validation Report soon being issued in which we plan to stratify the validation results wrt. illumination conditions. We will add this link to the text.

Author's changes to the manuscript:

At the end of Section 2.3 we will add the following sentence "An even broader assessment of the quality of the presented dataset can be found in PVIR (2019), in which the results are also stratified by illumination conditions among others."

Along with adding the reference:

PVIR – Product Validation and Intercomparison Report (PVIR) ESA Cloud_cci, 2019, Issue 6, Rev. 0; DD/MM/2019, available from http://www.esa-cloud-cci.org/?q=documentation_v3, 2019.

Referee comment:

p7. I3. Please, briefly explain how the collocation is carried out. In particular, what is the impact of the temporal mismatch between CALIOP and AVHRR? And what is the impact of the mismatch in footprint?

Author's response:

The collocations are done identically to Stengel et al. (2017). Most important facts are that all collocations are based on searching for the nearest neighbour in the Cloud_cci Level-3U data to each CALIOP observation. Given that a L3U grid box is usually of smaller size than 5 km, 5 km is the maximal allowed spatial mismatch. Most collocations have much lower spatial mismatches. Due to the similar orbital characteristics of NOAA-18 and NOAA-19 compared with CALIPSO, a very large set of collocations can be retrieved, which in turn also allows for very strict criteria and still infer a sound collocation database. In this context, the temporal mismatch criteria was set to a time window of ± 3 min. While the systematic deviations to CALIOP do not significantly depend on the match-up criteria, the random deviations do. We will summarize all this information in the text.

Author's changes to the manuscript:

We will add the following sentences: The collocations between CALIOP and the AVHRR-PM data were done as reported in Stengel et al. (2017) with the most important fact being that only those collocations were included for which the spatial and temporal mismatch was below 5 km and 3 minutes, respectively. These criteria were chosen as compromise between using best spatial and temporal matches, and allowing for a compositions of a sound data basis to be used in the validation. Important to note that the random deviations of AVHRR-PM to CALIOP depend on the defined criteria, while the systematic do most likely not.

Referee comment:

p7. I22-23. Why would improved identification of liquid clouds lead to reduced POD for ice clouds? This suggests that some ice clouds are now erroneously identified as liquid. Does that mean there are more "false alarms" in terms of liquid cloud detection?

Author's response:

That sentence is indeed misleading. For the v3 algorithm the probability of detection (POD) of liquid clouds is improved while the POD for ice clouds is slightly reduced, still leading to an overall increased (improved) HSS values.

Author's changes to the manuscript:

We will modify that sentence to: "Comparing the HSS score as overall measure for the correct cloud phase detection, v3 performs better than v2. The POD of liquid clouds is significantly improved in v3, while a small degradation in POD of ice clouds is found in v3 compared to v2."

Referee comment:

p8. I5-15. It would be helpful to consider the results from Tables 4, 5, and 6 through a visual comparison, as done in Figure 4. A scatter plot (or 2D histogram) of CTH, LWP, and IWP comparing the data set with the "truth" could help identify where biases are most likely to occur. For instance, the CTH bias of ice cloud could be mostly due to the highest clouds, even at high COT, as these might have a region of low extinction coefficient near cloud top, that would lead to higher CTH in CALIOP. A scatter plot could show this clearly. Similarly, LWP and IWP are highly skewed variables and the metrics presented could be affected by a few outliers. A scatter plot or 2D histogram (perhaps shown on a logarithmic scale) could indicate whether LWP and IWP estimates are typically good, or whether there is a consistent bias across cloud types of all LWP and IWP values.

Author's response:

Thank you for this suggestion. We have included 2d histograms for CTH, LWP and IWP.

Author's changes to the manuscript:

We have included 2d histograms for CTH, LWP and IWP.

Referee comment:

p9. p10. p11. Please rename standard deviation to "root mean squared error", which is presumably what is reported.

Author's response:

We actually mean the standard deviation of the error (which is basically identical to the bias-corrected root mean squared error). We will add to the captions of tables 4,5 and 6 that we mean the standard deviation of the error and the mean error.

Author's changes to the manuscript:

We will modify the captions of tables 4,5 and 6 to "...standard deviation of the error (Std), the mean error (bias)"

Cloud_cci AVHRR-PM dataset version 3: 35 year climatology of global cloud and radiation properties

Martin Stengel¹, Stefan Stapelberg^{1,2}, Oliver Sus^{1,2}, Stephan Finkensieper¹, Benjamin Würzler¹, Daniel Philipp^{3,1}, Rainer Hollmann¹, Caroline Poulsen⁴, Matthew Christensen^{4,5}, and Gregory McGarragh⁵

¹Deutscher Wetterdienst, Frankfurter Str. 135, 63067, Offenbach, Germany

²European Organisation for the Exploitation of Meteorological Satellites, Darmstadt, Germany

³Institute for Atmospheric and Environmental Sciences, Goethe University, Frankfurt am Main, Germany

⁴Science technology Facility Council, Rutherford Appleton Laboratory, Didcot, Oxfordshire, UK

⁵Department of Physics, University of Oxford, Clarendon Laboratory, Parks Road, Oxford OX1 3PU, UK

Correspondence to: Martin Stengel (martin.stengel@dwd.de)

Abstract. We present version 3 of the Cloud_cci AVHRR-PM dataset which contains a comprehensive set of cloud and radiative flux properties on a global scale covering the period of 1982 to 2016. The properties were retrieved from Advanced Very High Resolution Radiometer (AVHRR) measurements recorded by the afternoon (post meridiem, PM) satellites of the National Oceanic and Atmospheric Administration (NOAA) Polar Operational Environmental Satellites (POES) missions. The cloud properties in version 3 are of improved quality compared with the precursor dataset version 2, providing better global quality scores for cloud detection, cloud phase and ice water path based on validation results against A-Train sensors. Furthermore, the parameter set was extended by a suite of broadband radiative flux properties. They were calculated by combining the retrieved cloud properties with thermodynamic profiles from reanalysis and surface properties. The flux properties comprise upwelling and downwelling, shortwave and longwave broadband fluxes at the surface (bottom-of-atmosphere - BOA) and top-of-atmosphere (TOA). All fluxes were determined at AVHRR pixel level for all-sky and clear-sky conditions, which will particularly facilitate the assessment of the cloud radiative effect at BOA and TOA in future studies. Validation of the BOA downwelling fluxes against the Baseline Surface Radiation Network (BSRN) show a very good agreement. This is supported by comparisons of multi-annual mean maps with NASA's Clouds and the Earth's Radiant Energy System (CERES) products for all fluxes at BOA and TOA. The Cloud_cci AVHRR-PM version 3 dataset allows for a large variety of climate applications that build on cloud properties, radiative flux properties and/or the link between them.

For the presented Cloud_cci AVHRR-PMv3 dataset a Digital Object Identifier has been issued:

https://doi.org/10.5676/DWD/ESA_Cloud_cci/AVHRR-PM/V003 (Stengel et al., 2019).

Copyright statement. TEXT

1 Introduction

Clouds play a critical role for the Earth's weather and climate through their contribution to the Earth's water cycle and their impact on the Earth's energy budget. Clouds impact the energy budget through their interaction with radiation, i.e. clouds usually reflect more solar radiation back to space than the underlying surface and absorb and re-emit infrared (IR) radiation, leading to less IR radiation leaving the system than without clouds. Thus clouds significantly alter important components of the Earth's radiation budget: the shortwave and longwave broadband fluxes at the top-of-atmosphere (TOA) and at the surface (bottom-of-atmosphere - BOA hereafter). Analysing cloud coverage and properties, and quantifying the impact they have on the radiation budget is of crucial importance for understanding the Earth's climate and the potential feedback mechanisms in a changing climate.

Since the beginning of the meteorological satellite era at the end of the 1970's attempts have been made to construct global cloud climatologies (e.g. Schiffer and Rossow, 1983) that are of sufficient quality to enable climate studies. Until recently the measurement records of meteorological satellite sensors have grown now cover more than 40 years. Even though many difficulties exist when attempting to construct homogeneous and stable climate datasets, those multi-decadal satellite measurements provide the single most important source of measurements with global coverage. Some international efforts exist to regularly improve and extend long-term satellite-based climatologies that contain a comprehensive suite of cloud properties: The Pathfinder Atmospheres - Extended (PATMOS-x, Heidinger et al., 2014), The International Satellite Cloud Climatology Project (ISCCP, Young et al., 2018), The EUMETSAT Climate Monitoring Satellite Application Facility (CM SAF) cloud and radiation data record (CLARA-A2, Karlsson et al., 2016) and the European Space Agency funded Climate Change Initiative ECV Cloud project (Cloud_cci, Stengel et al., 2017). All of these climatologies make use of measurements of the Advanced Very High Resolution Radiometer (AVHRR), which is a passive imaging sensor with 5-6 spectral bands in the visible, near-infrared and thermal infrared. It is flown on the National Oceanic and Atmospheric Administration (NOAA) Polar Operational Environmental Satellites (POES) and on the EUMETSAT Meteorological operational satellite (Metop) series. There are newer passive sensors in space that also allow for constructing cloud datasets. These are part of research satellite missions by ESA (e.g. the (Advanced) Along-Track Scanning Radiometers on-board the European Remote Sensing Satellite (ERS-1/2) and the Environmental Satellite (Envisat)) and by the National Aeronautics and Space Administration (NASA) (e.g. Moderate Resolution Imaging Spectroradiometer (MODIS) on board the Terra and Aqua satellites). However, mentioned research missions are often characterized by a significantly shorter data record and less spatial coverage due to smaller swath widths.

For the MODIS cloud record however there is the potential to be combined with high quality TOA radiation measurements made by the Clouds and the Earth's Radiant Energy System (CERES) sensors mounted on board the same platforms (Terra and Aqua). In addition to the TOA radiation measurements, CERES BOA radiative fluxes are available based on simulations (Kato et al., 2013). Together with available clear-sky fluxes, this set-up provides an excellent basis for analysing the radiative effect of clouds on TOA and BOA energy balances, although the MODIS and CERES records exist only from the year 2000 onwards. Limitations to resolve small scale clouds and their radiative effect might arise from the coarse spatial resolution of CERES (footprint size of approximately 30 km) and from the fact that the clear-sky fluxes are exclusively based on clear-sky pixels

(and interpolation of clear-sky fluxes for gap filling on monthly scales), by which the spatio-temporal sampling is reduced and in which the meteorological conditions are likely to be biased.

The World Climate Research Programme's (WCRP) Global Energy and Water Exchanges (GEWEX) Surface Radiation Budget (SRB) dataset (Stackhouse et al., 2011) is generated by application of a different approach. Here, retrieved cloud properties are used together with reanalysis information and additional radiative transfer calculations in order to determine all-sky and clear-sky fluxes at the same time for each pixel. The latest release of the GEWEX SRB dataset (v3.0), however, only covers a period until 2007. It makes use of ISCCP DX data (Rossow and Schiffer, 1999), which provides information on a temporal resolution of 3 hours, but includes some deficiencies such as utilizing less spectral information compared to AVHRR-based data, and a relatively coarse spatial resolution. The GEWEX SRB data have been used to revisit the cloud radiative effect on the global scale (e.g. Allan, 2011).

Based on the rationale above it seems logical to construct a record that includes both cloud and radiation properties based on AVHRR, covers a longer time period than alternative records, provides information at finer spatial scales (about 5 km for AVHRR global area coverage - GAC - data) and makes use of all 5 available spectral bands from the visible through the near-infrared to the thermal infrared. The availability of the full suites of cloud and radiative flux properties will also make these data superior to the already existing AVHRR-based datasets mentioned above. The usefulness of these data is further enhanced by the incorporation of the latest AVHRR intercalibration information and cloud retrieval developments.

This paper documents the approaches that have been followed to generate such an AVHRR-based data record with cloud and broadband radiative flux properties, and discusses derived results. The dataset is named Cloud_cci AVHRR-PMv3 (v3 hereafter) and is a successor of AVHRR-PMv2 (v2 hereafter), which contained cloud properties for the period 1982-2014 (see Stengel et al. (2017) for more details), and was already used in numerous studies, e.g. in model evaluation on the global scale (Lauer et al., 2017; Stengel et al., 2018; Eliasson et al., 2018) and on regional scales (Keller et al., 2018; Baró et al., 2018).

Superior to AVHRR-PMv2, AVHRR-PMv3 covers a longer time period (1982-2016), holds cloud properties of improved quality and includes broadband radiative flux properties at TOA and BOA. Appendix A lists additional information about the AVHRR measurement record used. To estimate the radiative fluxes additional radiative transfer calculations were conducted that included additional reanalysis information of tropospheric profiles of temperature and gaseous components as well as surface properties (all interpolated to AVHRR temporal and spatial resolution). This approach is similar to the GEWEX SRB data, thus the retrieved cloud properties are ingested into the reanalysis profiles to represent real clouds with realistic properties at the correct time and place. This is considered a superior approach compared to using reanalysis (thus modelled) clouds directly. All of this information is then input to calculate the broadband fluxes. Although a considerable amount of reanalysis data is still required, this approach provides a means for quantifying the impact of true (retrieved) cloud properties on radiative fluxes at TOA and BOA in a realistic way. This also enables the collection of clear-sky fluxes at the same temporal frequency as all-sky fluxes as opposed to collecting and interpolating the clear-sky fluxes into cloudy areas as is done for the CERES datasets.

In this paper the Cloud_cci AVHRR-PMv3 dataset is summarized. The following Section 2 reports recent cloud retrieval developments and updates, shows product examples and presents validation results all incorporating equivalent results from

the precursor dataset version (v2). Section 3 introduces the radiative flux properties and the algorithms they are based on, and, as for cloud properties, presents product examples and evaluation results. Section 4 gives a summary.

2 Cloud properties

The set of cloud properties included in v3 is identical to v2 and is outlined in the upper part of Table 1, which also gives all cloud property abbreviations used throughout the paper. All data are collected on two processing levels: (a) Level-3U which represents daily composites of non-averaged data collected on a global latitude-longitude grid with 0.05° resolution and (b) Level-3C which represents monthly averages and monthly histograms on a global latitude-longitude grid with 0.5° resolution. Input to Level-3U and Level-3C products are pixel-based retrievals using the algorithms described below. Further Level-3U and Level-3C specifications, i.e. the separation of data into liquid/ice sublayers as well as the histograms binning remain identical to v2 (see Tables 4 and 5 of Stengel et al. (2017)). The propagation of derived pixel-level uncertainties into the higher level products Level-3U and Level-3C remains identical to Stengel et al. (2017) as well.

2.1 Algorithms

The retrieval system employed for cloud properties is the Community Cloud retrieval for CLimate (CC4CL), which is summarized in Stengel et al. (2017) and described in detail in Sus et al. (2018) and McGarragh et al. (2018). However, further developments have taken place since v2, of which the key elements are listed in the following paragraphs. These improvements are grouped according to the CC4CL subcomponents: cloud masking and cloud phase determination, which now both employ Artificial Neural Network (ANN) schemes and require spectral band adjustments (SBAs), and a component for retrieving the remaining cloud properties using an optimal estimation technique (e.g. Rodgers, 2000).

- **Cloud mask:** The ANN for cloud detection (ANN_{mask}) has been retrained using a much larger set of training data (approx. 10 times more collocation data used for v3 than for v2), which is composed of collocations between AVHRR measurements and cloud optical depth observed by the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP, Winker et al., 2009). In addition, a different set of channels for daytime conditions, the $3.7 \mu m$ channel is now included in the ANN scheme (exception: $1.6 \mu m$ is used for NOAA-16 for the period 04/2001 through 04/2003). Table B1 summarizes the ANN_{mask} input data as a function of illumination conditions, while Table B2 reports the empirical thresholds applied posterior to convert the ANN_{mask} output to a binary cloud mask. Downstream, cloud detection is complemented by an additional cirrus test based on $10.8 \mu m$ and $12.0 \mu m$ IR measurements as defined in Pavolonis et al. (2005). As the cloud detection was developed and fine-tuned for AVHRR on board the NOAA-19 satellites, SBAs are applied for other sensors, which is described in Appendix C. Cloud detection improvements compared to v2 are mainly found for daytime and twilight conditions in general, but in particular also for conditions with snow or ice covered surfaces and in cases of low-level liquid clouds over the sub-tropical and tropical oceans. Validation scores are presented in Section 2.3 reflecting the improvements on the global scale.

Table 1. Cloud_cci AVHRR-PMv3 cloud and radiation properties. ANN_{mask} = artificial neural network for cloud detection, ANN_{phase} = artificial neural network for cloud phase, SV = state vector, PP = postprocessed, PV = Pavlonis algorithm (Pavlonis and Heidinger, 2004; Pavlonis et al., 2005), OE = optimal estimation, BR = BUGSrad (radiative flux algorithm), TOA = top-of-atmosphere, BOA = bottom-of-atmosphere (surface), LW = longwave, SW = shortwave. Upper part of the table (cloud properties) has been adopted from Sus et al. (2018).

Variable name	Abbreviation	Unit	Origin	Comment
Cloud properties				
cloud mask /	CMA /	1	ANN _{mask}	binary cloud occurrence classification /
cloud fraction	CFC	%		Fraction of cloudy pixels
cloud phase /	CPH /	1	ANN _{phase}	binary cloud phase classification /
liquid cloud fraction	LCF	%		Fraction of liquid clouds
cloud top pressure	CTP	hPa	SV	OE retrieval result of cloud top pressure
cloud top height	CTH	km	PP	derived from CTP and atmospheric profile
cloud top temperature	CTT	Kelvin	PP	derived from CTP and atmospheric profile
cloud effective radius	CER	μm	SV	OE retrieval result of cloud effective radius
cloud optical thickness	COT	1	SV	OE retrieval result of cloud optical thickness
surface temperature	STEMP	Kelvin	SV	OE retrieval result of surface temperature
cloud water path	CWP	g m^{-2}	PP	derived from CER and COT (Stephens, 1978)
cloud albedo at $0.6 \mu\text{m}$	CLA _{0.6}	1	PP	derived from CER and COT
cloud albedo at $0.8 \mu\text{m}$	CLA _{0.8}	1	PP	derived from CER and COT
cloud effective emissivity	CEE	1	PP	derived from 10.8 and $12.0 \mu\text{m}$ data
Broadband flux properties				
TOA upwelling SW flux	SWF _{TOA} ^{up} , clearSWF _{TOA} ^{up}	W m^{-2}	BR	all-sky and clear-sky
TOA upwelling LW flux	LWF _{TOA} ^{up} , clearLWF _{TOA} ^{up}	W m^{-2}	BR	all-sky and clear-sky
BOA upwelling SW flux	SWF _{BOA} ^{up} , clearSWF _{BOA} ^{up}	W m^{-2}	BR	all-sky and clear-sky
BOA upwelling LW flux	LWF _{BOA} ^{up} , clearLWF _{BOA} ^{up}	W m^{-2}	BR	all-sky and clear-sky
BOA downwelling SW flux	SWF _{BOA} ^{down} , clearSWF _{BOA} ^{down}	W m^{-2}	BR	all-sky and clear-sky
BOA downwelling LW flux	LWF _{BOA} ^{down} , clearLWF _{BOA} ^{down}	W m^{-2}	BR	all-sky and clear-sky
Photosynthetic Active Radiation	PAR	W m^{-2}	BR	total and diff

Please note, retrievals of CER, COT, CWP and CLA are also provided during night-time, although as experimental products. Under these illumination conditions the associated uncertainty can be large and should be inspected, and any usage of these data should be done with caution.

- **Cloud top phase** determination, which in v2 was inferred from the cloud typing procedure of Pavlonis and Heidinger (2004) and Pavlonis et al. (2005), was replaced by a ANN approach for v3 (ANN_{phase}). The strategy for training the ANN_{phase} was very similar compared to the cloud detection approach, training the ANN_{phase} to emulate CALIOP cloud

top phase using AVHRR measurements as primary input data. The exact list of input data for the ANN_{phase} is given in Table B3. Table B4 lists the thresholds applied to convert the ANN_{phase} output to a binary cloud phase. As for cloud detection, SBAs are applied prior to the cloud phase determination (see Appendix C). Significant improvements are found for the cloud phase in v3 compared to v2 when analysing validation results against CALIOP as reported in Section 2.3.

- 5 • **OE retrieval of cloud properties:** The surface reflectance model was revised leading to a corrected handling of the solar zenith angle with most pronounced changes at large angles. Furthermore, bugs were fixed in the code that composes the look-up tables (LUTs) based on pre-calculated radiative transfer simulations. In particular the LUTs for channels with solar reflectance contribution changed considerably. This led to smaller CER retrievals for $3.7\mu m$ measurements, in particular for CER_{ice} . Introducing the utilization of the ice cloud single-scattering properties of Baum et al. (2014) (Baran et al. (2005) used before) further reduced the CER_{ice} . For AVHRR-PMv3 cloud optical properties are also retrieved during night-time, facilitated by a differential sensitivity of the radiation in the spectral bands $3.7\mu m$ and $10.8\mu m/12.0\mu m$ to COT and CER. Night-time COT and CER retrievals are considered as experimental products and only included in Level-3U products. All retrieved cloud properties are input to the calculation of the radiative fluxes as described in Section 3. As for v2, retrievals of COT and CER are used in v3 to determine LWP and IWP following Stephens (1978).

15 2.2 Cloud property examples

Figure 1 shows global maps of monthly mean CFC, LCF, COT and CER for June 2014 for v3 Level-3C data - along with the same data from v2. In general, global patterns look very similar with only minor differences between v3 and v2 for CFC and COT. LCF increased (more liquid clouds) from v3 to v2 after a fundamental change of the phase detection approach (see above). CER of v3 is significantly lower than in v2, which is mainly due to fixing a bug in some CC4CL LUTs and introducing alternative single scattering properties, as mentioned in Section 2.1, which only affected retrieved ice cloud properties.

Figure 2 presents the same comparison for CTP, LWP, IWP and $CLA_{0.6}$. Global patterns remain very similar again. Mean CTP is **higher** in v3 than in v2 in the Tropics, which is predominantly due to detecting more very-low level clouds above tropical oceans. While LWP remains similar in v3 compared to v2, IWP is significantly lower in v3 due to lower CER_{ice} (input to the IWP calculation). Unrealistically high LWP and IWP values in polar regions are reduced in v3 due to reduced CER. $CLA_{0.6}$ is slightly higher in v3 compared to v2 although the changes are relatively small.

Detailed validation was carried out for all cloud properties for which accurate reference data exist. The results of those efforts are presented in the next section, highlighting the quality of the v3 data.

2.3 Validation

Cloud_cci AVHRR-PMv3 CMA, CPH and CTH Level-3U products were collocated with equivalent CALIOP products which are assumed to be of superior quality. More specifically, the CAL_LID_L2_05kmCLay-Prov product was downloaded from the ICARE Data and Service Center (<http://www.icare.univ-lille1.fr>). **The collocations between CALIOP and the AVHRR-PM data were done as reported in Stengel et al. (2017) with the most important fact being that only those collocations were**




Figure 1. Examples of Level-3C (monthly means) Cloud_cci AVHRR-PMv3 data for cloud fraction CFC (a), liquid cloud fraction LCF (b), cloud optical thickness COT (c) and cloud effective radius CER (d). Same data is shown for v2 (e-h). **Difference maps are shown in panels i-l.** All data is from June 2014.




Figure 2. As Figure 1 but for CTP, LWP, IWP and CLA

included for which the spatial and temporal mismatch was below 5 km and 3 minutes, respectively. These criteria were chosen as compromise between using best spatial and temporal matches, and allowing for a compositions of a sound data basis to be used in the validation. Important to note that the random deviations of AVHRR-PM to CALIOP depend on the defined criteria, while the systematic do most likely not. To investigate the sensitivity of passive imager retrievals to the thinnest cloud

5 layers, the cloud optical depth profiles included in the CALIOP profiles were employed as in Karlsson and Johansson (2013); Stengel et al. (2013); Sus et al. (2018). Following this approach different scenarios for excluding optically thin cloud layers are investigated when discussing validation of CMA, CPH and CTH below.

In addition to the validation against CALIOP, Cloud_cci AVHRR-PMv3 LWP was collocated with AMSR-E observations of LWP (Wentz and Meissner, 2004), and IWP was collocated to DARDAR (raDAR/liDAR Delanoë and Hogan, 2008, 2010) observations of IWP. Passive microwave observations of AMSR-E over ocean and active observations of CALIOP and Cloud-sat in DARDAR are assumed to provide best reference data for LWP and IWP on global scales. All validation results are accompanied by the equivalent results for v2.

Table 2 reports the validation results for CMA for two scenarios: (1) considering all CALIOP reference pixels as cloudy for which the CALIOP COT is above 0.0 ($COT_{thres}=0.0$), and (2) considering only those CALIOP reference pixels as cloudy for which the CALIOP COT is above 0.15 ($COT_{thres}=0.15$). The latter scenario is added to account for the lack of sensitivity of AVHRR measurements to very optically thin clouds. For both scenarios, the scores are generally better for v3 than for v2. Heidke Skill Scores (Heidke, 1926), hitrates and probabilities of detections (PODs) are higher (thus better). The only degradation in the scores is found for the bias, which is slightly more negative in v3 compared to v2.

Table 3 reports the validation results for CPH, also for two scenarios: (1) using the cloud phase at the top of the uppermost cloud layer detected by CALIOP as reference ($COT_{lev}=0.0$), and (2) using the cloud phase at an optical depth of 0.15 into the cloud (top-down) as reference ($COT_{lev}=0.15$). Comparing the HSS score as overall measure for the correct cloud phase detection, v3 performs better than v2. The POD of liquid clouds is significantly improved in v3, while a small degradation in POD of ice clouds is found in v3 compared to v2. The liquid bias has increased for v3. Removing the thinnest cloud layers, thus accounting for the AVHRR sensor limitation, the improvement of v3 over v2 becomes even clearer. In this scenario, the cloud phase of 84.7 % of all clouds is correctly identified in v3 (according to hitrate scores). It is important to note that the CALIOP data used for validation of cloud detection and cloud phase determination excluded the data that was used for training the ANNs.

Table 4 reports the validation results for CTH. The validation is stratified by the phase of the cloud and by optical depth into the cloud (top-down) at which the reference CTH is taken from the CALIOP profile. In addition to COT_{lev} of 0.0 and 0.15, COT_{lev} of 1.0 is also included. Generally only few changes in validations scores are found between v3 and v2. While for liquid clouds the scores remain nearly the same, a small degradation in the CTH bias for ice clouds is found. The underestimation of CTH is stronger in v3 compared to v2. For example for the geometrical CTH from CALIOP ($COT_{lev}=0.0$) the bias degrades from -2.59 km to -3.54 km. One reason for this can be that the LUT-related bug fixes (see Section 2.1) led to smaller CER_{ice} . Smaller ice particles absorb less radiation coming from below the cloud putting the cloud lower in the atmosphere in the retrieval. In contrast to the bias, standard deviations are reduced for v3 amounting to 2.36 km compared to 2.51 km in v2. For

$COT_{lev}=0.15$ and $COT_{lev}=1.0$, very similar findings are made with both of these scenarios showing the reduction in bias and standard deviation with increasing COT_{lev} for ice clouds. This highlights the difficulties in correctly placing (vertically) optically thin clouds and cloud layers when using AVHRR measurements. **Figure 3 shows 2-dimensional frequency distributions of all data included in the CTH validation statistics for $COT_{lev}=1.0$ (panel a for liquid clouds and panel b for ice clouds).**




Figure 3. Two-dimensional frequency distributions of AVHRR-PMv3 cloud properties (panel a: cloud top height (CTH) of liquid clouds; panel b: cloud top height of ice clouds; c: liquid water path (LWP); d: ice water path (IWP)) collocated with corresponding reference products of CALIOP, AMSR-E and DARDAR.

- 5 Table 5 reports the validation results for LWP. Although the bias for v3 remains small when compared against AMSR-E, it is slightly increased compare to v2 from -1.9 g m^{-2} to -3.2 g m^{-2} . Standard deviations are slightly decreased for v3 (26.4 g m^{-2}) compared to v2 (27.1 g m^{-2}) and the correlation remains unchanged at 0.65. **Figure 3 (panel c) shows the 2-dimensional frequency distribution of all data included in the LWP validation statistics.**

- 10 Table 6 reports the validation results for IWP. The AVHRR-PM IWP generally shows an underestimation of IWP when DARDAR is considered as reference. This underestimation has increased for v3 as the bias has become larger and negative (-307.1 g m^{-2} for v3 compared to 33.3 g m^{-2} for v2). However, the standard deviation has decreased significantly from 1299.8 g m^{-2} to 900.9 g m^{-2} along with a clear increase in correlation from 0.42 to 0.63. **Figure 3 (panel d) shows the 2-dimensional frequency distribution of all data included in the IWP validation statistics.**

- 15 Despite the assumption that the reference data used is of higher quality than the Cloud_cci data, uncertainties and inaccuracies remain in the reference data as well, which should be kept in mind when interpreting the presented validation scores. However, summarizing the discussion above, the cloud properties included in AVHRR-PMv3 are considered to be of superior quality than the precursor version.

An even broader assessment of the quality of the presented dataset can be found in PVIR (2019), in which the results are also stratified by illumination conditions among others.

Table 2. Cloud mask (CMA) validation results for Cloud_cci AVHRR-PMv3 when compared against CALIOP. Validation results for AVHRR-PMv2 are also reported. Validation measures are the probabilities of detecting cloudy and clear scenes (Hitrate , $\text{POD}_{\text{cloudy}}$, $\text{POD}_{\text{clear}}$) and bias. In addition, the number of collocated pixels is given. The scores are separated into two cloud optical thickness thresholds ($\text{COT}_{\text{thres}}$) reflecting above which CALIOP COT the CALIOP pixel was classified cloudy.

Score		AVHRR-PMv3	AVHRR-PMv2
$\text{COT}_{\text{thres}} = 0.0$	HSS	0.68	0.64
	Hitrate [%]	79.23	78.17
	$\text{POD}_{\text{cloudy}}$ [%]	75.82	75.46
	$\text{POD}_{\text{clear}}$ [%]	91.98	88.31
	Bias [%]	-17.38	-16.89
	Number	16 139 764	16 139 764
$\text{COT}_{\text{thres}} = 0.15$	HSS	0.66	0.63
	Hitrate [%]	83.01	81.86
	$\text{POD}_{\text{cloudy}}$ [%]	83.29	82.79
	$\text{POD}_{\text{clear}}$ [%]	82.45	79.97
	Bias [%]	-5.35	-4.86
	Number	16 139 764	16 139 764

Table 3. Cloud phase (CPH) validation results for Cloud_cci AVHRR-PMv3 when compared against CALIOP. Validation results for AVHRR-PMv2 are also reported. Validation measures are the probabilities of detecting liquid and ice phase (Hitrate, POD_{liq} , POD_{ice} and bias of liquid cloud occurrence). In addition the number of collocated pixels is given. The scores are separated into two cloud optical depth levels (COT_{lev}) representing at which top-down COT into the cloud the reference CALIOP CPH was taken.

Score		AVHRR-PMv3	AVHRR-PMv2
$COT_{lev} = 0.0$	HSS	0.62	0.56
	Hitrate [%]	79.74	77.87
	POD_{liq} [%]	86.25	78.02
	POD_{ice} [%]	75.44	77.77
	Bias [%]	9.35	4.67
	Number	8 788 655	8 788 655
$COT_{lev} = 0.15$	HSS	0.69	0.62
	Hitrate [%]	84.70	80.99
	POD_{liq} [%]	82.33	74.06
	POD_{ice} [%]	87.16	88.20
	Bias [%]	-2.72	-7.44
	Number	8 435 631	8 435 631

Table 4. Cloud-top height (CTH) validation results for Cloud_cci AVHRR-PMv3 when compared against CALIOP. Validation results for AVHRR-PMv2 are also reported. Validation measures are **standard deviation of the error (Std)** and the **mean error (bias)**. In addition the number of collocated pixels is given. All scores are separated into liquid and ice clouds (both Cloud_cci dataset and CALIOP had to agree on phase) and into three cloud optical depth levels (COT_{lev}) representing at which top-down COT into the cloud the reference CALIOP CTH was taken.

Score		AVHRR-PMv3	AVHRR-PMv2
$COT_{lev} = 0.0$	Std _{liq} [km]	0.86	0.86
	Bias _{liq} [km]	-0.10	-0.11
	Number _{liq}	2 603 163	2 603 163
	Std _{ice} [km]	2.36	2.51
	Bias _{ice} [km]	-3.54	-2.59
	Number _{ice}	3 691 179	3 691 179
$COT_{lev} = 0.15$	Std _{liq} [km]	0.91	0.91
	Bias _{liq} [km]	-0.06	-0.08
	Number _{liq}	3 016 985	3 016 985
	Std _{ice} [km]	2.14	2.30
	Bias _{ice} [km]	-2.95	-2.00
	Number _{ice}	3 376 337	3 376 337
$COT_{lev} = 1.0$	Std _{liq} [km]	0.80	0.80
	Bias _{liq} [km]	0.05	0.04
	Number _{liq}	2 982 690	2 982 690
	Std _{ice} [km]	1.95	2.09
	Bias _{ice} [km]	-1.62	-0.84
	Number _{ice}	2 077 074	2 077 074

Table 5. Liquid water path (LWP) validation results for Cloud_cci AVHRR-PMv3 when compared against AMSR-E for the year 2008. Validation results for AVHRR-PMv2 for the same time period are also reported. Validation measures are **standard deviation of the error (Std)**, the **mean error (bias)** and correlation. In addition the number of collocated pixels is given.

Score	AVHRR-PMv3	AVHRR-PMv2
Std [$g\ m^{-2}$]	26.4	27.1
Bias [$g\ m^{-2}$]	-3.2	-1.9
Correlation	0.64	0.64
Number	183 022	183 022

Table 6. Ice water path (IWP) validation results for Cloud_cci AVHRR-PMv3 when compared against DARDAR for January to July 2008. Validation results for AVHRR-PMv2 for the same time period are also reported. Validation measures are **standard deviation of the error (Std), the mean error (bias)** and correlation. In addition the number of collocated pixels is given.

Score	AVHRR-PMv3	AVHRR-PMv2
Std [g m ⁻²]	900.9	1299.8
Bias [g m ⁻²]	-307.1	33.3
Correlation	0.63	0.42
Number	92 293	92 293

3 Radiation properties

In addition to the cloud properties described in the previous section, radiative broadband flux properties (shortwave and long-wave) at TOA and BOA, and for all-sky and clear-sky conditions, were calculated employing the BUGSrad scheme (Stephens et al., 2001, more details below). Furthermore, the photosynthetically active radiation was determined. A full list of radiation properties is given in the bottom part of Table 1. As for the cloud properties, all radiation properties are derived at pixel level, sub sampled to daily, global composites (Level-3U products) and aggregated to monthly Level-3C products.

3.1 Algorithm

BUGSrad uses a two-stream approximation along with correlated-k distribution methods for atmospheric radiative transfer (Fu and Liou, 1992). It has been used to investigate aerosol-cloud interactions (Christensen et al., 2017) and to assess the Earth’s energy budget using CloudSat observations (Stephens et al., 2012). BUGSrad is applied to a single column, plane-parallel atmosphere with ingested cloud properties (i.e. CER, COT, CTP) previously retrieved with CC4CL (see Section 2.1). BUGSrad uses 18 spectral bands in the electromagnetic spectrum (6 in the shortwave and 12 in the longwave) to compute the broadband fluxes. Atmospheric profiles for temperature and water vapour are taken from ERA-Interim. Visible and near-infrared surface albedo are based on spatiotemporally resolved MODIS climatologies - with all data being identical to the usage in CC4CL. Total solar irradiance is based on SOHO (Solar and Heliospheric Observatory) and SORCE (Solar Radiation and Climate Experiment) measurements acquired from http://disc.sci.gsfc.nasa.gov/SORCE/data-holdingsusingSOR3TSID_v017 and further processed by applying a bi-linear interpolation followed by a bias correction to SOHO measurements to match SORCE. For well-mixed radiatively important trace gases constant values are used (CH₄ = 1.8 ppm, N₂O = 0.26 ppm). For CO₂ a linearly time dependent concentration is used anchored at 380 ppm for the year 2006. To account for the effect of aerosols on the radiation, an aerosol optical depth of 0.05 was added to the extinction throughout the atmosphere. It is acknowledged that this value is under-representing heavy aerosol loadings which motivates the utilization of spatiotemporally resolved aerosol information for future dataset versions. **The reader is referred to ATBD-CC4CL-BBFlux (2019) for more details on the calculation of the broadband fluxes.**

Due to the angular dependence of the solar illumination together with the low sampling frequency of a single polar-orbiting AVHRR sensor, an angular dependence correction is applied to the shortwave radiation properties to make the data represent 24 hour averages. This is done by calculating the diurnal cycle of the solar zenith angle (SZA) for a given pixel on the day of observation. The diurnal cycle of SZA is then used to rescale the incoming and reflected solar radiation, adjust the surface albedo (using an empirical quadratic function of SZA) and the atmospheric path length for a given set of time stamps throughout the local day. Averaging these samples gives a suitable approximation for a true 24 hour mean, which is needed to determine true climatological means. This procedure is however only applied for Level-3C products, while Level-3U products hold the instantaneous, uncorrected fluxes representing the solar illumination at the pixel location and at the time of observation.

For longwave radiation, a diurnal cycle correction is applied over land based on a cosine fit to an observed mean diurnal cycle by applying CC4CL to geostationary Spinning Enhanced Visible and Infrared Imager (SEVIRI). The observed diurnal cycle is converted to a correction factor, which itself is a function of local observation time, to mimic a 24 hour mean.

In contrast to the cloud properties, the radiative fluxes in the presented dataset version are not accompanied by uncertainty estimates on pixel level. While the validation results presented below provide a general guidance to the quality of the radiative fluxes, users of the data are also encouraged to inspect the pixel-level uncertainties of the cloud properties as these are dominant input to the calculation of the fluxes.

3.2 Radiation property examples

Figure 4 shows examples of Cloud_cci AVHRR-PMv3 Level-3C data of SWF_{TOA}^{up} , LWF_{TOA}^{up} , SWF_{BOA}^{down} , LWF_{BOA}^{down} for all-sky and clear-sky conditions for June 2014. As a general description of these properties, high clear SWF_{TOA}^{up} is found in regions with high surface albedo while high values in SWF_{TOA}^{up} are additional visible in regions with high cloud fraction, and vice versa. Clear SWF_{TOA}^{up} and SWF_{TOA}^{up} depend on incoming solar flux, which, in the month of June, is highest in the tropics and Northern Hemisphere. Clear LWF_{TOA}^{up} is highest in regions with high surface temperatures and low water vapour amounts in the atmospheric column above. Higher water vapour loadings and in particular frequent occurrence of cold clouds significantly reduce the LWF_{TOA}^{up} , for example visible in the tropics and the mid-latitudes. SWF_{BOA}^{down} represents the downwelling solar radiation that is neither reflected nor absorbed by clouds or the atmosphere, thus is, roughly speaking, high where SWF_{TOA}^{up} is low and vice versa. SWF_{BOA}^{down} and clear SWF_{BOA}^{down} strongly depend on illumination conditions. LWF_{BOA}^{down} represents the downwelling radiation emitted by the atmosphere and clouds and is high in regions with high water vapour amounts and further increased when clouds are frequently present.

The product portfolio for radiative fluxes is complemented by SWF_{TOA}^{down} : the incoming solar radiation at the top of the atmosphere and SWF_{BOA}^{up} (LWF_{BOA}^{up}): the reflected solar (emitted terrestrial) radiation at the Earth's surface (not shown).




Figure 4. Examples of Level-3C (monthly means) Cloud_cci AVHRR-PMv3 data of $\text{SWF}_{\text{TOA}}^{\text{up}}$, $\text{LWF}_{\text{TOA}}^{\text{up}}$, $\text{SWF}_{\text{BOA}}^{\text{down}}$, $\text{LWF}_{\text{BOA}}^{\text{down}}$ (a-d). $\text{clearSWF}_{\text{TOA}}^{\text{up}}$, $\text{clearLWF}_{\text{TOA}}^{\text{up}}$, $\text{clearSWF}_{\text{BOA}}^{\text{down}}$, $\text{clearLWF}_{\text{BOA}}^{\text{down}}$ are shown in (e-h) representing the same fluxes as in (a-d) but for clear-sky conditions. All data is from June 2014.

3.3 Validation

3.3.1 BOA radiative fluxes

The Cloud_cci AVHRR-PMv3 BOA radiative fluxes $\text{SWF}_{\text{BOA}}^{\text{down}}$ and $\text{LWF}_{\text{BOA}}^{\text{down}}$ were compared against ground-based reference stations of the World Radiation Monitoring Center (WRMC) Baseline Surface Radiation Network (BSRN, Driemel et al., 2018). For this, monthly mean BSRN $\text{SWF}_{\text{BOA}}^{\text{down}}$ and $\text{LWF}_{\text{BOA}}^{\text{down}}$ values were calculated per station from all available observations and then compared to the nearest neighbouring Cloud_cci grid box. Figure 5 shows scatter plots for all monthly pairs found within the period of 2003 to 2016. The validation scores are reported in Table 7. An excellent agreement of the Cloud_cci with the reference BSRN measurements is found for both $\text{SWF}_{\text{BOA}}^{\text{down}}$ and $\text{LWF}_{\text{BOA}}^{\text{down}}$ products with correlations above 0.98. Standard deviations are 13.8 W m^{-2} for $\text{SWF}_{\text{BOA}}^{\text{down}}$ and 11.5 W m^{-2} for $\text{LWF}_{\text{BOA}}^{\text{down}}$. The comparisons further reveal positive biases in Cloud_cci: 1.9 W m^{-2} for $\text{SWF}_{\text{BOA}}^{\text{down}}$ and 7.6 W m^{-2} for $\text{LWF}_{\text{BOA}}^{\text{down}}$. Considering the $\text{LWF}_{\text{BOA}}^{\text{down}}$ bias, Nyeki et al. (2017) recently found indications that the measured fluxes at BSRN stations are biased low. They quantified this with 3.5 to 5.4 W m^{-2} which has the potential to explain more than 50 % of the bias found between Cloud_cci and BSRN for $\text{LWF}_{\text{BOA}}^{\text{down}}$. Figure 5 also shows equivalent validation for upwelling fluxes at those BSRN sites which provide upwelling measurements (much fewer stations than for downwelling fluxes). For $\text{LWF}_{\text{BOA}}^{\text{up}}$ the agreement of Cloud_cci to BSRN is again very good with standard deviation of 14.1 W m^{-2} , a bias of -3.0 W m^{-2} and a correlation of 0.99.

In general, the agreement of the Cloud_cci $\text{SWF}_{\text{BOA}}^{\text{down}}$, $\text{LWF}_{\text{BOA}}^{\text{down}}$ and $\text{LWF}_{\text{BOA}}^{\text{up}}$ with the BSRN stations is remarkable when considering that only one satellite sensor is used at a time, thus for many locations on Earth only two satellite overpasses (one daytime, one night-time) within 24 hours provide observations. The results are a confirmation that the developed and applied diurnal cycle correction works well, which is more important for the shortwave than for the longwave fluxes.

In contrast, for SWF_{BOA}^{up} more scatter is found in the comparisons to BSRN. Considering SWF_{BOA}^{up} is simply the SWF_{BOA}^{down} multiplied with the surface albedo, and the good validation results for SWF_{BOA}^{down} , this leads to the conclusions that either imperfect surface albedo was used in Cloud_cci, or, more likely, the difference in spatial scales might be the dominating source of the discrepancy found. Fine scale inhomogeneities in surface albedo in the vicinity of the BSRN stations will propagate into the results.

In addition to the BSRN stations, Cloud_cci BOA downwelling and upwelling fluxes were compared to the Clouds and the Earth's Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) surface flux product (Kato et al., 2013), i.e. by means of comparing multi-annual mean maps for the period 2003-2016 (Figures 6 and 7), with corresponding latitude-weighted global mean values given in Table 8.

The Cloud_cci multi-annual mean maps of SWF_{BOA}^{down} for the chosen period agree very well with the CERES products (panels a and b of Figure 6), also for the clear-sky fluxes (panels d and e of Figure 6). This is also supported by global mean values reported in Table 8 in which Cloud_cci is slightly biased high (+0.9 $W m^{-2}$ for SWF_{BOA}^{down} and +2.2 $W m^{-2}$ for clear SWF_{BOA}^{down}). Clear-sky fluxes in both products are mainly characterized by larger incoming solar radiation at the equator, scattering and absorption by atmospheric gases and aerosols, and the surface reflectivity and emissivity. The presence of clouds usually leads to a significant reduction of SWF_{BOA}^{down} locally being a function of optical thickness and cloud fraction over larger domains. The fact that the all-sky fluxes SWF_{BOA}^{down} agree very well with CERES validates the Cloud_cci cloud detection and corresponding cloud property retrievals, which thus can be assumed to be of high quality.

The Cloud_cci multi-annual mean maps of LWF_{BOA}^{down} (Figure 7) also agree well with CERES in terms of global patterns. The absolute values however show systematically higher values for Cloud_cci of about 8 to 9 $W m^{-2}$ for both all-sky and clear-sky values. **The positive bias is relative homogenous over the globe.** In relative terms the systematic differences amount to approximately 2 to 3 %. However, these differences lie within the expected range of the CERES accuracy (Rutan et al., 2015).

The Cloud_cci multi-annual mean maps of SWF_{BOA}^{up} exhibit larger systematic deviations (not shown) than for SWF_{BOA}^{down} . **The larger standard deviations retrieved from the solar reflected radiation is primarily related to variances in surface albedo which tend to have significant annual cycles.** Global mean values reported in Table 9 give negative biases of -2.7 and -4.6 $W m^{-2}$ for Cloud_cci which in relative terms correspond to negative deviations of more than 10 %. It remains uncertain which of the two products are more realistic as no real ground truth is available for SWF_{BOA}^{up} that represents spatial scales of satellite pixels (several kilometres). Repeating the validation of SWF_{BOA}^{up} against BSRN but using CERES gives comparable, large deviations (not shown) as found for Cloud_cci (see above). This is in agreement with findings of (Kratz et al., 2010) who reported systematic deviations between CERES and surface observations of SWF_{BOA}^{up} depending on time of day, meteorological condition and location.

Cloud_cci multi-annual maps of LWF_{BOA}^{up} are again closer to CERES (not shown). Global mean values (Table 9) deviate by approximately 2 $W m^{-2}$ only with larger values for Cloud_cci. In relative terms the differences are about 0.5 %.




Figure 5. Comparison of Cloud_cci bottom-of-atmosphere (BOA) shortwave (SW, panel a) and longwave (LW, panel b) downwelling fluxes with ground-based reference measurements taken at globally distributed Baseline Surface Radiation Network (BSRN) sites for which equivalent reference data was available. Panels (c) and (d) are as (a) and (b) but for upwelling fluxes. Shown are all monthly data pairs within the period 2003-2016.

Table 7. Validation results for monthly Cloud_cci AVHRR-PMv3 shortwave and longwave, downwelling and upwelling radiative fluxes at bottom-of-atmosphere (BOA) when compared against Baseline Surface Radiation Network (BSRN) sites within the period 2003-2016. Validation measures are standard deviation (Std), bias and correlation. In addition the number of data pairs is given.

Score	SWF _{BOA} ^{down}	LWF _{BOA} ^{down}	SWF _{BOA} ^{up}	LWF _{BOA} ^{up}
Std [W m^{-2}]	13.83	11.52	31.18	14.11
Bias [W m^{-2}]	1.99	7.60	-6.16	-3.02
Correlation	0.98	0.99	0.93	0.99
Number	4487	5627	1022	1182

Table 8. Multi-annual (2003-2016), latitude-weighted, global mean downwelling broadband shortwave and longwave fluxes (SWF, LWF) at bottom-of-atmosphere (BOA) inferred from Cloud_cci AVHRR-PMv3 dataset for all-sky and clear-sky (clear) conditions. The values are compared to equivalents inferred from Clouds and the Earth's Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) surface fluxes. All values are given in W m^{-2} . In addition, differences and relative differences (Cloud_cci-CERES) for all fluxes are reported. **For comparison, ERA-Interim values are listed as well.**

	SWF _{BOA} ^{down}	clearSWF _{BOA} ^{down}	LWF _{BOA} ^{down}	clearLWF _{BOA} ^{down}
Cloud_cci AVHRR-PMv3 [W m^{-2}]	188.2	246.1	353.4	325.0
CERES EBAF Ed.4.0 [W m^{-2}]	187.3	243.9	345.4	314.6
difference [W m^{-2}]	+0.9	+2.2	+8.0	+10.4
rel. difference	+0.5 %	+0.9 %	+2.3 %	+3.3 %
ERA-Interim [W m^{-2}]	185.8	245.4	342.4	315.7




Figure 6. Multi-annual (2003-2016) mean downwelling shortwave (SW) radiative fluxes at bottom-of-atmosphere (BOA) for all-sky conditions for Cloud_cci AVHRR-PMv3 (a) and CERES EBAF surface fluxes (b). Panels (d) and (e) show the same data but for clear-sky conditions. Panels (c) and (f) show difference plots Cloud_cci minus CERES.




Figure 7. Multi-annual (2003-2016) mean downwelling longwave (LW) radiative fluxes at bottom-of-atmosphere (BOA) for all-sky conditions for Cloud_cci AVHRR-PMv3 (a) and CERES EBAF surface fluxes (b). Panels (d) and (e) show the same data but for clear-sky only. Panels (c) and (f) show difference plots Cloud_cci minus CERES.

Table 9. As Table 8 but for BOA upwelling broadband fluxes.

	$\text{SWF}_{\text{BOA}}^{\text{up}}$	$\text{clearSWF}_{\text{BOA}}^{\text{up}}$	$\text{LWF}_{\text{BOA}}^{\text{up}}$	$\text{clearLWF}_{\text{BOA}}^{\text{up}}$
Cloud_cci AVHRR-PMv3 [W m^{-2}]	20.6	25.1	400.3	400.3
CERES EBAF Ed.4.0 [W m^{-2}]	23.3	29.7	398.8	398.1
difference [W m^{-2}]	-2.7	-4.6	+1.5	+2.2
rel. difference	-11.6 %	-15.5 %	+0.4 %	+0.5 %
ERA-Interim [W m^{-2}]	23.8	-	397.9	-

3.3.2 TOA radiative fluxes

The Cloud_cci TOA radiative fluxes $\text{SWF}_{\text{TOA}}^{\text{up}}$ and $\text{LWF}_{\text{TOA}}^{\text{up}}$ were compared against the CERES EBAF TOA Edition-4.0 data (Loeb et al., 2018). As for the BOA fluxes the comparison includes multi-annual mean maps for the period 2003-2016. Figure 8 shows the maps for SWF for all-sky and clear-sky ($\text{clearSWF}_{\text{TOA}}^{\text{up}}$) conditions. Cloud_cci global patterns are very similar to those of the CERES products. High $\text{SWF}_{\text{TOA}}^{\text{up}}$ values are found in regions with high surface albedo, e.g. deserts and polar regions, or with high cloud frequency, e.g. in mid-latitude storm track regions in both hemispheres, in the inner-tropical convergence zone and in regions with persistent marine stratocumulus clouds. Most prominent regions with low $\text{SWF}_{\text{TOA}}^{\text{up}}$ values are the subtropical subsidence regions (low cloud frequency) over the ocean (low surface albedo). It can also be seen that Cloud_cci provides slightly higher values in regions with high SWF (mainly land). The comparisons of the clear-sky fluxes give very similar results, with the exception that Cloud_cci has generally slightly lower values than CERES over ocean. The global mean values given in Table 10 reveal differences of 2.9 and -3.3 W m^{-2} for all-sky and clear-sky fluxes, respectively. The smaller values in Cloud_cci clear-sky is partly explained by the differences already found for $\text{SWF}_{\text{BOA}}^{\text{up}}$ (see previous section).

Figure 9 shows the results of an equivalent analysis for upwelling LWF at TOA. The global Cloud_cci patterns are again in very good agreement to CERES. High $\text{LWF}_{\text{TOA}}^{\text{up}}$ are mainly found in tropical and subtropical regions (high surface temperature) with low cloud frequency or in regions with mainly low level clouds (marine stratocumulus or trade cumulus regions), where the cloud top temperatures are relatively warm. On the contrary, $\text{LWF}_{\text{TOA}}^{\text{up}}$ is low in regions with cold surfaces (e.g. polar regions) and regions with a high frequency of cold clouds. Cloud_cci $\text{clearLWF}_{\text{TOA}}^{\text{up}}$ are dominated by surface temperatures, thus decreasing towards higher latitudes; generally showing a very good agreement to CERES. The difference maps however reveal that Cloud_cci has generally lower values than CERES for both all-sky and clear-sky conditions. This is also reflected in the global mean values given in Table 10. This difference is almost doubled when considering clear-sky fluxes, which is likely due to different sampling approaches. While for Cloud_cci all conditions are included (but removing the clouds when existent), CERES clear-sky TOA fluxes are determined by including clear-sky conditions only, which has the potential to bias TOA longwave fluxes high as clear-sky conditions have less water vapour (Sohn et al., 2010). This could be confirmed by a 3-months covering test run in which Cloud_cci $\text{clearLWF}_{\text{TOA}}^{\text{up}}$ was only averaged over clear-sky cases, which led to an increase by about 3 W m^{-2} for the global mean value.




Figure 8. Multi-annual (2003-2016) mean top-of-atmosphere (TOA) upwelling shortwave (SW) radiative fluxes for all-sky conditions for Cloud_cci AVHRR-PMv3 (a) and CERES EBAF TOA edition 4.0 (b). Panels (d) and (e) show the same data but for clear-sky conditions. Panels (c) and (f) show difference plots Cloud_cci minus CERES.




Figure 9. As Figure 8 but for TOA upwelling longwave (LW) fluxes.

4 Summary

As described in this paper, version 3 of the Cloud_cci AVHRR-PM dataset has been generated (and linked to a DOI, (Stengel et al., 2019)). In addition to clouds properties, this new version extends the product portfolio by BOA and TOA broadband radiative fluxes and covers the time period 1982 to 2016.

Table 10. Multi-annual (2003-2016), latitude-weighted, global mean broadband fluxes at the top-of-atmosphere (TOA) inferred from Cloud_cci AVHRR-PMv3 dataset for all-sky and clear-sky (clear) conditions. The values are compared to equivalents inferred from Clouds and the Earth’s Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) TOA Edition-4.0. All values are given in W m^{-2} . In addition, differences and relative differences (Cloud_cci-CERES) of all fluxes are reported. **For comparison, ERA-Interim values are listed as well.**

	$\text{SWF}_{\text{TOA}}^{\text{down}}$	$\text{SWF}_{\text{TOA}}^{\text{up}}$	$\text{clearSWF}_{\text{TOA}}^{\text{up}}$	$\text{LWF}_{\text{TOA}}^{\text{up}}$	$\text{clearLWF}_{\text{TOA}}^{\text{up}}$
Cloud_cci AVHRR-PMv3 [W m^{-2}]	340.5	101.9	50.0	236.4	261.1
CERES EBAF Ed.4.0 [W m^{-2}]	340.3	99.0	53.3	240.3	268.3
difference [W m^{-2}]	+0.2	+2.9	-3.3	-3.9	-7.2
rel. difference	+0.1 %	+2.9 %	-6.2 %	-1.6 %	-2.7 %
ERA-Interim [W m^{-2}]	344.2	101.4	53.6	244.6	264.1

The cloud properties in v3 are superior to v2 in many aspects. This is demonstrated by analyses of global validation results against CALIOP (used for cloud detection, cloud phase, cloud top height), AMSR-E (used for liquid water path) and DARDAR (combined CALIOP and CPR information, used for ice water path). Heidke Skill Scores have increased from 0.64 to 0.68 for cloud detection and from 0.56 to 0.62 for cloud phase assignment. The scores are generally sensitive to whether or not thin clouds are included in the statistical comparisons. The improvements for cloud detection and phase determination in v3 remain conclusive also for scenarios in which very thin clouds are excluded. The validation scores for cloud top height assignment remain nearly identical for liquid clouds, whereas for ice clouds lower standard deviations (2.36 km vs. 2.51 km) but larger negative biases (-3.54 km vs. -2.59 km) are found in v3. Similar results are found for scenarios in which the reference height is taken from below the geometrical top with penetrations optical depths of 0.15 and 1.0. Validation results for liquid water path show a slight reduction in standard deviation for v3 from 27.1 g m^{-2} to 26.4 g m^{-2} , accompanied by a slight increase in bias from -1.9 g m^{-2} to -3.2 g m^{-2} . Correlations remain unchanged at 0.65. Ice water path validation shows reductions of standard deviations for v3 from 1299.8 g m^{-2} to 900.9 g m^{-2} compared to v2 (reduction by 30 %). While the clearly increased correlation coefficient emphasises the improvement in v3 as well, the biases are somewhat larger in v3 compared to v2.

A new contribution to version 3 was the addition of top and bottom-of-atmosphere broadband radiative fluxes. Validation of v3 monthly mean downwelling radiative fluxes at BOA against BSRN stations reveals a very good agreement with low standard deviations of 13.8 W m^{-2} for shortwave and 11.5 W m^{-2} for longwave fluxes and correlation coefficients above 0.98 for both. While the bias for shortwave is small (1.9 W m^{-2}), a somewhat larger positive bias is found for longwave (7.6 W m^{-2}) which is mainly driven by moderate overestimations of larger flux values, but can potentially also partly be due to overestimations in the reference (BSRN).

Comparisons of v3 multi-annual mean values of upwelling and downwelling fluxes at BOA and TOA with CERES additionally emphasises the good quality of the Cloud_cci radiative fluxes in terms of relative spatial pattern and absolute values. Concerning the latter, global mean values of Cloud_cci agree with CERES within 3.3 % for downwelling fluxes at BOA with larger deviations found for longwave fluxes. In contrast, Cloud_cci upwelling longwave fluxes at BOA agree very well with

CERES (below 0.5 %), upwelling shortwave fluxes at BOA show deviations up to about 15 %, although the absolute differences are only 4.6 W m^{-2} at maximum. It however remains uncertain to which extent uncertainties in CERES products contribute to these deviations.

In contrast to the BOA, CERES products for TOA fluxes are mainly based on observational information, thus provide an excellent reference for validation. For all-sky fluxes Cloud_cci agrees to CERES within 3% for global mean values. The differences are increased when considering clear-sky fluxes. It is likely that the different approaches to estimate the mean clear-sky fluxes in Cloud_cci (including all conditions, but removing the clouds) and CERES (including only cloud-free conditions) contribute considerably to these differences.

In summary, Cloud_cci AVHRR-PMv3 represents a dataset of consistent cloud properties and radiative fluxes, which in many aspects is superior to the precursor version v2 as data quality was improved, the product portfolio extended and the covered time period prolonged. Cloud_cci AVHRR-PMv3 offers a large variety of applications including climatological analyses of cloud properties and radiative fluxes as well as their dependency to each other at time scales of several decades.

5 Data availability

For the presented dataset (Cloud_cci AVHRR-PMv3) a DOI has been issued: https://doi.org/10.5676/DWD/ESA_Cloud_cci/AVHRR-PM/V003 (Stengel et al., 2019), of which the landing page points to additional documentation and data download sites. A parallel dataset based on AVHRR on board the NOAA and EUMETSAT morning satellites exists (AVHRR-AMv3), for which a DOI has been issued as well: https://doi.org/10.5676/DWD/ESA_Cloud_cci/AVHRR-AM/V003. The AVHRR-AMv3 dataset provides the feasibility to be combined with AVHRR-PMv3 to increase sampling frequency. However, for the period of NOAA-12 and NOAA-15 the AVHRR-AMv3 dataset is of reduced quality due to the difficult twilight orbits of NOAA-12 and NOAA-15. The CC4CL retrieval system used to produce the data is version controlled and accessible at github: <https://github.com/ORAC-CC/orac/wiki>. The LUT creation code is available at https://github.com/ORAC-CC/create_orac_lut. Both are licensed under the GNU General Public License (GPL) version 3.

Appendix A: AVHRR measurement data

The AVHRR measurement record used as basis for the presented cloud climatology spans the AVHRR/2 and AVHRR/3 sensor generations on board NOAA-7, NOAA-9, NOAA-11, NOAA-14, NOAA-16, NOAA-18 and NOAA-19. Based on the original AVHRR measurements (Local Area Coverage) with 1 km spatial resolution and sampling distance, the Global Area Coverage (GAC) data is globally available, but with reduced spatial resolution and sampling distance. Only every fourth scanline is used and within one scanline four neighbouring pixels are averaged. The AVHRR sensor has an on-board black-body calibration mechanism for its infrared channels. No attempt is made to further recalibrate these measurements. For the visible channels, no calibration is performed on board AVHRR. A recalibration procedure for these channels was applied as a preparatory step based on Devasthale et al. (2017) with further application aspects reported in Schlundt et al. (2017).

Appendix B: Measurement input to the ANNs and the thresholds applied posterior

Table B1. Measurement input to the trained artificial neural network for cloud detection (ANN_{mask}), used for different illumination conditions: daytime, twilight and night-time. The subscript in the table's headline corresponds to the approximate central wavelengths of the channels: 0.6 μ m, 0.8 μ m, 1.6 μ m, 3.7 μ m, 10.8 μ m, 12.0 μ m. In addition to the measurement input, all ANNs require surface temperature, a snow-ice flag and a land-sea flag as input. R=reflectance, BT=brightness temperature

ANN_{mask}	R _{0.6}	R _{0.8}	R _{1.6}	R _{3.7}	BT _{3.7}	BT _{10.8}	BT _{12.0}	BT _{10.8} -BT _{12.0}	BT _{10.8} -BT _{3.7}
Day	✓	✓	-	✓	-	✓	✓	✓	-
Twilight	-	-	-	-	✓	✓	✓	✓	✓
Night	-	-	-	-	✓	✓	✓	✓	✓

Table B2. Empirical thresholds used to convert the output of the cloud mask ANNs to a binary cloud mask. Thresholds depend on illumination conditions and surface type.

Illumination	Surface type	Threshold
Day	Sea ice	0.4
Day	Land ice	0.3
Day	Sea	0.25
Day	Land	0.3
Night	Sea ice	0.45
Night	Land ice	0.35
Night	Sea	0.25
Night	Land	0.3
Twilight	Sea ice	0.5
Twilight	Land ice	0.35
Twilight	Sea	0.35
Twilight	Land	0.45

Table B3. Measurement input to the trained artificial neural network for cloud phase determination (ANN_{phase}), used for different illumination conditions: daytime, twilight and night-time. The subscript in the table’s headline corresponds to the approximate central wavelength of the channels: 0.6 μm , 0.8 μm , 1.6 μm , 3.7 μm , 10.8 μm , 12.0 μm . In addition to the measurement input, all ANNs require a surface type flag containing the values 0:sea,1:land,2:desert,3:sea-ice,4:snow.

ANN_{phase}	$R_{0.6}$	$R_{0.8}$	$R_{1.6}$	$R_{3.7}$	$BT_{3.7}$	$BT_{10.8}$	$BT_{12.0}$	$BT_{10.8}-BT_{12.0}$	$BT_{10.8}-BT_{3.7}$
Day	✓	✓	-	✓	-	✓	✓	✓	-
Twilight	-	-	-	-	✓	✓	✓	✓	✓
Night	-	-	-	-	✓	✓	✓	✓	✓

Table B4. Empirical thresholds used to convert the output of the cloud phase ANNs to a binary cloud phase. Thresholds depend on illumination conditions and surface types.

Illumination	Surface type	Threshold
Day	Sea ice	0.5
Day	Land ice	0.7
Day	Sea	0.55
Day	Land	0.7
Night	Sea ice	0.7
Night	Land ice	0.6
Night	Sea	0.5
Night	Land	0.65
Twilight	Sea ice	0.7
Twilight	Land ice	0.9
Twilight	Sea	0.65
Twilight	Land	0.50

Appendix C: Spectral band adjustment (SBA)

As the cloud detection and cloud phase determination were developed and fine-tuned primarily based on NOAA-19 AVHRR, adjustment factors (slope and offset) were inferred to make all considered AVHRR sensors mimic NOAA-19 AVHRR. The SBAs were inferred from a set of SCIAMACHY and IASI orbits with both of these sensors providing hyperspectral measurements throughout the visible (SCIAMACHY) and infrared (IASI) part of the spectrum, respectively. Using the spectral response functions (SRF) of AVHRR channels 0.6 μm , 0.8 μm , 10.8 μm and 12.0 μm the SCIAMACHY and IASI measurements were convolved to mimic synthetic AVHRR measurements in each footprint of the considered SCIAMACHY and IASI orbits. Using this procedure for all AVHRR sensors (the AVHRR SRFs differ among the individual satellites) and collecting the

synthetic AVHRR measurements in all considered footprints of SCIAMACHY and IASI, a database was composed allowing for linearly fitting all AVHRR sensors to AVHRR onboard NOAA-19. This SBA is applied prior to the application of the cloud detection and cloud phase procedures. No attempt is made to adjust channels 1.6 μm and 3.7 μm as the SCIAMACHY and IASI spectra do not cover the full AVHRR SRF of these channels. All inferred SBAs are given in Table C1. In v2 of the datasets, 5 no SBAs were applied among the AVHRR sensors. As the OE retrieval makes direct use of the SRF of the individual AVHRR sensors, the application of the SBA is not required for the OE retrieval.

Table C1. Linear regression coefficients (slope and offset) applied as spectral band adjustment to either measured reflectances (R_s) or brightness temperature (TBs) of all used AVHRR channels (Ch) and all used sensors to mimic NOAA-19 AVHRR. The subscript in the table's headline corresponds to the approximate central wavelengths of the channels: 0.6 μm , 0.8 μm , 1.6 μm , 3.7 μm , 10.8 μm , 12.0 μm . Reflectances in channels 0.6 μm , 0.8 μm and 1.6 μm are generally not used in twilight and night conditions.

	$R_{0.6}$ slope offset	$R_{0.8}$ slope offset	$R_{1.6}$ slope offset	$BT_{3.7}$ slope offset	$BT_{10.8}$ slope offset	$BT_{12.0}$ slope offset
day						
NOAA-7	1.009 -0.036	1.007 -0.007	1.000 0.000	1.000 0.000	1.000 -0.198	0.991 1.991
NOAA-9	1.009 -0.013	1.006 0.011	1.000 0.000	1.000 0.000	1.000 -0.215	0.988 2.770
NOAA-11	1.009 -0.010	1.005 -0.012	1.000 0.000	1.000 0.000	1.000 -0.170	0.989 2.443
NOAA-14	1.008 0.016	1.011 -0.026	1.000 0.000	1.000 0.000	1.001 -0.446	0.995 1.081
NOAA-16	1.006 -0.039	1.009 0.057	1.000 0.000	1.000 0.000	1.000 -0.095	0.997 0.561
NOAA-18	1.002 -0.013	1.015 0.066	1.000 0.000	1.000 0.000	1.000 -0.214	0.997 0.626
NOAA-19	1.000 0.000	1.000 -0.000	1.000 0.000	1.000 0.000	1.000 -0.000	1.000 0.000
twilight & night						
NOAA-7	-	-	-	1.000 0.000	1.000 -0.194	0.992 1.786
NOAA-9	-	-	-	1.000 0.000	1.000 -0.243	0.989 2.500
NOAA-11	-	-	-	1.000 0.000	1.000 -0.178	0.990 2.184
NOAA-14	-	-	-	1.000 0.000	1.001 -0.427	0.996 0.945
NOAA-16	-	-	-	1.000 0.000	1.000 0.022	0.997 0.511
NOAA-18	-	-	-	1.000 0.000	1.000 -0.209	0.997 0.542
NOAA-19	-	-	-	1.000 0.000	1.000 0.000	1.000 0.000

Author contributions. MS coordinated the generation of the presented dataset, contributed to key developments and drafted the manuscript. SF prepared the AVHRR measurement record used as input. MS, SS and OS developed the cloud detection and phase determination and implemented the processing system used for the generation of the multi-decadal data set. CP and GM further developed the optimal estimation

system. MC implemented the BUGSrad scheme used for the calculation of the radiative flux properties. MS, SS, BW and DP evaluated the data. All authors contributed to finalizing the manuscript.

Competing interests. The authors declare that no competing interests are present.

Acknowledgements. This work was supported by the European Space Agency (ESA) through the Cloud_cci project (contract No.: 4000109870/13/I-
5 NB). The availability of the AVHRR GAC measurement record through the NOAA CLASS archive (<https://www.class.noaa.gov>) and the University of Wisconsin is much appreciated, with the latter also kindly providing corresponding intercalibration coefficients for the visible and near-infrared channels of AVHRR. Furthermore, the authors would like to thank Luca Lelli (University of Bremen) for providing SCIAMACHY data used to determine the spectral band adjustment.

References

- Allan, R. P.: Combining satellite data and models to estimate cloud radiative effect at the surface and in the atmosphere, *Meteorological Applications*, 18, 324–333, <https://doi.org/10.1002/met.285>, 2011.
- ATBD-CC4CL-BBFlux: ESA Cloud_cci Algorithm Theoretical Baseline Document (ATBD) of CC4CL Broadband Radiative Flux Retrieval, issue 1, rev. 1, 14 october 2019 edn., available from <http://www.esa-cloud-cci.org/?q=documentation>, 2019.
- Baran, A. J., Shcherbakov, V. N., Baker, B. A., Gayet, J. F., and Lawson, R. P.: On the scattering phase-function of non-symmetric ice-crystals, *Quarterly Journal of the Royal Meteorological Society*, 131, 2609–2616, <https://doi.org/10.1256/qj.04.137>, 2005.
- Baró, R., Jiménez-Guerrero, P., Stengel, M., Brunner, D., Curci, G., Forkel, R., Neal, L., Palacios-Peña, L., Savage, N., Schaap, M., Tuccella, P., Denier van der Gon, H., and Galmarini, S.: Evaluating cloud properties in an ensemble of regional online coupled models against satellite observations, *Atmospheric Chemistry and Physics*, 18, 15 183–15 199, <https://doi.org/10.5194/acp-18-15183-2018>, 2018.
- Baum, B. A., Yang, P., Heymsfield, A. J., Bansemer, A., Cole, B. H., Merrelli, A., Schmitt, C., and Wang, C.: Ice cloud single-scattering property models with the full phase matrix at wavelengths from 0.2 to 100 μ m, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 146, 123 – 139, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2014.02.029>, 2014.
- Christensen, M. W., Neubauer, D., Poulsen, C. A., Thomas, G. E., McGarragh, G. R., Povey, A. C., Proud, S. R., and Grainger, R. G.: Unveiling aerosol–cloud interactions – Part 1: Cloud contamination in satellite products enhances the aerosol indirect forcing estimate, *Atmospheric Chemistry and Physics*, 17, 13 151–13 164, <https://doi.org/10.5194/acp-17-13151-2017>, 2017.
- Delanoë, J. and Hogan, R. J.: A variational scheme for retrieving ice cloud properties from combined radar, lidar, and infrared radiometer, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD009000>, 2008.
- Delanoë, J. and Hogan, R. J.: Combined CloudSat-CALIPSO-MODIS retrievals of the properties of ice clouds, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD012346>, 2010.
- Devasthale, A., Raspaud, M., Schlundt, C., Hanschmann, T., Finkensieper, S., Dybbroe, A., Hörnquist, S., Håkansson, N., Stengel, M., and Karlsson, K.: PyGAC: An open-source, community-driven Python interface to preprocess more than 30-year AVHRR Global Area Coverage (GAC) data, *GSICS Quarterly Newsl.*, 11, <https://doi.org/10.7289/V5R78CFR>, 2017.
- Driemel, A., Augustine, J., Behrens, K., Colle, S., Cox, C., Cuevas-Agulló, E., Denn, F. M., Duprat, T., Fukuda, M., Grobe, H., Haeffelin, M., Hyett, N., Ijima, O., Kallis, A., Knap, W., Kustov, V., Long, C. N., Longenecker, D., Lupi, A., Maturilli, M., Mimouni, M., Ntsangwane, L., Ogihara, H., Olano, X., Olefs, M., Omori, M., Passamani, L., Pereira, E. B., Schmithüsen, H., Schumacher, S., Sieger, R., Tamlyn, J., Vogt, R., Vuilleumier, L., Xia, X., Ohmura, A., and König-Langlo, G.: Baseline Surface Radiation Network (BSRN): Structure and data description (1992–2017), *Earth System Science Data Discussions*, 2018, 1–17, <https://doi.org/10.5194/essd-2018-8>, 2018.
- Eliasson, S., Karlsson, K. G., van Meijgaard, E., Meirink, J. F., Stengel, M., and Willén, U.: The Cloud_cci simulator for the ESA Cloud_cci climate data record and its application to a global and a regional climate model, *Geoscientific Model Development Discussions*, 2018, 1–34, <https://doi.org/10.5194/gmd-2018-212>, 2018.
- Fu, Q. and Liou, K.: On the correlated k-distribution method for radiative transfer in nonhomogeneous atmospheres, *Journal of the Atmospheric Sciences*, 49, 2139–2156, 1992.
- Heidinger, A. K., Foster, M. J., Walther, A., and Zhao, X. T.: The Pathfinder Atmospheres–Extended AVHRR Climate Dataset, *Bulletin of the American Meteorological Society*, 95, 909–922, <https://doi.org/10.1175/BAMS-D-12-00246.1>, 2014.
- Heidke, P.: Berechnung des Erfolges und der Güte der Windstärkevorhersagen im Sturmwarnungsdienst, *Geografiska Annaler*, 8, 301–349, 1926.

- Karlsson, K.-G. and Johansson, E.: On the optimal method for evaluating cloud products from passive satellite imagery using CALIPSO-CALIP data: example investigating the CM SAF CLARA-A1 dataset, *Atmospheric Measurement Techniques*, 6, 1271–1286, <https://doi.org/10.5194/amt-6-1271-2013>, 2013.
- Karlsson, K.-G., Anttila, K., Trentmann, J., Stengel, M., Meirink, J. F., Devasthale, A., Hanschmann, T., Kothe, S., Jääskeläinen, E., Sedlar, J., Benas, N., van Zadelhoff, G.-J., Schlundt, C., Stein, D., Finkensieper, S., Håkansson, N., and Hollmann, R.: CLARA-A2: The second edition of the CM SAF cloud and radiation data record from 34 years of global AVHRR data, *Atmospheric Chemistry and Physics Discussions*, 2016, 1–41, <https://doi.org/10.5194/acp-2016-935>, 2016.
- Kato, S., Loeb, N. G., Rose, F. G., Doelling, D. R., Rutan, D. A., Caldwell, T. E., Yu, L., and Weller, R. A.: Surface Irradiances Consistent with CERES-Derived Top-of-Atmosphere Shortwave and Longwave Irradiances, *Journal of Climate*, 26, 2719–2740, <https://doi.org/10.1175/JCLI-D-12-00436.1>, 2013.
- Keller, M., Kröner, N., Fuhrer, O., Lüthi, D., Schmidli, J., Stengel, M., Stöckli, R., and Schär, C.: The sensitivity of Alpine summer convection to surrogate climate change: an intercomparison between convection-parameterizing and convection-resolving models, *Atmospheric Chemistry and Physics*, 18, 5253–5264, <https://doi.org/10.5194/acp-18-5253-2018>, 2018.
- Kratz, D. P., Gupta, S. K., Wilber, A. C., and Sothcott, V. E.: Validation of the CERES Edition 2B Surface-Only Flux Algorithms, *Journal of Applied Meteorology and Climatology*, 49, 164–180, <https://doi.org/10.1175/2009JAMC2246.1>, 2010.
- Lauer, A., Eyring, V., Righi, M., Buchwitz, M., Defourny, P., Evaldsson, M., Friedlingstein, P., de Jeu, R., de Leeuw, G., Loew, A., Merchant, C. J., Müller, B., Popp, T., Reuter, M., Sandven, S., Senfleben, D., Stengel, M., Roozendaal, M. V., Wenzel, S., and Willén, U.: Benchmarking CMIP5 models with a subset of ESA CCI Phase 2 data using the ESMValTool, *Remote Sensing of Environment*, 203, 9 – 39, <https://doi.org/https://doi.org/10.1016/j.rse.2017.01.007>, <http://www.sciencedirect.com/science/article/pii/S003442571730007X>, 2017.
- Loeb, N. G., Doelling, D. R., Wang, H., Su, W., Nguyen, C., Corbett, J. G., Liang, L., Mitrescu, C., Rose, F. G., and Kato, S.: Clouds and the Earth’s Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) Top-of-Atmosphere (TOA) Edition-4.0 Data Product, *Journal of Climate*, 31, 895–918, <https://doi.org/10.1175/JCLI-D-17-0208.1>, 2018.
- McGarragh, G. R., Poulsen, C. A., Thomas, G. E., Povey, A. C., Sus, O., Stapelberg, S., Schlundt, C., Proud, S., Christensen, M. W., Stengel, M., Hollmann, R., and Grainger, R. G.: The Community Cloud retrieval for CLimate (CC4CL) – Part 2: The optimal estimation approach, *Atmospheric Measurement Techniques*, 11, 3397–3431, <https://doi.org/10.5194/amt-11-3397-2018>, 2018.
- Nyeki, S., Wacker, S., Gröbner, J., Finsterle, W., and Wild, M.: Revising shortwave and longwave radiation archives in view of possible revisions of the WSG and WISG reference scales: methods and implications, *Atmospheric Measurement Techniques*, 10, 3057–3071, <https://doi.org/10.5194/amt-10-3057-2017>, 2017.
- Pavolonis, M. J. and Heidinger, A. K.: Daytime Cloud Overlap Detection from AVHRR and VIIRS, *Journal of Applied Meteorology*, 43, 762–778, <https://doi.org/10.1175/2099.1>, 2004.
- Pavolonis, M. J., Heidinger, A. K., and Uttal, T.: Daytime Global Cloud Typing from AVHRR and VIIRS: Algorithm Description, Validation, and Comparisons, *Journal of Applied Meteorology*, 44, 804–826, <https://doi.org/10.1175/JAM2236.1>, 2005.
- PVIR: ESA Cloud_cci Product Validation and Intercomparison Report (PVIR), issue 6, rev. y, day month 2019 edn., available from http://www.esa-cloud-cci.org/?q=documentation_v3, 2019.
- Rodgers, C. D.: Inverse methods for atmospheric sounding: theory and practice, vol. 2, World scientific, 2000.
- Rossow, W. B. and Schiffer, R. A.: Advances in Understanding Clouds from ISCCP, *Bulletin of the American Meteorological Society*, 80, 2261–2288, [https://doi.org/10.1175/1520-0477\(1999\)080<2261:AIUCFI>2.0.CO;2](https://doi.org/10.1175/1520-0477(1999)080<2261:AIUCFI>2.0.CO;2), 1999.

- Rutan, D. A., Kato, S., Doelling, D. R., Rose, F. G., Nguyen, L. T., Caldwell, T. E., and Loeb, N. G.: CERES Synoptic Product: Methodology and Validation of Surface Radiant Flux, *Journal of Atmospheric and Oceanic Technology*, 32, 1121–1143, <https://doi.org/10.1175/JTECH-D-14-00165.1>, 2015.
- Schiffer, R. A. and Rossow, W. B.: The International Satellite Cloud Climatology Project (ISCCP): The First Project of the World Climate Research Programme, *Bulletin of the American Meteorological Society*, 64, 779–784, <https://doi.org/10.1175/1520-0477-64.7.779>, 1983.
- Schlundt, C., Finkensieper, S., Raspaud, M., Karlsson, K.-G., Stengel, M., and Hollmann, R.: ESA Cloud_cci Technical Report on AVHRR GAC FCDR generation, Issue 4, Rev. 1; 01 May 2017, available from <http://www.esa-cloud-cci.org/?q=documentation>, 2017.
- Sohn, B. J., Nakajima, T., Satoh, M., and Jang, H.-S.: Impact of different definitions of clear-sky flux on the determination of longwave cloud radiative forcing: NICAM simulation results, *Atmospheric Chemistry and Physics*, 10, 11 641–11 646, <https://doi.org/10.5194/acp-10-11641-2010>, 2010.
- Stackhouse, P., Gupta, S., Cox, S., Zhang, T., Mikovitz, J., and Hinkelman, L.: 24.5-year SRB data set released, *GEWEX News*, 21, 10–12, 2011.
- Stengel, M., Mieruch, S., Jerg, M., K.-G, K., Scheirer, R., Maddux, B., Meirink, J. F., Poulsen, C., Siddans, R., Walther, A., and Hollmann, R.: The Clouds Climate Change Initiative: Assessment of state-of-the-art cloud property retrieval schemes applied to AVHRR heritage measurements, *Remote Sensing of Environment*, 162, <https://doi.org/10.1016/j.rse.2013.10.035>, 2013.
- Stengel, M., Stapelberg, S., Sus, O., Schlundt, C., Poulsen, C., Thomas, G., Christensen, M., Carbajal Henken, C., Preusker, R., Fischer, J., Devasthale, A., Willén, U., Karlsson, K.-G., McGarragh, G. R., Proud, S., Povey, A. C., Grainger, R. G., Meirink, J. F., Feofilov, A., Bennartz, R., Bojanowski, J. S., and Hollmann, R.: Cloud property datasets retrieved from AVHRR, MODIS, AATSR and MERIS in the framework of the Cloud_cci project, *Earth System Science Data*, 9, 881–904, <https://doi.org/10.5194/essd-9-881-2017>, 2017.
- Stengel, M., Schlundt, C., Stapelberg, S., Sus, O., Eliasson, S., Willén, U., and Meirink, J. F.: Comparing ERA-Interim clouds with satellite observations using a simplified satellite simulator, *Atmospheric Chemistry and Physics*, 18, 17 601–17 614, <https://doi.org/10.5194/acp-18-17601-2018>, 2018.
- Stengel, M., Sus, O., Stapelberg, S., Finkensieper, S., Würzler, B., Philipp, D., Hollmann, R., and Poulsen, C.: ESA Cloud Climate Change Initiative (ESA Cloud_cci) data: Cloud_cci AVHRR-PM L3C/L3U PRODUCTS v3.0, Deutscher Wetterdienst (DWD), https://doi.org/10.5676/DWD/ESA_Cloud_cci/AVHRR-PM/V003, 2019.
- Stephens, G. L.: Radiation Profiles in Extended Water Clouds. II: Parameterization Schemes, *Journal of the Atmospheric Sciences*, 35, 2123–2132, [https://doi.org/10.1175/1520-0469\(1978\)035<2123:RPIEWC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2123:RPIEWC>2.0.CO;2), 1978.
- Stephens, G. L., Gabriel, P. M., and Partain, P. T.: Parameterization of Atmospheric Radiative Transfer. Part I: Validity of Simple Models, *Journal of the Atmospheric Sciences*, 58, 3391–3409, [https://doi.org/10.1175/1520-0469\(2001\)058<3391:POARTP>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<3391:POARTP>2.0.CO;2), 2001.
- Stephens, G. L., Li, J., Wild, M., Clayson, C. A., Loeb, N., Kato, S., L’Ecuyer, T., Stackhouse Jr, P. W., Lebsock, M., and Andrews, T.: An update on Earth’s energy balance in light of the latest global observations, *Nature Geoscience*, 5, 691, <https://doi.org/10.1038/ngeo1580>, 2012.
- Sus, O., Stengel, M., Stapelberg, S., McGarragh, G., Poulsen, C., Povey, A. C., Schlundt, C., Thomas, G., Christensen, M., Proud, S., Jerg, M., Grainger, R., and Hollmann, R.: The Community Cloud retrieval for CLimate (CC4CL) – Part 1: A framework applied to multiple satellite imaging sensors, *Atmospheric Measurement Techniques*, 11, 3373–3396, <https://doi.org/10.5194/amt-11-3373-2018>, 2018.

Wentz, F. J. and Meissner, T.: AMSR-E/Aqua Daily L3 Global Ascending/Descending .25x.25 deg Ocean Grids, Version 2. [subset used: 2008/01/01-2008/12/31]. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center., https://doi.org/https://doi.org/10.5067/AMSR-E/AE_DYOCN.002 [data accessed: 2017/09/18], 2004.

Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the
5 CALIPSO Mission and CALIOP Data Processing Algorithms, *Journal of Atmospheric and Oceanic Technology*, 26, 2310–2323, <https://doi.org/10.1175/2009JTECHA1281.1>, 2009.

Young, A. H., Knapp, K. R., Inamdar, A., Hankins, W., and Rossow, W. B.: The International Satellite Cloud Climatology Project H-Series climate data record product, *Earth System Science Data*, 10, 583–593, <https://doi.org/10.5194/essd-10-583-2018>, 2018.