

*Modifications made in the minor revision process:*

We had to change the DOI reference since we noticed, soon after the first revision process, that some values regarding the zooplankton parameter were incorrect. Specifically, we compared the published version of the database with another, older database version and we found discrepancies maybe due to an erroneous manual transcription from one version to the other. We corrected the database for those particular zooplankton values, we then re-checked the entire database in order to have a definitive version of our data and we published it on Zenodo. The database has now a new DOI (<http://doi.org/10.5281/zenodo.3516717>), which is reported in this revised manuscript both in the Abstract and in the Data availability section, as required.

# A long term (1965-2015) ecological marine database from the LTER-Italy site Northern Adriatic Sea: plankton and oceanographic observations

Francesco Acri<sup>1</sup>, Mauro Bastianini<sup>1</sup>, Fabrizio Bernardi Aubry<sup>1</sup>, Elisa Camatti<sup>1</sup>, Alfredo Boldrin<sup>1</sup>, Caterina Bergami<sup>2</sup>, Daniele Cassin<sup>1</sup>, Amelia De Lazzari<sup>1</sup>, Stefania Finotto<sup>1</sup>, Annalisa Minelli<sup>3\*</sup>, Alessandro Oggioni<sup>4</sup>, Marco Pansera<sup>1</sup>, Alessandro Sarretta<sup>5</sup>, Giorgio Socal<sup>1</sup>, Alessandra Pugnetti<sup>1</sup>

<sup>1</sup> CNR-ISMAR, Arsenale - Tesa 104, Castello 2737/F, 30122 Venezia, Italy

<sup>2</sup> CNR-ISMAR, Via Gobetti 101, 40129 Bologna, Italy

<sup>3</sup> CNR-IRBIM, Largo Fiera della Pesca 2, 60125 Ancona, Italy

<sup>4</sup> CNR-IREA, Via Bassini 15, 20133 Milano, Italy

<sup>5</sup> CNR-IRPI, Corso Stati Uniti 4, 35127 Padova, Italy

Correspondence to: [annalisa.minelli@gmail.com](mailto:annalisa.minelli@gmail.com)

## Abstract

In this paper we describe a 50 years (1965-2015) ecological database containing data collected in the Northern Adriatic Sea (NAS), one of the 25 research parent sites belonging to the Italian Long Term Ecological Research Network (LTER-Italy, <http://www.lteritalia.it>). LTER-Italy is a formal member of the international (<https://www.ilter.network>) and European (<http://www.lter-europe.net/>) LTER networks. The NAS is undergoing a process, led by different research institutions and projects, for the establishment of a marine ecological observatory, building on the existing facilities, infrastructures, and long-term ecological data. Along this process, the implementation of the Open Access and Open Science principles has started, by creating an open research lifecycle that involves sharing ideas and results (scientific papers), data (raw and processed), metadata, methods, and software. The present data paper is framed within this wider context. The database is composed of observations on abiotic parameters, phyto- and zooplankton abundances, collected during 299 cruises in different sampling stations, in particular in the Gulf of Venice: we describe here the sampling and analytical activities, the parameters, and the structure of the database. The database is available at <http://doi.org/10.5281/zenodo.3516717> <http://doi.org/10.5281/zenodo.3465097> (Acri et al., 2019), it was also uploaded in the DEIMS-SDR repository (Dynamic Ecological Information Management System - Site and Dataset Registry, <https://deims.org/>), which is the official sites and data registry for LTER International network.

## 1. Introduction

We describe in this paper a 50 years (1965-2015) ecological database containing data on plankton communities and related abiotic parameters, collected in the Northern Adriatic Sea (NAS). Plankton communities, which are at the base of aquatic ecosystem functioning, have a broad and diversified range of seasonal patterns, multi-annual trends, and shifts across different marine ecosystems: making available long term series of plankton and oceanographic observations provides unique and precious tools for depicting reliable patterns of average annual cycles and for detecting significant changes and trends in response to global or local pressures and impacts.

Open Data is nowadays considered a crucial issue in both scientific research and public administration and management. Wilkinson et al. (2016) conceived the “FAIR” data management principles, which states that data must be “Findable, Accessible, Interoperable and Reusable”. The open access to data is one crucial step of Open Science (<http://www.budapestopenaccessinitiative.org/read>, European Commission, 2016), which is a wider approach embracing transparency at all stages of the research process, from research ideas to papers, open access to data, codes, and software. Open Science is actually a democratic way of making freely available, for every researcher and stakeholder, research ideas, data, metadata, tools, and outcomes. From the researcher point of view, open practices have been reported to give advantage, first of all, to open new frontiers in science (Science|Business network’s cloud consultation group, 2019) and provide solutions to urgent societal problems (Palen et al., 2015; Tai and Robinson, 2018); moreover, it allows gaining more citations, media

50 attention, potential collaborators, and funding opportunities (Eisenbach, 2006; McKiernan et al., 2016, Tennant et al., 2019)  
51 and it is vital for leaving a heritage to future generations.

52  
53 Ecology, being a multidisciplinary science, can surely benefit from the Open Science approach, which is, however, a matter  
54 of interest and discussion among ecologists only since the last decade (Reichman et al., 2011). Yet, the cultural shift from  
55 "data ownership to data stewardship" is not widely accomplished and data sharing standards, both from a technical and ethical  
56 point of view, have just started to be established (Hampton et al., 2015).

57 The Open Science approach is fostered in the data management plans of the Long Term Ecological Research (LTER) networks,  
58 at the national, European (LTER-Europe: <http://www.lter-europe.net/>) and global level (International LTER, ILTER:  
59 <https://www.ilter.network>), being considered a crucial step to advance socio-ecological research and education (Mirtl et al.,  
60 2018). ILTER provides a globally distributed network of long-term research sites for multiple purposes and uses in the fields  
61 of ecosystem, biodiversity, and socio-ecological research, it currently consists of 44 national networks, managing more than  
62 700 sites worldwide (Haase et al., 2018; Mirtl et al., 2018). LTER-Italy ([www.lteritalia.it](http://www.lteritalia.it)), a formal component of ILTER and  
63 LTER Europe since 2006, consists of 79 research sites, organized in 25 parent sites, which include terrestrial, freshwater,  
64 transitional and marine ecosystems, managed and coordinated by public research, monitoring Institutions and Universities  
65 (Bergami et al., 2019).


66 The LTER marine component, which represents around 10% of global ILTER sites, focuses mainly on ecosystem structure  
67 and function, in response to a wide range of environmental forcing factors, using long-term, site-based research. As a result of  
68 the wide range and of the exceptional rate and intensity of human impacts, the scientific value of long-term ecological  
69 observations is more crucial than ever for effective assessment, management, and prediction of the state and pressure in the  
70 marine environment. The creation and maintenance of marine ecological observatories, able to arrange and maintain integrated,  
71 harmonized and coherent long-term ecological observations, is actually stressed as a relevant step at the European level, for  
72 sustaining European marine policies (Benedetti-Cecchi et al., 2018; European marine Board 2019).

73 The marine component of LTER-Italy is made up of eight parent sites, mainly representing transitional and coastal ecosystems.  
74 Among them, the NAS is a significant geographical zone for the establishment of a marine ecological observatory, due to the  
75 concomitant presence of sensitive habitats, numerous ongoing monitoring, and research activities, as well as of heavy and  
76 diversified human pressures and economic interests. For these main reasons, during the years 2017-18, the Italian national  
77 flagship project RITMARE ("Italian research for the sea", <http://www.ritmare.it/>), funded by the Italian Ministry of University  
78 and Research, dedicated a Research Line to the establishment of a marine ecological observatory in the NAS. Building on the  
79 existing facilities, infrastructures and long-term ecological data, it aims at enhancing the marine observational capacities and  
80 at activating synergies among the main conservation management questions and key ecological and oceanographic variables.  
81 Along this process, it appeared crucial to start applying the Open Science principles, by creating an open research lifecycle,  
82 which foresees sharing each step of the process, from ideas and results (scientific papers) to data (raw and processed), from  
83 metadata to methods and software. The ideas and plans for the development of the Open Science principles to the NAS  
84 ecological observatory, which we named project "EcoNAOS" (Ecological Northern Adriatic Open Science Observatory  
85 System), are thoroughly described by Minelli et al. (2018).

86 This data paper represents one relevant step of this wider activity. The database that we present is composed of observations  
87 on abiotic (physical and chemical) parameters and phyto- and zooplankton abundances, collected in 50 years (from 1965 to  
88 2015), during cruises which interested different sampling stations across the NAS, in particular in the Gulf of Venice. Here we  
89 describe the sampling and analytical activities, the parameters, and the structure of the database.

90  
91 **2. The LTER-Italy parent site Northern Adriatic Sea**

92



Created on Inkatlas. © OpenStreetMap contributors (openstreetmap.org). Map data Oct 27, 2017. 1:2000000

Figure 1 - The LTER-Italy parent site Northern Adriatic Sea, with its four research sites. 1: Gulf of Trieste; 2: Gulf of Venice; 3: Po Delta and Romagna Coast; 4: Senigallia-Susak Transect. The fixed point observatories at each research site are evidenced (see Ravaioli et al., 2016 for a full description). Base map credits: © OpenStreetMap contributors 2019. Distributed under a Creative Commons BY-SA License.

The NAS (Figure 1) is the northernmost basin of the Mediterranean Sea and one of its most productive areas. It is characterized by a shallow depth and by a dominant cyclonic circulation. The oceanographic and meteorological parameters show a marked seasonal and interannual variability. The major forcings of the system are represented by the remarkable river inputs along the Italian coast, the Eastern Adriatic Current-EAC, which brings high salinity and oligotrophic waters from the southern basin, and the notable sea-level range, relatively to the rest of Mediterranean area. The urban and industrial inputs and the hydrodynamic exchange between the NAS and the lagoons located along the Italian coast are also elements of ecological relevance. A trophic gradient, decreasing from northwest to southeast, is typically observed in the basin, in which the nutrient-rich waters coming from the rivers are mainly spread southward and eastward from the Italian coast (Bernardi Aubry et al., 2006; Solidoro et al., 2009). The NAS is subject to multiple anthropogenic impacts (e.g., nutrient inputs, coastal urbanization, fishing activity, tourism, and maritime trade). The basin has undergone overfishing (Fortibuoni et al., 2010), marked eutrophication (during the 70s; Giani et al., 2012), followed by a phase of oligotrophication (years 2000s; Mozetič et al., 2010) and by a recent increase of nutrient concentrations (since 2007; Totti et al., 2019). The NAS has also been subjected to frequent development of mucilage aggregates (Giani et al., 2005; De Lazzari et al., 2008), until the first decade of the 2000s.

The LTER-Italy parent site NAS includes four research sites (Gulf of Trieste, Gulf of Venice, Po Delta and Romagna Coast, Senigallia-Susak Transect; Figure 1), where meteo-oceanographic and biological data, mainly on plankton (Table 2), are gathered both during oceanographic cruises and at fixed point observatories. Detailed information can be found in the ILTER Dynamic Ecological Information Management System Site and Dataset Registry, DEIMS-SDR (<https://deims.org/92fd6fad-99cd-4972-93bd-c491f0be1301>) (Wohner et al., 2019). The database we describe here refers to an area of about 40000 km<sup>2</sup>, ranging between 43.7° and 45.8° North and 12.2° and 14.3° East (coordinate reference system: WGS84).

### 3. Description of the database

122 The database described in this data paper is composed of 108687 records. Each record is intended as a timestamped and  
123 georeferenced set of information, individuated by a row in the database. These observations belong to 22 datasets coming from  
124 299 oceanographic cruises, carried out from 1965 to 2015.

125 Due to the long time coverage, the collection and analysis system for many parameters changed in time, thus making the  
126 database very heterogeneous for what concerns data management and organization. The heterogeneity is mainly due to:

- 127 • Sampling frequency: e.g., data coming from CTD (Conductivity, Temperature, Depth) sensors, such as temperature,  
128 oxygen, and pH, are registered in real-time at each meter in depth; other parameters, like nutrients and phytoplankton,  
129 are sampled at a lower time-frequency and at variable depths. The overall depth coverage ranged between 0-63 m,  
130 the sampling frequency from monthly to seasonal;
- 131 • Data treatment: some data are basically raw, e.g., data registered by CTD are reported into the database as they are  
132 delivered from the instrument; other data need some elaboration to obtain specific parameters' value (e.g., nutrients,  
133 chlorophyll-*a*, plankton abundance);
- 134 • Methodologies and units of measurements: e.g., changes of methodologies due to the introduction of CTD  
135 measurements; change of the units of measure of salinity, which passed from g l<sup>-1</sup> to a dimensionless parameter.
- 136 • Data format: data collected between 1965 and 1990 were registered only on paper archives, while those from 1990  
137 onwards on spreadsheets.

138 In particular, methodological protocols and associated documentation changed through time. Several sensors are described and  
139 extensively documented through the GET-IT platform (Geoinformation Enabling ToolKIT starterkit®, see Section 5), where  
140 it is possible to visualize all the observations related to a specific instrument or method. Other protocols have undergone a  
141 deep metadatation process by analyzing ancillary historical metadata (Scovacricchi, 2017). In this case, it is not immediately  
142 possible to obtain data related to a specific protocol, but it is still possible to filter data by method by importing the .csv file in  
143 a spreadsheet.


### 144 3.1 Data sources and geographical coverage

145 Data sources for this database come mainly from oceanographic cruises that were carried out on 12 different research vessels,  
146 at the basin scale (Table 1). The other observations come from sampling stations located next to the fixed automatic sensors:  
147 in this case the cruises are named as the nearby sensor, i.e.: 576 observations at the Paloma buoy (Gulf of Trieste), 1284 at the  
148 Acqua Alta oceanographic tower (Gulf of Venice), 138 at the S1 buoy (Po Delta). The data were gathered in the frame of  
149 many different projects that are all mentioned in the database:  
150  
151


Operation period	Research Vessel (R/V)	Nr. of observations
1965-1966	Vercelli	861
1966	Sea Quinn	60
1966-1990	Bannock	997
1968-2002	D'Ancona	45357
1977	Marsili	23
1979-1980	Mysis	48
1979-1990	Vila Vilebita	139
1986-1988	Minerva	737
2003	Boreana	2158
2003-2015	Dallaporta	43689
2007-2015	Litus	1900
2012-2014	Urania	12718

152 Table 1 – Operation periods of the different research vessels between 1965 and 2015 and number of observations.  
153  
154

155 Until the early 1990s, GPS systems were not usually on board of research vessels. For this reason, oceanographers used to  
 156 refer to a fixed grid covering the entire research area and identified the sampling positions (stations) with the nodes of this  
 157 grid. An example of grids used for this purpose is reported in Figure 2 (Franco, 1972).  
 158 In Figure 3a, the geographical coverage of the entire database is shown. Red dots represent the real observation points, while  
 159 the nodes of the grid are evidenced with black crosses. Observations referring to a specific station were assigned to the  
 160 coordinates of the corresponding node on the grid even if the real position was not precisely located on the grid node. This  
 161 resulted in a cloud of points in the nearby of each sampling station. Since our main aim was to preserve most of the information  
 162 for each observation, we decided not to “correct” the position of these points (see an example in Figure 3b for the station  
 163 09/0E).  
 164 In the following years, when the GPS allowed a better precision of the sampling position, researchers often continued referring  
 165 to the nodes of the grid for the station names and they adopted a nomenclature coherent with the one of the original grid also  
 166 for new sampling stations. For example, the new sampling point located eastward of the “09/2E” station is named “10/2E”,  
 167 since it is located at the same longitude (2E), but different latitude of “09/2E” station (Figure 3b). In Figure 3c, a 3D view of  
 168 the entire database is shown.  
 169 Due to transcription errors occurred during the oldest cruises, some data were misplaced, falling on land or outside the NAS.  
 170 A Python script (available under GNU GPL v.3 license here: <https://github.com/CNR-ISMAR/econaos/tree/master>) has been  
 171 written in order to correct this kind of errors. The same script implemented also a routine to homogenize different names of  
 172 the same sampling station (e.g. station “020D” could appear as well as station “02-0D” or “02/0D” or “020D\_07/07/1968”).  
 173 We selected the name reported on the original stations’ network grid (Figure 2) and we created from these stations a vector  
 174 layer (black crosses in Figure 3). Finally, since some stations changed their name through time, in order to maintain coherence  
 175 with the same sampling point, we appointed them with the last, most recently used name.  
 176



177  
 178  
 179 Figure 2 - An example of sampling stations based on the regular grid created in the NAS for the cruises from 1966 to 1980  
 180 (from Franco, 1972).  
 181



182  
183  
184  
185  
186  
187

Figure 3 - (A) Geographical distribution of the observations: red dots for observations; black crosses for nodes of the grid. (B) Example of cloud distribution of observations around sampling station 09/2E and the naming of new sampling station 10/2E. (C) 3D view of the database. Base map credits: © OpenStreetMap contributors 2019. Distributed under a Creative Commons BY-SA License.

188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200

### 3.2 Parameters: history, time coverage, and sensors

Samples collected during each cruise, whatever the station of collection, were then analyzed in the laboratory by means of diverse techniques. Since 2000 analytical quality of nutrients and chlorophyll analyses is assessed through participation to the Quality Assurance of Information for Marine Environmental Monitoring In Europe (QUASIMEME; <http://www.quasimeme.org>) international laboratory proficiency-testing. The complete list of the parameters of the database is reported in Table 2, together with some descriptive elements, i.e.:

- Total number of observations,
- Temporal coverage (from the first to the last record),
- Method or sensor currently used,
- Current unit of measure.

Parameter	Number of observations	Temporal coverage	Current sensor	Unit of measure	Acronym in the database
Transparency	2322	1965-2015	Secchi Disk	m	Secchi
Temperature	107648	1965-2015	CTD	C	Temp
Salinity	107655	1965-2015	CTD	dimensionless	Sal
Density anomaly	99961	1965-2015	Derived from temperature and salinity	kg m <sup>-3</sup>	Dens
pH	70376	1965-2011	pH glass membrane and pH electrode	pH unit	pH

Alkalinity	492	1965-2002	Titrimetric titration	meq l <sup>-1</sup>	Alky
Oxygen	12791	1965-2012	Oxygen Polarographic sensor	cc l <sup>-1</sup>	Oxyg (ml/l)
N-NH3	11154	1965-2015	Automated nutrient analysis	µm dm <sup>-3</sup>	NH3 (microMol)
N-NO2	11232	1965-2015	Automated nutrient analysis	µm dm <sup>-3</sup>	NO2 (microMol)
N-NO3	11299	1965-2015	Automated nutrient analysis	µm dm <sup>-3</sup>	NO3 (microMol)
P-PO4	11191	1965-2015	Automated nutrient analysis	µm dm <sup>-3</sup>	PO4 (microMol)
Si-SiO4	11420	1965-2015	Automated nutrient analysis	µm dm <sup>-3</sup>	Si (microMol)
Chlorophyll- <i>a</i>	11541	1965-2015	Spectrofluorimeter	µg l <sup>-1</sup>	Chla (ug/l)
Pheopigments	6352	1979-2015	Spectrofluorimeter	µg l <sup>-1</sup>	Pheo (ug/l)
Total Phytoplankton	3463	1977-2015	Inverted microscope	Cells l <sup>-1</sup>	Phyto TOT (cell/ml)
Diatoms	3070	1977-2015	Inverted microscope	Cells l <sup>-1</sup>	Diato (cell/ml)
Dinoflagellates	3070	1977-2015	Inverted microscope	Cells l <sup>-1</sup>	Dino (cell/ml)
Coccolithophores	3070	1977-2015	Inverted microscope	Cells l <sup>-1</sup>	Cocco (cell/ml)
Others	3070	1977-2015	Inverted microscope	Cells l <sup>-1</sup>	Flag (cell/ml)
Total Zooplankton	372	1987-2015	Stereo microscope	Ind. m <sup>-3</sup>	Zoo (ind/m <sup>3</sup> )

Table 2 - Database parameters and main descriptive information.

Instruments and sensors changed over the 50 year period, due to technological and scientific progress. Furthermore, instruments are also subject to degradation and need to be replaced. It is essential to preserve the information about these instrument changes and upgrading, to track the reliability of the measurements.

In order to appropriately document data and guarantee the consistency of data within the database, we collected most ancillary information as possible on the changes occurred in time for each parameter measurement. To this purpose, a thorough review of historical sources (e.g. logbooks and manual transcription in spreadsheets) was carried out (Scovacicchi, 2017), working in cooperation with some researchers - now retired - who participated to the first cruises and referring as well to papers by Franco (1970, 1972 and 1982), which describe methods and instruments during a number of oceanographic cruises in the NAS from 1965 to 1979.

Plankton data are particularly sensitive to the skill of the operators, in particular during the microscope analyses of the samples. The change of the operators, which necessarily occurred during 50 years, actually could hamper the data comparison across time. To deal with this issue, internal education and recurring calibration of taxonomic competence were carefully considered, with training periods and intercalibrations phases.

Since 2006 the taxonomic revision of the phytoplankton species has been made according to "Algaebase" ([www.algaebase.org](http://www.algaebase.org)), the global algal database of taxonomic, nomenclatural and distributional information for the zooplankton the Marine Planktonic Copepods catalog (<https://copepodes.obs-banyuls.fr/en/links.php>, Razolus et al., 2005-2019) has been used. In the past, for phyto- and zooplankton analyses several texts and monographs were used (Berard-



221 Therriault et al., 1999; Harris et al., 2000; Heimdal, 1993; Hendey, 1964; Hustedt, 1930-1966; Pascher, 1915; Peragallo and  
 222 Peragallo, 1897-1908; Rampi and Bernhardt, 1980; Schiller, 1931-37; Thronsdon, 1993; Tomas, 1997).  
 223 The phytoplankton was gathered and analyzed with the same method (Utermohl, 1958) across the years. In the database we  
 224 report the total phytoplankton abundances and the following main groups: diatoms, dinoflagellates (naked and armoured cells),  
 225 coccolithophorids and “others”, which include the sum of cells belonging to cryptophyceans, crysophyceans,  
 226 prymnesiophyceans (except coccolithophorids), prasinophyceans and chlorophyceans, whose sizes lie between 4 and 20 µm  
 227 and often remain undetermined. Mesozooplankton was always identified under a stereo-microscope and expressed as the total  
 228 number of organisms per cubic meter. Compared to phytoplankton, the mesozooplankton data are much fragmented over time:  
 229 they cover a 28 year period, from 1987 to 2015, for a total of 372 observations.

#### 230 4. Database structure and analysis

231 The present version of the database is recorded in a unique spreadsheet (Figure 4), carrying information, for each record, about:

- 232 • Coordinates (longitude-latitude) of the sampling station;
- 233 • Sampling depth;
- 234 • Sampling station name;
- 235 • Cruise and R/V (Ship) name;
- 236 • Sampling date and time;
- 237 • Water column depth (Bot. Depth);
- 238 • Instrument/method used for each measurement and relative parameter value.


239

Long	Lat	Depth	Station	Cruise	Ship	YYYY-MM-DD	hh:mm:ss	Bot. Depth	Temp_sensor	Temp	Sal_sensor	Sal
12.68	45.33	0.5B	PP/1	VERCELLI	1965-04-12	9:33:00	23	Tilting thermometer	13.12	Morh Knudsen titration	29.61	
12.68	45.33	5B	PP/1	VERCELLI	1965-04-12	9:33:00	23	Tilting thermometer	12.35	Morh Knudsen titration	35.66	
12.68	45.33	10B	PP/1	VERCELLI	1965-04-12	9:33:00	23	Tilting thermometer	12.45	Morh Knudsen titration	35.43	
12.68	45.33	20B	PP/1	VERCELLI	1965-04-12	9:33:00	23	Tilting thermometer	12.14	Morh Knudsen titration	38.01	
12.86	45.28	0.5C	PP/1	VERCELLI	1965-04-12	12:20:00	29	Tilting thermometer	12.25	Morh Knudsen titration	35.44	
12.86	45.28	5C	PP/1	VERCELLI	1965-04-12	12:20:00	29	Tilting thermometer	12.24	Morh Knudsen titration	35.46	
12.86	45.28	10C	PP/1	VERCELLI	1965-04-12	12:20:00	29	Tilting thermometer	11.16	Morh Knudsen titration	37.79	
12.86	45.28	20C	PP/1	VERCELLI	1965-04-12	12:20:00	29	Tilting thermometer	12.3	Morh Knudsen titration	37.92	
12.48	45.40	0.5A	PP/2	VERCELLI	1965-04-28	6:42:00	16.4	Tilting thermometer	12.27	Morh Knudsen titration	33.04	
12.48	45.40	1A	PP/2	VERCELLI	1965-04-28	6:42:00	16.4	Tilting thermometer	12.37	Morh Knudsen titration	33.39	
12.48	45.40	5A	PP/2	VERCELLI	1965-04-28	6:42:00	16.4	Tilting thermometer	12.44	Morh Knudsen titration	35.39	
12.48	45.40	10A	PP/2	VERCELLI	1965-04-28	6:42:00	16.4	Tilting thermometer	12.23	Morh Knudsen titration	37.3	
12.68	45.33	0.5B	PP/2	VERCELLI	1965-04-28	9:10:00	22.3	Tilting thermometer	12.49	Morh Knudsen titration	32.9	
12.68	45.33	5B	PP/2	VERCELLI	1965-04-28	9:10:00	22.3	Tilting thermometer	12.43	Morh Knudsen titration	33.78	
12.68	45.33	10B	PP/2	VERCELLI	1965-04-28	9:10:00	22.3	Tilting thermometer	11.92	Morh Knudsen titration	37.21	
12.68	45.33	20B	PP/2	VERCELLI	1965-04-28	9:10:00	22.3	Tilting thermometer	10.5	Morh Knudsen titration	37.72	
12.86	45.28	0.5C	PP/2	VERCELLI	1965-04-28	11:20:00	31	Tilting thermometer	12.4	Morh Knudsen titration	34.2	
12.86	45.28	5C	PP/2	VERCELLI	1965-04-28	11:20:00	31	Tilting thermometer	12.09	Morh Knudsen titration	36.15	
12.86	45.28	8C	PP/2	VERCELLI	1965-04-28	11:20:00	31	Tilting thermometer	11.5	Morh Knudsen titration	37.38	
12.86	45.28	20C	PP/2	VERCELLI	1965-04-28	11:20:00	31	Tilting thermometer	10.42	Morh Knudsen titration	37.9	
12.48	45.40	0.5A	PP/3	VERCELLI	1965-05-13	6:47:00	16	Tilting thermometer	15.92	Morh Knudsen titration	33.66	
12.48	45.40	1A	PP/3	VERCELLI	1965-05-13	6:47:00	16	Tilting thermometer	15.8	Morh Knudsen titration	33.77	
12.48	45.40	5A	PP/3	VERCELLI	1965-05-13	6:47:00	16	Tilting thermometer	14.92	Morh Knudsen titration	33.51	
12.48	45.40	10A	PP/3	VERCELLI	1965-05-13	6:47:00	16	Tilting thermometer	11.34	Morh Knudsen titration	37.61	
12.68	45.33	0.5B	PP/3	VERCELLI	1965-05-13	9:23:00	21	Tilting thermometer	17.4	Morh Knudsen titration	33.84	
12.68	45.33	5B	PP/3	VERCELLI	1965-05-13	9:23:00	21	Tilting thermometer	15.66	Morh Knudsen titration	36.2	
12.68	45.33	10B	PP/3	VERCELLI	1965-05-13	9:23:00	21	Tilting thermometer	13.64	Morh Knudsen titration	37.3	
12.68	45.33	20B	PP/3	VERCELLI	1965-05-13	9:23:00	21	Tilting thermometer	11.83	Morh Knudsen titration	37.72	
12.86	45.28	0.5C	PP/3	VERCELLI	1965-05-13	12:15:00	31	Tilting thermometer	18.03	Morh Knudsen titration	33.01	

240  
 241  
 242 Figure 4 - An example of the database showing the fields for each observation.

243  
 244 Around 89% of the observations of the database refers to the years 1999-2015 and the remaining 11% covers the previous 33  
 245 years (see Figure 5a for details). This is mainly due to the adoption of CTD probes since 1999 for measuring abiotic parameters  
 246 at each meter depth, leading to an imbalance between the observations before (e.g. 778 in 1978) and after 1999 (e.g. 11359 in  
 247 2004). In Figure 5b observations from oceanographic cruises onboard of the different research vessels are shown (see also Table  
 248 1). The number of observations on abiotic parameters (nutrients, alkalinity, and transparency) is higher than the biotic  
 249 (chlorophyll-*a*, phytoplankton, and zooplankton abundances) ones up to the year 2000; since then, they become comparable  
 250 (Figure 5c).


251 The database presents a heterogeneous number of observations for each parameter, mainly due to: (i) parameter priority for  
 252 the specific research conducted, (ii) the instruments and analytical efforts required, and (iii) the specific funding programs and  
 253 resources.




(A)

Ship

BANNOCK	DALLAPORTA	MINERVA	URANIA
BOREANA	LITUS	MISYS	VERCELLI
D'ANCONA	MARSILI	SEA QUINN	VILA VILEBITA




(B)




(C)

254  
255  
256  
257


Figure 5 - Total number of observations over the whole period (A), clustered by research vessel (B), and by biotic (orange line) and abiotic (blue line) parameters (C).



(A)



(B)



(C)

Figure 6 - Distribution over the years of the temperature (A), chlorophyll-*a* (B) and phytoplankton (C) observations

259  
260  
261  
262

263 In Figure 6 we compare the total number of observations of one physical (temperature) and two biological (chlorophyll-*a* and  
264 phytoplankton abundance) parameters. All the three parameters were measured each year, although with different frequency.  
265 Temperature attains up to ~15000 records, while chlorophyll-*a* ~1200 records at most and phytoplankton ~300. The number  
266 of temperature data has a temporal distribution similar to the general one described in Figure 5a, where 89% of the observations  
267 occurred in the last 17 years, due to the adoption of CTD probes. Chlorophyll-*a* observations show instead peaks during the  
268 years 1987-1990, due to intense regional monitoring activities occurring in those years. The lowest number of phytoplankton  
269 observations is mainly due to the complex and time-consuming analytical procedure, which do not allow processing too many  
270 samples, and to the reduction of extensive monitoring activities since 2006.

## 271 5. Data visualization

272 The data management activities of the national flagship project RITMARE (Fugazza et al. 2014) allowed to develop two tools  
273 to enhance the deployment of a distributed Spatial Data Infrastructure (SDI) for Italian marine researchers community. SDI is  
274 an interoperable technological infrastructure for preservation, publication, and discovery of geospatial, modeled on standard  
275 (Open Geospatial Consortium - OGC, World Wide Web Consortium - W3C, and INSPIRE Directive 2007/2/EC) web services.  
276 In order to strengthen the RITMARE infrastructure, the Open Source software suite GET-IT (Oggioni et al., 2017; Menegon  
277 et al., 2017) and the customizable, template-driven metadata editor [EDI](#) (Pavesi et al. 2016; Tagliolato et al. 2016;  
278 [https://github.com/SP7-Ritmare/EDI-NG\\_client](https://github.com/SP7-Ritmare/EDI-NG_client)) have been developed and released as Open Source code. One of the nodes of  
279 the distributed SDI provides geospatial data collected by CNR-ISMAR marine researchers (<http://vesk.ve.ismar.cnr.it>).  
280 Following the OGC Sensor Web Enablement (SWE) web service, each instrument or procedure has to be filled out as a  
281 “sensor”, then observations can be provided, for a specific parameter, as OGC O&M (Observations and Measurements) web  
282 standards. Through the EDI interface, integrated within GET-IT software suite, a first core of sensors was already tested and  
283 uploaded in 2015 (Bastianini et al. 2015). A number of buoys (e.g. [ABATE - Seabird SBE 19 Plus V2](#)), laboratory instruments  
284 (e.g. [Spettrophotometer Perkin Elmer](#)), methods (e.g. [Titration Winkler](#)) and sensors, have been described for this study by  
285 using XML SensorML v2.0 language and their metadata, including manufacturer (provided as RDF, which stays for Resource  
286 Description Framework, Friends Of A Friends FOAF in Oggioni, 2019), owner and operator contacts, measured parameters,  
287 position, documentation, and history, can be easily visualized in separate dedicated landing pages (Figure 7). Currently, in the  
288 CNR-ISMAR GET-IT data node, 35 sensors have been described (<http://vesk.ve.ismar.cnr.it/sensors/>), for which it is possible  
289 to upload observations, collected from different stations in the NAS.  
290

# CONTROS HydroC™ (Piattaforma Acqua Alta - CONTROS HydroC)

The CONTROS HydroC™ CO2 sensor is a unique underwater dioxide sensor for in-situ and online measurements of dissolved CO2. The versatile HydroC™ CO2 is suitable for platform installations (e.g. ROV s or AUV s), long-term deployments (e.g. buoys and moorings) as well as for profiling applications (e.g. water sampling rosettes). Fields of application include: ocean acidification research, climate studies, air-sea gas exchange, limnology, fresh water control, aquaculture/fish farming, carbon capture and storage – monitoring, measurement and verification (CCS-MMV).

## Manufacturer Name

**CONTROS**

tel:+49-431-2609-5900

Wischhofstrasse 1-3 Geb.  
Kiel - Seefischmarkt  
D-24148  
DE

mailto:contact@contros.eu

http://www.contros.eu

## Model Number

HydroC

## Parameters

[Partial\\_pressure\\_of\\_carbon\\_dioxide\\_{pCO2}\\_in\\_the\\_atmosphere\\_by\\_infra-red\\_gas\\_analysis](#)

Unit of measure: uatm

Partial pressure of carbon dioxide {pCO2} in the atmosphere by infra-red gas analysis

## Position



## Contact

### Owner

**Istituzione Centro Previsioni  
Segnalazioni Maree**

+39 041 27 48 787

S.Marco 4090  
Venezia  
30124  
Italia

maree@comune.venezia.it



http://www.comune.venezia.it/maree/

### Operator

**Istituzione Centro Previsioni  
Segnalazioni Maree**

+39 041 27 48 787

S.Marco 4090  
Venezia  
30124  
Italia

maree@comune.venezia.it



http://www.comune.venezia.it/maree/

## Documentation

Documentazione Stazione dal sito ufficiale

application/pdf

## History

Today

### Updates

2015-02-23

Metadata creation


### Installation

2014-05-01

Date of installation

291  
292  
293  
294  
295

Figure 7 - Example of the sensor description provided by GET-IT. Information about manufacturer, owner and operator contacts, measured parameters, position, documentation, and history are displayed. Base map credits: © OpenStreetMap contributors 2019. Distributed under a Creative Commons BY-SA License.



296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311

Figure 8 - Example of the graph in output from a query into the database. Oxygen data at station C10 for the period 01/01/1992-01/01/1996 are displayed.

Since v1.3.17 GET-IT still does not allow the three-dimensional representation of data, we decided to upload into the software suite only surface values of each parameter and sampling operation. This part of the database can be queried and graphed, directly into GET-IT using developed tool, in order to showtime series of selected parameters (Figure 8). A total of 16017 observations have been uploaded.

Observations can upload using the graphical interface or, for the skilled people, using an XML language directly into SOS (Sensor Observation Service) web service. For the upload from the interface, data have to be formatted in a table with datetime and parameter value (Figure 9). Since the speed of the process largely depends on the browser used to upload data, most of the data have been uploaded, through a Python script, by formatting specific .xml files, containing information about the sensor's ID, sampling station, and date time and following SWE standard. In both cases, the data upload begins with the selection of the sensor we want to upload data from and, then, with the selection of the sampling station from the map, if already available, or by creating a new one.

Name	Sampling point			Sampled Feature (URI)	Action
	Lat.	Lon.	SRS		
PTF	45.3085	12.505	http://www.opengis.net/def/crs/EPSSG/0/4326	Adriatic Sea	Use
2E	44.73667	12.6305	http://www.opengis.net/def/crs/EPSSG/0/4326	Adriatic Sea	Use
C10	45.2507	12.7602	http://www.opengis.net/def/crs/EPSSG/0/4326	Adriatic Sea	Use

(A)

phenomenonTime	Temperature ITS-90_of
2013-10-13T15:56:22	18.680
2013-10-13T16:54:27	18.770
2014-02-24T10:23:55	12.260
2014-02-24T13:00:55	11.260
2014-02-24T15:28:55	11.020
2014-09-28T05:09:39	21.170
2014-09-28T06:06:43	21.210
2014-09-28T07:17:51	21.310
2014-09-28T08:05:10	21.510
2014-09-28T08:57:55	21.720

save data    reset table

Legend - Fields definition

phenomenonTime (Date and Time)  
 Temperature ITS-90\_of\_the\_water\_body\_by\_CTD\_or\_STD (Quantity) Unit of measure: degC

(B)

312 Figure 9 - Data upload from the graphic interface. Selection of the sampling station for the specific sensor (A) and format of  
 313 data to be uploaded into the SDI (B).  
 314

315 **6. Data availability**  
 316

317 The dataset is available at <http://doi.org/10.5281/zenodo.3516717><http://doi.org/10.5281/zenodo.3465097> (Aciri et al., 2019). It  
 318 was also uploaded in the DEIMS-SDR repository (Dynamic Ecological Information Management System - Site and Dataset  
 319 Registry, <https://deims.org/dataset/38d604ef-decb-4d67-8ac3-cc843d10d3ef>), which is the official sites and data registry for  
 320 LTER International network. The aim of DEIMS-SDR is to be a catalogue of in-situ observation or experimentation facilities;  
 321 it is implemented as a web-based information portal for integrated ecological information which comprises detailed  
 322 descriptions of sites where research is carried out, including the technical infrastructure, ecosystem properties and research  
 323 activities (see Wohner et al., 2019 for a full description). DEIMS-SDR provides a service which allows to associate a PID  
 324 (Persistent Identifier) to the uploaded dataset. Thanks to an agreement between the eLTER Research Infrastructure and the  
 325 EUDAT Collaborative Data Infrastructure (CDI), the dataset is automatically available also in the B2Share catalogue

326  
327  
328  
329  
330  
331

(<https://b2share.fz-juelich.de/records/e8d57102fd194bde957407ca290ad06a>) and, through this, in the EOSC (European Open Science Cloud) and GEOSS (Global Earth Observation System of Systems) catalogues. Since we opted for CC-BY license our data are immediately fully available for download and reuse upon citation, without embargo rules or any further limitations. Table 3 collects the list of columns, short name and extended name of each parameter and ancillary field composing the database.

Column number	Parameter short name (database)	Parameter extended name
1	Long	Longitude [decimal degrees]
1	Long	Longitude [decimal degrees]
2	Lat	Latitude [decimal degrees]
3	Depth	Depth [m]
4	Station	Name of sampled station
5	Station_updated_name	Updated name of sampling station, if present
6	Cruise	Cruise
7	Ship	Ship
8	YYYY-MM-DD	Date
9	hh:mm:ss	Time
10	Bot. Depth [m]	Water column depth [m]
11	Secchi [m]	Transparency [m]
12	Temp_sensor	Temperature sensor
13	Temp	Temperature [°C]
14	Sal_sensor	Salinity sensor
15	Sal	Salinity [dimensionless]
16	Dens	Density Anomaly [ $\text{kg m}^{-3}$ ]
17	pH_sensor	pH sensor
18	pH	pH [pH Units]
19	Oxyg_sensor	Oxygen sensor
20	Oxyg (ml/l)	Dissolved Oxygen concentration [ $\text{ml l}^{-1}$ ]
21	Ox%	Dissolved Oxygen saturation [%]
22	NH3_sensor	Ammonia sensor
23	NH3 (microMol)	Ammonia [ $\mu\text{m dm}^{-3}$ ]
24	NO2_sensor	Nitrite sensor
25	NO2 (microMol)	Nitrite [ $\mu\text{m dm}^{-3}$ ]



26	NO3_sensor	Nitrate sensor
27	NO3 (microMol)	Nitrate [ $\mu\text{m dm}^{-3}$ ]
28	Din (microMol)	Total Dissolved inorganic Nitrogen [ $\mu\text{m dm}^{-3}$ ]
29	PO4_sensor	Phosphate sensor
30	PO4 (microMol)	Phosphate [ $\mu\text{m dm}^{-3}$ ]
31	Si_sensor	Silicate sensor
32	Si (microMol)	Silicate [ $\mu\text{m dm}^{-3}$ ]
33	Chla_sensor	Chlorophyll- <i>a</i> sensor
34	Chla (ug/l)	Chlorophyll- <i>a</i> concentration [ $\mu\text{g l}^{-1}$ ]
35	Pheo (ug/l)	Phaeopigments concentration [ $\mu\text{g l}^{-1}$ ]
36	Alky	Alkalinity
37	Diato (cell/ml)	Diatoms abundance [ $\text{cell ml}^{-1}$ ]
38	Dino (cell/ml)	Dinoflagellates abundance [ $\text{cell ml}^{-1}$ ]
39	Flag (cell/ml)	Other Flagellates abundance [ $\text{cell ml}^{-1}$ ]
40	Cocco (cell/ml)	coccolithophorids abundance [ $\text{cell ml}^{-1}$ ]
41	Phyto TOT (cell/ml)	Total phytoplankton abundance [ $\text{cell ml}^{-1}$ ]
42	Zoo (ind/m <sup>3</sup> )	Total mesozooplankton organisms [ $\text{ind m}^{-3}$ ]

332  
333  
334  
335  
336

Table 3 - Correspondence between column number, short name and extended name of each parameter reported into the database.

## 7. Conclusions

337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356

The 50-year database of plankton and abiotic parameters in the NAS may contribute to an in-depth comprehension of plankton dynamics required not only to manage aquatic resources but also to predict and tackle future environmental changes. Long-term site-based studies on plankton may provide an invaluable opportunity to assess common or contrasting patterns of variability, to understand how those patterns change at different scales and to hypothesize about causes and consequences. Wide availability of the data on long-term variations of the planktonic system allows large scale studies that obviously go beyond the local use, representing a source of information for cross-system analysis, allowing comparison among ecosystems as well as new approaches in data analysis and in the development of water quality indicators. However, these potential uses appear constrained by issues that are intrinsic to long-term series and that are related to the obvious variations, across the years, of sampling coverage and frequency and of analytical methodologies. In this respect, it is crucial to appropriately document the data, collecting and making available most ancillary information as possible on the changes occurred in time for each parameter measurement. This process was thoroughly carried for the 50 years NAS dataset so that the potential users might know which could be the proper application and the limitations of the dataset. The open access to the 50-year dataset of abiotic data and plankton in the NAS was framed in a wider open science life-cycle approach undertaken in the EcoNAOS project (Minelli et al., 2018), with the purpose to develop a practical case study which could root the high and inspiring principles of Open Science into the scientific community, fostering as well a cultural shift. In EcoNAOS we involved, since its start, both LTER and data management researchers in a joint partnership. In particular, the elaboration of the 50-year datasets has been worked out by a small group of plankton ecologists and data management experts, with the aim of sharing and harmonizing as well the different experiences, needs and points of view. This participatory process is recognized to be crucial to contribute overcoming cultural differences, barriers and fragmentation that might represent an obstacle for Open Science (Björk, 2004; Janssen et al., 2012; Barry & Bannister, 2014). The constant interactions of

357 oceanographers and ecologists with experts on data management and analysis, geospatial standards and web services  
358 interoperability, creating a rich and multi-domain research group, has been necessary to make available and understandable  
359 the very detailed knowledge behind environmental surveys, samplings, analyses, methodologies, through the sound and fit-  
360 for-purpose technical solutions for data management and interoperability.  
361 Accessibility and interoperability concepts and practices are crucial elements for LTER networks because the more the time  
362 series are consistent, coherent and available, the more it is possible to reconstruct trends and dynamics and to identify and  
363 compare reliable trends. The consistency and the coherence of the dataset require careful efforts in supplying the proper  
364 metadata, which could document the methodological changes that occurred through the years, thus allowing the potential users  
365 to evaluate the restrictions as well as the most suitable uses of the dataset. The activity described in this data paper is fully in  
366 line with the data management plan of the LTER networks, at the national, European and global level, since one of the LTER  
367 mandates is actually to foster open sharing of LTER data (Mirtl 2010; Mirtl et al., 2018). The national LTER networks are  
368 fostered to adopt the aspects of open science that are currently feasible in the different research groups.  
369 Currently, a dynamic update and integration of the published dataset is not yet supported by specific tools nor integrated in  
370 automatic procedures; anyway, it is foreseen to go on with the promotion of a full open science approach to LTER also in the  
371 coming years and extend the dataset through the publication of updates and possibly through the integration of different long-  
372 term datasets.

### 373 **Author contribution**

374 Minelli A. led the whole process of dataset recovery, metadatation and publication. Oggioni A., Pugnetti A. and Sarretta A.  
375 were the reference persons for the entire activity. Acri F., Bastianini M., Bernardi Aubry F., Camatti E., Bergami C., Minelli  
376 A., Oggioni A., Sarretta A., Pugnetti A. contributed to the writing of this paper: drafting of the text and of the figures and  
377 tables, revisions and suggestions.  
378 Acri F., Bastianini M., Bernardi Aubry F., Camatti E., Boldrin A., Cassin D., De Lazzari A., Pansera M., Finotto S., Socal G.  
379 collected and analysed data over time, providing statistics and material (graphs and tables) for the paper.

### 380 **Competing interests**

381 Authors declare that there are no competing interests that might have influenced the performance or presentation of the work  
382 described in this manuscript.

### 383 **References**

- 384 Acri, F., Bastianini, M., Bernardi Aubry, F., Camatti, E., Bergami, C., Boldrin, A., De Lazzari, A., Finotto, S., Minelli, A.,  
385 Oggioni, A., Pansera, M., Sarretta, A., Socal, G. and Pugnetti, A.: LTER Northern Adriatic Sea (Italy) marine data from 1965  
386 to 2015 [Data set], Zenodo, <http://doi.org/10.5281/zenodo.3266246>, 2019.
- 387  
388 Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J., Bonino da  
389 Silva Santos, L., Bourne, P.E., Bouwman, J., Brookes, A.J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo,  
390 C.T., Finkers, R., Gonzalez-Beltran, A., Gray, A.J.G., Groth, P., Goble, C., Grethe, J.S., Heringa, J., 't Hoen, P.A.C., Hooft,  
391 R., Kuhn, T., Kok, R., Kok, J., Lusher, S.J., Martone, M.E., Mons, A., Packer, A.L., Persson, B., Rocca-Serra, P., Roos, M.,  
392 van Schaik, R., Sansone, S., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M.A., Thompson, M., van der Lei, J.,  
393 van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J. and Mons, B.: The FAIR Guiding  
394 Principles for scientific data management and stewardship, *Scientific data*, 3 <http://dx.doi.org/10.1038/sdata.2016.18>, 2016.
- 395  
396 European Commission: Open innovation, open science, open to the world - a vision for Europe, [RTD-  
397 PUBLICATIONS@ec.europa.eu](https://doi.org/10.2777/061652) [ISBN 978-92-79-57346-0], <https://doi.org/10.2777/061652>, 2016.
- 398  
399 Science|Business Network's Cloud Consultation Group: Why Open Science is the Future (and how to make it happen).  
400 Science|Business. Brussels. Report available here: [https://sciencebusiness.net/report/why-open-science-future-and-how-  
401 make-it-happen](https://sciencebusiness.net/report/why-open-science-future-and-how-make-it-happen), 2019.
- 402  
403 Palen, L., Soden, R., Anderson, T. J., and Barrenechea, M.: Success & scale in a data-producing organization: The socio-  
404 technical evolution of OpenStreetMap in response to humanitarian events. In Proceedings of the 33rd annual ACM conference  
405 on human factors in computing systems (pp. 4113-4122). ACM, <http://doi.org/10.1145/2702123.2702294>, 2015.
- 406

407 Tai, T., and Robinson, J.: Enhancing climate change research with open science. *Frontiers in Environmental Science*, 6, 115,  
408 <https://doi.org/10.3389/fenvs.2018.00115>, 2018.

409  
410 Eysenbach, G.: Citation advantage of open access articles. *PLoS biology*, 4(5), e157,  
411 <http://doi.org/10.1371/journal.pbio.0040157>, 2006.

412  
413 McKiernan, E. C., Bourne, P. E., Brown, C. T., Buck, S., Kenall, A., Lin, J., McDougall, D., Nosek, B.A., Ram, K., Soderberg,  
414 C.K., Spies, J.R., Thaney, K., Updegrave, A., Woo, K.H. and Yarkoni, T.: Point of view: How open science helps researchers  
415 succeed, *eLife* 2016;5:e16800, <https://doi.org/10.7554/eLife.16800>, 2016.

416  
417 Tennant, J. P., Crane, H., Crick, T., Davila, J., Enkhbayar, A., Havemann, J., Kramer, B., Martin, R., Masuzzo, P., Nobes, A.,  
418 Rice, C., Rivera-López, B. S., Ross-Hellauer, T., Sattler, S., Thacker, P. and Vanholsbeeck M.: Ten myths around open  
419 scholarly publishing. *PeerJ Preprints* 7:e27580v1, <http://doi.org/10.7287/peerj.preprints.27580v1>, 2019.

420  
421 Reichman, O. J., Jones, M. B., and Schildhauer, M. P.: Challenges and opportunities of open data in ecology, *Science*,  
422 331(6018), 703-705, <http://doi.org/10.1126/science.1197962>, 2011.

423  
424 Hampton, S. E., S. S. Anderson, S. C. Bagby, C. Gries, X. Han, E. M. Hart, M. B. Jones, W. C. Lenhardt, A. MacDonald, W.  
425 K. Michener, J. Mudge, A. Pourmokhtarian, M. P. Schildhauer, K. H. Woo, and N. Zimmerman: The Tao of open science for  
426 ecology, *Ecosphere* 6(7):120, <http://dx.doi.org/10.1890/ES14-00402.1>, 2015.

427  
428 Mirtl, M., Borer, E. T., Djukic, I., Forsius, M., Haubold, H., Hugo, W., Orenstein, D. E., Pauw, J.C., Peterseil, J., Shibata, H.,  
429 Wohner, C., Yu, X. and Haase, P.: Genesis, goals and achievements of long-term ecological research at the global scale: a  
430 critical review of ILTER and future directions, *Science of the Total Environment*, 626, 1439-1462,  
431 <https://doi.org/10.1016/j.scitotenv.2017.12.001>, 2018.

432  
433 Haase, P., Tonkin, J. D., Stoll, S., Burkhard, B., Frenzel, M., Geijzendorffer, I. R., Mirtl, M., Müller, F., Musche, M., Penner,  
434 J., Zacharias, S. and Schmeller, D.S.: The next generation of site-based long-term ecological monitoring: linking essential  
435 biodiversity variables and ecosystem integrity, *Science of the Total Environment*, 613, 1376-1384,  
436 <https://doi.org/10.1016/j.scitotenv.2017.08.111>, 2018.

437  
438 Bergami C., L' Astorina A. and Pugnetti A. (a cura di): *I Cammini della Rete LTER-Italia, Il racconto dell'ecologia in cammino*,  
439 Roma: CNR Edizioni. ISBN (online) 978888080304-1, ISBN (cartaceo) 978888080312-6,  
440 <http://doi.org/10.32018/978888080304-1>, 2018.

441  
442 Benedetti-Cecchi, L., Crowe, T., Boehme, L., Boero, F., Christensen, A., Grémare, A., Hernandez, F., Kromkamp, J. C.,  
443 Nogueira García, E., Petihakis, G., Robidart, J., Sousa Pinto, I. and Zingone, A.: Strengthening Europe's Capability in  
444 Biological Ocean Observations, Muñiz Piniella, Á., Kellett, P., Larkin, K., Heymans, J. J. [Eds.] *Future Science Brief 3 of the*  
445 *European Marine Board*, Ostend, Belgium. 76 pp. ISBN: 9789492043559 ISSN: 2593-5232, 2018.

446  
447 European Marine Board: *Navigating the Future V: Marine Science for a Sustainable Future*, Position Paper 24 of the European  
448 Marine Board, Ostend, Belgium, ISBN: 9789492043757, ISSN: 0167-9309  
449 <http://doi.org/10.5281/zenodo.2809392>, 2019.

450  
451 Minelli, A., Oggioni, A., Pugnetti, A., Sarretta, A., Bastianini, M., Bergami, C., Bernardi Aubry, F., Camatti, E., Scovacricchi,  
452 T. and Socal, G.: The project EcoNAOS: vision and practice towards an open approach in the Northern Adriatic Sea ecological  
453 observatory, *Research Ideas and Outcomes* 4: e24224, <https://doi.org/10.3897/rio.4.e24224>, 2018.

454  
455 Bernardi Aubry, F., Aciri, F., Bastianini, M., Pugnetti, A., and Socal, G.: Picophytoplankton contribution to phytoplankton  
456 community structure in the Gulf of Venice (NW Adriatic Sea), *International Review of Hydrobiology*, 91(1), 51-70,  
457 <https://doi.org/10.1002/iroh.200410787>, 2006.

458  
459 Solidoro, C., Bastianini, M., Bandelj, V., Codermatz, R., Cossarini, G., Melaku Canu, D., Ravagnan, E., Salon, S. and  
460 Trevisani, S.: Current state, scales of variability, and trends of biogeochemical properties in the northern Adriatic Sea, *J.*  
461 *Geophys. Res.*, 114 (2009), p. C07S91, <https://doi.org/10.1029/2008JC004838>, 2009.

462

463 Fortibuoni, T., Libralato, S., Raicevich, S., Giovanardi, O. and Solidoro, C.: Coding Early Naturalists' Accounts into Long-  
464 Term Fish Community Changes in the Adriatic Sea (1800–2000), *PLoS ONE* 5(11): e15502,  
465 <https://doi.org/10.1371/journal.pone.0015502>, 2010.  
466

467 Giani, M., Djakovac, T., Degobbis, D., Cozzi, S., Solidoro, C. and Fonda Umani, S.: Recent changes in the marine ecosystems  
468 of the northern Adriatic Sea, *Estuarine, Coastal, and Shelf Science*, Volume 115, 2012, Pages 1-13, ISSN 0272-7714,  
469 <https://doi.org/10.1016/j.ecss.2012.08.023>, 2012.  
470

471 Mozetič, P., Solidoro, C., Cossarini, G., Socal, G., Precali, R., Francé, J., Bianchi, F., De Vittor, C., Smoldaka, N. and Umani,  
472 S. F.: Recent trends towards oligotrophication of the northern Adriatic: evidence from chlorophyll a time series, *Estuaries and  
473 coasts*, 33(2), 362-375, <https://doi.org/10.1007/s12237-009-9191-7>, 2010.  
474

475 Totti, C., Romagnoli, T., Accoroni, S., Coluccelli, A., Pellegrini, M., Campanelli, A., Grilli, F. and Marini, M.: Phytoplankton  
476 communities in the northwestern Adriatic Sea: Interdecadal variability over a 30-years period (1988–2016) and relationships  
477 with meteorological drivers, *Journal of Marine Systems*, 193: 137-153, <https://doi.org/10.1016/j.jmarsys.2019.01.007>, 2019.  
478

479 Giani, M., Berto, D., Zangrando, V., Castelli, S., Sist, P., and Urbani, R.: Chemical characterization of different typologies of  
480 mucilaginous aggregates in the Northern Adriatic Sea, *Science of the Total Environment*, 353(1-3), 232-246,  
481 <https://doi.org/10.1016/j.scitotenv.2005.09.027>, 2005.  
482

483 De Lazzari, A., Berto, D., Cassin, D., Boldrin, A., and Giani, M.: Influence of winds and oceanographic conditions on the  
484 mucilage aggregation in the Northern Adriatic Sea in 2003–2006, *Marine Ecology*, 29(4), 469-482,  
485 <https://doi.org/10.1111/j.1439-0485.2008.00268.x>, 2008.  
486

487 Ravaioli, M., Bergami, C., Riminucci, F., Langone, L., Cardin, V., Di Sarra, A., Aracri, S., Bastianini, M., Bensi, M.,  
488 Bergamasco, A., Bommarito, C., Borghini, M., Bortoluzzi, G., Bozzano, R., Cantoni, C., Chiggiato, J., Crisafi, E., D'Adamo,  
489 R., Durante, S., Fanara, C., Grilli, F., Lipizer, M., Marini, M., Miserocchi, S., Paschini, E., Penna, P., Pensieri, S., Pugnetti,  
490 A., Raichich, F., Schroeder, K., Siena, G., Specchiulli, A., Stanghellini, G., Vetrano, A. and Crise, A.: The RITMARE Italian  
491 Fixed-Point Observatory Network (IFON) for marine environmental monitoring: a case study, *Journal of Operational  
492 Oceanography*, 9(sup1), s202-s214, <https://doi.org/10.1080/1755876X.2015.1114806>, 2016.  
493

494 Moedas, C.: H2020 Programme Guidelines on FAIR Data Management in Horizon 2020,  
495 [http://ec.europa.eu/research/participants/data/ref/h2020/grants\\_manual/hi/oa\\_pilot/h2020-hi-oa-data-mgt\\_en.pdf](http://ec.europa.eu/research/participants/data/ref/h2020/grants_manual/hi/oa_pilot/h2020-hi-oa-data-mgt_en.pdf), 2016.  
496

497 Council, B.: Freedom of Information Act 2000, Education, 18, 21. UK Government,  
498 <https://www.legislation.gov.uk/ukpga/2000/36/contents>, 2006.  
499

500 Swan A.: The Open Access citation advantage: Studies and results to date, Technical Report, School of Electronics &  
501 Computer Science, University of Southampton, [https://eprints.soton.ac.uk/268516/2/Citation\\_advantage\\_paper.pdf](https://eprints.soton.ac.uk/268516/2/Citation_advantage_paper.pdf), 2010.  
502

503 Bernius, S., and Hanauske, M.: Open access to scientific literature-increasing citations as an incentive for authors to make their  
504 publications freely accessible, In *System Sciences*, 2009 42nd Hawaii International Conference on (pp. 1-9),  
505 <http://doi.org/10.1109/HICSS.2009.335>, IEEE, 2009.  
506

507 Björk, B. C.: Open access to scientific publications - an analysis of the barriers to change?, *Information Research*, 2004, Vol.9,  
508 No.2, paper 170, <http://hdl.handle.net/10227/647>, 2004.  
509

510 Janssen, M., Charalabidis, Y., and Zuiderwijk, A.: Benefits, adoption barriers and myths of open data and open government,  
511 *Information systems management*, 29(4), 258-268, <https://doi.org/10.1080/10580530.2012.716740>, 2012.  
512

513 Barry, E., and Bannister, F.: Barriers to open data release: A view from the top, *Information Polity*, 19(1, 2), 129-152,  
514 [https://www.scss.tcd.ie/disciplines/information\\_systems/egpa/docs/2013/BarryBannister.pdf](https://www.scss.tcd.ie/disciplines/information_systems/egpa/docs/2013/BarryBannister.pdf), 2014.  
515

516 Picone, M., Orasi, A. and Nardone, G.: Sea Surface Temperature monitoring in Italian Seas: Analysis of long-term trends and  
517 short-term dynamics, *Measurement*, 129, pages 260-267, ISSN 0263-2241,  
518 <https://doi.org/10.1016/j.measurement.2018.07.033>, 2018.

520 Driemel, A., Fahrbach, E., Rohardt, G., Beszczynska-Möller, A., Boetius, A., Budéus, G., Cisewski, B., Engbrodt, R., Gauger,  
521 S., Geibert, W., Geprägs, P., Gerdes, D., Gersonde, R., Gordon, A. L., Grobe, H., Hellmer, H. H., Isla, E., Jacobs, S. S., Janout,  
522 M., Jokat, W., Klages, M., Kuhn, G., Meincke, J., Ober, S., Østerhus, S., Peterson, R. G., Rabe, B., Rudels, B., Schauer, U.,  
523 Schröder, M., Schumacher, S., Sieger, R., Sildam, J., Soltwedel, T., Stangeew, E., Stein, M., Strass, V. H., Thiede, J.,  
524 Tippenhauer, S., Veth, C., von Appen, W., Weirig, M., Wisotzki, A., Wolf-Gladrow, D. A. and Kanzow T.: From pole to pole:  
525 33 years of physical oceanography onboard R/V Polarstern, *Earth System Science Data*, 9(1), 211-220,  
526 <https://doi.org/10.5194/essd-9-211-2017>, 2017.

527

528 Morris, D., Pinnegar, J., Maxwell, D., Dye, S., Fernand, L., Flatman, S., Williams, O. J., and Rogers, S.: Over 10 million  
529 seawater temperature records for the United Kingdom Continental Shelf between 1880 and 2014 from 17 Cefas (United  
530 Kingdom government) marine data systems, *Earth System Science Data*, 10, 27-51, <https://doi.org/10.5194/essd-10-27-2018>,  
531 2018.

532

533 Nohe, A., Knockaert, C., Goffin, A., Dewitte, E., De Cauwer, K., Desmit, X., Vyverman, W., Tyberghein, L., Lagring, R. and  
534 Sabbe, K.: Marine phytoplankton community composition data from the Belgian part of the North Sea, 1968–2010, *Scientific*  
535 *data*, 5, <http://dx.doi.org/10.1038/sdata.2018.126>, 2018.

536

537 Davies, C. H., Ajani, P., Armbrrecht, L., Atkins, N., Baird, M. E., Beard, J., Bonham, P., Burford, M., Clementson, L., Coad,  
538 P., Crawford, C., Dela-Cruz, J., Doblin, M. A., Edgar, S., Eriksen, R., Everett, J. D., Furnas, M., Harrison, D. P., Hassler, C.,  
539 Henschke, N., Hoenner, X., Ingleton, T., Jameson, I., Keesing, J., Leterme, S. C., McLaughlin, M. J., Miller, M., Moffatt, D.,  
540 Moss, A., Nayar, S., Patten, N. L., Patten, R., Pausina, S. A., Proctor, R., Raes, E., Robb, M., Rothlisberg, P., Saeck, E. A.,  
541 Scanes, P., Suthers, I. M., Swadling, K. M., Talbot, S., Thompson, P., Thomson, P. G., Uribe-Palomino, J., van Ruth, P., Waite,  
542 A. M., Wright, S. and Richardson, A. J.: A database of chlorophyll a in Australian waters, *Scientific data*, 5, 180018,  
543 <http://dx.doi.org/10.1038/sdata.2018.18>, 2018.

544

545 Sparnocchia S., Ravaioli M. and Focaccia P.: La rete osservativa fissa del CNR, Internal Technical Report. CNR-ISMAR,  
546 <http://www.ismar.cnr.it/area-riservata/documenti/La%20rete%20osservativa%20fissa%20del%20CNR.pdf>, 2010.

547

548 Scovaccicchi, T.: Technical recovery of material and methods applied in the cruises of 1965, 1966, 1978/79 as described by P.  
549 Franco (1970, 1972, 1982), Internal Report, CNR-ISMAR Venezia (Italy), 2017.

550

551 Franco, P.: Oceanography of Northern Adriatic-Sea: 1. Hydrologic features-cruises July-August and October-November 1965,  
552 *Archivio di Oceanografia e Limnologia*, 16, 1, 1970.

553

554 Franco, P.: Oceanography of Northern Adriatic Sea: 2. Hydrologic features: cruises January-February and April-May 1966,  
555 *Archivio di Oceanografia e Limnologia*, 17 (suppl.), P 1-97. 1972. Illuss, 1972.

556

557 Franco, P.: Oceanography of Northern Adriatic Sea: Data from the cruises of the years 1978 and 1979, Istituto di biologia del  
558 mare, 1982.

559

560 Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N.: Diversity and Geographic Distribution of Marine Planktonic  
561 Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>, 2005-2019.

562

563 Berard-Therriault L., Poulin M. and Bossé L.: Guide d'identification du phytoplancton marin de l'estuaire et du golfe du Saint-  
564 Laurent. NRC Research Press, 387 pp., ISBN: 978-0-660-96057-9, 1999.

565

566 Harris, R.P., Wiebe, P.H., Lenz, J., Skjoldal, H.R. and M. Huntley.: ICES Zooplankton Methodology Manual, Academic Press,  
567 USA. pp. 684, <https://doi.org/10.1016/B978-0-12-327645-2.X5000-2>, 2000.

568

569 Heimdal B. R.: Modern Coccolithophorids in: Marine phytoplankton a guide to naked flagellates and coccolithophorids. Tanos  
570 editors, Academic Press: 147- 248., ISBN: 9780323138277, 1993.

571

572 Hendey, N. I.: An introductory account of the smaller algae of British coastal waters. Part V: Bacillariophyceae, Diatoms.  
573 *Fishery Invest. Lond. Ser. IV* 5, 317 pp., 1964.

574

575 Hustedt F.: Die Kiesealgen von Deutschland, Österreichs und der Schweiz mit Berücksichtigung der übrigen Länder Europas  
576 sowie der angrenzender Mehresgebiete. In : Rabenhorst's Kriptogamen-Flora von Deutschland, Österreichs und der Schweiz.

577 Akad; Verlag. m. b. H. Leipzig. 7 : Tl. 2. 920 pp. : Tl., 2 845 pp. ; Tl. 3, 816 pp., <https://doi.org/10.5962/bhl.title.1356>, 1930-  
578 1966.

579

580 Pascher A.: Clorophyceae. In: Die Susswasser Flora Deutschlands, Osterreichs und der Schweiz. Verlags von Gustav Fisher,  
581 Jena, Heft 5, 250 pp., ISBN: 1176110802, 1915.

582

583 Peragallo H. and Peragallo M.: Diatomees Marine de France et des Districts Maritimes Voisins. Micrographe Editeur Grez sur  
584 Loing (S. et M.), 419 pp., ISBN: 0270995986, 1897-1908.

585

586 Rampi L. and Bernhardt M.: Chiave per la determinazione tassonomica delle Peridinee Pelagiche Mediterranee: C.N.E.N.,  
587 Roma (RT/B10 (81)13): 1-98, 1980.

588

589 Schiller J: Dinoflagellatae (Peridineae) Monografischer Behandlung. In : Rabenhorst Kriptogamen-Flora von Deutschland,  
590 Osterreichs und der Schweiz. Verlag. m. b. H. Leipzig. 10 (3) -1, 1-617, (1931-1933), (10) 3-2, 1-590, (1933-1937),  
591 <https://doi.org/10.5962/bhl.title.1356>, 1931-1937.

592

593 Sournia A.: Atlas du phytoplancton marin. Editions du Centre National de la recherche Scientifique. (1), 1-219, (2) 1-297,  
594 ISBN-10: 2222038235, 1993.

595

596 Thronsdon J.: The planktonic marine flagellates in: Marine phytoplankton a guide to naked flagellates and coccolithophorids.  
597 Tanos editors, Academic Press: 7- 131, ISBN: 9780323138277, 1993.

598

599 Tomas, C. R.: Identifying Marine Phytoplankton. Academic Press, Arcourt Brace & Company, [https://doi.org/10.1016/B978-  
600 0-12-693018-4.X5000-9](https://doi.org/10.1016/B978-0-12-693018-4.X5000-9), 1997.

601

602 Utermöhl, H.: Zur Vervollkommung der quantitativen Phytoplankton-Methodik, Mitt. Int. Ver. Limnol., 9, 38 p., 1958.

603

604 Van Landingham, J.W.: A modification of the Knudsen method for salinity determination, J. Cons. Perm. Int. Explor. Mer 22,  
605 174-179, 1956.

606

607 Williams, J.: A small portable salinometer, Chesapeake Bay Inst. Tech. Rep. 23, 1961.

608

609 Anderson, D.H. and Robinson, R.J.: Rapid electrometric determination of the alkalinity of sea water using a glass electrode,  
610 Ind. Engng Chem. (analyt Edn) 18, 767-769, 1946.

611

612 Buch, K.: Das Kohlensaure Gleichgewichtssystem im Meerwasser, Merentutkimuslait. Julk. 151, 9-18, 1951.

613

614 Winkler, L. W.: Die bestimmung des im wasser gelösten sauerstoffes, Berichte der deutschen chemischen Gesellschaft, 21(2),  
615 2843-2854, 1988.

616

617 Cox, R.A.: International oceanographic tables, UNESCO, 1966.

618

619 Gieskes, J.M.: Effect of temperature on the pH of seawater, Limnol. Oceanogr. 14, 679-685, 1969.

620

621 Van Slyke, D. D.: Eine Methode zur quantitativen Bestimmung der aliphatischen Aminogruppen; einige Anwendungen  
622 derselben in der Chemi der Proteine, des Harns und der Enzyme, Berichte der Deutschen Chemischen Gesellschaft, 43 : 3170-  
623 3181, 1910.

624

625 Solorzano, L.: Determination of ammonia in natural waters by the phenolhypochlorite method, Limnol. Oceanogr. 14, 799-  
626 801, 1969

627

628 Shinn, M. B.: A colorimetric method for the determination of nitrite, Ind. Engng Chem., analyt. Edn, Vol. 13, pp. 33-5, 1941.

629

630 Bendschneider, K. and Robinson, R.J.: A new spectrophotometric method for the determination of nitrate in sea water, J. mar.  
631 Res. 11, 87-96, 1952.

632

633 Wood, E. D., Armstrong, F. A. J. and Richards, F. A.: Determination of nitrate in sea water by cadmium-copper reduction to  
634 nitrite, J. mar. boil. Ass. U.K. 47, 23-37, 1967.

635

636 Fossato, V. U.: Determinazione dei fosfati reattivi a basse concentrazioni nell'acqua di mare, Arch. Limnol. Oceanogr. 16, 95-  
637 97, 1968.

638

639 Murphy, J. and Riley, J. P.: A modified single solution method for the determination of phosphate in natural waters, Analyt.  
640 Chim. Acta 27, 31-36, 1962.

641

642 Mullin, J. B. and Riley, J. P.: The spectrophotometric determination of nitrate in natural waters with particular reference to sea  
643 water, Anal. Chim. Acta, 12: 464-480, 1955.

644

645 Fugazza, C., Basoni, A., Menegon, S., Oggioni, A., Pavesi, F., Pepe, M., Sarretta, A. and Carrara, P.: RITMARE: Semantics-  
646 aware Harmonisation of Data in Italian Marine Research, Procedia Computer Science 33: 261-265,  
647 <https://doi.org/10.1016/j.procs.2014.06.041>, 2014.

648

649 Oggioni, A., Tagliolato, P., Fugazza, C., Pepe, M., Menegon, S., Pavesi, F. and Carrara, P.: Interoperability in Marine Sensor  
650 Networks through SWE Services: The RITMARE Experience, <https://doi.org/10.4018/978-1-5225-0700-0.ch009>, 2017.

651

652 Menegon, G., Vianello, A., Oggioni, A., Tagliolato, P., Lanucara, S., Sarretta, A., Basoni, A., Zilioli, M., Pepe, M., Fugazza,  
653 C., Bastianini, M., Criscuolo, L., Pavesi, F. and Carrara, P.: SP7-Ritmare/starterkit v1.3.17,  
654 <https://doi.org/10.5281/zenodo.1101020>, 2017.

655

656 Pavesi, F., Basoni, A., Fugazza, C., Menegon, S., Oggioni, A., Pepe, M., Tagliolato, P. and Carrara, P.: EDI – A Template-  
657 Driven Metadata Editor for Research Data, Journal of Open Research Software 4 (1) <https://doi.org/10.5334/jors.106>, 2016.

658

659 Tagliolato, P., Oggioni, A., Fugazza, C., Pepe, M., and Carrara, P.: Sensor metadata blueprints and computer-aided editing for  
660 disciplined SensorML, IOP Conference Series: Earth and Environmental Science 34 (1): 012036-012036,  
661 <https://doi.org/10.1088/1755-1315/34/1/012036>, <http://doi.org/10.1088/1755-1315/34/1/012036>, 2016.

662

663 Oggioni, A.: oggioniale/RDF-FOAF-Manufacture-list: First release of RDF-FOAF-Manufacturers list,  
664 <https://doi.org/10.5281/zenodo.3247546>, 2019.

665

666 Bastianini, M., Bernardi Aubry, F., Bianchi, F., Boldrin, A., Camatti, E., Carrara, P., De Lazzari, A., Guerzoni, S., Menegon,  
667 S., Oggioni, A., Pugnetti, A., Sarretta, A., Socal, G., Tagliolato, P. and Vianello, A.: The LTER site Gulf of Venice and the  
668 project RITMARE: a case study for the recovery, search, view and sharing of long term ecological marine research data, AIOL  
669 - Associazione Italiana Oceanografia e Limnologia, Verbania Pallanza, 2015.

670

671 Vollenweider, R. A., Giovanardi, F., Montanari, G., and Rinaldi, A.: Characterization of the trophic conditions of marine  
672 coastal waters with special reference to the NW Adriatic Sea: proposal for a trophic scale, turbidity and generalized water  
673 quality index, Environmetrics: The official journal of the International Environmetrics Society, 9(3), 329-357,  
674 [http://doi.org/10.1002/\(SICI\)1099-095X\(199805/06\)9:3<329::AID-ENV308>3.0.CO;2-9](http://doi.org/10.1002/(SICI)1099-095X(199805/06)9:3<329::AID-ENV308>3.0.CO;2-9), 1998.

675

676 De Mauro, A., Greco, M., and Grimaldi, M.: A formal definition of Big Data based on its essential features, Library Review,  
677 65(3), 122-135, <https://doi.org/10.1108/LR-06-2015-0061>, 2016.

678

679 Pollock, D. and Michael, A.: News & Views: evaluating quality in Open Access journals, <https://deltathink.com/news-views-evaluating-quality-in-open-access-journals/>, 2018.

680

681

682 Chatterjee, R., Arun, G., Agarwal, S., Speckhard, B. and Vasudevan, R.: Using data versioning in database application  
683 development, In Software Engineering, 2004 ICSE Proceedings. 26th International Conference on (pp. 315-325), IEEE,  
684 <http://doi.org/10.1109/ICSE.2004.1317454>, 2004.

685

686 Hansell, D. A.: Dissolved Organic Matter, nutrients and CTD data collected from the R/V Oceanus cruise OC404-01 and R/V  
687 Weatherbird II cruise WB0409 from the Sargasso Sea from June 11, 2004 to July 01, 2004 (EDDIES project), [https://hdl.  
688 handle.net/1912/4848](https://hdl.handle.net/1912/4848), 2011.

689

690 Stadler, A., Nagel, C., König, G. and Kolbe, T. H.: Making interoperability persistent: A 3D geo database based on CityGML,  
691 In 3D Geo-information sciences (pp. 175-192), Springer, Berlin, Heidelberg, [https://doi.org/10.1007/978-3-540-87395-2\\_11](https://doi.org/10.1007/978-3-540-87395-2_11),  
692 2009.

693

694 McGavra, G., Morris, S. and Janée, G.: Technology Watch Report: Preserving Geospatial Data, Technical Report. Digital  
695 Preservation Coalition, 2009.

696

697 Chaturvedi, K., Smyth, C. S., Gesquière, G., Kutzner, T. and Kolbe, T. H.: Managing versions and history within semantic 3D  
698 city models for the next generation of CityGML, In Advances in 3D Geoinformation (pp. 191-206), Springer, Cham,  
699 [http://doi.org/10.1007/978-3-319-25691-7\\_11](http://doi.org/10.1007/978-3-319-25691-7_11), 2017.

700

701 Shaon, A. and Woolf, A.: Long-term preservation for spatial data infrastructures: A metadata framework and geo-portal  
702 implementation, D-Lib Magazine, 17(9/10), 1-14, <http://doi.org/10.1045/september2011-shaon>, 2011.

703

704 Berners-Lee, T., Fielding, R. and Masinter, L.: Uniform resource identifier (URI): Generic syntax (No. RFC 3986),  
705 <http://doi.org/10.17487/RFC2396>, 2004.

706

707 Halpin, P. N., Read, A. J., Best, B. D., Hyrenbach, K. D., Fujioka, E., Coyne, M. S., Crowder, L. B., Freeman, S. A. and  
708 Spoerri, C.: OBIS-SEAMAP: developing a biogeographic research data commons for the ecological studies of marine  
709 mammals, seabirds, and sea turtles, Marine Ecology Progress Series, 316, 239-246, <http://doi.org/10.3354/meps316239>, 2006.

710

711 Wohner, C., Peterseil, J., Poursanidis, D., Klimont, T., Wilson, M., Mirtl, M., and Nektarios, C.: DEIMS-SDR – A web portal  
712 to document research sites and their associated data, Ecological Informatics, <https://doi.org/10.1016/j.ecoinf.2019.01.005>,  
713 2019.