

Dear Editor,

we thank you for your (and referees') time spent helping us to improve the manuscript. We particularly appreciate the effort in this period of pandemic closures when running even standard business has become complicated for all.

Please find uploaded the new version, with all remarks by the Editor and Referee 2 implemented. In particular, we reopened a ticket on PANGAEA asking for the modifications requested by the Editor.

- (i) Column headers in CNR_DIN_WMED_20042017_original.csv and CNR_DIN_WMED_20042017_adjusted.csv are now the very same
- (ii) We uploaded a new readme file with more information and in particular with links to the reports and the cruise PIs along with explanation of the abbreviations. Abbreviations are explained accordingly in the supplementary material section A2 "Data product organization details".

Please be aware that as of today (16/06/2020), the ticket to PANGAEA is not yet resolved and we are waiting for PANGAEA feedbacks for final implementation.

We also implemented all minor edits requested. Please note that we added in the manuscript a new table, listing data sources and links to the reports (accessed June 2020). This new table has been numbered Table 1b. Accordingly, the old table 1 is now table 1a.

1 **Dissolved Inorganic Nutrients in the Western Mediterranean Sea (2004-2017)**

2 Malek Belgacem^{1,2}, Jacopo Chiggiato^{1,*}, Mireno Borghini¹, Bruno Pavoni², Gabriella Cerrati³,
3 Francesco Acri¹, Stefano Cozzi⁴, Alberto Ribotti⁵, Marta Álvarez⁶, Siv K. Lauvset⁷, Katrin Schroeder¹

4 ¹ CNR-ISMAR, Arsenale Tesa 104, Castello 2737/F, 30122 Venezia, Italy

5 ² Dipartimento di Scienze Ambientali Informatica e Statistica, Università Ca' Foscari Venezia,
6 Campus Scientifico Mestre, Italy

7 ³ ENEA, Department of Sustainability, S. Teresa, Marine Environmental center, 19032 Pozzuolo di
8 Lerici (SP), Italy

9 ⁴ CNR-ISMAR, Area Science Park – Basovizza, 34149 Trieste, Italy

10 ⁵ CNR-IAS, Loc. Sa Mardini snc, Torregrande, 9170 Oristano, Italy

11 ⁶ Instituto Español de Oceanografía, IEO, A Coruña, Spain

12 ⁷ NORCE Norwegian Research Centre, Bjerknes Centre for Climate Research, 5007 Bergen, Norway

13 *Corresponding author's email: jacopo.chiggiato@ismar.cnr.it

15 **Abstract**

16 Long-term time-series are a fundamental prerequisite to understand and detect climate shifts and
17 trends. Understanding the complex interplay of changing ocean variables and the biological
18 implication for marine ecosystems requires extensive data collection for monitoring, hypothesis testing
19 and validation of modelling products. In marginal seas, such as the Mediterranean Sea, there are still
20 monitoring gaps, both in time and in space. To contribute to filling these gaps, an extensive dataset of
21 dissolved inorganic nutrients observations (nitrate, NO_3^- ; phosphate, PO_4^{3-} ; and silicate, SiO_2) ~~have~~ has
22 been collected between 2004 and 2017 in the Western Mediterranean Sea and subjected to rigorous
23 quality control techniques to provide to the scientific community a publicly available, long-term,
24 quality controlled, internally consistent biogeochemical data product. The data product includes 870
25 stations of dissolved inorganic nutrients, including temperature and salinity, sampled during 24
26 cruises. Details of the quality control (primary and secondary quality control) applied are reported.

Formattato: Non Evidenziato

27 The data are available in PANGAEA (<https://doi.pangaea.de/10.1594/PANGAEA.904172>, Belgacem
28 et al. 2019)

29 **Keywords:** Mediterranean Sea, Dissolved Inorganic Nutrient, biogeochemistry.
30

31 **1 Introduction**

32 Dissolved inorganic nutrients play a crucial role in marine ecosystem functioning. They serve as
33 regulators of ocean biological productivity, and are trace elements for biogeochemical cycling as well
34 as for natural and anthropogenic sources and transport processes (Bethoux, 1989; Bethoux et al.,
35 1992). They are also non-conservative tracers, since their distribution vary according to both
36 biological (such as primary production and respiration) and physical (such as convection, advection,
37 mixing and diffusion) processes. Very schematically, inorganic nutrients are continuously consumed
38 by phytoplankton (due to primary production) in the sea surface and regenerated in the mesopelagic
39 layer by bacteria and animals (due to respiration). Moreover, the sinking of organic matter and its
40 decomposition increases the nutrient concentrations in the intermediate and deep-water masses over
41 time. To identify the limiting factors for biological production in the oceans, we need to understand
42 the underlying chemical constraints and especially the macro- and micronutrients spatial and temporal
43 variations. Dissolved inorganic nutrients may be used as tracers of water masses like salinity and
44 temperature, to assess mixing processes, and to understand the biogeochemical circumstances of their
45 formation regions. Understanding the complex interplay of changing ocean variables and the
46 biological implication for marine ecosystems is a difficult task and requires not only modelling, but
47 also extensive data collection for monitoring, hypothesis testing and validation. Monitoring gaps still
48 remain in both in time and space, especially for marginal seas such as the Arctic Ocean or the
49 Mediterranean Sea.

50 The Mediterranean Sea has been identified as a region significantly affected by ongoing climatic
51 changes, like warming and decrease in precipitation (Giorgi, 2006). In addition, it is a region
52 particularly valuable for climate change research because it behaves like a miniature ocean (Bethoux
53 et al., 1999) with a well-defined overturning circulation characterized by spatial and temporal scales
54 much shorter than for the global ocean, with a turnover of only several decades. Being an
55 intercontinental sea, and subjected to more terrestrial nutrient inputs (river runoff, submarine
56 groundwater discharge) and atmospheric deposition, the Mediterranean Sea has a nitrate to phosphate
57 N:P ratio that is anomalously high compared to the “classical” world's oceans Redfield ratio,
58 indicating a general P-limitation regime, which becomes stronger along a west-to-east gradient. The
59 Mediterranean Sea is therefore a potential model to study global patterns that will be experienced in
60 the next decades worldwide, not only regarding ocean circulation, but also the marine biota (Lejeune
61 et al., 2010). Several environmental variables can act as stressors for marine ecosystems, by which
62 climatically driven ecosystem disturbances are generated (Boyd, 2011). These changes affect, among
63 others, the distribution of biogeochemical elements (including inorganic nutrients) and the functioning
64 of the biological pump and CO₂ regulation.

65 Within this context, the aim of this paper is to compile an extensive dataset of dissolved inorganic
66 nutrient observations (nitrate, NO_3^- ; phosphate, PO_4^{3-} ; and silicate, SiO_2) collected between 2004 and
67 2017 in the Western Mediterranean Sea (WMED), to describe the quality control techniques and to
68 provide the scientific community with a publicly available, long-term, quality controlled, and
69 internally consistent biogeochemical data product, contributing to previously published Mediterranean
70 Sea datasets like the MEDAR/Medatlas (time period:1908–1999), (Fichaut et al., 2003) and the
71 Mediterranean Sea – Eutrophication and Ocean Acidification aggregated datasets v2018 (time period:
72 1911-2017) provided by EMODnet Chemistry (Giorgetti al.,2018) available at
73 <https://www.seadatanet.org/Products/Aggregated-datasets>.

74 Both original and quality-controlled data are available in PANGAEA:

75 <https://doi.pangaea.de/10.1594/PANGAEA.904172>

76 Coverage: 44°N-35°S; -6°W-14°E

77 Location Name: Western Mediterranean Sea

78 Date start: May 2004

79 Date end: November 2017

80 **2 Dissolved inorganic nutrient data collection**

81 **2.1. The CNR dissolved inorganic nutrient data in the WMED**

82 Long-term time-series, such as the OceanSites global time series (www.oceansites.org), are a
83 fundamental prerequisite to understand and detect climate shifts and trends. However, biogeochemical
84 time-series are still limited to the northern ~~western~~ Western Mediterranean Sea (MOOSE network,
85 Coppola et al., 2019). Yet, inorganic nutrients in the Mediterranean Sea has received more attention in
86 recent years, and various datasets have been compiled to understand its unique characteristics such as
87 the one build by the PERSEUS project Consortium (“Policy-oriented marine environmental research
88 in the southern European seas” - EU FP7 project GA #287600), that included 100 cruises collected
89 during the project’s lifetime, in addition to those from other projects like SESAME, EU FP7 project
90 GA #GOCE-036949), and data products such as the MEDAR/Medatlas. In addition to that, the data
91 assembly system EMODnet Chemistry, a leading infrastructure supported by pan-European directorate
92 General MARE set up (Martin Miguez et al., 2019, Tintoré et al.,2019).

93 The dataset presented here consists of 24 oceanographic cruises (Fig. 1, ~~and~~ Table 1a and Table 1b)
94 conducted in the WMED on board of research vessels run by the Italian National Research Council
95 (CNR) and the Science and Technology Organisation Centre for Maritime Research and
96 Experimentation (NATO-STO CMRE). All cruises were merged into a unified dataset with 870
97 nutrient stations and ~ 9666 data points over a period of 13 years (2004-2017). The overall spatial
98 distribution of the stations covers the whole WMED, but the actual distribution strongly varies

99 depending on the specific cruise and most of the data are collected along sections. At all stations,
100 pressure, salinity and temperature were measured with a CTD-rosette system consisting of a CTD SBE
101 911 plus and a General Oceanics rosette with 24 12L Niskin Bottles. Temperature measurements were
102 performed with the SBE-3/F thermometer with a resolution of 10^{-3} °C; conductivity measurements
103 were performed with the SBE-4 sensor with a resolution of $3 \cdot 10^{-4}$ S/m. The probes were calibrated
104 before and after each cruise. During all CNR cruises, redundant sensors were used for both
105 temperature and salinity measurements.

106 Seawater samples for dissolved inorganic nutrient measurements were collected during the CTD up-
107 cast at standard depths (with slight modifications according to the depth at which the deep chlorophyll
108 maximum was detected). The standard depths are usually 5, 25, 50, 75, 100, 200, 300, 400, 500, 750,
109 1000, 1250, 1500, 1750, 2000, 2250, 2500, 2750, 3000 m. No filtration was employed, nutrient
110 samples were immediately stored at -20 °C. Note that sample storage and freezing duration varied
111 greatly from one cruise to another (Table 3 shows ~~the~~ cruises where this exceeded 1 year).

112 **2.2. Analytical methods for inorganic nutrients**

113 For all cruises, nutrient determination (nitrate, orthosilicate and orthophosphate) was carried out
114 following standard colorimetric methods of seawater analysis, defined by Grasshoff et al. (1999) and
115 Hansen and Koroleff (1999). For inorganic phosphate, the method is based on the reaction of the ions
116 with an acidified molybdate reagent to yield a phosphomolybdate heteropoly acid, which is then
117 reduced to a blue-colored compound (absorbance measured at 880 nm). Inorganic nitrate is reduced
118 (with cadmium granules) to nitrite that react with an aromatic amine leading to the final formation of
119 the azo dye (measured at 550 nm). Then, the nitrite separately determined must be subtracted from the
120 total amount measured to get the nitrate concentration only. The determination of dissolved silicon is
121 based on the formation of a yellow silicomolybdic acid reduced with ascorbic acid to blue-colored
122 complex (measured at 820 nm).

123 Nutrient analysis was performed in three laboratories. From 2004 to 2013, all cruises nutrients were
124 analysed by ENEA, while for those of 2015 (cruise #23) and 2017 (cruise #24), nutrient
125 concentrations were analysed by CNR-ISMAR. Referring to Table 1S, four different models of
126 autoanalyzer were used. Measurements from the autoanalyzer were reported in $\mu\text{mol L}^{-1}$. Inorganic
127 nutrient concentrations were converted to the standard unit $\mu\text{mol kg}^{-1}$, using sample salinity from CTD
128 and a mean laboratory analytical temperature of 20°C. Data from nutrient analysis were then merged
129 to ancillary CTD bottle data.

130 **2.3. Reference inorganic nutrient data**

131 In addition to the data collected during the above-mentioned cruises, and in order to perform the
132 secondary quality control (described below), we identified five reference cruises (Table 2), based on
133 their spatial and temporal distribution ~~of the data~~ and the reliability of the measurements (see Fig. 2 –
134 Table.3S Fig.1S). Cruises 06MT20110405 and 06MT20011018 are the only two Mediterranean
135 cruises included in the publicly available Global Ocean Data Analysis Project version 2 (GLODAPv2,
136 Olsen et al. 2016). These cruises, conducted on board the R/V Meteor, provide a reliable reference
137 because nutrient analysis strictly followed the recommendation of the World Ocean circulation
138 experiment (WOCE) and the GO-SHIP protocols (Hydes et al., 2010; ,Tanhua et al., 2013). Cruises
139 29AH20140426 and 48UR20070528 are to be included in the CARIMED data product (personal
140 communication by M. Álvarez, in preparation but not yet available) and have undergone rigorous
141 quality control following GLODAP routines. Finally, 29AJ20160818 was carried out in the framework
142 of the MedSHIP programme (Schroeder et al., 2015) and its data are available at
143 <https://doi.org/10.1594/PANGAEA.902293> (Tanhua, 2019).

144 **3 Quality Assurance and quality control methods**

145 Combining inorganic nutrient data from different sources, collected by different operators, stored for
146 different amounts of time, and analysed by multiple laboratories, is not a straightforward task. This is
147 widely recognized in the biogeochemical oceanographic community. Since the 1990s, several studies

148 and programmes (e.g. World Ocean Database, World Ocean Atlas, WOCE) have been devoted to
149 facilitate the exchange of oceanographic data and develop quality control procedures to compile
150 databases by the estimation of systematic errors (Gouretski and Jancke, 2004) to increase the inter-
151 comparability, generate consistent data sets and accurately observe the long-term change.

152 An example of a first quality control procedure is the use of reference materials that are available for
153 salinity (IAPSO, salinity standard by OSIL) and temperature (SPRT, Standard Platinum Resistance
154 Thermometer). As for the inorganic carbon, total alkalinity (Dickson et al., 2003) and inorganic
155 nutrients (Aoyama et al., 2016), certified reference materials (CRM) have been recently made
156 applicable for oceanographic cruises. However, since CRM are not always available or used for
157 biogeochemical oceanographic data, Lauvset and Tanhua (2015) developed a secondary quality
158 control tool to identify biases in deep data. The method suggests adjustments that reduce cruise to
159 cruise biases, increase accuracy and allow for the inter-comparison between data from various sources.
160 This approach, based on a crossover and inversion method (Gouretski and Jancke, 2004; Johnson et
161 al., 2001), was used to generate the CARbon IN Atlantic ocean (CARINA, see Hoppema et al., 2009),
162 GLODAPv2.2019 (Olsen et al., 2019) and PACIFICA (Suzuki et al., 2013) data [products](#) [bases](#).

Codice campo modificato

163 **3.1 Primary Quality control**

164 Each individual cruise was first subjected to a primary quality control (1st QC) that included a check of
165 apparent and extreme outliers in CTD salinity, nitrate, phosphate and silicate. Each parameter included
166 a quality control flag, following standard WOCE flags (Table 3). Surface, intermediate and deep layer
167 were evaluated separately because nutrient observations evolve differently in each layer. The
168 coefficient of variation (CV, defined as standard deviation over mean) was computed for each depth
169 layer. Coefficients of variation in the surface (0-250 db) layer were high (nitrate -CV=1.16, phosphate
170 CV=1.005, silicate CV=0.75) due to air-sea interaction (Muniz et al., 2001) occurring in this layer
171 rendering it difficult to flag. These influences are of reduced importance in the intermediate (250-1000
172 db) layer (nitrate CV=0.23, phosphate CV=0.31, silicate CV=0.24) and the deep (>1000 db) layer

173 (nitrate CV=0.15, phosphate CV=0.22, silicate CV=0.14), decreasing the total variance. Flags in the
174 upper and intermediate layer were thus set based on outliers within pressure ranges defined according
175 to standard pressures (0-10, 10-30, 30-60, 60-80, 80-160, 160-260, 260-360, 360-460, 460-560,
176 1000 db).

177 Below 1000 db, flagging included an inspection of nitrate to phosphate (N:P) and nitrate to silicate (N:
178 Si) ratios. The Median and Median Absolute Deviation (MAD) was computed by classes of pressure:
179 we considered as outlier any atypical observation and any value that departs from the median by more
180 than three MADs in the different pressure ranges for each cruise.

181 An overview of the nutrient distribution is provided with scatter plots, showing also the flagged
182 measurements (Fig. 3). Each measurement was flagged 2 (“Acceptable/ measured”) or flagged 3
183 (“Questionable”): 4.1% of nitrate data, 3.37% of phosphate data, 3.16% of silicate data, and 0.07% of
184 CTD salinity data were considered outliers and flagged 3. As highlighted by Tanhua et al. (2010), the
185 primary QC can be subjective depending on the expertise of the person flagging the data, thus flagging
186 could bring in some uncertainties.

187 In order to have a first assessment of the precision of each cruise measurements, the standard deviation
188 of observations deeper than 1000 db was calculated along with averages and standard deviations for
189 each cruise and by subregions to have an overview about nutrient content variability in the deep layer
190 and about the observations spatial spread of individual cruises (Table 4). Following the subdivision of
191 Manca et al. (2004), the WMED has been divided into subregions (Fig.2S, Table 2S) according to the
192 general circulation patterns (details in Manca et al.,2004). Table 4 displays the comparison of standard
193 deviation of deep measurements for each cruise and within subregions. The overall standard deviation
194 between cruises in the deep layer varied between 0.51 and 1.41 $\mu\text{mol kg}^{-1}$ for nitrate, between 0.1 and
195 1.64 $\mu\text{mol kg}^{-1}$ for silicate and between 0.025 and 0.078 $\mu\text{mol kg}^{-1}$ for phosphate. Regional standard
196 deviation of nitrate measurements below 1000 db varied between 0.08 $\mu\text{mol kg}^{-1}$ in the Gulf of Lion
197 (DF2) with cruise #9 and 1.6 $\mu\text{mol kg}^{-1}$ in the Balearic Sea (DS2) observations of cruise #14.

198 Phosphate lowest regional standard deviation was $0.01 \mu\text{mol kg}^{-1}$ found in the observations of cruise
199 #9 in Gulf of Lion (DF2), cruise #10 in Balearic Sea (DS2) and Algerian West (DS3), cruise #14 and
200 cruise # 15 in Tyrrhenian South (DT3), cruise #18 in Algero-Provençal (DF1) and Sardinia Channel
201 (DI1) while the highest standard deviation was $0.1 \mu\text{mol kg}^{-1}$ in the observations of cruise #12 in
202 Algerian West (DS3). As for silicate, the lowest standard deviation was $0.02 \mu\text{mol kg}^{-1}$ observed in
203 cruise #9 measurements of Gulf of Lion subregion (DF2) and the highest deep standard deviation was
204 observed in cruise #6 in its all subregions together with cruise #5 measurement in Tyrrhenian North
205 (DT1) with $1.83 \mu\text{mol kg}^{-1}$ standard deviation.

206 Cruises #3, #6 and #9 had the largest spatial extension (see right side of Fig. 9) with a high number of
207 samples over more than seven subregions (Table 4) and the geographical variability of the distribution
208 in dissolved inorganic nutrients results thus in the largest standard deviations. Conversely, cruises
209 with smaller spatial coverages have lower standard deviations. Therefore, a relatively small spatial
210 coverage and high standard deviation is considered as indicative of data with low precision (Olsen et
211 al., 2016). This applies to cruises #1, #5, and #16. Despite the small spatial coverage, samples of
212 nitrate and phosphate of cruise #5 have an overall standard deviation of $1.35 \mu\text{mol kg}^{-1}$ and $0.07 \mu\text{mol}$
213 kg^{-1} , respectively, a high standard deviation- pointed out in also in the regional standard deviation of
214 deep measurements in Tyrrhenian North (DT1) and South (DT3) . Cruise #1, with few stations in
215 Tyrrhenian North (DT1) and South (DT3) subregions and 21 samples below 1000 db, has an overall
216 standard deviation of $1.25 \mu\text{mol kg}^{-1}$ for nitrate, $0.06 \mu\text{mol kg}^{-1}$ for phosphate and $1.64 \mu\text{mol kg}^{-1}$ for
217 silicate. The regional standard deviation was relatively high for nitrate ($0.51\text{-}1.32 \mu\text{mol kg}^{-1}$),
218 phosphate ($0.02\text{-}0.065 \mu\text{mol kg}^{-1}$) and silicate ($0.53\text{-}1.83 \mu\text{mol kg}^{-1}$). A comparison with the deviations
219 from e.g. cruise # 2, carried out in the same year and e.g. cruise #17 (with a similar cruise track),
220 confirms the lower precision of the data of cruise #1. Similar considerations apply to the quality of
221 nitrate samples ($0.87\text{-}1.02 \mu\text{mol kg}^{-1}$) and silicate ($0.87\text{-}0.9 \mu\text{mol kg}^{-1}$) from cruise #16, covering a

222 small area in Tyrrhenian North (DT1) and South (DT3), compared to cruise #17, carried out in the
223 same regions (right side of Fig. 9 and Table 4).

224 Deep silicate measurements of cruise #6 have twice the overall standard deviation of silicate data of
225 cruise #8 from the same year. Adding to that, in the seven subregions, the regional standard deviation
226 of deep silicate observations was the highest, between 1.04-2 $\mu\text{mol kg}^{-1}$ which was relatively high
227 compared to the surrounding cruises that have observations in the same subregions. This is again
228 suggestive of the limited precision. On the other hand, trying to explain the source of relatively high
229 standard deviations in specific cruises is not always straightforward, as they could stem from a variety
230 of sources, sampling, conservation and analysis. The bottom water in the WMED exhibits a high
231 nutrient content below 1000 db (Table 4), due to the longer residence time. Dividing the WMED into
232 subregions, has effectively removed the natural spatial change in nutrients, making the interpretation
233 of the standard deviation a matter of the precision of the measurements only.

234 | In Table 4, deep averages by subregions showed that overall nutrient concentration fluctuated around
235 | $7.4 \pm 0.9 \mu\text{mol kg}^{-1}$ for nitrate, $0.3 \pm 0.06 \mu\text{mol kg}^{-1}$ for phosphate and $7.7 \pm 0.8 \mu\text{mol kg}^{-1}$ for silicate,
236 | similar findings were reported by Manca et al. (2004). Comparing cruise averages in each region
237 | enabled the identification of “suspect” cruises. Cruise #24 has the lowest deep average in nitrate in
238 | ~~Algéro~~ Algero-Provençal (DF1), Tyrrhenian North (DT1) subregions and Sardinia Channel (D11). As
239 | for silicate of cruises #24 and #16 was very low compared to the overall regional average in Liguro-
240 | Provençal (DF3) and Tyrrhenian South (DT3) subregions. Deep average of phosphate did not show
241 | any outlier cruises in all subregions. Different reasons could explain the low precision in the samples,
242 | freezing is one. Although it is a valid preservation method (Dore et al., 1995~~6~~), the error is higher when
243 | samples were not analysed immediately (Segura-Noguera et al., 2011), so the storage time could
244 | influence.

245 3.2 Secondary Quality control: the crossover analysis

246 | The method used to perform the secondary QC on the WMED dissolved inorganic nutrient dataset ~~in~~
247 | ~~the WMED~~ makes use of the quality-controlled reference data, and the crossover analysis toolbox
248 | developed by Tanhua (2010^a) and Lauvset and Tanhua (2015). The computational approach is based
249 | on comparing the cruise data set to a high-quality reference data set to quantify biases, described in
250 | detail in Tanhua et al. (2010^b). Here, we summarize the technique with emphasis on inorganic
251 | nutrients. The first step consisted of selecting reference data, as described in section 2.3. The second
252 | step is the crossover analysis that was carried out using a MATLAB Toolbox (available online:
253 | https://cdiac.ess-dive.lbl.gov/ftp/oceans/2nd_QC_Tool_V2/) where crossovers are generated as
254 | difference between two cruises using the “running cluster” crossover routine. Each cruise is thus
255 | compared to the chosen set of reference cruises. For each crossover, samples deeper than 1000 db are
256 | selected within a predefined maximum distance set to 2° arc distance, defined as a crossing region, to
257 | ensure the quality of the offset with a minimum number of crossovers and to minimize the effect of the
258 | spatial change. The reason to select measurements deeper than 1000 db, is to remove the high
259 | frequency variability associated to mesoscale features, biological activity and the atmospheric forcing
260 | acting in the upper layers, that might induce changes in biogeochemical properties of water masses.
261 | On the other hand, also the deep Mediterranean cannot be considered truly “unaffected” by changes, as
262 | it is intermittently subjected to ventilation (Schroeder et al., 2016; Testor et al., 2018) and the real
263 | variability can be altered in adjusting data. The computational approach takes this into account, since
264 | weights are given to the less variant profile in the crossing region, according to the “confidence” in the
265 | determined offset of the compared profiles (i.e. the weighted mean offset of a given crossover-pair is
266 | weighted to the depth where the offsets of all compared profiles have the smallest variation (which
267 | indeed is strongly interlinked with the degree of variance of each profile) (for further details see
268 | Lauvset and Tanhua, 2015).

269 | Before identifying crossovers, each profile was interpolated using the piecewise cubic Hermite method
270 | and the distance criteria outlined in Lauvset and Tanhua (2015), their Table 1^a, ~~and~~ detailed in Key et

271 al. (2004). The crossover is a comparison between each interpolated profile of the cruise being
272 evaluated and the interpolated profile of the reference cruise. The result is a weighted offset (defined
273 as difference cruise/reference) and a standard deviation of the offset. The standard deviation is
274 indicative of the precision; however, it is important to note that this assumption only works because it
275 is a comparison to a reference, and the absolute offset is indicative of accuracy.

276 The third step consists in evaluating and selecting the suggested correction factor that was applied to
277 the whole water column. The correction factor was calculated from the weighted mean offset of all
278 crossovers found between the cruise and the reference data set, involving a somewhat subjective
279 process.

280 For inorganic nutrients, offsets are multiplicative so that a weighted mean offset > 1 means that the
281 measurements of the corresponding cruise are higher than the measurements of the reference cruise in
282 the crossing region and applying the adjustment would decrease the measured values. The magnitude
283 of an increase or a decrease is the difference of the weighted offset from 1. In general, no adjustment
284 smaller than 2% (accuracy limit for nutrient measurements) is applied (detailed description is found in
285 Hoppema et al., 2009; Lauvset and Tanhua, 2015; Olsen et al., 2016; Sabine et al., 2010; Tanhua et al.,
286 2010^b).

Formattato: Francese (Francia)

287 The last step is the computation of the weighted mean (WM) to determine the internal consistency and
288 quantify the overall accuracy of the adjusted product (Hoppema et al., 2009; Sabine et al., 2010;
289 Tanhua et al., 2009), with the difference that our assessment is based on the offsets with respect to a
290 set of reference cruises. This WM reflects the absolute weighted mean offset of the data set compared
291 to the reference data set, hence the smaller the WM the higher the internal consistency. The accuracy
292 was computed from the individual absolute weighted offsets. The WM, which will be discussed in
293 section 4.4., was computed using the individual weighted absolute offset (D) of number of crossovers

294 (L) and the standard deviation (σ): $WM = \frac{\sum_{i=1}^L D(i)/(\sigma(i))^2}{\sum_{i=1}^L 1/(\sigma(i))^2}$

295 **4 Results of the secondary QC and recommendations**

296 The results of the secondary QC revealed the necessary corrections for nitrate, phosphate and silicate.
297 Four cruises were not considered in the crossover analysis: cruises #7 and #11 do not have enough
298 stations > 1000 db (at least 3 to get valid statistics), while cruises #19 and #21 were outside the spatial
299 coverage of the reference cruises. Cruises that were not used for the crossover analysis are made
300 available in the original dataset but were not included in the final data product (see Supplementary
301 material – Part 2 (A2)).

302 Overall, we found a total number of 73 individual crossovers for nitrate, 72 for phosphate and 54 for
303 silicate. An example of the running cluster crossover output is shown in Fig.4. Results of the crossover
304 analysis is an adjustment factor for each cruise and each nutrient, that are shown in Table 5 and Fig. 5-
305 6-7. The adjustment factor was calculated from the weighted mean of absolute offset summarized in
306 Table 6 and Fig. 3S-4S-5S. Table 6 details the improvement of the weighted mean of absolute offset
307 by cruise prior to and after adjustments, the information is also displayed graphically in Fig. 3S-4S-5S.
308 Cruises are in chronological order in all figures and tables.

309 **4.1 Nitrate**

310 The crossover analysis suggests a significant adjustment for nitrate concentrations on 15 cruises,
311 between 0.94 and 0.98 (for adjustments <1) and between 1.02 and 1.34 (for adjustments >1) (Table 5
312 and Fig.5). Offsets suggest that the deep measurements of cruises #1, #3, #4, #5, #6, #8, #12, #13, #15,
313 #16, #23 and #24 need to be adjusted towards higher concentrations, when compared to the respective
314 reference (Fig.3S).

315 Nitrate observations of cruises #2, #9 and #10 on the other hand were higher than the reference cruises
316 and exhibit variation outside the accepted accuracy limit, thus requires a downward adjustment.

317 Finally, five cruises (#14, #17, #18, #20, and #22) were consistent with the reference data and no
318 adjustment was necessary. Considering the weighted mean of absolute offset after adjustments shown
319 in Table 6, two cruises (#5 and #24) required large correction factors but ~~still~~ remain outside the
320 accuracy threshold (Fig. 5). These cruises are considered in detail later (section 4.4).

321 4.2 Phosphate

322 For phosphate the crossover analysis suggests adjustments for 20 cruises, as shown in Fig. 6. Deep
323 phosphate measurements of 15 cruises (Table 6) appear to be lower than the respective reference
324 measurements (i.e. phosphate data of these cruises require an upward adjustment), while the data of
325 five cruises (#2, #3, #4, #6, #24) are higher (i.e. they need a downward adjustment) (Fig.4S). Applying
326 all the indicated adjustments, the large offsets of cruises #2, #3, #4, #6, #8, #9, #10, #18, #20, #23 and
327 #24 are reduced and ~~become~~ became consistent with the reference. Cruises #1, #5, #12, #13, #14, #15,
328 #16, #17, and #22 retain an offset even after applying the indicated adjustment. These cruises are
329 considered in detail later.

330 According to Olsen et al. (2016), if a temporal trend is detected in the offsets, no adjustments should
331 be applied. There is indeed a decreasing trend between 2008 and 2017 in the phosphate correction
332 factor (Fig. 6), and thus an increasing one in the weighted mean offset (Fig.4S), implying a temporal
333 increase of phosphate. Therefore, phosphate data of the cruises being part of the trend were not
334 flagged as questionable, except some cruises that are discussed further in section 4.4.

335 Comparing phosphate before and after adjustments, the corrections did minimise the difference with
336 the reference, while the actual variation with time was preserved (Fig.6). The temporal trend towards
337 higher phosphate concentrations in the Mediterranean Sea is considered to be real, even though
338 studies concerning the biogeochemical trends in the deep layers of the WMED are scarce (Pasqueron
339 et al., 2015). However, this variation could be consistent with the findings of Béthoux et al.(1998,
340 2002) and the modelling studies by Moon et al. (2016) and Powley et al. (2018) who indeed found an

341 increasing trend in phosphate concentrations over time, due to the increase in the atmospheric and
342 terrestrial inputs.

343 **4.3 Silicate**

344 The results of the crossover analysis for silicate suggests corrections for all cruises (Fig.7). The
345 crossovers indicate that deep silicate measurements are lower in the evaluated cruises than in the
346 corresponding reference cruises (i.e. they need to be adjusted upward) (Fig.5S). This is likely to be a
347 direct result of freezing the samples before analysis, since the reactive silica polymerizes when frozen
348 (Becker et al., 2019). After applying the adjustment (Table 5), as expected, the offsets are reduced
349 (Table 6), but five cruises (#1, #5, #6, #15, and #16) remain outside the accuracy envelope. Due to the
350 large offsets, these cruises will be discussed further in section 4.4.

351 **4.4 Discussion and recommendation**

352 Adjustments were evaluated for each cruise separately. As a general rule, no correction was applied
353 when the suggested adjustment is strictly within the 2% limit (indicated with NA in Table 5). The
354 average correction factors were 1.06 for nitrate, 1.14 for phosphate and 1.14 for silicate, respectively.
355 To verify the results, we re-ran the crossover analysis and re-computed offsets and adjustment factors
356 using the adjusted data (as shown in blue in Fig. 3S-4S-5S and Fig. 5-6-7). Most of the new
357 adjustments are within the accuracy envelope and few are outside the limit, except for the cruises
358 belonging to the above mentioned “phosphate-trend” and the other outlying cruises which are detailed
359 hereafter. By the application of adjustments, the deep-water offsets were reduced. This can be seen in
360 the decrease of the weighted mean offset between the data before adjustments (after 1st QC, Fig. 3S-
361 4S-5S, in grey) and the adjusted data (after 2nd QC, Fig. 3S-4S-5S, in blue).

362 Referring to the analysis detailed in section 3.2, the internal consistency of the nutrient data set has
363 improved and increased significantly after the adjustment, from 4% for nitrate, 19% for phosphate and
364 13% for silicate, to a more unified dataset with 3 % for nitrate, 6 % for phosphate and 3% for silicate.

365 A comparison between the original and the adjusted nutrient observations is shown in Fig. 8A-B-C,
366 indicating an improvement in the accuracy based on the reference ~~measurement data~~ and a relatively
367 reduced range particularly for phosphate (Fig. 8B). Figure 8. D-E scatterplots show that after the
368 quality control, nutrient stoichiometry slopes obtained from regressions, between tracers along the
369 water column ~~show demonstrate~~ a strong coupling and provide a nitrate to phosphate ratio of ~22.09
370 and a nitrate to silicate ratio of ~0.94. These values are consistent with nutrient ratios range found in
371 the WMED as reported in Lazzari et al. (2016); Pujo-Pay et al., (2011) and Segura-Noguera et al.
372 (2016).

373 The regression model is more accurate after adjustments with an improved r^2 for N:P (from 0.81 to
374 0.90) and for N: Si (from 0.85 to 0.87).

375 ~~One of the main reasons for an upward/ downward bias would be the lack of use of CRM for nutrients~~
376 ~~in all cruises as also noted in CARINA (Tanhua et al., 2009) or in the most recent global comparability~~
377 ~~study by Aoyama (2020).~~ ▲

Formattato: Inglese (Regno Unito)

378 In the following some details on the adjustment of specific cruises are given:

379 Cruise #2 [48UR20041006] needed an adjustment of 0.98 for nitrate, 0.9 for phosphate and 1.06 for
380 silicate. Most of the crossover profiles occur in the Tyrrhenian ~~sea~~Sea (Tyrrhenian North and
381 Tyrrhenian South subregions). After adjustment, the cruise is inside the 2% envelope.

382 Cruise #3 [48UR20050412] appeared to be outside the 2% envelope before adjustments. Its offsets
383 with five reference cruises, crossing the Tyrrhenian ~~sea~~Sea, Sardinia ~~channel~~Channel, Gulf of Lion
384 and Algero-Provençal subregions, showed that nitrate and silicate values to be relatively low, and thus
385 an adjustment of 1.08 and 1.15 was applied respectively. On the other hand, phosphate values were
386 relatively high, and a 0.93 adjustment was applied.

387 Cruise #4 [48UR20050529] correction factor estimate was based on five crossovers that covered five
388 subregions: Tyrrhenian South, Sardinian ~~channel~~Channel, Algerian East and West and the Alboran
389 ~~sea~~Sea. Table 4) show that there are no large differences between regional averages within the cruise
390 which justify an adjustment of 1.04 for nitrate, 0.85 for phosphate and 1.183 for silicate.

391 Cruise #8 [48UR20060928] was adjusted by 1.03 for nitrate, 1.14 for phosphate and 1.1 for silicate,
392 because it showed values to be low compared to four references. After adjustment, the data were
393 inside the acceptable range.

394 Cruise #9 [48UR20071005] values of nitrate were slightly outside the 2% envelope before
395 adjustments, similar to phosphate and silicate that were lower compared to the reference. The
396 adjustments of 0.97 for nitrate, 1.14 for phosphate and 1.115 for silicate suggested by the mean offset
397 against the reference cruises were recommended.

398 Cruise #13 [48UR20090508] has three crossovers in the common crossing zone that included
399 Tyrrhenian North, Tyrrhenian South and Sardinia Channel subregions. The crossover suggests that this
400 cruise has too low values and needs an adjustment of 1.05 for nitrate, 1.33 for phosphate and 1.15 for
401 silicate.

402 Cruise #14 [48UR20100430] has a mean offset with four reference cruises that suggests an adjustment
403 ~~factor of 1.34~~ for phosphate ~~of 1.34~~ and ~~1.123~~ for silicate ~~of 1.123~~. Nitrate did fall within the accuracy
404 envelope; ~~no adjustment was needed~~.

405 Cruise #10 [48UR20080318] has only three crossovers in the Algero-Provençal subregion, showing
406 that nitrate is too high compared to the reference while phosphate and silicate are slightly lower. We
407 therefore applied the adjustments of Table 5, since the deep averages ~~for in~~ each region (~~†~~Table 4) did
408 not show large regional difference.

409 | Cruise #17 [48UR20110421] crossover analysis did not suggest any correction for nitrate; ~~h~~However,
410 | ~~for phosphate and silicate~~ with an offset based on two crossovers in the Tyrrhenian North and South
411 | subregions, adjustments were recommended for phosphate (1.25) and silicate (1.12), for being lower
412 | than the reference cruises.

413 | Cruise #18 [48UR20111109] is similar to cruise #17, since it was suggested to ~~correct~~ adjust
414 | phosphate by 1.14 and silicate by 1.09, based on four crossovers in the Tyrrhenian North and South,
415 | Sardinia ~~channel~~ Channel and Algero-Provençal subregions.

416 | Cruise #20 [48UR20120111] has four crossovers over the Tyrrhenian North and South and Algero-
417 | Provençal subregions. Its measurements were slightly lower than the reference cruises suggesting a
418 | correction factor of 1.17 for phosphate and 1.08 for silicate.

419 | Cruise #22 [48UR20131015] has similar correction factors as cruise #20, based on three crossovers in
420 | the Sardinia ~~channel~~ Channel and Tyrrhenian North and South subregion, with measurements being
421 | lower than the reference.

422 | Cruise #23 [48QL20150804] showed nutrient values slightly lower than the reference cruises as well,
423 | suggesting small correction factors of 1.02 for both nitrate and phosphate and 1.08 for silicate ~~that, a~~
424 | correction factors that were based on offsets with five cruises.

425 | Below, we discuss the recommended flags in the final product (Table 3: see supplementary Materials
426 | Part-2 (A2)) assigned for some cruises that needed further consideration, since they required larger
427 | adjustment factors:

428 | Cruise #1 [48UR20040526]: The adjusted values are still lower than the reference (Fig.5-6-7-Fig.3S-
429 | 4S-5S) and are still outside the 2% accuracy range. This cruise had stations in the Sicily Strait,
430 | Tyrrhenian North and South and Ligurian East sub-regions- (Fig. 9, right side) and only 4 stations
431 | were deeper than 1000 db (those within the Tyrrhenian Sea). The low precision of this cruise has

432 | already been evidenced during the primary-1st QC (section 3.1). We recommend flagging this cruise as
433 | questionable (flag 3).

Formattato: Apice

434 | Cruise #5 [48UR20051116]: This cruise took place between Sicily Strait and the Tyrrhenian North and
435 | South (Fig. 9, right side). Nitrate, phosphate and silicate data were lower than those from other cruises
436 | (#3 and #4) run the same year (Fig. 5-6-7-Fig.3S-4S-5S) and are still biased after adjustments.
437 | Considering the limited precision and the low number of crossovers, it is recommended to flag the
438 | cruise as questionable (flag 3).

439 | Cruise #6 [48UR20060608]: This cruise had an offset with five cruises giving evidence that
440 | adjustments of 1.05 for nitrate, 0.86 for phosphate and 1.26 for silicate are needed. The silicate bias
441 | was reduced after adjustment but remained eds large with respect to the accuracy limit (Fig. 7-Fig. 5S).
442 | This cruise has a wide geographic coverage, with stations along 9 sections (Fig. 9, right side).
443 | Considering also the high standard deviation (Table 4), which is partially attributed to the spatial
444 | coverage of the cruise, there is still uncertainty about the quality of the samples. It is recommended to
445 | flag silicate data of cruise #6 as questionable (flag 3).

446 | Cruise #12 [48UR20081103]: Phosphate data have low accuracy with respect to the reference cruises
447 | (Fig. 6-Fig. 4S). This cruise has stations along a longitudinal section from ~~the~~ Sicily Strait to the
448 | Alboran Sea, which might explain the large standard deviation of deep phosphate samples (Table 4).
449 | Cruise #12 was given a correction of 1.08 for nitrate, 1.12 for silicate and 1.38 for phosphate. The
450 | mean offset from five crossovers computed within the Tyrrhenian South, Sardinia Channel, Algerian
451 | East, Algerian West and Alboran Sea subregions suggests that this cruise has lower nutrient values
452 | than the reference cruise. After adjustment, cruise #12 is within the acceptable range for nitrate and
453 | silicate but not for phosphate as highlighted in section 3.2. In addition, considering the relatively high
454 | number of stations >1000 db and a plausible trend in phosphate, it is recommended to flag the
455 | phosphate data as good/acceptable (flag 2).

456 | Cruise #15 [48UR20100731]: This cruise has^d 149 station along a similar track as cruise #12, but
457 shows larger offsets for phosphate and silicate (Fig. 6-7-Fig. 4S-5S), compared to cruise #12.
458 Considering that deep silicate data was not of low quality (small standard deviation, see Table 4), and
459 that deep phosphate fall within the “phosphate-trend” discussed above, these data are flagged
460 good/acceptable (flag 2).

461 Cruise #16 [48UR20101123]: The cruise shows large offsets for phosphate and silicate (Fig. 6-7- Fig.
462 4S-5S), similar to cruise #15. Considering that the overall cruise standard deviation of silicate samples
463 below 1000 db was relatively high (1.02 over 14 samples, see Table 4), and that it has only one
464 crossover between the Tyrrhenian North and South subregions (Table 6), and that when comparing
465 deep regional averages, this cruise had the lowest average silicate value, it is recommended to flag
466 silicate data of cruise #16 as questionable (flag 3). As for phosphate, the cruise is part of the
467 “phosphate-trend” and is therefore flagged good/acceptable (flag 2).

468 Cruise #24 [48QL20171023]: This cruise has the largest offset for nitrate even after adjustment. It is
469 very likely due to a difference between laboratories (calibration standards) concerning nitrate, which
470 needs to be flagged as questionable (flag 3) in the final product.

471 There are several sources of bias in the observation. One of the main reasons for an upward/
472 downward bias would be the difference in the nutrient’s chemical analytical method and the lack of
473 use of CRM in all cruises as also noted in CARINA (Tanhua et al., 2009) or in the most recent global
474 comparability study by Aoyama (2020). ▲

Formattato: Bordo: Superiore: (Nessun bordo), Inferiore: (Nessun bordo), A sinistra: (Nessun bordo), A destra: (Nessun bordo), Tra : (Nessun bordo)

Formattato: Inglese (Stati Uniti)

475 The eCruises discussed in this section were not removed from the final product but are retained along
476 with their recommended quality flag (Table 3) detailed above and in the supplementary material – Part
477 2 (A2)). We have done the evaluation of their overall quality but leave it up to the users how to
478 appropriately use these data.

479 4.5 Product assessment: Comparison with MEDATLAS

480 | Averages water mass biogeochemical properties have been computed from the adjusted product (Table
481 | 7), and compared to the MEDAR/Medatlas annual climatological profiles, downloaded from the
482 | Italian NODC website (<http://doga.ogs.trieste.it/medar/>) given by Manca et al. (2004), in order to
483 | evaluate and assess the new product. Since nutrient properties exhibit differences with depths, we
484 | compared average nutrient concentrations of the three main water masses in twelve subregions of the
485 | WMED (Table 7, Fig 2S).

486 | The results of Table 7 compares water mass biogeochemical properties with the reference climatology.
487 | The new product agrees well with the Medatlas climatology. However, there are some distinctions.
488 | The surface layer (0-150db) is characterized by a low nutrient content. The surface nitrate varies
489 | between 0.69 and 2.75 $\mu\text{mol kg}^{-1}$ with a maximum found in the Ligurian East (DF4) and the minimum
490 | in the Alboran Sea (DS1) subregions, similar values were recorded in the climatology (0.61- 3.00
491 | $\mu\text{mol kg}^{-1}$). The differences in nitrate averages in the surface layer are observed in the Gulf of Lion
492 | (DF2) where the new product is higher than the climatology and slightly lower in the Liguro-
493 | Provençal (DF3). As for, the surface content in phosphate, it varied between 0.04 and 0.16 $\mu\text{mol kg}^{-1}$
494 | with a maximum found in the Ligurian East (DF1) and a minimum in the Alboran Sea (DS1), alike the
495 | Medatlas climatology, where phosphate averages fluctuate between 0.05 and 0.19 $\mu\text{mol kg}^{-1}$. The new
496 | product is slightly lower compared to the climatology. As to the average surface in silicate, it varies
497 | between 1.36 and 2.91 $\mu\text{mol kg}^{-1}$ with a minimum found in the Ligurian East (DF4), the maximum in
498 | the Gulf of Lion (DF2)) while in the climatology, it varied between 1.27 and 2.31 $\mu\text{mol kg}^{-1}$ (the
499 | minimum in the Ligurian East (DF4) and the maximum in the Alboran Sea (DS1)). The new product is
500 | slightly higher in silicate.

501 | Overall, the differences in the surface layer are observed in the Gulf of Lion (DF2), the Liguro-
502 | Provençal (DF3) and; the Ligurian East (DF4) regions; which could be is-due to the intense variability
503 | of the vertical mixing occurring in the northern WMED compared to the other subregions.

504 In the intermediate layer, averages were computed from the depth of the salinity maximum (S_{\max})
505 $\pm 100\text{m}$ from a regional average profile, indicative of the Levantine Intermediate Water (LIW) core.
506 Nitrate average varied between 4.94 and $9.32 \mu\text{mol kg}^{-1}$ where the minimum content ~~was~~ recorded
507 in Sicily strait (DI3) and the maximum in the Algerian West (DS3) while ~~in the~~ in the Medatlas
508 climatology, nitrate was between 5.14 and $8.60 \mu\text{mol kg}^{-1}$. In average, the lowest content in nitrate
509 was in ~~the~~ Tyrrhenian North (DT1) ~~and~~ South (DT3), Sardinia Channel (DI1) and Sicily Strait (DI3)
510 while LIW of ~~the~~ Gulf of Lion (DF2), Liguro-Provençal (DF3), Ligurian East (DF4), Balearic Sea
511 (DS2), Algero-Provençal (DF1), Alboran Sea (DS1), Algerian West (DS3) and East (DS4) subregions
512 ~~was~~ relatively rich in nitrate. Compared to the Medatlas product, though the new product ~~was~~ slightly
513 higher mainly in the Gulf of Lion (DF2), Ligurian East (DF4) and Balearic Sea (DS2). As for
514 phosphate, LIW averages showed similar behavior as nitrate, the lowest phosphate content ($0.21 - 0.27$
515 $\mu\text{mol kg}^{-1}$) ~~was~~ observed in the Eastern subregions of WMED (~~Sicily Strait (DI3), Sardinia Channel~~
516 ~~(DI1), Tyrrhenian South (DT3) and North (DT1)~~), when the maximum concentrations ($0.4 - 0.37 \mu\text{mol}$
517 kg^{-1}) were reported in the Western subregions of the WMED (~~the Alboran Sea (DS1), Algerian West~~
518 ~~(DS3) and East (DS4), Balearic Sea (DS2) and Gulf of Lion (DF2)~~). The large differences between the
519 two products were in ~~the~~ Ligurian East (DF4) and the Alboran Sea (DS1), subregions of few numbers
520 of observations.

521 Concerning silicate, the lowest average concentration ($5.25 \mu\text{mol kg}^{-1}$) ~~was~~ observed in LIW core of
522 ~~the~~ Sicily Strait (DI3,) and the maximum concentrations ($8.66 - 8.77 \mu\text{mol kg}^{-1}$) ~~were~~ in ~~the~~ Alboran
523 Sea (DS1) and Gulf of Lion (DF2), similar values were recorded in the Medatlas climatology ($4.86 -$
524 $7.95 \mu\text{mol kg}^{-1}$). There are some discrepancies, where the new product ~~was~~ higher particularly in the
525 Gulf of Lion (DF2), Liguro-Provençal (DF3) and Algerian West (DS3) subregions. This difference is
526 explained by the limited number of observations within depth range in the new product compared to
527 the observations used in the climatology in these subregions.

528 Referring to Manca et al.,(2004), the LIW core salinity values are relatively more pronounced in Sicily
529 | s, Strait (DI3), Sardinia ~~channel~~ Channel (DI1) and in the Tyrrhenian South (DT3) ~~and~~; North (DT1)
530 | subregions, where nutrients were lower than the Western subregions (DS3, ~~DS4~~, DS1 , DF1, DS2,
531 | DF4, DF3, DF2). The averages of nutrient within the LIW core ties well with the Medatlas
532 | climatology averages (Table 7), except in subregions with important vertical mixing.

533 We have verified also average biochemical properties in the deep layer (below 1500db). The new
534 | product is slightly higher in nitrate averages ($7.74 - 8.37 \mu\text{mol kg}^{-1}$) than the Medatlas climatology
535 | ($7.12 - 8.06 \mu\text{mol kg}^{-1}$) (Table 7). The largest difference ~~is~~ was found in Tyrrhenian South (DT3) and
536 | North (DT1) subregions. This difference could be due to the fact that, we are comparing two different
537 | time periods (2004-2017 and 1908-2001). As for the deep layer phosphate, average concentrations
538 | ~~varied~~ varied between 0.35 and $0.37 \mu\text{mol kg}^{-1}$ and ~~were~~ were within the climatology limits ($0.31 - 0.40 \mu\text{mol kg}^{-1}$)
539 | ¹). In all subregions, there ~~was~~ was not large differences. Overall, phosphate ~~was~~ was in accordance with the
540 | Medatlas climatology. Similar to nitrate, deep average silicate in the new product ($8.64 - 9.21 \mu\text{mol kg}^{-1}$)
541 | ¹) ~~was~~ was higher than the climatology (7.51 to $9.04 \mu\text{mol kg}^{-1}$). The largest difference in average silicate
542 | ~~was~~ was observed in ~~the~~ Tyrrhenian North (DT1), South (DT3) and Liguro-Provençal (DF3) subregions.

543 We then used the Root Mean Squared Error (RMSE) as statistical index to quantify the difference
544 | between averaged regional profiles from the new products and Medatlas product. The climatology
545 | annual profiles were interpolated to the regional average profiles of the new product, and the average
546 | RMSE for each layer and subregion was calculated. Fig. 10 shows the regional evolution of RMSE in
547 | the main water masses for the three nutrients. For nitrate (Fig. 10 A), the RMSE ~~in the surface layer~~ in the surface layer
548 | ~~varied~~ varied between $0.12 \mu\text{mol kg}^{-1}$ ~~(in the Tyrrhenian North (DT1))~~ and $1.36 \mu\text{mol kg}^{-1}$ ~~(in the Gulf of~~
549 | ~~Lion (DF2))~~ ~~in the surface layer,~~ in the intermediate layer, the RMSE was between $0.07 \mu\text{mol kg}^{-1}$ ~~(in~~
550 | ~~the Sardinia Channel (DI1))~~ and $2.35 \mu\text{mol kg}^{-1}$ ~~(in the Gulf of Lion (DF2))~~ ~~in the intermediate layer,~~
551 | and was lower in the deep layer, between $0.11 \mu\text{mol kg}^{-1}$ ~~(in the Algerian East (DS4))~~ and $0.79 \mu\text{mol}$
552 | ~~kg⁻¹ (the Gulf of Lion (DF2)).~~ The RMSE decreases in the Algerian East (DS4), Tyrrhenian North

553 (DT1), Tyrrhenian South (DT3), Sardinia Channel (DI1) and Sicily Strait (DI3). This illustrates the
554 low difference between the two products.

555 For phosphate (Fig. 10 B), the RMSE ranges between $0.0022 \mu\text{mol kg}^{-1}$ (in the Tyrrhenian South
556 (DT3)) and $0.12 \mu\text{mol kg}^{-1}$ (in the Ligurian East (DF4)) in the surface layer, and is between 0.003
557 $\mu\text{mol kg}^{-1}$ (in the Liguro-Provençal subregion (DF3)) and $0.048 \mu\text{mol kg}^{-1}$ (in the Alboran Sea (DS1))
558 at intermediate depths, while in the deep layer RMSE varied between 0.0087 (in the Gulf of Lion
559 (DF2)) and $0.057 \mu\text{mol kg}^{-1}$ (in the Tyrrhenian North (DT1)).

560 Regarding Silicate RMSE (Fig. 10 C) in surface, it varied between $0.13 \mu\text{mol kg}^{-1}$ (in the Algero-
561 Provençal subregion (DF1)) and $3.5 \mu\text{mol kg}^{-1}$ (in the Ligurian East subregion (DF4)) in the surface
562 layer. A lower RMSE between $0.10 \mu\text{mol kg}^{-1}$ (in the Sardinia Channel (DI1)) and $2.54 \mu\text{mol kg}^{-1}$ (in
563 the Gulf of Lion (DF2)) was reported in the intermediate layer, and the results in deep layer, RMSE
564 were ranges between $0.33 \mu\text{mol kg}^{-1}$ (in the Algerian East (DS4)) and $1.43 \mu\text{mol kg}^{-1}$ (in the Liguro-
565 Provençal subregion (DF3)).

566 The best agreement between the two products was observed in the intermediate and deep layer. The
567 lowest RMSE was confined to the deep layer in most of the subregions while the highest difference
568 was found in the surface layer since it is subjected to intense vertical mixing mainly in the northern
569 WMED. Comparing averages in subregions, showed similar differences in nutrient between the two
570 products particularly in the Gulf of Lion (DF2), the Liguro-Provençal (DF3), Ligurian East (DF4) and
571 Algerian East (DS4), due to the relative high variability in nutrient concentrations in these subregions.
572 These differences are not significant as there is discrepancy on the number of observations used in the
573 two products. Overall, inorganic nutrients of the new product agree very well with the
574 MEDAR/Medatlas climatology. The main features of the spatial distribution in the inorganic nutrients
575 were in accordance with the findings of Manca et al., (2004), where the relative high content in
576 nutrient was found in the intermediate layer of the Algerian subregions (DF1, DS3, DS4) than in other
577 subregions (Table 7). Besides, the highest concentrations in deep layer silicate were reported in the

578 Algerian subregions in the two products ($9.21 \mu\text{mol kg}^{-1}$ (DS3) in the new product; $9.04 \mu\text{mol kg}^{-1}$
579 (DS4) in the climatology) ~~in the two products~~, which is indicative of the poor regional ventilation and
580 of the longer residence time of deep water especially in these subregions.

581 **5 Final remarks**

582 An internally consistent data set of dissolved inorganic nutrients has been generated for the WMED
583 (2004-2017). The accuracy envelope for nitrate and silicate was set to 2%, a predefined limit used in
584 GLODAP and CARINA data products. Regarding phosphate data, these were almost entirely outside
585 this limit, because of its natural variations and the overall very low concentrations in the WMED, a
586 highly P-limited basin. Using a crossover analysis (2nd QC toolbox) to compare cruises with respect to
587 reliable reference data, improved the accuracy of the measurements by bias-minimizing the individual
588 cruises. ~~t~~The new product was broadly in consistent with the earlier climatology MEDAR/Medatlas.

589 The publication of a quality-controlled extensive (spatially and temporally) database of inorganic
590 nutrients in the WMED was timely and fills a gap in information that prevented baseline assessments
591 on spatial and temporal variability of biogeochemical tracers in the Mediterranean. In combination
592 with older databases in the same region (e.g. bottle data available in the MEDAR/Medatlas database),
593 this new data productbase will thus constitute a pillar on which the Mediterranean marine scientific
594 community will be able to build on original research topics on biogeochemical fluxes and cycles and
595 their relation to hydrological changes that occurred in the period covered by the dataset. The dataset is
596 also relevant for the modelling community as it can be used as an independent data product to assess
597 reanalysis products or it can be assimilated in new reanalysis products.

598 **6 Data availability**

599 The final product is available as a .csv merged file from PANGAEA, and can be accessed at
600 <https://doi.pangaea.de/10.1594/PANGAEA.904172> (Belgacem et al. 2019).

601 Ancillary information is in the supplementary materials with the list of variables included in [the](#)
602 original and final product. Table [1a](#) and [Table 1b](#) summarizes all cruises included in the dataset. The
603 dataset include frequently measured stations and key transects of the WMED with in situ physical and
604 chemical oceanographic observations. As mentioned, two files are accessible, both include
605 oceanographic variables observed at the standard depths (see supplementary Materials Part-2).

606 - *Original dataset: CNR_DIN_WMED_20042017_original.csv*: This is the original dataset with
607 flag variable for each of the following parameter: CTD salinity, nitrate, phosphate and silicate
608 from the primary quality control (detailed in section 3.1).

609 - *Adjusted dataset: CNR_DIN_WMED_20042017_adjusted.csv*: This is the product after
610 primary quality control and after applying the adjustment factors from the secondary quality
611 control. Recommendations of section 4.4 are included, as well as quality flags.

612 **Author contribution:** MB, MA, SL, JC and KS substantially contributed to write the manuscript. SC,
613 GC and FA run the chemical analysis and contributed to the manuscript. MB coordinated the technical
614 aspects of most of the cruises. SC, GC, FA, AR, BP contributed in specific part of the manuscript.

615 **Acknowledgements.** The data have been collected in the framework of several of national and
616 European projects, e.g.: KM3NeT, EU GA #011937; SESAME, EU GA #GOCE-036949; PERSEUS,
617 EU GA #287600; OCEAN-CERTAIN, EU GA #603773; COMMON SENSE, EU GA #228344;
618 EUROFLEETS, EU GA #228344; EUROFLEETS2, EU GA # 312762; JERICO, EU GA #262584;
619 the Italian PRIN 2007 program “Tyrrhenian Seamounts ecosystems”, and the Italian RITMARE
620 Flagship Project, both funded by the Italian Ministry of University and Research. We thank Sarah
621 Jutterström from the Swedish Environmental Research institute for the invaluable help in Quality
622 Control discussions. We would like to express our appreciation to the INOCEN laboratory team at
623 IEO for their help and collaboration during MB’s stay there. The authors are deeply indebted to all
624 investigators and analysts who contributed to data collection at sea during so many years, as well as to
625 the PIs of the cruises (S. Aliani, M. Astraldi, M. Azzaro, M. Dibitto, G. P. Gasparini, A. Griffa, J.

626 Haun, L. Jullion, G. La Spada, E. Manini, A. Perilli, C. Santinelli, S. Sparnocchia), the captains and
627 the crews for allowing the collection of this enormous dataset; without them, this work would not have
628 been possible.

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646 **References**

647 Aoyama, M., Woodward, E., Malcolm, S., Bakker, K., Becker, S., Björkman, K., Daniel, A.,
648 Mahaffey, C., Murata, A., Naik, H., Tanhua, T., Rho, T., Roman, R. and Sloyan, B.: Comparability of
649 oceanic nutrient data. Poster Cluster Community Whitepaper, CLIVAR Open Science Conference on
650 "Charting the course for climate and ocean research", 18-25 September 2016, Qingdao (China), 12 pp.,
651 <http://hdl.handle.net/10261/17137>, 2016.

652 Aoyama, Michio.: Global certified-reference-material-or reference-material-scaled nutrient gridded
653 dataset GND13. Earth System Science Data, 12, 4, 487-499, 2020 [https://doi.org/10.5194/essd-12-487-](https://doi.org/10.5194/essd-12-487-2020)
654 [2020, 2020](https://doi.org/10.5194/essd-12-487-2020).

Formattato: Tipo di carattere: 11 pt

Formattato: Francese (Francia)

655

656 Becker, S., Aoyama, M., Woodward, E.M.S., Bakker, K., Coverly, S., Mahaffey, C., and Tanhua, T.:
657 GO-SHIP Repeat Hydrography Nutrient Manual: The precise and accurate determination of dissolved
658 inorganic nutrients in seawater, using Continuous Flow Analysis methods, In: The GO-SHIP Repeat
659 Hydrography Manual: A Collection of Expert Reports and Guidelines, 56 , 2019,
660 <http://dx.doi.org/10.25607/OBP-555>, 2019.

661 Belgacem, M., Chiggiato, J., Borghini, M., Pavoni, B., Cerrati, G., Acri, F; Cozzi, S., Ribotti, A.,
662 Álvarez, M., Lauvset, S. K., Schroeder, K.: Quality controlled dataset of dissolved inorganic nutrients
663 in the western Mediterranean Sea (2004-2017) from R/V oceanographic cruises. PANGAEA,
664 <https://doi.pangaea.de/10.1594/PANGAEA.904172>, 2019.

666 Bethoux, J. P.: Oxygen consumption, new production, vertical advection and environmental evolution
667 in the Mediterranean Sea, Deep Sea Research, Part A, Oceanographic Research Papers, 36(5), 769–
668 781, doi:10.1016/0198-0149(89)90150-7, 1989.

669 Bethoux, J. P., Morin, P., Madec, C. and Gentili, B.: Phosphorus and nitrogen behaviour in the
670 Mediterranean Sea, Deep Sea Research, Part A, Oceanographic Research Paper, 39(9), 1641–1654,
671 doi:10.1016/0198-0149(92)90053-V, 1992.

672 Bethoux, J. P., Gentili, B., Morin, P., Nicolas, E., Pierre, C. and Ruiz-Pino, D.: The Mediterranean
673 Sea : a miniature ocean for climatic and environmental studies and a key for the climatic functioning of
674 the North Atlantic, Progress in Oceanography, 44, 131–146, 1999.

675 Béthoux, J. P., Morin, P., Chaumery, C., Connan, O., Gentili, B. and Ruiz-Pino, D.: Nutrients in the
676 Mediterranean Sea, mass balance and statistical analysis of concentrations with respect to
677 environmental change, Marine Chemistry , 63(1–2), 155–169, doi:10.1016/S0304-4203(98)00059-0,
678 1998.

679 Béthoux, J. P., Morin, P. and Ruiz-Pino, D. P.: Temporal trends in nutrient ratios: Chemical evidence
680 of Mediterranean ecosystem changes driven by human activity, Deep Sea Research Part II Topical

- 681 Studies in Oceanography, 49(11), 2007–2016, doi:10.1016/S0967-0645(02)00024-3, 2002.
- 682 Boyd, P. W.: Beyond ocean acidification, *Nature Geoscience*, 4(5), 273–274, doi:10.1038/ngeo1150,
683 2011.
- 684 Coppola, L., Raimbault, P., Mortier, L., and Testor, P.: Monitoring the environment in the
685 northwestern Mediterranean Sea, *Eos*, 100, <https://doi.org/10.1029/2019EO125951>, 2019.
- 686 Dickson, A. G., Afghan, J. D. and Anderson, G. C.: Reference materials for oceanic CO₂ analysis: A
687 method for the certification of total alkalinity, *Marine Chemistry*, 80(2–3), 185–197,
688 doi:10.1016/S0304-4203(02)00133-0, 2003.
- 689 [Dore, J. E., Houlihan, T., Hebel, D. V., Tien, G., Tupas, L., Karl, D. M.: Freezing as a method of
690 sample preservation for the analysis of dissolved inorganic nutrients in seawater, *Marine
691 Chemistry*, 53\(3–4\), 173-185, 1996.](#)
- 692 Fichaut, M., Garcia, M. J., Giorgetti, A., Iona, A., Kuznetsov, A., Rixen, M. and Group, M.:
693 MEDAR/MEDATLAS 2002: A Mediterranean and Black Sea database for operational oceanography,
694 Elsevier Oceanography Series, 69, 645–648, doi:10.1016/S0422-9894(03)80107-1, 2003.
- 695 Giorgetti, A., Partescano, E., Barth, A., Buga, L., Gatti, J., Giorgi, G., Iona A., Lipizer, M.,
696 Holdsworth, N., Larsen, M.M., Schaap, D., Vinci, M., Wenzel, M.: EMODnet Chemistry Spatial Data
697 Infrastructure for marine observations and related information. *Ocean & Coastal Management*, 166, 9–
698 17, 2018.
- 699 Giorgi, F.: Climate change hot-spots, *Geophysical Research Letters*, 33(8), 1–4,
700 doi:10.1029/2006GL025734, 2006.
- 701 Gouretski, V. V. and Jancke, K.: Systematic errors as the cause for an apparent deep water property
702 variability: Global analysis of the WOCE and historical hydrographic data, *Progress in Oceanography*,
703 48(4), 337–402, doi:10.1016/S0079-6611(00)00049-5, 2000.
- 704 Grasshoff, K., Kremling K., Ehrhardt M.: *Methods of seawater analysis* (3rd ed.), Weinheim
705 Press, WILEY-VCH, 203-273, 1999.
- 706
- 707 Hansen, H. P. and Koroleff, F.: Determination of nutrients, *Methods of Seawater Analysis*, 159–228,
708 1999.
- 709 Hoppema, M., Velo, A., van Heuven, S., Tanhua, T., Key, R. M., Lin, X., Bakker, D. C. E., Perez, F.
710 F., Ríos, A. F., Lo Monaco, C., Sabine, C. L., Álvarez, M. and Bellerby, R. G. J.: Consistency of
711 cruise data of the CARINA database in the Atlantic sector of the Southern Ocean, *Earth System
712 Science Data*, 1(1), 63–75, doi:10.5194/essd-1-63-2009, 2009.
- 713 Hydes, D. J., Aoyama, M., Aminot, A., Bakker, K., Becker, S., Coverly, S., Daniel, A., Dickson,
714 A. G., Grosso, O., Kerouel, R., van Ooijen, J., Sato, K., Tanhua, T., Woodward, E. M. S. and
715 Zhang, J. Z.: Determination of Dissolved Nutrients (N, P, SI) in Seawater With High Precision and
716 Inter-Comparability Using Gas-Segmented Continuous Flow Analysers. In: *The GO-SHIP Repeat
717 Hydrography Manual: A Collection of Expert Reports and Guidelines. Version 1.* (eds Hood, E.M.,

Formattato: SpazioDopo: 0 pt,
Controlla righe isolate, Regola lo spazio
tra testo asiatico e in alfabeto latino,
Regola lo spazio tra caratteri asiatici e
numeri

718 C.L. Sabine, and B.M. Sloyan). IOCCP Report Number 14, ICPO Publication Series Number 134. 87
719 pp., <http://dx.doi.org/10.25607/OBP-555>, 2010.

720

721 ~~Dore, J. E., Houlihan, T., Hebel, D. V., Tien, G., Tupas, L., Karl, D. M.: Freezing as a method of~~
722 ~~sample preservation for the analysis of dissolved inorganic nutrients in seawater, *Marine*~~
723 ~~*Chemistry*, 53(3-4), 173-185, 1996.~~

724 Johnson, G. C., Robbins, P. E. and Hufford, G. E.: Systematic adjustments of hydrographic sections
725 for internal consistency, *Journal of Atmospheric Oceanic Technology*, 18(7), 1234–1244,
726 doi:10.1175/1520-0426(2001)018<1234:SAOHSF>2.0.CO;2, 2001.

727 Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero,
728 F. J., Mordy, C. and Peng, T. H.: A global ocean carbon climatology: Results from Global Data
729 Analysis Project (GLODAP), *Global Biogeochem. Cycles*, 18(4), 1–23, doi:10.1029/2004GB002247,
730 2004.

731 Lauvset, S. K. and Tanhua, T.: A toolbox for secondary quality control on ocean chemistry and
732 hydrographic data, *Limnology and Oceanography Methods*, 13(11), 601–608,
733 doi:10.1002/lom3.10050, 2015.

734 Lazzari, P., Solidoro, C., Salon, S. and Bolzon, G.: Spatial variability of phosphate and nitrate in the
735 Mediterranean Sea: A modeling approach, *Deep Sea Research Part I*, 108, 39–52,
736 doi:10.1016/j.dsr.2015.12.006, 2016.

737 Lejeusne, C., Chevaldonné, P., Pergent-Martini, C., Boudouresque, C. F. and Pérez, T.: Climate
738 change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea, *Trends in*
739 *Ecology and Evolution*, 25(4), 250–260, doi:10.1016/j.tree.2009.10.009, 2010.

740 Manca, B., Burca, M., Giorgetti, A., Coatanoan, C., Garcia, M. J., & Iona, A. : Physical and
741 biochemical averaged vertical profiles in the Mediterranean regions: an important tool to trace the
742 climatology of water masses and to validate incoming data from operational oceanography. *Journal of*
743 *Marine Systems*, 48(1-4), 83-116, 2004.

744 Martín Míguez, B., Novellino, A., Vinci, M., Claus, S., Calewaert, J. B., Vallius, H., Schmitt, T.,
745 Pititto, P., Giorgetti, A., Askew, N., Iona, S., Schaap, D., Pinardi, N., Harpham, Q., Kater, B.J.,
746 Populus, J., She, J., Vasilev Palazov, A., McMeel, O., Oset, P., Lear, D., Manzella, G.M.R., Gorringer,
747 P., Simoncelli, S., Larkin, K., Holdsworth, N., Dimitrios_Arvanitidis C., Molina-Jack M.E., Chaves-
748 Montero M.D.M. , Herman, P.M.J., and Hernandez F.: The European marine observation and data
749 network (EMODnet): visions and roles of the gateway to marine data in Europe. *Frontiers in Marine*
750 *Science*, 6, (2019).

751 Moon, J., Lee, K., Tanhua, T., Kress, N. and Kim, I.: Temporal nutrient dynamics in the
752 Mediterranean Sea in response to anthropogenic inputs, , 5243–5251,
753 doi:10.1002/2016GL068788.Received, 2016.

754 Muniz, K., Cruzado, A., Ruiz De Villa, C. and Villa, C. R. De: Statistical analysis of nutrient data
755 quality (nitrate and phosphate), applied to useful predictor models in the northwestern Mediterranean

756 Sea, Methodology, 17, 221–231, 2001.

757 Olsen, A., Key, R. M., Heuven, S. Van, Lauvset, S. K., Velo, A., Lin, X., Schirnack, C., Kozyr, A.,
758 Tanhua, T., Hoppema, M. and Jutterström, S.: The Global Ocean Data Analysis Project version 2 (
759 GLODAPv2) – an internally consistent data product for the world ocean, , 297–323,
760 doi:10.5194/essd-8-297-2016, 2016.

761 Olsen, A., Lange, N., Key,R., Tanhua, T., Alvarez, M. et al.: GLODAPv2.2019 -an update of
762 GLODAPv2. Earth Syst. Sci. Data, [2019](#), 11 (3), pp.1437 - 1461. [ff10.5194/essd-11-1437-2019ff.](#)
763 [ffhal-02315662](#), [2019](#).

764

765 Pasqueron, O., Fommervault, D., Migon, C., Ortenzio, F. D., Ribera, M. and Coppola, L.: [Deep-Sea](#)
766 [Research](#) – Temporal variability of nutrient concentrations in the northwestern Mediterranean sea (
767 DYFAMED time-series station), Deep. Res. Part I, 100, 1–12, doi:10.1016/j.dsr.2015.02.006, 2015.

768 Powley, H. R., Krom, M. D. and Van Cappellen, P.: Phosphorus and nitrogen trajectories in the
769 Mediterranean Sea (1950–2030): Diagnosing basin-wide anthropogenic nutrient enrichment, Progress
770 in Oceanography, 162, 257–270, doi:10.1016/j.pocean.2018.03.003, 2018.

771 Pujó-Pay, M., Conan, P., Oriol, L., Cornet-Barthaux, V., Falco, C., Ghiglione, J. F., Goyet, C.,
772 Moutin, T. and Prieur, L.: Integrated survey of elemental stoichiometry (C, N, P) from the western to
773 eastern Mediterranean Sea, Biogeosciences, 8(4), 883–899, doi:10.5194/bg-8-883-2011, 2011.

774 Sabine, C. L., Hoppema, M., Key, R. M., Tilbrook, B., Van Heuven, S., Lo Monaco, C., Metzl, N.,
775 Ishii, M., Murata, A. and Musielewicz, S.: Assessing the internal consistency of the CARINA data
776 base in the Pacific sector of the Southern Ocean, Earth System Science Data Discussions, 2(2), 195–
777 204, doi:10.5194/essd-2-195-2010, 2010.

778 Schroeder, K., Tanhua, T., Bryden, H., Alvarez, M., Chiggiato, J. and Aracri, S.: Mediterranean Sea
779 Ship-based Hydrographic Investigations Program (Med-SHIP), Oceanography, 28(3), 12–15,
780 doi:10.5670/oceanog.2015.71, 2015.

781 Schroeder, K., Chiggiato, J., Bryden, H. L., Borghini, M. and Ben Ismail, S.: Abrupt climate shift in
782 the Western Mediterranean Sea, Scientific Reports, 1–7, doi:10.1038/srep23009, 2016.

783 Segura-Noguera, M., Cruzado, A. and Blasco, D.: The biogeochemistry of nutrients, dissolved oxygen
784 and chlorophyll a in the Catalan Sea (NW Mediterranean Sea), Sci. Mar., 80(S1), 39–56,
785 doi:10.3989/scimar.04309.20a, 2016.

786 Segura-Noguera, M., Cruzado, A., & Blasco, D.: Nutrient preservation, analysis precision and quality
787 control of an oceanographic database of inorganic nutrients, dissolved oxygen and chlorophyll a from
788 the NW Mediterranean Sea. Scientia Marina, 75(2), 321-339, 2011.

789 Suzuki, T., Ishii, M., Aoyama, A., Christian, J. R., Enyo, K., Kawano, T., Key, R. M., Kosugi, N.,
790 Kozyr, A., Miller, L. A., Murata, A., Nakano, T., Ono, T., Saino, T., Sasaki, K., Sasano, D., Takatani,
791 Y., Wakita, M., and Sabine, C. L.: PACIFICA Data Synthesis Project, ORNL/CDIAC-159, NDP-092,

792 Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U. S. Department of
793 Energy, Oak Ridge, Tennessee, 2013.

794 | Tanhua, T.: Hydrochemistry of water samples during MedSHIP cruise Talpro. PANGAEA,
795 <https://doi.org/10.1594/PANGAEA.902293>, 2019.

796 Tanhua, T.: Matlab Toolbox to Perform Secondary Quality Control (2nd QC) on Hydrographic Data,
797 ORNL CDIAAC-158. Carbon Dioxide Inf. Anal. Center, Oak Ridge Natl. Lab. U.S. Dep. Energy, Oak
798 Ridge, Tennessee, 158, doi:10.3334/CDIAC/otg.CDIAC_158, 2010a.

799 | Tanhua, T., Brown, P. J. and Key, R. M.: CARINA : ~~nutrient~~-Nutrient data in the Atlantic Ocean,
800 Earth Science Data, 1, 7–24, doi:10.3334/CDIAC/otg.CARINA.ATL.V1.0, 2009.

801 Tanhua, T., Heuven, S. van, Key, R. M., Velo, A., Olsen, A. and Schimick, C.: Quality control
802 procedures and methods of the CARINA database, Earth System Science Data, 2, 35–49, 2010b.

803 Tanhua, T., Hainbucher, D., Schroeder, K., Cardin, V., Álvarez, M. and Civitarese, G.: The
804 Mediterranean Sea system: A review and an introduction to the special issue, Ocean Science, 9(5),
805 789–803, doi:10.5194/os-9-789-2013, 2013.

806 Testor, P., Bosse, A., Houpert, L., Margirier, F., Mortier, L., Legoff, H., Dausse, D., Labaste, M.,
807 Karstensen, J., Hayes, D., Olita, A., Ribotti, A., Schroeder, K., Chiggiato, J., Onken, R., Heslop, E.,
808 Moure, B., D’ortenzio, F., Mayot, N., Lavigne, H., de Fommervault, O., Coppola, L., Prieur, L.,
809 Taillandier, V., Durrieu de Madron, X., Bourrin, F., Many, G., Damien, P., Estournel, C., Marsaleix,
810 P., Taupier-Letage, I., Raimbault, P., Waldman, R., Bouin, M. N., Giordani, H., Caniaux, G., Somot,
811 S., Ducrocq, V. and Conan, P.: Multiscale Observations of Deep Convection in the Northwestern
812 Mediterranean Sea During Winter 2012–2013 Using Multiple Platforms, Journal of Geophysical
813 Research: Oceans, 123(3), 1745–1776, doi:10.1002/2016JC012671, 2018.

814 | Tintoré, J., Pinardi, N., Alvarez Fanjul, E., Balbin, R., Bozzano, R., Ferrarin, C.,... ~~&~~and Clementi, E.:
815 Challenges for Sustained Observing and Forecasting Systems in the Mediterranean Sea. Frontiers in
816 Marine Science, 6, 568, (2019).

817

Formattato: Italiano (Italia)

818 **Figure Captions**

819 **Figure 1.** Map of the Western Mediterranean Sea showing the biogeochemical stations (in blue) and
820 the five reference cruise stations (in red).

821 **Figure 2.** Overview of the reference cruise spatial coverage and vertical distributions of the inorganic
822 nutrients. Top left: geographical distribution map, top right: vertical profiles of nitrate in $\mu\text{mol kg}^{-1}$,
823 bottom left: vertical profiles of phosphate in $\mu\text{mol kg}^{-1}$, bottom right: vertical profiles of silicate in
824 $\mu\text{mol kg}^{-1}$.

825 **Figure 3.** Scatter plots of (A.) phosphate vs nitrate (in $\mu\text{mol kg}^{-1}$) and (B.) silicate vs. nitrate (in μmol
826 kg^{-1}). Data that have been flagged as “questionable” (flag=3) are in red, the colour bar indicates the
827 pressure (in dbar). The black lines represent the best linear fit between the two parameters, and the
828 corresponding equations and r^2 values are shown on each plot. Average resulting N:P ratio is 20.87,
829 average resulting N:Si ratio is 1.05 (whole depth).

830 **Figure 4.** An example of the calculated offset for silicate between cruise 48UR20131015 and cruise
831 29AJ2016818 (reference cruise). Above: location of the stations being part of the crossover and
832 statistics. Bottom left: vertical profiles of silicate data in ($\mu\text{mol kg}^{-1}$) of the two cruises that fall within
833 the minimum distance criteria (the crossing region), below 1000 dbar. Bottom right: vertical plot of
834 the difference between both cruises (dotted black line) with standard deviations (dashed black lines)
835 and the weighted average of the offset (solid red line) with the weighted standard deviations (dotted
836 red line).

837 **Figure 5.** Results of the crossover analysis for nitrate, before (grey) and after adjustment (blue). Error
838 bars indicate the standard deviation of the absolute weighted offset. The dashed lines indicate the
839 accuracy limit 2% for an adjustment to be recommended.

840 **Figure 6.** The same as Fig. 5 but for phosphate.

841 **Figure 7.** The same as Fig. 5 but for silicate.

842 **Figure 8.** Dataset comparison before (black) and after (blue) adjustment, showing vertical profiles of
843 (A.) nitrate (in $\mu\text{mol kg}^{-1}$), (B.) phosphate (in $\mu\text{mol kg}^{-1}$) and (C.) silicate (in $\mu\text{mol kg}^{-1}$). Scatter plots
844 of the adjusted data from all depths after 1st and 2nd quality control for (D.) phosphate vs nitrate (in
845 $\mu\text{mol kg}^{-1}$) and (E.) silicate vs. nitrate (in $\mu\text{mol kg}^{-1}$). The black lines represent the best linear fit
846 between the two parameters, and the corresponding equations and r^2 values are shown on each plot.
847 Average resulting N:P ratio is 22.09, average resulting N:Si ratio is 0.94 (whole depth).

848 **Figure 9.** Vertical profiles of the inorganic nutrients in the dataset after adjustments and spatial
849 coverage of each cruise (reference to cruise ID is above each map). The whole WMED adjusted
850 product is shown in black while the data of each individual cruise are shown in blue (flag=2) and
851 green (flag=3).

852 **Figure 10.** RMSE regional averages of water mass properties computed between the new adjusted
853 product and MEDAR/Medatlas climatology for nitrate (A.), phosphate (B.) and silicate (C.).

854 **Table captions**

855 **Table 1a.** Cruise summary table and parameters listed with number of stations and samples. Cruises
856 were identified with an ID number and expedition code ('EXPOCODE' of format
857 AABBYYYMMDD with AA: country code, BB: ship code, YYYY: year, MM: month, DD: day
858 indicative of cruise starting day).

859 ~~**Table 1b.** [Data Collection Metadata and Sources](#) and links to the reports (accessed June 2020).~~

860 **Table 2.** Cruise summary table of the reference cruises collection used in the secondary quality
861 control, collected from 2001 to 2016.

862 **Table 3.** WOCE flags used in the original data product and in the adjusted product.

863 **Table 4.** Average and Standard deviations of nitrate, phosphate and silicate measurements by cruise
864 and for each region with number of samples deeper than 1000db included in the 2nd QC. Average
865 storage time: the minimum storage time defined as time difference between the cruise ending day and
866 the 1st day of the laboratory analysis.

867 **Table 5.** Summary of the suggested adjustment for nitrate, phosphate and silicate resulting from the
868 crossover analysis. Adjustments for inorganic nutrient are multiplicative. NA: denotes not adjusted,
869 i.e. data of cruises that could not be used in the crossover analysis, because of the lack of stations or
870 data are outside the spatial coverage of reference cruises.

871 **Table 6.** Secondary QC toolbox results: improvements of the weighted mean of absolute offset per
872 cruise of unadjusted and adjusted data; (n) is the number of crossovers per cruise. The numbers in red
873 (less than 1) indicate that the cruise data are lower than the reference cruises. NA: not adjusted.

874 **Table 7.** Water mass properties and regional average concentrations of inorganic nutrients:
875 comparison between the new adjusted product and the MEDAR/Medatlas climatology (with standard
876 deviations and number of observations in brackets).


Formattato: Tipo di carattere:
Grassetto

Formattato: Allineato a sinistra,
Livello 1, Interlinea multipla 1,15 ri

Formattato: Tipo di carattere: Non
Grassetto

Formattato: Tipo di carattere:
Grassetto, Inglese (Stati Uniti)

877 **Figure 1**



878

879

880

881

882

883

884


885

886

887

888

889 **Figure 2**



890

891

892

893

894

895

896

897

898

899


900

901

902

903

904 **Figure 3**



905

906


907

908

909

910

911 **Figure 4**




912

913


914

915



916

917 **Figure 5**



918

919


920

921

922

923
924
925
926
927
928
929
930
931
932
933


Figure 6



934
935
936
937
938

939
940
941
942
943
944
945
946
947
948
949


Figure 7



950
951
952
953
954

955
956
957
958
959
960
961
962
963
964
965

Figure 8



966
967
968
969

970

971

972

973

974

975


976


977


978

979

980 **Figure 9**









983


984



985

986





989

990

991

992

993

994

995

996

997

Table 1a

Cruise ID (#)	Common Name	EXPOCODE	Research vessel (RV)	Date Start/End	Stations	Samples Nitrate ₃	Samples Phosphate ₄	Samples Silicate ₂	Maximum bottom depth (m)	Chief scientist
1	TRENDS2004/MEDGOOS8leg2	48UR20040526	Urania	26 MAY - 14 JUN 2004	36	255	253	255	3499	M. Borghini
2	MEDGOOS9	48UR20041006	Urania	6 - 25 OCT 2004	68	627	626	627	3610	M. Borghini
3	MEDOCC05/MFSTEP2	48UR20050412	Urania	12 APR - 16 MAY 2005	68	828	828	828	3598	M. Borghini
4	MEDGOOS10	48UR20050529	Urania	29 MAY - 10 JUN 2005	36	577	577	577	3505	A. Perilli
5	MEDGOOS11	48UR20051116	Urania	16 NOV - 3 DEC 2005	14	143	143	143	2810	A. Perilli, M. Borghini, M. Dibitetto
6	MEDOCC06	48UR20060608	Urania	8 JUN - 3 JUL 2006	66	787	785	787	2881	M. Borghini
7	SIRENA06	06A420060720	NRV Alliance	20 JUL - 6 AUG 2006	35	208	208	209	1854	J. Haun
8	MEDGOOS13/MEDBIO06	48UR20060928	Urania	28 SEP - 8 NOV 2006	37	519	520	520	2862	A. Ribotti
9	MEDOCC07	48UR20071005	Urania	5 - 29 OCT 2007	71	977	977	979	3497	A. Perilli, <u>M. Borghini</u> , <u>A. Ribotti</u>
10	SESAMEI4	48UR20080318	Urania	18 MAR - 7 APR 2008	11	164	164	164	2882	C. Santinelli
11	SESAMEIT5	48UR20080905	Urania	5 - 16 SEP 2008	12	74	74	74	536	S. Sparnocchia, G.P. Gasparini, M. Borghini
12	MEDCO08	48UR20081103	Urania	3 - 24 NOV 2008	24	342	350	348	2880	A. Ribotti
13	TYRRMOUNTS	48UR20090508	Urania	8 MAY - 3 JUN 2009	41	430	441	440	2559	G.P. Gasparini
14	BIOFUN010	48UR20100430	Urania	30 APR - 17 MAY 2010	26	405	405	405	3540	E. Manini, S. Aliani
15	VENUS1	48UR20100731	Urania	31 JUL - 25 AUG 2010	32	431	432	428	3544	G.P. Gasparini, M. Borghini
16	BONSIC2010	48UR20101123	Urania	23 NOV - 9 DEC 2010	18	144	143	143	3540	A. Ribotti
17	EUROFLEET11	48UR20110421	Urania	21 APR - 8 MAY 2011	28	277	275	277	3540	G.P. Gasparini, M. Borghini
18	BONIFACIO2011	48UR20111109	Urania	9 - 23 NOV 2011	13	180	180	181	3541	A. Ribotti, G. La Spada, M. Borghini
19	TOSCA2011	48MG20111210	Maria Grazia	10 - 20 DEC 2011	21	310	310	309	2728	M. Borghini
20	ICHNUSSA12	48UR20120111	Urania	11 - 27 JAN 2012	21	353	352	323	3551	A. Ribotti
21	EUROFLEET2012	48UR20121108	Urania	8 - 26 NOV 2012	53	429	434	434	2633	M. Borghini
22	ICHNUSSA13	48UR20131015	Urania	15 - 29 OCT 2013	37	405	404	405	3540	A. Ribotti
23	OCEANCERTAIN15	48QL20150804	Minerva Uno	4 - 29 AUG 2015	71	531	531	531	3513	J. Chiggiato
24	ICHNUSSA17/INFRAOCE17	48QL20171023	Minerva Uno	23 OCT - 28 NOV 2017	31	251	254	254	3536	A. Ribotti, S. Sparnocchia, M. Borghini

Formattato: Inglese (Regno Unito)

Table 1b

<u>Cruise ID (#)</u>	<u>Expedition original Name</u>	<u>PIs/ Chief scientist</u>	<u>Specific link* (accessed June 2020)</u>
<u>1</u>	<u>TRENDS2004/ MEDGOOS8leg2</u>	<u>M. Borghini</u>	<u>https://isramar.ocean.org.il/perseus_data/CruiseInfo.aspx?cruiseid=5821</u> <u>https://isramar.ocean.org.il/perseus_data/CruiseInfo.aspx?cruiseid=4935</u>
<u>2</u>	<u>MEDGOOS9</u>	<u>M. Borghini</u>	<u>Report submission in progress</u> <u>https://isramar.ocean.org.il/perseus_data/CruiseInfo.aspx?cruiseid=5823</u> <u>https://doi.org/10.17882/70340</u>
<u>3</u>	<u>MEDOCC05/ MFSTEP2</u>	<u>M. Borghini</u>	<u>http://ricerca.ismar.cnr.it/CRUISE_REPORTS/2005/URANIA_MEDOCC05.pdf</u> <u>https://isramar.ocean.org.il/perseus_data/CruiseInfo.aspx?cruiseid=4936</u>
<u>4</u>	<u>MEDGOOS10</u>	<u>A. Perilli</u>	<u>http://www.seaforecast.cnr.it/it/observation_it.htm</u> <u>https://doi.org/10.17882/70340</u>
<u>5</u>	<u>MEDGOOS11</u>	<u>A. Perilli, M. Borghini, M. Dibitto</u>	<u>http://ricerca.ismar.cnr.it/CRUISE_REPORTS/2005/URANIA_MEDGOOS11_05_REP.pdf</u> <u>https://doi.org/10.17882/70340</u>
<u>6</u>	<u>MEDOCC06</u>	<u>M. Borghini</u>	<u>http://www.seaforecast.cnr.it/reports/Medocc06CR.pdf</u> <u>https://seadata.bsh.de/Cgi-csr/retrieve_sdn2/viewReport.pl?csref=20106010</u>
<u>7</u>	<u>SIRENA06</u>	<u>J. Haun</u>	<u>Report submission in progress</u>
<u>8</u>	<u>MEDGOOS13/ MEDBIO06</u>	<u>A. Ribotti</u>	<u>http://www.seaforecast.cnr.it/reports/Mebio06-Medg13_CR.pdf</u> <u>https://doi.org/10.17882/70340</u>
<u>9</u>	<u>MEDOCC07</u>	<u>A. Perilli, M. Borghini, A. Ribotti</u>	<u>http://www.seaforecast.cnr.it/reports/Medocc07-MedCo07_Rapp.pdf</u> <u>https://isramar.ocean.org.il/perseus_data/CruiseInfo.aspx?cruiseid=5146</u>
<u>10</u>	<u>SESAMEIt4</u>	<u>C. Santinelli</u>	<u>https://isramar.ocean.org.il/perseus_data/CruiseInfo.aspx?cruiseid=5148</u> <u>https://emodnet-chemistry.maris.nl/search/details.php?step=0012004~0022017~0153~057104001~058tdin.ntra.phos.slca~00445~0056~00617~00734~0541&count=3592&page=1000&sort=0&header=no</u>
<u>11</u>	<u>SESAMEIT5</u>	<u>S. Spamocchia, G.P. Gasparini, M. Borghini</u>	<u>https://isramar.ocean.org.il/perseus_data/CruiseInfo.aspx?cruiseid=5147</u>
<u>12</u>	<u>MEDCO08</u>	<u>A. Ribotti</u>	<u>http://www.seaforecast.cnr.it/reports/MedCO08_Rapp.pdf</u>
<u>13</u>	<u>TYRRMOUNTS</u>	<u>G.P. Gasparini</u>	<u>Report submission in progress</u>
<u>14</u>	<u>BIOFUN010</u>	<u>E. Manini, S. Aliani</u>	<u>http://www.ismar.cnr.it/products/reports-campagne/2010-2019</u>
<u>15</u>	<u>VENUS1</u>	<u>G.P. Gasparini, M. Borghini</u>	<u>Report submission in progress</u>
<u>16</u>	<u>BONSIC2010</u>	<u>A. Ribotti</u>	<u>http://www.seaforecast.cnr.it/reports/Bonifacio2010Sic_Rapp.pdf</u>
<u>17</u>	<u>EUROFLEET11</u>	<u>G.P. Gasparini, M. Borghini</u>	<u>Report submission in progress</u>
<u>18</u>	<u>BONIFACIO2011</u>	<u>A. Ribotti, G. La Spada, M. Borghini</u>	<u>http://www.seaforecast.cnr.it/reports/Bonifacio2011_Rapp.pdf</u>
<u>19</u>	<u>TOSCA2011</u>	<u>M. Borghini</u>	<u>Report submission in progress</u>
<u>20</u>	<u>ICHNUSSA12</u>	<u>A. Ribotti</u>	<u>http://www.seaforecast.cnr.it/reports/Ichnussa2012_Rapp.pdf</u>
<u>21</u>	<u>EUROFLEET2012</u>	<u>M. Borghini</u>	<u>Report submission in progress</u>
<u>22</u>	<u>ICHNUSSA13</u>	<u>A. Ribotti</u>	<u>http://www.seaforecast.cnr.it/reports/Ichnussa2013_Rapp.pdf</u>
<u>23</u>	<u>OCEANCERTAIN15</u>	<u>J. Chiggiato</u>	<u>https://doi.pangaea.de/10.1594/PANGAEA.911046</u>
<u>24</u>	<u>ICHNUSSA17/ INFRAOCE17</u>	<u>A. Ribotti, S. Spamocchia, M. Borghini</u>	<u>Report submission in progress</u>

* The specific links are subjected to updates.

Table 2

Common name	EXPOCODE	Date Start/End	Stations	Nitrate ^{O3} Sample	Phosphate ^{O4} Sample	Silicate ^{O2} Sample	Source	Nutrient PI	Chief scientist
<i>M51/2</i>	06MT20011018	18 OCT - 11 NOV 2001	6	79	79	82	GLODAPv2	B. Schneider	W. Roether
<i>TRANSMED_LEGII</i>	48UR20070528	28 MAY- 12 JUN 2007	4	78	77	78	CARIMED (not yet available)	S. Cozzi, V. Ibello	M. Azzaro
<i>M84/3</i>	06MT20110405	5 - 28 APR 2011	20	339	343	-	GLODAPv2	G. Civitarese	T. Tanhua
<i>HOTMIX</i>	29AH20140426	26 APR- 31 MAY 2014	18	144	140	144	CARIMED (not yet available)	XA Álvarez- Salgado	J. Aristegui
<i>TALPro-2016</i>	29AJ20160818	18 - 28 AUG 2016	42	293	293	293	MedSHIP programme	L. Coppola	L. Jullion, K. Schroeder

Table 3

WOCE flag value	Interpretation in original dataset	Interpretation in adjusted product
2	Acceptable/ measured	Adjusted and acceptable
3	Questionable/not used	Adjusted and recommended questionable
9	not measured/no data	-

Table 4

Cruise ID	EXPCODE/ Region	Regional Avg Nitrate O_2 ($\mu\text{mol kg}^{-1}$)	std Nitrate O_2 ($\mu\text{mol kg}^{-1}$)	Regional Avg Phosphate O_2 ($\mu\text{mol kg}^{-1}$)	std Phosphate O_2 ($\mu\text{mol kg}^{-1}$)	Regional Avg Silicate O_2 ($\mu\text{mol kg}^{-1}$)	std Silicate O_2 ($\mu\text{mol kg}^{-1}$)	# samples	Avg storage (in days)
1	48UR20040526/		1.25		0.062		1.64	21	131
	DT1-Tyrrhenian North	6.07	1.32	0.26	0.065	6.92	1.83	16	
	DT3-Tyrrhenian South	7.03	0.51	0.31	0.02	7.66	0.53	5	
2	48UR20041006/		0.59		0.029		0.81	21	251
	DT1-Tyrrhenian North	7.68	0.53	0.41	0.031	8.74	0.75	15	
	DT3-Tyrrhenian South	8.17	0.60	0.41	0.025	9.31	0.87	6	
3	48UR20050412/		1.15		0.050		1.41	233	135
	DF2-Gulf of Lion	7.89	0.98	0.40	0.044	8.17	1.065	24	
	DF3-Liguro-Provençal	7.45	1.08	0.41	0.05	7.72	1.10	66	
	DS2-Balearic Sea	7.44	1.14	0.40	0.039	7.68	1.47	21	
	DF1-Algero-Provençal	7.87	1.16	0.41	0.043	8.88	1.96	42	
	DS3-Algerian West	7.7	0.816	0.39	0.048	8.14	0.941	23	
	DT1-Tyrrhenian North	6.57	1.065	0.36	0.047	7.41	1.15	21	
	DT3-Tyrrhenian South	6.52	1.12	0.36	0.05	7.56	1.42	22	
	DII-Sardinia Channel	7.22	1.065	0.40	0.04	8.08	1.11	14	
	4	48UR20050529/		1.13		0.057		1.08	
DS1-Alboran Sea		6.4	1.15	0.38	0.041	6.26	1.02	32	
DS3-Algerian West		7.6	1.13	0.41	0.06	7.33	0.99	73	
DS4-Algerian East		7.48	1.13	0.41	0.06	7.50	1.23	47	
DT1-Tyrrhenian North		7.24	0.44	0.42	0.03	7.91	0.56	16	
DT3-Tyrrhenian South		7.70	0.38	0.41	0.03	7.55	0.36	14	
DII-Sardinia Channel		7.58	1.08	0.43	0.049	7.42	0.82	23	
5		48UR20051116/		1.35		0.078		0.98	16
	DT1-Tyrrhenian North	5.68	1.26	0.19	0.08	6.30	0.92	10	
	DT3-Tyrrhenian South	6.71	1.51	0.20	0.06	6.86	1.065	5	
	DII-Sardinia Channel	6.29	0	0.26	0	7.53	0	1	
6	48UR20060608/		1.16		0.054		1.47	221	27
	DF2-Gulf of Lion	7.69	1.02	0.42	0.04	7.089	1.04	27	
	DF3-Liguro-Provençal	8.08	0.78	0.43	0.04	7.41	1.21	35	
	DS2-Balearic Sea	8.06	0.9	0.43	0.03	7.07	1.18	30	
	DF1-Algero-Provençal	7.97	1.16	0.44	0.05	7.34	1.32	61	
	DS3-Algerian West	8.39	0.9	0.42	0.03	8.5	2	28	
	DT3-Tyrrhenian South	6.39	1.28	0.36	0.06	6.86	1.7	26	
	DII-Sardinia Channel	8.04	0.85	0.43	0.04	7.77	1.25	14	
	06A420060720		-		-		-	-	
8	48UR20060928/		0.71		0.036		0.76	179	606
	DS2-Balearic Sea	7.97	0.17	0.33	0.017	7.84	0.27	4	
	DF1-Algero-Provençal	8.17	0.22	0.33	0.026	8.11	0.3	22	
	DS1-Alboran Sea	8.2	0.14	0.35	0.02	8.59	0.35	47	
	DS3-Algerian West	7.93	0.89	0.33	0.03	8.09	0.91	70	
	DS4-Algerian East	7.98	0.68	0.34	0.04	8.01	0.7	28	
	DT3-Tyrrhenian South	6.2	1.51	0.28	0.04	6.71	1.45	3	
	DII-Sardinia Channel	7.66	0.6	0.28	0.02	8.00	0.49	5	
	9	48UR20071005/		0.89		0.040		0.86	
DF2-Gulf of Lion		8.41	0.08	0.31	0.01	7.43	0.02	4	
DF3-Liguro-Provençal		8.17	1.08	0.31	0.03	7.64	1.08	81	
DS2-Balearic Sea		8.17	0.43	0.31	0.02	7.58	0.39	29	
DF1-Algero-Provençal		8.33	0.6	0.32	0.03	7.79	0.69	82	
DS4-Algerian East		8.41	0.2	0.33	0.018	7.90	0.26	19	
DT1-Tyrrhenian North		7.83	0.41	0.28	0.03	8.26	0.55	26	
DT3-Tyrrhenian South		7.49	1.22	0.28	0.05	7.71	1.26	38	
DII-Sardinia Channel		7.92	1.05	0.33	0.02	8.26	0.41	23	
10		48UR20080318/		0.51		0.026		0.34	66
	DF2-Gulf of Lion	8.54	0.6	0.35	0.03	8.62	0.43	5	
	DS2-Balearic Sea	9.12	0.18	0.38	0.01	8.40	0.21	9	
	DF1-Algero-Provençal	9.02	0.36	0.38	0.03	8.65	0.25	15	
	DS3-Algerian West	8.93	0.46	0.36	0.01	8.69	0.35	20	
	DS4-Algerian East	8.43	0.25	0.38	0.02	8.32	0.22	10	
	DII-Sardinia Channel	7.62	0.6	0.34	0.03	8.49	0.36	3	
	11*	48UR20080905		-		-		-	-
12	48UR20081103/		1.11		0.077		0.10	110	536

	<i>DS1-Alboran Sea</i>	6.4	1.21	0.21	0.06	7.20	1.43	26	
	<i>DS3-Algerian West</i>	7.58	0.9	0.27	0.1	7.89	0.9	30	
	<i>DS4-Algerian East</i>	7.15	1.04	0.23	0.04	7.38	0.9	35	
	<i>DT3-Tyrrhenian South</i>	7.44	0.5	0.22	0.05	8.28	0.4	10	
	<i>DII-Sardinia Channel</i>	7.40	1.23	0.17	0.04	8.09	0.45	9	
13	48UR20090508/		1.41		0.051		1.42	88	164
	<i>DT1-Tyrrhenian North</i>	5.95	1.55	0.24	0.05	6.28	1.58	46	
	<i>DT3-Tyrrhenian South</i>	6.76	0.77	0.24	0.03	7.37	0.77	29	
	<i>DII-Sardinia Channel</i>	7.62	1.1	0.28	0.05	7.76	0.9	13	
14	48UR20100430/		1.06		0.036		1.03	159	213
	<i>DS2-Balearic Sea</i>	7.66	1.6	0.25	0.03	7.38	1.75	33	
	<i>DF1-Algero-Provençal</i>	8.43	0.29	0.26	0.03	8.06	0.31	61	
	<i>DS3-Algerian West</i>	8.5	0.14	0.26	0.03	8.25	0.3	26	
	<i>DT1-Tyrrhenian North</i>	6.88	0.8	0.23	0.022	7.17	0.77	11	
	<i>DT3-Tyrrhenian South</i>	6.38	1.35	0.22	0.01	6.76	1.56	7	
	<i>DII-Sardinia Channel</i>	7.71	0.87	0.23	0.02	7.80	0.74	21	
15	48UR20100731/		1.34		0.053		1.14	149	213
	<i>DS1-Alboran Sea</i>	7.30	1.18	0.29	0.05	7.21	1.11	25	
	<i>DS3-Algerian West</i>	7.67	1.15	0.28	0.045	7.24	1.16	54	
	<i>DS4-Algerian East</i>	7.38	0.89	0.29	0.03	7.00	0.78	29	
	<i>DT1-Tyrrhenian North</i>	7.66	0.96	0.29	0.05	7.89	1.07	10	
	<i>DT3-Tyrrhenian South</i>	5.4	0.67	0.22	0.01	5.52	1.56	30	
	<i>DII-Sardinia Channel</i>	4.92	0	0.20	0	5.55	0	1	
16	48UR20101123/		1.02		0.045		1.02	14	170
	<i>DT1-Tyrrhenian North</i>	6.34	0.87	0.27	0.02	6.12	0.87	8	
	<i>DT3-Tyrrhenian South</i>	5.43	1.02	0.22	0.04	5.08	0.9	6	
17	48UR20110421/		0.62		0.029		0.52	56	160
	<i>DT1-Tyrrhenian North</i>	7.77	0.45	0.28	0.02	8.11	0.35	21	
	<i>DT3-Tyrrhenian South</i>	7.76	0.7	0.28	0.03	8.017	0.6	35	
18	48UR20111109/		0.68		0.025		0.70	77	74
	<i>DF3-Liguro-Provençal</i>	6.68	0	0.33	0	6.26	0	1	
	<i>DF1-Algero-Provençal</i>	8.17	0.5	0.32	0.01	8.16	0.66	43	
	<i>DT1-Tyrrhenian North</i>	7.26	0.93	0.29	0.02	8.15	1.03	12	
	<i>DT3-Tyrrhenian South</i>	7.61	0.37	0.30	0.02	8.18	0.35	11	
	<i>DII-Sardinia Channel</i>	7.64	0.45	0.29	0.01	8.08	0.41	10	
19*	48MG20111210		-		-		-	-	38
20	48UR20120111/		0.97		0.051		0.26	152	317
	<i>DF1-Algero-Provençal</i>	8.45	0.49	0.31	0.039	7.91	0.53	23	
	<i>DT1-Tyrrhenian North</i>	7.67	0.83	0.27	0.02	8.29	0.8	30	
	<i>DT3-Tyrrhenian South</i>	7.65	1.06	0.31	0.06	8.03	1.26	69	
	<i>DII-Sardinia Channel</i>	7.65	0.96	0.31	0.03	7.86	0.78	30	
21*	48UR20121108		-		-		-	-	72
22	48UR20131015/		1.03		0.043		0.79	98	76
	<i>DF1-Algero-Provençal</i>	8.54	0.64	0.33	0.02	7.96	0.38	36	
	<i>DS4-Algerian East</i>	7.67	1.28	0.27	0.04	6.82	1.07	8	
	<i>DT1-Tyrrhenian North</i>	6.47	0.83	0.24	0.025	7.12	0.84	10	
	<i>DT3-Tyrrhenian South</i>	7.81	0.71	0.30	0.03	8.09	0.65	28	
	<i>DII-Sardinia Channel</i>	7.32	0.99	0.27	0.02	7.47	0.89	16	
23	48QL20150804/		0.84		0.038		0.85	94	30
	<i>DF3-Liguro-Provençal</i>	8.51	0.96	0.39	0.03	8.06	0.85	23	
	<i>DS2-Balearic Sea</i>	7.75	0.66	0.36	0.02	7.86	0.81	20	
	<i>DF1-Algero-Provençal</i>	7.9	0.59	0.37	0.03	8.34	0.68	23	
	<i>DS3-Algerian West</i>	7.84	0.67	0.36	0.02	7.75	0.68	6	
	<i>DT1-Tyrrhenian North</i>	7.92	0.61	0.37	0.02	8.75	0.4	8	
	<i>DT3-Tyrrhenian South</i>	7.23	0.75	0.34	0.025	8.2	0.94	13	
	<i>DII-Sardinia Channel</i>	6.30	0	0.25	0	5.36	0	1	
24	48QL20171023/		0.68		0.055		1.24	55	30
	<i>DF3-Liguro-Provençal</i>	6.63	0.41	0.40	0.05	10.76	1.07	3	
	<i>DF1-Algero-Provençal</i>	5.14	0.7	0.43	0.02	7.94	1.19	6	
	<i>DT1-Tyrrhenian North</i>	4.98	0.58	0.36	0.02	8.10	0.87	9	
	<i>DT3-Tyrrhenian South</i>	5.43	0.5	0.36	0.04	9.03	0.87	26	
	<i>DII-Sardinia Channel</i>	5.16	0.76	0.41	0.07	7.58	1.17	11	

(*): cruise not included in the 2ndQC (Section 4.)

in bold: the overall standard deviation by cruise; in normal font: regional standard deviation by cruise

Table 5

Cruise ID	EXPOCODE	Nitrate Θ_3 (x)	Phosphate Θ_4 (x)	Silicate Θ_2 (x)
1	48UR20040526	1.14	1.23	1.21
2	48UR20041006	0.98	0.9	1.06
3	48UR20050412	1.08	0.93	1.15
4	48UR20050529	1.04	0.85	1.183
5	48UR20051116	1.19	1.34	1.232
6	48UR20060608	1.05	0.86	1.261
7	06A420060720*	-	-	-
8	48UR20060928	1.03	1.14	1.1
9	48UR20071005	0.97	1.14	1.115
10	48UR20080318	0.94	1.09	1.02
11	48UR20080905*	-	-	-
12	48UR20081103	1.08	1.38	1.12
13	48UR20090508	1.05	1.33	1.15
14	48UR20100430	NA	1.34	1.123
15	48UR20100731	1.13	1.25	1.262
16	48UR20101123	1.15	1.29	1.28
17	48UR20110421	NA	1.25	1.12
18	48UR20111109	NA	1.14	1.09
19	48MG20111210*	-	-	-
20	48UR20120111	NA	1.17	1.08
21	48UR20121108*	-	-	-
22	48UR20131015	NA	1.17	1.11
23	48QL20150804	1.02	1.02	1.08
24	48QL20171023	1.34	0.98	1.06

(*) cruise not included in the 2nd QC (Section 4.)

Table 6

Cruise ID	EXPOCODE	Nitrate Θ_2 [%]			Phosphate Θ_4 [%]			Silicate Θ_2 [%]		
		<i>n</i>	<i>unadjusted</i>	<i>adjusted</i>	<i>n</i>	<i>unadjusted</i>	<i>adjusted</i>	<i>n</i>	<i>unadjusted</i>	<i>adjusted</i>
1	48UR20040526	2	0.86	0.98	2	0.77	0.95	1	0.79	0.96
2	48UR20041006	2	1.02	1.00	2	1.10	0.99	1	0.94	0.99
3	48UR20050412	5	0.92	0.99	5	1.07	1.00	4	0.85	0.98
4	48UR20050529	5	0.96	1.00	5	1.15	0.98	4	0.82	0.99
5	48UR20051116	2	0.81	0.96	1	0.66	0.89	1	0.77	0.95
6	48UR20060608	5	0.95	1.00	5	1.14	0.99	4	0.74	0.93
7	06A420060720	0	-	-	0	-	-	0	-	-
8	48UR20060928	4	0.97	1.00	4	0.86	0.98	3	0.90	0.99
9	48UR20071005	5	1.03	1.00	5	0.86	0.98	4	0.88	0.99
10	48UR20080318	3	1.06	1.00	3	0.91	0.99	2	0.98	1.00
11	48UR20080905	0	-	-	0	-	-	0	-	-
12	48UR20081103	5	0.92	0.99	5	0.62	0.85	4	0.88	0.99
13	48UR20090508	3	0.95	1.00	3	0.67	0.90	2	0.85	0.98
14	48UR20100430	4	1.01	NA	4	0.66	0.88	3	0.88	0.99
15	48UR20100731	5	0.87	0.99	5	0.75	0.93	4	0.74	0.93
16	48UR20101123	1	0.85	0.98	1	0.71	0.91	1	0.72	0.92
17	48UR20110421	2	1.01	NA	2	0.75	0.94	1	0.88	0.99
18	48UR20111109	4	0.99	NA	4	0.86	0.98	3	0.91	0.99
19	48MG20111210	0	-	-	0	-	-	0	-	-
20	48UR20120111	4	1.01	NA	4	0.83	0.98	3	0.92	0.99
21	48UR20121108	0	-	-	0	-	-	0	-	-
22	48UR20131015	4	1.00	NA	4	0.83	0.97	3	0.89	0.99
23	48QL20150804	5	0.98	1.00	5	0.98	1.00	4	0.92	1.00
24	48QL20171023	3	0.66	0.88	3	1.02	1.00	2	0.94	0.99

red: data lower than reference


Table 7

Region/ Water mass	Nitrate O_3 ($\mu\text{mol kg}^{-1}$)		Phosphate O_4 ($\mu\text{mol kg}^{-1}$)		Silicate O_2 ($\mu\text{mol kg}^{-1}$)	
	Avg new Product	Avg Medar	Avg new Product	Avg Medar	Avg new Product	Avg Medar
<i>DF2- Gulf of Lion</i>						
surface water (0-150db)	2.68 \pm 2.53(68)**	1.7 \pm 1.1	0.15 \pm 0.06(68)	0.13 \pm 0.04	2.91 \pm 1.33(68)	1.72 \pm 0.64
LIW core (S_{max} depth range: 300-500db)	8.49 \pm 0.18(17)	6.13 \pm 0.32	0.38 \pm 0.02(17)	0.34 \pm 0.01	8.67 \pm 0.69(17)	6.12 \pm 0.61
Deep water (>1500db)	8.03 \pm 0.43(33)	7.64 \pm 0.31	0.37 \pm 0.01(33)	0.37 \pm 0.015	8.7 \pm 0.67(33)	7.95 \pm 0.06
<i>DF3- Liguro-Provençal</i>						
surface water (0-150db)	2.31 \pm 2.4(205)	3.0 \pm 2.6	0.12 \pm 0.07(205)	0.19 \pm 0.05	2.45 \pm 1.05(205)	2.16 \pm 1.05
LIW core (S_{max} depth range: 300-500db)	8.05 \pm 0.18(76)	7.74 \pm 0.13	0.36 \pm 0.01(76)	0.35 \pm 0.01	7.49 \pm 0.55(76)	6.26 \pm 0.60
Deep water (>1500db)	8.18 \pm 0.25(142)	7.79 \pm 0.04	0.37 \pm 0.02(142)	1.03 \pm 1.29	8.98 \pm 0.39(142)	7.60 \pm 0.21
<i>DF4- Ligurian East</i>						
surface water (0-150db)	0.7 \pm 0.69(228)	0.61 \pm 1.03	0.05 \pm 0.02(228)	0.18 \pm 0.02	1.37 \pm 0.45(228)	1.27 \pm 1.86
LIW core (S_{max} depth range: 300-500db)	6.8 \pm 0.4(23)	5.54 \pm 0	0.3 \pm 0.02(21)	0.36 \pm 0.06	5.86 \pm 0.9(24)	4.86 \pm 0
Deep water (>1500db)	-	-	-	-	-	-
<i>DS2- Balearic Sea</i>						
surface water (0-150db)	1.32 \pm 1.46(196)	1.19 \pm 1.5	0.08 \pm 0.04(196)	0.11 \pm 0.04	1.61 \pm 0.64(196)	1.54 \pm 0.78
LIW core (S_{max} depth range: 300-500db)	8.32 \pm 0.32(58)	6.92 \pm 0.12	0.37 \pm 0.02(60)	0.39 \pm 0.003	7.31 \pm 0.9(60)	7.55 \pm 0.62
Deep water (>1500db)	8.2 \pm 0.35(88)	-	0.37 \pm 0.01(88)	-	8.71 \pm 0.51(88)	8.45 \pm 0.8
<i>DF1- Algero-Provençal</i>						
surface water (0-150db)	0.87 \pm 0.85(372)	1.08 \pm 1.7	0.05 \pm 0.02(372)	0.07 \pm 0.05	1.42 \pm 0.3(372)	1.28 \pm 0.73
LIW core (S_{max} depth range: 300-500db)	8.07 \pm 0.34(126)	7.51 \pm 0.18	0.36 \pm 0.02(126)	0.34 \pm 0.008	6.84 \pm 0.95(126)	5.96 \pm 0.77
Deep water (>1500db)	8.36 \pm 0.27(300)	7.87 \pm 0.13	0.38 \pm 0.02(300)	0.38 \pm 0.001	9.01 \pm 0.33(300)	8.18 \pm 0.10
<i>DS1- Alboran Sea</i>						
surface water (0-150db)	2.75 \pm 2.87(299)	2.51 \pm 2.23	0.17 \pm 0.11(299)	0.16 \pm 0.07	2.07 \pm 1.38(299)	2.31 \pm 1.14
LIW core (S_{max} depth range: 400-600db)	8.89 \pm 0.4(77)	8.14 \pm 0.11	0.42 \pm 0.02(77)	0.37 \pm 0.008	8.77 \pm 1.66(76)	7.95 \pm 0.34
Deep water (>1500db)	7.72 \pm 0.81(65)	-	0.36 \pm 0.04(65)	-	8.98 \pm 0.63(65)	8.16 \pm 0
<i>DS3- Algerian West</i>						
surface water (0-150db)	1.8 \pm 1.88(254)	1.82 \pm 2.01	0.11 \pm 0.05(354)	0.11 \pm 0.06	1.71 \pm 0.68(354)	2.10 \pm 0.91
LIW core (S_{max} depth range: 400-600db)	9.33 \pm 0.08(70)	8.28 \pm 0.15	0.41 \pm 0(73)	0.38 \pm 0.012	8.1 \pm 0.53(72)	6.68 \pm 0.80
Deep water (>1500db)	8.37 \pm 0.27(246)	8.047 \pm 0.013	0.37 \pm 0.02(246)	0.36 \pm 0.006	9.22 \pm 0.35(246)	8.87 \pm 0.23
<i>DS4- Algerian East</i>						
surface water (0-150db)	0.94 \pm 0.77(170)	0.75 \pm 1.26	0.07 \pm 0.02(170)	0.05 \pm 0.03	1.53 \pm 0.12(170)	1.35 \pm 0.52
LIW core (S_{max} depth range: 400-600db)	8.5 \pm 0.25(43)	8.60 \pm 0.06	0.38 \pm 0.03(43)	0.38 \pm 0.008	7.27 \pm 0.67(42)	7.092 \pm 0.55
Deep water (>1500db)	7.94 \pm 0.24(132)	8.06 \pm 0.06	0.36 \pm 0.02(132)	0.38 \pm 0.006	8.73 \pm 0.38(132)	9.04 \pm 0.24
<i>DT1- Tyrrhenian North</i>						
surface water (0-150db)	1.03 \pm 1.14(231)	0.88 \pm 1.2	0.06 \pm 0.02(231)	0.09 \pm 0.03	1.64 \pm 0.52(231)	2.19 \pm 0.59
LIW core (S_{max} depth range: 400-600db)	5.95 \pm 0.49(43)	5.86 \pm 0.36	0.27 \pm 0.03(44)	0.308 \pm 0.02	7.06 \pm 0.08(44)	6.76 \pm 0.59
Deep water (>1500db)	7.75 \pm 0.37(194)	7.12 \pm 0.47	0.36 \pm 0.03(194)	0.40 \pm 0.02	9.19 \pm 0.47(194)	7.51 \pm 0.49
<i>DT3- Tyrrhenian South</i>						
surface water (0-150db)	1.21 \pm 1.38(711)	1.23 \pm 1.80	0.06 \pm 0.03(711)	0.061 \pm 0.04	1.58 \pm 0.61(711)	1.55 \pm 1.05
LIW core (S_{max} depth range: 300-500db)	6.2 \pm 0.28(225)	6.42 \pm 0.01	0.26 \pm 0.02(225)	0.254 \pm 0.005	6.28 \pm 0.65(224)	6.68 \pm 0.44
Deep water (>1500db)	7.88 \pm 0.4(227)	7.12 \pm 0.26	0.37 \pm 0.02(227)	0.31 \pm 0.007	9.04 \pm 0.52(227)	8.02 \pm 0.07
<i>DII- Sardinia Channel</i>						
surface water (0-150db)	1.22 \pm 1.39(271)	1.42 \pm 1.95	0.07 \pm 0.03(271)	0.064 \pm 0.03	1.57 \pm 0.68(271)	1.39 \pm 1.01
LIW core (S_{max} depth range: 300-500db)	6.52 \pm 0.17(89)	6.45 \pm 0.22	0.27 \pm 0.02(89)	0.250 \pm 0.01	6.36 \pm 0.67(89)	6.27 \pm 0.70
Deep water (>1500db)	7.91 \pm 0.62(107)	-	0.37 \pm 0.03(107)	0.32 \pm 0	8.64 \pm 0.91(107)	-
<i>DI3- Sicily Strait</i>						
surface water (0-150db)	0.87 \pm 0.68(583)	0.77 \pm 0.81	0.06 \pm 0.02(583)	0.063 \pm 0.02	1.53 \pm 0.29(583)	1.44 \pm 0.58
LIW core (S_{max} depth range: 200-400db)	4.95 \pm 0.47(80)	5.14 \pm 0.14	0.21 \pm 0.02(78)	0.194 \pm 0.004	5.26 \pm 0.79(81)	6.744 \pm 0.41
Deep water (>1500db)	-	-	-	-	-	-


**Average (Avg) \pm standard deviation of inorganic nutrient (the number observation within depth range) for three layers from the adjusted/new product and MEDATLAS vertical climatological profiles (called here Medar). Regions are defined according to Manca et al. (2004) (table 2S, Fig.2S)

1 **Supplementary material – Part 1**


2 **Figure 1S.** Overview of vertical inorganic nutrient profiles and spatial coverage of reference cruises.



5 **Figure 2S.** Map of the WMED showing the geographical limits of the MEDAR/Medatlas sub-regions
 6 defined in Table.2S according to Manca et al. (2004).




7
 8
 9 **Figure 3S.** Weighted mean offset for nitrate, before (grey) and after adjustment (blue). Error bars indicate
 10 the standard deviation of the absolute weighted offset. The dashed lines indicate the accuracy limit 2% for
 11 an adjustment to be recommended




12

13 **Figure 4S.** Same as Fig. 2S but for phosphate



14

15 **Figure 5S.** Same as Fig. 2S but for silicate.



16

Table 1S. Summary table of laboratories and instruments used for nutrient analysis.

Laboratory	Autoanalyzer	Detection limit
ENEA	continuous-flow system multichannel (Auto Analyzer Bran+Luebbe III Generation)	limit of 0.01 μM for nitrate+nitrite, 0.01 μM for phosphate and 0.05 for silicate
CNR-ISMAR Trieste	OI-Analytical (Flow Solution III) flow-segmented	limit of 0.01 μM for nitrate+nitrite, 0.01 μM for phosphate and 0.05 for silicate
CNR-ISMAR Venezia	Systea discrete analyzer EasyChem Plus	limit of 0.1 μM for nitrate, 0.01 μM for phosphate and 0.02 μM for silicate

Table 2S. Geographical limits of subregion referring to Manca et al.(2004).

Code	Region	Lat °N (min)	Lat °N(max)	Lon °E (min)	Lon °E (max)	Maximum pressure(db)
DF2	Gulf of Lions	42	43.36	1	6.18	2517
DF3	Liguro-Provençal	41	45	6.18	9.18	2728
DF4	Ligurian East	42.48	45	9.18	11	1299
DS2	Balearic Sea	38.30	42	-1	4.3	2741
DF1	Algero-Provençal	39.18	41	4.3	9.18	2891
DS1	Alboran Sea	35.0	37.3	-6	-1	2683
DS3	Algerian West	35.36	38.3	-1	4.3	2837
DS4	Algerian East	36.30	39.18	4.3	8.24	2890
DT1	Tyrrhenian North	39.18	42.48	9.18	16.16	3610
DT3	Tyrrhenian South	38	39.18	10	16.16	3551
DI1	Sardinia Channel	36.0	39.18	8.24	10	2455
DI3	Sicily Strait	35	38	10	15	664

Table 3S. Reference cruises and coefficient of variation of nitrate, phosphate and silicate below 1000db.

Reference cruise ID	EXPOCODE	std Nitrate O_3	std Phosphate O_4	std Silicate O_2	# samples
6	06MT20011018	0.064	0.179	0.035	26
22	48UR20070528	0.121	0.074	0.144	34
27	29AJ20160818	0.052	0.062	0.054	116
64	06MT20110405	0.073	0.071	-	42
17	29AH20140426	0.045	0.112	0.036	91

Supplementary material – Part 2

A1. Database Data product description

The ~~database data product~~ includes 870 stations sampled during 24 cruises between 2004 and 2017 in the Western Mediterranean Sea (~~the reference cruise 29AJ20160818 was included in this dataset as cruise #24 because, it is part of the CNR-ISMAR data collection program~~), mainly on board of research vessels owned by the Italian National Research Council. It includes bottle data combined with CTD data.

Formattato: Inglese (Stati Uniti)

In all stations, measurements were carried out with a CTD-rosette system consisting of a CTD SBE 911 plus and a General Oceanics rosette with 24 12-l Niskin Bottles at the observed depth of the bottle sample. Temperature measurements were performed with an SBE-3/F thermometer with a resolution of 10⁻³ °C and conductivity measurements were performed with an SBE-4 sensor with a resolution of 3·10⁻⁴ S/m. The probes were calibrated before and after the cruise. Except for salinity, no certified reference material (CRM) was used. CTD salinity was calibrated against measurements made with a salinometer.

Samples of nitrate, phosphate and silicate were frozen to -20°C and stored before being analysed in laboratories onshore.

Measurements were subjected to a rigorous quality control (primary and secondary quality control) and the dataset presented is the product adjusted after the application of quality control approaches.

A2. Database Data product organization details

Formattato: Inglese (Stati Uniti)

Cruise identification: To guarantee the comparability between measurements, an alphanumeric identification code (ID) together with an expedition code (Expocode) are defined a unique identifier. The list of the parameters included in the data product are detailed in table below:

Formattato: Inglese (Stati Uniti)

#	Variable in Dataset Short name in data files	Parameter	Unit/format	Method/ description	Comment	Original Dataset	Adjusted
1	EXPOCODE	Expedition code	12 digits 24 EXPOCODEs	12 digits ; Shipcode_yyyy_mm_dd yyyy_mm_dd: cruise starting day		✓	✓
2	CRUISE	Cruise ID	24 Cruise IDs	From 1 to 24		✓	✓
3	DATE	Event date	yyyy-mm-dd			✓	✓
4	TIME	Event time	hhmm			✓	✓
5	DAY	Day	dd			✓	✓
6	MONTH	Month	mm			✓	✓
7	YEAR	Year	yyyy			✓	✓
8	LATITUDE	Longitude				✓	✓
9	LONGITUDE	Latitude				✓	✓
10	STNNBR	Station number				✓	✓
11	BTLNBR	Niskin bottle number				✓	✓
12	CASTNO	Cast number				✓	✓
13	CTDPRS	Pressure	dbar	CTD pressure		✓	✓
14	DEPTH	Depth	Meters	Depth from pressure		✓	✓
15	CTDSAL	Salinity		CTD salinity	PSS-78	✓	✓
16	CTDSAL_FLAG_W	Salinity flag		WOCE flags		✓	✓
17	CTDTMP	Temperature	°C	CTD temperature	ITS-90	✓	✓
18	THETA	Potential temperature		Theta from CTDTMP & CTDSAL		✓	✓
19	NITRAT	Nitrate	μmol kg ⁻¹	standard colorimetric methods*		✓	✓
20	NITRAT_FLAG_W NITRAT_FLAG_re	Nitrate flag		WOCE flags		✓	✓
		Recommended nitrate flag		After 1 st quality control Flags after 2 nd QC Flag 2: adjusted and acceptable Flag 3: adjusted and recommended questionable	Details in Section 4.4		
21	PHPHPT	Phosphate	μmol kg ⁻¹	standard colorimetric methods*		✓	✓
22	PHSPHT_FLAG_W PHSPHT_FLAG_re	Phosphate flag		WOCE flags		✓	✓
		Recommended phosphate flag		After 1 st quality control Flags after 2 nd QC Flag 2: adjusted and acceptable Flag 3: adjusted and recommended questionable	Details in Section 4.4		
23	SILCAT	Silicate	μmol kg ⁻¹	standard colorimetric methods*		✓	✓
24	SILCAT_FLAG_W SILCAT_FLAG_W	Silicate flag		WOCE flags		✓	✓
		Recommended silicate flag		After 1 st quality control Flags after 2 nd QC Flag 2: adjusted and acceptable Flag 3: adjusted and recommended questionable	Details in Section 4.4		

* Standard colorimetric methods of seawater analysis (Grasshoff et al. (1999)).

- Data format

Original dataset: CNR_DIN_WMED_20042017_original.csv: This is the original dataset with 24 fields-parameter including flag variables of 24 cruises for nitrate, phosphate, silicate and CTD salinity from the primary quality control.

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Formattato ...

Adjusted dataset: *CNR_DIN_WMED_20042017_adjusted.csv*: This is the adjusted product with 24 [fields](#) [parameter](#), after removing outlier data (issued from primary quality control) and after applying adjustment [factors from the secondary quality control \(Crossover Analysis\)](#).

Formattato: Inglese (Stati Uniti)

Formattato: Inglese (Stati Uniti)