



# The Vulcan Version 3.0 High-Resolution Fossil Fuel CO<sub>2</sub> Emissions for the United States

Kevin R. Gurney<sup>1</sup>, Jianming Liang<sup>2</sup>, Risa Patarasuk<sup>2</sup>, Yang Song<sup>2</sup>, Jianhua Huang<sup>2</sup>, Geoffrey Roest<sup>1</sup>

<sup>1</sup>School of Informatics, Computing, and Cyber Systems, Northern Arizona University, Flagstaff, AZ, USA

<sup>2</sup>School of Life Sciences, Arizona State University, Tempe AZ USA

Correspondence to: Kevin R. Gurney (kevin.gurney@nau.edu)

**Abstract.** Estimates of greenhouse gas emissions, quantified at fine space and time scales, has become a critical component of new multi-constraint flux information systems in addition to providing relevant information to decisionmakers when considering GHG mitigation opportunities. The ‘Vulcan Project’ is an effort to estimate bottom-up fossil fuel emissions and CO<sub>2</sub> emissions from cement production (FFCO<sub>2</sub>) for the entire United States landscape at space and time scales that satisfy both scientific and policy needs. Here, we report on version 3.0 of the Vulcan emissions which quantifies FFCO<sub>2</sub> emissions for the U.S. at a spatial resolution of 1km x 1km and hourly temporal resolution for the 2010-2015 time period. We provide a complete description of the updated methods, data sources, results, and comparison to a global gridded FFCO<sub>2</sub> data product. We estimate FFCO<sub>2</sub> emissions for the year 2011 of 1589.3 TgC with a 95% confidence interval of 1299/1917 TgC (+18.3%/-20.6%), implying a one-sigma uncertainty of  $\sim \pm 10\%$ . We find that per capita FFCO<sub>2</sub> emissions are larger in states dominated by the electricity production and industrial sectors and smaller in states dominated by onroad and residential/commercial building emissions. The center of mass (CoM) of FFCO<sub>2</sub> emissions in the US are located in the state of Missouri with mean seasonality that moves on a NE/SW near-elliptical path. Comparison to ODIAC, a global gridded FFCO<sub>2</sub> emissions estimate shows large differences in both total emissions (100.1 TgC for year 2011) and spatial patterns. The spatial correlation ( $R^2$ ) between the two data products was 0.38 and the mean absolute difference at the individual gridcell scale was 80.04%. The Vulcan v3.0 FFCO<sub>2</sub> emissions data product offers an immediate high-resolution estimate of emissions in every city within the U.S., providing a large potential savings of time and effort for cities planning to develop self-reported city inventories. The Vulcan v3.0 annual gridded emissions data product can be downloaded from the Oak Ridge National Laboratory Distributed Active Archive Center (ORNL DAAC) (<https://doi.org/10.3334/ORNLDAAC/1741>, Gurney et al., 2019).

## 1 Introduction

- 2 Global emissions of carbon dioxide from the combustion of fossil fuels (FFCO<sub>2</sub>) comprise the largest net flux of
- 3 carbon into the Earth’s atmosphere and remain the primary driver of anthropogenic climate change (IPCC 2013;
- 4 USGCRP 2018). Improving our quantitative understanding of FFCO<sub>2</sub> fluxes remains a critical component of climate
- 5 change research and climate policy. For example, scientific understanding of the global carbon cycle and how it
- 6 interacts with climate change rests on accurate quantification of FFCO<sub>2</sub> emissions at multiple scales (LeQuere,



2018). This, in turn, improves the reliability of future projections of climate change and specifies the emissions reductions necessary to meet specific targets, such as limiting the rise of global mean temperature to 1.5 C (IPCC 2018). Understanding FFCO<sub>2</sub> sources also assists in understanding the composition, driving factors, and responsibility for emissions, making mitigation options better-targeted, equitable and ultimately more effective (Durant et al., 2011; Janssens-Maenhout et al., 2013; Bellassen et al., 2015).

Quantification of FFCO<sub>2</sub> emissions began as efforts to capture total emissions at the global and national spatial scales aiming to quantify anthropogenic fluxes to better understand the drivers of climate change and the global carbon cycle. Employing accounting approaches that rely on national statistics of energy production and consumption, a number of national and international institutions produce and archive estimates of FFCO<sub>2</sub> emissions, often disaggregated to economic sector and fuel type (see reviews by Andres et al., 2012; Macknick, 2011). In response to the advances in carbon cycle observations and modeling studies, many of these FFCO<sub>2</sub> inventory products began to increase their spatial and temporal resolution below the nation-state, often representing emissions in regularized gridded format (Marland et al., 1985; Andres et al., 1996; Olivier et al., 1999). Gridded output was especially important when used within systems that solve carbon fluxes through inversion of atmospheric transport constrained by atmospheric concentration measurements (Gurney et al., 2002; 2005; Peylin et al., 2011; Liu et al., 2014; Yadav et al., 2016; Gaubert et al., 2019). Most often these sub-national representations of FFCO<sub>2</sub> emissions used proxy information, such as population statistics or remotely-sensed nighttime lights, to distribute the national/global emissions to smaller space/time scales (Andres et al., 1999, Olivier et al., 2005; Rayner et al., 2010; Ghosh et al., 2010, Oda and Maksyutov, 2011; Ou et al., 2015). Recent research has employed a mixture of global “bottom-up” information such as powerplant databases with remote-sensing information (Wang et al., 2013; Oda et al., 2018 etc), sometimes within optimization frameworks to more mechanistically distribute emissions in space and time in addition to offering more formal uncertainty estimation (Asefi et al., 2014).

In addition to the globally gridded representations, research effort has also aimed at specific national and regional domains often with additional detail on the emitting process (Gregg and Andres, 2007; Gregg et al., 2009; Bun et al., 2007; 2018; Gately et al., 2017; Kurokawa et al., 2013; Ivanova et al., 2017; Denier et al., 2017; Cai et al., 2018) with some studies focussed on an individual sector or source type (Petron et al., 2008; Gately et al., 2013; Zheng et al., 2014; Wang et al., 2014; Liu et al., 2015). These national and regional efforts were often modeled after work in local air pollution inventories (Cooke et al., 1999; Baldesano et al., 2008; O’Hara et al., 2007; Hoesly et al., 2017). In addition to focusing on different national domains with unique datasets, many of these past research efforts reflect different methodological approaches to data interpretation, downscaling and modeling. Many of these efforts, however, follow the general approach established in the pioneering work of the Vulcan Project, the first attempt to generate a completely bottom-up space/time-explicit national estimate of all FFCO<sub>2</sub> emission sources (Gurney et al., 2009). The Vulcan Project, which estimated FFCO<sub>2</sub> emissions at the “native” resolution of emission points, lines, and polygons, produced US FFCO<sub>2</sub> emissions on a 10km x 10km spatial grid at hourly time resolution for the year 2002. Used in a variety of research and applied policy settings, the Vulcan Project has spawned additional efforts at downscaling into the urban domain, where resolution has gone to the scale of individual buildings and street



1 segments for whole urban areas (VandeWeghe and Kennedy 2007; Shu and Lam 2011; Zhou and Gurney 2011;  
 2 Gurney et al., 2012; Wilson et al., 2013; Pincetl et al., 2014; Patarasuk et al. 2016; Gurney et al., 2018; 2019b).  
 3 All of the FFCO<sub>2</sub> emissions reviewed thus far are categorized as “scope 1” or “in-boundary” emissions. They are an  
 4 accounting of emissions that reflects physical emission of CO<sub>2</sub> molecules from the geography resolved (e.g. gridcell,  
 5 state, province). This is in contrast to quantification of fluxes that assign emissions to consumptive activity such as  
 6 using electricity or consuming food (Davis and Caldeira, 2010). The two accounting perspectives are identical at the  
 7 whole-planet scale but diverge as one considers scales at the nation-state or below. Consumption-based FFCO<sub>2</sub>  
 8 emissions quantification has a long history at scales ranging from the nation-state to the city, but has only recently  
 9 begun to systematically resolve (e.g. in gridded form) FFCO<sub>2</sub> emissions below the nation-state scale (Jones and  
 10 Kammen, 2011; 2013; Zhang et al., 2014; Minx et al., 2013; Moran et al., 2018). The current study emphasizes in-  
 11 boundary emissions because these can be directly used with atmospheric monitoring, a critical element in  
 12 evaluation/validating the estimated fluxes and a motivation for the research reported here (NRC 2010).  
 13 In this paper, we introduce a significant update to the Vulcan Project estimation of high-resolution US fossil fuel  
 14 carbon dioxide emissions and CO<sub>2</sub> emissions from cement production (collectively referred to here as “FFCO<sub>2</sub>”).  
 15 We report here on improvements in methodology, resolution, uncertainty estimation, in addition to more  
 16 contemporaneous, multiyear output. We present some of the fundamental results of the Vulcan output and compare  
 17 to the only commensurate resolved FFCO<sub>2</sub> emissions data product covering the entire U.S. landscape, the ODIAC  
 18 global data product. We show results associated with a few zoomed urban locations, suggesting that the Vulcan  
 19 FFCO<sub>2</sub> emissions data product has a role to play in providing U.S. cities with a sub-city resolved scope 1 CO<sub>2</sub>  
 20 emissions inventory.  
 21 Version 3.0 of the Vulcan data product and associated documentation is publicly available and annual, gridded,  
 22 multiyear results can be downloaded from the Oak Ridge National Laboratory Distributed Active Archive Center  
 23 (ORNL DAAC) (<https://doi.org/10.3334/ORNLDAAC/1741>).  
 24 This paper is structured as follows: In section 2, we describe the data and model processes used to generate the  
 25 Vulcan version 3.0 (v3.0) FFCO<sub>2</sub> emissions data product including those used for spatial and temporal distribution.  
 26 In section 3.0, we present the results, the uncertainties and a series of descriptive statistics at various scales of  
 27 aggregation. In section 4, we compare Vulcan to the ODIAC data product, and discuss the potential use and  
 28 relevance of this work, known gaps and weaknesses, in addition to next steps and future work.

## 29 **2 Methods**

30 The Vulcan version 3.0 FFCO<sub>2</sub> emissions data product represents total FFCO<sub>2</sub> emissions resulting from the  
 31 combustion of fossil fuel (coal, petroleum and natural gas) and the CO<sub>2</sub> from cement production in the 50 United  
 32 States and District of Columbia for 2010-2015 time period (Gurney et al., 2019). It is constructed from numerous  
 33 public datasets that generate emissions magnitude, the spatial representation, and temporal representation of those  
 34 emissions. The FFCO<sub>2</sub> emissions are initially estimated at their “native” spatial and temporal resolution (e.g.



counties, points, lines, annual, hourly) depending upon the characteristics of the incoming data sources. Additional spatial and temporal “conditioning” (e.g. downscaling, interpolation, proxy surrogates), where needed, is used to arrive at an hourly representation for six complete calendar years (2010-2015) at the spatial resolutions of a US Census block-group or finer (e.g. points, lines). The FFCO<sub>2</sub> emissions are further processed to regularized hourly grids at a resolution of 1 km x 1 km, for the contiguous United States and Alaska. The FFCO<sub>2</sub> emissions represent all fossil fuel combustion extending 12 nautical miles from the coastal boundary of the United States.

## 2.1 Data and processing

The data sources for the FFCO<sub>2</sub> emissions estimation are organized here by data source *type* and/or the economic *sector* in accordance with original data collection/categorization (see Table 1). This paper describes the scientific methodology used to generate the Vulcan v3.0 FFCO<sub>2</sub> emissions but should be considered in combination with the published results for the earlier version 2.0 Vulcan results (Gurney et al., 2009; Zhou et al., 2011) and the Vulcan version 2.0 documentation (<http://vulcan.rc.nau.edu/assets/files/Vulcan.documentation.v2.0.online.pdf>).

Uncertainty quantification relies on the characterization of a 95% confidence interval (CI). Due to the considerable runtime of the Vulcan codebase, only the boundaries of the upper and lower CI are estimated (referred to as “hi” and “lo” CI bounds). Future versions of the Vulcan data product will quantify the complete uncertainty distribution of the Vulcan FFCO<sub>2</sub> emissions output.

**Table 1: Overview of data sources used in generating the space/time-resolved Vulcan v3.0 FFCO<sub>2</sub> emissions (footnotes provide acronym explanations).**

Sector/type	Emissions Data Source	Original spatial resolution/information	Spatial distribution	Temporal distribution
Onroad	EMFAC <sup>a</sup> CO <sub>2</sub> , EPA NEI <sup>b</sup> onroad CO <sub>2</sub>	County, road class, vehicle class	FHWA AADT <sup>c</sup>	CCS <sup>d</sup>
Electricity production	CAMD <sup>d</sup> CO <sub>2</sub> , DOE/EIA <sup>a</sup> fuel, EPA NEI point CO	Lat/lon, fuel type, technology	EPA/EIA NEI Lat/Lon, Google Earth	CAMD, EIA and EPA
Residential nonpoint buildings	EPA NEI nonpoint CO	County, fuel type	FEMA HAZUS <sup>d</sup> , DOE RECS NE-EUI <sup>b</sup>	eQUEST <sup>d</sup> model
Nonroad	NEI nonpoint CO	County, vehicle class	EPA spatial surrogates (vehicle class specific)	EPA temporal surrogates (by SCC)
Airport	EPA NEI point CO	Lat/lon, aircraft class	Lat/Lon	LAWA & OPSNET <sup>k</sup>
Commercial nonpoint buildings	EPA NEI nonpoint CO	County, fuel	FEMA HAZUS, DOE CBECS NE-EUI <sup>b</sup>	eQUEST model
Commercial point sources	EPA NEI point CO	Lat/lon, fuel type, combustion technology	EPA NEI Lat/Lon, Google Earth	eQUEST model
Industrial point sources	EPA NEI point CO	Lat/Lon, fuel type, combustion technology	EPA NEI Lat/Lon, Google Earth	EPA temporal surrogates (by SCC)
Industrial nonpoint buildings	EPA NEI nonpoint CO	County, fuel type	FEMA HAZUS, DOE MECS NE-EUI <sup>b</sup>	eQUEST model
Commercial Marine Vessels	EPA NEI nonpoint CO	County, fuel type, port/underway	EPA port and shipping lane shapefiles	Flat time structure
Railroad	EPA NEI nonpoint CO, EPA NEI point CO	County, fuel type, segment	EPA NEI rail shapefile and density distribution	Point records: EPA temporal surrogates (by SCC). Nonpoint: flat time structure
Cement	Portland Cement Association, USGS	Lat/lon	PCA lat/lon checked in Google Earth	Flat time structure

- 19 a. Emissions Factors Model  
20 b. Environmental Protection Agency, National Emissions Inventory  
21 c. Federal Highway Administration, Annual Average Daily Traffic  
22 d. Federal Emergency Management Agency



- 1 e. Continuous Count Stations
- 2 f. Clean Air Markets Division
- 3 g. Department of Energy/Energy Information Administration
- 4 h. Department of Energy Residential Energy Consumption Survey, non-electric energy use intensity
- 5 i. Quick Energy Simulation Tool
- 6 j. Source Classification Code
- 7 k. Los Angeles World Airport, The Operations Network
- 8 l. Department of Energy Commercial Energy Consumption Survey, non-electric energy use intensity
- 9 m. Department of Energy Manufacturing Energy Consumption Survey, non-electric energy use intensity

## 10 2.1.2 Nonpoint sources

11 The area or nonpoint source emissions (dominated by residential and commercial economic sectoral categories) are  
 12 stationary sources that are not inventoried at the individual facility-level and can be thought of as representing  
 13 “diffuse” or dispersed sources within a geographic area. Vulcan nonpoint FFCO<sub>2</sub> emissions are estimated using a  
 14 number of data sources. Foremost among these are the Environmental Protection Agency (EPA) National Emission  
 15 Inventory (NEI) nonpoint reporting for carbon monoxide (CO) emissions, version 2 for the year 2011 (USEPA  
 16 2015a). The NEI is a comprehensive inventory of all criteria air pollutants (CAPs) and hazardous air pollutants  
 17 (HAPs) across the United States (USEPA 2005a). The NEI now includes greenhouse gases for select sectors  
 18 (onroad, nonroad). The NEI is a data structure with which the EPA can meet mandates established by the Clean Air  
 19 Act (CAA). The CAP emissions, the component of emissions used by the Vulcan system (other than onroad,  
 20 nonroad, and electricity production), are collected under the Air Emissions Reporting Rule (40 CFR Part 51) (CFR,  
 21 2008). The NEI can be used to track progress, drive air quality modeling, enable emissions trading, and ensure  
 22 comprehensive reporting and compliance.

23 The emissions data within the NEI are collected from state, local, and tribal (SLT) agencies and augmented by  
 24 numerous federal data sets such as the Toxics Release Inventory (TRI), the Acid Rain Program (ARP), and the  
 25 Federal Highway Administration (FHWA) traffic counts.

26 The EPA provides recommendations to SLT agencies on how to collect nonpoint source emissions information and  
 27 the SLT agencies are given a number of options in forming the basis of the reported information (ERG 2001). The  
 28 EPA prefers emissions to be estimated by extrapolating from a sample set of data for the activity to the entire  
 29 population, but a number of other approaches are allowed including material balance, mathematical models, and  
 30 emission factors (EFs). This means that the method employed will vary by location. The EPA will augment the  
 31 submitted data as a result of recognized data gaps, QA/QC procedures, or in consultation with SLT agencies.

32 The 2011 NEIv2 nonpoint data used in the Vulcan emissions estimation is composed of two core data files. These  
 33 data files share common, required key fields. The fundamental nonpoint “unit”, as pertains to the Vulcan  
 34 calculations, is a reported combustion process emitting carbon monoxide (CO) identified by a single source  
 35 classification code (SCC) in a single US county burning an identified fossil fuel. The numerical SCC (USEPA,  
 36 1995) and FIPS values (which identifies the state and county via numerical ID) are critical common IDs. Reporting  
 37 associated with fugitive emissions (non-combustion), chemical or “in-process”, or resulting from the combustion of  
 38 biogenic fuel sources are removed. An exception to this is the in-process emissions associated with cement  
 39 production, however these emissions are generated with different data outlined in a later section. Fuels considered in



- 1 the Vulcan nonpoint FFCO<sub>2</sub> estimation along with their thermodynamic heat value, default CO emission factor (EF),
- 2 and CO<sub>2</sub> EF are provided in Table 2.

3 **Table 2: Heat value, carbon monoxide emission factor, and carbon dioxide emission factor for emission**  
 4 **sources. Square brackets denote instances in which different emission factors are used in application to point**  
 5 **versus nonpoint data.**

Sector	Fuel	HV (e6btu/unit)	Unit	CO EF (lbs/e9btu)/	CO <sub>2</sub> EF (tC/e9btu)
Electricity Production	Bituminous Coal	26.50 <sup>c</sup>	Tonne	247	25.4
Electricity Production	Subbituminous Coal	19.30 <sup>c</sup>	Tonne	344	25.9
Electricity Production	Bituminous/Subbituminous Coal	22.90 <sup>c</sup>	Tonne	295	25.9
Electricity Production	Coal	22.90 <sup>c</sup>	Tonne	29	25.9
Electricity Production	Anthracite	27.49 <sup>c</sup>	Tonne	24	28.2
Electricity Production	Lignite	14.29 <sup>c</sup>	Tonne	39	26.2
Electricity Production	Natural Gas	1032.00 <sup>h</sup>	e6ft3	63	14.5
Electricity Production	Distillate Oil	139.93 <sup>s</sup>	e3gal	36	19.8
Electricity Production	Residual Oil	149.97 <sup>s</sup>	e3gal	33	21.3
Electricity Production	Liquified Petroleum Gas (LPG)	94.00 <sup>s</sup>	e3gal	28	16.9
Electricity Production	Process Gas	1068.57 <sup>h</sup>	e6ft3	33	15.3
Electricity Production	Coke	30.82 <sup>s</sup>	tonne	21	27.6
Electricity Production	Distillate Oil (Diesel)/Diesel	137.06 <sup>s</sup>	e3gal	929	20.0
Electricity Production	Oil	138.69 <sup>h</sup>	e3gal	36	19.8
Electricity Production	Jet Fuel	120.19 <sup>s</sup>	e3gal	751	19.2
Electricity Production	Refinery Gas	1068.57 <sup>h</sup>	e6ft3	33	15.3
Industrial	Bituminous Coal	"	"	250	25.4
Industrial	Subbituminous Coal	"	"	343	25.9
Industrial	Bituminous/Subbituminous Coal	"	"	296	25.9
Industrial	Coal	"	"	289	25.9
Industrial	Natural Gas	"	"	81	14.5
Industrial	Anthracite	"	"	24	28.2
Industrial	Waste Oil	138.69 <sup>h</sup>	e3gal	14	20.0
Industrial	Distillate Oil	"	"	36	19.8
Industrial	Residual Oil	"	"	33	21.3
Industrial	Liquified Petroleum Gas (LPG) [nonpoint/point]	"	"	85 <sup>o</sup> /36	16.9
Industrial	Coke [nonpoint/point]	"	"	21/24	27.6
Industrial	Process Gas	"	"	33/10	15.3
Industrial	Kerosene	134.91 <sup>s</sup>	e3gal	37	19.5
Industrial	Jet Fuel	120.19 <sup>s</sup>	e3gal	54	19.2
Industrial	Gasoline	129.88 <sup>s</sup>	e3gal	60820	19.2
Industrial	distillate oil (no 2)	139.93 <sup>s</sup>	e3gal	36	19.8
Industrial	Distillate Oil (Diesel)	"	"	48	20.0
Industrial	Refinery Gas	"	"	33	15.3
Industrial	jet A fuel	120.19 <sup>s</sup>	e3gal	54	19.2
Industrial	jet naptha	120.19 <sup>s</sup>	e3gal	54	19.7
Industrial	Oil	"	"	36	19.8
Industrial	blast furnace gas	92 <sup>h</sup>	e6ft3	5554	56.3
Industrial	coke oven gas	574 <sup>h</sup>	e6ft3	1836	11.1
Industrial	Propane	90.42 <sup>s</sup>	e3gal	35 <sup>o</sup>	17.0
Commercial	Anthracite Coal	"	"	240	"
Commercial	Bituminous Coal	"	"	250	"
Commercial	Subbituminous Coal	"	"	343	"
Commercial	Bituminous/Subbituminous Coal	"	"	296	"
Commercial	Coal	"	"	530	"
Commercial	Natural Gas	"	"	81	"
Commercial	Distillate Oil	"	"	36	"
Commercial	Residual Oil	"	"	33	"
Commercial	Liquified Petroleum Gas (LPG) [nonpoint/point]	"	"	85 <sup>o</sup> /21	"
Commercial	Kerosene	"	"	37	"
Commercial	Diesel [nonpoint/point]	137.06 <sup>s</sup>	e3gal	36 <sup>o</sup> /929	20.0
Commercial	Gasoline	"	"	60820	"
Commercial	Jet fuel	"	"	19	"
Commercial	Process gas	"	"	33	"



Commercial	Propane	"	"	21 <sup>o</sup>	"
Commercial	Anthracite culm	27.49 <sup>c</sup>	tonne	12	"
Residential	Bituminous Coal	"	"	11441	"
Residential	Subbituminous Coal	"	"	15705	"
Residential	Bituminous/Subbituminous Coal	"	"	13573	"
Residential	Coal	"	"	13238	"
Residential	Anthracite	"	"	11028	"
Residential	Natural Gas [nonpoint/point]	"	"	39 <sup>o</sup> /63	"
Residential	Distillate Oil	"	"	36	"
Residential	Residual Oil	"	"	33	"
Residential	Liquified Petroleum Gas (LPG)	"	"	21	"
Residential	Kerosene	"	"	37	"
Residential	Propane	"	"	21	"
Railroad	Diesel	"	"	428 <sup>o</sup>	"
Railroad	Distillate Oil (diesel)	"	"	811	"
Marine vessels	Diesel	"	"	428 <sup>o</sup>	"
Marine vessels	Residual Oil	"	"	33	"
Marine vessels	Gasoline	"	"	60820	"
Nonroad	Gasoline	"	"	60820	"
Nonroad	Distillate Oil (diesel)	"	"	929	"
Nonroad	Liquified Petroleum Gas (LPG)	"	"	28	"

<sup>c</sup> Average heating value estimated at the state scale using a volume-weighted average. Source: Department of Energy/Energy Information Administration, Electric Power Monthly, 2011. Form EIA-423, "Monthly Cost and Quality of Fuels for Electric Plants Report," Federal Energy Regulatory Commission, FERC Form 423, "Monthly Report of Cost and Quality of Fuels for Electric Plants."

<sup>f</sup> Natural gas heat value is sourced to EPA 2018, Annex 2, Table A-46, page A-76. Petroleum fuels heat values sourced to EPA 2018, Annex 2, Table A-50, page A-83.

<sup>g</sup> Values from Table 3-5, "Compendium of Greenhouse Gas Emissions Methodologies for the Oil and Gas Industry, American Petroleum Institute, February 2005. There is an updated document and it is API 2009: Table 3-8 on page 3-21.

<sup>j</sup> All values reported in Gurney et al., (2009) unless specified otherwise.

<sup>o</sup> Value retrieved from self-reported data (see main text).

<sup>h</sup> [http://www.engineeringtoolbox.com/heating-values-fuel-gases-d\\_823.html](http://www.engineeringtoolbox.com/heating-values-fuel-gases-d_823.html)

Fossil fuel CO<sub>2</sub> emissions are created from NEI-reported county-scale CO reporting through the application of CO and CO<sub>2</sub> emission factors as follows:

$$E_{n,f}^{CO_2} = \frac{E_{n,f}^{CO}}{EF_{n,f}^{CO}} EF_{n,f}^{CO_2} \quad (1)$$

where  $E_{n,f}^{CO_2}$ , are the CO<sub>2</sub> emissions for a process  $n$  (e.g. industrial 10 MMBTU boiler, industrial gasoline reciprocating turbine) and fuel  $f$  (e.g. natural gas, bituminous coal);  $E_{n,f}^{CO}$  are the equivalent amount of CO emissions for a process  $n$  and fuel  $f$ ;  $EF_{n,f}^{CO}$  is the CO emission factor for a process  $n$  and fuel  $f$ ; and  $EF_{n,f}^{CO_2}$  is the CO<sub>2</sub> EF for a process  $n$  and fuel  $f$ . The CO EF is retrieved from two categories of source information: 1) "self-reported" values (supplied by state or federal air quality specialists submitting the CO emissions reporting: [ftp://newftp.epa.gov/air/nei/2011/doc/2011v2\\_supportingdata/nonpoint/](ftp://newftp.epa.gov/air/nei/2011/doc/2011v2_supportingdata/nonpoint/))<sup>1</sup> or 2) "default" values generated from a combination of values retrieved from the EPA WebFIRE EF database (<https://cfpub.epa.gov/webfire/>) and values accumulated through literature review (see Table 2 and table footnotes for details). The self-reported CO EF values are assessed for reliability and replaced by a default value if the self-reported value is less than 0.1 or greater than 5 times the identified default value.

<sup>1</sup> The file "NonPoint\_Activity2011V2.csv" is no longer archived and/or available from the United States Environmental Protection Agency.





1 The state total FFCO<sub>2</sub> emissions calculated as described above are compared to sector and fuel-specific fuel  
 2 consumption totals reported by the Department of Energy/Energy Information Administration (DOE/EIA) State  
 3 Energy Data System (DOE/EIA, 2018). The EIA SEDS consumption data are gathered to create a historical time  
 4 series of energy production, consumption, prices and expenditures for members of congress, federal and state  
 5 agencies and the general public in addition to supporting EIA energy modeling analysis. The consumption in energy  
 6 units are converted to FFCO<sub>2</sub> using CO<sub>2</sub> EFs for each fuel type category (natural gas, petroleum, coal) from values  
 7 supplied in Table 2. Because the EIA SEDS does not separately report nonpoint versus point sources for a given  
 8 sector/fuel combination, the sum of the Vulcan nonpoint and point (see next section) FFCO<sub>2</sub> emissions are compared  
 9 to the EIA/SEDS totals. Adjustment of the Vulcan state/sector/fuel totals are made to the nonpoint residential and  
 10 commercial sectors only and for natural gas and petroleum fuel (aggregate) only. This is driven by the understanding  
 11 that the survey sampling performed by the EIA SEDS in the industrial sector is more uncertain due to the variety of  
 12 fuel consumption circumstances and idiosyncratic contractual arrangements made between utilities/fuel suppliers  
 13 and industrial entities. Furthermore, industrial facilities have the capability to “stockpile” fuel, making use of annual  
 14 consumption data difficult to interpret without stockpile information. This, and the fact that the coal-based emissions  
 15 are small to non-existent in the residential and commercial sectors, is why adjustment is not made for coal fuel  
 16 values. The adjustments made to the nonpoint residential and commercial FFCO<sub>2</sub> emission amounts are shown in the  
 17 supplementary material, Table S1.

18 Sub-county distribution of the county/sector/fuel-specific FFCO<sub>2</sub> emissions to US Census block-groups uses the  
 19 total floor area (m<sup>2</sup>) of buildings (specific to a building class) within each US Census block-group combined with  
 20 estimates of energy use intensity (EUI). The general approach follows:

$$21 \quad TE_{n^3,f}^{bg} = TFA_{n^1}^{bg} \times EUI_{n^2,f}^{cd} \{n^1 \rightarrow n^2 \rightarrow n^3\} \quad (2)$$

22 where the total emissions,  $TE$ , associated with a building of type,  $n$ , using fuel,  $f$ , in a block-group,  $bg$ , is equal to the  
 23 product of the total floor area,  $TFA$ , and the energy use intensity,  $EUI$ , of buildings in a census division,  $cd$ . Because  
 24 the data sources have somewhat different building type classification schemes, a crosswalk between the various  
 25 categories must be achieved.

26 Building floor area is retrieved from HAZUS General Building Stock data collected and compiled by the Federal  
 27 Emergency Management Agency (FEMA, 2017). Using multiple sources including the US Census and the DOE, the  
 28 FEMA floor area provides an estimate of the building floor area for each US Census block-group specific to a  
 29 classification of building types in the residential, commercial and industrial sectors. The data sources are primarily  
 30 reflective of conditions in 2010.

31 The non-electric energy use intensity (NE-EUI; joules/m<sup>2</sup>) values are compiled by the DOE from building  
 32 consumption energy surveys in different regions of the US. The NE-EUI values were calculated from data in the  
 33 DOE/EIA Commercial Buildings Energy Consumption Survey (CBECS, 2016), Manufacturing Energy  
 34 Consumption Survey (MECS, 2010), and Residential Energy Consumption Survey (RECS, 2013) microdata which  
 35 represent regional (9 US Census Divisions) surveys of building energy consumption categorized by building type,





1 fuel, and age cohort. The three data sources represent survey conditions in 2012, 2009, and 2010, respectively. A  
 2 crosswalk is created linking the FEMA building types to the DOE/EIA building types (supplementary material,  
 3 Table S2). For the industrial sector, data is insufficient to support specificity to US Census Division. Hence, the  
 4 national average results are used but specific to industrial NAICS category and fuel category.

5 Where insufficient data existed to support Census Division-specific NE-EUI values in any of the three sectors, an  
 6 average was calculated using all other Division/building type/fuel-specific NE-EUI values.

7 The product of the total building area for a given Census block-group/sector/building type combination and the  
 8 sector/building type/fuel NE-EUI values act as a distributional fraction of the county total county/sector/fuel FFCO<sub>2</sub>  
 9 to each Census block-group. Hence this acts to provide a relative distribution of building FFCO<sub>2</sub> emission within a  
 10 US county only.

11 The time distribution of the annual FFCO<sub>2</sub> emissions for the nonpoint data source uses a building energy model,  
 12 eQuest, to generate simulated building energy consumption which, in turn, is used to represent hourly time patterns  
 13 (Hirsch & Associates, 2004). The eQuest simulations are based on a series of building prototypes which must be  
 14 related to the FEMA building typology (in turn, related to the final Vulcan building types – see Table S2) of the  
 15 Vulcan system. This relationship is shown in supplementary material, Table S3.

16 To capture the local weather/climate conditions, the eQuest model is additionally driven by the 1020 “TMY3  
 17 weather station datasets (<http://doe2.com/Download/Weather/TMY3/>) from the DOE (Marion and Urban, 1995).  
 18 The weather statistics reflect the 1991-2005 climatological mean conditions. The resulting simulations are used to  
 19 generate hourly fractional energy consumption for each of the weather station locations and for each of the building  
 20 types listed in Table S3. The closest weather station location to each of the Census block-group centroids is used to  
 21 assign these hourly fractional time series to a given block-group/building type combination.

22 *Uncertainty*

23 Nonpoint source uncertainty is applied to the reported CO emissions, the CO EF, and the CO<sub>2</sub> EF. For the reported  
 24 CO emissions, an uncertainty value of  $\pm 12.8\%$  was used, a value reported by Gately et al. (2017) for the residential  
 25 sector (which dominates the nonpoint sources) and based on a state-scale difference between ACES and EIA state  
 26 residential fuel consumption. We interpret this as a 95% confidence interval given that this is estimated from a  
 27 measure of difference by Gately et al. (2017).

28 For the EF uncertainty, the CO and CO<sub>2</sub> EFs were adjusted in combination such that the outcome achieves the hi and  
 29 lo CI, respectively. For example, the upper/lower CI bound for the CO EF was combined with the lower/upper CI  
 30 bound for the CO<sub>2</sub> EF to achieve the hi/lo FFCO<sub>2</sub> emissions output CI bound.

31 An uncertainty of  $\pm 20\%$  is applied to both the default and self-reported CO EFs regardless of fuel type. An  
 32 exception to this is for the “blast furnace gas” and “coke oven gas” fuel types in which the adjustment is  $\pm 35\%$   
 33 (Table 3). The CO EF adjustment is based on estimates of the range found in the WebFIRE database and the self-  
 34 reported CO emission factors. The CO<sub>2</sub> EF uncertainty for coal is derived from the work of Quick (2010) while



- 1 uncertainty for petroleum fuels and natural gas are derived from USEPA Greenhouse Gas Inventory, Annex 2
- 2 (USEPA, 2017).

3 **Table 3: Upper and lower confidence interval values for the CO and CO<sub>2</sub> emission factors.**

Sector	Fuel	CO EF lo/hi (lbs/e9btu)	CO <sub>2</sub> EF lo/hi (tC/e9btu)
Electricity Production	Bituminous Coal	296 / 196	24.7 / 26.1
Electricity Production	Subbituminous Coal	413 / 275	25.5 / 26.4
Electricity Production	Bituminous/Subbituminous Coal	354 / 236	25.1 / 26.8
Electricity Production	Coal	34.8 / 23.2	23.8 / 28.0
Electricity Production	Anthracite	28.8 / 19.2	26.3 / 30.0
Electricity Production	Lignite	46.8 / 31.2	25.4 / 27.1
Electricity Production	Natural Gas	75.6 / 50.4	13.8 / 15.2
Electricity Production	Distillate Oil	43.2 / 28.8	19.0 / 20.5
Electricity Production	Residual Oil	39.6 / 26.4	19.2 / 23.4
Electricity Production	Liquified Petroleum Gas (LPG)	33.6 / 22.4	15.9 / 17.9
Electricity Production	Process Gas	39.6 / 26.4	11.3 / 19.3
Electricity Production	Coke	25.2 / 16.8	25.9 / 29.2
Electricity Production	Distillate Oil (Diesel)/Diesel	1114 / 743	19.3 / 20.8
Electricity Production	Oil	43.2 / 28.8	17.8 / 21.8
Electricity Production	Jet Fuel	901 / 601	18.4 / 19.9
Electricity Production	Refinery Gas	39.6 / 26.4	11.3 / 19.3
Industrial	Bituminous Coal	300 / 200	*
Industrial	Subbituminous Coal	412 / 274	*
Industrial	Bituminous/Subbituminous Coal	355 / 237	*
Industrial	Coal	347 / 231	*
Industrial	Natural Gas	97.2 / 64.8	*
Industrial	Anthracite	28.8 / 19.2	*
Industrial	Waste Oil	16.8 / 11.2	18.0 / 22.1
Industrial	Distillate Oil	43.2 / 28.8	*
Industrial	Residual Oil		*
Industrial	Liquified Petroleum Gas (LPG) [nonpoint/point]	101 / 67.8° 43.4 / 28.9	*
Industrial	Coke [nonpoint/point]	25.2 / 16.8 43.4 / 28.9	*
Industrial	Process Gas	39.6 / 26.4 120 / 80.3	*
Industrial	Kerosene	44.4 / 29.6	18.8 / 20.3
Industrial	Jet Fuel	648 / 432	*
Industrial	Gasoline	72984 / 48656	18.3 / 20.0
Industrial	distillate oil (no 2)	*	19.0 / 20.5
Industrial	Distillate Oil (Diesel)	57.6 / 38.4	*
Industrial	Refinery Gas	39.6 / 26.4	*
Industrial	jet A fuel	648 / 423	18.4 / 19.9
Industrial	jet naptha	648 / 432	18.9 / 20.5
Industrial	Oil	*	*
Industrial	blast furnace gas	7498 / 3610	41.5 / 71.0
Industrial	coke oven gas	2479 / 1194	8.18 / 13.9
Industrial	propane	42.5 / 28.3°	16.0 / 18.1
Commercial	Anthracite Coal	*	*
Commercial	Bituminous Coal	300 / 200	"
Commercial	Subbituminous Coal	412 / 274	"
Commercial	Bituminous/Subbituminous Coal	355 / 237	"
Commercial	Coal	636 / 424	"
Commercial	Natural Gas	97.2 / 64.8	"
Commercial	Distillate Oil	*	"
Commercial	Residual Oil	*	"
Commercial	Liquified Petroleum Gas (LPG) [nonpoint/point]	102 / 67.8° 25.5 / 17.0	"
Commercial	Kerosene	*	"
Commercial	Diesel [nonpoint/point]	43.8 / 29.2° 111 / 74.3	*
Commercial	Gasoline	*	"



Commercial	Jet fuel	*	"
Commercial	Process gas	*	"
Commercial	Propane	25.2 / 16.8°	"
Commercial	Anthracite culm	14.4 / 9.62	"
Residential	Bituminous Coal	13729 / 9153	"
Residential	Subbituminous Coal	18846 / 12564	"
Residential	Bituminous/Subbituminous Coal	16288 / 10858	"
Residential	Coal	15886 / 10590	"
Residential	Anthracite	13234 / 8822	"
Residential	Natural Gas [nonpoint/point]	46.5 / 31.0° 79.6 / 50.4	"
Residential	Distillate Oil	*	"
Residential	Residual Oil	*	"
Residential	Liquified Petroleum Gas (LPG)	25.5 / 17.0	"
Residential	Kerosene	*	"
Residential	Propane	*	"
Railroad	Diesel	514 / 343°	"
Railroad	Distillate Oil (diesel)	973 / 649	"
Marine vessels	Diesel	514 / 343°	"
Marine vessels	Residual Oil	39.6 / 26.4	"
Marine vessels	Gasoline	72984 / 48656	"
Nonroad	Gasoline	72984 / 48656	"
Nonroad	Distillate Oil (diesel)	1115 / 743	"
Nonroad	Liquified Petroleum Gas (LPG)	33.6 / 22.4	"

### 2.1.3 Point data

The point emissions represent facilities with a physically identifiable emission “stack” or point location and exceed a specific criteria air pollution threshold (USEPA 2015c). The NEI point source data files are primarily comprised of processes associated with the industrial and airport sectors but emissions from the commercial, railroad, nonroad, and electricity production sectors are present as well (USEPA 2015a).

A number of key fields that define a point location for the purposes of the Vulcan FFCO<sub>2</sub> emissions estimation within the point database and include the state and county FIPS code, the “state facility identifier” (which identifies the individual emitting facility) and the tribal code (used in place of the FIPS in tribal lands). Each site or facility can have multiple emission points (different “stacks”), units (different buildings or portions of a complex facility or site), or emission processes (e.g. energy production, heaters, engines). Some of the emitting points/units/processes can have different geocoded locations and these are retained in the Vulcan processing. Hence, exact latitude and longitude is critical for allocation to the physical US landscape. Corrections to location information were made in urban domains associated with the Hestia Project : the Los Angeles Basin, Baltimore, Salt Lake City, and Indianapolis (e.g. Gurney et al., 2018; 2019b).

Each point emission record is also associated with an SCC which is used to retrieve the needed CO and CO<sub>2</sub> EFs to enact the same procedure outlined in the description of the nonpoint source processing. In the case of the point sources, no self-reported EFs are supplied. Separation is first made between airport point sources (processing of which is described in a later section) and non-airport point sources. The non-airport point sources are matched to a CO EF via the SCC from the EPA’s WebFIRE EF database as the first choice for the CO EF. Where no match is found, default CO EF values are used, themselves archived from literature review (see Table 2) and determined through a combination of the sector and fuel.



1 All point source emission records designated as industrial, railroad, and nonroad are distributed to hourly temporal  
 2 resolution from the 2011 annual total using SCC-specific temporal surrogate profiles provided by the EPAs  
 3 Clearinghouse for Inventories and Emissions Factors (CHIEF) (USEPA, 2015c). The temporal surrogate profiles are  
 4 constructed from monthly, weekly and diurnal cycles (data available at:  
 5 [ftp://newftp.epa.gov/air/emismod/2011/v3platform/ancillary\\_data/ge\\_dat\\_for\\_2011v3\\_temporal.zip](ftp://newftp.epa.gov/air/emismod/2011/v3platform/ancillary_data/ge_dat_for_2011v3_temporal.zip)). These  
 6 temporal surrogates are comprised of three cyclic time profiles (diurnal, weekly, monthly) specific to SCC that are  
 7 combined to generate hourly SCC-specific time fractions for an entire calendar year. Records which do not have an  
 8 SCC match are distributed as a constant hourly emission.

#### 9 *Uncertainty*

10 Point source uncertainty is applied to the reported CO emissions, the CO EF, and the CO<sub>2</sub> EF. For the reported CO  
 11 emissions, an uncertainty value of  $\pm 7.8\%$  is used, a value reported by Gately et al. (2017) for the industrial and  
 12 commercial sectors (which dominate the point sources) and based on a state-scale difference between ACES and  
 13 EIA state industrial+commercial fuel consumption. We interpret this as a 95% confidence interval given that this is  
 14 estimated from a measure of difference by Gately et al. (2017).

15 For the default EF uncertainty, the CO and CO<sub>2</sub> EFs were adjusted in combination in a fashion similar to that  
 16 described in the nonpoint source section and the same percentage numerical boundaries described there were used.  
 17 For the records that use the WebFIRE CO EFs, an uncertainty value of  $\pm 20\%$  is used for the 95% CI bounds.

#### 18 **2.1.4 Electricity Production**

19 Three sources of data are used to estimate the FFCO<sub>2</sub> emissions at electricity production facilities, all are geocoded  
 20 to a physical location. The first is the Environmental Protection Agency's Clean Air Markets Division (CAMD) data  
 21 (USEPA, 2015b). The second is the Department of Energy's Energy Information Administration (EIA) reporting  
 22 data (DOE/EIA, 2003). The third is the reporting done within the NEI point source reporting (described previously).  
 23 Overlap exists between these three data sources (corrected in the processing here) which is corrected according to  
 24 the prioritization in the order listed above. A detailed comparison made between the CAMD and EIA FFCO<sub>2</sub>  
 25 emissions along with greater detail regarding data sources, data processing and procedures can be found in Gurney  
 26 et al. (2016).

27 The CAMD data is collected under the Acid Rain Program (ARP), which was instituted in 1990 under Title IV of  
 28 the Clean Air Act (CFR 2008; USEPA 2005b; USEPA 2010). Though the CAMD dataset does not include all power  
 29 plants in the US, it accounts for a very large proportion. The CAMD data used in Vulcan are reported as hourly CO<sub>2</sub>  
 30 emissions monitored from an emitting stack or through a calculation, based on records of fuel consumption  
 31 (<ftp://ftp.epa.gov/dmdnload/emissions/hourly/monthly/>). The annual reporting is also used for additional information  
 32 related to the facility (<http://ampd.epa.gov/ampd>).

33 The EIA dataset is derived from the EIA reporting form 923, which reports monthly data on receipts and cost of  
 34 fossil fuel, fuel stocks, generation, consumption of fuel for generation, and environmental data at each power plant



1 (<http://www.eia.gov/electricity/data/eia923>). Fuel consumption is reported as a heat input value (e.g. british thermal  
 2 units). CO<sub>2</sub> emission factors are then utilized to calculate the quantity of CO<sub>2</sub> emitted. In order to maintain  
 3 consistency with the data source, the CO<sub>2</sub> emission factors used by the EIA are adopted to estimate the FFCO<sub>2</sub>  
 4 emissions from these facilities (DOE, 2011).

5 Some manual corrections are performed to the geocoordinates of both the CAMD and EIA electricity production  
 6 data, as a result of searching in Google Earth or via alternative online information resources (e.g. utility websites).

7 A hierarchy was employed given that there was overlap between the two datasets. This was performed at the unit  
 8 level given that a single facility might have individual power units reporting to CAMD and another only reporting to  
 9 the EIA. Where overlap did exist at this scale, preference was made to retain the CAMD data. Further details and  
 10 rationale can be found in Gurney et al. (2016).

11 The CAMD reporting data is archived at the hourly temporal scale and directly used in Vulcan. The EIA electricity  
 12 production reporting is resolved at the monthly scale. This is transformed into hourly reporting using a “flat” time  
 13 profile or a constant level such that the monthly integral matches the reported monthly emissions data. The  
 14 electricity production facilities reported in the NEI as point sources also use a flat time profile but instead of  
 15 distribution over each of the reported months, the emissions are held constant over an entire year.

16 Table 4 provides a summary of the electricity production data totals for the three data sources.

17 **Table 4: Summary information for 2011 electricity production facilities in Vulcan version 3.0.**

Data source	Number of facilities	Total FFCO <sub>2</sub> emissions (MtC/year)
CAMD	1479	592.1
EIA	2255	40.00
NEI	11832	8.87
Total	15566	641.0

## 18 *Uncertainty*

19 Gurney et al., (2016) found that one-fifth of US power plants had monthly FFCO<sub>2</sub> emission differences exceeding -  
 20 6.4%/+6.8% for the year 2009 (the closest analyzed year to the 2011 base year presented here). The emissions  
 21 distribution of the two datasets were not normally distributed nor were the differences. Hence, a typical gaussian  
 22 uncertainty estimate cannot be made – rather, the difference distribution was represented by quintiles of percentage  
 23 difference. Hence, these values cannot be cast within the context of other normally-distributed errors. However, we  
 24 conservatively consider the quintile value (the positive and negative tails) as a one-sigma value and ±13% as a 95%  
 25 CI boundary value.

## 26 **2.1.5 Onroad**

27 County scale FFCO<sub>2</sub> emissions are retrieved from the 2011 EPA NEIv1 onroad results (USEPA, 2011). The 2011  
 28 NEI onroad results report emissions for every US county by 13 vehicle types (designating vehicle class and fuel)  
 29 and 12 road types, including urban and rural distinctions. It is based on simulations using the Motor Vehicle



1 Emissions Simulator (MOVES) model with inputs supplied to a county database (CDB) by SLT agencies (USEPA,  
 2 2012; USEPA, 2015a). Version 1.0 of the 2011 NEI includes 1,363 CDB submissions out of a total of 3,234  
 3 counties. In order to generate results for all counties in the US, the EPA used multiple data and modeling tools to  
 4 estimate county-specific FFCO<sub>2</sub> emissions including identifying “representative” counties among the data supplied  
 5 by SLT agencies to best match those there were not reported.

6 The state of California did not report FFCO<sub>2</sub> to the 2011 NEI. Hence, the Vulcan onroad FFCO<sub>2</sub> emissions for  
 7 California used the 2011 results from the Emissions FACtors 2014 model (EMFAC2014), produced by the  
 8 California Air Resources Board (CARB, 2014). The EMFAC2014 model estimates vehicle miles traveled (VMT)  
 9 and FFCO<sub>2</sub> emissions for 27 vehicle types (reduced here to 13 via aggregation) using emissions rates  
 10 (FFCO<sub>2</sub>/distance traveled) and data on the California vehicle fleet and activity statistics such as VMT, speed  
 11 distributions, and idle times (CARB, 2015). Distribution to sub-state scales uses annual vehicle counts from the  
 12 Highway Performance Monitoring System (HPMS). The HPMS is a spatial road network database managed by the  
 13 FHWA to monitor and record Average Annual Daily Traffic (AADT) counts (FHWA, 2014). By considering  
 14 vehicle registration in combination with the HPMS data, EMFAC also accounts for inter-regional travel.

15 County-scale FFCO<sub>2</sub> emissions for all US states are spatially assigned to road segments via a road basemap that best  
 16 represents the entirety of the road surface occupied by onroad vehicles. Vulcan uses a combination of the 2011  
 17 Highway Performance Monitoring System (HPMS, 2017) road network and Open Street Map (OSM;  
 18 <http://download.geofabrik.de/>) road network. The Census Urbanized Areas boundary  
 19 (<https://www.fhwa.dot.gov/policyinformation/hpms/shapefiles.cfm>) was used to assign an urban/rural distinction to  
 20 each of the 7 original HPMS road classes making them compatible with the onroad NEI road classes (supplementary  
 21 material, Table S4).

22 The distribution of the county-scale road/vehicle-specific FFCO<sub>2</sub> emissions along the complete length of road class  
 23 in a county, is achieved through the use of the 2011 AADT data from the FHWA’s HPMS  
 24 (<http://www.fhwa.dot.gov/policyinformation/hpms/shapefiles.cfm>; state scale data files were used). AADT counts  
 25 are collected using short-term and continuous counting methods. Most data are collected by individual states and  
 26 reported to the FHWA, but some data are also collected by the FHWA directly. Very little AADT data was collected  
 27 on local roads (urban local, rural local). For those segments in our merged basemap that do not have an AADT  
 28 value, gap-filling was used (see supplementary material for details on gap-filling methods).

29 With a complete US map of AADT values and road segment length, the vehicle miles traveled (VMT) can be  
 30 estimated. The fraction of a non-local road class-specific road segment’s VMT within a county acts as the  
 31 distribution means to allocate county-scale onroad FFCO<sub>2</sub>. For local roads, given the paucity of AADT data, the  
 32 fraction of a road segment’s length out of all local roads within a county acts as the allocation method. Hence, the  
 33 local roads have no spatial gradients along the local roads (at the sub-county scale). However, there are FFCO<sub>2</sub>  
 34 emissions gradients in space that are determined by the spatial density of local roads.

35 In order to use the spatial distribution methods employed by the Vulcan system and be compatible with the NEI  
 36 results for the other US states, the vehicle class/county-specific California onroad FFCO<sub>2</sub> emissions must be



1 translated to the 6 vehicle classes and 14 road classes (7 of urban and rural sub-types) in the NEI. This is performed  
 2 via the use of Federal Highway Administration (FHWA) state-scale VMT data by road class and the proportion of  
 3 VMT by vehicle class. Details are provided in the supplementary material.

4 Carrying out the spatialization procedure across all counties in the United States, it became clear that there were  
 5 some mismatches between NEI road class VMT and the AADT on the HPMS road network. For example, there  
 6 were instances in which onroad FFCO<sub>2</sub> emissions were present in a county for a particular road class, but for which  
 7 no AADT data existed and vice-versa. These mismatches could be due to the demarcation of urban versus rural  
 8 roads. As noted previously the roads were divided into urban and rural classes based on the US Census Urbanized  
 9 Areas. This may differ from the choices made when state officials were generating the county database inputs for the  
 10 EPA (if the NEI estimate uses state-supplied data in the MOVES onroad emissions estimate, for example). While  
 11 the HPMS AADT data has an urban code, we used the US Census Urbanized Areas to divide a road classes so that  
 12 the urban/rural classification would be consistent between the OSM and HPMS basemaps.

13 In cases where emissions were reported for a road class in NEI, but for which there were no physical roads in our  
 14 AADT gap-filled basemap, the emissions reported in NEI were moved to the next closest road class with AADT.  
 15 The closest road class is the urban or rural counterpart within the same class-size, and the second-closest being the  
 16 road class that is the next class-size down. In cases where AADT was present for a road class, but no NEI FFCO<sub>2</sub>  
 17 emissions were reported for that road class, FFCO<sub>2</sub> emissions were redistributed from the next closest road class,  
 18 proportional to VMT. For example, if the NEI reports emissions for urban interstates, but VMT was estimated for  
 19 both urban and rural interstates, then the NEI reported emissions would be redistributed from urban interstates to  
 20 rural interstates proportional to the VMT in each road class.

21 In the state of California, the EMFAC results were crosswalked from county-scale, vehicle class-specific FFCO<sub>2</sub>  
 22 emissions to totals that include road class. These FFCO<sub>2</sub> emissions were distributed onto road segments in the same  
 23 manner as done for other states. However, unlike other states, there were no cases in which the EMFAC onroad  
 24 FFCO<sub>2</sub> emissions needed to be “shuffled” to partner road classes.

25 Hourly traffic volume data for the years 2011-2013 were obtained from the FHWA Continuous Count Stations  
 26 (CCS) dataset (previously known as the Automatic Traffic Recorder; ATR) (Jessberger, 2016). The CCS stations  
 27 measure hourly traffic volume at a fixed location in space and we use the station’s latitude and longitude as a unique  
 28 station identifier. Corrections were made to the Connecticut station coordinates based on data from the Connecticut  
 29 Department of Transportation (<http://www.ct.gov/dot/cwp/view.asp?a=1383&q=330402>).

30 For each station, the direction(s) and lane(s) of traffic are recorded but are aggregated to estimate the total traffic  
 31 volume moving through a station across all lanes and directions. The result is a single measure of traffic volume per  
 32 hour that is traveling through a station in any direction and on any lane with a unique location. Any station with  
 33 greater than six months total (either contiguous or not) of missing traffic monitoring data, were removed from the  
 34 dataset. This left a total of 5106 traffic volume monitoring stations in the year 2011, 5172 in 2012 and 5527 in 2013.  
 35 2011 contained 141 stations that were not present in either 2012 or 2013. 2012 contained 57 stations that were not  
 36 present in either 2011 or 2013. 2013 contained 511 stations that were not present in either 2011 or 2012. Each year






1 of the traffic monitoring data (for which there are no instances of gaps exceeding six months) are gap-filled  
2 individually, maintaining the cyclic integrity of the hour of day and day of the week. Details are provided in the  
3 supplementary material.

4 After combining the 2011, 2012 and 2013 CCS data into a single average year dataset, there were a total of 6047  
5 CCS measurement locations including Alaska and Hawaii. There are 5890 stations in the Continental US and these  
6 are used for the construction of the temporal profiles.

7 In order to distribute the temporal distribution measured at the gap-filled CCS measurement stations to all road  
8 segments in the US, interpolation/extrapolation of the traffic patterns is required. Given the paucity of traffic  
9 measurement stations relative to the total area of the US landscape and the fact that the temporal distribution of  
10 traffic is less related to road class than space, it was determined to aggregate the eight road classes to four,  
11 “temporal” road classes for purposes of spatial interpolation. There is evidence that interstates have unique traffic  
12 patterns from all other road classes due to the preponderance of interstate trucking commerce. Furthermore,  
13 interstate usage in cities is a mix of passenger vehicles and commercial trucking while rural interstates are  
14 dominated by commercial trucking. Hence, the road classes chosen for the purposes of temporal interpolation were:  
15 rural interstate, urban interstate, rural non-interstate, and urban non-interstate. Figure 1 shows the CCS measurement  
16 locations aggregated to these four temporal road classes.



**Figure 1: Distribution of CCS measurements stations separated into four road classes. a) rural interstate; b) urban interstate; c) rural non-interstate; d) urban non-interstate.**

Inverse Distance Weighted (IDW) interpolation was performed for each of the four temporal road classes separately, and only for grid cells that are occupied by roads of that road class. The two inputs are the gap-filled CCS traffic data, and the locations of road segments for each of the four road classes. The IDW used the default number of neighbors (all neighbors), and the default power function (2), making this an inverse distance squared method.

#### *Uncertainty*

The uncertainty in the onroad sector uses the results from Gately et al., (2017) which, in turn, references Gately et al., (2013) and Mendoza et al., (2013). This uncertainty was estimated at  $\pm 7.1\%$  for a presumed 1-sigma uncertainty. Here, we have assigned  $\pm 14.2\%$  to the 95% CI boundaries for all road types.



#### 1    **2.1.6    Nonroad**

2    The nonroad sector CO<sub>2</sub> emissions estimates are retrieved from the 2011 EPA NEIv2 which uses the NONROAD  
 3    model to estimate emissions ([ftp://ftp.epa.gov/EmisInventory/2011/2011neiv2\\_nonroad\\_byregions.zip](ftp://ftp.epa.gov/EmisInventory/2011/2011neiv2_nonroad_byregions.zip)) across a  
 4    large number of mobile sources that travel “off-road” (USEPA, 2015a) except locomotives, airplanes and  
 5    commercial marine vessels (CMV) which are taken up in separate sections in this document. The NONROAD  
 6    model results, in turn, are based on output from the National Mobile Inventory Model (NMIM) which relies on data  
 7    inputs from the National County Data base (NCD) (USEPA 2005c; 2005d). Both the NMIM and the NCD were  
 8    described previously (Gurney et al., 2009). The EPA updated data within the NCD from 12 SLT agencies along with  
 9    EPA default values to generate the results in the 2011 NEIv2 (for a description of these updates see  
 10    [ftp://ftp.epa.gov/EmisInventory/2011/doc/2011neiv2\\_supdata\\_nonroad](ftp://ftp.epa.gov/EmisInventory/2011/doc/2011neiv2_supdata_nonroad)).

11    As with the onroad sector, California presents a special case. The CO emissions are reported comprehensively using  
 12    California’s OFFROAD model ([www.arb.ca.gov/msei/offroad/offroad.htm](http://www.arb.ca.gov/msei/offroad/offroad.htm)) but no CO<sub>2</sub> was reported. Hence, we  
 13    scaled the California CO emissions by the mean SCC-specific CO<sub>2</sub>/CO ratio from all other US counties.

14    Spatial distribution uses the spatial surrogates generated by the EPA reflecting a series of spatial representations  
 15    such as the mines, golf course and agricultural land (The shapefiles can be found here:  
 16    [ftp://ftp.epa.gov/EmisInventory/emiss\\_shp2003/us/](ftp://ftp.epa.gov/EmisInventory/emiss_shp2003/us/) or  
 17    [ftp://ftp.epa.gov/EmisInventory/2011v6/v1platform/spatial\\_surrogates/shapefiles/](ftp://ftp.epa.gov/EmisInventory/2011v6/v1platform/spatial_surrogates/shapefiles/)). There were instances in which  
 18    nonroad FFCO<sub>2</sub> emissions could not be associated with a spatial entity due to missing data. These emissions are  
 19    spatialized by first aggregating all the unassociated sub-county emission elements to the county scale for a given  
 20    spatial shape (e.g., golf courses, mines) and then distributing these emissions evenly across the county.

21    The sub-annual temporal distribution of the nonroad FFCO<sub>2</sub> emissions uses SCC-specific temporal surrogate  
 22    profiles provided by the EPAs Clearinghouse for Inventories and Emissions Factors (CHIEF) (USEPA, 2015c). The  
 23    temporal surrogate profiles are constructed from monthly, weekly and diurnal cycles (data available at:  
 24    [ftp://newftp.epa.gov/air/emismod/2011/v3platform/ancillary\\_data/ge\\_dat\\_for\\_2011v3\\_temporal.zip](ftp://newftp.epa.gov/air/emismod/2011/v3platform/ancillary_data/ge_dat_for_2011v3_temporal.zip)).

25    These temporal surrogates are comprised of three cyclic time profiles (diurnal, weekly, monthly) specific to SCC  
 26    that are combined to generate hourly SCC-specific time fractions for an entire calendar year. There are 5 SCC codes  
 27    present in the NEI 2011 nonroad data file but not found in the temporal surrogate files (2260006035, 2265006035,  
 28    2267006035, 2270006035, 2268006035) - these were given a “flat” or constant time profile in the absence of any  
 29    specified temporal distribution.

#### 30    *Uncertainty*

31    Nonroad records other than those derived from the point source data files (which follow the point source uncertainty  
 32    estimation described in the point source section) are assigned a 95% CI boundary of  $\pm 3.8\%$  for the FFCO<sub>2</sub> emission  
 33    value. This was derived from examination of the range of carbon content and fuel density uncertainties as outlined in  
 34    EPA (2017), Annex 2, page A-86. This is consistent with the point source uncertainty for nonroad distillate fuel  
 35    consumption.



### 2.1.7 Airport

As described in the point source section, the airport FFCO<sub>2</sub> emissions are estimated from the 2011 NEI point source reporting for CO. The emission factors used (Table 5) convert the reported CO emissions to FFCO<sub>2</sub> and are specific to aircraft class and fuel, consistent with the reporting in the NEI which often listed multiple processes (aircraft class/fuel) for a single airport facility. The fuel type implied by the CO<sub>2</sub> EF values uses jet fuel except where explicitly indicated in the SCC description (NG, LPG, diesel, gasoline).

**Table 5: SCC, description, CO EF and CO<sub>2</sub> EF values for the airport sources.**

SCC	description	CO EF <sup>r</sup> (lbs/e6btu)	CO <sub>2</sub> EF (tC/e6btu)
2275060011	Aircraft /Air Taxi /Piston	0.751	0.019
2275060012	Aircraft /Air Taxi /Turbine	0.751	0.019
2275070000	Aircraft /Aircraft Auxiliary Power Units /Total	0.5396	0.019
2268008005	Airport Ground Support Equipment, CNG	0.081	0.014
2267008005	Airport Ground Support Equipment, LPG	0.085	0.017
2270008005	Airport Ground Support Equipment, Diesel	0.828	0.020
2265008005	Airport Ground Support Equipment, 4-Stroke Gasoline	30.34	0.019
2275020000	Aircraft /Commercial Aircraft /Total: All Types	1.083	0.019
2275050011	Aircraft /General Aviation /Piston	0.751	0.019
2275050012	Aircraft /General Aviation /Turbine	0.751	0.019
2275001000	Aircraft /Military Aircraft /Total	1.083	0.019
27505011	Aircraft /Civil /Jet Engine: Jet A	1.083	0.019
27505001	Aircraft /Civil /Piston Engine: Aviation Gas	0.751	0.019
27502011	Aircraft /Commercial /Jet Engine: Jet A	1.083	0.019
27501015	Aircraft /Military /Jet Engine: JP-5	1.083	0.019
2275060011	Aircraft /Air Taxi /Piston	0.751	0.019

<sup>r</sup> CO emission factors are retrieved from the Intergovernmental Panel on Climate Change Guidelines on National Greenhouse Gas Inventories (IPCC 2006). The values reflect a 50/50 mixture of new versus old aircraft fleet characteristics in addition to a 67/33 mixture of domestic and international flight characteristics. The emission factors for domestic aviation have been derived from an average of a number of typical aircraft. For domestic aircraft, the average fleet is represented by Airbus A320, Boeing 727, Boeing 737-400, Mc Donnell Douglas DC9 and MD 80 aircraft. The old fleet is represented by Boeing B737 and Mc Donnell Douglas DC9. For international traffic the average fleet is represented by Airbus A300, Boeing 767, B747 and Mc Donnell Douglas DC10, whilst the old fleet is represented by the Boeing B707, Boeing 747 and Mc Donnell Douglas DC8.


The airport FFCO<sub>2</sub> emissions are only associated with the taxi & takeoff/landing sequences. FFCO<sub>2</sub> emissions associated with non-aircraft processes such as building operations and non-aircraft mobile sources are reported as emissions in other sectors (e.g. commercial, nonroad). The airports are geocoded to the airport location in the NEI though some manual adjustments have been made to the original coordinates using manual inspection in Google Earth. The emission point, in these instances, is placed in the middle of the central runway.

Temporal distribution of the FFCO<sub>2</sub> airport emissions use a series of datasets. The Los Angeles World Airports (LAWA) dataset reports hourly flight volume for three airports in the LA Basin domain: Los Angeles International airport (LAX), Ontario airport (ONT), and Van Nuys airport (VNY) (Hastings, 2014). The Operations Network (OPSNET) dataset from the FAA reports total date-specific, daily flight volume (365 values) at specific airports (<https://aspm.faa.gov/opsnet/sys/Default.asp>). An hourly time profile was constructed by combining the LAWA diurnal profile and the OPSNET annual profile. The three LAWA airports constituted the diurnal cycle (Figure 2) at all US airports with the LAX assigned to international airports, the ONT to non-international airports and the VNY to local airports.

Airports were matched with a Federal Aviation Administration (FAA) international airport database (FAAINTL) by airport code to determine whether an airport is international



- 1 ([https://hub.arcgis.com/datasets/4782d6f5aa844591a16d46df635b7af4\\_1](https://hub.arcgis.com/datasets/4782d6f5aa844591a16d46df635b7af4_1)). Airports which could not be matched to
- 2 the OPSNET data by airport code/airport name were assigned a temporal invariant (“flat”) hourly time structure.



3  
 4 **Figure 2: Average hourly flight volume fractions at LAX, VNY and ONT**

5 FAAINTL, OPSNET, and two additional airport databases (the National Airport Atlas (NAA;  
 6 <https://catalog.data.gov/dataset/airports-of-the-united-states-direct-download>) and AIRNAV; [www.airnav.com](http://www.airnav.com))  
 7 were used to determine whether an airport was an airport or a helipad. The name/code of each airport was searched  
 8 in these airport databases. An airport which could not be identified in any of the aforementioned airport databases  
 9 would be categorized as a helipad. A temporally invariant time structure was applied to all helipads.

10 A portion of the Vulcan v3.0 CMV FFCO<sub>2</sub> emissions would be considered “bunker” fuel combustion (i.e. consumed  
 11 as part of international travel) under the IPCC reporting methodology within the UNFCCC process. Vulcan does not  
 12 separate bunker from non-bunker fuel consumption and a portion of the airport sector emissions (particularly  
 13 international air flights) would be considered as such were the IPCC reporting categorization applied here. No  
 14 attempt has been made to limit or separately report airport emissions that would be considered part of the bunker  
 15 fuel definition.

#### 16 *Uncertainty*

17 The uncertainty in the airport sector is derived from the point source processing as described previously (magnitude  
 18 and EF-based uncertainty) except that the CO<sub>2</sub> EFs are specific to the mix of aviation fuels associated with the  
 19 emission records and are based on uncertainty estimation from the USEPA (2017), Annex 2, pages 85 & 89.

#### 20 **2.1.8 Railroad**

21 The FFCO<sub>2</sub> emissions associated with railway activity are derived from the 2011 NEIv2 CO emissions reporting  
 22 which, in turn, were developed for the 2008 NEI (ERG 2011) and scaled to 2011 values (ERG 2012). Emissions  
 23 related to the railroad sector were reported as a mixture of nonpoint and point emissions and hence, these were  
 24 managed separately but combined when represented as spatial entities. The CO emissions were converted to FFCO<sub>2</sub>  
 25 following the procedures outlined in the nonpoint and the point sections, respectively.



1 The two NEI source categories imply different spatial representations, however. The point source railroad emissions  
 2 are associated with rail yards and related geo-specific locales and are placed in space according to the provided  
 3 latitude and longitude. The railroad FFCO<sub>2</sub> emissions associated with the nonpoint NEI reporting contain an ID  
 4 variable that links to a spatial element (rail line segment) in the EPA railroad GIS shapefile  
 5 ([https://www.epa.gov/sites/production/files/2015-06/railway\\_20140730.zip](https://www.epa.gov/sites/production/files/2015-06/railway_20140730.zip)). A large number of railroad emission  
 6 records have no railroad segment match and are spatialized using freight statistics described in supplementary  
 7 material.

8 The annual railroad FFCO<sub>2</sub> emissions are distributed to the hourly timescale with no additional temporal structure (a  
 9 “flat” time distribution), unless they originated from point source data for which the SCC-specific time profiles,  
 10 previously described, are used.

#### 11 *Uncertainty*

12 The uncertainty for the railroad emissions is directly inherited from the uncertainty estimation described for the  
 13 point and nonpoint source processing, respectively. The only difference related to the the CO magnitude uncertainty  
 14 ( $\pm 3.8\%$ ) which was derived from examination of the range of carbon content and fuel density uncertainties outlined  
 15 in EPA (2017), Annex 2, page A-86 for distillate fuels, the dominant fuel used in railroad.

#### 16 **2.1.9 Commercial Marine Vessels**

17 The FFCO<sub>2</sub> emissions associated with commercial marine vessels (CMV) rely on nonpoint NEIv2 CO emissions  
 18 reporting and follow the same emission factor-related conversion outlined in the nonpoint source section. CMV  
 19 includes vessels directly or indirectly involved in commerce or military activity. The emissions encompass  
 20 maneuvering, hoteling, cruise and reduced speed zone travel and are specific to geographically located ports and  
 21 shipping lanes that extend 12 nautical miles from the US shoreline. Private or “pleasure” craft are not included as  
 22 part of the CMV emissions but are captured in the nonroad reporting. As with the nonroad reporting, the EPA used a  
 23 mixture of SLT data submissions and default values, in collaboration with the Office of Transportation and Air  
 24 Quality to generate an estimate of CO emissions for CMV. A portion of the Vulcan v3.0 CMV FFCO<sub>2</sub> emissions  
 25 would be considered “bunker” fuel combustion (i.e. consumed as part of international trade) under the IPCC  
 26 reporting methodology within the UNFCCC process. Vulcan does not separate bunker from non-bunker fuel  
 27 consumption and a portion of the CMV sector emissions (particularly ship travel directed towards international  
 28 waters) would be considered as such were the IPCC reporting categorization applied here. No attempt has been  
 29 made to limit or separately report CMV emissions that would be considered part of the bunker fuel definition.

30 The spatialization utilized the EPA shapefiles that delineate US ports and US shipping lanes through spatial IDs  
 31 associated with the emission records ([https://www.epa.gov/sites/production/files/2015-06/ports\\_20140729.zip](https://www.epa.gov/sites/production/files/2015-06/ports_20140729.zip);  
 32 [https://www.epa.gov/sites/production/files/2015-06/shippinglanes\\_072914.zip](https://www.epa.gov/sites/production/files/2015-06/shippinglanes_072914.zip)). In the instance that no spatial entity  
 33 is identified for an emission record, a simple spatial alternative is employed whereby all the unlinked port (or



1 “underway”) emissions are summed within a county and evenly distributed to the shapes that are identified within  
 2 that county (either ports or shipping lanes).

3 The CMV sector has no data allowing for the designation of hourly time structure. Hence, the emissions are  
 4 temporally invariant over all hours of the year (“flat” distribution).

#### 5 *Uncertainty*

6 The uncertainty of the CMV emissions is directly inherited from the uncertainty estimation described for the point  
 7 and nonpoint source processing, respectively. The only difference related to the the CO magnitude uncertainty  
 8 ( $\pm 10.0\%$ ) which was derived from examination of the range of carbon content and fuel density uncertainties outlined  
 9 in EPA (2017), Annex 2, page A-87 for residual fuels, the dominant fuel used in CMV.

#### 10 **2.1.10 Cement**

11 CO<sub>2</sub> is emitted from cement manufacturing as a result of fuel combustion and as process-derived emissions  
 12 (Andrew, 2018). The emissions from fuel combustion are captured in the point source reporting. The process-  
 13 derived CO<sub>2</sub> emissions result from the chemical process that converts limestone to calcium oxide and CO<sub>2</sub>. This  
 14 occurs during “clinker” production (clinker is the raw material for cement which is produced by grinding the clinker  
 15 material).

16 Estimation of CO<sub>2</sub> emissions from clinker production utilizes two datasets. The first is the data provided by the  
 17 Portland Cement Association which provides the annual clinker capacity at individual facilities, postal addresses,  
 18 facility name, zip code and contact phone numbers (PCA, 2006). The capacity data reflects conditions for the  
 19 calendar year 2006. The other dataset utilized is the Minerals Yearbook produced by the United States Geological  
 20 Survey which provides the capacity factor (or percent utilization of capacity) on a statewide or multi-state basis  
 21 (some states are quantified individually, others are part of an aggregate) (USGS 2013). The product of capacity and  
 22 the capacity factor provides an estimate of clinker production.

23 Clinker production for 2011 is scaled from the Vulcan version 2.0 (CY 2002) estimate (Gurney et al., 2009) using  
 24 the relative annual capacity factor. The CO<sub>2</sub> emission factor used in the Vulcan Project is 0.59 metric tonnes  
 25 CO<sub>2</sub>/short ton of clinker produced (IPCC, 2006).

26 The geolocation for each of the individual facilities was achieved by entering the PCA document’s facility address  
 27 into Google Earth and visually inspecting the scene for the primary emitting stack of the cement facility. This  
 28 approach succeeded in locating all 105 facilities present in the PCA document.

29 The EPA estimates cement manufacturing in 2011 to account for 32.2 MtCO<sub>2</sub>/year (USEPA 2017). These estimates,  
 30 in turn, are based upon throughput estimates from the U.S. Geological Survey. Vulcan estimates a total of 34.6  
 31 MtCO<sub>2</sub>/year which compares well with the cement manufacturing estimate from the EPA.

32 The cement sector has no data allowing for the designation of hourly time structure. Hence, the emissions are evenly  
 33 distributed over all hours of the year (a “flat” distribution).





## 1 *Uncertainty*

2 The uncertainty in the cement emissions sector is currently prescribed as +/- 10% for the 95% CI. We use a  
 3 comparison between the facility-scale sum of clinker production in a state and the USGS state throughput (estimated  
 4 from the capacity factor and capacity). The mean percentage difference across all states and multistate aggregates  
 5 was 9.8%, which was rounded to 10% and interpreted as a 95% CI value.

## 6 **2.2 Multiyear estimation**

7 The multiyear (2010-2015) results were achieved using scale factors constructed from the EIA State Energy Data  
 8 System (SEDS) database (<http://www.eia.gov/state/seds/>). Ratios were constructed relative to the year 2011 in all  
 9 SEDS sector/fuel designations for each US state. The crosswalk from the EIA SEDS codes to a sector/fuel  
 10 designation is provided in supplementary material, Table S8.

11 Exceptions to the use of the EIA SEDS database were made for the electricity production, railroad and CMV  
 12 multiyear scaling. Electricity production FFCO<sub>2</sub> emissions are monitored on an hourly basis for all the output  
 13 derived from the CAMD data (92.4% of the total electricity production emissions) and on a monthly basis for all of  
 14 the EIA reported data (6.2% of the total electricity production emissions). The remaining NEI reported electricity  
 15 production emissions (1.4% of the total electricity production emissions) use the EIA SEDS multiyear ratios.

16 In the case of the railroad sector, state-scale EIA specific to distillate fuel oil sales to the railroad sector was used  
 17 ([http://www.eia.gov/dnav/pet/pet\\_cons\\_821dsta\\_a\\_epd0\\_val\\_mgal\\_a.htm](http://www.eia.gov/dnav/pet/pet_cons_821dsta_a_epd0_val_mgal_a.htm)) to construct the year-to-year ratios  
 18 relative to 2011. This data is used in generating the results in the EIA SEDS database but is aggregated and thus not  
 19 as specific to the railroad sector as needed. Large year-over-year ratio values were found for a few individual years  
 20 in low-population states (Nevada, Rhode Island, New Mexico, Hawaii). Values that exceeded 5.0 were replaced by  
 21 the year-specific US average ratio.

22 The procedure for the CMV FFCO<sub>2</sub> emissions is similar but combines the EIA data on distillate fuel oil sales for  
 23 “vessel bunkering use” ([http://www.eia.gov/dnav/pet/pet\\_cons\\_821dsta\\_a\\_epd0\\_vab\\_mgal\\_a.htm](http://www.eia.gov/dnav/pet/pet_cons_821dsta_a_epd0_vab_mgal_a.htm)) with residual  
 24 fuel oil sales for transportation ([http://www.eia.gov/dnav/pet/pet\\_cons\\_821rsda\\_a\\_eppr\\_vat\\_mgal\\_a.htm](http://www.eia.gov/dnav/pet/pet_cons_821rsda_a_eppr_vat_mgal_a.htm)). As with  
 25 the railroad sector application, large year-over-year ratios were filtered (those exceeding 5.0 were replaced by the  
 26 US national average).

27 The ratio values are applied to the annual totals in each of the sector/fuel categories specific to the state FIPS code to  
 28 generate a multiyear time series.

## 29 **3. Results**

30 Annual sector totals are provided in Table 6 for the 2010-2015 time period. Across all sectors, 2012 is the year with  
 31 the least emissions (1529.4 MtC/yr; 95% CI: 1249-1846 MtC/yr). While 2010 was the largest year for total  
 32 emissions (1638.2; 95% CI: 1338-1977 MtC/yr), the maximum value was primarily due to large FFCO<sub>2</sub> emissions in




the electricity production sector. The total FFCO<sub>2</sub> emissions (plus cement) in 2015, the most recent year in the time series, were 1543.7 MtC/year (95% CI: 1268, 1857), a decline driven almost entirely by electricity production FFCO<sub>2</sub> emissions. Electricity production is the largest emitting sector in all years, followed by the onroad and industrial sectors, respectively.

**Table 6: Annual sector specific FFCO<sub>2</sub> (and cement) emission totals for the United States, 2010-2015, estimated by Vulcan v3.0. (units: MtC/year)**

Sector\Year	2010	2011	2012	2013	2014	2015
Residential	92.0	89.2	78.5	91.3	95.3	88.0
Commercial	63.0	62.9	57.1	63.4	66.8	68.5
Industrial	230.6	228.4	227.2	233.8	237.4	231.4
Elec Prod	667.3	641.0	604.3	609.0	609.2	574.4
Onroad	452.0	440.6	436.6	443.1	448.6	452.4
Nonroad	63.6	62.7	61.8	62.9	64.0	64.6
Airport	19.8	19.6	20.5	22.3	22.3	21.8
Rail	11.9	11.9	12.6	13.7	15.1	14.6
CMV	28.4	23.3	20.9	18.5	16.2	18.1
Cement	9.5	9.7	9.8	9.8	9.8	9.8
Total	1638.2 (1338, 1977)	1589.3 (1299, 1917)	1529.4 (1249, 1846)	1567.9 (1284, 1889)	1584.6 (1300, 1908)	1543.7 (1268, 1857)

The order of the 2011 FFCO<sub>2</sub> emitting sectors (Figure 3) varies regionally (US Census Regions) with the electricity production sector accounting for the largest share in the Midwest (44%) and South (46%) while onroad emissions account for the largest share in the West (32%) and Northeast (29%). The sum of the commercial and residential sectors are a larger share of total emissions in the Northeast (22%) than in the other three regions (6%-11%). The industrial FFCO<sub>2</sub> emissions account for the largest industrial share in the West (19%) compared to the other three regions (13%-14%). Overall, 2011 FFCO<sub>2</sub> emissions are largest in the South (652 TgC), followed by the Midwest (434 TgC), the West (293 TgC) and the Northeast (200 TgC).




1  
2 **Figure 3: Sector-specific percentage share of 2011 Vulcan v3.0 FFCO<sub>2</sub> emissions for the United States by US**  
3 **Census Region: a) Northeast; b) Midwest; c) South; d) West.**

4 When examined at the state-scale, the apportioning of the FFCO<sub>2</sub> emitting sectors shows a relationship to total  
5 FFCO<sub>2</sub> per capita emissions (Figure 4). States with larger per capita emissions tend to be dominated by industrial  
6 and electricity production sector FFCO<sub>2</sub> emissions. States with lower per capita total FFCO<sub>2</sub> emissions tend to have  
7 lesser industrial and electricity production FFCO<sub>2</sub> emissions and a greater share of onroad and  
8 residential/commercial emissions. A few states are notable exceptions to this pattern. For example, the states of  
9 Alaska, Washington, and South Dakota have a relatively large portion of nonroad emissions while Rhode Island and  
10 Washington DC have a relatively large proportion of commercial sector FFCO<sub>2</sub> emissions. Tabular results at the  
11 state-scale are provided in supplementary material, Table S9.

12 Per capita emissions vary across the states, with the largest in the state of Wyoming (38.5 tC/person) and the  
13 smallest in Washington DC (2.11 tC/person) and California (2.81 tC/person). The median total per capita FFCO<sub>2</sub>  
14 emissions at the county-scale are 3.80 tC/person (see Figure S3 in supplementary material). It is worth noting that




- 1 the population statistics used here define the population as that residing within the state which will influence the
- 2 results for Washington DC where there is a large daytime non-resident population.



3  
 4 **Figure 4: Vulcan v3.0 FFCO<sub>2</sub> emissions sector share (left y-axis: %) by state and per capita FFCO<sub>2</sub> emissions**  
 5 **(right y-axis: tC/person) for year 2011.**


- 6 The Vulcan FFCO<sub>2</sub> emissions are quantified at the sub-national scale according to three general shape types: points
- 7 (e.g. electricity production, industrial point reporting), lines (e.g. onroad) and polygons (e.g. nonroad, residential).
- 8 For use in atmospheric transport modeling and ease of use in analysis, these results are gridded using a 1km x 1km
- 9 regular grid (Figure 5a). The importance of urban areas is clearly demonstrated in the complete US mapped
- 10 landscape along with the greater urbanization in the eastern half of the country and along the West coast. Interstates
- 11 and other large primary roadways are also evident across the US connecting large population centers. Normalization
- 12 by population (Landsat, 2017) offers a dramatically different perspective on U.S. FFCO<sub>2</sub> emissions, placing
- 13 greater emphasis on the western half of the country (Figure 5b).




**Figure 5: Vulcan v3.0 2011 FFCO<sub>2</sub> emissions for the United States. a) absolute emissions (1km x 1km resolution, tC); b) per capita emissions (0.1° x 0.1° resolution, tC; different resolution and projection required for integration with population data). Copyright © National Geographic Society, i-cubed.**

A center of mass (CoM) is a useful and compact metric to understand and illustrate the spatial changes in fossil fuel CO<sub>2</sub> emissions over time (Gregg et al., 2009). The CoM summarizes the distribution of emissions in the same way as the mean summarizes a probability distribution (Asefi et al., 2014). Figure 6 shows both the multiyear and monthly mean CoM. The multiyear CoM shows a general shift from the East to the West over the six years examined here with the CoM located in Missouri approximately 70 miles SW of St Louis, MO. The monthly mean results show a tendency to move along a NE/SW axis with wintertime movement towards the NE driven by greater heating needs associated with cold/continental conditions. Summertime movement is towards the SW associated with the rising air-conditioning demand during summer months. The months of May/June/July show movement towards the SE in May and June, a shift towards the North in July, before resuming the Western shift in August and September.

The 2011 monthly FFCO<sub>2</sub> emissions magnitude exhibits two maxima of roughly equal value over the course of the year: a winter maximum in the months of December and January and a summer maximum in the months of July and August. The maxima correspond to the northermost CoM position in the winter and near-southernmost CoM position in summer which are associated with the demand for heating in the winter, dominated by more northerly locations, and the demand for cooling in the summer, dominated by more southerly locations.



2  
3 **Figure 6: Vulcan v3.0 FFCO<sub>2</sub> emissions center of mass estimate. Red line/symbols denote 2010–2015 annual**  
4 **time series. Purple line/symbols denote monthly mean FFCO<sub>2</sub> emissions. This map was made in ArcMap™ by**  
5 **Esri using the OpenStreetMap basemap layer (Copyright © Esri, with data from OpenStreetMap**  
6 **contributors ©).**



7  
8 **Figure 7. Vulcan v3.0 2011 FFCO<sub>2</sub> emissions for the United States by month with 95% confidence interval.**  
9 **Units: TgC/month.**






#### 1    4. Discussion

2    The Vulcan approach to quantification of bottom-up granular FFCO<sub>2</sub> emissions established a method that has been  
3    since followed by other investigators with useful and instructive variations (e.g. Bun et al., 2019; Gately et al.,  
4    2017). Some of the differences are driven by differing national circumstances related to data availability and  
5    collection sources. However, other than the ACES data product, which covers only the NorthEast US domain, there  
6    is no other US-based granular estimate of FFCO<sub>2</sub> emissions with which to evaluate the results presented here. As  
7    noted in the introduction, however, numerous global gridded estimates of FFCO<sub>2</sub> emissions have been constructed  
8    starting in the 1990s. Currently only the ODIAC estimate is quantified at the same 1km x 1km resolution as found in  
9    Vulcan. Hence, we perform comparison to the ODIAC output over the Vulcan domain in the hope of providing  
10   insight into one or both of the emission estimates. We masked the ODIAC output with a mask that includes all land  
11   surface gridcells and all gridcells offshore for which Vulcan possesses a non-zero emission value. We estimate the  
12   ODIAC emissions to be 1453.7 TgC/year for the year 2011. The same mask applied to Vulcan results in FFCO<sub>2</sub>  
13   emissions of 1553.8 TgC/yr or a difference of 100.1 MtC/yr (7.6%). We also removed all CMV emissions from  
14   Vulcan due to the fact that the ODIAC 1km x 1km data product does not include any bunker fuels in the emissions.  
15   We make no adjustment to the Vulcan airport emissions, though a portion is also likely in the bunker fuel category.  
16   The inability to precisely isolate the bunker fuel amounts from Vulcan will result in comparison uncertainties but  
17   these are considered small relative to the scale at which the comparison is made.

18   At the individual gridcell spatial scale, further detail on differences between the two data products can be examined  
19   (Figure 8). Three different relationships appear in the spatial gridcell comparison with an correlation coefficient of  
20   0.69 and a slope of 0.43 (change in ODIAC/change in Vulcan). The first shows good correlation close to the 1:1 line  
21   for large emitting gridcells. These are gridcells dominated by power production facilities and hence, traced to  
22   common regulatory data reporting in the two data products. The second relationship evident in the paired gridcell  
23   comparison shows rough correspondence whereby Vulcan has a larger range of emission values to a narrower  
24   ODIAC range rotated clockwise from the 1:1 line. There is also a well-defined lower threshold of emissions in  
25   ODIAC (~11 tC/yr), likely tied to the threshold associated with low levels of nighttime lighting, a dominant driver  
26   of the ODIAC spatial distribution (Liang et al., 2019). The third discrete relationship is a non-correlated collection  
27   of paired gridcells in the upper range of ODIAC emissions for which the Vulcan counterparts exhibit midrange  
28   emission values.





**Figure 8. Comparison of log-transformed ODIAC (y-axis) and Vulcan v3.0 (x-axis) FFCO<sub>2</sub> emissions (units: Natural log tC).**

When presented explicitly in space, total ODIAC and Vulcan FFCO<sub>2</sub> emissions show similar spatial patterns at the domain-wide scale, characterized by large concentrations in urban centers across the US landscape, particularly along the Northeastern seaboard and the upper Midwest (Figure 9a, 9c). ODIAC exhibits large numbers of gridcells in rural areas across the Western U.S. with no emission value, likely due to the lack of a nighttime light signal in those areas. This is further demonstrated by the emission histogram (Figure 9b, 9d) whereby ODIAC has a distinct lower cutoff at 11.02 tC/yr (natural log of which is 2.4) compared to Vulcan which has a more continuously declining low value distribution. The maximum emission frequency bin for ODIAC is centered at 23.3 tC/yr whereas the equivalent value for Vulcan is 12.8 tC/yr. Vulcan gridcells in these areas have emission values but they are small in comparison to more populated areas and can be dominated by nonroad emissions which use large spatial proxies for distribution. In estimating the gridcell-scale relative emissions difference (GRD), these pairs are excluded. GRD values are high throughout the populated portions of the U.S., particularly in the Eastern half of the country. There are large spatially continuous areas in which ODIAC emissions exceed Vulcan and vice-versa. Large differences occur in urban centers, most notably in the Western U.S., with Vulcan often exceeding ODIAC emissions in the urban core but ODIAC exceeding Vulcan outside of the urban core in these cities. (e.g. Phoenix, Dallas, Los Angeles, St Louis).




1 To provide an average relative difference between the two data products, we calculate the gridcell absolute median  
2 relative difference,  $GAMRD$ , the median of a set of individual paired gridcell relative differences, where the  
3 differences are represented in absolute units (i.e. so all GRD values are positive).  $GAMRD$  is calculated as,

$$4 \quad GAMRD = med \left\{ \frac{abs(E_i^A - E_i^V)}{\frac{E_i^A + E_i^V}{2}} \right\} \quad (1)$$

5 where  $E$  represented emissions for the ODIAC ( $A$ ) and Vulcan ( $V$ ) for each  $i^{th}$  paired gridcell. We only include  
6 gridcell pairs in which neither of the emission values is zero. We find that the  $GAMRD$  between the ODIAC and  
7 Vulcan v3.0 FFCO<sub>2</sub> emissions at the 1 km x 1 km spatial scale is 80.04%.

8



**Figure 9.** Comparison of the ODIAC and Vulcan total FFCO<sub>2</sub> emissions as mapped distributions (left) and frequency histograms (right) for contiguous U.S. only: (a,b) ODIAC total FFCO<sub>2</sub> emissions; (c,d) Vulcan total FFCO<sub>2</sub> emissions; (e,f) GRD values ((ODIAC-Vulcan)/Vulcan) (Values larger than 99% and smaller than -99% were excluded from the GRD frequency distribution).




1 The other means by which to assess the Vulcan results is via comparison to recent work using  $^{14}\text{CO}_2$  measurements  
2 and an atmospheric inversion approach by Basu et al. (2019). In that study, the mean of the ensemble of atmospheric  
3  $\text{CO}_2$  inversion estimates was within 1.4% of the total US Vulcan estimate in the year 2011.


4 The increased resolution of the Vulcan v3.0 FFCO<sub>2</sub> emissions data product (1 km x 1 km) in comparison to the  
5 previous 10 km x 10 km Vulcan v2.0 data product raises the prospect of supplying information that resolves sub-city  
6 spatial scales (e.g. neighborhood) in a comprehensive fashion across the entire U.S. landscape. At this resolution,  
7 most urbanized areas in the U.S. would comprise domains much larger than the 1 km x 1 km resolution and, hence,  
8 have sub-domain information emissions content. In this way, the Vulcan v3.0 FFCO<sub>2</sub> emissions data product offers a  
9 scope 1, high-resolution inventory estimate for every urbanized area in the U.S. (Figure 10).

10  
11

12  
13







e) f)  
**Figure 10: Vulcan v3.0 FFCO<sub>2</sub> emissions for a selected group of U.S. urban areas. a) Washington DC; b) New York; c) Boston; d) Houston; e) San Francisco; f) Chicago. These maps were made in ArcMap™ by Esri using the World Imagery basemap layer (Copyright © Esri).**

After adopting the US census “urbanized area” boundary definition (<https://www.census.gov/programs-surveys/geography/guidance/geo-areas/urban-rural/2010-urban-rural.html>), total US FFCO<sub>2</sub> emissions within these urban area boundaries come to 45.1% of the total Vulcan FFCO<sub>2</sub> contiguous US emissions in 2011 (709.6 TgC). We narrow urban emissions to the sum of residential, commercial and onroad in an effort to eliminate emissions sectors that are often historically artifactual to location within a given urban area (e.g. power plants, industrial facilities) and hence, less directly related to urban residents and their emitting activities. The sum of these three sectors within the urbanized area boundary accounts for 65% of the these same three sectors at the national scale, slightly less than the proportion of the US population within the urbanized area boundary, 73%. This indicates that for the sum of the residential, commercial and onroad sectors, urban residents emit less per capita than non-urban residents in the contiguous US.

## 5 Data availability, policy and future updates

The sector-specific Vulcan v3.0 annual gridded emissions data product can be downloaded from the Oak Ridge National Laboratory Distributed Active Archive Center (ORNL DAAC) (<https://doi.org/10.3334/ORNLDAAC/1741>) and is distributed under Creative Commons Attribution 4.0 International (CC-BY 4.0, <https://creativecommons.org/licenses/by/4.0/deed.en>). The Vulcan v3.0 FFCO<sub>2</sub> emissions data product is provided in annual 1 km x 1 km NetCDF file formats, one file for each of the 6 years (2010–2015). The annual files vary in size (by sector) with the largest individual file being approximately 50 GB. Separate files are available for Alaska and the contiguous United States (Gurney et al., 2019).

Attempts will be made to update the Vulcan FFCO<sub>2</sub> emissions on a roughly bi-annual basis, depending upon support and the availability of data sources described in this study.



## 6 Conclusions

Fossil fuel carbon dioxide (FFCO<sub>2</sub>) emissions, spanning the years 2010-2015, at a spatial scale of 1km x 1km and an hourly temporal scale have been completed for the United States under the Vulcan Project. These Vulcan version 3.0 emissions are constructed through a bottom-up approach, gathering and combining data from multiple sources such as CO emissions reporting, direct flux measurements, and traffic monitoring. We describe the complete Vulcan version 3.0 methodology here, sector-by-sector in addition to the methods for uncertainty estimation.

We estimate FFCO<sub>2</sub> emissions for the year 2011 of 1589.3 TgC with a 95% confidence interval of 1299/1917 TgC (+18.3%/-20.6%), implying a one-sigma uncertainty of  $\sim \pm 10\%$ . The order of the 2011 FFCO<sub>2</sub> emitting sectors shows the electricity production sector accounting for the largest share in the Midwest (44%) and South (46%) while onroad emissions account for the largest share in the West (32%) and Northeast (29%). Overall, 2011 FFCO<sub>2</sub> emissions are largest in the South (652 TgC), followed by the Midwest (434 TgC), the West (293 TgC) and the Northeast (200 TgC).

We find that per capita FFCO<sub>2</sub> emissions are larger in states proportionately dominated by the electricity production and industrial sectors and smaller in states proportionately dominated by onroad and residential/commercial building emissions. The center of mass (CoM) of FFCO<sub>2</sub> emissions in the US are located in the state of Missouri with mean seasonality that extends towards the Northeast in the wintertime then moves towards the Southwest in the summer, likely reflecting the seasonal demand for heating and air-conditioning. Comparison to ODIAC, a global gridded FFCO<sub>2</sub> emissions estimate shows large differences in both total emissions (100.1 TgC for year 2011) and spatial patterns. The spatial correlation ( $R^2$ ) between the two data products was 0.38 and the mean absolute difference at the individual gridcell scale was 80.04%.

The Vulcan v3.0 FFCO<sub>2</sub> emissions data product offers an immediate high-resolution estimate of emissions in every city within the U.S., providing a large potential savings of time and effort for cities planning to develop self-reported city inventories. Research associated with comparison to existing self-reported urban inventories and the addition of indirect FFCO<sub>2</sub> emissions (scope 2 & 3) are underway.

**Author Contributions.** KRG conceived of the content, performed portions of the data collection, coding and analysis, and wrote the paper. JL, RP, YS, and JH performed portions of the data collection, coding and analysis. GR performed analysis.

**Competing Interests.** The authors declare that they have no conflict of interest.

**Financial Support.** This research was made possible through support from the National Aeronautics and Space Administration grant NNX14AJ20G and the NASA Carbon Monitoring System program, Understanding User Needs for Carbon Information project (subcontract 1491755).



## 1 References

- 2 Andres, R. J., Marland, G., Fung, I., & Matthews, E.: A  $1^\circ \times 1^\circ$  distribution of carbon dioxide emissions from fossil  
3 fuel consumption and cement manufacture, 1950-1990. *Global Biogeochemical Cycles*, 10(3), 419–429.  
4 <https://doi.org/10.1029/96GB01523>, 1996.
- 5 Andres, R. J., Fielding, D. J., Marland, G., Boden, T. A., Kumar, N., & Kearney, A. T.: Carbon Dioxide Emissions  
6 from Fossil Fuel Use, 1751-1950. *Tellus Series B-Chemical and Physical Meteorology*, 51(4), 759–765.  
7 <https://doi.org/10.1034/j.1600-0889.1999.t01-3-00002.x>, 1999.
- 8 Andres, R. J., Boden, T. A., Bréon, F., Ciais, P., Davis, S., Erickson, D., ... Treanton, K.: A synthesis of carbon  
9 dioxide emissions from fossil-fuel combustion. *Biogeosciences Discussions*, 9(1), 1299–1376.  
10 <https://doi.org/10.5194/bgd-9-1299-2012>, 2012.
- 11 Andrew, R. M.: Global CO<sub>2</sub> emissions from cement production, *Earth System Science Data*, 10, 195–217.  
12 <https://doi.org/10.5194/essd-10-195-2018>, 2018.
- 13 Asefi-Najafabady, S., Rayner, P. J., Gurney, K. R., McRobert, A., Song, Y., Coltin, K., ... Baugh, K.: A multiyear,  
14 global gridded fossil fuel CO<sub>2</sub> emission data product: Evaluation and analysis of results. *J. Geophys. Res.*, 119(17),  
15 10,213–10,231. <https://doi.org/10.1002/2013JD021296>, 2014.
- 16 Baldasano, J. M., Güereca, L. P., López, E., Gassó, S., & Jimenez-Guerrero, P.: Development of a high-resolution (1  
17 km × 1 km, 1 h) emission model for Spain: The High-Effective Resolution Modelling Emission System (HERMES).  
18 *Atmospheric Environment*, 42(31), 7215–7233. <https://doi.org/10.1016/j.atmosenv.2008.07.026>, 2008.
- 19 Basu, S., S.J. Lehman, J.B. Miller, A.E. Andrews, C. Sweeney, K.R. Gurney, P.P. Tans: Constraints on Fossil Fuel  
20 CO<sub>2</sub> Emissions from Measurements of <sup>14</sup>C in Atmospheric CO<sub>2</sub>, submitted to *Proc. Nat.Acad. Sci.*, 2019.
- 21 Bellassen, V., Stephan, N., Afriat, M., Alberola, E., Barker, A., Chang, J. P., ... Shishlov, I.: Monitoring, reporting  
22 and verifying emissions in the climate economy. *Nature Climate Change*, 5(4), 319–328.  
23 <https://doi.org/10.1038/nclimate2544>, 2015.
- 24 Bun, R., Gusti, M., Kujii, L., Tokar, O., Tsybrivskyy, Y., & Bun, A.: Spatial GHG inventory: Analysis of  
25 uncertainty sources. A case study for Ukraine. *Accounting for Climate Change: Uncertainty in Greenhouse Gas*  
26 *Inventories - Verification, Compliance, and Trading*, 63–74. [https://doi.org/10.1007/978-1-4020-5930-8\\_6](https://doi.org/10.1007/978-1-4020-5930-8_6), 2007.
- 27 Bun, R., Nahorski, Z., Horabik-Pyzel, J., Danylo, O., See, L., Charkovska, N., ... Kinakh, V.: Development of a  
28 high-resolution spatial inventory of greenhouse gas emissions for Poland from stationary and mobile sources.  
29 *Mitigation and Adaptation Strategies for Global Change*. <https://doi.org/10.1007/s11027-018-9791-2>, 2018.
- 30 Cai, B., Liang, S., Zhou, J., Wang, J., Cao, L., Qu, S., ... Yang, Z.: China high resolution emission database  
31 (CHRED) with point emission sources, gridded emission data, and supplementary socioeconomic data. *Resources*,  
32 *Conservation and Recycling*, 129 (November 2017), 232–239. <https://doi.org/10.1016/j.resconrec.2017.10.036>,  
33 2018.
- 34 California Air Resources Board (2014) EMFAC2014 Volume I – User’s Guide, v1.0.7, April 30, 2014, California  
35 Environmental Protection Agency Air Resources Board, Mobile Source Analysis Branch, Air Quality Planning &  
36 Science Division.
- 37 California Air Resources Board (CARB) “EMFAC2014 Volume III – Technical Documentation” (Publication  
38 v1.0.7, CARB, 2015). Data available from <https://www.arb.ca.gov/emfac/2014/>.
- 39 Code of Federal Regulations (2008), Protection of Environment, Environmental Protection Agency, 40 CFR Part 75,  
40 Continuous Emission Monitoring, Code of Federal Regulations, Revised as of January 24, 2008.  
41 [https://www.ecfr.gov/cgi-bin/text-](https://www.ecfr.gov/cgi-bin/text-idx?SID=4719db7a48cd26050b0732d0f9adc3ad&mc=true&node=pt40.2.51&rgn=div5)  
42 [idx?SID=4719db7a48cd26050b0732d0f9adc3ad&mc=true&node=pt40.2.51&rgn=div5](https://www.ecfr.gov/cgi-bin/text-idx?SID=4719db7a48cd26050b0732d0f9adc3ad&mc=true&node=pt40.2.51&rgn=div5). AERR summary:  
43 <https://www.epa.gov/air-emissions-inventories/air-emissions-reporting-requirements-aerr#rule-summary>
- 44 Commercial Building Energy Consumption Survey (2016) 2012 CBECS microdata files and information, U.S.  
45 Energy Information Administration. Data retrieved from:  
46 <https://www.eia.gov/consumption/commercial/data/2012/index.php?view=microdata> (Aug 1, 2018).





- 1 Cooke, W.F., C. Lioussé, C., Cachier, H., Feichter, J.: Construction of a  $1^\circ \times 1^\circ$  fossil fuel emission data set for  
2 carbonaceous aerosol and implementation and radiative impact in the ECHAM4 model, *J. Geophys. Res.*, 104 (D18),  
3 22137–22162, 1999.
- 4 Davis, S.J. and Caldeira, K.: Consumption-based accounting of CO<sub>2</sub> emissions, *Proc. Nat. Acad. USA*, doi:  
5 10.1073/pnas.0906974107, 2010.
- 6 Denier van der Gon, H. A. C., Kuenen, J. J. P., Janssens-Maenhout, G., Döring, U., Jonkers, S., & Visschedijk, A.:  
7 TNO\_CAMS high resolution European emission inventory 2000–2014 for anthropogenic CO<sub>2</sub> and future years  
8 following two different pathways. *Earth System Science Data Discussions*, (November), 1–30.  
9 <https://doi.org/10.5194/essd-2017-124>, 2017.
- 10 Department of Energy/Energy Information Administration (2003) *Electric Power Monthly March 2003*, Energy  
11 Information Administration, Office of Coal, Nuclear, and Alternate Fuels, U.S. Department of Energy, Washington  
12 D.C. 20585, DOE/EIA-0226 (2003/03).
- 13 Department of Energy/Energy Information Administration (2011) *Electric Power Annual 2010*, November 2011,  
14 available at [http://www.eia.gov/cneaf/electricity/epa/epa\\_sum.html](http://www.eia.gov/cneaf/electricity/epa/epa_sum.html).
- 15 Department of Energy/Energy Information Administration (2018) State Energy Consumption Estimates 1960  
16 through 2016, DOE/EIA-0214(2016), June 2018, Washington DC.
- 17 Durant, A.J., LeQuere, C., Hope, C., Friend, A.D.: Economic value of improved quantification in global sources and  
18 sinks of carbon dioxide, *Phil. Trans. R. Soc. A*, 369, 1967–1979, 2011.
- 19 Eastern Research Group, Inc.: *Introduction to Area Source Emissions Inventory Development*, Volume III, Chapter  
20 1, prepared for: Area Sources Committee, Emission Inventory Improvement Program, January 2001.
- 21 Eastern Research Group: *Documentation for Locomotive Component of the National Emissions Inventory*  
22 *Methodology*, prepared by: Eastern Research Group, ERG No.: 0245.02.401.001, Contract No.: EP-D-07-097, 2011.
- 23 Eastern Research Group: *Development of 2011 Railroad Component for National Emissions Inventory*,  
24 Memorandum from Heather Perez, Susan McClutchey, and Richard Billings/ERG, to Laurel Driver/US EPA,  
25 September 5, 2012.
- 26 Federal Emergency Management Agency (2017) HAZUS database. Retrieved from:  
27 <https://www.fema.gov/summary-databases-hazus-multi-hazard> (Aug 1, 2018).
- 28 Federal Highway Administration, Highway Statistics Series, *Highway Statistics 2011, User's Guide*, Policy and  
29 Government Affairs, Office of Highway Policy Information. Available at:  
30 <https://www.fhwa.dot.gov/policyinformation/statistics/2011/userguide.cfm> (2011).
- 31 Federal Highway Administration, Highway Statistics Series, *Highway Statistics 2013, User's Guide*, Policy and  
32 Government Affairs, Office of Highway Policy Information. Available at:  
33 <https://www.fhwa.dot.gov/policyinformation/statistics/2013/userguide.cfm> (2013).
- 34 Federal Highway Administration, *Highway Performance Monitoring System: Field Manual*, Office of Highway  
35 Policy Information, Office of Management & budget (OMB) Control No. 2125-0028, March 2014.  
36 <https://www.fhwa.dot.gov/policyinformation/hpms/shapefiles.cfm>
- 37 Gately, C. K., Hutyrá, L. R., Wing, I. S., & Brondfield, M. N.: A bottom up approach to on-road CO<sub>2</sub> emissions  
38 estimates: Improved spatial accuracy and applications for regional planning. *Environmental Science & Technology*,  
39 47(5), 2423–2430. <https://doi.org/10.1021/es304238v>, 2013.
- 40 Gately, C. K., & Hutyrá, L. R.: Large Uncertainties in Urban-Scale Carbon Emissions. *Journal of Geophysical*  
41 *Research: Atmospheres*, 122(20), 11,242–11,260. <https://doi.org/10.1002/2017JD027359>, 2017.
- 42 Gaubert, B., Stephens, B. B., Basu, S., Chevallier, F., Deng, F., Kort, E. A., ... Yin, Y.: Global atmospheric CO<sub>2</sub>  
43 inverse models converging on neutral tropical land exchange but disagreeing on fossil fuel and atmospheric growth  
44 rate. *Biogeosciences*, 16, 117–134. <https://doi.org/10.5194/bg-16-117-2019>, 2019.
- 45 Ghosh, T., Elvidge, C. D., Sutton, P. C., Baugh, K. E., Ziskin, D., & Tuttle, B. T.: Creating a global grid of  
46 distributed fossil fuel CO<sub>2</sub> emissions from nighttime satellite imagery. *Energies*, 3(12), 1895–1913.  
47 <https://doi.org/10.3390/en3121895>, 2010.



- 1 Gregg, J. S., & Andres, R. J.: A method for estimating the temporal and spatial patterns of carbon dioxide emissions
- 2 from national fossil-fuel consumption. *Tellus, Series B: Chemical and Physical Meteorology*, 60 B(1), 1–10.
- 3 <https://doi.org/10.1111/j.1600-0889.2007.00319.x>, 2008.
- 4 Gregg, J. S., Losey, L. M., Andres, R. J., Blasing, T. J., & Marland, G.: The temporal and spatial distribution of
- 5 carbon dioxide emissions from fossil-fuel use in North America. *Journal of Applied Meteorology and Climatology*,
- 6 48(12), 2528–2542. <https://doi.org/10.1175/2009JAMC2115.1>, 2009.
- 7 Gurney, K. R., Law, R. M., Denning, A. S., Rayner, P. J., Baker, D., Bousquet, P., ... Yuen, C. W.: Towards robust
- 8 regional estimates of CO<sub>2</sub> sources and sinks using atmospheric transport models. *Nature*, 415(6872), 626–630.
- 9 <https://doi.org/10.1038/415626a>, 2002.
- 10 Gurney, K. R., Chen, Y. H., Maki, T., Kawa, S. R., Andrews, A., & Zhu, Z.: Sensitivity of atmospheric CO<sub>2</sub>
- 11 inversions to seasonal and interannual variations in fossil fuel emissions. *Journal of Geophysical Research D:*
- 12 *Atmospheres*, 110(10), 1–13. <https://doi.org/10.1029/2004JD005373>, 2005.
- 13 Gurney, K.R., Mendoza, D., Zhou, Y., Fischer, M., de la Rue du Can, S., Geethakumar, S., Miller, C.: The Vulcan
- 14 Project: High resolution fossil fuel combustion CO<sub>2</sub> emissions fluxes for the United States, *Environ. Sci. Technol.*,
- 15 43(14), 5535–5541, doi:10.1021/es900806c, 2009.
- 16 Gurney, K. R., Razlivanov, I., Song, Y., Zhou, Y., Benes, B., & Abdul-Massih, M.: Quantification of fossil fuel
- 17 CO<sub>2</sub> emissions on the building/street scale for a large U.S. City. *Environmental Science and Technology*, 46(21),
- 18 12194–12202. <https://doi.org/10.1021/es3011282>, 2012.
- 19 Gurney, K. R., Liang, J., O’Keeffe, D. O., Patarasuk, R., Hutchins, M., Huang, J., Rao, P., and Song, Y.:
- 20 Comparison of Global Downscaled Versus Bottom-Up Fossil Fuel CO<sub>2</sub> Emissions at the Urban Scale in Four US
- 21 Urban Areas, *J. Geophys. Res.-Atmos.*, 124, 2823–2840, <https://doi.org/10.1029/2018JD028859>, 2018.
- 22 Gurney, K.R., Huang, J., and Coltin, K.: Bias present in US federal agency power plant CO<sub>2</sub> emissions data and
- 23 implications for the US clean power plan *Env. Res. Lett.*, 11, 064005, 2016.
- 24 Gurney, K.R., J. Liang, R. Patarasuk, Y. Song, J. Huang, G. Roest, G.: Vulcan Fossil Fuel Carbon Dioxide (FFCO<sub>2</sub>)
- 25 Emissions Data Product, version 3.0, 1 km grid, <https://doi.org/10.3334/ORNLDAAAC/1741>, 2019.
- 26 Gurney, K.R., Patarasuk, R., Liang, J., O’Keeffe, D., Rao, P., Song, Y.: The Hestia Fossil Fuel CO<sub>2</sub> Emissions
- 27 Dataset for the Los Angeles Basin, *Earth System Science Data*, 11, 1309–1335, [https://doi.org/10.5194/essd-11-](https://doi.org/10.5194/essd-11-1309-2019)
- 28 [1309-2019](https://doi.org/10.5194/essd-11-1309-2019), 2019b.
- 29 Hastings, Norene: personal communication (2014) Environmental Supervisor, LAWA, Environmental Services
- 30 division, January 2014.
- 31 Highway Performance Monitoring System: [https://catalog.data.gov/dataset/highway-performance-monitoring-](https://catalog.data.gov/dataset/highway-performance-monitoring-system-hpms-national)
- 32 [system-hpms-national](https://catalog.data.gov/dataset/highway-performance-monitoring-system-hpms-national), 2017.
- 33 Hirsch, J. & Associates (2004) *Energy Simulation Training for Design & Construction Professionals*. Retrieved
- 34 from: <http://doe2.com/download/equest/eQuestTrainingWorkbook.pdf> (Aug 1, 2018). eQuest model download
- 35 available from: <http://www.doe2.com/eQuest/> (Aug 1, 2018).
- 36 Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., ... Zhang, Q. (2018).
- 37 Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data
- 38 System (CEDS). *Geoscientific Model Development*. <https://doi.org/10.5194/gmd-11-369-2018>
- 39 Intergovernmental Panel on Climate Change “IPCC guidelines for national greenhouse gas inventories, Prepared by
- 40 the National Greenhouse Gas Inventories Programme” (IGES, Japan, 2006).
- 41 Intergovernmental Panel on Climate Change (2013). Working Group I Contribution to the IPCC Fifth Assessment
- 42 Report, Climate Change 2013: The Physical Science Basis.
- 43 IPCC. (2018). Special Report on 1.5 degrees: Summary for Policymakers. Retrieved from
- 44 [https://www.ipcc.ch/site/assets/uploads/sites/2/2019/05/SR15\\_SPM\\_version\\_report\\_LR.pdf](https://www.ipcc.ch/site/assets/uploads/sites/2/2019/05/SR15_SPM_version_report_LR.pdf)
- 45 Ivanova, D., Vita, G., Steen-Olsen, K., Stadler, K., Melo, P. C., Wood, R., & Hertwich, E. G.: Mapping the carbon
- 46 footprint of EU regions. *Environmental Research Letters*, 12(5). <https://doi.org/10.1088/1748-9326/aa6da9>, 2017.



- 1 Janssens-Maenhout, G., Petrescu, A. M. R., Muntean, M., & Blujdea, V.: Verifying Greenhouse Gas Emissions:
- 2 Methods to Support International Climate Agreements. *Greenhouse Gas Measurement and Management*, 1(2), 132–
- 3 133. <https://doi.org/10.1080/20430779.2011.579358>, 2013.
- 4 Jessberger, Steven: personal communication: Engineer, Federal Highway Administration, Office of Highway Policy
- 5 Information, Travel Monitoring and Surveys, 1200 New Jersey Avenue, S.E., HPPI-30, E83-418, Washington, DC
- 6 20590, USA, 202-366-5052, 202-366-7742), 2016.
- 7 Jones, C. M., & Kammen, D. M.: Quantifying carbon footprint reduction opportunities for U.S. households and
- 8 communities. *Environmental Science and Technology*. <https://doi.org/10.1021/es102221h>, 2011.
- 9 Jones, C., & Kammen, D. M.: Spatial Distribution of U.S. Household Carbon Footprints Reveals Suburbanization
- 10 Undermines Greenhouse Gas Bene fi ts of Urban Population Density. *Environmental Science & Technology*, 48(2),
- 11 895–902. <https://doi.org/10.1021/es4034364>, 2014.
- 12 Kurokawa, J., Ohara, T., Morikawa, T., Hanayama, S., Janssens-Maenhout, G., Fukui, T., ... Akimoto, H.:
- 13 Emissions of air pollutants and greenhouse gases over Asian regions during 2000-2008: Regional Emission
- 14 inventory in ASia (REAS) version 2. *Atmospheric Chemistry and Physics*, 13(21), 11019–11058.
- 15 <https://doi.org/10.5194/acp-13-11019-2013>, 2013.
- 16 This product was made utilizing the LandScan (2017)<sup>TM</sup> High Resolution global Population Data Set copyrighted by
- 17 UT-Battelle, LLC, operator of Oak Ridge National Laboratory under Contract No. DE-AC05-00OR22725 with the
- 18 United States Department of Energy. The United States Government has certain rights in this Data Set.
- 19 LeQuéré, C., Andrew, R., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., ... Zheng, B.: Global Carbon Budget
- 20 2018. *Earth System Science Data*. <https://doi.org/10.5194/essd-10-2141-2018>, 2018.
- 21 Liu, Z., Bambha, R. P., Pinto, J. P., Zeng, T., Boylan, J., Huang, M., ... Michelsen, H. A.: Toward verifying fossil
- 22 fuel CO<sub>2</sub> emissions with the CMAQ model: Motivation, model description and initial simulation. *Journal of the Air*
- 23 *and Waste Management Association*, 64(4), 419–435. <https://doi.org/10.1080/10962247.2013.816642>, 2014.
- 24 Liu, F., Zhang, Q., Tong, D., Zheng, B., Li, M., Huo, H., & He, K. B.: High-resolution inventory of technologies,
- 25 activities, and emissions of coal-fired power plants in China from 1990 to 2010. *Atmospheric Chemistry and*
- 26 *Physics*, 15(23), 13299–13317. <https://doi.org/10.5194/acp-15-13299-2015>, 2015.
- 27 Macknick, J.: Energy and CO<sub>2</sub> emission data uncertainties. *Carbon Management*. Future Science Ltd London, UK.
- 28 <https://doi.org/10.4155/cmt.11.10>, 2011.
- 29 Manufacturing Energy Consumption Survey (2010) 2010 MECS Survey Data, U.S. Energy Information
- 30 Administration. Retrieved from: <https://www.eia.gov/consumption/manufacturing/data/2010/#r10> (Aug 1, 2018).
- 31 Marland, G., Rotty, R. M., & Treat, N. L.: CO<sub>2</sub> from fossil fuel burning: global distribution of emissions. *Tellus B*,
- 32 37 B(4–5), 243–258. <https://doi.org/10.1111/j.1600-0889.1985.tb00073.x>, 1985.
- 33 Mendoza, D., Gurney, K. R., Geethakumar, S., Chandrasekaran, V., Zhou, Y., & Razlivanov, I.: Implications of
- 34 uncertainty on regional CO<sub>2</sub> mitigation policies for the U.S. onroad sector based on a high-resolution emissions
- 35 estimate. *Energy Policy*, 55, 386–395. <https://doi.org/10.1016/j.enpol.2012.12.027>, 2013.
- 36 Minx, J., Baiocchi, G., Wiedmann, T., Barrett, J., Creutzig, F., Feng, K., ... Hubacek, K.: Carbon footprints of cities
- 37 and other human settlements in the UK. *Environmental Research Letters*, 8(3). [https://doi.org/10.1088/1748-](https://doi.org/10.1088/1748-9326/8/3/035039)
- 38 [9326/8/3/035039](https://doi.org/10.1088/1748-9326/8/3/035039), 2013.
- 39 Moran, D., Kanemoto, K., Jiborn, M., Wood, R., Többen, J., & Seto, K. C.: Carbon footprints of 13 000 cities.
- 40 *Environmental Research Letters*, 13(6). <https://doi.org/10.1088/1748-9326/aac72a>, 2018.
- 41 National Research Council: *Verifying Greenhouse Gas Emissions: Methods to Support International Climate*
- 42 *Agreements*. Washington DC: The National Academies Press. <https://doi.org/10.17226/12883>, 2010.
- 43 Oda, T., & Maksyutov, S.: A very high-resolution (1km×1 km) global fossil fuel CO<sub>2</sub> emission inventory derived
- 44 using a point source database and satellite observations of nighttime lights. *Atmospheric Chemistry and Physics*.
- 45 <https://doi.org/10.5194/acp-11-543-2011>, 2011.



- 1 Oda, T., Maksyutov, S., & Andres, R. J.: The Open-source Data Inventory for Anthropogenic CO<sub>2</sub>, version 2016  
2 (ODIAC2016): A global monthly fossil fuel CO<sub>2</sub> gridded emissions data product for tracer transport simulations and  
3 surface flux inversions. *Earth System Science Data*. <https://doi.org/10.5194/essd-10-87-2018>, 2018.
- 4 Ohara, T., Akimoto, H., Kurokawa, J., Horii, N., Yamaji, K., Yan, X., ... Asian, A.: An Asian emission inventory of  
5 anthropogenic emission sources for the period 1980 ? 2020 To cite this version : and Physics An Asian emission  
6 inventory of anthropogenic emission sources for the period 1980 – 2020. *Atmospheric Chemistry and Physics*,  
7 7((16)), 4419–4444, 2007.
- 8 Olivier, J. G. J., Bloos, J. P. J., Berdowski, J. J. M., Visschedijk, A. J. H., & Bouwman, A. F.: A 1990 global  
9 emission inventory of anthropogenic sources of carbon monoxide on 1° × 1° developed in the framework of  
10 EDGAR/GEIA. *Chemosphere - Global Change Science*, 1(1–3), 1–17. [https://doi.org/10.1016/S1465-](https://doi.org/10.1016/S1465-9972(99)00019-7)  
11 [9972\(99\)00019-7](https://doi.org/10.1016/S1465-9972(99)00019-7), 1999.
- 12 Olivier, J. G. J., Van Aardenne, J. A., Dentener, F. J., Pagliari, V., Ganzeveld, L. N., & Peters, J. A. H. W.: Recent  
13 trends in global greenhouse gas emissions: regional trends 1970–2000 and spatial distribution of key sources in 2000.  
14 *Environmental Sciences*, 2(2–3), 81–99. <https://doi.org/10.1080/15693430500400345>, 2005.
- 15 Ou, J., Liu, X., Li, X., Li, M., & Li, W.: Evaluation of NPP-VIIRS nighttime light data for mapping global fossil  
16 fuel combustion CO<sub>2</sub> emissions: A comparison with DMSP-OLS nighttime light data. *PLoS ONE*, 10(9), e0138310.  
17 <https://doi.org/10.1371/journal.pone.0138310>, 2015.
- 18 Patarasuk, R., Gurney, K. R., O’Keeffe, D., Song, Y., Huang, J., Rao, P., ... Ehleringer, J. R.: Urban high-resolution  
19 fossil fuel CO<sub>2</sub> emissions quantification and exploration of emission drivers for potential policy applications. *Urban*  
20 *Ecosystems*, 19(3), 1013–1039. <https://doi.org/10.1007/s11252-016-0553-1>, 2016.
- 21 Peylin, P., Houweling, S., Krol, M. C., Karstens, U., Rödenbeck, C., Geels, C., ... Heimann, M.: Importance of  
22 fossil fuel emission uncertainties over Europe for CO<sub>2</sub> modeling: Model intercomparison. *Atmospheric Chemistry*  
23 *and Physics*, 11(13), 6607–6622. <https://doi.org/10.5194/acp-11-6607-2011>, 2011.
- 24 Petron, G., Tans, P., Frost, G., Chao, D., and Trainer, M.: High resolution emissions of CO<sub>2</sub> from power generation  
25 in the USA, *J. Geophys. Res.*, 113 doi:10.1029/2007/JG000602, 2008.
- 26 Pincetl, S., Chester, M., Circella, G., Fraser, A., Mini, C., Murphy, S., ... Sivaraman, D.: Enabling Future  
27 Sustainability Transitions: An Urban Metabolism Approach to Los Angeles Pincetl et al. Enabling Future  
28 Sustainability Transitions. *Journal of Industrial Ecology*, 18(6), 871–882. <https://doi.org/10.1111/jieec.12144>, 2014.
- 29 Portland Cement Company, Economic Research Department: *U.S. and Canadian Portland Cement Industry Plant*  
30 *Information Summary*, Portland Cement Association, Skokie, IL, 2006.
- 31 Quick, J. C.: Carbon dioxide emission factors for U.S. coal by origin and destination. *Environmental Science and*  
32 *Technology*, 44(7), 2709–2714. <https://doi.org/10.1021/es9027259>, 2010.
- 33 Rayner, P. J., Raupach, M. R., Paget, M., Peylin, P., & Koffi, E.: A new global gridded data set of CO<sub>2</sub> emissions  
34 from fossil fuel combustion: Methodology and evaluation. *Journal of Geophysical Research Atmospheres*, 115(19),  
35 1–11. <https://doi.org/10.1029/2009JD013439>, 2010.
- 36 Residential Energy Consumption Survey (2013) 2009 RECS Survey Data, U.S. Energy Information Administration.  
37 Retrieved from: <https://www.eia.gov/consumption/residential/data/2009/index.php?view=microdata> (Aug 1, 2018).
- 38 Shu, Y., & Lam Nina, N. S. N. (2011). Spatial disaggregation of carbon dioxide emissions from road traffic based  
39 on multiple linear regression model. *Atmospheric Environment*, 45(3), 634–640.  
40 <https://doi.org/10.1016/j.atmosenv.2010.10.037>.
- 41 United States Environmental Protection Agency (1995) *FIRE Version 5.0 Source Classification Codes and Emission*  
42 *Factor Listing For Criteria Air Pollutants*, Office of Air Quality Planning and Standards, Research Triangle Park,  
43 NC 27711, EPA-454/R-95-012, August 1995.
- 44 United States Environmental Protection Agency (2005a), *Emissions Inventory Guidance for Implementation of*  
45 *Ozone and Particulate Matter National Ambient Air Quality Standards (NAAQS and Regional Haze Regulations)*,  
46 Emissions Inventory Group, Emissions, Monitoring and Analysis Division, Office of Air Quality Planning and  
47 Standards, U.S. Environmental Protection Agency, Research Triangle Park, NC 27711, EPA-454/R-05-001, August.



- 1 United States Environmental Protection Agency (2005b), *Plain English Guide to the Part 75 Rule*, U.S.
- 2 Environmental Protection Agency, Clean Air Markets Division.
- 3 United States Environmental Protection Agency (2005c), *EPA's National Mobile Inventory Model (NMIM), A*
- 4 *consolidated emissions modeling system for MOBILE6 and NONROAD*, Office of Transportation and Air Quality,
- 5 Assessment and Standards Division, U.S. Environmental Protection Agency, EPA420-R-05-024, December.
- 6 United States Environmental Protection Agency (2005d), *User's Guide for the Final NONROAD2005 Model,*
- 7 *Assessment and Standards*, Division Office of Transportation and Air Quality U.S. Environmental Protection
- 8 Agency, December.
- 9 United States Environmental Protection Agency (2010) *Draft Part 75 Emissions Monitoring Policy Manual*, U.S.
- 10 Environmental Protection Agency, Clean Air Markets Division, Washington, D.C., April 2010.
- 11 United States Environmental Protection Agency (2011) *2011 National Emissions Inventory, version 1 Technical*
- 12 *Support Document*, June 2014 – Draft. Office of Air Quality Planning and Standards. Retrieved from:
- 13 <https://www.epa.gov/air-emissions-inventories/2011-national-emissions-inventory-nei-technical-support-document>
- 14 (August 12, 2018).
- 15 U.S. Environmental Protection Agency (2012) *Motor Vehicle Emission Simulator (MOVES): User Guide for*
- 16 *MOVES2010b* Office of Transportation and Air Quality, EPA-420-B-12-001b. Retrieved from:
- 17 <https://nepis.epa.gov/Exe/ZyPDF.cgi?Dockey=P100EP28.pdf> (August 12, 2018).
- 18 United States Environmental Protection Agency (2015a) *2011 National Emissions Inventory, version 2 Technical*
- 19 *Support Document*, U.S. Environmental Protection Agency, Office of Air Quality Planning and Standards, Air
- 20 Quality Assessment Division, Emissions Inventory and Analysis Group, Research Triangle Park, North Carolina,
- 21 August 2015. [www.epa.gov/air-emissions-inventories/2011-national-emissions-inventory-nei-data](http://www.epa.gov/air-emissions-inventories/2011-national-emissions-inventory-nei-data)
- 22 United States Environmental Protection Agency (2015b), 40 DFR Part 60, EPA-HQ-OAR-2013-0602; FRL-XXXX-
- 23 XX-OAR, RIN 2060-AR33, Carbon Pollution Emission Guidelines for Existing Stationary Sources: Electric Utility
- 24 Generating Units, August 3, 2015.
- 25 United States Environmental Protection Agency (2015c) *Technical Support Document (TSD) Preparation of*
- 26 *Emissions Inventories for the Version 6.2, 2011 Emissions Modeling Platform*, U.S. Environmental Protection
- 27 Agency, Office of Air Quality Planning and Standards, Air Quality Assessment Division, contacts: Alison Eyth, Jeff
- 28 Vukovich, August 2015. Retrieved from [https://www.epa.gov/air-emissions-modeling/2011-version-62-technical-](https://www.epa.gov/air-emissions-modeling/2011-version-62-technical-support-document)
- 29 [support-document](https://www.epa.gov/air-emissions-modeling/2011-version-62-technical-support-document) (July 27, 2018).
- 30 United States Environmental Protection Agency (2017) *Inventory of U.S. Greenhouse Gas Emissions and Sinks*
- 31 *1990-2015*, EPA 430-P-17-001.
- 32 United States Environmental Protection Agency (2018) *Inventory of U.S. Greenhouse Gas Emissions and Sinks*
- 33 *1990-2016*, EPA 430-R-18-003.
- 34 United States Geological Survey (2013) *2011 Minerals Yearbook: Cement*, U.S. Department of the Interior, U.S.
- 35 Geological Survey. December 2013.
- 36 USGCRP, 2018: Second State of the Carbon Cycle Report (SOCCR2): A Sustained Assessment Report [Cavallaro,
- 37 N., G. Shrestha, R. Birdsey, M. A. Mayes, R. G. Najjar, S. C. Reed, P. Romero-Lankao, and Z. Zhu (eds.)]. U.S.
- 38 Global Change Research Program, Washington, DC, USA, 878 pp., <https://doi.org/10.7930/SOCCR2.2018>.
- 39 VandeWeghe, J. R., & Kennedy, C.: A Spatial Analysis of Residential Greenhouse Gas Emissions in the Toronto
- 40 Census Metropolitan Area. *Journal of Industrial Ecology*, 11(2), 133–144. <https://doi.org/10.1162/jie.2007.1220>,
- 41 2007.
- 42 Wang, R., Tao, S., Ciais, P., Shen, H. Z., Huang, Y., Chen, H., ... Piao, S. L.: High-resolution mapping of
- 43 combustion processes and implications for CO<sub>2</sub> emissions. *Atmospheric Chemistry and Physics*, 13(10), 5189–5203.
- 44 <https://doi.org/10.5194/acp-13-5189-2013>, 2013.
- 45 Wang, J., Cai, B., Zhang, L., Cao, D., Liu, L., Zhou, Y., ... Xue, W.: High resolution carbon dioxide emission
- 46 gridded data for China derived from point sources. *Environmental Science and Technology*, 48(12), 7085–7093.
- 47 <https://doi.org/10.1021/es405369r>, 2014.





- 1 Wilson, J., Spinney, J., Millward, H., Scott, D., Hayden, A., & Tyedmers, P.: Blame the exurbs, not the suburbs:  
2 Exploring the distribution of greenhouse gas emissions within a city region. *Energy Policy*, 62, 1329–1335.  
3 <https://doi.org/10.1016/j.enpol.2013.07.012>, 2013.
- 4 World Business Council for Sustainable Development: *CO<sub>2</sub> accounting and reporting standard for the cement*  
5 *industry*, version 2.0, June 2005.
- 6 Yadav, V., Michalak, A. M., Ray, J., & Shiga, Y. P.: A statistical approach for isolating fossil fuel emissions in  
7 atmospheric inverse problems. *Journal of Geophysical Research*, 121(20), 12,490–12,504.  
8 <https://doi.org/10.1002/2016JD025642>, 2016.
- 9 Zhang, Y., Wang, H., Liang, S., Xu, M., Liu, W., Li, S., ... Bi, J.: Temporal and spatial variations in consumption-  
10 based carbon dioxide emissions in China. *Renewable and Sustainable Energy Reviews*, 40, 60–68.  
11 <https://doi.org/10.1016/j.rser.2014.07.178>, 2014.
- 12 Zheng, B., Huo, H., Zhang, Q., Yao, Z. L., Wang, X. T., Yang, X. F., ... He, K. B.: High-resolution mapping of  
13 vehicle emissions in China in 2008. *Atmospheric Chemistry and Physics*, 14(18), 9787–9805.  
14 <https://doi.org/10.5194/acp-14-9787-2014>, 2014.
- 15 Zhou, Y., and Gurney, K.R.: Spatial relationships of sector-specific fossil fuel CO<sub>2</sub> emissions in the United States,  
16 *Glob. Biogeochem. Cycles*, 25, GB3002, doi:10.1029/2010GB003822, 2011.