

PROTEVS-MED field experiments: Very High-Resolution Hydrographic Surveys in the Western Mediterranean Sea

5

Pierre Garreau¹, Franck Dumas², Stéphanie Louazel², Stéphanie Correard², Solenn Fercoq², Marc Le Menn², Alain Serpette², Valérie Garnier¹, Alexandre Stegner³, Briac Le Vu³, Andrea Doglioli⁴, Gerald Gregori⁴

10

¹ IFREMER, Univ. Brest, CNRS UMR 6523, IRD, Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM, 29280, Plouzané, France

² SHOM, Service Hydrographique et Océanographique de la Marine, 13 rue de Chatellier, CS592803, 29228 Brest CEDEX 2, France

15 ³ Laboratoire de Météorologie Dynamique (LMD), CNRS UMR 8539, Ecole Polytechnique, 91128 Palaiseau, France

⁴ Aix Marseille Univ., Université de Toulon, CNRS, IRD, MIO UM 110, 13288 Marseille, France

Correspondence to: Pierre Garreau (pierre.garreau@ifremer.fr)

20

Abstract: From 2015 to 2018 four field experiments (7 legs) have been performed in the Western Mediterranean Basin during winter or early spring. The main objectives were the assessment of high-resolution modelling, the observation of mesoscale structure and associated ageostrophic dynamics. Thanks to the intensive use of a towed vehicle undulating in the upper oceanic layer between 0 and 400 meter depth (a SeaSoar), a large amount of very high resolution hydrographic transects (total length about 10.000 km) have been performed, observing mesoscale dynamics (slope current and its instabilities, anticyclonic eddies, submesoscale coherent vortices, frontal dynamics, convection events, strait outflows) and submesoscale processes like stirring, mixed layer or symmetric instabilities. When available, the data were completed with velocities recorded by Vessel Mounted Acoustic Doppler Current Profiler (VMADCP) and by surface salinity and temperature recorded by ThermosalinoGraph (TSG). Classical full depth CTD (Conductivity, Temperature, Depth) stations have also been performed giving the background hydrography of the deeper layers when focusing on peculiar structures. In 2017, a free fall profiler (an MVP-200) has been deployed to manage even higher horizontal resolution. In 2018, another free fall profiler (a rapidCast) has been tested. When available, biological sensors (chlorophyll-a, turbidity, dissolved oxygen etc.) have been carried out. They provided useful complementary observations about the circulation. This data set is an unprecedented opportunity to investigate the very fine-scale processes as the Mediterranean Sea is known for its intense and

contrasted dynamics. It should be useful for modellers (who reduce the grid size below a few hundred meters) and expect to properly resolve finer scale dynamics. Likewise, theoretical work could also be
40 illustrated by in situ evidence embedded in this data set. The data are available through SEANOE repository (<https://doi.org/10.17882/62352>; Dumas et al., 2018).

1 Introduction

45

Progress in numerical modelling and conceptual approaches both emphasized the importance of fine scale processes in connecting the ocean interior to the atmosphere, driving the energy cascade to small scales and reversely (McWilliams, 2016) and shaping the biochemical cycles and biodiversity distribution (Lévy et al., 2012; Lévy et al., 2018). For instance, in the Northwestern Mediterranean Sea, chorophyll-a filaments near the external boundary of the North Current in the Ligurian Sea are
50 generated by frontal instabilities (Niewiadomska et al., 2008) and coherent vortices may act efficiently both as biological barriers and drivers of plankton diversity (Bosse et al., 2017; Rousselet et al., 2019). As all these scales interactions are ubiquitous, it is of crucial importance to develop an observation strategy to get insight simultaneously of the large-scale dynamics, the mesoscale and the submesoscale
55 processes. Unfortunately, it is not straightforward to reach this objective using conventional cruise strategies leading to a lack of in situ observations of fine scale processes. Due to their synoptic view, satellite observations partly fill the gap between large and finer scale dynamics. Since many years, remotely sensed observations of surface temperature, ocean colours or altimetry exhibit a large spectrum of processes with various cut off scale (from around 70 km for the altimetry down to some
60 tens of meters for imagery). Some of these limits will be pushed back soon by the future Surface Water and Ocean Topography (SWOT) satellite, expecting to provide substantial improvement for small scales processes having a sea surface height signature (d'Ovidio et al., 2019).

The data presented hereafter are a contribution to very high-resolution observations of the top oceanic
65 layer and are freely available on SEANOE repository: <https://doi.org/10.17882/62352> (Dumas et al., 2018). Long transects of the first 400 m below the surface was sampled with a horizontal resolution in the range of two kilometres in the Western Mediterranean Sea. The hydrographic and dynamics background of this region is given in section 2. The objectives and implementation of the surveys are presented in section 3. The details of the measurements (platforms, sensors, methodology, metrology,
70 data control, ancillary data) are reported in section 4. To illustrate the potential of the dataset, an overview of the observed processes are displayed in section 5. Lessons learned during surveys and summary are displayed in section 6.

2 Oceanic contexts of the Western Mediterranean Sea

75

The Mediterranean Sea is often referred as a “laboratory basin” exhibiting many processes that are met pervasively and are of primary interest in the functioning of the global ocean (Robinson and Golnaraghi, 1994). It thus provides the opportunity to investigate a large panel of oceanic features in a relatively restrained and accessible area. Therefore, the PROTEVS-MED cruises potentially caught a multitude of physical processes in the Western Mediterranean Sea (figure 1).

80

2.1 Surface and intermediate circulation

The basin or sub-basin dynamics is largely driven by the thermohaline circulation. The Mediterranean Sea is a semi enclosed evaporation basin including areas of intermediate to deep convection. Light (fresh) Atlantic Water (AW), inflowing through the Gibraltar Strait generally circulates along the continental slope in both western and eastern basins (Millot and Taupier Letage, 2005). The slope current is unstable along the Algerian Coast and generates anticyclonic eddies called Algerian Eddies (AE) that spread AW in the southern half of the western basin called Algerian Basin (Escudier et al., 2016a; Puillat et al.; 2002). In the northern part of the western basin, the AW composed East Corsica Current (ECC) and West Corsica Current (WCC) join in the Ligurian sea to form the Northern Current (NC) that flows along the slope until the Balearic Sea (Millot et al., 1999; Send et al., 1999). The existence and the strength of a return branch of this current along the North Balearic Front (NBF) between Menorca Island and Corsica Island is still under debate despite the generally accepted concept of a Northern (cyclonic) Gyre, in agreement with the doming of isopycnals in the central part of this sub-basin. The Levantine Intermediate Water (LIW) formed in winter in the eastern basin, entering into the western basin through the Sicilian Strait, follows more or less the same cyclonic circulation pattern (Millot and Taupier Letage, 2005). It spreads out into the northern part of the western basin between 400 and 800 m depths and is found sporadically within the Algerian Basin. In the western basin, this important water mass, marked by a relative subsurface maximum of temperature and salinity, was already quoted as water type in Sverdrup et al. (1942).

100




Figure 1: Map of PROTEVS-MED cruises. Dynamical features, North Balearic Front (NBF), East Corsica Current (ECC), West Corsica Current (WCC) and Northern current (NC) are depicted in grey.

2.2 Mesoscale structures

105 As in the global ocean the mesoscale dynamics is ubiquitous within the Mediterranean basin; it plays a major role in redistributing water masses and has been evidenced by remote sensing for a long time (Millot et al., 1990). In the Western Basin, the first internal radius of deformation spans in the range of 6 km in the Northern Gyre and of 16 km in the Algerian Basin (Escudier et al., 2016b). It is an indicator of the typical size of the mesoscale activity because the scale of surface intensified eddy in geostrophic balance range a few deformation radii. As a result of ocean-atmosphere exchanges, of large structures instabilities or of flow-topographies interactions, Submesoscale Coherent Vortices, hereafter named SCV (Mc Williams, 1985), whose sizes are currently close to the local radius of deformation, have been observed in the Western Mediterranean (Testor and Gascard 2003; Bosse et al., 2015; 2016). Eddies, meanders, filaments and fronts are typically smaller than in the world ocean.

115

2.3 Sub mesoscale structures.

120 There are strong interactions between mesoscale structures thus generating intense stirring, layered structures and patchy ocean areas. Air-Sea exchanges are marked by frequent strong events (Tramontane and Mistral gusts for instance in the northwestern basin); they interact with the NC and mesoscale structures and generate sinks or sources of potential vorticity, thus leading to ageostrophic dynamics (Bosse 2015; Estournel et al., 2016; Giordani et al., 2017; Testor et al., 2018).

125 Besides, the north western mediterranean basin is known to be the place of deep convection events which has been studied for a long time and even taken as one of the paradigms of deep oceanic convection (Medoc Group et al., 1970; Schott et al., 1996; Houpert et al., 2016; Testor et al., 2018) Marshall and Schott, 1999). Both modelling (Jones and Marshall, 1993; 1997) and observations (Bosse et al., 2016; Margirier et al., 2017) show that deep convection is highly favourable to the production of fine scale structures at submesoscale whether they are due to deepening of the mixed layer during winter or to postconvection restratification.

130 2.4 Previous high-resolution observations

The finest part of the mesoscale dynamics often escapes the usual sampling strategy (CTD arrays, glider deployments) because of being short lived, small in size and quickly advected. The development in the last decade of gliders fleet revealed nevertheless the mesoscale variability in the Western Mediterranean Basin.

135 Recent field experiments based on the multi-platform integrated monitoring program MOOSE (Coppola et al., 2019) or on intensive targeted experiment HYMEX (Estournel et al., 2016) have revisited the hydrography and the dynamics of the northwestern part of the western basin. A strategy of regular and repeated gliders lines as well as dedicated deployments allowed to characterize the variability of the dynamics and to describe crossed fine scale structures. Bosse et al. (2015; 2016) inventoried the
140 Submesoscale Coherent Vortices (SCV) and their contributions to water mass redistribution. With data issued from the same strategy, Margirier et al. (2017) characterized the convection plumes in the Gulf of Lions. Testor et al. (2018) summarized the observations of convection during the dedicated experiment HYMEX. Multi-platform strategies including gliders, mooring, combined cruises (Ruiz et al., 2009; Pascual et al., 2017; Petrenko et al., 2017; Knoll et al., 2017; Onken et al., 2018; Troupin et al., 2019), or colocation with altimetric tracks (Borrione et al., 2016; Heslop et al., 2017; Aulicino et al.,
145 2018; Aulicino et al., 2019; Carret et al., 2019) can provide part of the missing synoptic view.

The capability of changing the glider's trajectory at any time has not often been used in a small-scale context because its horizontal velocity remains low (in the range of 15-30 km/day), preventing any
150 rapid assessment of a detected small structure. Despite this lack of synopticity, Cotroneo et al. (2015; 2019) adapted a glider trajectory to a remote sensed observed Algerian Eddy and Current. Conversely the SeaSoar horizontal velocity is 10 times faster than the glider one. This towed vehicle can handle a turning radius of 2 nautical miles (i.e. gyration speed of 10 degree/min) when the ship changes direction. It allows a strategy based on long exploratory transects as the ship velocity is close to its

155 transit velocity and of intensive sampling on particular detected structures. Due to heavy logistics
involvement, the use of SeaSoar remained scarce in the Western Mediterranean Sea. Allen et al. (2001;
2008) observed an oblate lens of 20 km radius, 150 m thick, centred at 250 m depth during the
OMEGA-2 field experiment in fall 1996. Salat et al. (2013) reported Seasoar transects in the Gulf of
160 Lions after the convection in spring 2009. The Seasoar was also used during one leg of the ELISA field
experiment devoted to the Algerian Eddies (Taupier-Letage et al., 2003) but only the mesoscale features
have been reported.

A free fall recovered platform, the Moving Vessel Profiler (MVP-200) has a lighter logistics but
requires a lower vessel velocity to reach depths equivalent to those reached with the SeaSoar: that is to
say 2-4 knots to go down 400 m deep. In the Western Mediterranean Sea, the MVP was deployed
165 during OSCAHR cruise allowing a detailed study of a cyclonic structure in the Ligurian Sea (Rousselet
et al., 2019) and in situ estimation of the sea surface height for a comparison with along track satellite
data (Meloni et al., 2019).

3 Objectives and achievement of the field experiments.

170

The main scientific objectives of the cruises were threefold:

- to assess the large-scale circulation features of the Western Mediterranean Basin, evaluating the water
masses and the fluxes at different key points in the basin (the Ligurian Sea, the Balearic Sea, the
Northern Tyrrhenian Sea, the North Balearic front area and the Algerian Basin). The goal was the
175 assessment of operational numerical simulation of the circulation performed for the Navy.

- to identify and follow peculiar mesoscale structures such as surface eddies, modal structures composed
of Winter Intermediate Water (WIW), submesoscale coherent vortices (SCV), meanders and filaments
and explore their signatures on the sea surface height (altimetry) and their acoustic impact (i.e. through
their modulation of the sound propagation speed).

180 - to observe and characterize the submesoscale dynamics such as ageostrophic stirring, symmetric
instabilities, mixed layer instabilities, subduction and convection.

Clearly, the main part of the present dataset is not devoted to track any climatic change in water mass
properties; the SeaSoar, the MVP or the rapidCast are rapid moving platforms leading to acquire less
185 precise temperature, conductivity and above all deduced salinity data than standardized CTD's protocol.
MVP and rapidCAST are equipped with unpumped sensors and the three tools reach high ascending or
descending velocity (above 2m/s) that leads to inescapable thermal lag issues across sharp fronts.
Readers interested in this topic should only use the classical CTD stations data.

190 Four cruises were conducted between 2015 to 2018 by the "Service Hydrographique et
Océanographique de la Marine" (SHOM) in the western mediterranean basin, during winter or early
spring, managing mainly the towed undulating vehicle SeaSoar to investigate the sub-surface (0-400m)
layer. When the deployment of this vehicle was either unsafe (over shallow water) or even impossible

195 (due to rough meteorological conditions, breakdown of winch or vehicle) or when complementary observations were requested (e.g. go below 400m or getting water samples for biochemical analysis), CTD casts were rather performed. Ship board routinely acquired data (Vessel Mounted Acoustic Doppler Current Profiler -VMADCP- and ThermoSalinoGraph -TSG-) were also included in this database.

200 We present here in a synthetic dataset all data recorded during the cruises (figure 2; table 1). Complementary data used to the cruise design, to adapt on field the strategy or to interpret results (altimetric tracks, remote sensed sea surface temperature or chlorophyll-a) are available on CMEMS servers (<http://marine.copernicus.eu>). An eddy detection tools called AMEDA (Le vu et al., 2017) has also been used to detect and track structures during the cruises. Surface and Argo drifters were also

205 dropped during the cruises and data are available on companion datasets (table 2).

PROTEVS-MED	2015_leg1 07-01 / 24-01	2015_leg2 16-04 / 03-05	2016 22-03 / 04-04	2017_leg1 27-01 / 07-02	2017_leg2 11-02 / 23-02	swot_2018_le g1 23-04 / 26-03	swot_2018_le g2 30-04 / 18-05
SEASOAR	2290 km 1137 profiles	329 km 263 profiles	2090 km 1369 profiles	1858 km 706 profiles	620 km 1162 profiles	615 km 411 profiles	2830 km 2381 profiles
MVP				153 km 813 profiles	188 km 708 profiles		
RAPIDCAST						22 km 92 profiles	167 km 71 profiles
CTD	62 profiles	151 profiles	47 profiles	17 profiles	27 profiles	1 profile	12 profiles
LADCP	56 profiles	137 profiles	47 profiles	18 profiles	26 profiles		3 profiles
XBT	30 profiles	22 profiles		39 profiles	1 profile		
VMADCP 38kHz	2676 km					1071 km	4118 km
VMADCP 150kHz	3036 km	3688 km		2705 km	5384 km	1068 km	4128 km
TSG	3756 km	3861 km	3744 km	2999 km	3011 km	1212 km	4984 km

Table 1: Summary of performed transects, profiles and routinely acquired data. Cumulated length of transects and total numbers of vertical profiles are displayed.

210

The first cruise called **PROTEVS-MED 2015_leg1** took place from 7 to 24 January 2015 on board of the R/V *Pourquoi Pas?*. Its main objective was the dynamics of the NC from the Ligurian Sea to the Gulf of Lions and the associated meso- and submeso-scale processes. Attention has been paid to cross-

215 slope transects in the Gulf of Lions, in order to examine the behaviour of the North Current and the exchanges across the shelf break. An intensive survey of the North Current between Toulon and Nice was performed.

220 The second leg, **PROTEVS-MED 2015_leg2**, was carried out on the R/V *Beautemps-Beaupré* from 16 April to 3 May 2015. It started in the Balearic Sea and investigated the slope current from Ligurian Sea to Balearic Sea. During this cruise, the SeaSoar trawl failed early, just after three transect acquisitions in the Balearic Sea describing the hydrology relative to the cyclonic circulation and its associated mesoscale structures. This has therefore led to carry out mostly CTD stations and VMADCP 150 kHz records. In particular, a dense array of CTD casts was then performed within the North Current between Nice and Toulon. The **PROTEVS-MED 2015_leg2** survey was characterized by a proliferation of
225 jellyfish, the CTD measurements are to be taken with caution as they caused an excessive smoothing of temperature and salinity. When too large differences appeared between the values at the ascent and descent, the profiles have been flagged 4 (bad value that can be corrected)

230 The second campaign **PROTEVS-MED 2016** took place on the R/V *Beautemps-Beaupré* from 22 March to 4 April 2016. It was designed to focus on the origin of the NC where the flows through the Corsica Channel and the WCC join. The behaviour and the origin of the WCC was also explored along the western coast of Corsica. Besides unveiling part of the complex hydrological structure of the NBF in early spring, the **PROTEVS-MED 2016** survey allowed to capture an Algerian Eddy in interaction with the NBF. Garreau et al. (2018) described in details its original double core structure: a
235 superposition of two water masses of different origin spinning together. The survey provides also scenes and insights about the way both components of the Northern Current merge together to the north of the Corsica Channel during early spring.

240 The third campaign **PROTEVS-MED 2017** held from 27 January to 7 February (**leg1**) and from 11 February to 23 February (**leg2**) on board the R/V *Atalante*. This survey was devoted to explore eddies detected by altimetry in the North Balearic Front and to assess an eddy tracking tool (Le Vu et al., 2017). Transects across the North Balearic Front revealed the complexity of this transition zone. In order to escape rough sea state consecutive to a strong Mistral gust, part of the cruise was dedicated to the investigation of the Balearic Sea and the outflow of coastal fresher and colder water mass from the
245 Gulf of Lions. Back to the deep-sea area, partial convection and WIW formation were recorded. An SCV was thoroughly observed, north of the Balearic front.

250 The fourth and last field experiment **PROTEVS-MED-SWOT 2018** was conducted in the framework of SWOT Preparatory Phase from 23 April to 26 April (**leg1**) and from 30 April to 18 May (**leg2**) south from Balearic Islands on board the R/V *Beautemps-Beaupré*. The first leg (**leg1**) performed a general overview of the oceanic situation followed by a more intensive survey (**leg2**) planned on the basis of daily release of near real-time satellite imagery, altimetry, and lagrangian analyses, performed on land by use of a dedicated package (<http://www.mio.univ-amu.fr/SPASSO/>, as in Nencioli et al., 2011; de Verneil et al., 2017). Satellite data of altimetry, sea surface temperature and ocean colour revealed
255 ubiquity throughout the cruise period of very fine oceanic structures such as dipolar structures or

260 tenuous fronts. A special focus was stressed on the south of Mallorca where fronts were detected by altimetry-derived currents and diagnosis (e.g. Finite Singular Lyapunov Exponents; d'Ovidio et al., 2004), by contrasted surface chlorophyll-a concentrations and confirmed by high frequency flow cytometry analyses of phytoplankton performed onboard (data not included in the present dataset). A Lagrangian strategy was specifically set up in order to study the structure and growth rate (at 24 hours time scale) of the various phytoplankton groups as defined by flow cytometry measurements as in Marrec et al. (2018). Last, it is noticeable that a companion campaign (PRE-SWOT) managed by IMEDEA-SOCIB held in the same area and during the same period on board of R/V *Garcia Del Cid* (not included in the present dataset, see Barceló-Lull et al., 2018).

265




Figure 2 : Map of PROTEVS-MED cruises and instruments deployed; TSG is available for all the ship track

4 Data, Methods and Quality Controls

270 **4.1 CTD casts and LADCP**

The CTD casts were performed with the Sea-Bird SBE-9 instrument mounted in a General Oceanics 12-places rosette frame fitted with 12 Niskin bottles. Sometimes a RDI 150 kHz current profiler was also implemented on the rosette and then LADCP (Lower Acoustic Doppler Current Profiler) performed measurements during the cast. Standard hydrographic procedures for CTD casts were applied. When
275 available, LADCP recorded data were processed following the inversion method of Visbeck (2002).

4.2 Seasoar deployments

The SeaSoar is a towed undulating vehicle designed and built by Chelsea Instruments. Two Sea-Bird SBE-9 (with SBE-3 temperature and SBE-4 conductivity sensors) instruments were mounted on either
280 side of the SeaSoar. When available, a WET Labs WetStar chlorophyll-a fluorometer, both oxygen sensor (SBE-43) and optical properties sensor (WET Labs C-Star) were deployed. The SeaSoar was trawled at 9 knots by a profiled cable. It was undulating between the surface and 400m below the surface under optimal conditions with a horizontal resolution in the range of 2 km. Rough sea states, lateral currents, strong vertical shears can degrade the performance of the vehicle and reduce the vertical
285 range of exploration between 20 and 360m. As the software allows real time visualisation of the ongoing transect, it is a perfect tool to scan the upper oceanic layer where meso- and submesoscale dynamics are the most intense. A total of 10.000 kilometres of transects crossing numerous and various structures has been yet recorded during the cruises, giving the unique opportunity to explore fine scales patterns of the upper layer of the western basin.

290

4.3 MVP deployments

During PROTEVS-MED 2017 surveys a Moving Vessel Profiler (MVP200) - a computer controlled winching system that can deploy and recover a sensor from a ship that is underway - was deployed for finer transects. The sensor was an AML CTD embedded in a free fall fish. At 2-4 knots, it was possible
295 to monitor the 0-400 m layer with a horizontal resolution less than 1 km. To remove spurious salinity values due to bubbles when the instrument is surfacing, the minimum pressure for valid record was set to 1 decibar. A peculiar transect has been monitored using successively SEASOAR, MVP and CTD casts given the opportunity to compare the three techniques.

300 **4.4 RapidCast deployments**

During PROTEVS-MED 2018 a free fall CTD system, called rapidCast (Teledyne Marine ; <http://www.teledynemarine.com/rapidcast>) was tested for three transects near Balearic Islands. It was equipped with the “rapidCTD - Underway Profiler” proposed by Valeport. A bluetooth communication allowed real-time evaluation of each profile when the probe is surfacing near the ship deck. This system

305 sampled the water layer from 0 to 400m with a navigation speed in the range of 5-6 knots and a resolution similar to the SeaSoar (about 2 km).

4.5 TSG

310 During the cruises, a Seabird SBE-21 ThermoSalinoGraph recorded the sea surface temperature and conductivity. The inlet was equipped with a SBE 38 thermometer. The recorded sea surface temperature and salinity contributed to the Global Ocean Surface Underway Data (GOSUD) program (Gosud, 2016). The metrological traceability and the data treatment are insured according to the procedures described in Gaillard et al. (2015) which explains the delayed mode processing of datasets and presents an overview of the resulting quality. The calibrations are complemented with rigorous adjustments on
315 water samples leading to reach a salinity accuracy of about 0.01 or less.

4.6 VMADCP

The hardwares used, their configurations and the way they are carried out are similar on R/V Pourquoi Pas ?, R/V L'Atalante and R/V Beutemps Beaupré. The VMADCPs are 150 kHz and 38 kHz Ocean
320 Surveyor by RDI Teledyne. They are both monobloc antennas using beam forming process to form four beams oriented towards 30° from the vertical. Nominally, they emitted a ping per second from which ensemble are built to get less noisy profiles. Two ensembles are routinely processed:

- a Short-Term Average (hereafter noted STA) which gathers and averages the pings of two-minute window. It makes ensemble of 120 pings at least,
- 325 - a Long-Term Average (noted LTA) made of 600 pings or averaged over ten minutes.

The series of geometric transformations necessary to pass from beam coordinate along beam data to absolute geographic coordinate and geophysical velocity are performed thanks to VMDAS software from RDI Teledyne. It combines the position (latitude and longitude) from the DGPS Aquarius and Octans central with the PHINS inertial navigation system from IXSEA (that provides vessels attitude
330 data: pitch, roll, heaving) to provide synchronised single ping earth coordinates data (file .ENX) and short- and long-term ensemble (STA / LTA).

This native format (.STA/.LTA) were also processed with WinAdcp in order to extract and provide only significant (i.e. with a satisfactory signal/noise ratio) data that are additionally formatted to text file.

335 Note that data processed by CASCADE software can be requested on Sismar repository, collecting and processing progressively all VMADCP from french research vessels (<https://sextant.ifremer.fr/record/60ad1de2-c3e1-4d33-9468-c7f28d200305/en/index.htm>).

4.7 Data metrological traceability and calibration

340 SBE 9 temperature and conductivity sensors deployed on all CTD were calibrated before and after each campaign or at least once a year in the SHOM's thermo-regulated baths, whose temperature can be

stabilized to less than 1 mK (peak to peak) during control and calibration operation. Such a procedure allows the monitoring of sensors drifts between calibrations and the detection of anomalies. In the cases sensors have kept a good linearity, which is the most common, data are corrected with offset-slope coefficients. Figure 3a show the review of corrections applied on data at 15 °C, after the calibrations of SBE 3 sensors used for PROTEVS-MED campaigns. Figure 3b shows the review of corrections applied at 40 mS cm⁻¹ after the calibrations of SBE 4 sensors.

The temperature of the thermo-regulated bath is monitored with SBE 35 which is used as laboratory reference temperature sensors. They are linked to the International Temperature Scale of 1990 (ITS-90) thanks to calibrations performed once a year in a triple point of water cell and in a melting point of Gallium. These reference cells are regularly calibrated by the French National Metrology Institute (NMI) LNE-CNAM. The calibration expanded uncertainty of SBE 3 sensors is between 1.8 and 2.3 mK according to the residual linearity errors of SBE 3's.

Conductivity calibration of SBE 4 sensors is made in the same bath during the temperature calibration. Seawater samples are taken in the bath and tested against Autosal and Portasal salinometers. The calibration procedure and the propagation of uncertainties to the calculated salinities from SBE 9 data are described in Le Menn (2011). Practical salinity expanded uncertainty varies from 0.0032 to 0.0034. In 2015, the SHOM laboratory took part in the JCOMM intercomparison for seawater salinity measurements (JCOMM, 2015) showing that Autosal and Portasal measurements are within ± 0.001 compared to other participating laboratories. Note that the same process was done in the framework of an international network for the TSG data of the French Research Vessel (see §4.5 and table 2). Unfortunately, the MVP and the rapidCast sensors were not available for such a common process and were calibrated directly by the constructor, but a comparison with the in situ records with calibrated SBE sensors can be carried out. As the optical properties and oxygen concentration were used as tracer only, no calibration process was performed.




Figure 3 (a) review of corrections applied on data at 15 °C, after the calibrations of SBE 3 sensors. (b) review of corrections applied on data at 40 mS cm⁻¹, after the calibrations of SBE 4 sensors.

4.8 Data processing levels

Three levels of processing are available for each dataset:

- 375 - Level 0 (L0) consists in the direct output of sensors at full temporal resolution;
- Level 1 (L1) displaying data in ascii (.csv) or netcdf (.nc) files are only processed from the software of the constructor, keeping the full resolution and computing the derived variables into standard units. Recent instrumental system (AML and Valport probes) directly provides level 1 files. L1 files are corrected from eventual drift of sensors;
- 380 - Level 2 (L2) is proposed as gridded, controlled and resampled data in netcdf files (.nc). Gridded dataset for salinity and temperature have been resampled vertically every meter removing spikes, spurious values, density inversions when they persist after the first process supplied by the sensor manufacturer. They are then positioned as vertical profiles at the mean geographical position of the considered up or down record.

385

Temperature and salinity data were also compared to the historical data in the neighbourhood of the profiles or transects using the validated CORA database distributed by the Copernicus Marine and Environment Service (Cabanne et al., 2013; Szekely et al., 2017). PROTEVS-MED data are not yet included in this database but will be transmitted for a future release.

390 All gridded profiles or transects have been plotted for a visual quality check and are available as “quick looks” on the repository. Level 1 (L1) and (or) level 2 (L2) dataset are released in the present database. Level 0 (L0) remain available in constructor format upon request to the data providing institution (SHOM; data-support@shom.fr).

395 4.9 Companion datasets.

During the field experiments, surface drifters with holey socks located at 100m, 75m, 50m or 15 m depth were deployed given the opportunity at the beginning of their track to perform a Lagrangian survey of observed structures.

400 Few Argo floats were also dropped and experienced first PROTEVS-MED dedicated mission with high temporal resolution (daily cycle) and parking depths adjusted to the observations to maintain the drifters as long as possible within the targeted structures (typically 100 m deep). After the drifter left the structure, it used the usual Argo standard procedure in the Mediterranean (i.e. a 5-day cycle and a parking depth of 350 m). Already stored in dedicated and accessible stable repositories, they can be
405 found using their WMO identifiers (World Meteorological Organisation) (see table 2). Ancillary data can be found on different repositories selecting date and locations corresponding to PROTEVS-MED surveys.

	ARGO float WMO identifier http://www.ifremer.fr/argoMonitoring/	SVP surface drifter WMO identification number (holey sock depth) http://www.jcommops.org/dbcp/	TSG ThermoSalinoGraph (platform identifier) http://www.gosud.org
PROTEVS-MED 2015_leg1	6901707 6901708	6100536 (50m) 6100537 (50m) 6100538 (50m) 6100539 (50m) 6100540 (50m)	FMCY
PROTEVS-MED 2015_leg2		6100863 (50m) 6100864 (50m) 6100865 (50m) 6100866 (50m) 6100867 (50m)	FABB
PROTEVS-MED 2016		6101525 (75m) 6101526 (75m) 6101527 (75m) 6101528 (75m) 6101529 (50m) 6101530 (50m) 6101531 (50m) 6101532 (50m) 6101533 (50m) 6101534 (50m) 6101535 (50m) 6101536 (50m) 6101537 (50m) 6101538 (50m) 6101539 (50m) 6101540 (50m) 6101541 (50m) 6101542 (50m) 6101631 (50m)	FABB
PROTEVS-MED 2017_leg1	6902764 6902765 6902767	6101634 (50m) 6101639 (50m) 6101635 (50m) 6101637 (50m) 6101636 (50m) 6101643 (50m) 6101641 (50m) 6101647 (50m)	FNCM
PROTEVS-MED 2017_leg2		6101648 (50m) 6101633 (50m) 6101638 (50m) 6101640 (50m) 6101632 (50m) 6101642 (50m)	FNCM
PROTEVS-MED-SWOT 2018_leg1	6902844	6101669 (50m) 6102612 (50m) 6101677 (100m) 6102613 (100m)	FABB

PROTEVS-MED-SWOT 2018_leg2		6101671 15m 6101678 15m 6101672 15m 6102615 15m 6101670 50m 6101674 50m	FABB
-------------------------------	--	--	------

410 table 2 : WMO index of Argo-float and surface drifters (SVP) dropped during PROTEVS-MED surveys. Surface temperature and
 415 salinity recorded by ThermoSalinoGraph (TSG) are tagged by ship identifier. All data are available from the CORIOLIS website
<http://www.coriolis.eu.org/Data-Products/Data-Delivery/Data-selection> by entering the WMO numbers in the field 'Platform
 codes', adjusting the time period of interest (e.g., 01/01/2018 to 30/06/2019), and clicking on 'refresh'. The web interface displays
 the trajectories of the buoys, profilers or TSG and can be used to find additionally opportunity data. The data can then be
 downloaded in NetCDF format.

5 Overview of the selected observations

420 When deployed together, VMADCP and SeaSoar provided a unique synoptic view along transects. It
 was also possible to observe simultaneously the density and the velocity fields in the sub-surface layer,
 showing the importance of the geostrophy even at fine scale (in the range of 10 km). The temperature
 and salinity fields were patchier than expected but the thermal expansion and the saline contraction
 coefficient of sea water often compensate and lead to a smoother density (and thus dynamical) field.
 425 The structure of observed anticyclonic eddies appeared also more complex than formerly described and
 were commonly composed of many different water masses. Eddies with similar altimetric or surface
 thermal signatures can be very different hydrographically. For instance, a dual-core anticyclonic eddy
 observed east of Menorca Island in March 2016 (figure 4a), was composed of a superposition of Winter
 Intermediate Water and Atlantic Water and in May 2018 a three-cores eddy has been detected (figure
 4b). The exact process of their formation remains to be investigated. One can invoke the coalescence of
 430 pre-existing eddies, the extraction of water masses from neighbouring structures or ageostrophic
 processes. Measurements reveal submesoscale (ageostrophic) dynamics both in the eddy cores
 (upwelling/downwelling) and at eddy edges (symmetric instabilities, frontogenesis). Intra-pycnocline
 structures, subducted, stirred or locally formed were commonly observed at the edge of gyres.

435 In the Northern Current, stirring appeared in both tracers and velocity fields and an SCV formed by
 LIW detached in front of Toulon (figure 4c) was observed as confirmed by observed swirling velocities
 on VAMDCP records. It was also topped by a surface cyclonic gyre. Currents were routinely recorded
 and a particular attention was paid to the Northern Current dynamics as shown figure 4d.


440 The fine structure of the NBF showed the interaction between the front and the SCV's generated in the
 deep convection area. In the NBF, a shift between a surface layer front and a deep front is revealed. As
 long as most of the experiments presented here were performed during late winter or early spring in the
 vicinity of deep convection area, they frequently showed small scale structures that are likely formed by
 convection. For instance, from the north to the south on figure 4e one can observe successively the -
 probably partial - convection area, the formation of SCVs composed of WIW in the mesoscale

adjustment area around the convection chimney, the surface thermal front and finally a deeper front at
445 40.6° N.

Over the abyssal plain near the bottom, classical CTD station highlighted different Western Mediterranean Deep Waters (WMDW); besides, under the LIW layer, where the profiles foster double diffusion process, staircases in temperature and salinity were commonly observed as in Onken and Brambilla (2003) (figure 4f).

450 MVP transects performed in February 2017, in front of Cap Creus, across the Blanes Canyon, and off Barcelona showed cold water diving along the Catalan shelf and slope (figure 4g). The WIW observed along the Catalan slope was relatively fresh and cold water originating from the Gulf of Lion's shelf. The WIW was progressively entrained and mixed with the AW and LIW while flowing southwards, in a less intense way than dense water cascading described by Durrieu de Madron et al. (2013).

455 Similar and extensive quick looks of all Seasoar, MVP, or rapidCast transects and XBT, CTD profiles are plotted and available as additional resources on the data repository.



460 **Figure 4 : Overview of some transects or profiles recorded during the PROTEVS-MED fields experiments: (a) Dual core eddy in NBF in 2016 (39.96N:4.74E- 39.93N:6.20E), (b) Three layered eddy in Algerian Basin (38.5N:4.87E – 38.75N:5.88E ; 2018), (c) SCV of LIW front of Toulon (42.17N:6.14E – 42.78N:6.10E ; 2015), (d) North Current position and intensity (2015), (e) cross frontal (NBF) transect (40.35N:6.43E - 41.92:4.01E ; 2017), (f) staircase in temperature front of Sardinia (2016), vein of cold water from the Gulf of Lion on Catalan sea (40.68N:7.5E ; 2017). The reader will find similar quick looks of the transects for all the surveys into the data repository.**

465 **6 Conclusions**

The PROTEVS-MED dataset available through an unrestricted unique repository is an unprecedented opportunity for the community to approach the fine-scale dynamics in the Western Mediterranean Sea and more largely the submesoscale dynamics associated with strong mesoscale dynamics. In the framework of the high-resolution altimetry this dataset can help to characterize the scales of fine structures in the Western Mediterranean Sea and to design combined experiments using high resolution in-situ measurements (Seasoar or MVP) and altimetry with the future SWOT satellite (d'Ovidio et al., 2019). It also complements the repeated glider lines maintained in the framework of the MOOSE observatory (Coppola et al, 2019) and is useful to design future combined multi-platform experiments.

During these campaigns, one had the opportunity to deploy different instruments to obtain temperature, salinity and possibly other parameters profiles. Some transects have been performed successively using CTD and Seasoar (all surveys) or using CTD, Seasoar and MVP-200 (PROTEVS-MED 2017). Easy to manage, the rapidCast has only been tested and used as instant spare in 2018 when the Seasoar failed. It produced a similar result to the Seasoar in temperature and salinity. The SeaSoar is heavy to manage, needs a consequent research vessel for the winch system, a constant watch on its navigation and calm sea state for launch and recovery. Once deployed, the machine can stay at sea for days. Thanks to the required ship velocity (about 9 knots), the SeaSoar remains a perfect platform to identify mesoscale structures before examining them in detail. It explores the oceanic surface layer down to 400 m deep which is sometimes a little bit too short in the mediterranean context, missing deeper part of AEs or deep SCVs but sufficient to describe important surface and subsurface dynamical features. For the same depth range an MVP-200 requires a ship velocity about 2-4 knots and is then more devoted to short transect with higher horizontal resolution. As it is a free-fall platform, its setting is lighter, despite regular inspection of the cable and winch every 10 hours. In any case, when exploring a structure in detail, a CTD network remains necessary, at least to have a valid reference level for the thermal wind equation and hydrography.

Despite the suspected lack of accuracy of the sensors due to the velocity of the platforms (Seasoar, MVP, rapidCast), this experiment of fast and high-resolution sampling revealed fine oceanic patterns never described before in the Western Mediterranean. In situ observations of ageostrophic dynamics remain rare and the synergy between these observations and theory, and then between these observations and modelling, should be very fruitful. These data should contribute to the knowledge of small scales and fill some of the gaps in observing system in the Mediterranean Sea (Tintore et al., 2019). As numerical modelling gain in resolution (in the range of few hundred meters), the simulation of submesoscale processes (layering, subduction, stirring, vertical velocities) is therefore expected and this dataset, providing data at similar scales, is an opportunity to validate the secondary simulated circulation.

505 **Authors contributions**

Louazel S., Correard S., Garreau P., and Dumas F. designed and conducted the field experiments as PI. Marc Le Menn managed the calibration and the metrological traceability of SBE sensors. Valerie Garnier has carefully checked the dataset. All co-authors carried them out, participated to the cruise or
510 processed the data. Garreau P. and Dumas F. prepared the manuscript and the data with contributions from all co-authors.

Competing interests.

The authors declare that they have no conflict of interest.

515 **Acknowledgements**

The authors acknowledge that the French Government Defense procurement and technology agency (Délégation Générale de l'Armement) funded extensively through Protevs and Protevs II “programme d'étude amont” all the campaigns reported here. They thank the technical team at the French Naval
520 Hydrologic and Oceanographic Service (SHOM), the crews of the French Navy ship *Beautemps-Beaupré*, of the R/V *Pourquoi Pas ?* and the R/V *L'Atalante* for their contribution to the field experiments.

525

Data Availability and Repository

Data are freely available on SEANOE repository (<https://doi.org/10.17882/62352>; Dumas 2018). Some parts of the data are already under investigations or publications; the authors would appreciate
530 collaboration proposals. For a first overview quick looks of all SeaSoar, MVP, or rapidCast transects and XBT, CTD profiles are available in catalogues on the repository.

SeaSoar, MVP, rapidCast, CTD, LADCP and XBT data are stored in both CSV (ASCII) and Netcdf files for “L1” (directly extracted from the instrument or constructor software), in Netcdf for “L2” (resampled every meter) files;

535 For TSG, the present database provides only L1 files; L2 (validated and resampled data) are available on dedicated repository (see table 2).

For the sake of simplicity, VMADCP files were concatenated over each cruise duration to provide a single file per cruise; for a given cruise, the data are a function of time and depth within the single file dedicated to the cruise

540

Data are displayed by cruises and instruments and the syntax is :

instrument_data-type_cruises_starting-date-of-record_index.file-type , where

instruments = ctd,seasoar,ladcp,xbt,rapidcast,mvp,vmadcp_XXX

545 **data-type** = “L1” or “L2”

cruises = cruise and leg name

date = the date of the first record in the file.

index = sequential index of this kind of profile recorded during the cruise.

file-type = csv(.csv) or netcdf(.nc)

550

Additionally, data extracted from on-board automatic acquisition are provided in Netcdf file for the ship navigation. Future PROTEVS-MED experiments are scheduled and results will be added to the repository.

555 **References**

Allen, J.T., Smeed, D.A., Tintoré, J., Ruiz, S.: Mesoscale subduction at the Almeria–Oran front: Part 1: Ageostrophic flow. *Journal of Marine Systems* 30, 263–285. [https://doi.org/10.1016/S0924-7963\(01\)00062-8](https://doi.org/10.1016/S0924-7963(01)00062-8), 2001.

560

Allen, J. T., Painter, S. C., & Rixen, M.: Eddy transport of Western Mediterranean Intermediate Water to the Alboran Sea. *Journal of Geophysical Research: Oceans*, 113(C4), C04024. <https://doi.org/10.1029/2007JC004649>, 2008.

565 Aulicino, G., Cotroneo, Y., Ruiz, S., Sánchez Román, A., Pascual, A., Fusco, G., Tintoré, J., Budillon, G.: Monitoring the Algerian Basin through glider observations, satellite altimetry and numerical simulations along a SARAL/AltiKa track. *Journal of Marine Systems* 179, 55–71. <https://doi.org/10.1016/j.jmarsys.2017.11.006>, 2018.

570 Aulicino, G.; Cotroneo, Y.; Olmedo, E.; Cesarano, C.; Fusco, G.; Budillon, G.: In Situ and Satellite Sea Surface Salinity in the Algerian Basin Observed through ABACUS Glider Measurements and BEC SMOS Regional Products. *Remote Sens.* 11, 1361. <https://doi.org/10.3390/rs11111361>, 2019.

575 Barceló-Llull, B., Pascual A., Díaz Barroso L., Sánchez-Román A., Casas B., Muñoz C., Torner M., Alou E., Cutolo E., Mourre B., Allen J., Aulicino G., Cabornero A., Calafat N., Capó E., Cotroneo Y., Cyr F., Doglioli A., d'Ovidio F., Dumas F., Fernández J.G., Gómez Navarro L., Gregori G., Hernández-Lasheras J., Mahadevan A., Mason E., Miralles A., Roque D., Rubio M., Ruiz I., Ruiz S., Ser-Giacomi

- 580 E. and Toomey T.: PRE-SWOT Cruise Report. Mesoscale and sub-mesoscale vertical exchanges from
multi-platform experiments and supporting modeling simulations: anticipating SWOT launch
(CTM2016-78607-P). 138 pp. <https://digital.csic.es/handle/10261/172644>, 2019.
- 585 Borrione, I., Falchetti, S., Alvarez, A.: Physical and dynamical characteristics of a 300m-deep
anticyclonic eddy in the Ligurian Sea (Northwest Mediterranean Sea): Evidence from a multi-platform
sampling strategy. *Deep Sea Research Part I: Oceanographic Research Papers* 116, 145–164.
<https://doi.org/10.1016/j.dsr.2016.07.013>, 2016.
- 590 Bosse, A., Testor, P., Mortier, L., Prieur, L., Taillandier, V., d’Ortenzio, F., Coppola, L.: Spreading of
Levantine Intermediate Waters by submesoscale coherent vortices in the northwestern Mediterranean
Sea as observed with gliders. *J. Geophys. Res. Oceans* 120, 1599–1622.
<https://doi.org/10.1002/2014JC010263>, 2015.
- 595 Bosse, A., Testor, P., Houpert, L., Damien, P., Prieur, L., Hayes, D., Taillandier, V., Durrieu de
Madron, X., d’Ortenzio, F., Coppola, L., Karstensen, J., Mortier, L.: Scales and dynamics of
Submesoscale Coherent Vortices formed by deep convection in the northwestern Mediterranean Sea. *J.*
Geophys. Res. Oceans 121, 7716–7742. <https://doi.org/10.1002/2016JC012144>, 2016.
- 600 Bosse, A., Testor, P., Mayot, N., Prieur, L., D’Ortenzio, F., Mortier, L., Goff, H.L., Gourcuff, C.,
Coppola, L., Lavigne, H., Raimbault, P.: A submesoscale coherent vortex in the Ligurian Sea: From
dynamical barriers to biological implications. *Journal of Geophysical Research: Oceans* 122, 6196–
6217. <https://doi.org/10.1002/2016JC012634>, 2017.
- 605 Cabanes, C., Grouazel A., von Schuckmann K., Hamon M., Turpin V., Coatanoan C., Paris F.,
Guinehut S., Boone C., Ferry N., de Boyer Montégut C., Carval T., Reverdin G., Pouliquen S., and Le
Traon P.Y.: The CORA dataset: validation and diagnostics of in-situ ocean temperature and salinity
measurements, *Ocean Sci.*, 9, 1-18, <https://doi.org/10.5194/os-9-1-2013>, 2013.
- 610 Carret, A., Birol, F., Estournel, C., Zakardjian, B., Testor, P.: Synergy between in situ and altimetry
data to observe and study Northern Current variations (NW Mediterranean Sea). *Ocean Science* 15,
269–290. <https://doi.org/10.5194/os-15-269-2019>, 2019.
- 615 Coppola, L., P. Raimbault, L. Mortier, and P. Testor : Monitoring the environment in the northwestern
Mediterranean Sea, *Eos*, 100, <https://doi.org/10.1029/2019EO125951> 2019.
- 620 Cotroneo, Y., Aulicino, G., Ruiz, S., Pascual, A., Budillon, G., Fusco, G., Tintoré, J.: Glider and
satellite high resolution monitoring of a mesoscale eddy in the algerian basin: Effects on the mixed
layer depth and biochemistry. *Journal of Marine Systems*. 62,73-88.
<https://doi.org/10.1016/j.jmarsys.2015.12.004>, 2019.

- 625 Cotroneo, Y., Aulicino, G., Ruiz, S., Sánchez Román, A., Torner Tomàs, M., Pascual, A., Fusco, G., Heslop, E., Tintoré, J., Budillon, G.: Glider data collected during the Algerian Basin Circulation Unmanned Survey. *Earth System Science Data* 11, 147–161. <https://doi.org/10.5194/essd-11-147-2019>, 2019.
- 630 Durrieu de Madron, X., Houpert, L., Puig, P., Sanchez-Vidal, A., Testor, P., Bosse, A., Estournel, C., Somot, S., Bourrin, F., Bouin, M.N., Beauverger, M., Beguery, L., Calafat, A., Canals, M., Cassou, C., Coppola, L., Dausse, D., D’Ortenzio, F., Font, J., Heussner, S., Kunesch, S., Lefevre, D., Le Goff, H., Martín, J., Mortier, L., Palanques, A., Raimbault, P.: Interaction of dense shelf water cascading and open-sea convection in the northwestern Mediterranean during winter 2012. *Geophysical Research Letters* 40, 1379-1385., 2013.
- 635 de Verneil, A. de, Rousselet, L., Doglioli, A.M., Petrenko, A.A., Moutin, T.: The fate of a southwest Pacific bloom: gauging the impact of submesoscale vs. mesoscale circulation on biological gradients in the subtropics. *Biogeosciences* 14, 3471–3486. <https://doi.org/10.5194/bg-14-3471-2017>, 2017.
- 640 d’Ovidio, F., V. Fernández, E. Hernández-García, and C. López. Mixing structures in the Mediterranean Sea from finite-size Lyapunov exponents, *Geophys. Res. Lett.*, **31**, L17203, doi:[10.1029/2004GL020328](https://doi.org/10.1029/2004GL020328), 2004.
- 645 d’Ovidio, F., Pascual, A., Wang, J., Doglioli, A.M., Jing, Z., Moreau, S., Grégori, G., Swart, S., Speich, S., Cyr, F., Legresy, B., Chao, Y., Fu, L., Morrow, R.A.: Frontiers in Fine-Scale in situ Studies: Opportunities During the SWOT Fast Sampling Phase. *Front. Mar. Sci.* 6. <https://doi.org/10.3389/fmars.2019.00168>, 2019.
- 650 Dumas, F., Garreau, P., Louazel, S., Correard, S., Fercoq, S., Le Menn, M., Serpette, A., Garnier, V., Stegner, A., Le Vu, B., Doglioli, A., Gregori, G.: PROTEVS-MED field experiments: Very High Resolution Hydrographic Surveys in the Western Mediterranean Sea. SEANOE. <https://doi.org/10.17882/62352>, 2018.
- 655 Escudier, R., Mourre, B., Juza, M., Tintoré, J.: Subsurface circulation and mesoscale variability in the Algerian subbasin from altimeter-derived eddy trajectories. *J. Geophys. Res. Oceans* 121, 6310–6322. <https://doi.org/10.1002/2016JC011760>, 2016a.
- 660 Escudier, R., Renault, L., Pascual, A., Bresseur, P., Chelton, D., Beuvier, J.: Eddy properties in the Western Mediterranean Sea from satellite altimetry and a numerical simulation. *Journal of Geophysical Research: Oceans* 121, 3990–4006. <https://doi.org/10.1002/2015JC011371>, 2016b.
- Estournel, C., Testor, P., Taupier-Letage, I., Bouin, M.-N., Coppola, L., Durand, P., Conan, P., Bosse, A., Brilouet, P.-E., Beguery, L., Belamari, S., Béranger, K., Beuvier, J., Bourras, D., Canut, G.,

- 665 Doerenbecher, A., Durrieu de Madron, X., D’Ortenzio, F., Drobinski, P., Ducrocq, V., Fourrié, N.,
Giordani, H., Houpert, L., Labatut, L., Lebeau-pin Brossier, C., Nuret, M., Prieur, L., Roussot, O.,
Seyfried, L., Somot, S.: HyMeX-SOP2: The Field Campaign Dedicated to Dense Water Formation in
the Northwestern Mediterranean. *Oceanography* 29, 196–206. <https://doi.org/10.5670/oceanog.2016.94.2016>.
- 670 Gaillard, F., Diverres, D., Jacquin, S., Gouriou, Y., Grelet, J., Menn, M.L., Tassel, J., Reverdin, G.: Sea
surface temperature and salinity from French research vessels, 2001–2013. *Scientific Data* 2, 150054.
<https://doi.org/10.1038/sdata.2015.54>, 2015.
- 675 Garreau, P., Dumas, F., Louazel, S., Stegner, A., Le Vu, B.: High-Resolution Observations and
Tracking of a Dual-Core Anticyclonic Eddy in the Algerian Basin. *Journal of Geophysical Research: Oceans* 123, 9320–9339. <https://doi.org/10.1029/2017JC013667>, 2018
- 680 Giordani, H., Lebeau-pin-Brossier, C., Léger, F., Caniaux, G.: A PV-approach for dense water formation
along fronts: Application to the Northwestern Mediterranean. *Journal of Geophysical Research: Oceans*
122, 995–1015. <https://doi.org/10.1002/2016JC012019>, 2017.
- Gosud: GOSUD -Global Ocean Surface Underway data. SEANOE. <https://doi.org/10.17882/47403>
2016.
- 685 Heslop, E.E., Sánchez-Román, A., Pascual, A., Rodríguez, D., Reeve, K.A., Faugère, Y., Raynal, M.:
Sentinel-3A Views Ocean Variability More Accurately at Finer Resolution. *Geophysical Research
Letters* 44, 12,367-12,374. <https://doi.org/10.1002/2017GL076244>, 2017.
- 690 Houpert, L., Durrieu de Madron, X., Testor, P., Bosse, A., D’Ortenzio, F., Bouin, M.N., Dausse, D., Le
Goff, H., Kunesch, S., Labaste, M., Coppola, L., Mortier, L., Raimbault, P.: Observations of open-ocean
deep convection in the northwestern Mediterranean Sea: Seasonal and interannual variability of mixing
and deep water masses for the 2007-2013 Period. *Journal of Geophysical Research: Oceans*.
<https://doi.org/10.1002/2016JC011857>, 2016.
- 695 JCOMM, 2015: Pilot intercomparison project for seawater salinity measurements, final report. World
Meteorological Organization (WMO) ; Intergovernmental Oceanographic Commission : *JCOMM
technical report* n° 84.
https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&cad=rja&uact=8&ved=2ahUKEwiw19DZ1drkAhUN6OAKHTZ5BVUQFjAAegQIBRAB&url=https%3A%2F%2Flibrary.wmo.int%2Findex.php%3Fvl%3Dnotice_display%26id%3D17098&usg=AOvVaw0vKzR29aSHFtFLIagpY0v
700 [o](https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&cad=rja&uact=8&ved=2ahUKEwiw19DZ1drkAhUN6OAKHTZ5BVUQFjAAegQIBRAB&url=https%3A%2F%2Flibrary.wmo.int%2Findex.php%3Fvl%3Dnotice_display%26id%3D17098&usg=AOvVaw0vKzR29aSHFtFLIagpY0v), 2015.
- Jones, H., Marshall, J.: Convection with Rotation in a Neutral Ocean: A Study of Open-Ocean Deep
Convection. *J. Phys. Oceanogr.* 23, 1009–1039. [https://doi.org/10.1175/1520-0485\(1993\)023<1009:CWRIAN>2.0.CO;2](https://doi.org/10.1175/1520-0485(1993)023<1009:CWRIAN>2.0.CO;2), 1993.

705

Jones, H., Marshall, J.: Restratification after Deep Convection. *J. Phys. Oceanogr.* 27, 2276–2287. [https://doi.org/10.1175/1520-0485\(1997\)027<2276:RADC>2.0.CO;2](https://doi.org/10.1175/1520-0485(1997)027<2276:RADC>2.0.CO;2), 1997.

710

Knoll, M., Borrione, I., Fiekas, H.-V., Funk, A., Hemming, M.P., Kaiser, J., Onken, R., Queste, B., Russo, A.: Hydrography and circulation west of Sardinia in June 2014. *Ocean Science* 13, 889–904. <https://doi.org/10.5194/os-13-889-2017>, 2017.

715

Le Menn, M.: About uncertainties in practical salinity calculations. *Ocean Science*, 7(5), 651–659. <https://doi.org/10.5194/os-7-651-2011>, 2011.

Le Vu, B., Stegner, A., Arsouze, T.: Angular Momentum Eddy Detection and Tracking Algorithm (AMEDA) and Its Application to Coastal Eddy Formation. *J. Atmos. Oceanic Technol.* 35, 739–762. <https://doi.org/10.1175/JTECH-D-17-0010.1>, 2017.

720

Lévy, M., Ferrari, R., Franks, P.J.S., Martin, A.P., Rivière, P.: Bringing physics to life at the submesoscale. *Geophysical Research Letters* 39. <https://doi.org/10.1029/2012GL052756>, 2012.

725

Lévy, M., Franks, P. J., & Smith, K. S.: The role of submesoscale currents in structuring marine ecosystems. *Nature communications*, 9(1), 4758, 2018.

Marshall, J., Schott, F. : Open-ocean convection: Observations, theory, and models. *Rev. Geophys.* 37, 1–64. <https://doi.org/10.1029/98RG02739>, 1999.

730

McWilliams, J.C.: Submesoscale, coherent vortices in the ocean, *Rev. Geophys.*, 23(2), 165, doi:10.1029/RG023i002p00165, 1985.

735

McWilliams James C.: Submesoscale currents in the ocean. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences* 472, 20160117. <https://doi.org/10.1098/rspa.2016.0117>, 2016.

Medoc Group : Observation of Formation of Deep Water in the Mediterranean Sea 1969. *Nature* 227, 1037–1040. <https://doi.org/10.1038/2271037a0> ,1970.

740

Meloni, M., Bouffard, J., Doglioli, A.M., Petrenko, A.A., Valladeau, G.: Toward science-oriented validations of coastal altimetry: application to the Ligurian Sea. *Remote Sens. Environ.*, 224, 275–288, <https://doi.org/10.1016/j.rse.2019.01.028>, 2019.

745

Margirier, F., Bosse, A., Testor, P., L'Hévéder, B., Mortier, L., Smeed, D.: Characterization of Convective Plumes Associated With Oceanic Deep Convection in the Northwestern Mediterranean

- From High-Resolution In Situ Data Collected by Gliders. *Journal of Geophysical Research: Oceans* 122, 9814–9826. <https://doi.org/10.1002/2016JC012633>, 2017.
- 750 Marrec, P., Grégori, G., Doglioli, A.M., Dugenne, M., Della Penna, A., Bhairy, N., Cariou, T., Hélias Nunige, S., Lahbib, S., Rougier, G., Wagener, T., Thyssen M.: Coupling physics and biogeochemistry thanks to high resolution observations of the phytoplankton community structure in the North-Western Mediterranean Sea. *Biogeosciences*, 15, 1579-1606, <https://doi.org/10.5194/bg-15-1579-2018>, 2018.
- 755 Millot, C., Taupierletage, I., Benzohra, M.: The Algerian Eddies. *Earth-Sci. Rev.* 27, 203–219. [https://doi.org/10.1016/0012-8252\(90\)90003-E](https://doi.org/10.1016/0012-8252(90)90003-E), 1990.
- Millot, C.: Circulation in the Western Mediterranean Sea. *Journal of Marine Systems* 20, 423–442. [https://doi.org/10.1016/S0924-7963\(98\)00078-5](https://doi.org/10.1016/S0924-7963(98)00078-5), 1999.
- 760 Millot, C., Taupier-Letage, I. : Circulation in the Mediterranean Sea, in : *The Mediterranean Sea, Handbook of Environmental Chemistry*, edited by: Saliot, A., Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 29–66. <https://doi.org/10.1007/b107143>, 2005.
- 765 Nencioli, F., d’Ovidio, F., Doglioli, A.M., Petrenko, A.A. : Surface coastal circulation patterns by in-situ detection of Lagrangian coherent structures. *Geophysical Research Letters* 38. <https://doi.org/10.1029/2011GL048815>, 2011
- 770 Niewiadomska, K., Claustre, H., Prieur, L., d’Ortenzio, F.: Submesoscale physical-biogeochemical coupling across the Ligurian current (northwestern Mediterranean) using a bio-optical glider. *Limnology and Oceanography* 53, 2210–2225. https://doi.org/10.4319/lo.2008.53.5_part_2.2210, 2008.
- 775 Onken, R., Brambilla, E. : Double diffusion in the Mediterranean Sea: Observation and parameterization of salt finger convection. *Journal of Geophysical Research: Oceans* 108. <https://doi.org/10.1029/2002JC001349>, 2003.
- 780 Onken, R., Fiekas, H.-V., Beguery, L., Borrione, I., Funk, A., Hemming, M., Hernandez-Lasheras, J., Heywood, K.J., Kaiser, J., Knoll, M., Mourre, B., Oddo, P., Poulain, P.-M., Queste, B.Y., Russo, A., Shitashima, K., Siderius, M., Thorp Küsel, E.: High-resolution observations in the western Mediterranean Sea: the REP14-MED experiment. *Ocean Science* 14, 321–335. <https://doi.org/10.5194/os-14-321-2018>, 2018.
- 785 Pascual, A., Ruiz, S., Olita, A., Troupin, C., Claret, M., Casas, B., Mourre, B., Poulain, P.-M., Tovar-Sanchez, A., Capet, A., Mason, E., Allen, J.T., Mahadevan, A., Tintoré, J.: A Multiplatform Experiment to Unravel Meso- and Submesoscale Processes in an Intense Front (AlborEx). *Front. Mar. Sci.* 4. <https://doi.org/10.3389/fmars.2017.00039>, 2017.

- Petrenko, A.A., Doglioli, A.M., Nencioli, F., Kersalé, M., Hu, Z., d'Ovidio, F.: A review of the LATEX project: mesoscale to submesoscale processes in a coastal environment. *Ocean Dynam.*, 67:513, 790 <https://doi.org/doi:10.1007/s10236-017-1040-9>, 2017.
- Puillat, I., Taupier-Letage, I., Millot, C.: Algerian Eddies lifetime can near 3 years. *Journal of Marine Systems* 31, 245–259. [https://doi.org/10.1016/S0924-7963\(01\)00056-2](https://doi.org/10.1016/S0924-7963(01)00056-2), 2002.
- 795 Robinson, A.R., Golnaraghi, M.: The Physical and Dynamical Oceanography of the Mediterranean Sea, in: Malanotte-Rizzoli, P., Robinson, A.R. (Eds.), *Ocean Processes in Climate Dynamics: Global and Mediterranean Examples*. Springer Netherlands, Dordrecht, pp. 255–306. https://doi.org/10.1007/978-94-011-0870-6_12, 1994.
- 800 Rousselet L., Doglioli, A.M., de Verneil, A., Pietri, A., Della Penna, A., Berline, L., Marrec, P., Gregori, G., Thyssen, M., Carlotti, F., Barillon, S., Simon-Bot, F., Bonal, M., d'Ovidio, F. and Petrenko, A.A.: Vertical motions and their effects on a biogeochemical tracer in a cyclonic structure finely observed in the Ligurian Sea. *J.Geophys.Res.*, 124, <https://doi.org/10.1029/2018JC014392>, 2019.
- 805 Ruiz, S., Pascual, A., Garau, B., Faugère, Y., Alvarez, A., Tintoré, J.: Mesoscale dynamics of the Balearic Front, integrating glider, ship and satellite data. *Journal of Marine Systems, Coastal Processes: Challenges for Monitoring and Prediction* 78, S3–S16. <https://doi.org/10.1016/j.jmarsys.2009.01.007>, 2009.
- 810 Salat, J., Emelianov, M., Frail, E., Latasa, M.: After deep water formation: sinking and spreading or reorganising phase, including upwelling? *Rapp. Comm. int. Mer Médit.*40. http://www.ciesm.org/online/archives/abstracts/pdf/40/Pg_0175.pdf, 2013.
- Schott, F., Visbeck, M., Send, U., Fischer, J., Stramma, L., Desaubies, Y. : Observations of Deep Convection in the Gulf of Lions, Northern Mediterranean, during the Winter of 1991/92. *J. Phys. Oceanogr.* 26, 505–524. [https://doi.org/10.1175/1520-0485\(1996\)026<0505:OODCIT>2.0.CO;2](https://doi.org/10.1175/1520-0485(1996)026<0505:OODCIT>2.0.CO;2), 1996.
- 815 Send, U., Font, J., Krahnemann, G., Millot, C., Rhein, M., Tintoré, J.: Recent advances in observing the physical oceanography of the western Mediterranean Sea. *Progress in Oceanography* 44, 37–64. 820 [https://doi.org/10.1016/S0079-6611\(99\)00020-8](https://doi.org/10.1016/S0079-6611(99)00020-8), 1999.
- Sverdrup, H. U., Johnson, M. W. & Fleming, R. H.: *The oceans: their physics, chemistry, and general biology*. Englewood Cliffs, NJ: Prentice-Hall, 1942.
- 825 Szekely, T., Gourrion, J., Pouliquen, S., Reverdin, G.: CORA, Coriolis Ocean Dataset for Reanalysis. SEANOE. <http://doi.org/10.17882/46219>, 2016.

- 830 Taupier-Letage, I., Puillat, I., Millot, C., Raimbault, P.: Biological response to mesoscale eddies in the Algerian Basin. *Journal of Geophysical Research: Oceans* 108. <https://doi.org/10.1029/1999JC000117>, 2003.
- 835 Testor, P., Gascard, J.-C. : Large-Scale Spreading of Deep Waters in the Western Mediterranean Sea by Submesoscale Coherent Eddies. *J. Phys. Oceanogr.* 33, 75–87. [https://doi.org/10.1175/1520-0485\(2003\)033<0075:LSSODW>2.0.CO;2](https://doi.org/10.1175/1520-0485(2003)033<0075:LSSODW>2.0.CO;2), 2003.
- Testor, P., Gascard, J.-C.: Post-convection spreading phase in the Northwestern Mediterranean Sea. *Deep Sea Research Part I: Oceanographic Research Papers* 53, 869–893. <https://doi.org/10.1016/j.dsr.2006.02.004>, 2006.
- 840 Testor, P., Bosse, A., Houpert, L., Margirier, F., Mortier, L., Legoff, H., Dausse, D., Labaste, M., Karstensen, J., Hayes, D., Olita, A., Ribotti, A., Schroeder, K., Chiggiato, J., Onken, R., Heslop, E., Mourre, B., D’Ortenzio, F., Mayot, N., Lavigne, H., de Fommervault, O., Coppola, L., Prieur, L., Taillandier, V., Durrieu de Madron, X., Bourrin, F., Many, G., Damien, P., Estournel, C., Marsaleix, P.,
- 845 Ducrocq, V., Conan, P.: Multiscale Observations of Deep Convection in the Northwestern Mediterranean Sea During Winter 2012-2013 Using Multiple Platforms. *J. Geophys. Res. Oceans* 123,1745-1776 <https://doi.org/10.1002/2016JC012671>, 2018.
- 850 Tintore, J., Pinardi, N., Álvarez-Fanjul, E., Aguiar, E., Álvarez-Berastegui, D., Bajo, M., Balbin, R., Bozzano, R., Nardelli, B.B., Cardin, V., Casas, B., Charcos-Llorens, M., Chiggiato, J., Clementi, E., Coppini, G., Coppola, L., Cossarini, G., Deidun, A., Deudero, S., D’Ortenzio, F., Drago, A., Drudi, M., El Serafy, G., Escudier, R., Farcy, P., Federico, I., Fernández, J.G., Ferrarin, C., Fossi, C., Frangoulis, C., Galgani, F., Gana, S., García Lafuente, J., Sotillo, M.G., Garreau, P., Gertman, I., Gómez-Pujol, L., Grandi, A., Hayes, D., Hernández-Lasheras, J., Herut, B., Heslop, E., Hilmi, K., Juza, M., Kallos, G.,
- 855 Korres, G., Lecci, R., Lazzari, P., Lorente, P., Liubartseva, S., Louanchi, F., Malacic, V., Mannarini, G., March, D., Marullo, S., Mauri, E., Meszaros, L., Mourre, B., Mortier, L., Muñoz-Mas, C., Novellino, A., Obaton, D., Orfila, A., Pascual, A., Pensieri, S., Pérez Gómez, B., Pérez Rubio, S., Perivoliotis, L., Petihakis, G., de la Villéon, L.P., Pistoia, J., Poulain, P.-M., Pouliquen, S., Prieto, L., Raimbault, P., Reglero, P., Reyes, E., Rotllan, P., Ruiz, S., Ruiz, J., Ruiz, I., Ruiz-Orejón, L.F., Salihoglu, B., Salon, S., Sammartino, S., Sánchez Arcilla, A., Sánchez-Román, A., Sannino, G., Santoleri, R., Sardá, R.,
- 860 Schroeder, K., Simoncelli, S., Sofianos, S., Sylaios, G., Tanhua, T., Teruzzi, A., Testor, P., Tezcan, D., Torner, M., Trotta, F., Umgiesser, G., von Schuckmann, K., Verri, G., Vilibic, I., Yucel, M., Zavatarelli, M., Zodiatis, G.: Challenges for Sustained Observing and Forecasting Systems in the Mediterranean Sea. *Front. Mar. Sci.* 6. <https://doi.org/10.3389/fmars.2019.00568>, 2019.
- 865 Troupin, C., Pascual, A., Ruiz, S., Olita, A., Casas, B., Margirier, F., Poulain, P.-M., Notarstefano, G., Torner, M., Fernández, J.G., Rújula, M.À., Muñoz, C., Alou, E., Ruiz, I., Tovar-Sánchez, A., Allen, J.T., Mahadevan, A., Tintoré, J.: The AlborEX dataset: sampling of sub-mesoscale features in the Alboran Sea. *Earth System Science Data* 11, 129–145. <https://doi.org/10.5194/essd-11-129-2019>, 2019.

Visbeck, M.: Deep Velocity Profiling Using Lowered Acoustic Doppler Current Profilers: Bottom Track and Inverse Solutions. *J. Atmos. Oceanic Technol.* 19, 794–807. [https://doi.org/10.1175/1520-0426\(2002\)019<0794:DVPULA>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0794:DVPULA>2.0.CO;2), 2002.