

Sval_Imp: A gridded forcing dataset for climate change impact research on Svalbard

Thomas Vikhamar Schuler¹ and Torbjørn Ims Østby^{1,2}

¹Department of Geosciences, University of Oslo, Norway

²now at: Lyse AS, Stavanger, Norway

Correspondence: Thomas V Schuler (t.v.schuler@geo.uio.no)

Abstract. We present Sval_Imp, a high resolution gridded dataset designed for forcing models of terrestrial surface processes on Svalbard. The dataset is defined on a 1km grid covering the archipelago of Svalbard, located in the Norwegian Arctic (74-82°N). Using a hybrid methodology combining multi-dimensional interpolation with simple dynamical modelling, the atmospheric reanalyses ERA-40 and ERA-interim by the European Centre for Medium-Range Weather Forecasting have been
5 downscaled to cover the period 1957-2017 at steps of 6h. The dataset is publicly available from a data repository. In this paper, we describe the methodology used to construct the dataset, present the organization of the data in the repository and discuss the performance of the downscaling procedure. In doing so, the dataset is compared to a wealth of data available from operational as well as project-based measurements. The quality of the downscaled dataset is found to vary in space and time, but generally represents an improvement compared to unscaled values, especially for precipitation. Whereas operational records are biased to
10 low-elevations around the fringes, we stress the hitherto under-used potential of project-based measurements at higher elevation and in the interior of the archipelago for evaluating atmospheric models. For instance, records of snow accumulation on large ice masses may represent measures of seasonally-integrated precipitation in regions sensitive to the downscaling procedure, thus providing added value.

Sval_Imp (Schuler, 2018) is publicly available from the Norwegian Research Data Archive NIRD, a data repository (<https://doi.org/10.11582/2018.00006>).
15

Copyright statement. Sval_Imp is licensed under the Creative Commons Attribution 4.0 International License (CC-BY 4.0). In essence you are free to copy, distribute, and adapt the work, as long as you attribute the work to its origin and abide by the other license terms.

1 Introduction

The non-linearity of many surface processes poses challenges on appropriateness of atmospheric forcing for impact studies
20 in terms of accuracy and precision (e.g., Liston and Elder, 2006). Especially in mountainous areas, the variability of surface systems is typically governed by spatial scales not resolved in regional climate models and adjustments have to be made to overcome this (e.g., Fiddes and Gruber, 2014). A variety of methods has been developed for this purpose, differing in terms of

data requirements and computational cost. While empirical-statistical scaling requires reference data for training and assumes a temporal robustness of the employed statistical relations (e.g., Ehret et al., 2012; Maraun, 2013), dynamic downscaling by means of high-resolution atmospheric modelling has high computational costs (e.g., Gutmann et al., 2016).

In this paper, we present Sval_Imp (Schuler, 2018), a high resolution gridded dataset obtained using a hybrid methodology combining multi-dimensional interpolation with simple dynamical modelling. The dataset is defined on a 1km grid covering the archipelago of Svalbard, located in the Norwegian Arctic (74-82°N, 10-35°E). The atmospheric reanalyses ERA-40 and ERA-interim by the European Centre for Medium-Range Weather Forecasting have been downscaled to cover the period 1957-2017 at 6 hours temporal resolution. The dataset comprises the near-surface variables required to compute the surface energy balance, namely air temperature, precipitation, relative humidity, wind speed, and downwelling components of shortwave (solar) and longwave (thermal) radiation. Sval_Imp is publicly available from the Norwegian Research Data Archive NIRD, a data repository (<https://doi.org/10.11582/2018.00006>). In the following, we describe the methodology used to derive the dataset, present the organization of the data in the repository and discuss the performance of the downscaling procedure. For the latter, the dataset is compared to a wealth of data available from long-term operational as well as short-term scientific records of meteorological and glaciological measurements. Operational records are biased to low-elevations around the fringes of the archipelago. Therefore, we stress the hitherto under-used potential of project-based measurements in the interior, high-elevation regions for evaluating atmospheric models. For instance, records of snow accumulation on large ice masses may represent measures of seasonally-integrated precipitation in regions sensitive to the downscaling procedure, thus providing added-value. Sval_Imp has been employed entirely or in parts by a range of projects for forcing process models of the surface energy and mass balances of glaciers (Østby et al., 2017), precipitation patterns and meltwater production in the Kongsfjord area (Pramanik et al., 2018), for assimilation of remotely-sensed snow cover using a snow distribution model (Aalstad et al., 2018) as well as for assessing growing conditions for fungi (Botnen et al., in prep.). Further, the dataset has been used to assess changes and trends in climate conditions of Svalbard (Hanssen-Bauer et al., 2019).

2 Methodology

To generate fields of near-surface air temperature, precipitation, relative humidity, wind and downwelling shortwave and longwave radiation, we have downscaled the ERA-40 and ERA-interim reanalyses of the European Centre for Medium-Range Weather Forecasts (Uppala et al., 2005; Dee et al., 2011). The reanalysis data is provided at 6-hour intervals and has been retrieved on a $0.75^\circ \times 0.75^\circ$ spatial grid covering the periods 1957-2002 (ERA-40) and 1979-2017 (ERA-interim). These data have been downscaled to a 1km grid covering the region of interest (Fig. 1) using the scheme described in the following to produce Sval_Imp, a high-resolution, 6-hour dataset of precipitation, temperature, relative humidity, windspeed as well as downwelling shortwave and longwave radiation fluxes.




Figure 1. The map shows the Svalbard archipelago, shading indicates surface elevation, glacierized areas are outlined in blue. The locations of meteorological stations are represented by red diamonds, the numbers refer to the station names: 1 - Etonbreen, 2 - Janssonhaugen, 3 - Gruvefjellet, 4 - Kapp Heuglin, 5 - Rijpfjorden, 6 - Svalbard Airport, 7 - Isfjord Radio, 8 - Verlegenuken, 9 - Hornsund, 10 - Kvitøya, 11 - Holtedahlfonna, 12 - Ny-Ålesund, 13 - Hopen. The black crosses indicate the grid points of the ERA reanalyses, and the grey countourlines indicate the topography of Svalbard in the ERA reanalyses at 0, 100, 200 and 300 m asl.

2.1 Downscaling

Precipitation is often heavily biased in coarsely-resolved reanalyses, especially in environments with pronounced topography, where it typically is too low and lacks spatial detail (Schuler et al., 2008). This is associated with the smoothed representation of the actual topography in the large-scale model used for the reanalysis (Fig. 1), leading to an underestimate of orographic precipitation. Figure 1 shows that ERA greatly generalizes the high resolution topography, representing Svalbard as a wide and flat bump, that exceeds sea-level far off the actual coastlines, while surface elevation in the interior does not exceed 400 m asl. In contrast, the highest elevation in our gridded topography map is 1600 m asl. The roughness of the actual topography that gave rise to the name of the main island 'Spitsbergen' (sharp tops) is not represented by the smoothed topography used for the ERA reanalyses. We assume that this is the main reason for the poor performance of reanalyzed precipitation. To account for orographic enhancement, we use a linear theory (LT) of orographic precipitation (Smith and Barstad, 2004).

The other required climate variables are downscaled to the 1 km grid largely following the TopoSCALE methodology (Fiddes and Gruber, 2014) which also builds on the assumption that weaknesses in the representation of topography at the coarse scale are mainly responsible for the misfit between coarse scale and point observations. TopoSCALE exploits the relatively high vertical resolution of the reanalysis data to downscale variables to the elevation of the actual topography, based on the properties of the vertical structure in the reanalysis. The downscaled fields preserve the horizontal gradients present in ERA, but include additional features caused by the real topography which were not present in the ERA products. In doing so, we add spatial detail to the reanalysis fields that is consistent with the temporal evolution of atmospheric conditions of the reanalysis. This approach is assumed to outperform simpler bias corrections, since transient properties of the atmosphere are accounted for. For example, transient lapse rates including inversions in the reanalysis data will be preserved in the downscaled product.

In our application, we modified the TopoSCALE methodology regarding downscaling of direct shortwave radiation and air temperature, as described in Sections 2.3 and 2.4.

2.2 Precipitation

The LT-model describes an air parcel as it moves across a prescribed surface topography. The air parcel is characterized by its temperature, stability, wind direction and speed. Terrain induced uplift of the air parcel results in condensation and eventually precipitation of moisture downstream of the uplift. This model has been successfully evaluated using precipitation gauges (Barstad and Smith, 2005) and snow measurements (Schuler et al., 2008; Østby et al., 2017) and applied for downscaling precipitation (e.g. Crochet et al., 2007; Jarosch et al., 2012; Roth et al., 2018). The linear theory utilizes a Boussinesq description of mountain wave to derive a transfer function that, for given wind conditions, relates the orographically enhanced precipitation to terrain topography.

By spectral decomposition and algebraic manipulation, Smith and Barstad (2004) derived the following transfer function

$$\hat{P}(k, l) = \frac{C_w i \sigma \hat{h}(k, l)}{(1 - i m H_w)(1 + i \sigma \tau_f)(1 + i \sigma \tau_c)}. \quad (1)$$

85 relating $\hat{P}(k, l)$, the Fourier-transform of the precipitation enhancement, to the Fourier-transform of terrain elevation $\hat{h}(k, l)$,
with k and l being the horizontal wave numbers. This relation depends on the uplift sensitivity factor C_w , thickness of the moist
layer H_w , the intrinsic frequency $\sigma(k, l) = Uk + Vl$ (U and V being the east and north components of the wind vector), and
the conversion and fallout time scales τ_c and τ_f , respectively. In Equation (1), the vertical wave number m controls the depth
90 and tilt of the forced air uplift and is a function of the moist Brunt-Väisälä frequency, N_m , a quantity describing atmospheric
stability.

Precipitation rates are obtained by retransforming \hat{P} and adding it to the background precipitation P_∞ , that accounts for
large-scale frontal and convective precipitation separate from orographic precipitation P_{oro} . P_∞ has been corrected for the
orographic effect already present in the ERA reanalyses $P_{\text{oro}}(h_{\text{ERA}})$ by estimating this effect applying Equation 1 to the large-
scale topography h_{ERA} and removing the result from the ERA precipitation $P_\infty = P_{\text{ERA}} - P_{\text{oro}}(h_{\text{ERA}})$ (Schuler et al., 2008).

95 Total precipitation, P_{total} , is then

$$P_{\text{total}}(x, y) = \max \left[f \iint \hat{P}(k, l) e^{i(kx + yl)} dk dl + P_\infty, 0 \right]. \quad (2)$$

Since the theory assumes saturated conditions, we account for reduced orographic enhancement at lower humidity by adopting
a correction factor f proposed by Sinclair (1994)

$$f = \begin{cases} \left(\frac{RH - 0.8}{0.2} \right)^{1/4} & : RH \geq 0.8 \\ 0 & : \text{otherwise} \end{cases} \quad (3)$$

100 which suppresses orographic enhancement when $RH < 0.8$.

Instead of treating N_m , τ_f and τ_c as adjustable, constant parameters, we exploit the evolution of the moisture bearing layer
of the atmosphere, described in the reanalyses to derive transient values. In doing so, we remove calibration parameters from
our method and enable weather dependent variation of N_m , τ_f and τ_c . Values of N_m are calculated following

$$N_m^2 = \frac{g}{\bar{T}} (\Gamma_m - \Gamma_e) \quad (4)$$

105 where g is gravitational acceleration and \bar{T} is vertically averaged air temperature weighted by the moisture content at several
pressure levels (Jarosch et al., 2012). Environmental lapse-rates Γ_e are derived from air temperature at 700 hPa and 850 hPa
and corresponding geopotential heights. Γ_m is the moist adiabatic lapse rate, calculated according to Stone and Carlson (1979)
using vertically averaged values of atmospheric properties from the reanalyses weighted by moisture content. This follows
the convention that a positive lapse rate represents cooling with increasing elevation. Barstad and Smith (2005) report that
110 typical values of N_m range between 0 s^{-1} , representing an atmosphere with no stratification, and 0.01 s^{-1} representing a
stably stratified atmosphere. To avoid conditions inconsistent with the assumptions of the theory, we limit N_m to this range .
The quantities H_w , C_w and m are derived from N_m (Smith and Barstad, 2004).

Advection time scales τ_c and τ_f are assumed equal and $\tau = \tau_c = \tau_f = H_w/v$ is derived from the thickness of the moist
layer H_w and accounting for a typical hydrometeor fall speed v , which is taken as constant but allowed to take different
115 values for solid and liquid hydrometeors. This phase transition is determined by a threshold temperature of 273 K, hence




Figure 2. Example plot of longterm mean annual precipitation (m) over 1979-2017, downsampled from ERA-interim.

$v(T \leq 273K) = 1 \text{ ms}^{-1}$ for solid and $v(T \geq 273K) = 2 \text{ ms}^{-1}$ for liquid precipitation. In Equation 1, terrain elevation h is the only gridded variable, the other variables represent averages over the volume of the described air parcel. To characterize this air parcel, we first vertically average the values defined at the nodes of the horizontal domain over the 700 hPa and 850 hPa pressure-levels weighted by the moisture content of the individual layers. These vertically averaged values are then horizontally averaged over an area defined by a 200 km buffer around the 200 m contour of the reanalysis topography, the latter roughly outlining the extend of the archipelago (Fig. 1).

This setup is then applied to each 6h timestep and the resulting timeslices are progressively added to the record.

2.3 Air temperature

The downscaling for near-surface air temperature at 2 m level (T_2) closely follows the TopoSCALE procedure Fiddes and Gruber (2014), thereby we assume that the vertical structure of the free atmosphere determines the distribution of T_2 with terrain elevation. For each 6h-time step, T_2 is derived from a three-dimensional interpolation of the vertical air temperature

structure of the large-scale reanalysis to the location of the grid nodes representing the high-resolution terrain elevation. We notice, that for a melting snow or ice surface, skin temperature is bounded to 273 K, influencing the near surface air temperature and resulting in reduced along-surface air temperature lapse-rates (e.g., Marshall et al., 2007). To account for this effect, we apply a simple horizontal interpolation of the two-dimensional $T2_{\text{ERA}}$ field where $T2 > 273$ K, instead of interpolating a three-dimensional data volume to the surface elevation of the high resolution topography. This strategy is motivated by the discovery of unrealistic warm air temperatures at higher elevations in a test application. The occurrence of this unphysical temperature inversion was restricted to the melting period, caused by extrapolation of surface inversions close to a snow/ ice surface the temperature of which is capped at 273 K. To avoid this effect, we assume that where $T2 > 273$ K, the $T2$ of the reanalysis is consistent with a melting surface and hence more realistic than a free-atmosphere interpolation.




Figure 3. Example plot of mean air temperature (2 m) in $^{\circ}\text{C}$, over the period 1979-2017, downscaled from ERA-interim.

2.4 Radiation

Downscaling of shortwave and longwave downwelling radiative fluxes (SW and LW , respectively) was conducted by adopting the TopoSCALE methodology (Fiddes and Gruber, 2014) with a few adjustments. The shortwave radiation at surface level




Figure 4. a) Example plot of 1979-2017 incident shortwave radiation (W m^{-2}), downscaled from ERA-interim; b) comparison between measured (Maturilli et al., 2013) and downscaled daily values of SW at Ny-Ålesund. The color indicates the density of points from red (high) to green (low).

of the reanalysis is projected to the high-resolution topography in a three-step procedure: first the surface SW flux is separated into direct and diffuse components; second, the direct component is corrected for the elevation difference between the reanalysis surface and the high-resolution topography considering an effective atmospheric transmissivity that is derived from top-of-atmosphere and surface fluxes; third a topographic correction is applied to account for effects of slope and aspect of the high-resolution topography, as well as shading by surrounding topography. To compute direct solar radiation we apply the relationship of Kumar et al. (1997) for atmospheric attenuation rather than the one given by Fiddes and Gruber (2014). Solar geometry variables such as solar zenith and azimuth, and topographic shading due to local slope and aspect are calculated following Reda and Andreas (2004). Cast shadow and hemispherical obstructions caused by surrounding topography are calculated following Ratti (2001). The longwave surface flux is downscaled by correcting for the elevation difference between reanalysis and high-resolution grids using an atmospheric emissivity. This emissivity is estimated by accounting for a clear-sky component that depends on humidity and air temperature and a cloud component that is estimated from the difference between the clear-sky component and the reanalysis longwave flux. Further terrain effects are incorporated through multiplication with the sky-view factor.

2.5 Relative humidity and windspeed

Similar to our assumption about T_2 over a melting surface, we suggest that windspeed and RH in the boundary layer are more affected by surface rather than by free-atmosphere conditions and we hence apply a simple two-dimensional interpolation of




Figure 5. a) Example plot of 1979-2017 downwelling longwave radiation (W m^{-2}), downscaled from ERA-interim; b) comparison between measured (Maturilli et al., 2013) and downscaled daily values of LW at Ny-Ålesund. The color indicates the density of points from red (high) to green (low).

155 the near-surface values of the reanalysis, instead of interpolating a three-dimensional data volume to the surface elevation of the high resolution topography.

3 Performance evaluation

Østby et al. (2017) and Vikhamar-Schuler et al. (2019) have conducted thorough evaluations of the Sval_Imp dataset using data from meteorological stations. Here, we summarize their main results in the subsequent subsections and refer to Østby et al. (2017) for details. Furthermore, we present additional evaluation of the precipitation using snow measurements. Typically, meteorological records are available as daily mean values and 6-hourly, downscaled variables have been temporally aggregated to match the time step of measurements. Precipitation in reanalysis is not constrained by data assimilation, giving rise to uncertainty in timing as well as amount. Our downscaling aims for reducing the bias in precipitation amount but does not treat the timing. For a performance evaluation, we therefore use monthly precipitation sums which are regarded as robust against timing mismatches, whereas a higher temporal resolution could penalize a method that otherwise is successful in reducing the underestimation of the reanalysis.

3.1 Precipitation

At the operational weather stations (Fig. 1), downscaled precipitation (Fig. 2) is overestimated by 5 to 25 mm per month at the weather stations, with a slightly higher bias during winter (Table 1). This is also consistent with the findings of Vikhamar-Schuler et al. (2019) who evaluated the performance at 6 weather stations for several 30-year reference periods (1961-1990,

1971-2000, and 1988-2017). These biases are partly caused by too low precipitation measurements, the values of which are heavily affected by wind-induced undercatch, especially for solid precipitation. Førland and Hanssen-Bauer (2000) suggest that for solid precipitation, the actual precipitation at wind-exposed sites may be up to 80% higher than the gauge record. A newly developed correction scheme for Norwegian mountain environments (Wolff et al., 2015) supports the finding of large
175 undercatch for solid precipitation, however, this corrections has not yet been applied for arctic conditions in Svalbard.

In addition to gauge measurements from low-elevation stations along the coast, we also used snow survey transects across Austfonna, a large ice cap in NE Svalbard (Fig. 1) to evaluate the Sval_Imp precipitation. We suggest that snow deposition on large glacier areas, measured at the end of the winter, represents seasonally integrated precipitation. While these measurements do not allow temporal resolution below one snow season (typically Oct-May), they provide useful information about spatial
180 precipitation patterns, in areas and elevations not covered by the operational meteorological stations. This approach builds on the implicit assumption of negligible sublimation, such that accumulated snow water equivalent represents the sum of precipitation over the winter season. Svalbard has high air humidity throughout the year, limiting the potential for sublimation. In an energy-balance study, Østby et al. (2017) estimated sublimation to about $0.016 \text{ m w.e. year}^{-1}$, which is 1-2 orders of magnitude smaller than typical annual precipitation sums. There is generally good agreement concerning the spatial pattern
185 across the Austfonna ice cap (Fig. 6), where snow accumulation reveals a distinctive SE-NW asymmetry (Taurisano et al., 2007; Schuler et al., 2007; Dunse et al., 2009) caused by orographic enhancement of precipitation coming from the SE sector (i.e., the Barents Sea), although in individual years the downscaled values underestimate the measured accumulation, especially in 2007 (Fig. 6). Even though, there is considerable scatter between observed and downscaled winter precipitation, there is positive correlation, indicating that the spatial pattern is matched and in most years there is no systematic bias, showing that
190 the overall precipitation amount is adequately represented. On the other hand, the unscaled ERA-interim winter precipitation shows almost no spatial variation and considerably underestimates observed values (Fig. 2).

Østby et al. (2017) found that the winter mass balance of Hansbreen was not well reproduced both in terms of spatial pattern as well as accumulation amount. Aas et al. (2016) similarly reported the lowest performance for Hansbreen although they had used a much more complex precipitation scheme than the one presented here. The generally low performance of several
195 precipitation distribution schemes compared to the Hansbreen record has been interpreted to result from local conditions at Hansbreen where the spatial distribution of snow is caused by wind redistribution rather than by the spatial precipitation pattern (Grabiec et al., 2006).

3.2 Air temperature

Downscaled air temperatures (Fig. 3) are compared to observations at the meteorological stations listed in Table 1 mostly for the
200 period after 2004. Despite altitude differences of up to 100 m between measuring site and corresponding grid node in the model, no altitude correction is performed, due to unknown lapse rates. In general the agreement is good between downscaled ERA and observed air temperatures, with biases mostly below 1.5 K (Tab. 1). Despite a small bias for mean annual temperatures, there is a clear seasonal bias, with ERA temperatures too warm during winter and too cold during summer (Table 1). Although the biases are negative during summer, ERA is too warm over the glaciers during summer, when 2-m air temperatures are above




Figure 6. a) Map view of Sval_Imp precipitation accumulated over Oct 1998 to Mar 1999, overlaid by coloured circles, indicating the measured snow water equivalent (SWE) by Sand et al. (2003). b) scatterplot comparing the 1999 measurements to winter precipitation according to Sval_Imp (crosses) and to ERA-interim (dots). c) similar scatterplot as in b) but only for the Austfonna dataset that provides multi-temporal coverage (Taurisano et al., 2007; Dunse et al., 2009).

205 freezing. These findings are consistent with those of Vikhamar-Schuler et al. (2019) who evaluated differences in seasonal mean values for different 30-year periods 1961-1990, 1971-2000 and 1988-2017.

At Svalbard Airport, the performances of downscaled ERA-40 and ERA-interim are investigated for the entire model period. Over 1957-1979 only monthly measured air temperatures are available at Svalbard Airport, where downscaled ERA-40 has a monthly root mean square error (RMSE) of 2.3 °C. For the 1979-2002 period the reanalysis products overlap with monthly
 210 RMSE of 1.8 °C and 1.5 °C at Svalbard Airport for ERA-40 and ERA-interim, respectively. We attribute the lower performance prior to 1979 to the lack of satellite observations to constrain sea surface temperatures and sea ice cover in the reanalysis. Since the Svalbard Airport air temperature record and other sites at the west coast are likely incorporated into the reanalysis, the quality of the reanalysis in the pre-satellite era is possibly even lower in remote areas with no observations. Due to sparsity of available data in the overlap period 1979-2002, we have not evaluated the performance of Sval_Imp for other variables than
 215 air temperature. However, Østby et al. (2017) have evaluated the effect of this dataset discontinuity by simulating glacier mass balance using both, ERA-40 and ERA-interim to investigate whether this discontinuity could be responsible for a notable drop

in simulated mass balance around year 1980. They found that the ERA-40 based simulation yields an about 0.13 m w.e. higher mass balance than the one based on ERA-interim, but ERA-40 based simulations still show a 0.2 m drop of mass balance between 1970 and 1990, larger than that caused by the data set discontinuity. This suggests that the change in mass balance regime was not caused by the heterogeneity of our composite forcing. Nevertheless, we cannot rule out the possibility that this change was caused by the discontinuity inherent in both reanalyses due to the availability of satellite observations after 1979. The annual observed air temperature trend for the period 1957-2013 at Svalbard Airport is 0.70 ± 0.22 °C decade⁻¹, while the downscaled ERA data has an insignificantly lower warming trend of 0.67 ± 0.19 °C decade⁻¹ at Svalbard Airport.

3.3 Radiation

To evaluate the quality of Sval_Imp radiation components, we use available records from two stations, roughly 300 km apart from each other. The record from Ny-Ålesund is from a daily serviced Baseline Surface Radiation Network station (Maturilli et al., 2013) whereas the Austfonna measurements are collected by an autonomously recording weather station (Schuler et al., 2014).

In Ny-Ålesund the model largely reproduces observations both for short and longwave radiation (Figs. 5 and 4, Tab. 1). During winter, downwelling longwave radiation is slightly underestimated, while there is no bias during summer. Since there is no air temperature bias in Ny-Ålesund during winter, the underestimation of longwave radiation is indicative of a too thin cloud cover in the reanalysis. In general, the representation of clouds are among the major issues of the reanalysis (Aas et al., 2016). Downwelling shortwave radiation is overestimated by 7 W m^{-2} over the summer season in Ny-Ålesund. There is a much better agreement with radiation observations in Ny-Ålesund than on Etonbreen (Fig. 1), in northeastern Svalbard. This is to be expected, since radio soundings and other observation data from Ny-Ålesund are assimilated into ERA-interim. Therefore, cloud cover at Ny-Ålesund is much better represented by the reanalysis than at Austfonna. On Etonbreen during summer, Sval_Imp underestimates downwelling shortwave radiation by 40 W m^{-2} while downwelling longwave radiation is overestimated by 12 W m^{-2} , both indicative of a too thick atmosphere or too many clouds in the reanalysis. However, these biases could also be partly explained by measurement uncertainty caused by rime on the sensor or by sensor tilt. The latter issue is caused by the fact that the ice foundation of an autonomous weather stations may melt and deform causing tilt and thereby large errors, especially at high solar zenith angles (Bogren et al., 2016).

3.4 Relative humidity and windspeed

For relative humidity the reanalysis represents the seasonality well, and in late summer both the humidity and the biases are of largest magnitude. At the two coastal stations at Hopen and Rjipfjorden, the downscaled reanalysis is too dry whereas it is too humid at the two higher elevation stations. The coarse land mask of the reanalysis and the poor representation of sea ice are most likely the main causes for these biases.

Wind speeds are reproduced reasonably well, including the seasonal cycle (Tab. 1). Biases are within $\pm 1.5 \text{ m s}^{-1}$ with no clear seasonal trend. It is likely that the biases are caused by site specific effects, such as deceleration of air flow in the lee of a topographic obstacle or acceleration due to channelizing through valleys.

Table 1. Meteorological stations used for validation of the downscaled reanalysis and their elevation (second line). N indicates the number of daily averages used in the validation, except for precipitation for which monthly sums have been evaluated. Seasonal biases in meteorological variables (downscaled minus observational averages) at all sites are averaged over each site’s observation period. Shown are air temperature T (K), relative humidity, RH (%), wind speed, WS (ms^{-1}), shortwave radiation, SW , and longwave radiation, LW (both in Wm^{-2}) and precipitation, P (mm). Column headings S and W denote summer (Jun-Aug) and winter (Sep-May), respectively. Positive numbers indicate that the model results are larger than the observations. The second row at each site is the bias between the raw ERA data and the observations.

Location	ΔT			ΔRH			ΔWS			ΔSW			ΔLW		ΔP		
	S	W	N_T	S	W	N_{RH}	S	W	N_{WS}	S	W	N_{rad}	S	W	S	W	N_P
Etonbreen [†]	0.2	1.3	3295	-2.2	-5.4	2738	0.3	0.2	2913	-40	-10	3240	12	-14	-	-	0
369 m asl	<i>1.1</i>	<i>1.9</i>		<i>-2.2</i>	<i>-5.4</i>		<i>0.3</i>	<i>0.2</i>		<i>-37</i>	<i>-9</i>		<i>17</i>	<i>-16</i>	-	-	
Janssonhaugen	-1.1	0.3	910	-	-	0	-1.8	-1.3	945	-	-	0	-	-	-	-	0
270 m asl	<i>-1.0</i>	<i>-0.6</i>		-	-		<i>-1.8</i>	<i>-1.3</i>		-	-		-	-	-	-	
Gruvefjellet	0.2	0.8	2555	3.1	-2.9	2555	0.0	-0.2	2551	-	-	0	-	-	-	-	0
464 m asl	<i>0.7</i>	<i>0.9</i>		<i>3.1</i>	<i>-2.9</i>		<i>0.0</i>	<i>-0.2</i>		-	-		-	-	-	-	
Kapp Heuglin	-0.0	0.7	2099	-	-	0	-0.0	1.0	2112	-	-	0	-	-	-	-	0
18 m asl	<i>0.3</i>	<i>1.0</i>		-	-		<i>-0.0</i>	<i>1.0</i>		-	-		-	-	-	-	
Rijpfjorden	-0.3	0.9	1495	5.7	2.6	1495	0.8	0.9	1304	-	-	0	-	-	-	-	0
10 m asl	<i>0.0</i>	<i>0.6</i>		<i>5.7</i>	<i>2.6</i>		<i>0.8</i>	<i>0.9</i>		-	-		-	-	-	-	
Svalbard Airport	-2.4	-0.4	12777	-	-	0	-1.3	-1.0	12724	-	-	0	-	-	25	26	199*
28 m asl	<i>-2.4</i>	<i>-2.2</i>		-	-		<i>-1.3</i>	<i>-1.0</i>		-	-		-	-	-	-	
Isfjord Radio	-1.6	-1.2	1666	-	-	0	-	-	0	-	-	0	-	-	-	-	0
13 m asl	<i>-1.5</i>	<i>-0.5</i>		-	-		-	-		-	-		-	-	-	-	
Verlegenuken	-0.5	0.0	986	-	-	0	-1.9	-1.7	1700	-	-	0	-	-	-	-	0
8 m asl	<i>-0.3</i>	<i>0.5</i>		-	-		<i>-1.9</i>	<i>-1.7</i>		-	-		-	-	-	-	
Hornsund	-0.2	0.4	4635	-	-	0	-0.0	0.4	4473	-	-	0	-	-	5	19	95*
10 m asl	<i>-0.0</i>	<i>0.8</i>		-	-		<i>-0.0</i>	<i>0.4</i>		-	-		-	-	-	-	
Kvitøya	-0.1	-0.1	740	-	-	0	-0.9	-1.3	702	-	-	0	-	-	-	-	0
10 m asl	<i>0.3</i>	<i>0.6</i>		-	-		<i>-0.9</i>	<i>-1.3</i>		-	-		-	-	-	-	
Holtedahlfonna [†]	1.3	-	317	-	-	0	-0.7	-	265	-	-	0	-	-	-	-	0
688 m asl	<i>3.2</i>	-		-	-		<i>-0.7</i>	-		-	-		-	-	-	-	
Ny-Ålesund	-1.7	-0.0	12666	7.6	3.2	12708	0.7	0.8	12349	-7	4	3652	2	9	23	18	212*
8 m asl	<i>-1.6</i>	<i>-1.2</i>		<i>7.6</i>	<i>3.2</i>		<i>0.7</i>	<i>0.8</i>		<i>-8</i>	<i>3</i>		<i>0</i>	<i>15</i>	-	-	
Hopen	1.9	5.2	13178	-	-	0	0.3	-0.1	13044	-	-	0	-	-	7	10	306*

*: Number of months

†: Station located on glacier.

Table 2. Overview file structure of the Sval_Imp_v1 dataset

Sval_Imp_v1												
Stationary fields	Svalbard_DEM_mask_pcorr.nc											
Variables	T2		P		WS		RH		SW		LW	
Reanalysis	ERA40	ERAint	ERA40	ERAint	ERA40	ERAint	ERA40	ERAint	ERA40	ERAint	ERA40	ERAint
Number of files	540	468	540	468	540	468	540	468	540	468	540	468
Size (GB)	22.8	19.8	33.1	28.8	34.1	29.8	25.6	22.6	21.7	19.0	31.3	27.5

250 4 Dataset structure

The downloadable dataset comprises individual files for each of the variables precipitation, air temperature, relative humidity, windspeed, incident shortwave and downwelling longwave radiation. The records are organized in one file per month for each of the reanalysis periods; ERA-40: September 1957 to August 2002, and ERA-interim: January 1979 to December 2017. Each file contains the discovery metadata, and the complete metadata to locate the stack of fields in space and time. The

255 grid is regular and rectangular in UTM33X projection, the coordinates of which are defined by the vectors X (m easting in UTM33X, 448 elements) and Y (m northing in UTM33X, 548 elements). In geographical coordinates the grid is non-regular and therefore the location of each grid node is defined, rendering Latitude and Longitude (in decimal degrees) each a 448×548 matrix. The timestamp is given in days since 1 January 1900 using a standard Gregorian calendar having 365 days per year, i.e. without accounting for leap years. In addition, there is one file containing the stationary fields, i.e. the surface topography

260 and land-ocean mask. The file format is netCDF according to the CF conventions (<http://cfconventions.org/>), with all required metadata included. The metadata adhere to ISO19115 geospatial metadata standards and the Directory Interchange Format (DIF) requirements of the Global Change Master Directory GCMD (<https://gcmd.nasa.gov/DocumentBuilder/defaultDif10/guide/index.html>), and global attributes comply to the Attribute Convention for Data Discovery ACDD (http://wiki.esipfed.org/index.php/Attribute_Convention_for_Data_Discovery_1-3).

265 Table 2 gives an overview over the number of files and their sizes for the different epochs (ERA-40, ERA-interim) and variables contained in the dataset. The naming convention for the individual files is <EPOCH>_<VAR>_<YYYYMM>.nc, where EPOCH is either "ERA40" or "ERAi", VAR is an abbreviation of the variable of interest (one of "precip", "temp", "RH", "windspeed", "SWi" or "LWi") and YYYYMM identifies year and month, for instance *ERAi_temp_200410.nc* is the name of the file containing air temperature from ERA-interim for October 2004.

270 5 Conclusions

We present a gridded dataset of near-surface, meteorological variables at 1km resolution covering the Svalbard archipelago. The set of variables enables application of energy balance models and comes at a time steps of 6 h. The high-resolution grids are derived from coarse-scale reanalyses ERA-40 for the period 1957-2002 and from ERA-interim for 1979-2017. We describe

the intermediate-complexity downscaling procedure used to generate this dataset. Furthermore, we evaluate the performance of
275 the downscaled data using a suite of different meteorological and glaciological measurements and refer to several applications
of this dataset in different disciplines, all of them requiring longterm coverage at small spatial scales.

6 Data availability

The dataset is openly available from the National e-Infrastructure for Research Data (NIRD) archive at
<https://doi.org/10.11582/2018.00006> and referred to as *Svalbard impact assessment forcing dataset, version 1*, (Schuler, 2018).

280 ERA-40 (Uppala et al., 2005) and ERA-interim (Dee et al., 2011) data were retrieved from the ECMWF Public Datasets
web interface at <https://apps.ecmwf.int/datasets/>.

Weather station data are provided by the Norwegian Meteorological Institute and are available at <https://eklima.met.no>, and
by the University Centre of Svalbard at <https://www.unis.no/resources/weather-stations/>. Radiation from the BRSN-station in
Ny-Ålesund are provided by Maturilli et al. (2014).

285 *Author contributions.* TVS retrieved reanalysis data, developed code and performed the downscaling for precipitation, air temperature,
RH and windspeed; TIØ developed code and performed downscaling of radiation components. Evaluation performance was conducted
collaboratively. TVS was responsible for formatting, providing metadata and archiving the dataset; he led writing of the manuscript, TIØ
contributed to writing.

Competing interests. There are no competing interests present.

290 *Acknowledgements.* We greatly acknowledge the European Centre for Medium-range Weather Forecast, ECMWF for access to their reanal-
ysis products ERA-40 and ERA-interim. The Norwegian Meteorological Institute has been pioneering and is exemplary for open access
to historical data, especially through their *eklima*-service. Field work on Austfonna has been supported by NFR through IPY-Glaciodyn,
EU-ice2sea and ESA-CRYOVEX, along with KD scholarship for TIØ and mobility funds from SVALI.

References

- 295 Aalstad, K., Westermann, S., Schuler, T. V., Boike, J., and Bertino, L.: Ensemble-based assimilation of fractional snow-covered area satellite retrievals to estimate the snow distribution at Arctic sites, *The Cryosphere*, 12, 247–270, <https://doi.org/10.5194/tc-12-247-2018>, 2018.
- Aas, K. S., Dunse, T., Collier, E., Schuler, T. V., Berntsen, T. K., Kohler, J., and Luks, B.: The climatic mass balance of Svalbard glaciers: a 10-year simulation with a coupled atmosphere–glacier mass balance model, *The Cryosphere*, 10, 1089–1104, <https://doi.org/10.5194/tc-10-1089-2016>, 2016.
- 300 Barstad, I. and Smith, R. B.: Evaluation of an Orographic Precipitation Model, *Journal of Hydrometeorology*, 6, 85–99, <https://doi.org/10.1175/JHM-404.1>, 2005.
- Bogren, W. S., Burkhart, J. F., and Kylling, A.: Tilt error in cryospheric surface radiation measurements at high latitudes: a model study, *The Cryosphere*, 10, 613–622, <https://doi.org/10.5194/tc-10-613-2016>, 2016.
- Botnen, S., Mundra, S., Kauserud, H., and Eidesen, P. B.: Glacier retreat in the High Arctic: Opportunity or threat for ectomycorrhizal fungi?,
305 in prep.
- Crochet, P., Jóhannesson, T., Jónsson, T., Sigurðsson, O., Björnsson, H., Pálsson, F., and Barstad, I.: Estimating the Spatial Distribution of Precipitation in Iceland Using a Linear Model of Orographic Precipitation, *Journal of Hydrometeorology*, 8, 1285–1306, <https://doi.org/10.1175/2007JHM795.1>, 2007.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer,
310 P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- 315 Dunse, T., Schuler, T. V., Hagen, J. O., Eiken, T., Brandt, O., and Høgda, K. A.: Recent fluctuations in the extent of the firn area of Austfonna, Svalbard, inferred from GPR, *Annals of Glaciology*, 50, 155–162, <https://doi.org/10.3189/172756409787769780>, 2009.
- Ehret, U., Zehe, E., Wulfmeyer, V., Warrach-Sagi, K., and Liebert, J.: HESS Opinions "Should we apply bias correction to global and regional climate model data?", *Hydrology and Earth System Sciences*, 16, 3391–3404, <https://doi.org/10.5194/hess-16-3391-2012>, 2012.
- Fiddes, J. and Gruber, S.: TopoSCALE v.1.0: downscaling gridded climate data in complex terrain, *Geoscientific Model Development*, 7,
320 387–405, <https://doi.org/10.5194/gmd-7-387-2014>, 2014.
- Førland, E. J. and Hanssen-Bauer, I.: Increased Precipitation in the Norwegian Arctic: True or False?, *Climatic Change*, 46, 485–509, <https://doi.org/10.1023/A:1005613304674>, 2000.
- Grabiec, M., J., L., P., G., and J., J.: Distribution of snow accumulation on some glaciers of Spitsbergen, *Polish Polar Research*, 27, 309–326, 2006.
- 325 Gutmann, E., Barstad, I., Clark, M., Arnold, J., and Rasmussen, R.: The Intermediate Complexity Atmospheric Research Model (ICAR), *Journal of Hydrometeorology*, 17, 957–973, <https://doi.org/10.1175/JHM-D-15-0155.1>, 2016.
- Hanssen-Bauer, I., Førland, E., Hisdal, H., S.Mayer, Sandø, A., Sorteberg, A., Adakudlu, M., Andresen, J., Bakke, J., Beldring, S., Benestad, R., van der Bilt, W., Bogen, J., Borstad, C., Breili, K., Breivik, Ø., Børshem, K., Christiansen, H., Dobler, A., Engeset, R., Frauenfelder, R., Gerland, S., Gjeltén, H., Gundersen, J., Isaksen, K., Jaedicke, C., Kierulf, H., Kohler, J., Li, H., Lutz, J., Melvold, K., Mezghani,
330 A., Nilsen, F., Nilsen, I., Nilsen, J., Pavlova, O., Ravndal, O., Risebrobakken, B., Saloranta, T., Sandven, S., Schuler, T., Simpson, M.,

- Skogen, M., Smedsrud, L., Sund, M., Vikhamar-Schuler, D., Westermann, S., and Wong, W., eds.: Climate in Svalbard 2100 - a knowledge base for climate adaptation, vol. 1/2019, Norwegian Environment Agency (Miljødirektoratet), Norwegian Centre for Climate Services, https://cms.met.no/site/2/klimaservicesenteret/climate-in-svalbard-2100/_attachment/14428?_ts=169fd13ff23, 2019.
- Jarosch, A. H., Anslow, F. S., and Clarke, G. K. C.: High-resolution precipitation and temperature downscaling for glacier models, *Climate Dynamics*, 38, 391 – 409, <https://doi.org/10.1007/s00382-010-0949-1>, 2012.
- Kumar, L., Skidmore, A., and Knowles, E.: Modelling topographic variation in solar radiation in a GIS environment, *International Journal of Geographical Information Science*, 11, 475–497, <https://doi.org/10.1080/136588197242266>, 1997.
- Liston, G. E. and Elder, K.: A Meteorological Distribution System for High-Resolution Terrestrial Modeling (MicroMet), *Journal of Hydrometeorology*, 7, 217–234, <https://doi.org/10.1175/JHM486.1>, 2006.
- 335 Maraun, D.: Bias Correction, Quantile Mapping, and Downscaling: Revisiting the Inflation Issue, *Journal of Climate*, 26, 2137–2143, <https://doi.org/10.1175/JCLI-D-12-00821.1>, 2013.
- Marshall, S. J., Sharp, M. J., Burgess, D. O., and Anslow, F. S.: Near-surface-temperature lapse rates on the Prince of Wales Icefield, Ellesmere Island, Canada: implications for regional downscaling of temperature, *International Journal of Climatology*, 27, 385–398, <https://doi.org/10.1002/joc.1396>, 2007.
- 345 Maturilli, M., Herber, A., and König-Langlo, G.: Climatology and time series of surface meteorology in Ny-Ålesund, Svalbard, *Earth System Science Data*, 5, 155–163, <https://doi.org/10.5194/essd-5-155-2013>, 2013.
- Maturilli, M., Herber, A., and König-Langlo, G.: Basic and other measurements of radiation from the Baseline Surface Radiation Network (BSRN) Station Ny-Ålesund in the years 1992 to 2013, reference list of 253 datasets, <https://doi.org/10.1594/PANGAEA.150000>, <https://doi.org/10.1594/PANGAEA.150000>, supplement to: Maturilli, M et al. (2014): Surface radiation climatology for Ny-Ålesund, Svalbard (78.9° N), basic observations for trend detection. *Theoretical and Applied Climatology*, 120(1-2), 331-339, <https://doi.org/10.1007/s00704-014-1173-4>, 2014.
- 350 Østby, T. I., Schuler, T. V., Hagen, J. O., Hock, R., Kohler, J., and Reijmer, C. H.: Diagnosing the decline in climatic mass balance of glaciers in Svalbard over 1957–2014, *The Cryosphere*, 11, 191–215, <https://doi.org/10.5194/tc-11-191-2017>, 2017.
- Pramanik, A., Van Pelt, W., Kohler, J., and Schuler, T.: Simulating climatic mass balance, seasonal snow development and associated fresh-water runoff in the Kongsfjord basin, Svalbard (1980-2016), *Journal of Glaciology*, 64, 943–956, <https://doi.org/10.1017/jog.2018.80>, 2018.
- 355 Ratti, C.: Urban analysis for environmental prediction, Ph.D. thesis, University of Cambridge Department of Architecture, 2001.
- Reda, I. and Andreas, A.: Solar position algorithm for solar radiation applications, *Solar Energy*, 76, 577 – 589, <https://doi.org/https://doi.org/10.1016/j.solener.2003.12.003>, 2004.
- 360 Roth, A., Hock, R., Schuler, T. V., Bieniek, P. A., Pelto, M., and Aschwanden, A.: Modeling Winter Precipitation Over the Juneau Icefield, Alaska, Using a Linear Model of Orographic Precipitation, *Frontiers in Earth Science*, 6, 20, <https://doi.org/10.3389/feart.2018.00020>, 2018.
- Sand, K., Winther, J.-G., Maréchal, D., Bruland, O., and Melvold, K.: Regional Variations of Snow Accumulation on Spitsbergen, Svalbard, 1997-99, *Hydrology Research*, 34, 17–32, <https://doi.org/10.2166/nh.2003.0026>, <https://doi.org/10.2166/nh.2003.0026>, 2003.
- 365 Schuler, T.: Svalbard impact assessment forcing dataset, version 1, <https://doi.org/10.11582/2018.00006>, 2018.
- Schuler, T., Dunse, T., Østby, T., and Hagen, J.: Meteorological conditions on an Arctic ice cap—8 years of automatic weather station data from Austfonna, Svalbard, *International Journal of Climatology*, 34, 2047–2058, <https://doi.org/10.1002/joc.3821>, 2014.

- Schuler, T. V., Loe, E., Taurisano, A., Eiken, T., Hagen, J. O., and Kohler, J.: Calibrating a surface mass-balance model for Austfonna ice cap, Svalbard, *Annals of Glaciology*, 46, 241–248, <https://doi.org/10.3189/172756407782871783>, 2007.
- 370 Schuler, T. V., Crochet, P., Hock, R., Jackson, M., Barstad, I., and Jóhannesson, T.: Distribution of snow accumulation on the Svartisen ice cap, Norway, assessed by a model of orographic precipitation, *Hydrological Processes*, 22, 3998–4008, <https://doi.org/10.1002/hyp.7073>, 2008.
- Sinclair, M. R.: A Diagnostic Model for Estimating Orographic Precipitation, *Journal of Applied Meteorology*, 33, 1163–1175, [https://doi.org/10.1175/1520-0450\(1994\)033<1163:ADMFE0>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<1163:ADMFE0>2.0.CO;2), 1994.
- 375 Smith, R. B. and Barstad, I.: A Linear Theory of Orographic Precipitation, *Journal of the Atmospheric Sciences*, 61, 1377–1391, [https://doi.org/10.1175/1520-0469\(2004\)061<1377:ALTOOP>2.0.CO;2](https://doi.org/10.1175/1520-0469(2004)061<1377:ALTOOP>2.0.CO;2), 2004.
- Stone, P. H. and Carlson, J. H.: Atmospheric Lapse Rate Regimes and Their Parameterization, *Journal of Atmospheric Sciences*, 36, 415–423, 1979.
- Taurisano, A., Schuler, T. V., Hagen, J. O., Eiken, T., Loe, E., Melvold, K., and Kohler, J.: The distribution of snow accumulation
380 across the Austfonna ice cap, Svalbard: direct measurements and modelling, *Polar Research*, 26, 7–13, <https://doi.org/10.1111/j.1751-8369.2007.00004.x>, 2007.
- Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D. C., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Berg, L. V. D., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E.,
385 Hoskins, B. J., Isaksen, L., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Quarterly Journal of the Royal Meteorological Society*, 131, 2961–3012, <https://doi.org/10.1256/qj.04.176>, 2005.
- Vikhamar-Schuler, D., Førland, E., Lutz, J., and Gjelten, H.: Evaluation of downscaled reanalysis and observations for Svalbard: Background report for Climate in Svalbard 2100, Tech. rep., Norwegian Centre for Climate Services, 2019.
- 390 Wolff, M. A., Isaksen, K., Petersen-Øverleir, A., Ødemark, K., Reitan, T., and Brækkan, R.: Derivation of a new continuous adjustment function for correcting wind-induced loss of solid precipitation: results of a Norwegian field study, *Hydrology and Earth System Sciences*, 19, 951–967, <https://doi.org/10.5194/hess-19-951-2015>, 2015.