

First of all, we are happy, that the comments on the manuscript are generally positive. As some of the reviewers wrote, the compilation and quality control of a dataset like the one described here, really was a lot of work. We want to thank the four anonymous reviewers and Jianxi Huang, who did a short comment, for their statements. Here, we respond to all comments given and explain which changes we applied to the manuscript and dataset. Reviewer comments are black, the authors' response is green and changes to the text are blue.

RC1

I appreciate that the editor has offered this opportunity to comment. In this manuscript, Reichenau et al. present a comprehensive dataset of vegetation states, fluxes of matter and energy, weather, agricultural management, and soil properties from four intensively monitored crop sites in Western Germany. The authors are to be commended for undertaking an ambitious project and providing types of data useful to the agriculture and biogeochemical modeling communities. In this manuscript, all kinds of information regarding the development of the dataset are demonstrated. I especially appreciate that the authors specified the detailed uncertainty and quality assurance of each dataset, which is not usually seen in many published datasets. I think this dataset and paper are worth publishing in ESSD. However, there are some major or minor information needs to be clarified before its acceptance and publication.

Thank you for the positive feedback. It was indeed a lot of work to compile and quality-check everything.

1. These sites are/were intensively agricultural sites. As an ecosystem modeler, considering the purpose of initializing crop/ecosystem models, the pre-management vegetation and soil conditions (e.g., soil C and N content, potential vegetation type and its states) should be used to initialize models. However, the data regarding vegetation and soil properties provided in this paper are quite contemporary, say time coverage is from 2007 to 2017. The initial conditions of vegetation and soils were missed in these sites. Is it possible to add some of the background information to make the dataset more complete?

The whole area of investigation has been inhabited and agriculturally used for more than 2000 years. Because of the very long period of agricultural land-use, it can be assumed, that there are no persisting effects of the conversion to agricultural area. Therefore, the soil C- and N-contents shown represent the current states as influenced by current management with the usual variations due to plant growth (seasonality), fertilization, and tillage. The data given in the paper is all measured data we know about.

In the context of a comment in RC4 (see below), we added information on the duration of agricultural use of the sampled fields.

2. Under section 5, please specify the beginning and ending date of the 15 time series. I went over the spreadsheet and found the temporal coverages of these time series are not even. Still, it is amazing to see the 30-min interval data!

For the readers' convenience, we added the periods of the EC timeseries to Table 5. Furthermore, the datafiles were changed. They now do always start at 00:00 of the first day and end at 23:59 of the last day.

3. Under section 7, four different gap-filling methods were actually used. However, the reasoning of each method every time it is used at different sites is not explicit enough. Under what circumstances each method was exactly used? And why?

In the course of the project, different methods were applied to fill the gaps in the meteorological data. There were two project-wide efforts for meteo gap-filling. The latest used the EOF method. This is the method with the best evaluation. Therefore, this data was used where available. It could not

be applied earlier than 2011 since the number of available stations was too small then. In the earlier years, first the insertion method and later the more sophisticated regression method 2 was applied, which reflects a methodological progress in the project. Gap-filling in Hürtgenwald was not done with the EOF method, since this station does not belong to the centrally maintained stations of the project.

To make this clear to the readers, short explanations were added in section 7.1.1.

Minor comments:

The current organization of the paragraphs is not good enough. There is no first-line indentation where it is needed. Bullet points also do not have indentations. Some sloppy spelling and wrongly used punctuation and marks. Also, no line number, which makes it hard to specify comments.

We are sorry for the inconvenience experienced by reviewer 1. In general, we used the template provided by the Journal. However, we have to admit, that we were not consequent with the formatting of enumerations and bullet lists. This will be corrected during the typesetting process of the revised manuscript (if accepted for publication) as will be the indentation. We have no explanation why the line numbers were missing. They are present in the pdf-document on the journal's website.

We will check the revised manuscript for compliance with the journal's typesetting guidelines.

Section 2.1 Line 11: within the parenthesis, % was used while degree was used in other places. Be consistent.

We thank the reviewer for the thorough review!

We changed the units accordingly.

Line 19: data is a plural term!

Here, we would like to refer to <http://dataabinitio.com/?p=497> and point to the third paragraph. The author points out that, due to evolution of language, data nowadays can be used as singular. Since there are about three times more google hits with the singular form, we kept this.

Section 2.2 Line 25, degree sign is used here. Be consistent.

This was unified for comment Section 2.1 line 11 above.

Section 2.3 Last line: what is the temporal range for the mean annual temperature of 9.7 degrees in this site?

The temporal range for this data is unknown. It was taken from Schulz (2004) since this was the only literature with data on Merzenhausen climate. Unfortunately, there is no way to ask Elke Schulz for details on the data, because she retired recently and has no public contact address.

We added the citation to the sentence on climate.

Section 3 Line 10: data is a plural term!

See above.

Line 15: remove "and" before "can thus be"

Sentence was rephrased

Section 4.1 Line 4: remove "given"

done

Section 5.1 Line 20: please use subscript for H<sub>2</sub>O and CO<sub>2</sub>.

done

Section 7.1.1 Line 30: please specify the years for the sites for which the EOF method was applied.

Already done in context of major comment 3.

Page 16 Line 6: please correct the “ and ” signs.

done

RC2

### Summary and general comments

Reichenau et al. presents a comprehensive dataset related to vegetation states, energy and water fluxes, agricultural management, and soil properties from four agricultural sites in Western Germany. I appreciate the efforts the authors made in detailing all these datasets; however, there are also some confusions and issues that the authors should clarify in further revisions.

Thank you for the positive feedback.

- (1) Scientific relevance: As the selected sites are from the same area and are all agricultural-based sites, I would suggest the authors further sharpen their research motivation by stating how these datasets can facilitate understanding agricultural ecosystems and then generalize to the general ecosystem. Right now, it seems we can use these datasets to understand all aspects of ecological settings as the introduction specifies. In addition, as flux datasets from eddy covariance flux towers are also available, the authors need to point out what are the new items we can find in this dataset. What is missing before and what would be new data that are important to further gain insights of ecological systems? In addition, the provision of these data from only this area also raises the question whether these data are representative to investigate issues such as impacts of climate change on agricultural ecosystems? The authors may also want to clarify.

We respond to this comment and to a comment in RC4 where the reviewer states “the importance of the current dataset is not clear”.

The intention of the compilation and publication of this dataset is to provide coherent and comprehensive data on agricultural land. It is intended to provide modelers and remote sensing scientists a set of relevant types of data collected in a 12-year project. Parameterized models or remote sensing algorithms resulting from the usage of the data can then be used for whatever purpose the model or algorithm is applied to. This encompasses everything that requires a good representation of vegetation, be it studies on hydrology, balances of energy or mass, food production, etc. A generalization to non-agricultural ecosystems was not in our scope. Furthermore, the data is not assumed to be representative for agricultural ecosystems in general, since these show a large variability in terms of crops, management, soil, and climate. Therefore, detailed data from agricultural ecosystems differing in the named properties are required for large-scale analyses, e.g. on the continental or national scale. This illustrates the need for this kind of datasets from different places in order to develop realistic information on the regional differences of states and fluxes. By this means, the whole dataset is the reviewer’s “new item”.

We added a statement that this kind of data is required from different agricultural regions in the introduction.

Since it is usually hard to find fitting data for the same sites, we wanted to provide a convenient overall collection of well-documented quality-controlled data, which is ready to use. Let’s take the example of a modeler interested in simulating fluxes in a highly fragmented agricultural area with small fields. In order to accomplish this, flux data is required together with data on vegetation, soil, weather, and management, not only for the field with flux measurements but also for surrounding fields for extrapolation purposes. This kind of data is not available from the Fluxnet or Ameriflux networks where ancillary data is (if at all) only offered for the flux-measurement site itself. In this dataset, such data is available. Especially for the Selhausen site, the detailed soil map is a unique feature. Furthermore, this dataset includes six unpublished flux time-series from the site Merken. These are unpublished measurements from three nearby fields measured simultaneously at two heights, a setup which is rarely available.

We added a statement on the previously unpublished two-height flux data for the Merken site and on the differences in data availability between Fluxnet/Ameriflux and this dataset in the introduction and mentioned the former in the abstract.

- (2) Datasets: there are quite many missing data shown in Excel as NA. I am not sure whether it makes the whole data series. The authors also need to document why these missing data are there and if gap filling algorithms were used (at the least the manuscript provides gap filling algorithms) and/or needs to be used.

It is true that there are many data-points marked unavailable. On the one hand, this is due to maintaining a consistent structure of the data-tables: (1) If a certain measured variable is available for one site, the corresponding column in the tables is there for all sites. (2) For flux data, we used preexisting file formats, which define columns for magnitudes that were not measured in our case. Both cases cause columns totally filled with NA. On the other hand, some data had to be removed from the dataset due to quality control procedures.

We added this clarification in a new section 3.1.

Gap-filling was solely used for meteorological data because this is required to run a simulation model. Since flux data is sometimes provided with gaps filled, we added a statement in section 5.1 that no gap-filling was applied. We did not add such statements to the sections on vegetation data, because we do not want to start an unnecessary discussion on whether the indirect calculation of a biomass value should be called gap-filling. We consider this unnecessary, since the methods applied are described in detail in section 4.2. In case of soil data, we abstain from adding a statement on gap filling, since the method used by Brogi et al. (2019) to define soil units include an interpolation process. Because it is far from the topic of this paper, and because the methods are described in detail, we do not want to discuss this here.

Since flux data is sometimes provided with gaps filled, we added a statement in section 5.1 that no gap-filling was applied

In addition, data documents from all these sites are not consistent, for example, EC folder in all the sites except Huertgenwald. Management data from the Merken site only have a couple of years data. Why some years of data are missing? Some sites do not have meteo.csv?

Overall, the data are not as detailed as those described in the manuscript.

In this comment, the reviewer states that the documents for the sites were not consistent. From our point of view, all the information the reviewer is missing is given in the text. Since there were no flux measurements in Huertgenwald, there is no folder for this data. The same applies to meteorological data for Merken as described starting in line 28 in section 7.1. The availability of management data is given per data-type in Table 3.

We extended the statements on data availability in the sections on flux data and crop management data.

It is also hard to understand some of the excel documents since no metadata are provided to explain the variables? For example, what is u, v, w in the fluxes\_MEF01\_EC\_001\_2011?

Regarding the variables, the metadata in question is given in Tables A2 and A3, which are referred to in the text in section 5.4.

- (3) While the authors attempt to detail the datasets, it is necessary for the authors to provide at least several application examples that how these datasets can be used. Particularly, as the authors trying to understand “patterns in soil-vegetation-atmosphere-systems”, it would be helpful to see some research results from this dataset. This would also help clarify how these datasets can be used even if there are missing data. I do not expect complicated application examples, but generally, for example vegetation states and temperature/precipitation can be used to understand climate impacts on agricultural ecosystems.

The paper includes several references to publications where the data has been used. These papers illustrate several applications of the data, e.g.

LAI: Brogi et al. (2020), Reichenau et al. (2017), Ney and Graf (2018), Ahrends et al. (2014)

biomass and phenology: Schmidt et al. (2012), Korres et al. (2013)

management: Schmidt et al. (2012), Korres et al. (2013)

weather: Schmidt et al. (2012), Korres et al. (2013), Sakai (2016)

fluxes: Klosterhalfen et al. (2017), Schmidt et al. (2012), Ney and Graf (2018), Ahrends et al. (2014), Wienecke et al. (2018), Eder et al (2015)

soil: Borneman et al. (2011), Korres et al. (2013), Jakobi et al. (2020), Meyer et al. (2017)

For complete references please see the manuscript.

References to papers where the data was used were added to the end of each data-source section.

RC3

The authors generated a comprehensive dataset at four agricultural sites for the development and validation of hydro-ecological land-surface models, and as well as the remote sensing products. Thank the authors for the efforts on the data collection. As the authors stated that one of the goals of this dataset is for the validation of the remote sensing products, there are still some places that need the authors' clarification and consideration. I recommend a minor revision before publication.

Thank you for the comment from a remote sensing point of view.

Section 4, P5 L14: The leaf area index was measured by the destructive approach at a very small sampling area, e.g., 40 x 40 cm. I fully understand that this was due to the limitation of labor, while the spatial representativeness of the measurements can make the direct validation of the remote sensing products (10 m\_ 1km) to be challenging. The inhomogeneity of the vegetation states over the study area may need to be evaluated by the UAV data or very high spatial resolution data if available, especially at the same periods of the measurements. The corresponding high-resolution data can be very important for bridging the gap between the point-scale field measurements and satellite-level remote sensing products, as the two websites below. Even the information about the availability of the high-resolution data with good quality can also be very helpful for the users.  
[http://w3.avignon.inra.fr/valeri/fic\\_htm/database/main.php](http://w3.avignon.inra.fr/valeri/fic_htm/database/main.php) ,  
<http://calvalportal.ceos.org/web/olive/site-description>

As the reviewer states correctly, the limited area sampled is due to the limited work capacity together with the aim to sample multiple points per field or multiple fields. Furthermore, farmers would not permit to harvest the area of several 10 m pixels (100 m<sup>2</sup> each). However, especially in the data from the Selhausen site, several sampling points fall within the area of one remote sensing pixel thus enabling an estimate of the variability within the pixel.

In the context of remote sensing, the LAI required for calibration/evaluation is the average LAI of a pixel's area. Based on the above this usually cannot be measured destructively. Unfortunately, no data was collected via UAV. However, a way to connect field measurements to lower-resolution remote sensing products may be made via high resolution remote sensing. E.g. Brogi et al. (2020) calibrated the algorithm of Ali et al. (2015) based on NDVI derived from 5 m resolution RapidEye level 3A data and LAI data from the dataset presented in this manuscript. Reichenau et al. (2016) showed that realistic statistical distribution of LAI over a larger area could be derived without calibration. However, in that case ground truth is required to prove this. The resulting 5 m resolution LAI data can then be used to bridge the gap to lower-resolution datasets.

The example was added in the section on vegetation measurements.

Besides, the temporal frequency of the LAI measurements collected from 2007 to 2017 may need to be clearly explained in the manuscript for the readers. The publicly available LAI measurements at the seasonal cycle are rare and valuable for the validation of the current remote sensing products in time series.

The LAI data has been collected with frequencies from one to three weeks during the growing season. In the years from 2015 to 2017, this was done on overflight days of Sentinel 1 and Radarsat 2. There are also entries for dates without vegetation. These were included to have a zero-point for the calibration of remote sensing algorithms.

This information has been added to the LAI section. A statement on data frequency and measurements on satellite overflight days was added to the abstract.

For the linkage of the vegetation and soil properties with remote sensing data, in addition to the canopy structure parameters such as LAI, not quite sure whether there are also leaf and soil spectrum, leaf chlorophyll and water content, and photos of the vegetation canopy at different

growing stages available. If yes, these data would be very useful for the radiation transfer modeling and analysis over the agricultural ecosystem.

Unfortunately, leaf and soil spectra were not obtained. The photos of the canopy taken during the harvest of plants on the fields were taken for documentation purpose only. There was no standardized protocol for this. Therefore, it remains unclear whether they can be applied in the context of remote sensing applications. The vegetation water content, however, can be calculated from the difference of fresh and dry weight of the biomass.



RC4

I appreciate the authors' efforts in compiling this dataset. As a modeler, I understand the importance of such a dataset for model parameterization and validation. Data collection and compiling are usually two of the most time-consuming processes. It's good to see this dataset been organized following a consistent protocol.

Thank you for this positive feedback.

However, the importance of the current dataset is not clear. Such data (e.g. flux, managements) can be found from different sources. For example, Ameriflux (<https://ameriflux.lbl.gov/>) provides multi-year, quality-controlled data collected from hundreds of eddy-flux sites worldwide, and the management information for the cropland sites can be obtained from the website and the related publications. The authors did provide a description of the uniqueness of the dataset, but it sounds the data has limited impacts on at local area.

Please see our response to comment (1) of reviewer 2 (RC2), where we included the response on this comment.

Honestly, I am also not very satisfied with many missing data (NAs) in this dataset, especially in the management information category. The dataset barely provides a three-year continuous management records. I understand that this is labor-intensive work, but the current dataset is relatively short-term coverage.

Regarding the missing management data, this is not due to the time needed to collect it. It is simply due to farmers not reporting their data. Since that was, of course, a voluntary contribution of the farmers, there is no way to get that data. Please also refer to our comments on missing data in the response to reviewer 2 (RC2).

We added section 3.1 as a combined response to this comment and RC2.

Are the authors planning to continue the measurements and regularly update the dataset?

Unfortunately, the project has ended and there are no follow-up projects funding the continuation of the sampling. Solely field F01 in Selhausen is undergoing ongoing sampling since it has become an ICOS site.

Another concern is the lack of land use history information. For example, land use change plays a pivotal role in altering soil property. How long have the sites been converted to cropland? Generally, conversion from natural vegetation to cropland triggers rapid decomposition of soil organic matters, which may last for decades. Have the lagged impacts from land use change gone? This background information is essential, which should be provided before the data can be used in modeling.

The area of the Rur catchment is ancient agricultural land. There has been agriculture in the region in the time of the Romans or even earlier. Close to the field in Merzenhausen, a grave from the Bronze Age has been found. Therefore, it is justified to assume that effects of conversion to arable land do not persist.

We added general information on this in the section 2. More detailed statements were added in the subsections of the respective sites.

SC1

This manuscript provides a comprehensive dataset collected at four agricultural sites within the Rur catchment in Western Germany. The dataset includes various variables, e.g., vegetation states, carbon and energy fluxes, meteorological variables, agricultural management, and soil properties. The topic is an important one and the manuscript is well written.

We thank Jianxi Huang for these extra comments.

#### Major Comments

1. The authors have provided detailed description of the development and information about the dataset, e.g., the quality flags and the uncertainty. However, a discussion/conclusion section is missing, where the main advantages/disadvantages or the cautions in using the dataset of the provided dataset should be highlighted to the readers. I believe that highlight would greatly benefit the potential users of the dataset.

When writing the manuscript, we decided not to add a discussion section. We assume that researchers who use a dataset like the one described in this manuscript are familiar with the requirements and caveats in scientific processes like model validation or the calibration of models and remote sensing algorithms. Since the manuscript is already quite long, we think that this is not the right place for a discussion of basic scientific processes.

2. The manuscript seems incomplete with reference section missing.

In the pdf document on the ESSD website, there is a references section beginning on page 21.

Some basic methods or theory used in the data development process should be cited, e.g., the barometric formula used in Page 16 Line 30.

In our opinion, we cited all resources underlying the process of data collection, evaluation, analysis, and quality control. The barometric formula is derived from basic physical laws. It has been widely used for a long time and thus has no specific reference.

#### Minor Comments

1. Page 2 Line 19. Why use capital letters in “Monitoring, Modelling and Data”?

The capital letter originated from the project’s name. Nevertheless, in this place they should be lowercase.

Corrected

2. Page 2 Line 21. “TERENO (Terrestrial Environmental Observatories)” the abbreviation should be in the bracket?

We think it is a question of style either to give the explanation or the acronym in brackets.

3. Page 3 Line 12. What is “WRB”? please explain.

Done

4. Page 3 Line 12. “carbon (NEE), water (LE), and energy (H)”, please explain the meaning of the abbreviations first before using them, e.g., LE the latent heat flux, H the sensible heat flux.

This comment refers to page 10 line 17. We used the common symbols for the fluxes in capital letters. However, the symbols can also be interpreted as abbreviations.

Additional explanations were added.

5. Page 11 Line 21. TYPO. “CO2” should be “CO<sub>2</sub>”. Please also check throughout the manuscript.

Corrected

6. Page 15 Line 11. TYPO. "On short timescales"?

Removed

7. Page 16 Line 6. TYPO. ""bad", or "suspicious""

Already corrected based on a comment by reviewer 1

Done

8. Page 16 Line 30. Reference should be cited for the barometric formula?

Please see our response to the major comment 1 above.

# A comprehensive dataset of vegetation states, fluxes of matter and energy, weather, agricultural management, and soil properties from intensively monitored crop sites in Western Germany

5 Tim G. Reichenau<sup>1</sup>, Wolfgang Korres<sup>1</sup>, Marius Schmidt<sup>2</sup>, Alexander Graf<sup>2</sup>, Gerhard Welp<sup>3</sup>, Nele Meyer<sup>3,4</sup>, Anja Stadler<sup>5</sup>, Cosimo Brogi<sup>2</sup>, Karl Schneider<sup>1</sup>

<sup>1</sup>Institute of Geography, University of Cologne, Germany

<sup>2</sup>Institute of Bio- and Geosciences 3: Agrosphere (IBG-3), Jülich Research Centre, Jülich, Germany

<sup>3</sup>Institute of Crop Science and Resource Conservation (INRES), Soil Science and Soil Ecology, University of Bonn, Bonn, Germany

10 <sup>4</sup>Department of Forest Sciences, University of Helsinki, Helsinki, Finland

<sup>5</sup>Institute of Crop Science and Resource Conservation (INRES), Crop Science, University of Bonn, Bonn, Germany

*Correspondence to:* Tim Reichenau (tim.reichenau@uni-koeln.de)

**Abstract.** The development and validation of hydroecological land-surface models to simulate agricultural areas requires extensive data on weather, soil properties, agricultural management, and vegetation states and fluxes. However, this comprehensive data is rarely available since measurement, quality control, documentation and compilation of the different data types is costly in terms of time and money. Here, we present a comprehensive dataset, which was collected at four agricultural sites within the Rur catchment in Western Germany in the frame of the Transregional Collaborative Research Centre 32 “Patterns in Soil-Vegetation-Atmosphere-Systems: Monitoring, Modelling and Data Assimilation” (TR32). Vegetation-related data comprises fresh and dry biomass (green and brown, predominantly per organ), plant height, green and brown leaf area index, phenological development state, nitrogen and carbon content (overall > 17000 entries), and masses of harvest residues and regrowth of vegetation after harvest or before planting of the main crop (> 250 entries). Vegetation data including LAI was collected in frequencies of one to three weeks, in the years 2015 until 2017 mostly during overflights of the Sentinel 1 and Radarsat 2 satellites. Fluxes of carbon, energy, and water (> 180000 half-hourly records) for a variety of agricultural plants. In addition, masses of harvest residues and regrowth of vegetation after harvest or before planting of the main crop are included (> 250 entries). fluxes of carbon, energy, and water (> 180000 half-hourly records) measured using the eddy covariance technique for a variety of agricultural plants are included. Three flux timeseries have simultaneous data from two different heights. Data on agricultural management includes sowing and harvest dates, and information on cultivation, fertilization and agrochemicals (27 management periods). The dataset also includes gap-filled weather data (> 200000 hourly records) and soil parameters (particle size distributions, carbon and nitrogen contents; > 800 records). This data can also be useful for development and validation of remote sensing products. The dataset (Reichenau et al., [2019/2020](#)) is hosted at the TR32 database (<https://www.tr32db.uni-koeln.de/data.php?dataID=18869>) and has the DOI [10.5880/TR32DB.39](https://doi.org/10.5880/TR32DB.39).

## 1 Introduction

System states and processes at the land surface are of major interest in the context of climate change, hydrological and biogeochemical research. In order to understand the processes in their spatial context and to provide information for larger areas, remote sensing and simulations are heavily applied methods. In this context, it is crucial to understand the fluxes mediated by the vegetation at the land surface. Dependencies of processes on vegetation states and properties and on environmental conditions are often investigated using models, while their spatial variability is inferred using remote sensing techniques. In this context, well-documented and quality controlled, comprehensive field measurements of vegetation-related variables are essential for research tasks like model development, calibration, parameterization, and validation or as ground truth for remote sensing products. These variables include biomass per organ differentiated between living (green) and senescent or diseased (brown) material, leaf area index (LAI) and the phenological state of the vegetation. For a simulation, additional information on site conditions such as vegetation composition, soil texture, weather, and, in the case of agro-ecosystem models, agricultural management is required (Kersebaum et al., 2015). However, there is a scarcity of such datasets (Jones et al., 2017). This is of special relevance, since especially the crops grown and their properties differ between regions due to different soils and climate. Thus, detailed data on the named variables is required for different agricultural regions. With the publication of the data described in this article, we contribute a new coherent dataset on agro-ecosystems that includes all of the mentioned variables. The data was collected on conventionally managed fields cultivated by ordinary farmers working at the sites for many years. Thus, it represents conditions and usual practices representative for the intensively used agricultural region to the west of Cologne in Germany. The dataset comprises data from four sites. It consists of almost 1500 records of vegetation parameters and more than 200000 entries of weather data complemented by 15 flux datasets (eddy covariance), management information for 27 management periods, and soil information for all four sites. In contrast to the ancillary data often available with flux data from the Fluxnet or Ameriflux databases, vegetation and soil data in this dataset are also available for other fields in the region, enabling extrapolation of field-scale results to the region. Since collecting field-data is very time consuming and expensive there are not many datasets of this size.

The data was collected in the Rur catchment, located at the Belgium-German-Netherlands border within the frame of the Transregional Collaborative Research Centre 32 (TR32, Vereecken et al., 2010, Simmer et al., 2015) “Patterns in Soil-Vegetation-Atmosphere-Systems” funded by the German national science foundation (Deutsche Forschungsgemeinschaft, DFG). TR32 ran from 2007 until 2018. The project’s main focus was on the combination of Monitoring, Modelling and Data Assimilation to assess the role of patterns in Soil-Vegetation-Atmosphere-Systems across scales. The monitoring efforts of TR32 were accompanied by the long-term research program TERENO (Terrestrial Environmental Observatories) of the Helmholtz Association (Bogena, 2016), which made additional instrumentation available for TR32. The data presented in this paper is highly valuable for many applications such as outlined in the publication list of TR32 (<http://www.tr32.de>).

Here, we describe the observation sites, the structure of the dataset, and provide information on the observation and measurement methods. Furthermore, we illustrate the quality assurance procedures. With the provision of this dataset, we want

to document our measurement and quality control strategy and provide the scientific community with a comprehensive dataset for further applications.

## 2 Observation sites

All observation sites are located within the Rur catchment located at the Belgium-German-Netherlands border (Fig. 1). The catchment is divided into a fertile loess plain (“Jülicher Börde” and “Zülpicher Börde”) in the north and the low mountain range of the Eifel in the south. The fertile loess plain has a mean elevation of about 100 m above sea level. The land use here is 47 % arable land with the main crops winter wheat, sugar beet, and maize. The area has been inhabited since prehistoric times. Since there are confirmed signs of agriculture from 2000 years ago (Kalis, 1983), it can be assumed, that the soil has been influenced by anthropogenic activities for several thousand years. The warm temperate mid-latitude climate has an annual precipitation of about 700 mm and mean annual air temperature of about 10 °C. The major soils are Haplic Luvisols and Cumulic Anthrosols near the drainage lines, both with silty loamy textures. Soils close to the river Rur are Gleysols and Fluvisols with silty loamy and loamy sandy textures.

The low mountain range in the southern part of the catchment is characterized by a rolling topography. With a mean elevation of about 690 m above sea level and a mean annual precipitation of about 1400 mm, it is dominated by forest and grassland. The major soils are Fluvisols, Gleysols (along the Rur and its tributaries), Eutric Cambisols and Stagnic Gleysols with a silty loamy texture.

Location and numbering of the sites and fields are shown in Fig. 1. Terrain properties of each field are given in Table 1. Permissions to take samples from the fields were given by the respective farmers.

### 2.1 Selhausen

The intensively used cropping-site Selhausen is located in the east of the fertile loess plain (50°52'00" N 6°27'01" E). Crops are grown on gentle slopes (~~< 3°0 to 4°~~). The altitude ranges from 102 until 110 m a.s.l. According to the World Reference Base for Soil Resources (IUSS Working Group WRB, 2015), main soil reference groups are (gleyic) Cambisol and (gleyic) Luvisol. A westbound dip terrace slope cuts through the site with a NNW-SSE strike separating areas with little gravel in the west (fields SE F01, F02, F10, Fig. 1) from areas with more gravel (fields SE F04, F05, F11 – F14). Fields SE F03, F06 – F09, and F15 show a high content of gravel in the east but low content in the west.

The climate exhibits an annual precipitation of 698 mm and a mean annual temperature of 9.9 °C (average for 1961–2008, German weather service station Juelich Kernf.-Anlage, Stat-ID 2474, about 5 km north-west).

The Selhausen site was equipped with eddy covariance stations and meteorological sensors since 2007. Because it is the main agricultural observation site of TR32, numerous ancillary data from the site is available and was presented in the literature (e.g. Busch et al., 2014; Hoffmeister et al., 2016; Korres et al., 2010; Prolingheuer et al., 2014; Schiedung et al., 2017; von Hebel et al., 2018; Bornemann et al., 2011; Ney and Graf, 2018; Schmidt et al., 2012). Beginning in 1895 historical maps

document agricultural land-use for field F01. Based on this information and the general findings, that there has been agriculture in the region for several thousand years, it can be assumed, that effects of conversion of the fields to agricultural area does not have persisting effects on current states or processes.

## 2.2 Merken

5 The Merken site (5°50'47" N 6° 24'04" E) is located 4.5 km to the south west of Selhausen. Therefore, soil texture and meteorological conditions are similar. The area is dominated by agricultural fields. The elevation ranges from 107 to 115 m a.s.l. with slopes of less than 1°. The groundwater at the site is heavily influenced by a nearby open-pit mine. Additional information on the site is presented by Graf et al. (2011). From the farmers in the region it is known, that the region was under agricultural use for at least 100 years. Based on the same information as for Selhausen, it can be assumed, that effects of  
10 conversion of the fields to agricultural area does not have persisting effects on current states or processes.

## 2.3 Merzenhausen

The Merzenhausen site (50°55'47" N 6°17'46" E) is located 13 km to the north west of Selhausen at an altitude of 105 m with a slope of less than 1°. The area is dominated by agricultural fields. Mean annual temperature is 9.7 °C and mean annual precipitation is 750 mm (Schulz, 2004). The soil at the sampling location is described as an Orthic or Haplic Luvisol (Heitmann-  
15 Weber et al., 1994; Schulz, 2004). We don't have detailed information on the land-use history of this field. However, since tombs from the Bronze Age have been found close to the site, concerning effects of land-use conversion, the same assumptions as for Selhausen and Merken apply.

## 2.4 Hürtgenwald

The observation site Hürtgenwald (50°43'26" N 6°22'8" E) is located in the northern part of the low mountain range of the  
20 Eifel. The altitude ranges from 360 to 375 m a.s.l. with varying slopes. The hilly terrain is dominated by forest, pasture and arable land. The reference soil groups are described as Cambisol or Arenosol (Geological Survey of North Rhine-Westphalia). According to long-term private meteorological measurements (www.huertgenwaldwetter.de), the annual precipitation is 946 mm (2000–2018) and the annual mean temperature is 9.4 °C (1998–2018). For the site Hürtgenwald, is known that since the end of World War II, there has not been any forest on the fields. Earlier, they might have been used for forestry. At least since  
25 1953, the fields were used agriculturally alternating between arable land and grassland.

## 3 Conventions and dataset structure

The vegetation data is structured in management periods, which are defined by a combination of the observation site, the field, the crop and the year. A dataset identifier is assigned to each management period as, for example, "SEF05WW15" which describes a management period at the site Selhausen (SE) on field five (F05) where winter wheat (WW) was harvested in the

year 2015 (15). A management period can be either the growing period of a crop or the between-cropping period, where the field is fallow. The fallow period can be discontinuous and refers to the periods before planting and after harvesting of a crop. Data on fluxes and agricultural management ~~can be matched to the management periods~~ ~~are identified~~ by the site, the field, and the year ~~and can thus be assigned to the management periods~~. Meteorological data is given per site. Soil parameters are available for several points at a site. All measurement locations are identified by their positions and are assigned to fields. Fields are defined by field boundaries with a specific land use and homogeneous agricultural management. In the dataset and throughout this text, sites and land-use types are abbreviated as shown in Table 2, while the field numbering is shown in Fig. 1.

Additional conventions:

- 10 • For a crop, the given year is the year when the crop was harvested.
- ~~Throughout the dataset, the symbol NA is used for missing or unknown data.~~
- Time and date are in UTC.
- Coordinates are given in UTM (Zone 32N, WGS84)

The dataset (Reichenau et al., ~~2019~~2020) is provided as a zip-File containing text files in a separate folder for each site as shown in Fig. 2. Details on the data format are described below. An overview of management periods and available data is presented in Table 3.

### 3.1 Missing data

Missing (or unknown) data is denoted by the symbol NA throughout the dataset. There are two main causes of missing data:

- 20 1. Since the data described in this document are mostly results from field measurements, some numbers are missing due to instrument failure or quality issues (see section on quality assurance). For data on agricultural management, data availability depends on the willingness of the farmers to report their activities.
2. Consistency between sites and management periods: If a certain measured variable is available for one site, the variable is also listed in the respective data table of the other sites to keep the data format consistent. If there is no data for the respective variable at a site, all datapoints in that column are marked NA.
- 25 3. Consistency with predefined file formats: For flux data, we used preexisting file formats, which define columns for variables that were not measured in our case, which causes columns totally filled with NA.

## **4 Vegetation data**

### **4.1 Data source and methods**

The vegetation data contains information on fresh and dry biomass, development state, growth height, canopy density, row spacing, and tissue nitrogen and carbon content. Data on biomass is either ~~given~~ differentiated by organ (brown and green leaves and stems, respectively, and fruit) or undifferentiated as overall aboveground biomass (named “biomass\_undiff”).



Furthermore, data on the undifferentiated biomass categories “harvest residues” or “green sprouts” may be included in a record. Harvest residues are understood as the aboveground residues after harvesting which can be material lying on the ground or stubbles left standing. Green sprouts are defined as plants growing between the harvest residues or on an otherwise fallow field. This can be weeds or regrowing crops (especially cereals). In addition, an undifferentiated biomass-category named “biomass\_other” may contain biomass of roots, weeds or the like (specified in the database-column “other\_descr”).

Vegetation data was collected from 2007 to 2017 on different sites and fields (see Table 3). Biomass and leaf area from at least three points in the field were determined destructively. For row crops, also the number of plants in a certain distance of the row was determined. For cereals, plants were taken from 40 or 50 cm in three different rows. Triticale in Hürtgenwald was not sown in rows. Thus, plants from an area of at least 40 x 40 cm were collected. For crops with large individual plants like maize or sugar beet, and for rapeseed, the number of plants per square meter was determined from the row spacing and the number of plants per meter. At least three individual plants were collected at each point. In the field, canopy height and row spacing were measured at each sampling location before cutting the plants. The position in the field was determined using a GPS device. In addition, the phenological development state of the crop was assessed using the BBCH scale (Meier et al., 2009).

After being transported to the lab in airtight bags, the fresh weight (FW) of the plant sample was determined. An aliquot of 150 g or at least one individual plant was further analyzed. In case of a per-organ analysis, the sample was separated into fruit (understood as the harvested organ like ear, beet, etc.), green or brown stems (shoots), and green or brown leaves. A leaf or stem was classified as brown in case 50 % of its surface were not green. A functional definition of a leaf was applied for cereal leaves where only the leaf blade was considered as leaf, while the leaf sheath was assigned to the stem. Blossoms were defined as fruit. For Maize, the male blossoms on top of the plant were assigned to the stem, and only the female blossoms and the maize cobs that evolve from them were defined as fruit.

The leaf area was determined using either a LI-3000A Area Meter with a LI-3050A Belt Conveyer (LI-COR Biosciences, Lincoln, NE, USA) or a flatbed scanner (Epson GT-15000, Seiko Epson Corp., Suwa, Japan) together with the public domain image analysis software ImageJ (<https://imagej.nih.gov/ij/>). In a comparison using the same samples, both methods were shown to give equivalent results. Before determining the dry weight (DW), samples were dried in a drying oven at 105 °C for at least three days. For some samples, aliquots of the dried plant material were homogenized in a mortar and subsequently ground in a ball mill to determine the total content of carbon and nitrogen with an elemental analyzer (CNS Elemental Analyzer Vario EL, Elementar Analysensysteme GmbH, Hanau, Germany). This also includes nine records of C- and N-contents of harvest residues. Upscaling to a square meter of the field was accomplished in a two-step process: from the weighed aliquot to the sample collected in the field and from the sample to a square meter of the field based on the harvested area or the plant density (for MA, SB, RA). Dry weight and LAI were scaled up in proportion to fresh weight.

Additional information:

- The frequency of data collection ranges from one to three weeks. The number of measurements in each management period can be seen in Table 3.

- In the years 2015 until 2017, vegetation data was sampled on overflight days of a radar satellite (Sentinel 1, Radarsat 2).
- Per-organ data of crops for fields at a particular site without organ-specific measurements may be estimated from organ-specific biomass measurements for fields of the same crop on this site assuming equal proportions of the total above ground biomass. The validity of this approach depends on similarity of soil and management conditions.
- Prior to 2011, harvest residues and green sprouts were not sampled in the field. Therefore, these entries are always set undefined (NA) in the years 2007 until 2010. LAI is undefined instead of zero, where no LAI was reported in the field protocol.
- During the management periods HWF04HR15 and SEF04HR16, the fields were fallow. Therefore, all **biomass vegetation data** is zero. These **management periods and other** entries **containing only zeroes** are included in the dataset to document dates, where the field observations showed no biomass on the field. Explicitly distinguishing no biomass from undefined / no data (NA) provides important information for calibration or validation of remote sensing products.

Fig. 3 exemplarily shows dry weights and leaf area index of winter wheat from field F08 at the Selhausen site in 2009 (dataset identifier SEF08WW09). For this management period, three samples per field were collected at each of the 14 dates beginning in December 2008. The last samples were taken on 2009-07-27, one day before harvest on 2009-07-28. The graphs nicely show that the exponential growth phase in April comes along with higher variability between the points in the field in terms of green biomass and LAI. With the beginning of senescence in late May, brown biomass and LAI emerge, showing even higher variability. This is a result of small-scale spatial variability of soil and vegetation properties and terrain under field conditions, which is important information for model evaluation.

Using the data in remote sensing applications often results in scale problems. Since only small patches of 40 x 40 cm or three rows of 50 cm length could be harvested, there is a scale gap between the ground truth and the pixel size of a remote sensing scene, which often have edge lengths of more than 100 m. A possible way to bridge this gap can be high-resolution remote sensing products. For the estimation of LAI, Brogi et al. (2020) calibrated the algorithm of Ali et al. (2015) based on NDVI derived from 5 m resolution RapidEye level 3A data. LAI data from this was used as ground truth. Reichenau et al. (2016) showed that realistic statistical distribution of LAI over a larger area could even be derived without calibration. However, in that case ground truth is required to prove this. The resulting 5 m resolution LAI data can then be spatially aggregated to bridge the gap to lower-resolution datasets.

Examples of the application of the vegetation data can be found in Ahrends et al. (2014), Brogi et al. (2020), Korres et al. (2013), Ney and Graf (2018), Reichenau et al. (2016), Schmidt et al. (2012).

## 4.2 Quality assurance

The first step of the quality assurance procedure for the vegetation data was a rigorous documentation of the measuring process. In addition to written documentation on any phenomena, which might have affected the measurement (in the field and in the lab), a photographic documentation of the samples in the field and in the lab enables a visual inspection and provides

independent evidence in case of any doubts. Transcribing the analog protocols into a spreadsheet-based (MS Excel) digital field protocol provides a first test of data consistency. Possible errors, inconsistencies or incomplete data are reported automatically and the personnel entering the data is prompted to check the entries. Transcribing the data from the analog protocol to a digital data set is done as soon as possible to be able to trace possible errors. Keeping analog field protocols provides a double documentation of the valuable measurements and observations. In a second step, tests on consistency and plausibility were applied which ensure that

- coordinates are in UTM projection, and timestamps are in UTC,
- naming of crops, sites, and points follows conventions,
- values are in plausible ranges,
- missing values are set unknown (NA),
- the right upscaling method is set for a crop throughout a management period,
- there are no duplicate coordinates for points in a field at the same date.

A third step comprises statistical tests, which result in a quality flag for each value in the dataset (see below). These tests were applied using an R-Script (R Core Team, 2017) which reads from the digital field protocols, assigns the quality flags, and finally writes the files provided in the dataset.

**Quality flags.** The quality flags can take the values 1 to 5:

1. High quality (all tests could be applied and no problems were identified; no problems were identified in the field)
2. Good quality (a test could not be applied; information is missing to ensure high quality)
3. Unusual water content (a specific flag concerning the measured water content of the sample which may hint at problems with biomass measurements)
4. Suspicious (a test or a documented issue in the protocol showed possible problems)
5. Low quality (a value is known to have problems but is of interest as an evidence of the real conditions, e.g. root biomass)

The flags were set based on the criteria explained below. After evaluation of all tests, the flag with the highest value was assigned. Obvious erroneous data was removed from the dataset. There are no flags for the carbon and nitrogen contents of the plant tissue.

**General flagging.** Weight measurements below 1 gram were generally flagged as good quality (2) instead of high quality (1), as it is quite likely to lose material from samples, which will have a larger relative effect than for high biomass. All harvest residues are generally flagged suspicious (4). This is due to the fact that precise collection of only the aboveground material is rather difficult and error prone. It is even more difficult to extract the below ground biomass. Therefore, root biomass (given as “biomass\_other”) is generally flagged as low quality (5).

**Loss of material.** In most cases, a sample from the field had to be differentiated into fractions (organs, harvest residues, green sprouts). For larger samples, only a part (aliquot) was analyzed in the lab (see section 4.1). For organ-specific analysis, this aliquot is the sum of all organs. In case of undifferentiated biomass, the aliquot is the sum of the biomass categories

biomass\_undiff, harvest\_residues, green\_sprouts and biomass\_other. During the process of sample partitioning some material might get lost causing a difference between the aliquot and the sum of its components (median 1 %). Differences up to 5 % were accepted independent of their sign. Larger differences result in higher values of the quality flag (Table 4). Higher flags are set in case the sum of its components exceeded the aliquot, because this cannot be explained by losing material.

5 **Reconstruction of missing values.** If an aliquot was available but the FW of one of its components was missing, this FW was recalculated from the difference of the aliquot and the sum of the available FWs. Due to the missing value the loss of material during sample partitioning cannot be determined. Instead, it is contained in the recalculated value, which is therefore flagged as suspicious (4). In this case, the test against the aliquot is not applicable. Thus, the other FWs were flagged as good quality (2).

10 **Comparison of fresh and dry weight.** The comparison of FW and dry weight (DW) can reveal errors in the biomass data. In a first step, it was tested whether DW exceeds FW (Figure 4). For brown leaves and stems, FW and DW were compared directly, while for the other biomass categories, for this test the FW was reduced by 5 % assuming that percentage of minimal water content. In case DW exceeded the resulting FW, it was checked whether the sum of fresh weights was less than 95 % of the aliquot, which hints at a possible error in the FWs (see above). In that case, the error cannot be attributed to neither FW  
15 nor DW and both were removed from the dataset. If the sum of fresh weights was more than 95 % of the aliquot, the error was attributed to the DW, which consequently was removed from the dataset and the corresponding FW was flagged good quality (2).

In a second step it was checked whether the relative water contents  $(FW-DW)/FW$  of green stems, green leaves and fruit are within the range of usual values. This can hint at problems with the DW and FW, which were not identified based on either of  
20 the weight values alone. At first, it was assumed, that living plant tissue has at least a water content of 50 % and DW and FW of green stems or leaves were flagged suspicious (3) if the relative water content was below 50 %. In addition, a “usual course” of the relative water content (Fig. 5) was defined for fruit, green leaves and green stems for winter wheat, winter barley, rapeseed, maize, and sugar beet, respectively. In order to define a lower and upper boundary of the usual water content, the following steps were executed:

- 25
1. Use all water content data for a respective crop and organ.
  2. Exclude outliers by removing all values outside of the 10 % and 90 % percentiles in a running 21-days window.
  3. In each time window, determine the corridor of two standard deviations above and below the mean.

Owing to the low number of data for some crops and organs, and to its scattering, the upper and lower boundaries of the corridor show a lot of scatter. Since there is tendency towards lower water content with progressing phenological development,  
30 the limits of the usual course were defined as follows:

4. Lower limit: For each day in the direction of time only include the lower boundary of the corridor, if it is lower than the value at the previous day. Otherwise, keep the value of the previous day as the lower limit of the usual course.
5. Upper limit: For each day in reverse direction of time only include the upper boundary of the corridor, if it is higher than the value at the following day. Otherwise, keep the value of the following day as the upper limit of the usual course.

For water contents outside of the upper or lower limits, FWs and DWs were assigned the “unusual water content” flag (3). However, this data might also result from particularly dry or wet conditions at a point in a field in a certain year.

**Reported issues.** All issues observed in the field or in the lab which may have had an influence on the results, were translated into flags. For samples reported as dirty, FW and DW were flagged suspicious (4). For humid or wet samples, samples, which might not have been completely dried, and samples which were not analyzed on the same day, only the FW was flagged suspicious (4) since DW is not affected. In case the number of plants per meter was required for upscaling (MA, SB, RA) but missing, this value was derived from other points or dates in the same management period and field. Since this propagates linearly to LAI and to all biomasses per square meter and, since the germination rate is variable in space, all FWs and LAI were flagged suspicious (4).

10 **Propagation of quality flags.** FW and DW are connected by the upscaling process from the aliquot to the sample (see section 4.1), because the upscaling factor derived from FW is also applied to DW. Therefore, flags were propagated from FW to DW and in case of leaves also to LAI.

15 **Coordinates.** To ensure the validity of the location coordinates it was ensured that reported coordinates of a given measurement are within the given field and that no duplicate coordinates are assigned to different measurements at the same date. If it was not possible to correct implausible coordinates, they were removed. In 2008, measurement locations within each field were predefined and marked with flags. Consequentially, coordinates were not recorded explicitly. Since destructive sampling employed in this study prevents repetitive sampling of the exact same location, the prescribed coordinates represent the sampling location less accurate than those recorded directly at the sampling points. Thus, coordinates for 2008 were flagged good quality (2) instead of high quality (1).

## 20 **4.3 Uncertainty**

Uncertainty of biomass data is difficult to estimate. Sources of error exist in all steps of sampling and analysis, including harvest of the samples in the field (incomplete harvest), loss of material and water during handling of the sample, and the unsystematic error of the scales. The error of incomplete harvest cannot be quantified based on the existing data. However, the relative error can be assumed to be rather small for high biomass. The error of handling the sample in the lab (separation of the sample) can be assessed by comparing the weight of the aliquot that was separated by organ with the sum of the organ-weights. Of 1176 organ-specific records in the dataset, 229 have a valid aliquot. The other records either show missing values, or do only have a single organ, or were weighed in total without taking an aliquot. 164 records show a loss of material during separation, while 20 show an increase. The mean loss is 2.6 % of the aliquot (max. 15 %). The mean increase is 2.9 % (max. 17 %). The average error for the (un-)packing steps associated with transport and drying cannot be quantified based on the available data. However, since activities are similar, it can be assumed to be of a similar relative magnitude. The maximum error of the scales used in the lab was 0.1 g. Since leaf area can be measured quite precisely, the relative uncertainty of LAI depends primarily on the accuracy of the leaf weight used for upscaling. Since these are connected linearly in the upscaling process, it equals the relative uncertainty of biomass. A further source of error is the upscaling from the sample taken in the

field to a square meter. For row crops (see section 4.1), the error of the measured row spacing or plant density within the rows propagates linearly into the upscaled result. In order to reduce this error, the median of all row distances measured on a field in a management period was used for upscaling. As the sowing machine settings do not change within a field, the resulting error is considered small. In the field, plant height was measured with a folding rule. The reading accuracy is assumed to be 5 1 cm which is less than the natural variability of plant height.

The uncertainty of carbon and nitrogen contents of the plant tissues was determined by analyzing differences of 1034 duplicate measurements (two aliquots of the same sample). For carbon content, the mean difference of the samples was 0.6 %. For nitrogen content, the mean difference was 1.1 %. The largest differences occurred for root tissues.

Concerning the uncertainty of phenological states in the BBCH system, principal growth stages (first digit) can be assumed to 10 be correct, while secondary growth stages (second digit) may have an error. Since this depends on the observer, it cannot be generally quantified.

#### 4.4 Data format

Vegetation data is supplied per site in a UTF-8 coded csv-file named “vegetation\_” followed by the two-letter site abbreviation (Table 2). Column separator is the semicolon (;). A description of columns and units is presented in Table A1. The no-data 15 symbol is NA. The files have two header lines, of which the first contains the variable names while the second contains the units.

Phenological development (column bbch) may be given as a single number or as a range, if the development state could not be exactly identified in the field. Before sowing and after harvest, the land-use is set to harvest residues (HR) independent of the presence of residues on the surface of the field.

## 20 5 Fluxes of carbon, water and energy

### 5.1 Data source and methods

The dataset contains 15 time series of flux measurements (Table 5). Net fluxes of carbon (net ecosystem exchange, NEE), water (latent energy, LE), and energy (sensible heat flux, H) at the surface were measured at the sites Selhausen, Merzenhausen and Merken using state of the art eddy covariance systems. There were no flux measurements in Hürtgenwald. Wind 25 components and sonic temperature were measured with a three dimensional sonic anemometer (CSAT3, Campbell Scientific, Inc., Logan, UT, USA). Measurements of water vapor (H<sub>2</sub>O) and carbon dioxide (CO<sub>2</sub>) density were carried out using an open-path infrared gas analyzer (IRGA, model LI7500, LI-COR Inc., Lincoln, NE, USA). The previously unavailable data from Merken, contains data for three fields where the each-EC-tower wereas equipped with two sets of sensors at different heights (Table 5). The lower measurement height is usually more representative of the respective land use type. However, the upper 30 level has provided an even better energy balance closure, than the already good one of the lower level.

Measurements were taken with a sampling rate of 20 Hz and were aggregated to intervals of 30 minutes. Processing of raw measurements was accomplished as shown in Fig. 1 of Mauder et al. (2013) using the processing software shown in Table 5. The number of decimal places in the datafiles were kept as they were in the output of the processing software. No gap-filling was applied to the data.

- 5 Examples of the application of the flux data can be found in Ahrends et al. (2014), Eder et al. (2015), Klosterhalfen et al. (2017), Ney and Graf (2018), Schmidt et al. (2012), and Wienecke et al. (2018).

## 5.2 Quality assurance

Quality control was accomplished according to the “TERENO” scheme for quality and uncertainty assessment presented by Mauder et al. (2013). Deviating from this description, before 2011 the software TK2 (Mauder and Foken, 2011) was applied following the process described in section 2.3 of Schmidt et al. (2012). The software ECpack 2.5.20 (Van Dijk et al., 2004) was applied for the data from Merken (Table 5). The software TK uses flagging to indicate the quality of data. Flag values and their meanings are shown in Table 6.

Since flux data from Merken 2009 (MK09) was processed with the ECpack software, the concept of quality assurance differs from the other sites. ECpack provides tolerance values which can be used to rate the quality of data (Table 7). Values outside the lower and upper boundaries given in Table 7 are considered invalid. In addition, data can be filtered using the tolerance values. A tolerance is assigned to the lower and upper boundary of each variable, respectively. To evaluate the quality of the data in the valid ranges, tolerances have to be linearly interpolated between the boundaries. The most obvious tolerance violations have already been eliminated by a post-processing scheme. Tolerance limits were set sufficiently wide to retain most of the values, which still might be useful, in the dataset. For some variables, considering a value to be invalid causes the whole record to be invalid. These variables are assigned to group A in Table 7. If any value of group B is considered invalid, only the values of group B are invalid.

## 5.3 Uncertainty

Uncertainty information for fluxes per data-point is available for sensible heat flux, latent heat flux, NEE, and friction velocity. The kind of uncertainty information differs between the different software tools used for data processing (Table 5). For TK3, relative random errors and relative noise errors for friction velocity, sensible and latent heat flux, and for net ecosystem exchange are given in the respective columns (see Table A3) in the datafiles. For datasets processed with TK2 this information is not available. A rough estimate of the general uncertainty for these measurements may be obtained from statistics of the errors included given in the TK3-processed data. For other variables included in the TK output, the uncertainty is quantified from the instrument errors given by the respective manufacturers (Table 11). The uncertainties of CO<sub>2</sub> and water contents of the air (variables a and CO<sub>2</sub>) strongly depend on calibration. Detailed information can be obtained from the manual of the infrared gas analyzer (LiCOR LI7500, LI-COR Inc., Lincoln, NE, USA). However, the accuracy of the absolute measurements is of minor importance for the eddy-covariance method since it depends on relative changes. The other software tool, ECpack,

calculates 95 % confidence intervals per data-point for fluxes and several other variables. These so-called tolerances are given in the respective columns (see Table A2) in the datafiles. Additional information on uncertainties of eddy-covariance measurements is presented by Mauder et al. (2013).

#### 5.4 Data format

5 Flux data is provided in a UTF-8 coded csv-file per field and year. The filename consists of “fluxes” followed by the two-letter site abbreviation (Table 2), the field ID (Fig. 1), “EC”, a station identifier, and the year. The elements of the filename are separated by underscores (e.g. fluxes\_SEF01\_EC\_001\_2016.csv). Column separator is the semicolon (;). A description of columns and units is presented in Table A2 and Table A3 for the TK and ECpack software, respectively. The no-data symbol is NA. The files have two header lines, of which the first contains the variable names while the second contains the units.

## 10 6 Soil properties

### 6.1 Data source and methods

Soil property data includes particle size distribution of the fine soil (<2 mm), proportion of coarse material (gravel, >2 mm), bulk density, and soil carbon and nitrogen contents. The availability of data differs from site to site (Table 8).

All particle sizes were analyzed following DIN ISO 11277. Therefore, they follow the definition of particle size classes of  
15 DIN 14688. Particles larger than 2 mm are considered gravel. To recalculate particle sizes to the USDA system, which is assumed for many pedotransfer functions, refer to e.g. Nemes et al. (1999). All data on particle sizes and soil carbon or nitrogen-contents refers to fine soil after the removal of gravel. Therefore, percentages of sand, silt, and clay refer to fine soil, while the percentage of gravel refers to total soil mass. Bulk density was determined gravimetrically. Total C concentrations in soil samples were determined by elemental analysis. Based on previous analyses it can be assumed that all samples were  
20 free of carbonates. Hence, total C concentrations are in accordance with those of SOC.

**Selhausen.** Soil data for Selhausen originates from different sources. Particle sizes for three depths in field SE F08 were analyzed at the Laboratory for Physical Geography, University of Cologne. For the ploughing horizon of field SE F00, particle sizes were analyzed at the Institute of Crop Science and Resource Conservation, Bonn University (Bornemann et al., 2011). This data has a high spatial resolution that enables analysis of small-scale heterogeneity. A third dataset consists of horizon-  
25 specific particle size data from 100 randomly chosen points from a 1 km<sup>2</sup> area that includes most fields with vegetation data. The samples were analyzed at the Soil Physical Laboratory of IBG-3, Jülich Research Centre, using a Sedimat 4–12 apparatus (UGT, Umwelt Geräte Technik GmbH, Münchenberg, Germany). From this data and extensive EMI measurements, Brogi et al. (2019) generated a map of soil units, which groups the abovementioned 100 sampling locations into 18 geophysics-based soil units composed of two to twelve sampling locations. These soil units are also provided with quantitative description  
30 (layering, texture, total carbon and nitrogen content) of the soil profile. In the files containing information on the soil and vegetation samples, a column (soil\_unit) establishes the link to the respective soil unit. For several fields, total carbon and



nitrogen contents for three depths were determined from composite samples at the Laboratory for Physical Geography, University of Cologne, using a CNS Elemental Analyzer (Vario EL, Elementar Analysensysteme GmbH, Hanau, Germany). If data was available for several dates, a date after harvest but before the next fertilizer application was preferred if possible. This is noted in the comments column of the data-table. Soil carbon and nitrogen data are assigned to a field instead of a  
5 specific location, because a composite sample containing equal fractions of material from several points was analyzed.

From the 100 sampling points of the 1 km<sup>2</sup> area, carbon and nitrogen content for two horizons, (Ap and Bw) were determined for composite samples from all sampling points within a soil unit, respectively. Therefore, this data is given per soil units. To determine nitrogen and carbon content, a standard combustion method was used at the Geography Institute of the Ruhr University Bochum using a CNS Elemental Analyzer (Vario Max, Elementar Analysensysteme GmbH, Hanau, Germany). All  
10 samples were collected between 6th and 15th February 2017. It has to be noted that samples were collected regardless of the agricultural management.

Due to temporal and spatial variability, this data has to be understood as snapshots and cannot be transferred to other points in space or time.

**Merken.** Particle size data for the Merken site is only available from a composite sample based on samples from all fields.  
15 This data is assumed to be valid for all fields due to small spatial heterogeneity of the soil at the site. The analysis was carried out at the Soil Physical Laboratory of IBG-3, Jülich Research Centre. Field-specific carbon and nitrogen contents for three soil depths were measured from composite samples as described for Selhausen.

**Merzenhausen.** For the field in Merzenhausen, soil texture, bulk density, and contents of carbon and nitrogen were determined for the Ap horizon at a single point in the field following the methodology described by Bornemann et al. (2011). Bulk density  
20 was quantified from three independent 100 cm<sup>3</sup> samples. Since no data was collected for other soil horizons in the frame of the TR32 project, we include data published by Pütz (1993) for the sake of completeness.

**Hürtgenwald.** For Hürtgenwald, particle sizes were analyzed at the Institute of Crop Science and Resource Conservation, Bonn University, while bulk densities were determined at the Laboratory for Physical Geography, University of Cologne.

[Applications of the soil data can be found in Bornemann et al. \(2011\), Brogi et al. \(2019\), Brogi et al. \(2020\), Jakobi et al. \(2020\), Korres et al. \(2013\), Meyer et al \(2017\).](#)  
25

## 6.2 Quality assurance

For the determination of particle sizes, bulk density, and carbon and nitrogen contents, at least two samples from each point were analyzed in parallel. This was not the case for the 1 km<sup>2</sup> data from Selhausen where single analyses were carried out. In this case, the weight of the sample was taken before and after the texture analysis. The analysis was repeated if the final weight  
30 was lower than 95% of the initial weight. If at the second iteration the value was again lower than 95%, the analysis was repeated for a third time.

### 6.3 Uncertainty

To quantify the uncertainty of particle size fractions, data of a repeatedly analyzed sample was evaluated at Bonn University. The results show coefficients of variation (CV) of 2.0 % for sand, 2.4 % for silt, 2.5 % for clay, and 3.5 % for gravel. Since such repeated estimates were not performed at the University of Cologne, it is assumed, that the uncertainty of their measurements is of the same magnitude. At Jülich Research Centre, particle sizes were automatically analyzed with a Sedimat (see above), which has uncertainties in the calculation of the particle size fractions that are comparable to those obtained in above mentioned analysis performed in Bonn. For bulk density, a CV of 10 % was determined from the analysis of multiple adjacent samples from the same horizon (Bonn University). For the soil unit data from Selhausen, uncertainties for particle size fractions and layer depth are given in the respective columns (Table A6) in the datafiles. The CNS Elemental analyzers used to determine soil carbon and nitrogen contents show uncertainties of  $\pm 0.01$  % for carbon and  $\pm 0.002$  % for nitrogen.

### 6.4 Data format

Soil data is provided in a UTF-8 coded csv-file per site named “soil\_” followed by the two-letter site abbreviation (Table 2). Column separator is the semicolon (;). A description of columns and units is presented in Table A5. The no-data symbol is NA. Soil unit data for Selhausen is provided in a UTF-8 coded csv-file named “soil\_units\_SE.csv”. Column separator is the semicolon (;). A description of columns and units is presented in Table A6. The no-data symbol is NA. The files have two header lines, of which the first contains the variable names while the second contains the units.

## 7 Meteorological data

### 7.1 Data source and methods

The dataset was assembled with the aim to provide the data usually required to run a hydroecological crop growth model. Therefore, the dataset includes gap-filled hourly meteorological data of air pressure (AirPres, Pa), air temperature (AirTemp, K), relative air humidity (AirHum, %), wind speed (Wind,  $\text{m s}^{-1}$ ), precipitation (Precip,  $\text{mm h}^{-1}$ ), cloudiness, (Cloudiness, 1/8), and global radiation (Globrad,  $\text{W m}^{-2}$ ). The meteorological data starts about one year earlier than the vegetation data, to provide data for model spin-up concerning water pools in the vadose zone.

The availability of meteorological field data varies between the sites as well as in time. The temporal availability of data increased significantly with time due to the setup of the meteorological stations of the TERENO Eifel/Lower Rhine Valley observatory in 2011. In the earlier years, only a few meteorological stations were run near the sites. Table 9 shows a list of all meteorological stations used in this study. Methods to fill gaps in the time series vary between years and stations. The gap-filling methods are explained in the following sections. The data source for each year and site is presented in Table 10. In most cases, information on the measurement devices and raw data with gaps can be obtained from the data sources shown in the table.

Meteorological data for the site **Hürtgenwald** is provided for the years 2014 through 2016. Meteorological measurements started on 21 April 2015 (station 20 in Table 9). In 2016, additional stations were set up nearby (station 21 in Table 9). Data for earlier dates was generated using the regression gap-filling method (see section 7.1.1) for all variables but AirPres, where gaps were filled using the barometric formula (Eq. 1). The first year for HW consists of reconstructed data only.

5 Data for the site **Merzenhausen** is provided for the years 2009 through 2017. Local measurements are available for the whole period (stations 1 and 15 in Table 9). For the years 2009 and 2011, gaps were filled using the “regression” method. From 2011 on, the EOF method was used (see section 7.1.1).

Data for **Selhausen** is provided for the years 2007 through 2017. Local measurements are available for the whole period, starting on 27 May 2007 (stations 10, 11, and 19 in Table 9). For the years 2007 until 2010, the regression method was used  
10 to fill gaps. From 2011 on, the EOF method was applied.

For the site **Merken**, no meteorological data is available. Since the distance to Selhausen is only about 4 km and the difference in elevation is about 10 m, it can be assumed, that the weather was very similar to that in Selhausen. Therefore, it is suggested to use Selhausen meteorological data when working with Merken vegetation or flux data.

Since cloudiness is not available for any of the sites, but required in some ecohydrological models (e.g. the DANUBIA  
15 simulation system, for an application see e.g. Korres et al., 2013), data on cloudiness from the German National Weather Services’ station Aachen (distances to HW, ME, SE 37, 37, and 42 km, respectively) was used. Since there is no reliable method to adjust cloudiness data to remote stations, the data was used without modifications.

Information on the conditions at the locations of the meteorological stations, especially in the past, are not fully available. Therefore, precipitation data is given as measured at the stations. Since the data was not corrected for shielding effects,  
20 precipitation can be assumed to be slightly underestimated.

Fig. 6 shows an excerpt of the meteorological data for the Selhausen site for the period May to July 2011. The graphs show a period where there are no breaks or shifts in the continuous curves, which is the usual case in the weather timeseries (for a discussion of inhomogeneities, compare section 7.1.3). In the middle of June, the example data shows a noticeable period of two days with low radiation and temperatures together with rather high wind speed and high cloud cover. All variables show  
25 a reduced diurnal cycle, which confirms the consistency of the timeseries’ of the separate variables, which is an important prerequisite for a good reproduction of real processes in a simulation. ~~On short timescales~~

[Applications of the meteorological data can be found in Korres et al. \(2013\), Ney and Graf \(2018\), Sakai et al. \(2016\), and Schmidt et al. \(2012\).](#)

### 7.1.1 Gap filling

30 In the course of the TR32 project, an increasing number of meteorological stations was set up in the Rur catchment. Therefore, different methods were chosen for different periods to fill gaps in the meteorological data.

**Insertion method.** For this simple approach (method 0 in Table 10), data of a nearby station was simply inserted into gaps of the reference station's timeseries. This method was applied in the beginning of the project when only a few stations were set up.

5 **Regression method 1.** This method (method 1 in Table 10) was applied to fill gaps ~~in the years 2007 through 2010 in Selhausen and for the whole period~~ in Hürtgenwald. This method was applied since Hürtgenwald was not included in the central tr32 gap-filling efforts with the EOF method (see below) since it was not an official set of TR32 stations. A simple linear regression was set up between the available data of the station with gaps and a nearby station for each variable, respectively. The slope of the regression was then applied to the data of the nearby station to fill the gap. In case of a data gap at the nearby station, data from a further station was used. In the seldom cases where no data was available at any station, the gap was filled based  
10 on linear interpolation. No gaps longer than four hours had to be filled this way.

**Regression method 2.** For variant 2 of the regression method (method 2 in Table 10), which was applied for the year 2010 in Selhausen and for the years 2009 and 2010 in Merzenhausen, the data of a reference station was correlated with data of the closest remote station using a reduced major axis regression (Webster, 1997). If the coefficient of determination was higher than 0.9 the data of the remote station was inserted into the data gap without further processing (same as insertion method). In  
15 case  $R^2$  was lower than 0.9, the slope of the regression (for AirTemp also the offset) was applied before inserting data into the data gap. For AirHum, the method was applied to dewpoint temperatures, which were converted back to relative humidity after gap-filling. This method was the first central gap-filling effort in the project. Due to the lack of enough stations, the EOF method was not applicable in that period.

**EOF method.** This method (method 3 in Table 10) was applied for the sites Merzenhausen and Selhausen from 2011 on, as  
20 soon as enough stations were available in the TR32 set of meteorological stations. It utilizes empirical orthogonal functions (EOF) to describe the relation between variables at several meteorological stations. The approach was originally introduced by Beckers and Rixen (2003) and adapted for station time series by Graf (2017); further information on EOF computation on similar data can be found in Graf et al. (2012). Since the approach does not depend on the regular spatial arrangement of the  
25 pixels, it can easily be transferred to a network of stations. In contrast to the original approach, this method works on the z-transform of each time series (normalization by dividing the deviations from the mean by the standard deviation), which ensures that stations where the variable has a low amplitude receive the same importance as a predictor as others with a larger amplitude. The following steps were accomplished for each variable separately. Shortwave incoming (global) and photosynthetically active radiation, however, were treated jointly due to their close linear relation:

0. Prior to gap-filling, remove all values rated “bad”, or “suspicious”.
- 30 1. Delete an additional 10 % (randomly selected) of the available data per station and set them aside for cross-validation purposes
2. z-transform the data for each station and variable, respectively
3. Replace all missing values by zeroes

4. Compute the EOFs and reconstruct the time series of each station and variable using only the first EOF (“truncated reconstruction”)
5. Fill all gaps with the reconstruction and repeat step 4 with the filled time series. Repeat the procedure until no data point is changed from one iteration to the next by more than 1 %, or if the change between iterations starts to increase again in at least one data point, or if a maximum of 1000 iterations is reached.
6. Use the dataset with the new preliminary fillers to initialize at step 4 again, but this time using the first two EOFs. Continue as in step 5. After this has converged too, use the first 3 EOFs and so on, until 10 EOFs are used.
7. Re-transform results to absolute values (reverse step 2).
8. Use the cross-validation dataset set aside in step 1 to determine the number of EOFs at which the prediction is optimal (minimum RMSE between validation data and prediction). Repeat the whole procedure up to this number of EOFs starting with step 2 (i.e. without removing cross-validation data).

An advantage of this approach is, that the EOF method exploits the same underlying statistics as multiple linear regression would, but does not need to be re-evaluated each time a predictor variable becomes unavailable. The method was applied to 10-minute resolution data from stations 1 to 18 (Table 9). Results were aggregated to hourly resolution.

### 15 **Gap-filling of cloudiness data**

Gaps in cloudiness data were filled using the “na.approx” method in the R-package “zoo” (Zeileis and Grothendieck, 2005).

#### **7.1.2 Adjustment of atmospheric pressure**

For the sites and years where the EOF-Method was not applied, air pressure (AirPres, in hPa) data was transformed between stations by using the barometric formula:

$$20 \text{ AirPres} = \text{AirPres}_r \left( 1 - \left( 0.0065 \frac{\Delta h}{\text{AirTemp}} \right) \right)^{5.255} \quad (1)$$

Where  $\Delta h$  is the elevation difference between Stations (m), AirTemp is the air temperature (K), and AirPres is the atmospheric pressure at the remote station (hPa).

#### **7.1.3 Inhomogeneities**

- A closer look at the time series of meteorological data reveals differences in general characteristics between different years.
- 25 This is mainly due to different instruments or different calibration of instruments. By these means, synthetic breaks in the time series are generated that can disturb the analysis of real phenomena. This is particularly a problem when using the data with models, which deterministically transform weather data into plant growth and into exchange fluxes of matter and energy.
- Several breaks can easily be identified from graphical visualizations of the data. Fig. 7 a shows a shift in air pressure measured in Selhausen from 2009 to 2010 using different instruments. A similar effect can be observed in the Merzenhausen data. Fig.
- 30 7 b illustrates different maxima of relative humidity in 2015 in Hürtgenwald, which are due to differences in instrument calibration. This effect can also be found in the data for Merzenhausen and Selhausen. Other obvious breaks refer to lower

extrema of air temperature (SE and ME), maxima of global radiation (ME), maxima of wind speed (SE), and changing temporal variability of wind speed (HW). Often, these breaks coincide with a change in the main source station (Table 10). Other less noticeable breaks may be included in the time series.

The removal of such breaks in the time series is known as homogenization in the literature. Several methods have been developed to detect the breaks and correct for inconsistencies. However, most of these methods were designed for monthly or annual data (Venema et al., 2012), and are not applicable to subdaily data (Aguilar et al., 2003, Auer et al., 2005, Wijngaard et al., 2003). Since methods for data on higher temporal resolutions would involve dealing with non-linear atmospheric processes (Della-Marta and Wanner, 2006), the world meteorological organization does not yet make any recommendations on how to homogenize this data. Nevertheless, the following literature might help finding an appropriate homogenization method for the intended application of the data. Temperature: Vincent et al. (2002), Brandsma and Können (2006), Della-Marta and Wanner (2006), Kuglitsch et al. (2009), Mestre et al. (2011), Trewin (2013); Precipitation: Beaulieu et al. (2008), Beaulieu et al. (2009); Both: Domonkos and Coll (2017).

## 7.2 Quality assurance

Time series of meteorological data were checked for plausibility of the recorded data. Values outside of a plausible range were removed from the dataset. Periods of repeated identical (but plausible) values were removed. To ensure good quality of gap-filling, the gap-filling methods were applied to periods with good quality measurements.

## 7.3 Uncertainty

Measurement uncertainties of weather variables are given as instrument errors in Table 11. It has to be mentioned that especially for precipitation, the instrument error is much smaller than systematic errors. For a discussion of such errors, compare Dengel et al. (2018).

Additional uncertainty occurs when gaps in timeseries are filled based on data from other stations. Because different methods and data sources were used, uncertainty was determined separately for the different sites and years. For the years 2007 to 2010, uncertainty was estimated by deriving a fill-value from remote stations for each available value at the reference station using the respective method shown in Table 10. Bias and root mean square error (RMSE) were calculated from the differences (Table 12). These results for the Selhausen site are assumed to be transferable to the other sites.

For the EOF method, which was applied for the years from 2011 to 2017, an extra run on a dataset copy with artificial gaps was used to determine worst-case uncertainty estimates. These artificial gaps were inserted for the Merzenhausen site for the 2.5 consecutive days with the highest mean for the respective variable (relative humidity: lowest mean) for all sensors at the site. The artificial gaps were then filled and the differences to the measured data were evaluated in terms of bias and RMSE (Table 12). By selecting an extreme situation for gap-filling, uncertainties for the EOF-method are a worst-case estimate. Inserting arbitrary gaps would probably give lower uncertainty values. In addition to this, when comparing uncertainty estimates between different periods, it has to be taken into account that the analysis for the EOF method was applied to raw

10-min data while the evaluation for the years 2007 to 2010 is based on hourly data, which generally results in slightly lower RMSE values. Again, we assume that results can be transferred to the Selhausen site.

For precipitation in the period 2007 to 2010 and for global radiation in 2010, uncertainty estimates cannot be given since the raw data is no longer available. Data from the German weather service (DWD) was used for global radiation in 2007 to 2009  
5 and for air pressure in 2010. Since this data was without gaps, there is no gap-filling uncertainty for these variables.

## 7.4 Data format

Weather data is provided in a UTF-8 coded csv-file per site named “meteo\_” followed by the two-letter site abbreviation (Table 2) and the span of years available. Column separator is the semicolon (;). A description of columns and units is presented in Table A7. The no-data symbol is NA. The files have two header lines, of which the first contains the variable names while  
10 the second contains the units.

## 8 Crop management data

### 8.1 Data source and methods

Crop management data was inquired from the farmer of the respective fields by means of a questionnaire. However, the information given by the farmers was only reported for 27 out of 58 management periods and is often incomplete (Table 3).

15 The following information was inquired:

- Sowing date
- Sowing density, row spacing, seed spacing in a row, seed weight, sowing depth, cultivar
- Fertilization date, amount, and product
- Cultivation date and type
- 20 • Growth regulator application date, amount, and product
- Fungicide/Insecticide/Herbicide application date, amount and product
- Harvesting date
- Dry weight of yield after harvest
- Information on residues left on the field

25 All fertilization data was recalculated to kilograms nitrogen per hectare. Since for some products, nitrogen content is not explicitly stated, the following assumptions were made: It was assumed that KAS (calcium ammonium nitrate) contains 27 mass-% nitrogen. Furthermore, it was assumed that Sulfan contains 24 % nitrogen. AHL (urea ammonium nitrate solution, UAN) was assumed to have a density of 1.3 kg l<sup>-1</sup>. All fields were managed conventionally.

Applications of the management data can be found in Korres et al. (2013) or Schmidt et al. (2012).

## 8.2 Quality assurance

Some of the fields were equipped with automatic camera systems, which took hourly photos. Management information gathered from the farmers were checked against these photos.

## 8.3 Uncertainty

- 5 Accuracy of management data is based on the reliability of the information provided by the farmers. Since there is no way to check information on fertilizer or agrochemical types and amounts, an uncertainty cannot be assigned.

## 8.4 Data format

Management data is provided in a UTF-8 coded csv-file per management period. The filename starts with “management\_” followed by the ID of the management period (e.g. management\_SEF08WW09.txt). The file can contain data on management activities in the fallow period before or after harvest. If no management information is available, the file contains a comment only. There are no management files for management periods denominated “harvest residues” (HR).

Each record is structured in the same way: date; keyword; additional information. The elements of the record are separated by a semicolon (;). The record starts with the date in YYYY-MM-DD format, where day may be replaced by “xx” if the exact date is unknown. In the second position, the record contains a keyword that defines the management activity. Keywords refer to basic crop related activities (“Sowing”, “Harvesting”, “Fertilizer”, “Cutting”), soil management (“Plow”, “Rotary harrow”, “Harrow”, “Roller”, “Cultivator”, “Tyre Packer”), and application of agrochemicals (“Herbicide”, “Growth control”, “Fungicide”, “Insecticide”, “Co Formulant”). After the keyword, one or more pieces of additional information may follow in a semicolon-separated list:

- Fertilizer: amount of fertilizer in kilograms nitrogen per hectare; information on the product and its contents (may also be a semicolon-separated list)
- Application of agrochemicals: amount of agrochemical per area; information on the product and its contents (may also be a semicolon-separated list)
- Sowing: sowing density; row spacing; seed spacing; weight of seeds; sowing depth; cultivar

Unknown information is indicated by the no-data symbol NA. Units are given with the data. Comments start with “#”. Comments can contain additional information on yield, management of harvest residues, additional contents of agrochemical, etc.

## Data availability

The dataset (Reichenau et al., [20192020](#)) can be downloaded from the TR32 database (<https://www.tr32db.uni-koeln.de/data.php?dataID=18869>) or using the DOI [10.5880/TR32DB.39XXXX](#). The dataset is provided as a zip-compressed container. All files are plain text files organized in a folder per site as show in Fig. 2 and as explained in section 3. Technical



details on file formats and data structure within files is presented for the different kinds of data in sections 4.4, 5.4, 6.5, 7.4, and 8.4.

## **Appendix:**

The tables in the appendix describe the datafiles in terms of their column order, variables, units, and datatypes.

## **5 Author contributions**

TR designed and compiled the dataset, and did the quality control and processing of the vegetation data. AS and AG did the gap filling of meteorological data and developed the methods. MS processed the eddy covariance data and collected the management data. WK and KS were responsible for the collection of the vegetation data and some soil data. GW, NM, and CB took and analyzed soil samples. The manuscript was prepared by TR with contributions of all co-authors.

## **10 Competing interests**

The authors declare that they have no conflict of interest.

## **Acknowledgements**

This study was supported by the Deutsche Forschungsgemeinschaft through the Transregional Collaborative Research Center 32 – Patterns in Soil-Vegetation-Atmosphere Systems: Monitoring, Modelling and Data Assimilation. In addition, support was received through the “Terrestrial ENvironmental Observatories” (TERENO). We thank the numerous student helpers for their help with the field campaigns. Special thanks go to the farmers who granted access to their fields and to our student helpers Michael Holthausen (gap filling of meteorological data), Tobias Bothe (gap filling and soil sampling), and Nils Eingrüber (consistency check of soil data). We thank Marius Schmidt and Florian Steininger for collecting the management data from the farmers, Victor Venema for discussions and literature on the homogenization of meteorological [data](#), Ulrike Schwedler for cartography, and Constanze Curdt for advice concerning all aspects of data management. [Finally we want to thank five reviewers for their helpful comments.](#)

## **References**

Aguilar, E., Auer, I., Brunet, M., Peterson, T. C. and Wieringa, J.: Guidelines on climate metadata and homogenization, WMO/TD No. 1186, Geneva, Switzerland., 2003.


- [Ahrends, H., Haseneder-Lind, R., Schween, J., Crewell, S., Stadler, A. and Rascher, U.: Diurnal Dynamics of Wheat Evapotranspiration Derived from Ground-Based Thermal Imagery, \*Remote Sens.\*, 6\(10\), 9775–9801, doi:10.3390/rs6109775, 2014.](#)
- 5 [Ali, M., Montzka, C., Stadler, A., Menz, G., Thonfeld, F. and Vereecken, H.: Estimation and Validation of RapidEye-Based Time-Series of Leaf Area Index for Winter Wheat in the Rur Catchment \(Germany\), \*Remote Sens.\*, 7\(3\), 2808–2831, doi:10.3390/rs70302808, 2015.](#)
- Auer, I., Böhm, R., Jurković, A., Orlik, A., Potzmann, R., Schöner, W., Ungersböck, M., Brunetti, M., Nanni, T., Maugeri, M., Briffa, K., Jones, P., Efthymiadis, D., Mestre, O., Moisselin, J.-M., Begert, M., Brazdil, R., Bochnicek, O., Cegnar, T., Gajić-Čapka, M., Zaninović, K., Majstorović, Ž., Szalai, S., Szentimrey, T. and Mercalli, L.: A new instrumental precipitation dataset for the greater alpine region for the period 1800-2002: Precipitation Dataset: European Greater Alpine Region, *Int. J. Climatol.*, 25(2), 139–166, doi:10.1002/joc.1135, 2005.
- 10 Beaulieu, C., Seidou, O., Ouarda, T. B. M. J., Zhang, X., Boulet, G. and Yagouti, A.: Intercomparison of homogenization techniques for precipitation data: Homogenization of Precipitation, *Water Resour. Res.*, 44(2), doi:10.1029/2006WR005615, 2008.
- 15 Beaulieu, C., Seidou, O., Ouarda, T. B. M. J. and Zhang, X.: Intercomparison of homogenization techniques for precipitation data continued: Comparison of two recent Bayesian change point models: Homogenization with Bayesian Change Point, *Water Resour. Res.*, 45(8), doi:10.1029/2008WR007501, 2009.
- Beckers, J. M. and Rixen, M.: EOF Calculations and Data Filling from Incomplete Oceanographic Datasets\*, *J. Atmospheric Ocean. Technol.*, 20(12), 1839–1856, doi:10.1175/1520-0426(2003)020<1839:ECADFF>2.0.CO;2, 2003.
- 20 [Bogena, H. R.: TERENO, \*J. Large Scale Res. Facil. JLSRF\*, 2, doi:10.17815/jlsrf-2-98, 2016.](#)
- [Bogena, H. R.: TERENO: German network of terrestrial environmental observatories, \*J. Large-Scale Res. Facil. JLSRF\*, 2, doi:10.17815/jlsrf-2-98, 2016.](#)
- Bornemann, L., Herbst, M., Welp, G., Vereecken, H. and Amelung, W.: Rock Fragments Control Size and Saturation of Organic Carbon Pools in Agricultural Topsoil, *Soil Sci. Soc. Am. J.*, 75(5), 1898, doi:10.2136/sssaj2010.0454, 2011.
- 25 Brandsma, T. and Können, G. P.: Application of nearest-neighbor resampling for homogenizing temperature records on a daily to sub-daily level, *Int. J. Climatol.*, 26(1), 75–89, doi:10.1002/joc.1236, 2006.
- Brogi, C., Huisman, J. A., Pätzold, S., von Hebel, C., Weihermüller, L., Kaufmann, M. S., van der Kruk, J. and Vereecken, H.: Large-scale soil mapping using multi-configuration EMI and supervised image classification, *Geoderma*, 335, 133–148, doi:10.1016/j.geoderma.2018.08.001, 2019.
- 30 [Brogi, C., Huisman, J. A., Herbst, M., Weihermüller, L., Klosterhalfen, A., Montzka, C., Reichenau, T. G. and Vereecken, H.: Simulation of spatial variability in crop leaf area index and yield using agroecosystem modeling and geophysics-based quantitative soil information, \*Vadose Zone J.\*, 19\(1\), doi:10.1002/vzj2.20009, 2020.](#)
- Busch, S., van der Kruk, J. and Vereecken, H.: Improved Characterization of Fine-Texture Soils Using On-Ground GPR Full-Waveform Inversion, *IEEE Trans. Geosci. Remote Sens.*, 52(7), 3947–3958, doi:10.1109/TGRS.2013.2278297, 2014.

- Della-Marta, P. M. and Wanner, H.: A Method of Homogenizing the Extremes and Mean of Daily Temperature Measurements, *J. Clim.*, 19(17), 4179–4197, doi:10.1175/JCLI3855.1, 2006.
- Dengel, S., Graf, A., Grünwald, T., Hehn, M., Kolari, P., Löfvenius, M. O., Merbold, L., Nicolini, G. and Pavelka, M.: Standardized precipitation measurements within ICOS: rain, snowfall and snow depth: a review, *Int. Agrophysics*, 32(4), 607–617, doi:10.1515/intag-2017-0046, 2018.
- Domonkos, P. and Coll, J.: Homogenisation of temperature and precipitation time series with ACMANT3: method description and efficiency tests: Homogenisation of time series with ACMANT3, *Int. J. Climatol.*, 37(4), 1910–1921, doi:10.1002/joc.4822, 2017.
- Eder, F., Schmidt, M., Damian, T., Träumner, K. and Mauder, M.: Mesoscale Eddies Affect Near-Surface Turbulent Exchange: Evidence from Lidar and Tower Measurements, *J. Appl. Meteorol. Climatol.*, 54(1), 189–206, doi:10.1175/JAMC-D-14-0140.1, 2015.
- Graf, A.: Gap-filling meteorological variables with Empirical Orthogonal Functions, *Geophys. Res. Abstr.*, 19, 2017.
- Graf, A., Prolingheuer, N., Schickling, A., Schmidt, M., Schneider, K., Schüttemeyer, D., Herbst, M., Huisman, J. A., Weihermüller, L., Scharnagl, B., Steenpass, C., Harms, R. and Vereecken, H.: Temporal Downscaling of Soil Carbon Dioxide Efflux Measurements Based on Time-Stable-Spatial Patterns, *Vadose Zone J.*, 10(1), 239, doi:10.2136/vzj2009.0152, 2011.
- Graf, A., Herbst, M., Weihermüller, L., Huisman, J. A., Prolingheuer, N., Bornemann, L. and Vereecken, H.: Analyzing spatiotemporal variability of heterotrophic soil respiration at the field scale using orthogonal functions, *Geoderma*, 181–182, 91–101, doi:10.1016/j.geoderma.2012.02.016, 2012.
- von Hebel, C., Matveeva, M., Verweij, E., Rademske, P., Kaufmann, M. S., Brogi, C., Vereecken, H., Rascher, U. and van der Kruk, J.: Understanding Soil and Plant Interaction by Combining Ground-Based Quantitative Electromagnetic Induction and Airborne Hyperspectral Data, *Geophys. Res. Lett.*, 45(15), 7571–7579, doi:10.1029/2018GL078658, 2018.
- Heitmann-Weber, B., Mittelstaedt, W. and Fuhr, F.: The degradation of anilazine and dihydroxy-anilazine at various soil depths of an orthic luvisol, *J. Environ. Sci. Health Part B*, 29(2), 247–264, doi:10.1080/03601239409372878, 1994.
- Hoffmeister, D., Waldhoff, G., Korres, W., Curdt, C. and Bareth, G.: Crop height variability detection in a single field by multi-temporal terrestrial laser scanning, *Precis. Agric.*, 17(3), 296–312, doi:10.1007/s11119-015-9420-y, 2016.
- IUSS Working Group WRB: World Reference Base for Soil Resources 2014, update 2015. FAO, Rome, Italy, 2015.
- Jakobi, J., Huisman, J. A., Schrön, M., Fiedler, J., Brogi, C., Vereecken, H. and Bogena, H. R.: Error Estimation for Soil Moisture Measurements With Cosmic Ray Neutron Sensing and Implications for Rover Surveys, *Front. Water*, 2, doi:10.3389/frwa.2020.00010, 2020.
- Jones, J. W., Antle, J. M., Basso, B., Boote, K. J., Conant, R. T., Foster, I., Godfray, H. C. J., Herrero, M., Howitt, R. E., Janssen, S., Keating, B. A., Munoz-Carpena, R., Porter, C. H., Rosenzweig, C. and Wheeler, T. R.: Toward a new generation of agricultural system data, models, and knowledge products: State of agricultural systems science, *Agric. Syst.*, 155, 269–288, doi:10.1016/j.agsy.2016.09.021, 2017.

- [Kalis, A. J.: Die menschliche Beeinflussung der Vegetationsverhältnisse auf der Aldenhovener Platte \(Rheinland\) während der vergangenen 2000 Jahre, in Archäologie in den rheinischen Lößböden: Beiträge zur Siedlungsgeschichte im Rheinland, vol. 24, Rheinland-Verlag, Köln, 1983.](#)
- 5 Kersebaum, K. C., Boote, K. J., Jorgenson, J. S., Nendel, C., Bindi, M., Frühauf, C., Gaiser, T., Hoogenboom, G., Kollas, C., Olesen, J. E., Rötter, R. P., Ruget, F., Thorburn, P. J., Trnka, M. and Wegehenkel, M.: Analysis and classification of data sets for calibration and validation of agro-ecosystem models, *Environ. Model. Softw.*, 72, 402–417, doi:10.1016/j.envsoft.2015.05.009, 2015.
- [Klosterhalfen, A., Herbst, M., Weihermüller, L., Graf, A., Schmidt, M., Stadler, A., Schneider, K., Subke, J.-A., Huisman, J. A. and Vereecken, H.: Multi-site calibration and validation of a net ecosystem carbon exchange model for croplands, \*Ecol. Model.\*, 363, 137–156, doi:10.1016/j.ecolmodel.2017.07.028, 2017](#)
- 10 Korres, W., Koyama, C. N., Fiener, P. and Schneider, K.: Analysis of surface soil moisture patterns in agricultural landscapes using Empirical Orthogonal Functions, *Hydrol. Earth Syst. Sci.*, 14(5), 751–764, doi:10.5194/hess-14-751-2010, 2010.
- Korres, W., Reichenau, T. G. and Schneider, K.: Patterns and scaling properties of surface soil moisture in an agricultural landscape: An ecohydrological modeling study—[ScienceDirect](#), *J. Hydrol.*, 498, 89–102, 2013.
- 15 Kuglitsch, F. G., Toreti, A., Xoplaki, E., Della-Marta, P. M., Luterbacher, J. and Wanner, H.: Homogenization of daily maximum temperature series in the Mediterranean, *J. Geophys. Res.*, 114(D15), doi:10.1029/2008JD011606, 2009.
- Mauder, M. and Foken, T.: Documentation and Instruction Manual of the Eddy-Covariance Software Package TK3, Universität Bayreuth, Abt. Mikrometeorologie, Bayreuth., 2011.
- Mauder, M., Cuntz, M., Drüe, C., Graf, A., Rebmann, C., Schmid, H. P., Schmidt, M. and Steinbrecher, R.: A strategy for quality and uncertainty assessment of long-term eddy-covariance measurements, *Agric. For. Meteorol.*, 169, 122–135, doi:10.1016/j.agrformet.2012.09.006, 2013.
- 20 Meier, U., Bleiholder, H., Buhr, L., Feller, C., Hack, H., Heß, M., Lancashire, P. D., Schnock, U., Stauß, R., van den Boom, T., Weber, E. and Zwerger, P.: The BBCH system to coding the phenological growth stages of plants – history and publications –, *J. Cultiv. Plants*, 61(2), 41–52, 2009.
- 25 Mestre, O., Gruber, C., Prieur, C., Caussinus, H. and Jourdain, S.: SPLIDHOM: A Method for Homogenization of Daily Temperature Observations, *J. Appl. Meteorol. Climatol.*, 50(11), 2343–2358, doi:10.1175/2011JAMC2641.1, 2011.
- [Meyer, N., Bornemann, L., Welp, G., Schiedung, H., Herbst, M. and Amelung, W.: Carbon saturation drives spatial patterns of soil organic matter losses under long-term bare fallow, \*Geoderma\*, 306, 89–98, doi:10.1016/j.geoderma.2017.07.004, 2017.](#)
- 30 Nemes, A., Wosten, J. H. M., Lilly, A. and Voshaar, J. H. O.: Evaluation of different procedures to interpolate particle-size distributions to achieve compatibility within soil databases, *GEODERMA*, 90(3–4), 187–202, 1999.
- Ney, P. and Graf, A.: High-Resolution Vertical Profile Measurements for Carbon Dioxide and Water Vapour Concentrations Within and Above Crop Canopies, *Bound.-Layer Meteorol.*, 166(3), 449–473, doi:10.1007/s10546-017-0316-4, 2018.

- Prolingheuer, N., Scharnagl, B., Graf, A., Vereecken, H. and Herbst, M.: On the spatial variation of soil rhizospheric and heterotrophic respiration in a winter wheat stand, *Agric. For. Meteorol.*, 195–196, 24–31, doi:10.1016/j.agrformet.2014.04.016, 2014.
- Pütz, T.: Lysimeterversuche zum Verlagerungsverhalten von Methabenzthiazuron und gelöstem organischen Kohlenstoff in einer Parabraunerde, Aufbau von zwei Klimameßstationen und Untersuchungen zur Validierung des Lysimetersystems, 5  
Forschungszentrum Zentralbibliothek, Jülich. [online] Available from: <http://gso.gbv.de/DB=2.1/PPNSET?PPN=152374191>, 1993.
- R Core Team: R, R Foundation for Statistical Computing, Vienna, Austria., 2017.
- Reichenau, T. G., Korres, W., Montzka, C., Fiener, P., Wilken, F., Stadler, A., Waldhoff, G. and Schneider, K.: Spatial  
10 Heterogeneity of Leaf Area Index (LAI) and Its Temporal Course on Arable Land: Combining Field Measurements, Remote Sensing and Simulation in a Comprehensive Data Analysis Approach (CDA), edited by D. Hui, PLOS ONE, 11(7), e0158451, doi:10.1371/journal.pone.0158451, 2016.
- Reichenau, T. G., Korres, W., Schmidt, M., Graf, A., Welp, G., Meyer, N., Stadler, A., Brogi, C. and Schneider, K.: A comprehensive dataset of vegetation states, fluxes of matter and energy, weather, agricultural management, and soil properties 15  
from intensively monitored crop sites in Western Germany, [online] Available from: <https://www.tr32db.uni-koeln.de/data.php?dataID=18869>, doi: 10.5880/TR32DB.39, 20192020.
- Sakai, T., Iizumi, T., Okada, M., Nishimori, M., Grünwald, T., Prueger, J., Cescatti, A., Korres, W., Schmidt, M., Carrara, A., Loubet, B. and Ceschia, E.: Varying applicability of four different satellite-derived soil moisture products to global gridded crop model evaluation, Int. J. Appl. Earth Obs. Geoinformation, 48, 51–60, doi:10.1016/j.jag.2015.09.011, 2016.
- 20 Schad, P., Krasilnikov, P. V. and Arnold, R.: German Soil Classification, in A handbook of soil terminology, correlation and classification, edited by P. V. Krasilnikov, J. J. Ibáñez Martí, R. Arnold, and S. Shoba, pp. 122–130, Earthscan, London ; Sterling, VA., 2009.
- Schiedung, H., Bornemann, L. and Welp, G.: Seasonal Variability of Soil Organic Carbon Fractions Under Arable Land, *Pedosphere*, 27(2), 380–386, doi:10.1016/S1002-0160(17)60326-6, 2017.
- 25 Schmidt, M., Reichenau, T. G., Fiener, P. and Schneider, K.: The carbon budget of a winter wheat field: An eddy covariance analysis of seasonal and inter-annual variability, *Agric. For. Meteorol.*, 165, 114–126, doi:10.1016/j.agrformet.2012.05.012, 2012.
- Schulz, E.: Influence of site conditions and management on different soil organic matter (som) pools, *Arch. Agron. Soil Sci.*, 50(1), 33–47, doi:10.1080/03650340310001627577, 2004.
- 30 Simmer, C., Thiele-Eich, I., Masbou, M., Amelung, W., Bogen, H., Crewell, S., Diekkrüger, B., Ewert, F., Hendricks Franssen, H.-J., Huisman, J. A., Kemna, A., Klitzsch, N., Kollet, S., Langensiepen, M., Löhnert, U., Rahman, A. S. M. M., Rascher, U., Schneider, K., Schween, J., Shao, Y., Shrestha, P., Stiebler, M., Sulis, M., Vanderborght, J., Vereecken, H., van der Kruk, J., Waldhoff, G. and Zerenner, T.: Monitoring and Modeling the Terrestrial System from Pores to Catchments, *Bull. Am. Meteorol. Soc.*, 96(10), 1765–1787, doi:10.1175/BAMS-D-13-00134.1, 2015.

- Trewin, B.: A daily homogenized temperature data set for Australia, *Int. J. Climatol.*, 33(6), 1510–1529, doi:10.1002/joc.3530, 2013.
- Van Dijk, A., Moene, A. F. and De Bruin, H. A. R.: The principles of surface flux physics: theory, practice and description of the ECPACK library, Meteorology and Air Quality Group, Wageningen University, Wageningen, the Netherlands., 2004.
- 5 Venema, V. K. C., Mestre, O., Aguilar, E., Auer, I., Guijarro, J. A., Domonkos, P., Vertacnik, G., Szentimrey, T., Stepanek, P., Zahradnick, P., Viarre, J., Müller-Westermeier, G., Lakatos, M., Williams, C. N., Menne, M. J., Lindau, R., Rasol, D., Rustemeier, E., Kolokythas, K., Marinova, T., Andresen, L., Acquaotta, F., Fratianni, S., Cheval, S., Klancar, M., Brunetti, M., Gruber, C., Prohom Duran, M., Likso, T., Esteban, P. and Brandsma, T.: Benchmarking homogenization algorithms for monthly data, *Clim. Past*, 8(1), 89–115, doi:10.5194/cp-8-89-2012, 2012.
- 10 Vereecken, H., Kollet, S. and Simmer, C.: Patterns in Soil–Vegetation–Atmosphere Systems: Monitoring, Modeling, and Data Assimilation, *Vadose Zone J.*, 9(4), 821, doi:10.2136/vzj2010.0122, 2010.
- Vincent, L. A., Zhang, X., Bonsal, B. R. and Hogg, W. D.: Homogenization of Daily Temperatures over Canada, *J. Clim.*, 15(11), 1322–1334, doi:10.1175/1520-0442(2002)015<1322:HODTOC>2.0.CO;2, 2002.
- Webster, R.: Regression and functional relations, *Eur. J. Soil Sci.*, 48(3), 557–566, doi:10.1111/j.1365-2389.1997.tb00222.x, 15 1997.
- [Wieneke, S., Burkart, A., Cendrero-Mateo, M. P., Julitta, T., Rossini, M., Schickling, A., Schmidt, M. and Rascher, U.: Linking photosynthesis and sun-induced fluorescence at sub-daily to seasonal scales, \*Remote Sens. Environ.\*, 219, 247–258, doi:10.1016/j.rse.2018.10.019, 2018.](https://doi.org/10.1016/j.rse.2018.10.019)
- Wijngaard, J. B., Klein Tank, A. M. G. and Können, G. P.: Homogeneity of 20th century European daily temperature and 20 precipitation series: Homogeneity of European Climate Series, *Int. J. Climatol.*, 23(6), 679–692, doi:10.1002/joc.906, 2003.
- Zeileis, A. and Grothendieck, G.: zoo: S3 Infrastructure for Regular and Irregular Time Series, *J. Stat. Softw.*, 14(6), doi:10.18637/jss.v014.i06, 2005.



5 **Figure 1:** Left: Locations of the observation sites in the Rur catchment in Germany. Right: Locations of the fields at the observation sites with two-digit field IDs. At the Selhausen site (3), field 12 is a part of field 11. On the aerial photo of the Merken site (2), a part of field 01 is on the area of an open pit mine. At the time of field measurements, the mine was about 2.5 km away from the field. Map-data: GADM ([gadm.org/license.html](http://gadm.org/license.html)), OpenStreetMap (Open Database Licence (ODbL) 1.0). Aerial photography: Land NRW (2019) Datenlizenz Deutschland - Namensnennung - Version 2.0 ([www.govdata.de/dl-de/by-2-0](http://www.govdata.de/dl-de/by-2-0)).




Figure 2: Folder structure of the dataset. A slash (“/”) denotes a directory.

5






Figure 3: Dry weight (DW) and leaf area index (LAI) of winter wheat on field F08 at the Selhausen site in 2009. Dataset identifier is SEF08WW09.






**Figure 4: Flow chart of the decision process of quality assurance when fresh weight (FW) was found larger than dry weight (DW).**



5 **Figure 5: Temporal course of sugar beet leaves' relative water content (percentage of fresh weight, all available data). Black lines show the upper and lower limit of the "usual course" (definition in the text). Blue circles denote measurements outside the usual course and are therefore assigned the unusual water content flag (3). This data may still be valid because of heterogeneous conditions in a field (e.g. because of earlier drying).**



**Figure 6:** Excerpt of the gap-filled hourly meteorological data for the Selhausen site for the period from May until July 2011.




Figure 7: Breaks in time series of meteorological measurements. a) Air pressure in Selhausen, b) Air humidity in Hürtgenwald.

**Table 1: Terrain properties of the fields. Coordinates are for centroids, projection is UTM 32N (WGS 1984).**

site	field	UTM N	UTM E	elev. (m)	area (ha)	slope (°)
Selhausen (SE)	F00	5638377	320584	105	1.2	1.5
	F01	5638008	320341	103	9.7	0.4
	F02	5637780	320428	104	7.4	0.4
	F03	5638056	320643	105	4.0	1.1
	F04	5638122	320826	109	1.9	0.7
	F05	5637987	320860	110	2.4	0.9
	F06	5637683	320723	107	2.4	1.5
	F07	5638251	320613	105	2.4	1.4
	F08	5638568	320538	104	6.5	1.4
	F09	5638818	320403	102	2.6	0.8
	F10	5638362	320408	103	2.2	0.7
	F11	5638671	320699	106	4.8	1.0
	F12	5638617	320713	107	3.1	0.7
	F13	5638478	320742	108	0.8	0.4
	F14	5638434	320754	109	0.6	0.5
F15	5638329	320600	105	0.7	1.9	
Merzenhausen (ME)	F01	5645502	310014	105	7.7	0.6
Merken (MK)	F01	5636968	317442	93	0.7	0.7
	F02	5635985	316781	108	5.3	0.6
	F03	5636161	317011	116	6.1	0.4
	F04	5635973	317223	114	4.3	0.4
	F05	5635738	317217	115	1.1	0.5
Hürtgenwald (HW)	F01	5622785	314460	360	8.4	2.4
	F04	5621961	314387	373	6.8	1.1
	F05	5621879	314156	374	5.7	2.6

**Table 2: Abbreviations for sites and land-use types**

site	abbreviation
Selhausen	SE
Merken	MK
Merzenhausen	ME
Hürtgenwald	HW

land-use type	abbreviation
Catch crop	CC
Harvest residues*	HR
Maize	MA
Rapeseed	RA
Spelt	SP
Sugar beet	SB
Triticale	TC
Winter barley	WB
Winter wheat	WW

\* period before sowing and after harvest independent of the actual presence of residues on the field

**Table 3: Data availability for vegetation data, fluxes, and management data (“X” data available, “-“ no data available). For an explanation of vegetation data categories, refer to section 4. For crops, the year refers to harvest. Concerning vegetation data, number of points gives the maximum number of points in the field measured on the same date. In case of harvest residues, green sprouts and other biomass, data is only marked available if at least one value unequal zero is available.**

Dataset							Flux	Management						Vegetation									
Identifier	site	field	crop	variety / cultivar	year	preceding crop	EC data	sowing date	harvesting date	fertilization data	cultivation	sowing density	sowing depth	number of points (max)	number of dates	biomass per organ	aboveground biomass	LAI	C/N content	plant height	harvest residues (!= 0)	Green sprouts (!= 0)	Other (!= 0)
SEF08WW08	SE	F08	WW	Raspail	2008	SB	X	X	X	X	-	-	-	3	10	X	-	X	X	X	-	-	X
SEF01SB08	SE	F01	SB	-	2008	-	-	X	-	X	-	X	-	3	8	X	-	X	X	X	-	-	-
SEF14MA08	SE	F14	MA	-	2008	-	-	X	-	X	-	-	-	3	8	X	-	X	X	X	-	-	X
SEF11RA08	SE	F11	RA	-	2008	WB	-	X	-	X	-	X	-	3	9	X	-	X	X	X	-	-	X
SEF08WW09	SE	F08	WW	Raspail	2009	WW	X	X	X	X	X	X	-	3	14	X	-	X	X	X	-	-	-
SEF07SB09	SE	F07	SB	Pauletta	2009	-	-	X	X	X	X	X	X	3	11	X	-	X	X	X	-	-	-
SEF10MA09	SE	F10	MA	Agro Lux	2009	-	-	X	X	X	X	X	X	3	8	X	-	X	X	X	-	-	-
SEF13RA09	SE	F13	RA	-	2009	-	-	X	X	X	X	-	X	3	13	X	-	X	X	X	-	-	X
SEF15WB09	SE	F15	WB	Laverda	2009	-	-	-	-	-	-	-	-	3	8	X	-	X	X	X	-	-	-
MKF05MA09	MK	F05	MA	Ronaldinho	2009	-	-	X	X	X	X	-	-	3	7	X	-	X	X	X	-	-	-
MKF01RA09	MK	F01	RA	NK-Fair	2009	-	-	X	X	X	X	X	X	3	9	X	-	X	X	X	-	-	-
MKF04SB09	MK	F04	SB	Beretta KWS	2009	-	X*	X	X	X	X	X	X	3	10	X	-	X	X	X	-	-	-
MKF03WB09	MK	F03	WB	Fridericus	2009	-	X*	X	X	X	-	X	X	3	8	X	-	X	X	X	-	-	-
MKF02WW09	MK	F02	WW	Hattrick	2009	-	X*	X	X	X	X	-	X	3	10	X	-	X	X	X	-	-	-
SEF07WW10	SE	F07	WW	-	2010	SB	-	-	-	-	-	-	-	6	10	X	-	X	X	X	-	-	-
SEF08SB10	SE	F08	SB	Supero	2010	WW	-	X	X	X	X	-	X	7	12	X	-	X	X	X	-	-	-
SEF09MA10	SE	F09	MA	-	2010	-	-	-	-	-	-	-	-	3	1	X	-	X	-	X	-	-	-
SEF12RA10	SE	F12	RA	-	2010	-	-	-	-	-	-	-	-	5	8	X	-	X	X	X	-	-	-

SEF01WB10	SE	F01	WB	-	2010	-	-	-	-	-	-	-	5	8	X	-	X	-	X	-	-	-
MKF04WW10	MK	F04	WW	-	2010	SB	-	-	-	-	-	-	3	2	X	-	X	-	X	-	-	-
MKF02WB10	MK	F02	WB	-	2010	WW	-	-	-	-	-	-	3	2	X	-	X	-	X	-	-	-
MEF01HR11	ME	F01	HR	-	2011	-	X	-	-	-	-	-	12	1	-	-	-	X	-	X	-	-
MEF01WW11	ME	F01	WW	Potenzial	2011	SB	X	X	X	X	-	X	X	12	10	X	-	X	X	X	-	-
MEF01HR12	ME	F01	HR	-	2012	-	X	-	-	-	-	-	12	1	-	-	-	-	-	X	-	-
MEF01WW12	ME	F01	WW	Tobak	2012	WW	X	X	X	X	-	X	X	12	12	X	-	X	X	X	-	-
SEF04WW13	SE	F04	WW	-	2013	-	-	-	-	-	-	-	6	10	X	-	X	X	X	-	-	-
SEF01HR15	SE	F01	HR	-	2015	-	X	-	-	-	-	-	3	3	-	-	-	-	X	X	-	-
SEF01WW15	SE	F01	WW	Premio	2015	-	X	X	X	X	-	-	3	6	X	-	X	-	X	-	-	-
SEF03HR15	SE	F03	HR	-	2015	-	-	-	-	-	-	-	3	5	-	-	-	-	-	X	X	X
SEF03WW15	SE	F03	WW	-	2015	-	-	X	X	X	-	X	X	3	6	X	X	X	-	X	-	-
SEF02SB15	SE	F02	SB	-	2015	-	-	-	-	-	-	-	3	7	X	-	X	-	X	-	-	X
SEF04HR15	SE	F04	HR	-	2015	-	-	-	-	-	-	-	3	5	-	-	-	-	-	X	X	X
SEF04SP15	SE	F04	SP	-	2015	WW	-	-	-	-	-	-	3	4	X	-	X	-	X	-	-	-
SEF05HR15	SE	F05	HR	-	2015	-	-	-	-	-	-	-	3	3	-	-	-	-	-	X	X	-
SEF05WW15	SE	F05	WW	-	2015	-	-	-	-	-	-	-	3	4	X	X	X	-	X	-	-	-
HWF01HR15	HW	F01	HR	-	2015	-	-	-	-	-	-	-	3	2	-	-	-	-	X	X	-	-
HWF01MA15	HW	F01	MA	Silage maize	2015	-	-	X	X	X	X	X	3	7	X	-	X	-	X	-	-	X
HWF04HR15	HW	F04	HR	-	2015	-	-	-	-	-	-	-	0	1	-	-	-	-	-	-	-	-
HWF04TC15	HW	F04	TC	Winter TC	2015	-	-	X	X	X	X	-	X	3	3	X	X	X	-	X	-	X
SEF01WB16	SE	F01	WB	-	2016	WW	X	X	X	X	X	-	3	7	X	-	X	-	X	-	-	-
SEF01HR16	SE	F01	HR	-	2016	-	X	-	-	-	-	-	3	3	-	-	-	-	X	X	X	-
SEF01CC16	SE	F01	CC	-	2016	WB	X	X	X	X	X	-	3	2	-	X	-	-	X	-	-	-
SEF03HR16	SE	F03	HR	-	2016	WW	-	-	-	-	-	-	3	3	-	-	-	-	-	X	-	-
SEF03SB16	SE	F03	SB	-	2016	WW	-	-	-	-	-	-	3	7	X	-	X	-	X	-	-	-
SEF04HR16	SE	F04	HR	-	2016	-	-	-	-	-	-	-	0	2	-	-	-	-	-	-	-	-



SEF04SB16	SE	F04	SB	Kleist	2016	SP	-	X	-	X	X	X	X	3	7	X	-	X	-	X	-	-	-
SEF05HR16	SE	F05	HR	-	2016	WW	-	-	-	-	-	-	-	3	3	-	-	-	-	X	X	-	-
SEF05WW16	SE	F05	WW	-	2016	WW	-	-	-	-	-	-	-	3	7	X	X	X	-	X	-	-	-
SEF06HR16	SE	F06	HR	-	2016	-	-	-	-	-	-	-	-	4	2	-	-	-	-	X	X	-	-
SEF06WB16	SE	F06	WB	-	2016	-	-	-	-	-	-	-	-	4	7	X	-	X	-	X	-	-	-
HWF01HR16	HW	F01	HR	-	2016	-	-	-	-	-	-	-	-	3	4	-	-	-	-	X	X	-	-
HWF01MA16	HW	F01	MA	-	2016	MA	-	-	X	X	X	X	X	3	4	X	-	X	-	X	-	-	X
HWF05HR16	HW	F05	HR	-	2016	-	-	-	-	-	-	-	-	0	1	-	-	-	-	X	-	-	-
HWF05TC16	HW	F05	TC	-	2016	-	-	X	X	-	X	-	X	3	4	X	X	X	-	X	X	-	-
SEF06WB17	SE	F06	WB	-	2017	WB	-	-	-	-	-	-	-	6	1	X	-	X	-	X	-	-	-
MEF01HR17	ME	F01	HR	-	2017	-	X	-	-	-	-	-	-	3	1	-	-	-	-	X	-	-	-
MEF01WW17	ME	F01	WW	-	2017	-	X	X	-	-	X	-	-	3	7	X	-	X	-	X	-	-	X

\* Data from two heights

**Table 4: Quality flags set in case the sum of its components differed from the aliquot.**

	< 5 %	5 %–10 %	10 %–15 %	> 15 %
sum < aliquot	high quality (1)	good quality (2)	suspicious (4)	low quality (5)
sum > aliquot	high quality (1)	suspicious (4)	suspicious (4)	low quality (5)

**Table 5: Locations, processing software, and instrument heights, and temporal extent of eddy covariance stationsmeasurements. Coordinates are UTM zone 32N (WGS1984). For information on quality indicators see section 5.2.**

site	field	year	Identifier	UTM N	UTM E	elevation (m a.s.l.)	processing software	quality indicator	height (cm)*	<u>start</u> <sup>*2</sup> (y-m-d)	<u>end</u> <sup>*2</sup> (y-m-d)
SE	F08	2007	SEF08_SE_EC000_fluxes_2007	5638560	320543	103	TK2	flags	245	<u>2007-05-25</u>	<u>2007-12-25</u>
SE	F08	2008	SEF08_SE_EC000_fluxes_2008	5638560	320543	103	TK2	flags	245	<u>2008-01-08</u>	<u>2008-12-31</u>
SE	F08	2009	SEF08_SE_EC000_fluxes_2009	5638560	320543	103	TK2	flags	245	<u>2009-01-01</u>	<u>2009-12-31</u>
SE	F08	2010	SEF08_SE_EC000_fluxes_2010	5638560	320543	103	TK2	flags	245	<u>2010-01-01</u>	<u>2010-10-14</u>
SE	F01	2015	SEF01_SE_EC001_fluxes_2015	5638010	320380	103	TK3.1	flags	245	<u>2015-01-01</u>	<u>2015-12-31</u>
SE	F01	2016	SEF01_SE_EC001_fluxes_2016	5638010	320380	103	TK3.1	flags	245	<u>2016-01-01</u>	<u>2016-12-31</u>
ME	F01	2011	MEF01_ME_EC001_fluxes_2011	5645497	310059	93	TK3.1	flags	198	<u>2011-05-10</u>	<u>2011-12-31</u>
ME	F01	2012	MEF01_ME_EC001_fluxes_2012	5645497	310059	93	TK3.1	flags	198	<u>2012-01-01</u>	<u>2012-12-31</u>
ME	F01	2017	MEF01_ME_EC001_fluxes_2017	5645497	310059	93	TK3.1	flags	198	<u>2017-01-01</u>	<u>2017-12-31</u>
MK	F02	2009	MKF02_MK_ECJ11_fluxes_2009	5635998	316798	116	ECpack 2.5.20	tolerances	240	<u>2009-04-14</u>	<u>2009-08-27</u>
MK	F02	2009	MKF02_MK_ECJ1u_fluxes_2009	5635998	316798	116	ECpack 2.5.20	tolerances	590	<u>2009-04-07</u>	<u>2009-08-27</u>
MK	F03	2009	MKF03_MK_ECS4l_fluxes_2009	5636165	317010	114	ECpack 2.5.20	tolerances	260	<u>2009-04-07</u>	<u>2009-09-03</u>
MK	F03	2009	MKF03_MK_ECS4u_fluxes_2009	5636165	317010	114	ECpack 2.5.20	tolerances	596	<u>2009-04-21</u>	<u>2009-09-07</u>
MK	F04	2009	MKF04_MK_ECS3l_fluxes_2009	5635956	317204	115	ECpack 2.5.20	tolerances	248	<u>2009-04-09</u>	<u>2009-09-29</u>
MK	F04	2009	MKF04_MK_ECS3u_fluxes_2009	5635956	317204	115	ECpack 2.5.20	tolerances	604	<u>2009-04-23</u>	<u>2009-09-15</u>

\* height: instrument height of anemometer and IRGA (above ground)

\*2 first and last day with valid data, y:year, m: month, d: day

**Table 6: Flag values set by the TK software and their meanings.**

flag	meaning
0	high quality data, use in fundamental research possible
1	moderate quality data, no restrictions for use in long term observation programs
2	low data quality, gap filling necessary

**Table 7: Acceptable value ranges of ECpack results and tolerance-values at the lower and upper boundary. For the meaning of the group-column refer to the text.**

variable	lower boundary	tolerance at lower boundary	upper boundary	tolerance at upper boundary	group
Mean(u)	0	0.2	200	1	A
Mean(w)	0	0.05	20	0.1	A
Mean(TSon)	273.15	0.1	350	0.1	A
Mean(rhoV)	0	2E-4	0.2	2E-4	B
Mean(qCO2)	4E-4	1E-5	1E-3	1E-5	B
U_dir	0	180	360	180	A
Hsonic	0	25	1000	100	A
SumLvE	0	50	1000	200	B
Ustar	0	0.2	1.0	0.3	A
SumFCO2	0	5E-7	5E-6	1E-6	B

**Table 8: Availability of soil information per site. C/N: carbon and nitrogen content are not always both available. Due to the absence of carbonates, C-content is expected to equal SOC.**

Site	particle sizes fine soil	gravel	bulk density	SOC	C and/or N
HW	X	X	X	-	-
MK	X	-	-	-	X
ME	X	X	X	X	X
SE	X	X	X	-	X

**Table 9: Meteorological Stations, their positions, available data and data source**

ID	station	UTM northing	UTM easting	period used	AirHum	AirPres	AirTemp	Precip	Globrad	Wind	Cloud	data source
1	ME_BCK_001	5645555	310095	2011–2016	X	X		X	X	X		TEODOOR <sup>1</sup>
2	RO_AKRW_003	5611891	309102	2011–2016	X	X	X	X		X		TEODOOR <sup>1</sup>
3	RO_BKY_010	5611219	309322	2012–2016	X	X	X	X	X	X		TEODOOR <sup>1</sup>
4	RO_EC_001	5611250	309312	2011–2016	X	X	X	X	X	X		TEODOOR <sup>1</sup>
5	RU_BCK_002	5652036	312165	2011–2016	X	X	X	X	X	X		TEODOOR <sup>1</sup>
6	RU_BCK_003	5637669	318956	2011–2016	X	X	X	X		X		TEODOOR <sup>1</sup>
7	RU_BCK_004	5668397	301947	2012–2016	X	X	X	X		X		TEODOOR <sup>1</sup>
8	RU_BCDKR_001	5599172	313945	2011–2016	X	X	X	X	X	X		TEODOOR <sup>1</sup>
9	RU_K_002	5642873	317452	2013–2016	X	X	X	X	X	X		TEODOOR <sup>1</sup>
10	SE_BDK_999	5638335	320536	2009–2016	X	X	X	X	X	X		TEODOOR <sup>1,3</sup>
11	SE_EC_001	5638012	320375	2011–2016	X	X	X	X	X	X		TEODOOR <sup>1</sup>
12	WU_BKY_010	5597950	310540	2012–2016	X	X	X	X	X	X		TEODOOR <sup>1</sup>
13	WU_EC_002	5597955	311089	2013–2016	X	X	X		X	X		TEODOOR <sup>1</sup>
14	WU_K_002	5597960	311091	2014–2016	X	X	X	X		X		TEODOOR <sup>1</sup>
15	ME_EC_001	5645497	310059	2011–2016	X	X	X	X	X	X		TEODOOR <sup>1</sup>
16	RU_K_001	5643013	317883	2007–2016	X	X		X	X	X		TEODOOR <sup>1</sup>
17	RU_EC_001	5637813	318969	2011–2016	X	X	X		X	X		TEODOOR <sup>1</sup>
18	WU_EC_001	5598173	310739	2011–2014	X	X	X		X	X		TEODOOR <sup>1</sup>
19	SE_EC_002	5638375	320591	2010	X	X	X	X	X	X		TEODOOR <sup>1</sup>
20	HW_BK_001	5622292	314567	2015–2016	X	X	X	X	X	X		GLOBE <sup>4</sup>
21	HW_BK_002	5621923	314600	2016	X	X	X	X	X	X		GLOBE <sup>4,5</sup>
22	10501	5629698	295161	2007–2010		X					X	DWD <sup>6,7</sup>
23	10505	5631617	290318	2011–2016							X	DWD <sup>6,8</sup>
24	H827	5616739	317991	2014–2015	X		X	X		X		DWD <sup>6</sup>
25	SE_EC_000	5638537	320558	2007–2009	X	X	X	X	X	X		TR32DB <sup>2</sup>

<sup>1</sup> <http://teodoor.icg.kfa-juelich.de/ibg3searchportal2/>, Eifel/Lower Rhine Valley Observatory<sup>2</sup> <http://www.tr32db.de><sup>3</sup> includes data from stations SE\_BK\_001 and SE\_BDK\_002 from TEODOOR<sup>1</sup><sup>4</sup> <https://datasearch.globe.gov/><sup>5</sup> consists of three stations

---

<sup>6</sup> Deutscher Wetterdienst, DWD, [ftp://ftp-cdc.dwd.de/pub/CDC/observations\\_germany/climate/hourly/](ftp://ftp-cdc.dwd.de/pub/CDC/observations_germany/climate/hourly/)

<sup>7</sup> DWD station Aachen, old location

<sup>8</sup> DWD station Aachen, new location



**Table 10: Source of meteorological data given as station IDs as defined in Table 9. Station IDs in parenthesis are stations used for gap-filling. Colum “Met” shows the method used for gap-filling as explained in the text.**

year	AirPres	GlobRad	AirTemp	AirHum	Wind	Precip	Cloud	Met
Hürtgenwald (HW)								
2014	4 (24)	4 (24)	4 (24)	4 (24)	4 (24)	4 (24)	23	1
2015	20 (4, 24)	20 (4, 24)	20 (4, 24)	20 (4, 24)	20 (4, 24)	20 (4, 24)	23	1
2016	20 (4)	20 (21)	20 (21)	20 (21)	20 (21)	20 (21)	23	1
Selhausen (SE)								
2007	25 (22)	22	25 (16)	25 (16)	25 (16)	16	22	0
2008-2009	25 (22)	22	25 (16)	25 (16)	25 (10,16)	16	22	0
2010	22	11 (19, 10, 16)	11 (19, 10, 16)	10 (16)	11 (19, 10, 16)	16	22	2
2011-2017	10 (1–18)	11 (1–18)	10 (1–18)	10 (1–18)	11 (1–18)	10 (1–18)	23	3
Merzenhausen (ME)								
2009-2010	22	15 (1, 16)	15 (1, 16)	15 (1, 16)	15 (1, 16)	15 (1, 16)	22	2
2011-2017	1 (2–18)	15 (1–18)	1 (2–18)	1 (2–18)	1 (2–18)	1 (2–18)	23	3

**Table 11: Instrument uncertainties for meteorological measurements at stations 1 to 21 and 25 (for station IDs compare Table 9).**

variable	Uncertainty
Air pressure (Pa)	p_ref (flux data): 1 % for relative humidity AirPres (weather data): 0 to 30°C: $\pm 0.5$ hPa; -52 to +60°C: $\pm 1$ hPa
global radiation ( $\text{W m}^{-2}$ )	$\pm 5$ % to $\pm 10$ % for daily sums
Air temperature (°C)	$\pm 0.2 - 0.4$ °C
Air humidity (%)	Accuracy at 20°C: $\pm 2$ % RH (0 to 90 % Relative Humidity); $\pm 3$ % RH (90 to 100 % Relative Humidity)
Wind speed ( $\text{m s}^{-1}$ )	Offset error: $< \pm 8.0 \text{ cm s}^{-1}$ (u, v), $< \pm 4.0 \text{ cm s}^{-1}$ (z) Gain Error: Wind Vector within $\pm 5^\circ$ of horizontal: $< \pm 2\%$ of reading Wind Vector within $\pm 10^\circ$ of horizontal: $< \pm 3\%$ of reading Wind Vector within $\pm 20^\circ$ of horizontal: $< \pm 6\%$ of reading
Precipitation ( $\text{mm h}^{-1}$ )	$< 3$ %
Cloudiness (1/8)	Uncertainty unknown
Wind direction (°)*	$\pm 0.7^\circ$ at $1 \text{ m s}^{-1}$ for horizontal wind

\* included in flux datafiles

**Table 12: Uncertainty estimates for weather timeseries gap-filling expressed as pairs of bias (B) and RMSE (R). Missing data is denoted by NA; “-“ marks cases where there were no gaps or no data. Methods are described in the text. For abbreviations see Table A7.**

Period	AirPres		GlobRad		AirTemp		AirHum		Wind		Precip		InLW	
	(Pa)		(W m <sup>-2</sup> )		(K)		(%)		(m s <sup>-1</sup> )		(mm)		(W m <sup>-2</sup> )	
	B	R	B	R	B	R	B	R	B	R	B	R	B	R
2007 to 2009	24.7	27.2	-	-	0.1	0.9	-5.7	9.5	-0.4	0.9	NA	NA	-	-
2010	-	-	NA	NA	-0.3	0.8	-5.7	8.0	0.1	0.8	NA	NA	-	-
2011 to 2017	-7.6	12.2	0.6	78.8	0.1	0.9	0.3	1.2	-1.6	2.5	-0.1	0.3	-5.8	24.6

**Table A1: Columns in the vegetation data files**

#Col.	variable	units	data type	description
1	dataset	-	character	Dataset name
2	test_site	-	character	Site ID
3	field	-	character	Field ID (per site)
4	land_use	-	character	Land-use ID
5	Date	-	date	Date of field measurement (YYYY-MM-DD)
6	time	-	time	UTC-time of field measurement (hh:mm)
7	UTM_northing	m	numeric	UTM Northing (WGS84, 32N)
8	UTM_northing_FLAG	-	numeric	UTM Northing quality flag
9	UTM_easting	m	numeric	UTM Easting (WGS84, 32N)
10	UTM_easting_FLAG	-	numeric	UTM Easting quality flag
11	canopy_height	cm	numeric	Height of the canopy
12	bbch	BBCH*	character	Phenological development state (BBCH scale)
13	num_plants_m2	m <sup>-2</sup>	numeric	Number of plants per m <sup>2</sup> (calculated)
14	LAI_green	m <sup>2</sup> m <sup>-2</sup>	numeric	Green LAI
15	LAI_green_FLAG	-	numeric	Green LAI quality flag
16	LAI_brown	m <sup>2</sup> m <sup>-2</sup>	numeric	Brown LAI
17	LAI_brown_FLAG	-	numeric	Brown LAI quality flag
18	FW_green_leaves	g m <sup>-2</sup>	numeric	Fresh weight of green leaves
19	FW_green_leaves_FLAG	-	numeric	Fresh weight of green leaves quality flag
20	DW_green_leaves	g m <sup>-2</sup>	numeric	Dry weight of green leaves
21	DW_green_leaves_FLAG	-	numeric	Dry weight of green leaves quality flag
22	FW_brown_leaves	g m <sup>-2</sup>	numeric	Fresh weight of brown leaves
23	FW_brown_leaves_FLAG	-	numeric	Fresh weight of brown leaves quality flag
24	DW_brown_leaves	g m <sup>-2</sup>	numeric	Dry weight of brown leaves
25	DW_brown_leaves_FLAG	-	numeric	Dry weight of brown leaves quality flag
26	FW_green_stems	g m <sup>-2</sup>	numeric	Fresh weight of green stems/tillers/stalks
27	FW_green_stems_FLAG	-	numeric	Fresh weight of green stems/tillers/stalks quality flag
28	DW_green_stems	g m <sup>-2</sup>	numeric	Dry weight of green stems/tillers/stalks

29	DW_green_stems_FLAG	-	numeric	Dry weight of green stems/tillers/stalks quality flag
30	FW_brown_stems	g m <sup>-2</sup>	numeric	Fresh weight of brown stems/tillers/stalks
31	FW_brown_stems_FLAG	-	numeric	Fresh weight of brown stems/tillers/stalks quality flag
32	DW_brown_stems	g m <sup>-2</sup>	numeric	Dry weight of brown stems/tillers/stalks
33	DW_brown_stems_FLAG	-	numeric	Dry weight of brown stems/tillers/stalks quality flag
34	FW_fruit	g m <sup>-2</sup>	numeric	Fresh weight of harvest organ (e.g. fruit, beet)
35	FW_fruit_FLAG	-	numeric	Fresh weight of harvest organ quality flag
36	DW_fruit	g m <sup>-2</sup>	numeric	Dry weight of harvest organ (e.g. fruit, beet)
37	DW_fruit_FLAG	-	numeric	Dry weight of harvest organ quality flag
38	FW_biomass_undiff	g m <sup>-2</sup>	numeric	Fresh weight of aboveground biomass not separated into organs
39	FW_biomass_undiff_FLAG	-	numeric	Fresh weight of aboveground biomass not separated into organs quality flag
40	DW_biomass_undiff	g m <sup>-2</sup>	numeric	Dry weight of aboveground biomass not separated into organs
41	DW_biomass_undiff_FLAG	-	numeric	Dry weight of aboveground biomass not separated into organs quality flag
42	FW_harvest_residues	g m <sup>-2</sup>	numeric	Fresh weight of harvest residues
43	FW_harvest_residues_FLAG	-	numeric	Fresh weight of harvest residues quality flag
44	DW_harvest_residues	g m <sup>-2</sup>	numeric	Dry weight of harvest residues
45	DW_harvest_residues_FLAG	-	numeric	Dry weight of harvest residues quality flag
46	FW_green_sprouts	g m <sup>-2</sup>	numeric	Fresh weight of green sprouts (growing between harvest residues)
47	FW_green_sprouts_FLAG	-	numeric	Fresh weight of green sprouts quality flag
48	DW_green_sprouts	g m <sup>-2</sup>	numeric	Dry weight of green sprouts (growing between harvest residues)
49	DW_green_sprouts_FLAG	-	numeric	Dry weight of green sprouts quality flag

50	FW_other	g m <sup>-2</sup>	numeric	Fresh weight of other biomass (e.g. weeds)
51	FW_other_FLAG	-	numeric	Fresh weight of other biomass quality flag
52	DW_other	g m <sup>-2</sup>	numeric	Dry weight of other biomass (e.g. weeds)
53	DW_other_FLAG	-	numeric	Dry weight of other biomass quality flag
54	other_descr	-	character	type of biomass measured as "biomass_other"
55	N_green_leaves	mass%	numeric	Relative nitrogen content of green leaves
56	C_green_leaves	mass%	numeric	Relative carbon content of green leaves
57	N_brown_leaves	mass%	numeric	Relative nitrogen content of brown leaves
58	C_brown_leaves	mass%	numeric	Relative carbon content of brown leaves
59	N_green_stems	mass%	numeric	Relative nitrogen content of green stems/tillers/stalks
60	C_green_stems	mass%	numeric	Relative carbon content of green stems/tillers/stalks
61	N_brown_stems	mass%	numeric	Relative nitrogen content of brown stems/tillers/stalks
62	C_brown_stems	mass%	numeric	Relative carbon content of brown stems/tillers/stalks
63	N_fruit	mass%	numeric	Relative nitrogen content of harvest organ (e.g. fruit, beet)
64	C_fruit	mass%	numeric	Relative carbon content of harvest organ (e.g. fruit, beet)
65	N_biomass_undiff	mass%	numeric	Relative nitrogen content of aboveground biomass not separated into organs
66	C_biomass_undiff	mass%	numeric	Relative carbon content of aboveground biomass not separated into organs
67	N_harvest_residues	mass%	numeric	Relative nitrogen content of harvest residues
68	C_harvest_residues	mass%	numeric	Relative carbon content of harvest residues
69	N_green_sprouts	mass%	numeric	Relative nitrogen content of green sprouts (growing between harvest residues)

70	C_green_sprouts	mass%	numeric	Relative carbon content of green sprouts (growing between harvest residues)
71	N_other	mass%	numeric	Relative nitrogen content of other biomass (e.g. weeds)
72	C_other	mass%	numeric	Relative carbon content of other biomass (e.g. weeds)
73	is_cn_field_mean	-	logical	Have C- and N-contents been measured from a composite sampled from all points in the field?
74	soil_unit	-	character	Assignment to a soil unit of Brogi et al. (2019), only Selhausen
75	comment	-	character	Comment

---

\* see Meier et al. (2009) and references therein

**Table A2: Columns of flux datafiles processed with the software ECPack. With the exception of the timestamps, all datatypes are numeric.**

#Col.	variable	Units	description
1	Datetime(end)	-	UTC-time end of interval (YYYY-MM-DD hh:mm)
2	#Samples	-	Number of records aggregated to data in the current row
3	Mean(u)	m s <sup>-1</sup>	horizontal wind component (coordinate system turned into mean wind)
4	TolMean(u)	m s <sup>-1</sup>	Estimate of 95% confidence intervals for horizontal wind component u
5	Mean(v)	m s <sup>-1</sup>	horizontal wind component orthogonal to v (almost zero due to rotation of coordinate system)
6	TolMean(v)	m s <sup>-1</sup>	Estimate of 95% confidence intervals for horizontal wind component v
7	Mean(w)	m s <sup>-1</sup>	vertical wind (after planar fit rotation)
8	TolMean(w)	m s <sup>-1</sup>	Estimate of 95% confidence intervals for vertical wind speed
9	Mean(TSon)	K	air temperature, calculated from sonic temperature, pressure and H <sub>2</sub> O density
10	TolMean(TSon)	K	Estimate of 95% confidence intervals for air temperature
11	Mean(rhoV)	kg m <sup>-3</sup>	H <sub>2</sub> O density
12	TolMean(rhoV)	kg m <sup>-3</sup>	Estimate of 95% confidence intervals for average H <sub>2</sub> O density
13	Mean(qCO2)	kg kg <sup>-1</sup>	CO <sub>2</sub> mixing ratio
14	TolMean(qCO2)	kg kg <sup>-1</sup>	Estimate of 95% confidence intervals for average CO <sub>2</sub> mixing ratio
15	Std(u)	m s <sup>-1</sup>	Standard deviation of horizontal wind component u
16	TolStd(u)	m s <sup>-1</sup>	Estimate of 95% confidence intervals for horizontal wind component u
17	Std(v)	m s <sup>-1</sup>	Standard deviation of horizontal wind component v
18	TolStd(v)	m s <sup>-1</sup>	Estimate of 95% confidence intervals for horizontal wind component v
19	Std(w)	m s <sup>-1</sup>	Standard deviation of vertical wind speed
20	TolStd(w)	m s <sup>-1</sup>	Estimate of 95% confidence intervals for vertical wind speed
21	Std(TSon)	K	Standard deviation of sonic temperature



---

22	TolStd(TSon)	K	Estimate of 95% confidence intervals for standard deviation of air temperature
23	Std(q)	kg kg <sup>-1</sup>	Standard deviation of specific humidity
24	TolStd(q)	kg kg <sup>-1</sup>	Estimate of 95% confidence intervals for specific humidity
25	Std(qCO2)	kg kg <sup>-1</sup>	Standard deviation of CO <sub>2</sub> mixing ratio
26	TolStd(qCO2)	kg kg <sup>-1</sup>	Estimate of 95% confidence intervals for standard deviation of average CO <sub>2</sub> mixing ratio
27	Cov(u*v)	m <sup>2</sup> s <sup>-2</sup>	Covariance of wind components u and v
28	TolCov(u*v)	m <sup>2</sup> s <sup>-2</sup>	Estimate of 95% confidence intervals for covariance of wind components u and v
29	Cov(v*w)	m <sup>2</sup> s <sup>-2</sup>	Covariance of wind components v and w
30	TolCov(u*w)	m <sup>2</sup> s <sup>-2</sup>	Estimate of 95% confidence intervals for covariance of wind components u and w
31	Cov(u*w)	m <sup>2</sup> s <sup>-2</sup>	Covariance of wind components u and w
32	TolCov(v*w)	m <sup>2</sup> s <sup>-2</sup>	Estimate of 95% confidence intervals for covariance of wind components v and w
33	RhoSon	kg m <sup>-3</sup>	Air density from the ultrasonic anemometer
34	Tol(RhoSon)	kg m <sup>-3</sup>	Estimate of 95% confidence intervals for air density from the ultrasonic anemometer
35	U_vect	m s <sup>-1</sup>	In this processing scheme, identical to Mean(u)
36	Tol(U_vect)	m s <sup>-1</sup>	Estimate of 95% confidence intervals for U_vect
37	U_dir	°	Wind direction in geographical coordinate system
38	Tol(U_dir)	°	Estimate of 95% confidence intervals for wind direction in geographical coordinate system
39	HSonic	W/m <sup>2</sup>	Sensible heat flux including planar fit, Moore and Schotanus correction
40	Tol(HSonic)	W m <sup>-2</sup>	Estimate of 95% confidence intervals for sensible heat flux
41	SumLvE	W m <sup>-2</sup>	Latent heat flux including planar fit, Moore and WPL correction
42	Tol(SumLvE)	W m <sup>-2</sup>	Estimate of 95% confidence intervals for latent heat flux
43	Ustar	m s <sup>-1</sup>	Friction velocity including planar fit and Moore correction
44	Tol(Ustar)	m s <sup>-1</sup>	Estimate of 95% confidence intervals for friction velocity
45	SumFCO2	kg m <sup>-2</sup> s <sup>-1</sup>	CO <sub>2</sub> flux without consideration of storage flux

---

---

46 Tol(SumFCO2)

kg m<sup>-2</sup> s<sup>-1</sup>

Estimate of 95% confidence intervals for CO<sub>2</sub> flux without  
consideration of storage flux

---

**Table A3: Columns of flux datafiles processed with the software TK. With the exception of the timestamps, all datatypes are numeric.**

#Col.	variable	units	Description
1	T_begin	-	UTC-time beginning of interval (YYYY-MM-DD hh:mm)
2	T_end	-	UTC-time end of interval (YYYY-MM-DD hh:mm)
3	u	m s <sup>-1</sup>	Horizontal wind speed (coordinate system turned into mean wind)
4	v	m s <sup>-1</sup>	Horizontal wind speed (zero due to rotation of coordinate system)
5	w	m s <sup>-1</sup>	Vertical wind speed
6	Ts	°C	Sonic temperature
7	Tp	°C	[no data]
8	a	g m <sup>-3</sup>	H <sub>2</sub> O content of the air (LI7500)
9	CO2	mmol m <sup>-3</sup>	CO <sub>2</sub> content of the air
10	T_ref	°C	Air temperature
11	a_ref	g m <sup>-3</sup>	Reference H <sub>2</sub> O content of the air (HMP45C)
12	p_ref	hPa	Air pressure
13	Var[u]	m <sup>2</sup> s <sup>-2</sup>	Variance of horizontal wind speed
14	Var[v]	m <sup>2</sup> s <sup>-2</sup>	Variance of horizontal wind speed
15	Var[w]	m <sup>2</sup> s <sup>-2</sup>	Variance of vertical wind speed
16	Var[Ts]	°C <sup>2</sup>	Variance of sonic temperature
17	Var[Tp]	°C <sup>2</sup>	[no data]
18	Var[a]	g <sup>2</sup> m <sup>-6</sup>	Variance of H <sub>2</sub> O content of the air
19	Var[CO2]	mmol <sup>2</sup> m <sup>-6</sup>	Variance of CO <sub>2</sub> content of the air
20	Cov[u'v']	m <sup>2</sup> s <sup>-2</sup>	Covariance of wind components u and v
21	Cov[v'w']	m <sup>2</sup> s <sup>-2</sup>	Covariance of wind components v and w
22	Cov[u'w']	m <sup>2</sup> s <sup>-2</sup>	Covariance of wind components u and w
23	Cov[u'Ts']	°C m s <sup>-1</sup>	Covariance of wind component u and sonic temperature
24	Cov[v'Ts']	°C m s <sup>-1</sup>	Covariance of wind component v and sonic temperature
25	Cov[w'Ts']	°C m s <sup>-1</sup>	Covariance of wind component w and sonic temperature
26	Cov[u'Tp']	°C m s <sup>-1</sup>	[no data]
27	Cov[v'Tp']	°C m s <sup>-1</sup>	[no data]
28	Cov[w'Tp']	°C m s <sup>-1</sup>	[no data]
29	Cov[u'a']	g s <sup>-1</sup> m <sup>-2</sup>	Covariance of wind component u and H <sub>2</sub> O content of the air
30	Cov[v'a']	g s <sup>-1</sup> m <sup>-2</sup>	Covariance of wind component v and H <sub>2</sub> O content of the air
31	Cov[w'a']	g s <sup>-1</sup> m <sup>-2</sup>	Covariance of wind component w and H <sub>2</sub> O content of the air

---

32	Cov[u'CO2']	mmol m <sup>-2</sup> s <sup>-1</sup>	Covariance of wind component u and CO <sub>2</sub> content of the air
33	Cov[v'CO2']	mmol m <sup>-2</sup> s <sup>-1</sup>	Covariance of wind component v and CO <sub>2</sub> content of the air
34	Cov[w'CO2']	mmol m <sup>-2</sup> s <sup>-1</sup>	Covariance of wind component w and CO <sub>2</sub> content of the air
35	Nvalue	-	Number of samples the aggregated 30-min-value is based on
36	dir	°	Wind direction
37	ustar	m s <sup>-1</sup>	Friction velocity
38	HTs	W m <sup>-2</sup>	Sensible heat flux
39	HTp	W m <sup>-2</sup>	[no data]
40	LvE	W m <sup>-2</sup>	Latent heat flux
41	z/L	-	Stability parameter (positive values denote stable boundary layer) based on sonic temperature
42	z/L-virt	-	Stability parameter (positive values denote stable boundary layer) based on virtual temperature
43	Flag(ustar)	-	Quality flag for ustar time series. Refer to the flag info in the "general info" sheet
44	Flag(HTs)	-	Quality flag for sensible heat time series. Refer to the flag info in the "general info" sheet
45	Flag(HTp)	-	[no data]
46	Flag(LvE)	-	Quality flag for latent heat time series. Refer to the flag info in the "general info" sheet
47	Flag(wCO2)	-	Quality flag for NEE time series. Refer to the flag info in the "general info" sheet
48	T_mid	-	UTC-time middle of interval (YYYY-MM-DD hh:mm)
49	FCstor	mmol m <sup>-2</sup> s <sup>-1</sup>	CO <sub>2</sub> storage of the air column below the measurement height
50	NEE	mmol m <sup>-2</sup> s <sup>-1</sup>	Net ecosystem exchange of CO <sub>2</sub>
51	Ftprint_trgt_1	%	Cumulative source contribution of the target area
52	Ftprint_trgt_2	%	Cumulative source contribution of adjacent areas of the same type as the target area
53	Ftprnt_xmax	m	Distance between EC-tower and the point of the maximum source contribution
54	r_err_ustar	%	Relative random error of ustar
55	r_err_HTs	%	Relative random error of HTs

---

---

56	r_err_LvE	%	Relative random error of LvE
57	r_err_co2	%	Relative random error of CO <sub>2</sub> flux
58	noise_ustar	%	Relative noise error of ustar
59	noise_HTs	%	Relative noise error of HTs
60	noise_LvE	%	Relative noise error of LvE
61	noise_co2	%	Relative noise error of CO <sub>2</sub> flux

---

**Table A4: Size ranges of particle size classes (DIN 14688)**

particle size class	abbreviation	size range (mm)
coarse material / gravel	Gr	> 2
coarse sand	CSa	0.63–2
medium sand	MSa	0.2–0.63
fine sand	FSa	0.063–0.2
sand	Sa	0.063–2
coarse silt	CSi	0.02–0.063
medium silt	MSi	0.0063–0.02
fine silt	FSi	0.002–0.0063
silt	Si	0.002–0.063
clay	Cl	≤ 0.002

**Table A5: Columns of soil datafiles. For particle size classes see Table A4.**

#Col.	variable	units	datatype	Description
1	Site	-	character	Site ID
2	Field	-	character	Field ID (per site)
3	UTM_northing	m	numeric	UTM Northing (WGS84, 32N)
4	UTM_easting	m	numeric	UTM Easting (WGS84, 32N)
5	Depth	cm	character	Sampling depth (layer)
6	horizon	-	character	Soil horizon (see Schad et al., 2009)
7	CSa	mass%	numeric	Percentage of coarse sand particles in fine soil
8	MSa	mass%	numeric	Percentage of medium sand particles in fine soil
9	FSa	mass%	numeric	Percentage of fine sand particles in fine soil
10	Sa	mass%	numeric	Percentage of sand particles in fine soil (CSa+MSa+FSa)
11	CSi	mass%	numeric	Percentage of coarse silt particles in fine soil
12	MSi	mass%	numeric	Percentage of medium silt particles in fine soil
13	FSi	mass%	numeric	Percentage of fine silt particles in fine soil
14	Si	mass%	numeric	Percentage of silt particles in fine soil (GSi+MSi+FSi)
15	Cl	mass%	numeric	Percentage of clay particles in fine soil
16	date_part_siz	-	date and time	sampling date for particle size distribution (in the field, YYYY-MM-DD)
17	Gr	mass%	numeric	Percentage of coarse material / gravel in soil sample
18	bulk_dens	g cm <sup>-3</sup>	numeric	Bulk density
19	date_bulk_dens	-	date and time	sampling date for bulk density (in the field, YYYY-MM-DD)
20	SOC	mass%	numeric	Soil organic carbon content
21	tot_C	mass%	numeric	Total carbon content
22	tot_N	mass%	numeric	Total nitrogen content
23	date_CN	-	date and time	Sampling date for C- and N-content (in the field, YYYY-MM-DD)
24	soil_unit	-	character	Assignment to a soil unit of Brogi et al. (2019), only Selhausen
25	comment	-	character	Comment

**Table A6: Columns of the soil units datafile. This data exists for the site Selhausen only. For particle size classes see Table A4.**

#Col.	variable	units	datatype	description
1	soil_unit	-	character	Assignment to a soil unit of Brogi et al. (2019)
2	horizon	-	character	Soil horizon (see Schad et al., 2009)
3	max_depth	cm	numeric	Maximum depth of the soil horizon found in the corresponding soil unit
4	Sa	mass%	numeric	Percentage of sand particles in fine soil
5	Si	mass%	numeric	Percentage of silt particles in fine soil
6	Cl	mass%	numeric	Percentage of clay particles in fine soil
7	Gr	mass%	numeric	Percentage of coarse material / gravel in soil sample
8	tot_C	mass%	numeric	Total carbon content
9	tot_N	mass%	numeric	Total nitrogen content
10	CV_max_depth	%	numeric	Uncertainty of max_depth (coefficient of variation)
11	CV_Sa	%	numeric	Uncertainty of Sa (coefficient of variation)
12	CV_Si	%	numeric	Uncertainty of Si (coefficient of variation)
13	CV_Cl	%	numeric	Uncertainty of Cl (coefficient of variation)
14	CV_Gr	%	numeric	Uncertainty of Gr (coefficient of variation)



**Table A7: Columns of weather datafiles. With the exception of the timestamps, all datatypes are numeric.**

#Col.	variable	units	Description
1	Date & Time begin (UTC)	-	UTC-time beginning of interval (YYYY-MM-DD hh:mm)
2	Date & Time end (UTC)	-	UTC-time end of interval (YYYY-MM-DD hh:mm)
3	AirPres	Pa	Air pressure
4	GlobRad	W m <sup>-2</sup>	Global Radiation
5	AirTemp	K	Air temperature
6	AirHum	%	Relative humidity of the air
7	Wind	m s <sup>-1</sup>	Wind speed
8	Precip	Mm	Precipitation
9	SurfaceTemp	K	Surface temperature*
10	InLW	W m <sup>-2</sup>	Incoming longwave radiation
11	Cloudiness	1/8	Cloud cover

\* contains no data, included for compatibility purposes