



# 1 **Glacial lake inventory of High Mountain Asia (1990–2018)** 2 **derived from Landsat images**

3 Xin Wang<sup>1,2</sup>, Xiaoyu Guo<sup>1</sup>, Chengde Yang<sup>2</sup>, Qionghuan Liu<sup>3</sup>, Junfeng Wei<sup>1</sup>, Yong Zhang<sup>1</sup>,  
4 ShiyinLiu<sup>4</sup>, Ynalín Zhang<sup>1</sup>, Zongli Jiang<sup>1</sup>, Zhiguang Tang<sup>1</sup>

5 <sup>1</sup>School of Resource Environment and Safety Engineering, Hunan University of Science and Technology,  
6 Xiangtan, 411100, China

7 <sup>2</sup>State Key Laboratory of Cryospheric Science, Northwest Institute of Ecology and Environmental  
8 Resources, Chinese Academy of Sciences, Lanzhou 730000, China

9 <sup>3</sup>Key Laboratory of Land Surface Pattern and Simulation, Institute of Geographic Sciences and Natural  
10 Resources Research, Chinese Academy of Sciences, Beijing 100101, China

11 <sup>4</sup>Institute of International Rivers and Eco-security, Yunnan University, Kunming, 650000, China

12 *Correspondence to:* Xin Wang (xinwang\_hn@163.com)

13 **Abstract.** There is currently no glacial lake inventory data set for the entire High Mountain Asia (HMA)  
14 area. The definition and classification of glacial lakes remain controversial, presenting certain obstacles  
15 to extensive utilization of glacial lake inventory data. This study integrated glacier inventory data and  
16 426 Landsat TM/ETM+/OLI images, and adopted manual visual interpretation to extract glacial lake  
17 boundaries within a 10-km buffer from glacier terminals using ArcGIS and ENVI software, normalized  
18 difference water index maps, and Google Earth images. The theoretical and methodological basis for all  
19 processing steps including glacial lake definition and classification, lake boundary delineation, and error  
20 assessment are discussed comprehensively in the paper. Moreover, detailed information regarding the  
21 coding, location, perimeter and area, area error, type, time phase, source image information, and  
22 sub-regions of the located lakes is presented. It was established that 26,089 and 28,953 glacial lakes in  
23 HMA, with sizes of 0.0054–5.83 km<sup>2</sup>, covered a combined area of 1692.74 ± 231.44 and 1955.94 ±  
24 259.68 km<sup>2</sup> in 1990 and 2018, respectively. The data set now is available at the National Special  
25 Environment and Function of Observation and Research Stations Shared Service Platform (China) at  
26 <http://dx.doi.org/10.12072/casnw.064.2019.db> (Wang et al., 2019a).

## 27 **1 Introduction**

28 Under the background of climate warming and the consequent widespread mass loss of glaciers in  
29 alpine regions, increasing volumes of glacial meltwater are being released. This results in glacial lake  
30 expansion and extended areas of low-lying terrain (e.g., depressions and troughs) left behind by



1 retreating glaciers in which water can accumulate and new glacial lakes can form (Clague and Evans,  
2 2000; Mool et al., 2001; Song et al., 2016). As both a water resource and a source of flash flood/debris  
3 flow hazards, glacial lakes participate in several natural processes, e.g., regional energy and water cycles  
4 (Slemmons et al., 2013), act as both indicators and containers of environmental information (Wang et al.,  
5 2016, 2019b), and drive hillslope erosion and landscape evolution (Cook et al., 2018) in the alpine  
6 cryosphere. On the one hand, glacial lakes act as temporary storage for the meltwater resource because  
7 a considerable amount of meltwater is retained by glacial lake expansion, e.g., approximately  $0.2 \% a^{-1}$   
8 of the total glacial meltwater was reserved in the glacial lakes from 1990 to 2010 in the Tien Shan  
9 Mountains in Central Asia (Wang et al., 2013). On the other hand, given the worldwide expansion in  
10 lake area in recent decades, the potential will increase for glacial lakes to develop into glacial lake  
11 outburst floods and related debris flows that could threaten downstream residents, infrastructure, and  
12 regional ecological and environmental security (Huggel et al., 2002; ICMOD, 2011; Bolch et al., 2012;  
13 Haebeli et al., 2016). Thus, glacial lakes perform important roles both in the meltwater cycle and in  
14 glacier hazard evolution in the cryosphere.

15 Following the rapid development of remote sensing technology and computer science, remote  
16 sensing imagery acquired by various satellites and sensors has been used widely in glacial lake research.  
17 In particular, Landsat imagery has become the most important data source for dynamic investigation of  
18 glacial lakes because of its wide coverage, continuous and long-term temporal sequence, and  
19 accessibility. Based on remote sensing data, both the distribution and the characteristics of change of  
20 glacial lakes in the mountains and watersheds in the High Mountain Asia (HMA) region have been  
21 widely reported. For example, multi-source remote sensing imagery has been used to compile glacial  
22 lake inventories for regions of the Tibetan Plateau (Zhang et al., 2015), Tien Shan Mountains (Wang et  
23 al., 2013), Himalaya (Gardelle et al., 2011; Nie et al., 2017), Hengduan Mountains (Wang et al., 2017),  
24 Uzbekistan (Petrov et al., 2017), and Pakistan (Senese et al., 2018). These inventories have proved an  
25 important data resource both for recognizing the spatiotemporal characteristics of glacial lakes and for  
26 understanding the response of glacial lakes to the effects of climate change in these regions.

27 Automatic and semi-automatic glacial lake boundary vectorization approaches have been used  
28 most widely in regional glacial lake investigations because of their higher efficiency and objectivity in  
29 comparison with manual visual vectorization. In such research, water bodies are usually determined  
30 based on the characteristics of different remote sensing bands and computer-dependent algorithms, e.g.,  
31 the normalized difference water index (NDWI), band ratio, support vector machine, decision tree,



1 spectral transformation, object-oriented classification, global–local iterative scheme, active contour  
2 model, and random forest (Gardelle et al., 2011, Huggel et al., 2002, Li et al., 2011; Veh et al., 2018,  
3 Zhang et al., 2018). However, manual post-processing is often required to calibrate the uncertainties  
4 that could easily be produced by the above approaches. Furthermore, the labour costs associated with  
5 rectification of lake boundary errors increase sharply with increasing complexity of study area terrain  
6 (Yang et al., 2019). With consideration of the accuracy, efficiency, and time overheads associated with  
7 the various vectorization approaches, a manual vectorization approach was adopted for investigation of  
8 the glacial lakes on the Tibetan Plateau (Zhang et al., 2015) despite the labour requirements and  
9 anticipated additional errors produced by individual subjectivity (Nie et al., 2017; Yang et al., 2019;  
10 Song et al., 2014).

11 Controversies and knowledge gaps remain regarding available glacial lake inventories for different  
12 alpine cryosphere regions, which present certain obstacles to extensive utilization of glacial lake  
13 inventory data. The main problems relate to regional differences in lake development, inconsistent  
14 specifications of lake definition, and the adoption of various approaches to lake interpretation (Yao et  
15 al., 2018). There is no existing comprehensive glacial lake inventory for the entire HMA and  
16 knowledge regarding the spatiotemporal characteristics of glacial lakes in this region is incomplete.  
17 The objectives of this study were to fill this knowledge gap by producing a glacial lake inventory data set  
18 for HMA derived from a Landsat imagery series, and to provide fundamental data for water resource  
19 evaluation, assessment of glacial lake outburst floods, and glacier hydrology research in the mountain  
20 cryosphere region.


## 21 **2 Study area**

22 The HMA area mainly comprises the Tibetan Plateau and surrounding alpine ranges. The area is  
23 divided into 13 sub-regions in version 5 of the Randolph Glacier Inventory (RGI 5.0), i.e., the  
24 Himalaya area (Western Himalaya, Central Himalaya, and Eastern Himalaya), Hengduan Mountains,  
25 Southern and Eastern Tibet, Inner Tibetan Plateau, Karakoram and Western Kun Lun, Qilian Shan and  
26 Eastern Kun Lun, Hindu Kush, Pamir, Alay and Western Tien Shan, Eastern Tien Shan, and Altay and  
27 Sayan (Arendt et al., 2014; Pfeffer et al., 2014). The boundaries of the 13 sub-regions and outlines of  
28 the glaciers in HMA derived from RGI 5.0 are shown in Figure . This region covers an area that  
29 extends approximately 40° in longitude and 27° in latitude, which is characterized by tremendously  
30 complex topographic conditions with widespread distribution of mountain glaciers. According to the  
31 Climatic Research Unit Time Series v4.02 data set



1 ([http://data.ceda.ac.uk/badc/cru/data/cru\\_ts/cru\\_ts\\_4.02/](http://data.ceda.ac.uk/badc/cru/data/cru_ts/cru_ts_4.02/)), the air temperature of the different  
2 sub-regions in HMA increased at an average annual rate of 0.002–0.054 °C a<sup>-1</sup> during 1990–2018 (Fig.  
3 1). The annual rate of change of precipitation in HMA during 1990–2018 varied from –9.9 to 4.2 mm  
4 a<sup>-1</sup> with a small average rate of increase of 0.3 mm a<sup>-1</sup>.

5 The HMA area has the largest surviving glaciers of any region other than Antarctica. As reported  
6 in RGI 5.0, there were 97,974 modern glaciers in our study area, covering a total area of approximately  
7 98,768.86 km<sup>2</sup>. Together, these glaciers produced an average meltwater volume of 110–150 km<sup>3</sup> a<sup>-1</sup>,  
8 which was the primary source of both lake basin formation and water supply for the development of  
9 glacial lakes. Over recent decades, glaciers in most areas of HMA appear to have experienced  
10 widespread mass wastage and area shrinkage (Bolch et al., 2012; Yao et al., 2012; Kääb, et al., 2012;  
11 Brun, et al., 2017). However, the so-called “Karakoram Anomaly” refers to a region that is a prominent  
12 exception, which is characterized by glaciers with stable or positive mass balance (Hewitt, 2005;  
13 Gardelle et al., 2013; Kääb, et al., 2015).






1 Figure 1. Location of sub-regions, rate of change of air temperature (1990–2018), and buffer area within  
2 10 km of glacier terminals for glacial lake inventory of High Mountain Asia.

### 3 **3 Data source**

4 We developed our glacial lake inventory of HMA based on 426 high-quality images selected from  
5 more than 1800 Landsat images with 30-m spatial resolution derived from the websites of the United  
6 States Geological Survey (<https://www.usgs.gov/>) and Geospatial Data Cloud (<http://www.gscloud.cn/>).  
7 To ensure the accuracy of glacial lake boundary extraction, the following criteria were applied to  
8 imagery selection. First, the cloud coverage in an image had to be <10 %. Second, for areas with no  
9 eligible or only low-quality imagery (because of snow or shadows) in the given year, acceptable images  
10 from years closest to the given year were chosen as replacements (Fig. 2). Third, images acquired in  
11 summer or autumn (July–November), when lake areas were believed near or at their maximal extent,  
12 were set as optimal choices to minimize the impact produced by seasonal area changes of the glacial  
13 lakes (Fig. 2). Based on the above criteria, 204 and 222 Landsat images were selected to represent 1990  
14 and 2018, respectively, which completely covered the buffer area within 10 km of the glaciers  
15 terminals acquired from the Second Chinese Glacier Inventory (<http://westdc.westgis.ac.cn>) and RGI  
16 5.0 ([https://www.glims.org/RGI/rgi50\\_dl.html](https://www.glims.org/RGI/rgi50_dl.html)). Among the selected images, those acquired during  
17 July–November accounted for 91 % of the total number of selected images, while those acquired  
18 during September–October accounted for 87 % of the total number. In addition, a Shuttle Radar  
19 Topography Mission digital elevation model with spatial resolution of 1"  
20 (<http://imagicode.de/map/demsearch.php>) was used to derive the elevation of the glacial lakes.

21



1  
2 Figure 2. Time phases of remote sensing images selected for High Mountain Asian glacial lake  
3 inventory.  
4

## 5 4 Glacial lake inventory methods

### 6 4.1 Outline of workflow

7 The methods and workflow adopted in this study to produce a glacial lake inventory mainly  
8 included collation of knowledge and formulation of the specifications of the glacial lake inventory, data  
9 pre-processing, manual vectorization of glacial lakes, interactive checking and error controlling, and  
10 attribute database assignment (Fig. 3).

11 (1) Collation of available knowledge regarding glacial lake inventories. As much literature as  
12 possible relevant to the investigation and recording of glacial lakes was collected. The various  
13 definitions and classifications of glacial lakes, as well as the methods adopted previously for glacial  
14 lake boundary extraction and assessment of the extent of glacial lake distributions, were summarized  
15 and normative rules formulated for the HMA glacial lake inventory, as explained further in Sect. 3.2.

16 (2) Formulation of the specifications of lake identification. First, a working group of four leading  
17 experts in the field was founded in 2014 to discuss and formulate the specifications of the glacial lake  
18 inventory. Current knowledge regarding identification of lakes from Landsat imagery (e.g., pixel colour,  
19 lake shape, and lake background features) and specifications of vectorization (e.g., viewing scale on a  
20 computer screen vectorization of mixed pixels) was discussed and unified operating criteria were



1 compiled to guide the glacial lake inventory operatives. Novice vectorization operatives were trained  
2 until their vectorization results met the pre-specifications of the inventory.


3 (3) Pre-processing of remote sensing data. Pre-processing of the Landsat imagery included false  
4 colour compositing and calculation of NDWI maps. The false colour composite images were based on  
5 combinations of the operating bands of 7, 5, and 2 or 4, 3, and 2 for Landsat TM/ETM/ETM+ images  
6 and 5, 4, and 3 for Landsat OLI images. The preliminary lake extent was extracted automatically from  
7 each image over the entire HMA area using the NDWI based on the near infrared band (NIR) and green  
8 band (GREEN), which represent the minimum and maximum water reflectance, respectively  
9 (McFeeters, 1996; Zhai et al., 2015; Li et al., 2016; Zhang et al., 2018):

10 
$$\text{NDWI} = \frac{B_{\text{GREEN}} - B_{\text{NIR}}}{B_{\text{GREEN}} + B_{\text{NIR}}} \quad (1)$$

11 where  $B_i$  is the spectral band of Landsat imagery. The NDWI maps were calculated for each selected  
12 Landsat image using different region-specific thresholds, which typically were in the range  $-0.10$  to  
13  $0.20$  for lake surfaces. When manual vectorization was performed on a false colour composite image,  
14 the NDWI maps of potential glacial lakes were overlaid to assist in glacial lake identification.

15 (4) Manual vectorization and entering of attribute data. The inventory work was performed during  
16 2014–2019. Seven groups were formed to conduct lake boundary vectorization of the 13 HMA  
17 sub-regions. After vectorization of a glacial lake, it was required that manual attribute items (e.g., data  
18 source and lake type) be input concurrently.

19 (5) Interactive checking and accuracy control. First, glacial lakes were discerned via human–  
20 computer interaction, i.e., potential glacial lakes were revealed by the NDWI maps or identified  
21 visually from the false colour composite images. Second, glacial lake boundary vectorization results  
22 were checked interactively by another vectorization operative to eliminate misclassified areas of  
23 shadow and ice and to add areas of glacial lakes evidently omitted in the boundary extraction process.  
24 This checking process also minimized the subjective judgment errors of the operatives. Third, attribute  
25 items such as glacial lake classification, new/disappeared lakes, and separated/coalesced lakes were  
26 checked interactively. In this process, Google Earth imagery was used as an important auxiliary  
27 reference data source for error examination.



1  
2

Figure 3. Flow chart of HMA glacial lake inventory.

### 3 4.2 Illustration of key methods

#### 4 4.2.1 Definition of glacial lakes

5 The definition of a glacial lake determines the type of cryosphere water body that will be recorded  
 6 as a glacial lake. There are multiple definitions of a glacial lake based on different perspectives (Mool  
 7 et al., 2001; Yao et al., 2018). When glacial lake inventories are undertaken, most emphasize the  
 8 elementary role of glaciation in the formation of glacial lakes (Clague and Evans, 2000; Qin et al., 2016;  
 9 Mool et al., 2001). The remarkable difference is whether the period of glaciation or the supply source  
 10 of glacial lakes is given greatest attention. Some studies that focused on the former proposed that a  
 11 glacial lake is a natural water body formed by alpine glacier movement since the Last Glacial





1 Maximum, i.e., ancient or modern glaciers (Liu et al., 1988; Costa and Schuster, 1988). However, other  
2 studies emphasized the relation of glacial lakes to meltwater in glaciated areas (Wang et al., 2013;  
3 Weicai et al., 2014; Zhang et al., 2015). The glacial lake inventory data compiled in this study are  
4 intended for use both in water source evaluation and in assessment of environmental change in the  
5 alpine cryosphere. Thus, lakes related to glaciers or to glaciation in the alpine cryosphere were all  
6 recorded as glacial lakes.

7 Most Quaternary glaciers have disappeared and the remaining relics are incomplete, which makes  
8 it difficult to recover a continuous and complete glaciation range in alpine regions. Thus, it is of great  
9 importance to ensure the range of glaciation in an alpine region when conducting a glacial lake  
10 inventory based on remote sensing data. The most practical approach might be to specify an indicator  
11 threshold to define the glaciation extent according to relevant findings of existing glacier relics in a  
12 typical region. On the one hand, the glaciation frontier can usually be indicated by a specified lowest  
13 elevation threshold, which is generally closely related to the regional climatic context caused by the  
14 elevation effect. However, the lowest elevation threshold might vary enormously with respect to  
15 different regions because regional climatic settings differ. For instance, the lowest elevations of 1700 m  
16 in Austria (Buckel et al., 2018), 2000 m in Pakistan (Senese et al., 2018), 3000 m in Nepal and Bhutan  
17 (Mool et al., 2001), and 3500 m in Peru (Hanshaw and Bookhagen, 2014) were used as specified  
18 elevation thresholds to record glacial lakes. On the other hand, defining glaciation extent within a  
19 specific distance from glacier terminals could be more suitable for the establishment of a glacial lake  
20 inventory in relatively large-scale regions with complex regional climate, because the differing climate  
21 within large-scale regions can be indicated approximately by the lowest elevation of individual glacier  
22 terminals. Some studies adopted distances of 2, 3, or 10 km from modern glacier terminals as  
23 thresholds with which to define areas of glacial lakes (Petrov et al., 2017; Veh et al., 2018; Wang et al.,  
24 2012, 2013). Distances of 2, 5, 10, and 20 km were considered by Zhang et al. (2015). They found that a  
25 distance of 10 km from a modern glacier terminal might be a reasonable guide to glaciation extent and a  
26 threshold suitable for a glacial lake inventory of the Tibetan Plateau. This was supported by the finding  
27 that the most distant glacierized boundary of the Little Ice Age was up to 10 km from the modern  
28 glaciers in the Himalaya area (Wang et al., 2012, Nie et al., 2017). Additionally, to record glacial lakes  
29 more precisely, combined distance and elevation thresholds have been used simultaneously to define  
30 areas of glacial lakes in special small regions, e.g., lakes at elevations above 1500 m and within 2 km of  
31 modern glaciers were recorded as glacial lakes in Uzbekistan (Petrov et al., 2017). In this study, given



1 the large scale of the HMA region with its complex climatic context and extremely varied terrain, the  
2 data set compiled included glacial lakes within a buffer zone of 10 km from modern glacier terminals,  
3 which covered an area of approximately  $1.19 \times 10^6 \text{ km}^2$  according to the Second Glacier Inventory of  
4 China and RGI 5.0 (Fig. 1).

#### 5 **4.2.2 Classification of glacial lakes**

6 In glaciation regions, the characteristics of glacial lakes, which include the phase of lake formation,  
7 lake basin topography, dam material constituents, geometrical relationship with modern glaciers, and  
8 source of water supply (or combinations thereof), have been employed as the basis for glacial lake  
9 classification systems (Huggel et al., 2002; Liu et al., 1988; Mool et al., 2001; Yao et al., 2018). For  
10 instance, based on lake basin topography, lakes in an inventory of the Hindu Kush–Himalaya region  
11 were classified as erosion lakes, valley trough lakes, cirque lakes, blocked lakes, lateral and end  
12 moraine-dammed lakes, and supraglacial lakes (Liu et al., 1988; Mool et al., 2001). Recently, Yao et al.  
13 (2018) presented a reasonably complete classification schema for glacial lake inventory and study of  
14 glacial lake hazards that included six classes and eight sub-classes based mainly on the mechanism of  
15 glacial lake formation, lake basin topography, and the geometrical relationship with modern glaciers.

16 Generally, it is a little difficult to distinguish glacial lake type in terms of material properties,  
17 topographic features, and phase of lake formation using remote sensing imagery. Moreover, most of the  
18 standards mentioned above were found inapplicable in previous studies of glacial lake classification in  
19 large-scale regions such as HMA because of the lack of sufficient amounts of remote sensing data with  
20 appropriate resolution. In this study, the hydrologic relationship between glacial lakes and modern  
21 glaciers was adopted as a classification criterion because the present data set is intended to provide  
22 fundamental data for water resource evaluation and glacier hazard assessment. Consequently, glacial  
23 lakes were divided into just two types: glacier-fed lakes and non-glacier-fed lakes. This classification  
24 was based on whether the hydrological flow of the modern glacier and topographic features of the lake  
25 basin allowed a lake to receive meltwater from the modern glacier. To achieve reliable classification  
26 results, glacial lakes were distinguished with the assistance of 3D digital terrain imagery from Google  
27 Earth, a Shuttle Radar Topography Mission digital elevation model, and glacier outlines from RGI 5.0.



### 1 4.2.3 Extraction of lake boundary


2 This study adopted automatic glacial lake extraction and manual glacial lake boundary  
3 vectorization to determine glacial lake boundaries. In the NDWI-based automatic lake boundary  
4 extraction approach, two bands were selected to facilitate a ratio calculation to maximize the difference  
5 between water and non-water objects in the remote sensing imagery based on a given threshold. The  
6 given threshold was determined subjectively with consideration of how much detailed information of  
7 the lake water bodies was captured precisely. The given threshold was varied to account for various  
8 factors such as the differences in Landsat sensors (i.e., TM, ETM/ETM+, and OLI), time phase of  
9 images, quality of images, and complexity of surface features. To achieve the optimal threshold for  
10 lake water body recognition, the candidate threshold was debugged iteratively for each image. In  
11 practice, because the area of the glacial lakes was usually small (see next paragraph) and the spectral  
12 features of the lake water bodies were varied, the threshold had to be set to allow capture of the greatest  
13 number of water body pixels, which consequently resulted in simultaneous acquisition of more  
14 non-lake-water-body noise information. It also resulted in more effort in the subsequent manual  
15 modification to reduce noise information using methods such as algorithms to eliminate mountain  
16 shadows (Gardelle et al., 2011).

17 Manual visual vectorization distinguishes lake boundaries by identifying the unique texture,  
18 colour, and other characteristics of glacial lakes in false colour composite images based on available  
19 professional knowledge and accumulated experience in vectorization operations. Even though it was  
20 regarded a time-consuming and labour-intensive process, it was also considered an attractive approach  
21 because of its consistency, high level of quality control, and reasonably simple operational procedure,  
22 given the varied quality of Landsat images for the large-scale HMA region. In this study, the manual  
23 visual vectorization process was generally found more suitable in terms of effort and precision for  
24 generating a glacial lake inventory data set of the HMA region in comparison with automatic glacial  
25 lake extraction. Therefore, manual visual vectorization in conjunction with NDWI maps was the main  
26 method adopted to extract glacial lake boundaries to minimize the deficiencies produced by individual  
27 subjectivity of the operatives.

28 The minimum number of pixels used to extract a glacial lake water body was found inconsistent in  
29 the available literature. For example, arbitrary threshold areas of 0.0027 km<sup>2</sup> (three lake water body  
30 pixels) (Zhang et al., 2015) and 0.0081 km<sup>2</sup> (nine lake water body pixels) (Nie et al., 2017) have been  
31 used in earlier glacial lake investigations. Moreover, minimum threshold areas of 0.01 km<sup>2</sup>



1 (approximately 10 lake water body pixels), 0.02 km<sup>2</sup> (approximately 22 lake water body pixels) and  
2 0.1km<sup>2</sup> (~111 lake water body pixels) have also been set to evaluate the level of risk of glacial lake  
3 outburst floods in Himalaya and the Tien Shan Mountains (Petrov et al., 2017, Wang et al., 2013;  
4 Worni et al., 2013; Bolch et al., 2011; Allen et al., 2019). Theoretically, one pure pixel of a lake water  
5 body could be recorded as a glacial lake. However, a glacial lake is generally not represented by one  
6 pure pixel unless perfectly aligned with the raster grid; usually, it would be surrounded partly or fully  
7 by 1–8 mixed lake water body pixels (Fig. 4a, b). Consequently, manual delineation was required for  
8 approximately 1/2, 1/4, or 3/4 of the peripheral mixed pixels surrounding pure lake water body pixels  
9 (Fig. 4c). In this study, the minimum glacial lake area recorded was set at 0.0054 km<sup>2</sup> (e.g., 3–4 pure  
10 lake water body pixels with approximately 12 mixed boundary pixels) because a lake area covering  
11 fewer than three pure lake water pixels could possibly have an error of >100 % (see Sect. 5).



12  
13 Figure 4. Sketches showing a pure water body pixel and surrounding potential mixed water body pixels:  
14 (a) a pure water body pixel, (b) a pure water body pixel surrounded by four potential mixed water body  
15 pixels, and (c) three pure water body pixels with seven surrounding potential mixed water body pixels.

#### 16 4.2.4 Input of attribute items

17 Eight attribute items were input into the HMA glacial lake inventory: lake coding, location  
18 (longitude, latitude, and elevation), perimeter, area, type, area error, time phase, source image



1 information, and sub-region of located lake. (1) We encoded each glacial lake based on its central  
2 location using the same coding format used by the National Snow and Ice Data Centre to encode  
3 glaciers. The code can be expressed as “GLmmmmmmEnnnnnN”, where m and n represent the results  
4 of the longitude and latitude of each glacial lake centroid multiplied by 1000, respectively, GL is the  
5 abbreviation of glacial lake, and E and N represent eastings and northings, respectively. (2) The  
6 location information of each glacial lake was labelled as the geographic coordinates of the centroid of  
7 the shape of each glacial lake, calculated using ArcMap software. The lake elevation was defined as the  
8 average elevation of a buffer zone of 30-m radius centred on the glacial lake centroid, which was  
9 derived from the Shuttle Radar Topography Mission digital elevation model. (3) The area and  
10 perimeter of each lake were calculated using ArcMap based on the unified geography coordinate  
11 system of GCS\_WGS\_1984 and the Asia\_North\_Albers\_Equal\_Area\_Conic projection system,  
12 respectively, to avoid errors caused by projection deformation. (4) The error of lake area was calculated  
13 using Eqs. (2) and (3) (Sect. 5). (5) Lake type, which was input manually, was defined as either  
14 glacier-fed lake or non-glacier-fed lake (see Sect. 4.2.2). (6) Lake time phase was the acquisition date  
15 of the original Landsat image, which was recorded as the month and year for each lake. (7) Source  
16 image information referred to the image number of the Landsat images used to extract the glacial lake  
17 boundary. (8) The sub-region to which each lake belonged identified the regional location within the  
18 HMA area. Each lake was assigned based on shp. file data of the boundaries of the 13 HMA  
19 sub-regions, obtained from the National Snow and Ice Data Centre, using the ArcMap spatial analysis  
20 tool.

## 21 **5 Error assessment**

22 The errors associated with glacial lake extraction from remote sensing imagery using manual visual  
23 delineation are generally related to components of the quality of the images (e.g., spatiotemporal  
24 resolution, cloud coverage, and mountain shadows), experience, operative subjectivity, and the threshold  
25 area of the inventory (Gardelle et al., 2011; Hall et al., 2003; Paul et al., 2004; Salerno et al., 2012; Zhang  
26 et al., 2015). It has been reported that the area error of glacial lake boundary extraction based on remote  
27 sensing images can be approximately  $\pm 0.5$  pixels depending on the quality of the imagery (Fujita et al.,  
28 2009; Salerno et al., 2012). Furthermore, the area error of glacial lake delineation attributable to manual  
29 delineation can be assumed to follow a Gaussian distribution (Hanshaw and Bookhagen, 2014). Hence,  
30 the theoretical maximum area error of glacial lake boundary extraction is the half-area of the edge pixels



1 because pure lake water body pixels are usually surrounded by mixed pixels (Fig. 4b, c). The lake area  
2 error within one standard deviation ( $1\sigma$ ) can be expressed as follows (Hanshaw and Bookhagen, 2014):


$$3 \quad \text{Error}(1\sigma) = \frac{P}{G} \times \frac{G^2}{2} \times 0.6872 \quad (2)$$

$$4 \quad E = \frac{\text{Error}(1\sigma)}{A} \times 100 \% \quad (3)$$

5 where  $P$  is the perimeter of the glacial lake (m),  $G$  is the spatial resolution of the remote sensing  
6 imagery (30 m in this data set), 0.6872 is the revised coefficient under  $1\sigma$  (i.e., approximately 69 % of  
7 peripheral pixels are subjected to errors),  $E$  is the relative error of the glacial lake, and  $A$  is the total  
8 area of the glacial lake.

9 The resulting calculated error indicated that the total absolute area error of HMA glacial lakes was  
10 approximately  $\pm 231.44$  and  $\pm 259.68$  km<sup>2</sup> and the average relative error was  $\pm 13.8$  and  $\pm 13.3$  % in 1990  
11 and 2018, respectively. The relative area errors of each lake varied from 2–85 %, and a significant  
12 power exponential relationship was found between the relative area error and the sizes of the glacial  
13 lakes ( $E = 0.050A^{-0.45}$ ,  $R^2 = 0.96$ ,  $\alpha < 0.001$ ) (Fig. 5a). Small-sized lakes (i.e., area  $\leq 0.01$  km<sup>2</sup>,  
14 which accounted for 2 % of the total lake area in HMA) had the largest average relative area error of  
15 44.6 % (Fig. 5b). Medium-sized lakes (i.e., area of 0.01–0.1 km<sup>2</sup>, which accounted for 34 % of the total  
16 lake area in HMA) had an average relative area error of 22.1 % (Fig. 5c). Large-sized lakes (i.e., area  $\geq$   
17 0.1 km<sup>2</sup>, which accounted for 64 % of the total lake area in HMA) had the smallest average relative  
18 area error of 4.1 % (Fig. 5d). In summary, smaller glacial lakes in the HMA region had larger relative  
19 area errors, and vice versa.

20 To further verify the accuracy of the manual delineation of glacial lake boundaries, nine lakes  
21 located within the HMA region were surveyed using a portable GPS device (Trimble GeoXH6000)  
22 during July–August 2018 (Fig. 6). The lakes selected for field survey covered areas of 0.01–2.97 km<sup>2</sup>.  
23 The field-based lake boundaries were compared with those obtained via manual delineation (i.e.,  
24 derived from Landsat OLI imagery acquired during 2018). It was found that the area error (i.e., the  
25 percentage difference of the absolute area encircled by the manually delineated lake boundary and that  
26 derived by the GPS survey) varied from 5.5–25.5 %. Moreover, it was determined that the average  
27 horizontal distance deviation between the two types of boundary varied from 4.5–33.5 m (Table 1).  
28 Overall, the horizontal deviations were largely confined to one pixel, and the average accuracy of the  
29 delineation of glacial lake boundaries was within  $\pm 0.5$  pixels ( $\pm 15$  m).




1  
 2 Figure 5. Relationships of relative area error against size of glacial lakes in HMA: (a) relationship for  
 3 glacial lakes of all sizes and (b)–(d) relationships for glacial lakes of specific size.

4  
 5 Table 1. Horizontal deviations between lake boundaries obtained by manual delineation and field  
 6 survey using a portable GPS device (Trimble GeoXH6000)

Name (labelled in Figure 6 )	Lake ID	Lake size (km <sup>2</sup> )	Horizontal deviations of delineation boundary (m)			Area error(%)
			minimum	maximum	average	
Qiongyong Cuo (A)	GL090225E28890N	0.08	-7.6	9.1	4.5	5.5
Passu Lake (B)	GL074878E36457N	0.15	-10.9	12.0	6.0	6.8
Longbasa Lake (C)	GL088071E27950N	1.49	-22.7	-8.4	12.4	/
Zongge Cuo (D)	GL087654E28113N	1.48	-26.8	24.9	13.5	6.1
Unnamed (E)	GL088151E28010N	0.01	-4.7	4.8	3.2	16.3
Unnamed (F)	GL088257E28011N	0.58	-12.9	12.4	4.6	/
Unnamed (G)	GL088240E28005N	0.40	-20.8	15.9	7.1	/
Large Laigu Lake (H)	GL096818E29298N	2.97	-36.7	-6.4	15.3	/
Small laigu Lake (I)	GL096832E29294N	1.02	-32.8	17.6	9.8	/

7 Note: “/” indicates the sample lake boundary was only partly surveyed using the handheld GPS device.

8




1  
2 Figure 6. Glacial lakes in the HMA region surveyed in summer 2018 (the background maps of surveyed  
3 lakes are the Landsat OLI images; red lines denote lake boundary obtained by field survey; blue lines  
4 denote outlines derived from satellite imagery).

## 5 **6 Distribution and changes of HMA glacial lakes**

6 As indicated by the achieved HMA glacial lake inventory, 28,953 ( $1955.94 \pm 259.68 \text{ km}^2$ ) glacial  
7 lakes were identified in 2018 and their distribution had considerable spatial heterogeneity (Fig. 7). The  
8 greatest concentration of glacial lakes was in Eastern Himalaya ( $308.23 \pm 37.29 \text{ km}^2$ , which accounted  
9 for 15.8 % of the total area of glacial lakes) and Inner Tibet ( $299.67 \pm 38.75 \text{ km}^2$ , which accounted for  
10 15.3 % of the total area of glacial lakes). Relatively few glacial lakes were found distributed in Eastern  
11 Kun Lun and Qilian Shan ( $38.85 \pm 6.04 \text{ km}^2$ , which accounted for 2.0 % of the total area of glacial lakes)  
12 and Eastern Tien Shan ( $40.55 \pm 6.98 \text{ km}^2$ , which accounted for 2.1 % of the total area of glacial lakes).  
13 The HMA glacial lakes are located within the elevation range of 1600–6300 m. An approximate normal  
14 distribution was presented both for the lakes of the entire HMA region and for the lakes in most  
15 sub-regions. More than 46 % of the HMA lake area has survived within the vertical range of 4500–5400  
16 m, with the peak lake area of  $256.13 \pm 34.25 \text{ km}^2$  (accounting for 13.1 % of the total area) in the range of  
17 5000–5200 m. The elevation band of peak lake area in the different sub-regions varied from 2400–2600  
18 m in Altay and Sayan to 5200–5400 m in Central Himalaya, Karakoram, and Western Kun Lun.





1  
 2 Figure 7. Distribution and change of glacial lake area in the entire HMA and its 13 sub-regions from  
 3 1990–2018.

4  
 5 The HMA glacial lakes experienced widespread areal expansion during 1990–2018 with an  
 6 average rate of increase in area of 15.5 % (Fig. 7). The rate of change of area varied widely between  
 7 different sub-regions and different 200-m elevation bands. The glacial lakes in Altay and Sayan  
 8 experienced the most rapid expansion in area during 1990–2018 with an average rate of increase of  
 9 45.6 %, whereas the rate of change was only 7.2 % in Pamir. Glacial lakes have tended to develop to  
 10 higher elevations during recent decades with the maximum distribution elevation of 6078 m in 1990  
 11 rising to 6247 m in 2018. The rate of change of glacial lakes in the different 200-m elevation bands  
 12 presented a large average trend against elevation, rising as a whole during 1990–2018 (Fig. 7). The lake  
 13 area expanded most from the elevation of approximately 5000 m and the rate of expansion reached  
 14 approximately 35 % above 5800 m in the entire HMA region, although it differed between different  
 15 sub-regions. Lake area showed a notable rate of increase with elevation in most sub-regions, e.g.,  
 16 Hissar Alay and Western Tien Shan, Hindu Kush, Eastern Himalaya, Hengduan Shan, Eastern Tien



1 Shan, and Altay and Sayan. Although the rate of expansion varied violently, no observable trends were  
2 discovered in Karakoram and Western Kun Lun, Western Himalaya, and Inner Tibet. The rate of  
3 expansion in Central Himalaya and Southern and Eastern Tibet was found to have decreased with  
4 increasing elevation (Fig. 7).

## 5 **7 Data availability**

6 The data set developed in this study comprised two .shp file documents containing the glacial lake  
7 inventory of the HMA region in 1990 and 2018. The data now can be accessed on the National Special  
8 Environment and Function of Observation and Research Stations Shared Service Platform (China) at  
9 <http://dx.doi.org/10.12072/casnw.064.2019.db> (Wang et al., 2019a).

## 10 **8 Conclusions**

11 A glacial lake inventory of the HMA region was realized based on satellite remote sensing data  
12 and GIS techniques. Eight attribute items were recorded in the glacial lake inventory data set of the  
13 HMA region. Lake area error was assessed carefully with respect to theoretical analysis of lake  
14 boundary pixels and actual boundaries derived by GPS field-based surveys. On average, the deviations  
15 between the delineation of lake boundaries derived using the two methods were within  $\pm 0.5$  pixels ( $\pm 15$   
16 m). The relative area errors of each lake in 2018 varied from 2–85 %, and the average relative area  
17 errors of  $\pm 13.3$  % in the entire HMA region were characterized by increase in the relative area error  
18 with decreasing lake size.

19 Overall, 28,953 glacial lakes with a total area of  $1955.93 \pm 259.68$  km<sup>2</sup> were catalogued in 2018 in  
20 the HMA region. Glacial lakes survived in all 13 sub-regions of HMA from the elevation of 1600 to  
21 6300 m. Glacial lakes were found concentrated in the sub-regions of Eastern Himalaya and Inner Tibet  
22 and at elevation bands of 4500–5400 m. The HMA glacial lakes have experienced widespread  
23 expansion in area with an average rate of increase in area of 15.5 %. Lake area expanded most in the  
24 higher elevation bands during 1990–2018. The data set has been developed as basic data for cryosphere  
25 hydrology research; however, it is expected that it could support practical utilization and management  
26 of water resources and assessment of glacier-related hazards in the HMA region.

## 27 **9 Acknowledgements**

28 The study was funded by the National Natural Science Foundation of China (No. 41771075, No.  
29 41571061, and No. 41271091). The authors are grateful to the lake boundary vectorization operators  
30 who were not written as authors, there are Yao Chao, Chen Shiyin, Zhu Xiaoxi, Li Ruijia, Huang Rong,



1 Peng Xin, Xiang Lili, Yi Ying, Liu Yanlin, Fu Yongqiao, Ran Weijie, Gu Ju. We thank James Buxton  
2 MSc from Liwen Bianji, Edanz Group China ([www.liwenbianji.cn/ac](http://www.liwenbianji.cn/ac)), for editing the English text of  
3 this manuscript.

#### 4 **10 References**

- 5 Allen, S.K., Zhang, G., Wang, W., Yao, T., and Bolch, T.: Potentially dangerous glacial lakes across the  
6 Tibetan Plateau revealed using a large-scale automated assessment approach. *Science Bulletin*, 64,  
7 435-445, doi.org/10.1016/j.scib.2019.03.011, 2019.
- 8 Arendt, A., Bliss, A., Bolch, T., Pfeffer, W. T., Cogley, J. G., and Gardner, A. S.: Randolph Glacier  
9 Inventory – A Dataset of Global Glacier Outlines: Version 5.0, Boulder Colorado, USA, 2015.
- 10 Bajracharya, S. R., Mool, P. K., and Shrestha, B. R.: Impact of climate change on Himalayan glaciers and  
11 glacial Lakes, Kathmandu, Nepal: International Centre for Integrated Mountain Development, 1-119,  
12 2007.
- 13 Bolch, T., Kulkarni, A., Kääb, A., Huggel, C., Paul, F., Cogley, J. G., Frey, H., Kargel, J. S., Fujita, K.,  
14 Scheel, M., Bajracharya, S., and Stoffel, M.: The State and Fate of Himalayan Glaciers, *Science*, 336  
15 (6079), 310–314 , doi:10.1126/science.1215828 , 2012.
- 16 Bolch, T., Peters, J., Yegorov, A., Pradhan, B., Buchroithner, M., and Blagoveshchensky, V.:  
17 Identification of potentially dangerous glacial lakes in the northern Tien Shan, *Natural Hazards* 59,  
18 1691-1714, doi:10.1007/s11069-011-9860-2, 2011.
- 19 Brun, F., Berthier, E., Wagnon, P., Kääb, A., and Treichler, D.: A spatially resolved estimate of High  
20 Mountain Asia glacier mass balances,2000-2016, *Nat. Geosci.*, 10(9), doi:10.1038/NGEO2999  
21 668-673,2017.
- 22 Buckel, J., Otto, J. C., Prasicek, G., and Keuschnig, M.: Glacial lakes in Austria - Distribution and  
23 formation since the Little Ice Age, *Global*, 164, 39-51. doi:10.1016/j.gloplacha.2018.03.003, 2018.
- 24 Chaohai, L., and Sharma, C. K.: Report on first expedition to glaciers and glacier lakes in the Pumqu  
25 (Arun) and Poiqu (Bhote-SunKosi) River Basin, Xizang (Tibet), China. Beijing: Science Press,  
26 1-192,1988.
- 27 Chengde, Y., Xin, W., Junfeng, W., Qionghuan, L., Anxin, L., Yong, Z., and Zhiguang, T.: Chinese  
28 glacial lake inventory based on 3S technology method, *J Geogr Sci*, 74, 544-556,  
29 doi:10.11821/dlxb201903011, 2019.



- 1 Chunqiao, S., Bo, H., Linghong, K., and Richards, K. S.: Remote sensing of alpine lake water
- 2 environment changes on the tibetan plateau and surroundings: A review, *Int. J. Remote Sens.*, 92, 26-37,
- 3 doi:10.1016/j.isprsjprs.2014.03.001, 2014.
- 4 Chunqiao, S., Yongwei, S., Linghong, K., Yong, N., and Jida, W.: Glacial lake evolution in the
- 5 southeastern Tibetan Plateau and the cause of rapid expansion of proglacial lakes linked to
- 6 glacial-hydrogeomorphic processes, *Journal of Hydrology*, 540, 504-514,
- 7 doi:10.1016/j.jhydrol.2016.06.054, 2016.
- 8 Clague, J. J., and Evans, S. G.: A review of catastrophic drainage of moraine-dammed lakes in British
- 9 Columbia, *Quaternary Science Reviews*, 19, 1763-1783, doi:10.1016/s0277-3791(00)00090-1, 2000.
- 10 Cook, K. L., Andermann, C., Gimbert, F., Adhikari, B. R., and Hovius, N.: Glacial lake outburst floods as
- 11 drivers of fluvial erosion in the Himalaya, *Science*, 362, 53–57, doi:10.1126/science.aat4981, 2018.
- 12 Costa, J. E., and Schuster, R. L.: The formation and failure of natural dams, *Geological Society of*
- 13 *America Bulletin*, 100, 1054-1068, doi:10.1130/0016-7606(1988)1002.3.CO;2, 1988.
- 14 Fujita, K., Akiko, S., Takayuki, N., Yamaguchi, S., and Sharma, R. R.: Recent changes in Imja glacial
- 15 lake and its damming moraine in the Nepal Himalaya revealed by in situ surveys and multi-temporal
- 16 ASTER imagery, *Environmental Research Letters*, 4, 045205, doi:10.1088/1748-9326/4/4/045205,
- 17 2009.
- 18 Gardelle, J., Yves, A., and Etienne, B.: Contrasted evolution of glacial lakes along the Hindu Kush
- 19 Himalaya mountain range between 1990 and 2009, *Global*, 75, 47-55,
- 20 doi:10.1016/j.gloplacha.2010.10.003, 2011.
- 21 Gardner, A. S., Moholdt, G., Graham, C. J., Wouters, J., Arendt, A. A., Wahr, J., Berthier, E., Hock, R.,
- 22 Pfeffer, W. T., Kaser, G., Ligtenberg, S. M. R., Bolch, T. Sharp, M. J., Hagen, J. O., Broeke, M. R. V. D.,
- 23 and Paul, F.: A Reconciled Estimate of Glacier Contributions to Sea Level Rise: 2003 to 2009, *Science*,
- 24 340, 852–857, doi:10.1126/science.1234532, 2013.
- 25 Hall, D. K., Klaus J. B., Wolfgang S., Bindschadler, R. A., and Chien, J. Y. L.: Consideration of the
- 26 errors inherent in mapping historical glacier positions in Austria from the ground and space (1893–2001),
- 27 *Remote Sens. Environ.*, 86, 566-577, doi:10.1016/s0034-4257(03)00134-2, 2003.
- 28 Hanshaw, M. N., and Bookhagen, B.: Glacial areas, lake areas, and snow lines from 1975 to 2012: status
- 29 of the Cordillera Vilcanota, including the Quelccaya Ice Cap, northern central Andes, Peru, *The*
- 30 *Cryosphere*, 8, 359-376, doi:10.5194/tc-8-359-2014, 2014.



- 1 Hewitt, K.: The Karakoram Anomaly? Glacier Expansion and the ‘Elevation Effect,’ Karakoram
- 2 Himalaya, *Mountain Research and Development*, 25, 332-340, doi:10.1659/0276-4741(2005)025
- 3 [0332:TKAGEA]2.0.CO;2, 2005.
- 4 Huggel, C., Kääb, A., Haeberli, W., Teysseire, P., and Paul, F.: Remote sensing based assessment of
- 5 hazards from glacier lake outbursts: a case study in the Swiss Alps, *Canadian Geotechnical Journal*, 39,
- 6 316-330, doi:10.1139/t01-099, 2002.
- 7 Kääb, A., Berthier, E., Nuth, C., Gardelle, J., and Arnaud, Y.: Contrasting patterns of early
- 8 twenty-first-century glacier mass change in the Himalayas, *Nature*, 488(7412), 495-498,
- 9 doi:10.1038/nature11324, 2012.
- 10 Kääb, A., Treichler, D., Nuth, C., and Berthier, E.: Brief Communication: Contending estimates of
- 11 2003-2008 glacier mass balance over the Pamir–Karakoram–Himalaya, *The Cryosphere*, 9, 557–564, doi:
- 12 10.5194/tcd-8-5857-2014, 2015.
- 13 Li, J., Sheng, Y., and Luo, J.: Automatic extraction of Himalayan glacial lakes with remote sensing,
- 14 *Journal of remote sensing*, 15(1), 29-43, doi:10.1631/jzus.C0910717, 2011.
- 15 Li, Y., Gong, X., Guo, Z., Xu, K., Hu, D., and Zhou, H.: An index and approach for water extraction
- 16 using Landsat–OLI data, *Int. J. Remote Sens.*, 37(16), 3611-3635. doi: 10.1080/01431161.2016.1201228,
- 17 2016.
- 18 Mcfeeters, S. K.: The use of the Normalized Difference Water Index (NDWI) in the delineation of open
- 19 water features, *Int. J. Remote Sens.*, 17(7), 1425-1432, doi: 10.1080/01431169608948714, 1996.
- 20 Mei-meí, Z., Fang, C., and Bang-sen, T.: An automated method for glacial lake mapping in High
- 21 Mountain Asia using Landsat 8 imagery. *J. Mt. Sci.* 15(1), 13-24, doi: 10.1007/s11629-017-4518-5,
- 22 2018.
- 23 Mool, P. K., Bajracharya, S. R., Joshi, S. P.: Inventory of glaciers, glacial lakes and glacial lake outburst
- 24 floods, Nepal. International Centre for Integrated Mountain Development, Kathmandu, Nepal, 2001.
- 25 Nie, Y., Sheng, Y., Liu, Q., Liu, L., Liu, S., Zhang, Y., and Song, C.: A regional-scale assessment of
- 26 Himalayan glacial lake changes using satellite observations from 1990 to 2015, *Remote Sens. Environ.*,
- 27 189, 1-13, doi:10.1016/j.rse.2016.11.008, 2017.
- 28 Paul, F., Haggel, C., and Kääb, A.: Combining satellite multispectral image data and a digital elevation
- 29 model for mapping debris-covered glaciers, *Remote Sens. Environ.*, 89, 510-518,
- 30 doi:10.1016/j.rse.2003.11.007, 2004.



- 1 Petrov, M. A., Sabitov, T. Y., Tomashevskaya, I. G., Glazirin, G. E., Chernomorets, S. S., Savernyuk, E.
- 2 A., and Stoffel, M.: Glacial lake inventory and lake outburst potential in Uzbekistan, *Sci Total Environ*
- 3 592, 228-242, doi:10.1016/j.scitotenv.2017.03.068, 2017.
- 4 Pfeffer, W., Tad, A. A., Arendt, A., Bolch, T., Cogley, J. G., and Gardner, A. S.: The Randolph Glacier
- 5 Inventory: a globally complete inventory of glaciers, *Journal of Glaciology*, 60, 537-552,
- 6 doi:10.3189/2014JoG13J176, 2017.
- 7 Salerno, F., Thakuri, S., D'Agata, C., Smiraglia, C., Manfredi, E. C., Viviano, G., and Tartari, G.: Glacial
- 8 lake distribution in the Mount Everest region: Uncertainty of measurement and conditions of formation,
- 9 *Global*, 92-93,30-39, doi:10.1016/j.gloplacha.2012.04.001, 2012.
- 10 Senese, A., Maragno, D., Fugazza, D., Soncini, A., D'Agata, C., Azzoni, R. S., and Diolaiuti, G. A.:
- 11 Inventory of glaciers and glacial lakes of the Central Karakoram National Park (CKNP-Pakistan),
- 12 *Journal of Maps* ,14, 189-198, doi:10.1080/17445647.2018.1445561, 2018.
- 13 Slemmons, K. E., Saros, J. E., and Simon, K.: The influence of glacial meltwater on alpine aquatic
- 14 ecosystems: a review, *Environ Sci Process Impacts*, 15(10), 1794-806, doi: 10.1039/c3em00243h, 2013.
- 15 Veh, G., Korup, O., Roessner, S., and Walz, A.: Detecting Himalayan glacial lake outburst floods from
- 16 Landsat time series, *Remote Sens. Environ*, 207, 84–97, doi:10.1016/j.rse.2017.12.025, 2018.
- 17 Wang, W., Xiang, Y., Gao, Y., Lu, A. and Yao, T.: Rapid expansion of glacial lakes caused by climate
- 18 and glacier retreat in the Central Himalayas, *Hydrological Processes*, 29(6): 859-874, 2014.
- 19 Wang, X., Chai, K., Liu, S., Wei, J., Jiang, Z., and Liu, Q.: Changes of glaciers and glacial lakes implying
- 20 corridor-barrier effects and climate change in the Hengduan Shan, southeastern Tibetan Plateau, *Journal*
- 21 *of Glaciology*, 63, 535-542, doi:10.1017/jog.2017.14, 2017.
- 22 Wang X., Guo X., Yang C., Liu Q., Wei J., Zhang Y., Liu S., Zhang Y., Jiang Z., and Tang Z., Glacial
- 23 lake inventory of High Mountain Asia, National Special Environment and Function of Observation and
- 24 Research Stations Shared Service Platform, doi:10.12072/casnw.064.2019.db, 2019a.
- 25 Wang, X., Ding, Y., and Zhang, Y.: The influence of glacier meltwater on the hydrological effect of
- 26 glacial lakes in mountain cryosphere, *Journal of lake Sciences*, 31(3), 609-620, 2019b.
- 27 Wang, X., Ding, Y., Liu, S., Jiang, L., Wu, K., Jiang, Z., and Guo, W.: Changes of glacial lakes and
- 28 implications in Tian Shan, central Asia, based on remote sensing data from 1990 to 2010, *Environmental*
- 29 *Research Letters* , 8, 044052, doi:10.1088/1748-9326/8/4/044052, 2013.



- 1 Wang, X., Liu, S., Guo, W., Yao, X., Jiang, Z., and Han, Y.: Using Remote Sensing Data to Quantify
- 2 Changes in Glacial Lakes in the Chinese Himalaya, *Mountain Research and Development*, 32(2),
- 3 203-212, doi: 10.1659/mrd-journal-d-11-00044.1, 2012.
- 4 Wang, X., Liu, Q., Liu, S., Wei, J., and Jiang, Z.: Heterogeneity of glacial lake expansion and its
- 5 contrasting signals with climate change in Tarim Basin, Central Asia, *Environmental Earth Sciences*, 75,
- 6 696, doi:10.1007/s12665-016-5498-4, 2016.
- 7 Yao, X., Liu, S., Han, L., Sun, M., and Zhao, L.: Definition and classification system of glacial lake for
- 8 inventory and hazards study, *J. Geogr. Sci.*, 28(2), 193-205, 2018.
- 9 Yao, T., Thompson, L., Wei, Y., Wusheng, Y., Yang, G., Xuejun, G., Xiaoxin, Y., Keqin, D., Huabiao,
- 10 Z., Baiqing, X., Jiancheng, P., Anxin, L., Yang, X., Dambaru B. K., and Daniel J.: Different glacier status
- 11 with atmospheric circulations in Tibetan Plateau and surroundings, *Nat. Clim. Chang*, 2 (9), 663–667,
- 12 doi:10.1038/nclimate1580, 2012.
- 13 Zhai, K., Wu, X., Qin, Y., and Du, P.: Comparison of surface water extraction performances of different
- 14 classic water indices using OLI and TM imageries in different situations. *Geo-spatial Information*
- 15 *Science*, 18(1), 32-42, doi: 10.1080/10095020.2015.1017911, 2015.
- 16 Zhang, G., Bolch, T., Alien, S., Linsbauer, A., Chen, W., and Wang, W.: Glacial lake evolution and
- 17 glacier–lake interactions in the Poiqu River basin, central Himalaya, 1964–2017, *Journal of Glaciology*,
- 18 65(251), 347–365, doi:10.1017/jog.2019.13, 2019.
- 19 Zhang, G., Yao, T., Xie, H., Wang, W., and Yang, W.: An inventory of glacial lakes in the Third Pole
- 20 region and their changes in response to global warming, *Global*, 131, 148–157, doi:
- 21 10.1016/j.gloplacha.2015.05.01, 2015.