

# Answers to the referees on the article "In-situ airborne measurements of atmospheric and sea surface parameters related to offshore wind parks in the German Bight"

Astrid Lampert<sup>1</sup>, Konrad Bärfuss<sup>1</sup>, Andreas Platis<sup>2</sup>, Simon Siedersleben<sup>3</sup>, Bughsin Djath<sup>4</sup>, Beatriz Cañadillas<sup>5</sup>, Robert Hunger<sup>1</sup>, Rudolf Hankers<sup>1</sup>, Mark Bitter<sup>1</sup>, Thomas Feuerle<sup>1</sup>, Helmut Schulz<sup>1</sup>, Thomas Rausch<sup>1</sup>, Maik Angermann<sup>1</sup>, Alexander Schwithal<sup>1</sup>, Jens Bange<sup>2</sup>, Johannes Schulz-Stellenfleth<sup>4</sup>, Thomas Neumann<sup>5</sup>, and Stefan Emeis<sup>3</sup>

<sup>1</sup>Institute of Flight Guidance, Technische Universität Braunschweig, Braunschweig, Germany

<sup>2</sup>Eberhard Karls University, Tübingen, Germany

<sup>3</sup>Karlsruhe Institute of Technology, Garmisch-Partenkirchen, Germany

<sup>4</sup>Helmholtz Center for Material and Coastal Research, Geesthacht, Germany

<sup>5</sup>UL International, Oldenburg, Germany

**Correspondence:** Astrid Lampert (Astrid.Lampert@tu-braunschweig.de)

## 1 Answers to referee 1

Dear Reviewer,

We thank you for your kind comments and for taking the time to consider our submitted manuscript. We have incorporated your advice into the revised manuscript that will be resubmitted. Below you will find our direct responses to the comments in normal letters. The comments are given in italic. Our changes to the text are additionally presented in quotation marks.

*This paper provides a description of a relatively unique and comprehensive dataset collected using an aircraft platform that contains measurements to evaluate the impact of offshore wind turbines on atmospheric properties. I believe that the paper is very appropriate for ESSD and that the material, in general, is well described. I did have several comments on the material in the paper that may help to improve the usefulness of the paper to readers outside of the team that collected these measurements. These are included below.*

The authors would like to thank the referee for this positive overall judgement.

*Specific Comments:*

15 - *Line 74: In the conversion to static temperature, are the pressure and temperature measurements co-located? It may be helpful to have some photographs or diagrams of the payload configuration on the aircraft to better understand how everything is laid out.*

We added a figure showing the nose boom with the sensors and added in the text: "the central sensor package is contained in

the nose boom (Fig. 1)."

20

- Line 102: *How does this 1.2 K uncertainty vary with temperature? How linear is this relationship? Is the 20 C value listed the instrument temperature, the air temperature or the surface temperature? Also, what impact does the vertical structure of temperature between the sensor and surface have on the quality of the measurement?*

We changed the text to: " It has an accuracy of  $\pm 1.2$  K at 20° C surface temperature and a temporal resolution of 20 Hz. If no  
25 clouds are between the sensor and the surface, the surface temperature measurements are not influenced by the atmospheric temperature or humidity distribution."

- Line 107: *Significantly more information could be provided on how surface deflection is calculated using aircraft attitude corrections.*

30 We added in the text: "From the point measurements in the scanner's coordinate system  $\begin{pmatrix} v_{x_{body}} \\ v_{y_{body}} \\ v_{z_{body}} \end{pmatrix}$ , aircraft attitude corrections using Eulerian angles  $\Psi, \Theta, \Phi$  are applied to rotate aircraft body fixed coordinates into the geodetic coordinate system (positive directions East, North, Up), which then are geolocated by applying the aircraft position  $\begin{pmatrix} p_{x_{body}} \\ p_{y_{body}} \\ p_{z_{body}} \end{pmatrix}$  in the following manner:

$$\begin{pmatrix} p_{x_{geo}} \\ p_{y_{geo}} \\ p_{z_{geo}} \end{pmatrix} = R_{geo}^{body}(-\Psi, -\Theta, -\Phi) \begin{pmatrix} v_{x_{body}} \\ v_{y_{body}} \\ v_{z_{body}} \end{pmatrix} + \begin{pmatrix} p_{x_{body}} \\ p_{y_{body}} \\ p_{z_{body}} \end{pmatrix} \quad (1)$$

Subsequently, the surface deflection  $\eta$  is calculated out of the georeferenced point cloud using mean sea level."

35

- Lines 120-126: *Given that the camera images are not publicly available and that they are not included with the main dataset described here, does it make sense to provide more than one sentence on them? I'm not sure that the current configuration aligns with ESSD policies about data availability (not that I disagree with not making the imagery public).*

The authors would like to present the whole measurement system of the campaign. If the information about the camera images  
40 is not available in the text, the readers will not know about these additional data. So we would like to leave the section in the text, together with the information in the data availability section.

- Lines 128-137: *It would be very useful to show some statistics on the regions sampled (e.g. distributions of flight altitudes, distributions of distance from a known shore point, distributions of distance from known wind-farm points).*

45 We added a column with the main flight altitude in the tables. We think that the map is the most useful illustration of the locations of the flight.

- *Figure 2: What are the flight tracks that end abruptly at the coastline? There are several clusters that clearly are going into/out of an airport, but then there are also several singular lines that don't seem to go anywhere.*

50 We added in the figure caption: "Flight tracks end when the data acquisition was shut down."

- *Lines 130-131: It might be useful to show a wind rose with these directions on one or both of the maps.*

We think that the wind rose is too much information for the map. Anyway measurements were not restricted to the indicated wind directions. This was the general idea, but we have several examples (e.g. Flight 5) where we chose the wind park differently due to other reasons (different flight pattern, etc.). We changed the text to: "Generally, flights were performed downwind of Amrumbank West for a wind direction sector of 80° to 200°, and downwind of Godewind for a sector from 160° to 350°. However, there are exceptions for particular reasons (e.g. during Flight 5 for consecutively probing the wakes of both wind parks, Flight 6 for investigating the changes of the wind field above the wind park)."

60 *It may also be useful to show the extent of German-controlled airspace.*

We included the permitted airspace in the figure.

- *Section 4.2: Without going into evaluation/analysis of the data, it might be useful to show an example dataset from one of these CROSS flights. One figure that illustrates the flight pattern followed and some of the structure that might be observed in the key quantities measured might be insightful for the reader.*

65

We added references for each flight pattern in the sections. Further, we added an example of the coastal effect.

- *Section 5: For all of these quantities, it might be interesting to pick a few altitudes (e.g. 100, 200, 500 m) and come up with distributions of the mean quantities at these altitudes to plot. This will provide insight into the heterogeneity of the conditions sampled (or lack thereof). One could also imagine looking at scatter plots comparing these mean values (e.g. mean 50 m temperature vs. stability or similar).*

70

According to the analysis shown in other publications based on the data set (Platis et al., 2018; Siedersleben et al., 2018a, b; Platis et al., 2019a; Cañadillas et al., 2019; Siedersleben et al., 2019), the parameters strongly depend on each other. E.g. stable conditions are associated with flow from land if the land surface is warmer than the sea surface. Therefore, we think that scatter plots and statistics that are not taking into account the meteorological situation will not be useful for further analyses.

75

*As an example, instead of showing the mean temperature profile and the range of values around that mean, perhaps it would be insightful to include the distribution of some "stability statistic" (e.g. LTS, etc.). Also, while the mean profiles are interesting, maybe it would also be interesting to show a time-height plot of all of these profiles to demonstrate when the flights took place and whether there were clusters of flights that had similar conditions.*

80

The interaction of the atmospheric boundary layer with wind parks and coastal effects is very complex, and scatter plots or time series alone do not take into account the interactions. Results of detailed scientific analyses can be found in Platis et al.

(2018); Siedersleben et al. (2018a, b); Platis et al. (2019a); Cañadillas et al. (2019); Siedersleben et al. (2019). As an example, we now include Fig. ??, which shows the development of the wind speed at the altitude 120 m depending on the fetch length for all profiles (the distance from the coast along the wind direction). No systematic behaviour is obvious. The large scatter indicates that more parameters are necessary to understand the development of wind speed with fetch length. One of them is stability.

- Line 185: What is "cut-in speed"?

90 We changed the text to: "The typical cut-in speed at which offshore wind turbines start producing power is around  $3 \text{ m s}^{-1}$ ."

- Section 5.3: Here again, it might be more insightful to look at some statistic (e.g. ratio of wind at 50 m to wind at 100 m) as a function of other variables and/or location.

As discussed above, we would prefer to leave statistical information to dedicated studies, as the interactions require complex analyses of different parameters. We added an example of the modification of the wind speed by coastal effects:

"The profiles of temperature and wind speed are modified by coastal effects (e.g. Dörenkämper et al. (2015); van der Laan et al. (2017)). During the WIPAFF campaign, climb and descent flights were performed on the way to the wind park measurement area and back. As an example of current and future research, the modification of the wind by coastal effects was investigated. Figure 2 shows the difference of the wind speed at hub height (120 m) for each profile minus the wind speed at hub height obtained during the profile closest to the coast. The fetch length is defined as the mean length that the air travelled above open water along the wind direction. Only flights are included where it was possible to determine the fetch length (not from North and West, as the distances to the next coast lines are too large). There is a large scatter in the data. Figure 3 shows the same data points. However, they are grouped by wind direction. There is still large scatter in the different data sets. However, depending on wind direction, the wind speed either increases or decreases with fetch length. This shows that there are more parameters required to explain the modification of wind speed besides the fetch length. A weakness of this analysis is that the profiles were not obtained along the mean wind direction. So air masses do not have the same origin, and besides the fetch length, variability along the coast line influences the results. More investigation is required to understand and parameterize the coastal effect."

- Line 189: Am I understanding correctly that these profiles include all data, and aren't necessarily at a single location?  
110 This makes these very difficult to interpret.

Yes, all available data of the vertical profiles are included, irrespective of the location. The figures intend to provide a rough overview of the meteorological conditions encountered during the campaigns. However, other ways of presentation (e.g. scatter plots) seem even more difficult to interpret, see above.

115 - Lines 191-192: This could be a very interesting 3-component plot (e.g. a scatter plot of wind direction vs. strength, color-coded by stability).

As described above, we included an example of the coastal effect. Fig. ?? shows the same information as Fig. ??, with addi-

tional information on the wind direction. The linear regressions are very different for each wind sector than for all data points. However, the scatter of the data points is still very large.

120

- *Lines 195-196: For unstable conditions, wouldn't you expect that mixing would result in an adiabatic profile that has an increasing RH with height?*

We agree with the referee, and changed the sentence to: "For unstable conditions, an enhanced water vapour mixing ratio directly above the water surface was present. "

125

- *Line 196: Is the increasing humidity with altitude for stable profiles the result of layered advection impacting temperature, or are there moisture plumes being advected? Or something else?*

We changed the sentence to: "For stable conditions, humidity was often increased at higher altitudes, which in most cases is most likely caused by advection of air masses with higher water vapour mixing ratio."

130

- *Section 6: I would have liked to have seen a bit more information in this conclusion. Some discussion on other complementary datasets, how these are expected to be used, etc. could be useful.*

We added in the conclusion section: "The unique data has been the base for different studies, proving for the first time directly the horizontal extension of wakes downwind of offshore wind parks (Platis et al., 2018), quantifying the recovery of wind speed in dependence of stability (Cañadillas et al., 2019; Platis et al., 2019a), and for the validation of the WRF mesoscale model (Siedersleben et al., 2018a, b, 2019), which can then be used for larger scales and scenario calculations."

*Also, surprisingly, there was no mention of anything going wrong/not working/etc. during the campaign, which would be very remarkable for a campaign of this extent. Did anything go wrong that the reader should know about?*

140 We changed the text to: " Altogether, 41 measurement flights were conducted during different seasons, wind direction, wind speed and stability. An overview of the flights performed during WIPAFF and meteorological conditions is shown in Tab. 1 and 2. A map with all flight paths flown during WIPAFF is provided in Fig. 3. During the flights, no instrument failures occurred. Only during one flight, the data acquisition had to be re-started (Flight 35)."


145 *Technical Corrections:*

- *None at this time. The grammar, while sometimes different than I would have personally used, is perfectly suitable and readable. Perhaps the editorial team finds reasons to reword/correct, but I did not come across anything that required being corrected/changed.*


Thank you!

## 150 References

- Cañadillas, B., Foreman, R., Barth, V., Siedersleben, S.K., Lampert, A., Platis, A., Djath, B., Schulz-Stellenfleth, J., Bange, J., Emeis, S., and Neumann, T.: Offshore wind farm wake recovery: Airborne measurements and its representation in engineering models, accepted to *Wind Energy*, 2019.
- Corsmeier, U., Hankers, R., and Wieser, A.: Airborne turbulence measurements in the lower troposphere onboard the research aircraft Dornier 128-6, *D-IBUF, Meteorol. Z.*, 10, 315-329, 2001.
- Dörenkämper, M., Optis, M., Monahan, A., and Steinfeld, G.: On the Offshore Advection of Boundary-Layer Structures and the Influence on Offshore Wind Conditions, *Boundary-Layer Meteorol* 155, 459–482, <https://doi.org/10.1007/s10546-015-0008-x>, 2015.
- van der Laan, M.P., Peña, A., Volker, P., Hansen, K.S., Sørensen, N.N., Ott, S., and Hasager, C.B.: Challenges in simulating coastal effects on an offshore wind farm, *IOP Conf. Series: Journal of Physics: Conf. Series*, 854, 012046, doi:10.1088/1742-6596/854/1/012046, 2017.
- 160 Platis, A., Siedersleben, S., Bange, J., Lampert, A., Bärfuss, K., Hankers, R., Cañadillas, B., Foreman, R., Schulz-Stellenfleth, J., Djath, B., Neumann, T., and Emeis, S.: First in situ evidence of wakes in the far field behind offshore wind farms, *Scientific Reports*, doi 10.1038/s41598-018-20389-y, 2018.
- Platis, A., Hundhausen, M., Siedersleben, S., Emeis, S., Lampert, A., Bärfuss, K., Djath, B., Schulz-Stellenfleth, J., Neumann, T., Cañadillas, B., and Bange, J.: Evaluation of a simple analytical model for offshore wind farm wake recovery by in-situ data and WRF simulations, *Wind Energy*, under review, 2019.
- 165 Platis, A., Siedersleben, S., Hundhausen, M., Bärfuss, K., Cañadillas, B., Schulz-Stellenfleth, J., Djath, B., Lampert, A., Neumann, T., Bange, J., and Emeis, S.: Long-range modifications of the wind field by offshore wind parks - results of the project WIPAFF, submitted to *Meteorologische Zeitschrift*, 2019.
- Siedersleben, S.K., Platis, A., Lundquist, J.K., Lampert, A., Bärfuss, K., Cañadillas, B., Djath, B., Schulz-Stellenfleth, J., Neumann, T., Bange, J., and Emeis, S.: Evaluation of a Wind Farm Parametrization for Mesoscale Atmospheric Flow Models with Aircraft Measurements, *Meteorologische Zeitschrift*, PrePub DOI 10.1127/metz/2018/0900, 2018a.
- 170 Siedersleben, S.K., Lundquist, J.K., Platis, A., Bange, J., Bärfuss, K., Lampert, A., Cañadillas, B., Neumann, T., and Emeis, B.: Micrometeorological impacts of offshore wind farm as seen in observations and simulations, *Env. Res. Lett.*, 13, 124012, 2018b.
- Siedersleben, S.K., Platis, A., Lundquist, J.K., Djath, B., Lampert, A., Bärfuss, K., Cañadillas, B., Schultz-Stellenfleth, J., Bange, J., Neumann, T., and Emeis, B.: Observed and simulated turbulent kinetic energy (WRF 3.8.1) overlarge offshore wind farms, *Geosci. Model Dev. Discuss.*, <https://doi.org/10.5194/gmd-2019-100>, in review, 2019.
- 175




**Figure 1.** Nose boom and instrumentation of the research aircraft Dornier 128-6 D-IBUF. Photo: Mark Bitter, TU Braunschweig



**Figure 2.** Changes of the wind speed at hub height (120 m) from the profile closest to the coast line to the other vertical profiles depending on fetch length. The vertical profiles of Flight 1, 3, 4, 5, 6, 7, 8, 9, 10, 13, 18, 19, 24, 30, 31, 32, 35, 36, 37, 38, 40, and 41 are included. Excluded were flights with wind direction from North or West, where no fetch length can be determined.





**Figure 3.** Changes of the wind speed at hub height as in Fig. 2. Different sectors of the wind direction are indicated in different colours.

**Table 1.** Overview of the WIPAFF measurement flights 1–20. The flight patterns are MEANDER(M), CROSS (C) or ABOVE (A) as indicated in Sect. 4. The wind parks are Godewind (GD) or Amrumbank West (AM). Information on cloud conditions is not always available (n.a.). Sentinel 1A and 1B satellite overpasses on the same day are indicated as well.

flight number	date	flight time [UTC] take off – landing	wind park	flight pattern	main flight altitude [m]	wind speed [m s <sup>-1</sup> ]	wind dir [°]	cloud conditions	satellite [UTC]
1	6 Sep 2016	12:13-15:20	AM	M	90	7	190	n.a.	-
2	7 Sep 2016	07:27-10:43	AM	M, C	90	4	210	n.a.	-
3	7 Sep 2016	12:06-13:59	AM	C	90, 120, 150	4	190	n.a.	-
4	8 Sep 2016	08:39-12:23	AM	M, C	60, 90, 120, 150	8	120	Ci	1A 17:09
5	9 Sep 2016	09:00-12:40	GD, AM	M	90, 120	6	240	n.a.	-
6	9 Sep 2016	13:42-17:10	AM	A	300	6	250	Cu	
7	10 Sep 2016	07:43-11:13	AM	M,C	60, 90, 120, 150, 200	7	190	clear sky	1A 05:41
8	10 Sep 2016	12:17-15:58	AM	M,C	60, 90, 120, 150, 200	4	190	Ci	-
9	30 Mar 2017	13:56-17:02	GD	M	120	15	240	As, Ci	1B 17:16
10	31 Mar 2017	13:36-16:59	GD	A	60, 90, 120, 200, 250	13	180	As, Ci	-
11	5 Apr 2017	13:42-16:33	GD	M,A	120, 200, 250	14	310	Sc	1A 17:17
12	6 Apr 2017	13:29-16:20	GD	M,A	120, 200, 250	10	310	Sc, As	-
13	9 Apr 2017	10:36-14:05	GD	M,C	60, 120, 200, 250	7	220	clear sky	-
14	9 Apr 2017	14:31-17:16	GD	C	60, 120, 200, 250, 350	4	200	clear sky	-
15	11 Apr 2017	09:15-13:09	GD	A	250, 300	8	280	Cu, As	-
16	11 Apr 2017	14:07-17:07	GD	M	120, 200	8	260	St, showers	1B 17:16
17	13 Apr 2017	11:23-14:40	GD	M,A	120, 250, 300	13	290	Cu	1B 05:48
18	17 May 2017	11:31-14:27	AM	C	90, 120, 150, 200, 250	8	110	Sc	-
19	17 May 2017	15:16-17:45	AM	M	220	12	120	Ac	1B 17:16
20	23 May 2017	07:53-10:41	GD	M,A	120, 250	6	250	Ci, Ac, Sc	-