

Dear Editor,

Please find a detailed response to the reviews of the manuscript essd-2019-23. We have copied below the comments of both referees, with our responses in black below.

We also attached a modified version of the manuscript, with differences highlighted in red and blue, at the end of this document.

Best Regards,
Kévin Fourteau on behalf of all co-authors.

Review 1 by Christo Buizert

Fourteau et al. publish detailed historical firm pycnometry records from three sites (Vostok, Greenland Summit and Law Dome DE-08) that were never published due to the passing of Jean-Marc Barnola. A porosity parameterization based on these data (published in Goujon et al. 2003) is widely used, and the paper describes the underlying data. These are very valuable observations that need to be published, and I commend Fourteau et al. for taking the effort to do so. I only have some recommendations for technical corrections.


I am very intrigued to see the differences in the close-off density at these three sites (Fig. 3). This suggests highest air content at DE08, and lowest at Vostok. This is indeed the observation of Martinerie et al. (1994). Do you think this is the underlying mechanism of the Martinerie et al. relationship between air content and temperature? This is certainly what is implied by Fig. 5. I think it would be appropriate to address this briefly in the paper.

Martinerie et al (1994) report that not only air content increases with temperature, but also that the porous volume at close-off (V_i in their article) increases with temperature. This is consistent with the pycnometry data presented here, with warmer sites closing at lower densities and therefore with a larger porous volume. This is however inconsistent with the results of Schaller et al (2017), essentially showing that the porous volume at close-off is site-independent. Further work is required to explain this discrepancy, and goes beyond the scope of this paper.

We will address this point in the paper **P5L30**:

“It is interesting to note that the pycnometry data indicate that a cold site like Vostok reaches pore close-off at a higher density than a warm site like DE08-2. This is consistent with the results of Martinerie et al. (1994) that indicate an increase of porous volume at close-off with temperature based on air content measurements.”

Would it be possible to show the newer pycnometry data from the “Lock-In” site (Fourteau et al. 2019) somewhere together with the data from the three old sites (for example as a third panel in Fig. 3)? One would expect them to look more like Vostok than like Summit or DE08. Is that indeed the case? Showing these data would also give us a way to compare the consistency of the older and newer data.



As seen in the Figure of this response, the Lock-In pycnometry data are closer to the Summit values. However, one should be aware that the Lock-In data have not been submitted to an alpha correction similar to the datasets presented here, as this was not part of the experimental protocol during the Lock-In measurements in 2018. However, the Lock-In pycnometry data have been confirmed using tomography scanning of some of the samples.

On the other hand, if one uses air content data to derive the porous volume at close-off, the close-off at Lock-In is expected to happen at a porosity closer to the one of Vostok. This indicates that a discrepancy subsists between the close porosity data and the measured air content values.

These are for sure interesting questions that deserve dedicated work. However, we think that discussing them in this short article will obscure our main goal here, which is to make these historical data accessible, along with the protocol used to obtain them.

[Could you give some examples of mechanisms that can cause system drift in the pycnometry measurements? Are you talking about vapor freezing onto the surfaces, snow getting stuck, etc? Or is it something else?](#)

We were not able to specifically identify the physical mechanism at the origin of this system drift, and J.M. Barnola notebooks did not provide any hypothesis for it neither. Our understanding is that this missing mechanism is one of the reasons why J.M. Barnola did not publish the data in the 90's.

Figure 1 of the article shows that the alpha coefficient tends to be lower than one. This indicates that the system drift usually leads to an overestimation of the inaccessible volume V_s . This overestimation could be due to an overestimation of the volume chamber V_2 . One could imagine that this overestimation of V_2 comes from the presence of firm dust or frozen surfaces in the chamber, that would decrease its actual volume. However, this is largely speculative at this point as we do not have any supporting observations. We will write clearly in the text that we do not know the physical mechanism responsible for the system drift, **P4L30**:

“One should note that we were not able to identify the physical mechanism at the origin of the pycnometry system drift. Further work should be dedicated to this topic.”

We will correct the technical points below, following the recommendations of the referee.

P1L8: remove “intend to”

P1L20: remove plural “s” from “precipitations”

P2L6 and P2L7: replace “reach” with “connect”

P2L6: airtight is not the right word, given that fugitive gases like Ne still escape. I think you can just remove the phrase “and are therefore airtight”.

P4L4: “on the field” should be “in the field”

P5L2: “depending of” should be “depending on”

Review 2 by Johannes Freitag

Fourteau et al present historical porosity data of firn from three polar sites originally measured by J.-M. Barnola with the means of gas pycnometry several years ago. These data were the base of the important closed porosity-density parametrization widely used in firn gas models. Fourteau et al. provide the raw data and give insight into the original data processing. They confirm the use of the dataset for the closed-porosity parametrization first introduced in Goujon et al. (2003) and highlight its limitation. I appreciate the reworking of this fundamental data set.

The manuscript is well written with an appropriate introduction and description of the method. The data processing is described in required detail to enable the reader to follow the analysis. The data set is accessible via the given identifier and complete. I highly recommend the manuscript for publication. In my opinion the data set will encourage further methodological improvements and investigations on the cut-bubble-effect. Maybe it would be worthwhile to set up a future study where different methods like X-ray tomography and pycnometry are applied on the same set of samples? We strongly agree that a joint study of pycnometry and large-scale tomography is currently one of the best option to study the cut-bubble effect in firn samples, and to check the consistency between pycnometry and tomography-based data.

Specific comments: 1) The names of the columns in the data set are a little bit misleading. The pore volumes are named by "Poros_xx" like "Poros_closed_cm_3_" which might be interpreted as abbreviation for porosity (no units) instead of pore volumes (unit in cm₃). I would suggest names like "Pore_vol_xx". 2) The column "Pores_frac" should be renamed to "ClosedPorosRatio" as it is defined in the manuscript. 3) In the data sets there are some non-physical values like negative pore volumina or closed pore ratios larger than 1. I would prefer to assign them to the physical limits (0 or 1). Unfortunately, it is ESSD policy to have the data registered before the submission of the paper and PANGAEA policy not to modify datasets once they are registered. We will therefore not be able to modify the data submitted to the PANGAEA database. We will add a sentence in the article highlighting the difference in naming between the database and the article **P9L10**:
"Finally, we made these data publicly available on the PANGAEA database (Fourteau et al., 2019a). Note that the naming convention used in the database is different from the one used in the article."

We however do not agree that the data should be clipped to only have closed porosity ratios between 0 and 1. This would artificially reduce the experimental dispersion of the pycnometry method and introduce a bias in the data for fully open samples (respectively fully closed), as positive errors would no longer be statistically compensated by negative errors (respectively negative errors compensated by positive errors). We will add a sentence clarifying the presence of non-physical values in the dataset **P7L3**:
"Finally, because of experimental dispersion, some firn samples were measured with a closed pore volume below zero or above the total porous volume. Potential users of the data should be aware that these values are not physically sound, and reflects the experimental errors of the pycnometry method."

Historical porosity data in polar firn

Kévin Fourteau¹, Laurent Arnaud¹, Xavier Faïn¹, Patricia Martinerie¹, David M. Etheridge², Vladimir Lipenkov³, and Jean-Marc Barnola^{†1}

¹Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, F-38000 Grenoble, France

²Climate Science Centre, CSIRO Oceans and Atmosphere, Aspendale, Victoria, Australia

³Climate and Environmental Research Laboratory, Arctic and Antarctic Research Institute, St. Petersburg, 199397, Russia

[†] deceased, September 21, 2009

Correspondence: laurent.arnaud@univ-grenoble-alpes.fr

Abstract. In the 1990's, closed and open porosity volumes of firn samples have been measured by J.-M. Barnola using the technique of gas pycnometry, on firn from three different polar sites. They are the basis of a parameterization of closed porosity in polar firn, first introduced in Goujon et al. (2003) and used in several firn physics models (e.g. Buizert et al., 2012). However, these data and their processing have not been published in their own right yet. In this short article, we detail how they were processed by J.-M. Barnola, and how the closed porosity parameterization was obtained. We show that the original data processing only partially accounts for the presence of re-opened bubbles in the samples. Since the proper correction to apply for this effect is hard to estimate, we also processed the data without including a correction for re-opened bubbles. Finally, we ~~intend to make~~ made these pycnometry data available, in order to be used by the glaciology community, notably for the study of polar ice formation and of the composition of gas records in ice cores. They are hosted on the PANGAEA database: <https://doi.pangaea.de/10.1594/PANGAEA.907678> (Fourteau et al., 2019a).

Copyright statement. TEXT

1 Introduction

The enclosing of atmospheric air in the ice of polar regions is of great importance for the study of past climates. Indeed, ice cores drilled in the polar regions have the unique characteristics of containing bubbles of air from past atmospheres. They have thus been used to reconstruct the atmosphere history in major greenhouse gas concentrations for the last 800,000 years (Lüthi et al., 2008; Loulergue et al., 2008). However, in order to properly interpret the gas records from ice cores, it is necessary to understand the trapping of air in the ice (~~Schwander and Stauffer, 1984; Stauffer et al., 1985; Schwander et al., 1993; Rommelaere et al., 1997; Trudinger et al., 1997, 2002~~). (Schwander and Stauffer, 1984; Stauffer et al., 1985; Schwander et al., 1993; Rommelaere et al., 1997; Trudinger et al., 1997, 2002). The snow at the surface of polar ice sheets is a porous material, and interstitial air can freely exchange with the atmosphere (Stauffer et al., 1985). Snow strata are progressively buried under new ~~precipitations~~ precipitation and are compressed due to the weight of the younger snow above. This buried, metamorphosed, and compacted snow is referred to as firn. With time the

firn strata are further buried and their interstitial porous networks shrink. Depending on the local temperature and accumulation conditions, some of the pores start to pinch and encapsulate the interstitial air at depths between 50 and 100m below the surface (Witrant et al., 2012). The porous network then continues to close until all the interstitial air is isolated from the atmosphere. The firn then becomes airtight ice with enclosed bubbles of atmospheric air.

5 One way to characterize the closing of the porous network and the trapping of gases is to measure the closed and the open pore volumes. Closed pores are pores that no longer ~~reach-connect~~ to the atmosphere, ~~and are therefore airtight~~. On the other hand, open pores ~~reach-up-connect~~ to the atmosphere, through an interconnected porous network. Moreover, data of closed and open pore volumes are required for the usage of gas trapping models (Rommelaere et al., 1997; Goujon et al., 2003; Buizert et al., 2012; Witrant et al., 2012). That is why closed and open volumes have been measured along various firn columns from
10 Greenland and Antarctica (Schwander and Stauffer, 1984; Schwander et al., 1993; Schaller et al., 2017). Such measurements were notably performed in the 1990's by J.-M. Barnola on firn cores drilled at the three polar sites of Vostok (Antarctica), Summit (Greenland) and DE08-2 (Antarctica), using the technique of gas pycnometry (Schwander and Stauffer, 1984; Stauffer et al., 1985). The three sites have very different characteristics, from the cold and low-accumulation site of Vostok to the high-accumulation site of Law Dome's DE08-2. Their respective accumulation rates and temperatures
15 are given in Table 1. The three sites have been exploited for the range of environmental information they contain, from recent decades to more than 400,000 years before present (~~Barnola et al., 1987; Schwander et al., 1993; Trudinger et al., 2002~~) (Barnola et al., 1987; Schwander et al., 1993; Trudinger et al., 1997). The obtained porosity data have been widely used to parameterize closed porosity as a function of density in firn physics models (~~Trudinger et al., 2002; Goujon et al., 2003; Buizert et al., 2012~~) (Trudinger et al., 1997; Goujon et al., 2003; Buizert et al., 2012; Witrant et al., 2012). Unfortunately, J.-M. Barnola passed
20 away before publishing the data in the peer-reviewed literature. For transparency, and to recognize J.-M. Barnola's effort, care and foresight in undertaking the measurements at three remote sites, we decided to make them available. Our goal is also to provide an explanation of how the closed porosity parameterization proposed in Goujon et al. (2003) was derived. Moreover, measuring closed porosity is labor intensive and requires a large amount of firn material. As a result, this type of data is rather scarce. We hope that making these data available will help the ice core community to better understand the trapping of gases
25 in polar ice.

2 The pycnometry method

The technique used by J.-M. Barnola to measure the closed and open porosity volumes in firn samples is the gas pycnometry method (Schwander and Stauffer, 1984; Stauffer et al., 1985). The pycnometry apparatus is composed of two airtight chambers of known volumes V_1 and V_2 with a valve between them allowing to either connect or isolate the chambers. A scheme is
30 provided in the Supplementary Material of Fourteau et al. (2019b). A pressure gauge is joined to the chamber V_1 to monitor its internal pressure. For the measurements, a firn sample is placed in the first chamber V_1 , while the second one is isolated and vacuum-pumped. Placing a firn sample in the chamber V_1 renders a volume V_s inaccessible to the gases. The pressure P in V_1 is recorded. Then, the two chambers are connected, allowing the gas in chamber V_1 to expand in a larger volume.

Table 1. Characteristics of the three investigated polar sites

Site Name	Location	Accumulation($\text{g cm}^{-3} \cdot \text{yr}^{-1}$)	Temperature ($^{\circ}\text{C}$)
Vostok	East Antarctic plateau	2.2 ^a	-56 ^a
Summit	Central Greenland	20.9 ^b	-31 ^b
DE08-2	Coastal East Antarctica	110 ^c	-19 ^d

a: Lipenkov et al. (1997)

b: Schwander et al. (1993)

c: Etheridge et al. (1996)

d: Etheridge et al. (1992)

The pressure after expansion P' is then recorded. The system was designed to minimize the pressure drop, in order to avoid rupturing recently closed and still fragile pores. The volume V_s can be related to the recorded pressure by:

$$V_s = V_1 - \frac{R}{1-R} V_2 \quad (1)$$

where $R = P'/P$.

- 5 The protocol followed by J.-M. Barnola was to first execute an expansion without any sample in the first chamber, then a second expansion with the sample in. In this case, Equation 1 can be rearranged as:

$$V_s = V_2 \left(\frac{R_0}{1-R_0} - \frac{R_1}{1-R_1} \right) \quad (2)$$

where R_0 and R_1 are respectively the pressure ratio in the cases without and with the firm sample in V_1 .

- 10 The volume inaccessible to gases is composed of the ice phase and of the closed pores phase. Therefore, one can deduce the closed and open porosity volumes:

$$\begin{aligned} V_{cl} &= V_s - V_{ice} \\ V_o &= V_{cyl} - V_s \end{aligned} \quad (3)$$

- where V_{cl} and V_o are the closed and open porosity volumes, V_{ice} the volume of the ice phase and V_{cyl} the volume of the firm sample. The volume of the ice phase is deduced from the mass M of the sample knowing that $V_{ice} = M/\rho_{ice}$, where ρ_{ice} is the density of pure ice. The density of pure ice is estimated using the temperature relationship $\rho_{ice} = 0.9165(1 - 1.53 \times 10^{-4} T)$, where ρ_{ice} is expressed in g cm^{-3} and T is the temperature expressed in $^{\circ}\text{C}$ (Bader, 1964; Goujon et al., 2003). The volume of the firm sample V_{cyl} is measured geometrically with calipers.

Note that in the pycnometry experiment, all the pores reaching the edge of the sample are considered as open. This means that some pores that are closed in the firm column (they do not reach the atmosphere) will be considered open during the pycnometry

measurement. This is known as the cut-bubble effect and leads to an underestimation of the closed porosity (Martinerie et al., 1990; Schaller et al., 2017).

The firm samples used for the pycnometry measurements are cylindrical samples of about 4 to 5cm in height and diameter. They were produced by machining on a lathe and trimmed with a drop saw in order to produce well-shaped cylinders. The measurements were performed in environments with a good temperature stability, to limit the effect of temperature variations. Finally, in order to avoid post-coring effects, all the samples were measured directly ~~on~~in the field, shortly after the drilling of the firm core.

3 Processing of the data

For each of the three sites we retrieved a computer file containing the expansion ratios R_0 and R_1 , the mass of the firm samples, their volumes, and the temperature during the experiment. We also retrieved the source codes that J.-M. Barnola used to process the data. Finally, we have the experiment notebooks of J.-M. Barnola.

3.1 Original data processing

In this section, we aim to reproduce the data processing performed by J.-M. Barnola. This is done for two reasons. First, the data have been used to derive a parameterization of the closed porosity in polar firm (Goujon et al., 2003), and it is therefore important to understand how they were processed. Second, the original processing includes corrections for experimental biases that were observed by J.-M. Barnola and that have to be taken into account.

The original source codes indicate that the processing included a correction to account both for a pycnometry system drift and the cut-bubble effect. This correction is based on the idea that the pycnometry method should ideally indicate fully open samples at low density ($\rho < 0.72\text{g.cm}^{-3}$) and fully closed ones at high density ($\rho > 0.86\text{g.cm}^{-3}$). A correcting factor α to be applied to the inaccessible volume V_s can thus be determined for each of these low and high-density samples. The α factors are such that the measurements of low-density firm samples (respectively high-density firm samples) yields a fully open (respectively closed) porosity. In the case of low-density firm, α is computed as $1 - V_{c1}/V_{ice}$, and in the case of high density firm $\alpha = V_{cy1}/V_s$. J.-M. Barnola observed that the α factors are linearly related to the empty expansion ratio R_0 (Figure 1). This might come as a surprise as R_0 is measured before the sample is inserted in the apparatus. Our understanding is that it reflects that α and R_0 are simultaneously affected by system drifts. By monitoring the evolution of R_0 , one is able to estimate the correcting factor α to be applied. However, as seen in Figure 1, the linear relationship is not the same for high density and low density firm samples. J.-M. Barnola derived intermediate linear relationships, in black in Figure 1. Thus, for each firm sample (including mid-density samples), a correcting factor is determined thanks to the preceding empty expansion and applied to the inaccessible volume V_s . Finally, in all of the three measurement campaigns the volume V_2 has been estimated to be of 7.2cm^3 (value found directly in J.-M. Barnola's processing codes). One should note that we were not able to identify the physical

mechanism at the origin of the pycnometry system drift. Further work should be dedicated to this topic.

Vostok measurement campaign:

5 The Vostok measurements were performed on the BH3 firn core, drilled during the austral summer 1991/92. For this measurement campaign, J.-M. Barnola observed a bias due to the measurement of sample volumes with a caliper, depending ~~of~~on the pressure applied by the caliper. He therefore proposed to apply a volume correction of 0.9% to the firn samples that were measured by applying a too weak pressure with the caliper. He also applied a 4°C correction to the recorded temperatures, in order to account for the heat dissipation of heated elements towards the temperature sensor. For the Vostok campaign measurements α is computed following $\alpha = -40.2145R_0 + 39.22304$, shown in black in the left panel of Figure 1. These three
10 corrections were found hard coded in the Vostok processing source code, and have been corroborated by the notebooks.

Summit measurement campaign:

15 The firn porosity measurement of Summit were performed in the framework of the 1989 Eurocore project. For this campaign the same volume correction of 0.9% was applied to all the samples. Moreover, a weighting bias was found by J.-M. Barnola and is taken into account by applying a 0.9983 correction factor to the measured mass. Finally, the α correction is given by $\alpha = -45.045045R_0 + 43.820459$, in black in the middle panel of Figure 1. It is interesting to note that for Summit the correction chosen by J.-M. Barnola seems to be primarily based on the low density α only.

20

DE08-2 measurement campaign:

The DE08-2 measurements were performed during the austral summer 1992/93. For DE08-2, no volume or mass correction is reported in the original processing code. The α correction is given by $\alpha = -37.2577R_0 + 36.4204$, in black in the right panel
25 of Figure 1. Again, the correction chosen by J.-M. Barnola appears to be primarily based on the low density α .

From these data, we can deduce the closed and open porosity volumes. Figure 2 displays closed porosity and closed porosity ratio values against total porosity. Closed porosity is defined as the volume fraction occupied by the closed pores in the firn sample, total porosity is defined as the volume fraction of all pores and the closed porosity ratio is the ratio of the closed pores
30 volume over the total porous volume. We chose to use volume fractions instead of porous volumes per gram of firn, as the former are not sensitive to temperature and therefore renders the comparison between sites easier. Yet, the volume fraction data can easily be converted to porous volumes per gram of firn using the density of pure ice. It is interesting to note that the pycnometry data indicate that a cold site like Vostok reaches pore close-off at a higher density than a warm site like DE08-2. This is consistent with the results of Martinerie et al. (1994) that indicate an increase of the porous volume at close-off with
35 temperature based on air content measurements. Finally, because of experimental dispersion, some firn samples were measured

with a closed pore volume below zero or above the total porous volume. Potential users of the data should be aware that these values are not physically sound, and reflect the experimental errors of the pycnometry method.




Figure 1. In each panel: the blue dots are the correcting factors α computed for low density firm as a function of the preceding R_0 , the orange dots are the correcting factors α computed for high density firm as a function of R_0 , the black line is the linear relation originally derived by J.-M. Barnola, and the blue line is the linear regression based solely on the low density α




Figure 2. Porosity data obtained using J.-M. Barnola processing chain. Left panel: closed porosity against total porosity. The dashed line represents where the closed porosity equals the total porosity. Right panel: closed porosity ratio over total porosity. The dashed line indicates where the closed porosity ratio equals one.

3.2 Uncertainty analysis

- Unfortunately, the original data we retrieved do not allow us to perform a systematic uncertainty analysis. Indeed, we did not retrieve a quantification of the uncertainties of the raw measurements, such as the sample's mass or volume. We are therefore not able to propagate the uncertainties of the raw measurements to the final derived quantities, such as the density or the closed porosity ratio.

However, a recent pycnometry campaign was conducted for an East Antarctic firn core, including a dedicated uncertainty analysis (Section S1.3 of the Supplement in Fourteau et al., 2019b). The measurements were performed with the same pycnometry apparatus and with the same sample size as the data presented in this article. We can therefore expect the data of JM Barnola to be affected by similar uncertainties. Fourteau et al. (2019b) quantified the errors associated with the measurements performed in the cold room and used to derive the density and the closed porosity of the samples. This includes the errors on the mass of the sample, its radius and height, the pressures of the chambers, and the volumes of the chambers. These errors were then propagated to obtain an estimation of the uncertainty of the density and closed porosity of the samples. Fourteau et al. (2019b) analysis indicates that the uncertainty on density is fairly constant over the entire range of measurements with a value of 0.0082 g cm^{-3} , that is to say an uncertainty of about 0.009 on density relative to pure ice. This represents a relative uncertainty of about 1% on the derived density. Note that this is of the same order as the correction applied by J.-M. Barnola to the volumes of the Vostok samples. Contrary to density, the uncertainty of the closed porosity ratio is not constant over the entire range of data, and increases from about 0.02 for low-density samples to about 0.2 for high-density samples. For both quantities, the dominant contribution to the final uncertainty is the uncertainty of the measured sample volume.

3.3 A new data processing

We identify one major issue in the processing elaborated by J.-M. Barnola. In the case of high density firn samples, determining a correcting factor with $\alpha = V_{\text{cyl}}/V_{\text{s}}$ both encapsulates the effect of a system drift and of cut-bubbles. Indeed, the assumption under which this factor is computed is that the pycnometry experiment should measure a fully closed sample at high density, de facto including a cut-bubble correction. However, the correction to be applied for cut-bubbles is not the same at all densities (Schaller et al., 2017). It thus explains why the high and low density α relationship with R_0 might differ. On the other hand, the low density α does not include any cut-bubble correction, and therefore should only account for system drifts. We therefore propose to correct the data using a linear regression between R_0 and the low density α only. These corrections are displayed as blue lines in Figure 1. As shown in the figure, the new corrections mainly differ in the Vostok case. The closed porosity and closed porosity ratio after applying this new correction are displayed in Figure 3.

It is important to note that these data are not corrected for cut bubbles, and therefore do not directly represent the amount of closed pores in the firn column. We decided not to correct the data for cut bubbles in this article. Indeed, the appropriate corrections are hard to estimate and potentially site dependent (Schaller et al., 2017). Recently, Schaller et al. (2017) reported a fraction of re-open bubbles of up to 60% for similar size B53 firn samples (East Antarctic plateau). Further research is needed to fully solve this problem.




Figure 3. Same as Figure 2, with the new α correction.

4 The Barnola parameterization for closed porosity

The firm densification and gas trapping model of Goujon et al. (2003) uses a parameterization of closed porosity proposed by J.-M. Barnola (Equation 9 of Goujon et al., 2003). This Barnola parameterization relates the closed porosity to the total porosity with:

$$5 \quad P_{\text{closed}} = \gamma P_{\text{total}} \left(\frac{P_{\text{total}}}{P_{\text{close-off}}} \right)^{-7.6} \quad (4)$$

where P_{closed} is the closed porosity, P_{total} the total porosity, $P_{\text{close-off}}$ the close-off porosity that can be estimated using air content measurements or a temperature regression (Martinerie et al., 1994), and γ a factor valued at 0.37.

We are confident that the Barnola closed porosity parameterization was deduced from the pycnometry data described in Section 3.1, with the original processing chain. Indeed, there is a clear linear relationship between the logarithm of the closed porosity ratio and the logarithm of the total porosity normalized by the porosity at mean close-off deduced from air content data. This ~~is~~ relation, displayed in Figure 4, is consistent with the Barnola parameterization. The comparison between the experimental closed porosities and the Barnola parameterization is also displayed in Figure 5. It is therefore important to acknowledge that the Barnola parameterization is based on data that are not fully corrected for cut-bubbles. Future users of this parameterization should be aware of this potential limitation. However, since we are not able to properly estimate the corrections to be applied for cut bubbles, we cannot propose a new law replacing the Barnola parameterization at this point.

5 Conclusions

We evaluated the pycnometry data from three polar sites obtained in the 1990's by J.-M. Barnola. Based on original computer files, including raw data and processing source codes, we were able to reproduce the processing chain developed by J.-M.




Figure 4. Relationship between the logarithm of the closed porosity ratio and the logarithm of the total porosity. The solid black line corresponds to the Barnola parameterization. The closed porosity data were obtained with the original processing chain.




Figure 5. Measured closed porosity in the Vostok, Summit and DE08-2 firns with the original processing chain. The corresponding Barnola parameterizations are displayed as solid lines.

Barnola, including experimental bias corrections. We found that these data have not been fully corrected for the cut-bubble effect. We also confirm that the closed porosity data deduced from those pycnometry experiments were used to derive the Barnola closed porosity parameterization, first introduced in Goujon et al. (2003). Consequently, this parameterization suffers from the incomplete cut-bubble correction of the pycnometry data. More work is needed to quantify the amount of re-open
 5 bubbles in firn samples, but recent work highlighted a fraction of re-opened bubbles reaching up to 60% (Schaller et al., 2017).

Finally, we ~~intend to make~~ made these data publicly available ([Fourteau et al., 2019a](#)) on the [PANGAEA database](#) ([Fourteau et al., 2019a](#)). Note that the naming convention used in the database is different from the one used in the article. The three sites studied in this article are characterized by a wide range of accumulation rates and temperatures. Such type of data are crucial to understand
 10 the age, amount and composition of the air enclosed in polar ice sheets. They could be useful for future studies focusing on

the effect of the climatic conditions on pore closure and gas trapping, as well as to interpret long term ice core atmospheric records.

Code availability. The codes used to process the data were developed using python3. They will be provided upon direct request to the corresponding authors.

- 5 *Data availability.* The pycnometry datasets generated with the new correction methodology are hosted on the PANGAEA database: <https://doi.pangaea.de/10.1594/PANGAEA.907678> (Fourteau et al., 2019a). The datasets generated using the original JM Barnola correction methodology will be provided upon direct request to the corresponding authors.

- 10 *Author contributions.* The pycnometry measurements were performed by JMB with the help of DME and VL. The original data files and notebooks were retrieved by LA, XF and PM. The reconstruction of the original processing chain was done by LA, XF, KF and PM. The codes to process the data were developed by KF. All authors contributed to the interpretation of the data. The manuscript was written by KF with the help of all the co-authors.

Competing interests. The author declare having no competing interests.

Disclaimer. TEXT

- 15 *Acknowledgements.* We are grateful to Jakob Schwander for his help for the pycnometry measurements of the Summit site. We thank the 1991/92 Vostok, 1989 Eurocore, and 1992/93 DE08-2 at Law Dome ice core programs, as well as the field personal that contributed to these successful shallow drilling operations. We finally acknowledge Olivier Magand for his help retrieving the original notebooks, and his tests of the pycnometry method.

References

- Bader, H.: Density of ice as a function of temperature and stress, Cold Regions Research and Engineering Laboratory, US Army Material Command, 1964.
- Barnola, J.-M., Raynaud, D., Korotkevich, Y. S., and Lorius, C.: Vostok ice core provides 160,000-year record of atmospheric CO₂, *Nature*, 5 329, 408, <https://doi.org/10.1038/329408a0>, 1987.
- Buizert, C., Martinerie, P., Petrenko, V. V., Severinghaus, J. P., Trudinger, C. M., Witrant, E., Rosen, J. L., Orsi, A. J., Rubino, M., Etheridge, D. M., Steele, L. P., Hogan, C., Laube, J. C., Sturges, W. T., Levchenko, V. A., Smith, A. M., Levin, I., Conway, T. J., Dlugokencky, E. J., Lang, P. M., Kawamura, K., Jenk, T. M., White, J. W. C., Sowers, T., Schwander, J., and Blunier, T.: Gas transport in firm: multiple-tracer characterisation and model intercomparison for NEEM, Northern Greenland, *Atmos. Chem. Phys.*, 12, 4259–4277, 10 <https://doi.org/10.5194/acp-12-4259-2012>, 2012.
- Etheridge, D. M., Pearman, G. I., and Fraser, P. J.: Changes in tropospheric methane between 1841 and 1978 from a high accumulation-rate Antarctic ice core, *Tellus B Chem. Phys. Meteorol.*, 44, 282–294, <https://doi.org/10.3402/tellusb.v44i4.15456>, 1992.
- Etheridge, D. M., Steele, L. P., Langenfelds, R. L., Francey, R. J., Barnola, J.-M., and Morgan, V. I.: Natural and anthropogenic changes in atmospheric CO₂ over the last 1000 years from air in Antarctic ice and firm, *J. Geophys. Res. Atmos.*, 101, 4115–4128, 15 <https://doi.org/10.1029/95JD03410>, 1996.
- Fourteau, K., Arnaud, L., Faïn, X., Martinerie, P., Etheridge, D., Lipenkov, V., and Barnola, J.-M.: Gas pycnometry firm porosity data of firm ice cores from three polar sites, <https://doi.pangaea.de/10.1594/PANGAEA.907678>, 2019a.
- Fourteau, K., Martinerie, P., Faïn, X., Schaller, C. F., Tuckwell, R. J., Löwe, H., Arnaud, L., Magand, O., Thomas, E. R., Freitag, J., Mulvaney, R., Schneebeli, M., and Lipenkov, V. Y.: Multi-tracer study of gas trapping in an East Antarctic ice core, *Cryosphere Discuss.*, 2019, 1–34, 20 <https://doi.org/10.5194/tc-2019-89>, 2019b.
- Goujon, C., Barnola, J.-M., and Ritz, C.: Modeling the densification of polar firm including heat diffusion: application to close-off characteristics and gas isotopic fractionation for Antarctica and Greenland sites, *J. Geophys. Res. Atmos.*, 108, 4792, <https://doi.org/10.1029/2002jd003319>, 2003.
- Lipenkov, V. Y., Salamatin, A. N., and Duval, P.: Bubbly-ice densification in ice sheets: II. Applications, *J. Glaciol.*, 43, 397–407, 25 <https://doi.org/10.1017/S0022143000034973>, 1997.
- Loulergue, L., Schilt, A., Spahni, R., Masson-Delmotte, V., Blunier, T., Lemieux, B., Barnola, J.-M., Raynaud, D., Stocker, T. F., and Chappellaz, J.: Orbital and millennial-scale features of atmospheric CH₄ over the past 800,000 years, *Nature*, 453, 383–386, <https://doi.org/10.1038/nature06950>, 2008.
- Lüthi, D., Floch, M. L., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura, K., 30 and Stocker, T. F.: High-resolution carbon dioxide concentration record 650,000– 800,000 years before present, *Nature*, 453, 379–382, <https://doi.org/10.1038/nature06949>, 2008.
- Martinerie, P., Lipenkov, V. Y., and Raynaud, D.: Correction of air-content measurements in polar ice for the effect of cut bubbles at the surface of the sample, *J. Glaciol.*, 36, 299–303, <https://doi.org/10.3189/002214390793701282>, 1990.
- Martinerie, P., Lipenkov, V. Y., Raynaud, D., Chappellaz, J., Barkov, N., and Lorius, C.: Air content paleo record in the Vostok ice core (Antarctica): A mixed record of climatic and glaciological parameters, *J. Geophys. Res. Atmos.*, 99, 10 565–10 576, 35 <https://doi.org/10.1029/93JD03223>, 1994.

- Rommelaere, V., Arnaud, L., and Barnola, J.-M.: Reconstructing recent atmospheric trace gas concentrations from polar firn and bubbly ice data by inverse methods, *J. Geophys. Res. Atmos.*, 102, 30 069–30 083, <https://doi.org/10.1029/97jd02653>, 1997.
- Schaller, C. F., Freitag, J., and Eisen, O.: Critical porosity of gas enclosure in polar firn independent of climate, *Clim. Past*, 13, 1685–1693, <https://doi.org/10.5194/cp-13-1685-2017>, 2017.
- 5 Schwander, J. and Stauffer, B.: Age difference between polar ice and the air trapped in its bubbles, *Nature*, 311, 45–47, <https://doi.org/10.1038/311045a0>, 1984.
- Schwander, J., Barnola, J.-M., Andrié, C., Leuenberger, M., Ludin, A., Raynaud, D., and Stauffer, B.: The age of the air in the firn and the ice at Summit, Greenland, *J. Geophys. Res. Atmos.*, 98, 2831–2838, <https://doi.org/10.1029/92jd02383>, 1993.
- 10 Stauffer, B., Schwander, J., and Oeschger, H.: Enclosure of air during metamorphosis of dry firn to ice, *Ann. Glaciol.*, 6, 108–112, <https://doi.org/10.3189/1985AoG6-1-108-112>, 1985.
- Trudinger, C. M., Enting, I. G., Etheridge, D. M., Francey, R. J., Levchenko, V. A., Steele, L. P., Raynaud, D., and Arnaud, L.: Modeling air movement and bubble trapping in firn, *J. Geophys. Res. Atmos.*, 102, 6747–6763, <https://doi.org/10.1029/96jd03382>, 1997.
- Trudinger, C. M., Etheridge, D. M., Rayner, P. J., Enting, I. G., Sturrock, G. A., and Langenfelds, R. L.: Reconstructing atmospheric histories from measurements of air composition in firn, *J. Geophys. Res. Atmos.*, 107, ACH 15–1–ACH 15–13, <https://doi.org/10.1029/2002JD002545>, 2002.
- 15 Witrant, E., Martinerie, P., Hogan, C., Laube, J. C., Kawamura, K., Capron, E., Montzka, S. A., Dlugokencky, E. J., Etheridge, D., Blunier, T., and Sturges, W. T.: A new multi-gas constrained model of trace gas non-homogeneous transport in firn: evaluation and behaviour at eleven polar sites, *Atmos. Chem. Phys.*, 12, 11 465–11 483, <https://doi.org/10.5194/acp-12-11465-2012>, 2012.