

A global mean sea-surface temperature dataset for the Last Interglacial (129-116 kyr) and contribution of thermal expansion to sea-level change

Chris S.M. Turney^{1,2*}, Richard T. Jones^{3†}, Nicholas P. McKay⁴, Erik van Sebille^{5,6}, Zoë A. Thomas^{1,2}, Claus-Dieter Hillenbrand⁷, Christopher J. Fogwill^{1,8}

¹Palaeontology, Geobiology and Earth Archives Research Centre, School of Biological, Earth and Environmental Sciences, University of New South Wales, Australia

²ARC Centre of Excellence in Australian Biodiversity and Heritage, School of Biological, Earth and Environmental Sciences, University of New South Wales, Australia

³Department of Geography, Exeter University, Devon, EX4 4RJ, UK

⁴School of Earth and Sustainability, Northern Arizona University, Flagstaff, Arizona 86011, USA

⁵Grantham Institute & Department of Physics, Imperial College London, London, UK

⁶Institute for Marine and Atmospheric Research Utrecht, Utrecht University, Utrecht, Netherlands

⁷British Antarctic Survey, High Cross, Madingley Road, Cambridge CB3 0ET, UK

⁸School of Geography, Geology and the Environment, Keele University, ST5 5BG, UK

†Deceased.

Correspondence to: Chris Turney (c.turney@unsw.edu.au)

Abstract. A valuable analogue for assessing Earth’s sensitivity to warming is the Last Interglacial (LIG; 129-116 kyr), when global temperatures (0 to +2°C) and mean sea level (+6 to 11 m) were higher than today. The direct contribution of warmer conditions to global sea level (thermsteric) are uncertain. We report here a global network of LIG sea surface temperatures (SST) obtained from various published temperature proxies (e.g. faunal/floral assemblages, Mg/Ca ratios of calcareous plankton, alkenone U^K₃₇). We summarise the current limitations of SST reconstructions for the LIG and the spatial temperature features of a naturally warmer world. Because of local δ¹⁸O seawater changes, uncertainty in the age models of marine cores, and differences in sampling resolution and/or sedimentation rates, the reconstructions are restricted to mean conditions. To avoid bias towards individual LIG SSTs based on only a single (and potentially erroneous) measurement or a single interpolated data point, here we report average values across the entire LIG. Each site reconstruction is given as an anomaly relative to 1981-2010, corrected for ocean drift and where available, seasonal estimates provided (189 annual, 99 December-February, and 92 June-August records). To investigate the sensitivity of the reconstruction to high temperatures, we also report maximum values during the first five millennia of the LIG (129-124 kyr). We find mean global annual SST anomalies of 0.2 ± 0.1°C averaged across the LIG and an early maximum peak of 0.9 ± 0.1°C respectively. The global dataset provides a remarkably coherent pattern of higher SST increases at polar latitudes than in the tropics (demonstrating the polar amplification of surface temperatures during the LIG), with comparable estimates between different proxies. Polewards of 45° latitude, we observe annual SST anomalies averaged across the full LIG of >0.8 ± 0.3°C in both hemispheres with an early maximum peak of >2.1 ± 0.3°C. Using the reconstructed SSTs suggests a mean LIG global thermsteric sea level rise of 0.08 ± 0.1 m and a peak contribution of 0.39 ± 0.1 m respectively (assuming warming penetrated to 2000 m depth). The data provide an important natural baseline for a warmer world, constraining the contributions of Greenland and Antarctic ice sheets to global sea level during a geographically widespread expression of high sea level, and can be used to test the next inter-comparison of models for projecting future climate change. The dataset described in this paper, including summary temperature and thermsteric sea-level reconstructions, are available at <https://doi.pangaea.de/10.1594/PANGAEA.904381> (Turney et al., 2019).

1 Introduction

The timing and impacts of past, and future, abrupt and extreme climate change remains highly uncertain. A key challenge is that historical records of change are too short (since CE 1850) and their amplitude too small relative to projections for the next century (IPCC, 2013;PAGES2k Consortium et al., 2017), raising concerns over our ability to successfully plan for future change. While a wealth of geological, chemical, and biological records (often referred to as ‘natural archives’ or ‘palaeo’) indicate that large-scale and often multi-millennial duration shifts in the Earth system took place in the past (Thomas, 2016;Steffen et al., 2018;Lenton et al., 2008;Thomas et al., 2020), there are limited global datasets of such events. A comprehensive database of environmental

52 conditions during periods of warmer-than-present-day is essential for constraining uncertainties surrounding
53 projected future change, including sea level rise, extreme weather events and the climate-carbon cycle. In this
54 regard, the Last Interglacial (LIG), an interval spanning approximately 129,000 to 116,000 years ago, is of great
55 value (Dutton et al., 2015). Described as a ‘super-interglacial’ (Turney and Jones, 2010;Overpeck et al., 2005),
56 the LIG was one of the warmest periods of the last 800 kyr, experiencing relatively higher polar temperatures
57 compared to the global mean (‘polar amplification’) (Past Interglacials Working Group of PAGES,
58 2016;Hoffman et al., 2017;Turney and Jones, 2010;Capron et al., 2017), with the most geographically
59 widespread expression of high global mean sea level in the recent geological record (GMSL, +6.6 to +11.4 m)
60 (Dutton et al., 2015;Grant et al., 2014;Kopp et al., 2009;Rohling et al., 2017), abrupt shifts in regional
61 hydroclimate (Wang et al., 2008;Thomas et al., 2015), and elevated atmospheric CO₂ concentrations (relative to
62 the pre-industrial period) of ~290 ppm (Köhler et al., 2017;Schneider et al., 2013;Barnola et al., 1987;Petit et
63 al., 1999), suggesting non-linear responses in the Earth system to forcing (Steffen et al., 2018;Thomas,
64 2016;Dakos et al., 2008;Thomas et al., 2020). Importantly, there remain considerable debate over the
65 contribution of sources to the highstand in global sea level (Dutton et al., 2015;Rohling et al., 2019). Previous
66 work has suggested ocean thermal expansion contributed some 0.4 m (McKay et al., 2011), while Greenland Ice
67 Sheet melt is estimated at some 2 m (NEEM Community Members, 2013) and melting mountain glaciers ~0.6
68 m (Dutton et al., 2015), implying Antarctic mass loss >3.6 m (Fogwill et al., 2014;Turney et al., 2020;DeConto
69 and Pollard, 2016;Dutton et al., 2015;Rohling et al., 2019). Constraining the different contributions to GMSL
70 during the LIG requires a comprehensive ocean temperature database to precisely quantify the role of ocean
71 thermal expansion, compare to climate model-generated temperature estimates, and use these temperature
72 estimates to drive ice sheet models (Fogwill et al., 2014;Mercer, 1978;DeConto and Pollard, 2016;Sutter et al.,
73 2016;Hoffman et al., 2017;Clark et al., 2020).

74
75 Quantified temperature reconstruction data for the LIG are often drawn from disparate publications and
76 repositories (usually reported alongside other Late Pleistocene data). To obtain reliable temperature
77 reconstructions, it has until recently proved necessary to determine a global estimate of the magnitude of
78 warming using only a selected number of “high-quality” records; the resulting temperature reconstructions of
79 LIG temperatures ranged from 0.1 to >2°C warmer than present (CLIMAP, 1984;White, 1993;Hansen,
80 2005;Rohling et al., 2008;Turney and Jones, 2010). With the ever-increasing number of quantified temperature
81 reconstructions of the LIG reported in individual publications, it is crucial that these datasets are brought
82 together to derive a comprehensive reconstruction of global change during the LIG. A further consideration is
83 that in contrast to terrestrial sequences, marine records typically provide a continuous record of LIG conditions
84 (Turney and Jones, 2010;Turney and Jones, 2011), providing an opportunity to determine the sensitivity of
85 GMSL to sea-surface temperature (SST) conditions during the interglacial (including early maximum
86 temperatures). Given a possible warming of 2°C (Turney and Jones, 2010;Fischer et al., 2018), the LIG
87 potentially provides insights into the drivers of sea level rise and the long-term impacts under a global
88 temperature target set out in the 2016 Paris Climate Agreement (Schellnhuber et al., 2016).

89
90 Here we present version 1.0 of the Last Interglacial SST database (Turney et al., 2019). This database builds on
91 the previously published 2010 data compilation of (Turney and Jones, 2010), and includes substantially more
92 records. Importantly, the micro-organisms used to determine SSTs move along with the currents and encounter
93 a range of temperatures during their life cycle (van Sebille et al., 2015;Doblin and van Sebille, 2016;von
94 Gyldenfeldt et al., 2000). As a result, previous workers have suggested ocean drift of micro-organisms can have
95 a major influence on reconstructed environmental change (van Sebille et al., 2015;Monroy et al., 2017;Kienast
96 et al., 2016;Hellweger et al., 2016;Rembauville et al., 2016;Viebahn et al., 2016;Nooteboom et al.) and
97 potentially explains the divergence between laboratory culture and core-top calibrations (Anand et al.,
98 2003;Müller et al., 1998;Prahl et al., 2003;Sikes et al., 2005;Segev et al., 2016;Elderfield and Ganssen, 2000),
99 and palaeoclimate estimates and model outputs (Otto-Bliesner et al., 2013;Bakker and Renssen, 2014;NEEM
100 Community Members, 2013;Lunt et al., 2013), including the recently recognised historic (Anthropocene)
101 change in modern plankton communities which has major implications for calibration studies (Jonkers et al.,
102 2019). The influence of ocean currents has not been explored (or corrected for) in previous studies of the LIG
103 (Hoffman et al., 2017;Capron et al., 2014;Turney and Jones, 2010) and is important for obtaining correct
104 absolute SSTs. This descriptor describes the contents of the database, the criteria for inclusion, and quantifies
105 the relation of each record with instrumental temperature, including the estimated impact of ocean current drift
106 on individual sites and global averages. The current database includes a large number of metadata fields to
107 facilitate the reuse of the data and identification of key records for future investigations into the LIG. Specific
108 criteria were developed to gather all published proxy records that meet key objective and reproducible criteria.
109 The database will be updated yearly as newly reported records are published.

110 2 Methods

111 2.1 Global Compilation

112 We have compiled a global network of published quantified SSTs using faunal and floral assemblages, Mg/Ca
113 and Sr/Ca ratios of calcareous organisms, and $U^{K_{37}}$ estimates across the period of record interpreted as
114 representing the LIG. In many instances, we used the period represented by low ^{18}O values in benthic
115 foraminifera shells (the lightest isotopic values during 90-150 kyr representing minimum global ice volume),
116 although in some sequences, $\delta^{18}O$ values were reported and we relied on other complimentary proxies; for
117 instance, the $CaCO_3$ content of sediments as a measure of glacial-interglacial variability (Turney and Jones,
118 2010;Cortese et al., 2013) (Figures 1 and 2). Whilst the age control points defining the plateaus in $\delta^{18}O$ and
119 other proxies are not absolutely dated with chronological uncertainties of one to two millennia (Martinson et al.,
120 1987;Lisiecki and Raymo, 2005), it is important to note that we are not aiming to resolve centennial and
121 millennial-scale variability through the interglacial. We acknowledge that some individual SST estimates may
122 not fall within the LIG or have been excluded (due to these chronological uncertainties) but we consider the
123 averaging of values across the full interglacial provides a robust value for each record and ultimately the
124 regional and global reconstructions.

125
126 We have therefore not attempted to generate a time series of sea surface temperatures through the LIG. Previous
127 studies have highlighted that individual site $\delta^{18}O$ changes in benthic foraminifera (for instance, during
128 deglaciation) may be offset by several millennia as a result of local deep-water temperature and $\delta^{18}O$ seawater
129 variations (Govin et al., 2015;Waelbroeck et al., 2008) (Figure 2). In an attempt to bypass some of these issues,
130 other studies have attempted alignment of marine records to speleothem-dated, ice core reconstructions (Hoffman
131 et al., 2017) but modelled age uncertainties can be on the order of millennia (e.g. Hoffman et al. Fig. S7) while
132 the assumed synchronicity of extra-regional changes has challenges; for instance, more than half of reported
133 Pacific marine cores (those from the Northern Hemisphere) in a recent study were correlated to the Antarctic
134 EPICA Dome C δD (Hoffman et al., 2017), with warming in the south known to lead the north by one to two
135 millennia (Hayes et al., 2014;NEEM Community Members, 2013;Kim, 1998;Rohling et al., 2019). The
136 development of accurate and precise age estimates for LIG records is urgently needed to resolve the timing of
137 global climate change but will require a considerable future international effort (Govin et al., 2015). Given the
138 relatively large chronological uncertainties associated with comparing global SST time series (Hoffman et al.,
139 2017;Govin et al., 2015;Capron et al., 2017) we have therefore not attempted to generate a time-series of changes
140 within the LIG but instead determine average temperatures as a robust estimate of mean climatic conditions.
141 Whilst not offering precisely-dated geochronological frameworks, the global ice minima as represented by the
142 $\delta^{18}O$ plateau and/or associated proxy measures of interglacial conditions are sufficiently well-defined in all marine
143 records to accommodate local deep-water temperature and $\delta^{18}O$ variations, sampling resolution and/or
144 sedimentation rates to identify the LIG, thereby maximising the number of records that have reported quantified
145 SSTs across the interglacial (Cortese et al., 2013;Govin et al., 2015); a minimum of three SST values across the
146 LIG in each record were required for inclusion in our dataset. This is not to downplay the significance of
147 millennial-scale climate variability across the LIG (Galaasen et al., 2014;Rohling et al., 2002;Tzedakis et al.,
148 2018;Jones et al., 2017) but our approach does provide some benefits. Whilst our approach sacrifices temporal
149 control, it does minimise the uncertainty on zonal and global temperature averages.

150
151 To quantify the temperature difference between the LIG and present day, we do not compare the LIG estimates
152 to the relatively poor observational coverage of earlier periods, including the nineteenth century (pre-industrial)
153 (Hoffman et al., 2017) or the long-term annual means calculated from 1900-1997 (Capron et al., 2014), both of
154 which have considerable uncertainties given the limited network of ‘observations’ prior to the satellite era
155 (Brohan et al., 2006;Huang et al., 2020). Here instead we report SSTs expressed as anomalies relative to global
156 ‘modern’ instrumental and satellite observations across the period 1981-2010 obtained from HadISST (Rayner
157 et al., 2003). Each LIG temperature record is linked to at least one literature source, the citation of which
158 includes author(s), year of publication and typical archiving information (e.g. journal, volume, issue, pages,
159 publisher and place of publication). Where multiple temperature estimates have been published over time from
160 the same site, we chose the most recent publication for inclusion in the database (so long as the data were not
161 flagged as erroneous) (Figure 3). Note that alkenone proxies are interpreted as providing annual SST estimates.

162
163 Here we use the mean temperature estimates to constrain the role of thermal expansion in global sea level rise
164 across the LIG and provide boundary conditions for future modelling studies investigating the impact of warming
165 on polar ice sheets. To determine the greatest possible contribution of warming to ocean thermal expansion and
166 ice sheet melt, we used the published age models to identify the maximum annual SST within the first 5 kyr of
167 the LIG (i.e. 129-124 kyr). For the purposes of this sensitivity analysis, the maximum temperatures were assumed
168 to be synchronous globally, a scenario we recognise as unlikely but does provide an upper limit for warming in

169 the ‘early’ LIG. To provide an upper estimate on the magnitude of warming in polar waters over the deglaciation,
170 we also report here the difference between late Marine Isotope Stage 6 mean SSTs (~140-135 kyr) and the
171 maximum early LIG SSTs for ocean cores in the mid to high-latitudes. To calculate the anomaly relative to present
172 day, we utilise SSTs from the nearest 0.5° latitude x 0.5° longitude averaged across the period 1981-2010 (Rayner
173 et al., 2003). For the uncertainties calculated for the regional and global SST anomalies, we incorporate the
174 uncertainties from the proxies (reported in the database), and the uncertainties associated with estimating regional
175 and global temperatures from limited spatial coverage. To achieve this we propagated the SST uncertainties for
176 each measurement through each of the averaging steps (i.e. temporal to grid cell to zonal to area-weighted global)
177 in our ocean-area-weighted average (McKay et al., 2011). We used quoted uncertainty estimates for each study
178 where reported; if not available, we applied proxy-specific uncertainty estimates. Although the impact of the
179 spatial coverage was not explored in this study, it has been previously estimated using the same approach (McKay
180 et al., 2011). In that study, the uncertainty associated with the limited spatial range of the oceanographic proxies
181 was estimated by calculating 1000 random one-year global SST anomalies over the twentieth century, and
182 compared to averages derived using only the palaeoceanographic network. No systematic biases were identified
183 with a 1 σ uncertainty estimated to be <0.1°C. In this study, we have expanded the spatial network, and consider
184 $\pm 0.1^\circ\text{C}$ to be a reasonable, high-end estimate.

185
186 The database comprises six worksheets of data comprising maximum annual temperatures during the early LIG
187 (defined here as the maximum temperature reported within the first five millennium of the LIG; 129-125 kyr),
188 mean annual temperature, the Marine Isotope Stage 6/5 SST difference, December to February temperature
189 (DJF; Northern Hemisphere winter and Southern Hemisphere summer), June to August temperature (JJA;
190 Northern Hemisphere summer and Southern Hemisphere winter), and summary statistics (see Supplementary
191 Information):

- 192 - The early maximum and mean annual SST dataset comprises 189 marine sediment and coral records
193 from latitudes spanning from 55.55°S (radiolaria assemblage transfer function reconstruction obtained
194 from site V18-68) (CLIMAP, 1984) to 72.18°N (planktonic foraminifera assemblage modern analogue
195 technique from site V27-60) (Vogelsang et al., 2001)
- 196 - The mean December-February SST dataset comprises 99 marine sediment records from latitudes
197 spanning from 61.24°S (diatoms transfer function reconstruction obtained from site PS58/271-1) (Esper
198 and Gersonde, 2014) to 72.18°N (planktonic foraminifera assemblage modern analogue technique from
199 site V27-60) (Vogelsang et al., 2001).
- 200 - The mean June-August SST dataset comprises 92 marine sediment records from latitudes spanning
201 from 54.55°S (radiolaria assemblage transfer function reconstruction obtained from site V18-
202 68)(CLIMAP, 1984) to 72.18°N (planktonic foraminifera assemblage modern analogue technique from
203 site V27-60) (Vogelsang et al., 2001).

204
205 In total, the Last Interglacial SST database comprises a total of 203 unique sites described in 100 publications.
206

207 **2.2 Ocean Drift**

208 Crucially, modern calibration relationships are an average developed using a selected number of locations that
209 will not necessarily capture the range of “signal drift”. This drift is caused by the fact that planktic SST
210 recorders can be transported over considerable distances in the water column before being deposited, which
211 particularly applies to all those sites that lie under strong boundary currents or near major ocean fronts (van
212 Sebille et al., 2015). Unfortunately, Ocean General Circulation Models (OGCMs) typically have insufficient
213 spatial resolution to capture mesoscale features that are critical for modelling the lateral drift of particles
214 (Nooteboom et al., 2020). To investigate the impact of drift on SST reconstructions, we therefore used
215 contemporary ocean circulation as a first-order approximation for the LIG. Whilst we acknowledge that there
216 was likely a weakening of the Atlantic Meridional Overturning Circulation (AMOC) during the early LIG
217 (Shackleton et al., 2020;Turney et al., 2020;Thomas et al., 2020;Jones et al., 2017), subsequent recovery after
218 127 kyr appears to have established a global circulation comparable to present day as suggested by recent ocean
219 $\delta^{13}\text{C}$ modelling results across the mid-interglacial (Bengtson et al., 2020). We performed an experiment with
220 virtual particles in an eddy-resolving ocean model (the Japanese Ocean model For the Earth Simulator or OFES)
221 (Masumoto et al., 2004), which has a 1/10° horizontal resolution and near-global coverage between 75°S and
222 75°N (van Sebille et al., 2012). Utilising the 3D velocity field of the model, we used the Parcels code
223 (oceanparcels.org) (Lange and van Sebille, 2017) to compute the trajectories of more than 170,000 virtual
224 planktic particles that end up at each of the sites by tracking them backwards in time, first simulating the sinking
225 to these sites at 200 m/day and subsequently the advection at 30 m depth for a lifespan of 30 days; coral SSTs
226 were not corrected for drift. Given the lifespan of most organisms that have been used to generate a temperature
227 signal (Jonkers et al., 2015;Bijma et al., 1990), we consider a 30-day drift provides a reasonable estimate of the

228 drift distance. Previous work has demonstrated comparable uncertainties between different models (van Sebille
229 et al., 2015), providing confidence in the use of the OFES for the purposes of this study.

230
231 During the 30-day lifespans, we recorded the temperatures along the trajectories and compared those to the local
232 temperature at 30 m water depth at the site where the particles would end up on the ocean floor. This resulted in
233 daily temperature anomalies along the trajectories, which were averaged through the lifespan and over the 840
234 virtual particles that ended up at each site, and then subtracted from the reported LIG estimates (Figure 1 and
235 Database). With the recent recognition that core-top calibrations may be incorrect given historic changes in
236 marine communities that have accompanied anthropogenic warming (Jonkers et al., 2019), it should be noted
237 that SST proxy calibrations based on regional core-top calibrations may give an incorrect absolute value, an
238 aspect that will form the focus of future work.

239 240 **2.3 Hemispheric and Global Calculations**

241 Global mean SST anomalies were calculated by averaging anomalies in a 10° latitude \times 10° longitude grid, then
242 averaged globally after weighting for the area of ocean in each grid cell (Figure 5). The uncertainty calculated
243 for global SST anomalies incorporates uncertainties in the SST proxies as reported in the original studies, which
244 typically ranges from 1 to 2°C , and is then propagated through subsequent steps in the analysis. Additional
245 uncertainty associated with estimating global anomalies from limited spatial coverage, and the potential impacts
246 of age uncertainty or averaging non-synchronous data are not considered here. Consequently, the derived
247 estimates do not capture all of uncertainty in global and zonal SST anomalies, however, the zonal consistency of
248 the results suggest that the signal is large enough to overcome these unquantified sources of uncertainty.
249 Furthermore, whilst some regions may exhibit substantial differences arising from drift (Figure 4), taken
250 globally the mean annual temperature estimates are comparable (Figure 5). The new LIG SST dataset allows us
251 to report the estimated thermosteric contribution for LIG sea levels using the method reported by (McKay et al.,
252 2011). We use the above temperature changes to calculate the thermosteric contribution to LIG sea levels by
253 using the Thermodynamic Equation of Seawater 2010 (TEOS-10). To provide an estimate of thermosteric sea
254 level rise, we explored a range of scenarios where warming penetrated different ocean depths: 700 m, 2000 m
255 (approximately the upper half of the ocean) and 3500 m (the whole ocean). We determined the change in the
256 specific volume of the warmed water column of each a 10° latitude \times 10° grid cell while holding the salinity
257 constant and neglecting changes in ocean area. Here absolute temperature is considered, as specific volume is
258 more sensitive to temperature changes at warmer temperatures.

259 **3 Results and Discussions**

260 **3.1 Quality Control**

261 The Last Interglacial SST database is derived from published articles that have already been peer-reviewed. To
262 generate the database, we undertook a comprehensive check to remove duplicate records, erroneous location
263 information and other errors. In addition to ensuring consistency of data processing and any recalculations (for
264 instance, sea-surface temperature anomalies relative to the period CE 1981-2010), we also checked uncertain
265 metadata reported for individual sites, and directly communicated with selected article authors and/or other
266 experts as part of the record-validation process.

267 268 269 **3.2 Ocean Circulation**

270 A challenge for the Last Interglacial is determining what influence (if any) ocean circulation had on the
271 temperatures experienced (and reconstructed) by organisms that are used to generate SST reconstructions.
272 Addressing this issue is an important objective of the current study but we found the magnitude of temperature
273 offset (bias) is limited to only a few key locations (Fig. 1), with similar final reconstructions for individual sites,
274 latitudinally-averaged and globally average temperatures (Figures 4 and 5, and Table 1). This provides an
275 important check of our temperature recalculations. As a sensitivity test, we therefore explored virtual planktic
276 particles that ‘live’ for 30 days to investigate whether a prolonged period of drift made a discernible difference
277 (data not reported here). Only a few species have been suggested as living for a longer period of time. For
278 instance, in laboratory experiments the planktic foraminifer *Neogloboquadrina pachyderma* sinistral has been
279 shown to survive up to 230 days (Spindler, 1996) but this species may be an exception due to its ability to
280 survive in sea ice (Dieckmann et al., 1991).

281
282 Using 30-days’ drift to simulate the travelling time/lifespans of virtual planktic particles in the upper part of the
283 water column, we quantified the inherited temperature signal of flora/fauna at each site in the database. The
284 virtual microorganisms with a 30-day ‘lifespan’ travelled from a few tens to a few hundreds of kilometres. The

285 temperature offsets are almost all positive in the tropical East Pacific, the North Atlantic and South China Sea,
286 meaning that the planktic particles originated from warmer climates and hence record a higher temperature
287 estimate than local conditions would suggest; with the opposite effect observed in the western tropical Pacific
288 and Southern Ocean (Figure 1). The offset can be substantial – with values ranging from -6.9°C for site MD98-
289 2162 at 4.7°S in the tropical West Pacific (Visser et al., 2003) and up to 3.5°C in site RC13-110 on the Equator
290 (Pisias and Mix, 1997) – with the largest changes associated with boundary currents and major ocean fronts.
291 Intriguingly, these values are comparable to the difference previously reported for Mg/Ca foraminifera core-top
292 calibration with those obtained from laboratory-cultured Mg/Ca calibrations (Elderfield and Ganssen,
293 2000;Hönisch et al., 2013). Both the uncorrected and 30-day drift temperatures are provided in the database.
294 These temperature reconstructions led to statistically indistinguishable global temperature (and thermosteric sea
295 level change; Figure 5). Users of the database are therefore able to use either the authors’ original sea-surface
296 temperature determinations or our drift-corrected estimates, as required.
297

298 **3.3 Proxy and Seasonal Effects**

299 To evaluate potential biases in our analysis, we further subsampled our database by proxy type (Figure 4). The
300 large network of sites and proxies do not appear to demonstrate any significant offset in annual reconstructions
301 (at least within the uncertainty of the reconstructions), although there is a tendency for alkenone temperatures to
302 be at the upper end of the range, implying there may be a seasonal bias, as reported previously (Hoffman et al.,
303 2017). Importantly, we also compiled seasonal quantified temperature estimates that have been reported as the
304 seasonal warmest or coolest months in the year (taken here to represent June-August and December-February
305 depending on the hemisphere being considered). Our result suggests that any bias, if real, is smaller than the
306 uncertainties at the global or zonal level reported here. Intriguingly, the warmest month estimates for the high
307 latitudes in both hemispheres have more muted warming than the mean annual estimates while the low to mid
308 latitudes exhibit considerably cooler estimates (Table 1). In contrast to the alkenone estimates for the annual
309 estimates, the more muted response of foraminifera, radiolaria and diatoms for the seasonal reconstructions
310 implies they are influenced by a larger part of the seasonal cycle. We therefore consider that seasonal
311 reconstructions should be treated as conservative estimates of temperature for the LIG.
312

313 **3.4 Average and Early Temperatures during the Last Interglacial**

314 We find global average annual temperatures across the full duration of the LIG were only marginally warmer
315 than present day. We derive a global mean annual temperature anomaly of $0.2 \pm 0.1^{\circ}\text{C}$, the same value obtained
316 after correcting for drift (Table 1). These values, however, mask considerable zonal differences, with
317 significantly cooler mean annual uncorrected temperatures (i.e. not corrected for drift) within 23.5° of the
318 equator ($-0.3 \pm 0.2^{\circ}\text{C}$) and amplified warming polewards (Figure 5). Ideally, we would have a dense network of
319 records in the mid- to high-latitudes for investigating the impact of warming surrounding polar ice sheets but
320 unfortunately the number of sites and their spatial distribution do appear to have an impact on the reconstructed
321 values. Comparison of the SST anomalies poleward of 45° and 50° latitude (Table 1) shows substantial
322 differences, most notably in the Southern Hemisphere where a large increase in zonally averaged SST occurs
323 alongside a decrease in the number of records polewards of 50°S (Table 1). For instance, the drift-corrected
324 SSTs for the LIG are $0.8 \pm 0.3^{\circ}\text{C}$ ($n=13$) and $2.7 \pm 1.1^{\circ}\text{C}$ ($n=3$) polewards of 45°S and 50°S respectively. It
325 should also be noted that whilst the Northern Hemisphere polar estimates are similar for both latitudinal ranges,
326 the majority of sites are in the North Atlantic, with limited representation in the Pacific Ocean (Figure 1). We
327 therefore recommend that when considering mid- to high-latitude zonal SST averages, the values derived from
328 records polewards of 45° are more likely robust but acknowledge these may be conservative estimates (with
329 considerably larger warming further to the south). We therefore estimate uncorrected ‘polar’ warming in the
330 Northern Hemisphere to be $2.0 \pm 0.4^{\circ}\text{C}$, and in the Southern Hemisphere, $0.2 \pm 0.3^{\circ}\text{C}$ (Table 1). Correcting for
331 drift decreased the northern estimate to $1.5 \pm 0.4^{\circ}\text{C}$ and increased in the south to a mean annual SST to 0.8 ± 0.3
332 $^{\circ}\text{C}$.
333

334 The maximum temperatures of the early LIG were up to $0.9 \pm 0.1^{\circ}\text{C}$ warmer than 1981-2010, regardless of
335 whether the values were corrected for drift (Table 1 and Figure 6). Similar to the mean SSTs of the LIG, there
336 appears to have been considerable zonal differences in the uncorrected values: $0.1 \pm 0.2^{\circ}\text{C}$ within 23.5° of the
337 equator, $3.2 \pm 0.4^{\circ}\text{C}$ polewards of 45°N , and $1.5 \pm 1.1^{\circ}\text{C}$ polewards of 45°S . After correcting for drift, the
338 estimated SST in the north changed to $2.8 \pm 0.4^{\circ}\text{C}$ and in the south, to $2.1 \pm 1.1^{\circ}\text{C}$. The latter estimate from the
339 Southern Hemisphere is $\sim 2^{\circ}\text{C}$ (relative to 1981-2010), potentially providing an important constraint for future
340 Antarctic ice-sheet model simulations for the LIG (Turney et al., 2020;Golledge et al., 2015). These data
341 support previous work which have reported substantial polar temperature amplification during the LIG,
342 particularly in the Northern Hemisphere (described in the literature as ‘Arctic amplification’) (Overpeck et al.,
343 2006;Mercer, 1978;Mercer and Emiliani, 1970;Thomas et al., 2020;Miller et al., 2010). The global temperature
344 pattern closely follows insolation changes across this period, during which the Earth’s greater eccentricity led to

345 reduced radiation over the equator and more intense high latitude spring-summer insolation (Figure 2)
346 (Overpeck et al., 2006;Hoffman et al., 2017). Comparison to Marine Isotope Stage 6 SSTs appears to show the
347 greatest warming in the northeast Atlantic and south Atlantic (Figure 7), suggesting Greenland and the West
348 Antarctic ice sheets would have been particularly vulnerable to warming in the early interglacial (Clark et al.,
349 2020;Turney et al., 2020;Dutton et al., 2015;Mercer, 1978) though we cannot resolve the relative timing of mass
350 loss in this analysis (Rohling et al., 2019;Hayes et al., 2014). Recent work suggests the earliest warming took
351 place in the Atlantic (and Indian) Ocean sectors of the Southern Ocean (Chadwick et al., 2020), consistent with
352 our findings. However, our observed polar warming is larger than some climate model simulations, implying the
353 latter are failing to capture one or more key feedbacks (e.g. carbon, sea-ice and ice-sheet feedbacks) in the
354 climate system (Bakker et al., 2013;Otto-Bliesner et al., 2013;Thomas et al., 2020;Clark et al., 2020;Fogwill et
355 al., 2015).

356

357 **3.5 Thermal Expansion Contribution to Last Interglacial Sea Level**

358 The LIG is characterised by higher GMSL than present day (+6.6 to +11.4 m) (Grant et al., 2014;Dutton et al.,
359 2015;Turney and Jones, 2010;Rohling et al., 2017;Rohling et al., 2019). Here we quantified the contribution of
360 the relatively high temperatures on global sea levels through ocean thermal expansion for warming down to
361 2000 m ocean depth (Table 2). We find that through the LIG, the average SSTs contribution to thermosteric sea
362 level was negligible, approximately 0.05 ± 0.10 m uncorrected for ocean drift and 0.08 ± 0.10 m corrected for
363 drift, consistent with a recent reconstruction of near-modern global ocean heat content and negligible
364 thermosteric sea level rise (Shackleton et al., 2020). But for the early LIG (129-124 kyr), using our maximum
365 SST estimate, we obtained high-end estimate of thermal expansion to GMSL of 0.36 ± 0.10 m (uncorrected) and
366 0.39 ± 0.10 m (drift corrected). These quantified estimates are comparable to a previously reported value of 0.4
367 ± 0.3 m (McKay et al., 2011) which used the same methodology as here but a smaller network of SST records.
368 However, we should recognise that the depth of ocean warming is uncertain, and could have extended deeper
369 than 2000 m. If we assume warming penetrated the full ocean depth (down to 3500 m), we obtained a maximum
370 early LIG thermosteric sea level rise of 0.67 ± 0.10 m (uncorrected) and 0.72 ± 0.10 m (drift corrected) (Table
371 2). The recently reported early LIG (~129 ka) peak in global ocean heat content reconstructed from isotopic
372 ratios of atmospheric trace gases has determined a maximum thermal expansion of 0.7 ± 0.3 m (Shackleton et al.,
373 2020). To achieve ~0.7 m of thermosteric sea level rise during the early interglacial peak in temperatures, we
374 have to use both our maximum estimate of temperature rise, and our maximum estimate of the depth of
375 warming. A recent modelling-proxy estimate proposed a range of 0.08 to 0.51 m for peak LIG warmth centred
376 on 125 kyr (Hoffman et al., 2017), which is more consistent with our results. Even though 125 ka is later than
377 the peak in global ocean heat content, this is effectively the same event but represents the age uncertainties in
378 the marine records. Although some uncertainty remains in the amplitude of thermal expansion between these
379 studies, it is clear that the sustained high global sea levels across the LIG and the limited role of warming on
380 thermal expansion implies a greater contribution from ice sheets, mountain glaciers, permafrost and
381 hydrological change. With the greatest warming relative to Marine Isotope Stage 6 in Atlantic basin (Figure 7),
382 our results are consistent with previous studies suggesting substantial mass loss from Greenland and the West
383 Antarctic Ice Sheet early in the Last Interglacial (Clark et al., 2020;Turney et al., 2020;Dutton et al.,
384 2015;Mercer, 1978;Hayes et al., 2014;Rohling et al., 2019).

385 **4 Data Availability**

386 The Last Interglacial SST database is provided as an Excel workbook in Supplementary Information and on the
387 PANGAEA Data Publisher at <https://doi.pangaea.de/10.1594/PANGAEA.904381> (Turney et al., 2019); the data
388 is also available on the NCEI-Paleo/World Data Service for Paleoclimatology at
389 <https://www.ncdc.noaa.gov/paleo/study/26851>. This release comprises a single Excel file, tab delimited. We
390 welcome contributions from authors of additional or clarifying information. These will be incorporated into any
391 subsequent iteration of the database. When using data in this compilation, the original data collector(s) as well
392 as the data compiler(s) will be credited. Given the typically large uncertainties in the absolute dating of each
393 individual record, no attempt has been made to develop individual time series, and only mean values across the
394 Last Interglacial have been compiled. For simplicity we record the 1σ (68%) confidence interval in the site
395 temperature reconstructions. The inclusion of key metadata allows users to interrogate individual records for
396 their own appropriate screening criteria.


398 During the Last Interglacial (LIG; 129-116 kyr), global temperatures were up to 2°C warmer than present day
399 with marked polar amplification and global sea levels between 6.6 and 11.4 m higher than present day, offering a
400 powerful opportunity to obtain key insights into the drivers of future change (a so-called ‘process analogue’). The
401 contributions of different sources to the LIG sea level highstand remain highly uncertain, however. As a result of
402 relatively warmer surface temperatures, ocean thermal expansion has previously been estimated to have
403 contributed 0.4 ± 0.3 m. To more precisely constrain this contribution to global mean sea level we report a new
404 comprehensive database of quantified SSTs estimates derived from faunal and floral assemblages, Mg/Ca and
405 Sr/Ca ratios of calcareous organisms, and U^{K}_{37} estimates from records spanning 55.55°S to 72.18°N. Here we
406 have calculated maximum annual SSTs during the early interglacial (129-124 kyr) and mean annual SSTs through
407 the LIG (129-116 kyr; 189 sites) alongside mean December-February (99 records) and June-August (92 records)
408 values. Temperatures are reported as anomalies relative to the period CE 1981-2010. To estimate the temperature
409 footprint arising from ocean circulation we also report SST anomalies corrected for 30-day drift, to simulate the
410 travelling time/lifespans of virtual planktic particles in the upper part of the water column. Our reconstruction
411 suggests an early LIG maximum global mean annual SST of $0.9 \pm 0.1^{\circ}\text{C}$ and an average warming across the LIG
412 of $0.2 \pm 0.1^{\circ}\text{C}$. However, these values are strongly driven by polar warming of several degrees, with little to no
413 warming in the tropics. We find the influence of warming on ocean thermal expansion to have had a limited
414 influence on global mean sea levels across the full LIG, but with a likely range of between 0.39 ± 0.1 m and
415 0.72 ± 0.10 m early in the interglacial. Our findings therefore imply a relatively greater contribution of ice sheets,
416 mountain glaciers, permafrost and hydrological change to global sea level during the LIG, likely driven by polar
417 amplification of temperatures. We hope this database may provide a springboard for future studies that can bring
418 to bear new geochronological methods (e.g. tephra) to constrain the age models of individual sequences to sub-
419 millennial uncertainty, something currently not possible for most reported marine sequences. An improved
420 network of high-resolution, well-dated and quantified LIG climate reconstructions (particularly in data-sparse
421 locations) will enable precise integration of ice sheet, marine and terrestrial records to better understand Earth
422 system responses to high-latitude warming. The Southern Ocean and North Pacific are regions where major
423 knowledge gaps currently exist.

424
425 **Supplement.** The supplementary figures and version 1.0 of the database (Excel file) can be accessed via the
426 *Earth System Science Data* discussion page of this manuscript.

427
428 **Author contributions.** RTJ and CSMT conceived the research; CT, NPM, EvS, and ZT designed the methods
429 and performed the analysis; CT wrote the paper with substantial input from all authors.

430
431 **Competing interests.** The authors declare that they have no conflict of interest.


432
433 **Acknowledgements.** It is with great sadness that our close friend and colleague Richard T. Jones was not alive
434 to see the publication of this study. Without Richard this work would not have been possible. He is sorely
435 missed. CSMT and CJF were supported by their Australian Research Council (ARC) fellowships
436 (FL100100195 and FT120100004). We would also like to acknowledge the important role of the International
437 Ocean Discovery Program (IODP), the Australian and New Zealand International Ocean Discovery Consortium
438 (ANZIC), and the previous scientific ocean drilling programs, the results from which underpin this study and
439 without whom this analysis would not have been possible. We are grateful to the four reviewers and editor for
440 helping improve the first draft of this manuscript.



441
442
443
444
445
446


Figure 1: Last Interglacial proxy-based annual sea surface temperature dataset and modelled inherited signal.

Histogram showing the number of Last Interglacial records of annual sea surface temperature binned by 10° latitude (panel a) with virtual microfossil temperature offsets defined as the difference between along-trajectory recorded temperatures and local temperatures (panel b) and distance (panel c) travelled in the Japanese Ocean model For the Earth Simulator (OFES; run between CE 1981 and 2010) determined for 30-day ‘lifespans’ (van Sebille et al., 2015).




447
448
449
450
451
452
453
454
455

Figure 2: Relationships between $\delta^{18}\text{O}$ plateau and sea surface temperatures and environmental changes across the Last Interglacial. (a) Insolation changes calculated from ref. (Laskar et al., 2004). Sea surface temperatures (dashed purple lines) across the Last Interglacial (light orange shading) compared to the benthic foraminifera $\delta^{18}\text{O}$ (solid green lines) for selected sites in different ocean basins: (b) M23414 (North Atlantic) (Kandiano et al., 2004), (c) MD85674 (equatorial Indian Ocean) (Bard et al., 1997), (d) GeoB 3603-2 (southern Indian Ocean) (Schneider et al., 1999), and (e) MD06-2986 (southern Pacific Ocean) (Cortese et al., 2013). (f) The probabilistic reconstructed global sea level curve is reported by (Kopp et al., 2009); heavy lines mark median projections, dashed lines the 16th and 84th percentiles, and dotted lines the 2.5th and 97.5th percentiles.



456
457
458
459
460
461
462


Figure 3: Simplified scheme for the generation of the Last Interglacial sea-surface temperature database providing an overview of the data collection and processing. The numbered boxes set out the stages required to generate a global database of surface temperatures from marine records: 1. Location; 2. Last Interglacial and modern SSTs (including drift calculation); and 3. Metadata including method of temperature reconstruction and associated uncertainty. Grey boxes indicate additional processing of data from the original publications, generating new outputs (which are provided in the database).



463
 464
 465
 466
 467


Figure 4: Quality-control plot of latitudinal distribution of proxy mean annual Last Interglacial sea-surface temperature anomalies. Estimates given relative to the modern period (1981-2010) (Rayner et al., 2003) with no drift correction (upper panel) and 30-days drift (lower panel). Lower panel shows drift-corrected SSTs as open symbols with the uncorrected SSTs given as filled symbols. Uncertainties on upper panel given at 1σ .

Sea Surface Temperature



468
469
470
471
472
473
474
475


Figure 5: Global and zonal mean annual sea-surface temperature (SST) anomalies and thermosteric sea level change across the full Last Interglacial. Temperature anomalies reported as uncorrected (panels a and c respectively) and after applying 30-day (panels b and d respectively) temperature offsets arising from ocean current drift. Uncertainty for zonal average reconstructions given at 1σ . Here ocean warming is assumed to have penetrated to 2000 m depth, on average. Temperature estimates relative to the modern period (CE 1981-2010).



476
477
478
479
480
481
482
483

Figure 6: Global and zonal mean annual sea-surface temperature (SST) anomalies and thermosteric sea level change during the early Last Interglacial. Temperature anomalies reported as uncorrected (panels a and c respectively) and after applying 30-day (panels b and d respectively) temperature offsets arising from ocean current drift. Uncertainty for zonal average reconstructions given at 1σ . Here ocean warming is assumed to have penetrated to 2000 m depth, on average. Temperature estimates relative to the modern period (CE 1981-2010).

484



485

486

487

488

Figure 7: Mid- to high-latitude sea surface temperature (SST) difference between late Marine Isotope Stage 6 and maximum values of the early Last Interglacial (Stage 5). Map made using Generic Mapping Tools (GMT) (Wessel et al., 2013).

	Global SST (°C)	Tropical SST (23.5°N to 23.5°S)	SST Polewards of 45°N	SST Polewards of 50°N	SST Polewards of 45° S	SST Polewards of 50°S
Maximum Early LIG (n)	(189)	(87)	(22)	(20)	(13)	(3)
<i>Uncorrected</i>	0.9	0.1	3.2	3.8	1.5	3.7
<i>30-day drift</i>	0.9	0.1	2.8	3.2	2.1	3.7
<i>1σ</i>	0.1	0.2	0.4	0.4	0.3	1.1
Mean (n)	(189)	(87)	(22)	(20)	(13)	(3)
<i>Uncorrected</i>	0.2	-0.3	2.0	2.8	0.2	2.7
<i>30-day drift</i>	0.2	-0.3	1.5	2.3	0.8	2.7
<i>1σ</i>	0.1	0.2	0.4	0.4	0.3	1.1
DJF (n)	(99)	(35)	(16)	(15)	(14)	(9)
<i>Uncorrected</i>	-0.6	-0.7	-0.1	0.0	-0.3	0.8
<i>30-day drift</i>	-0.7	-0.9	-0.5	-0.7	0.3	1.0
<i>1σ</i>	0.2	0.3	0.4	0.5	0.3	0.3
JJA (n)	(92)	(35)	(20)	(19)	(4)	(1)
<i>Uncorrected</i>	-0.4	-1.1	1.3	1.3	-1.9	0.1
<i>30-day drift</i>	-0.5	-1.2	0.9	0.7	-1.2	-0.2
<i>1σ</i>	0.2	0.3	0.4	0.4	0.4	1.1

490

Table 1: Annual and seasonal temperature estimates for the Last Interglacial. DJF: December to February; JJA: June to August. Temperature anomalies relative to the period CE 1981-2010. Maximum early temperature is defined as the maximum annual temperature recorded during the estimated first five millennia of the Last Interglacial.

495

	Global sea level (m)		
	700 m depth	2000 m depth	3500 m depth
Maximum			
Early LIG			
(n=189)			
<i>Uncorrected</i>	0.12	0.36	0.67
<i>30-day drift</i>	0.13	0.39	0.72
<i>1σ</i>	0.10	0.10	0.10
Mean			
(n=189)			
<i>Uncorrected</i>	0.00	0.05	0.10
<i>30-day drift</i>	0.01	0.08	0.15
<i>1σ</i>	0.10	0.10	0.10

500 Table 2: Annual temperature contributions to sea level during the Last Interglacial for different warming depths.

References

- 505 Anand, P., Elderfield, H., and Conte, M. H.: Calibration of Mg/Ca thermometry in planktonic foraminifera from a sediment trap time series, *Paleoceanography*, 18, 10.1029/2002pa000846, 2003.
- Bakker, P., Stone, E. J., Charbit, S., Gröger, M., Krebs-Kanzow, U., Ritz, S. P., Varma, V., Khon, S., Lunt, D. J., Mikolajewicz, U., Prange, M., Renssen, H., Schneider, B., and Schulz, M.: Last interglacial temperature evolution – a model inter-comparison, *Clim Past*, 9, 605-619, 10.5194/cp-9-605-2013, 2013.
- 510 Bakker, P., and Renssen, H.: Last interglacial model-data mismatch of thermal maximum temperatures partially explained, *Clim Past*, 10, 1633-1644, 10.5194/cp-10-1633-2014, 2014.
- Bard, E., Rostek, F., and Sonzogni, C.: Interhemispheric synchrony of the last deglaciation inferred from alkenone palaeothermometry, *Nature*, 385, 707-710, 1997.
- Barnola, J. M., Raynaud, D., Korotkevich, Y. S., and Lorius, C.: Vostok ice core provides 160,000 year record of atmospheric CO₂, *Nature*, 329, 408-413, 1987.
- 515 Bengtson, S. A., Menviel, L. C., Meissner, K. J., Missiaen, L., Peterson, C. D., Lisiecki, L. E., and Joos, F.: Lower oceanic $\delta^{13}\text{C}$ during the Last Interglacial compared to the Holocene, *Clim. Past Discuss.*, 2020, 1-27, 10.5194/cp-2020-73, 2020.
- Bijma, J., Erez, J., and Hemleben, C.: Lunar and semi-lunar reproductive cycles in some spinose planktonic foraminifers, *Journal of Foraminiferal Research*, 20, 117-127, 1990.
- 520 Brohan, P., Kennedy, J. J., Harris, I., Tett, S. F. B., and Jones, P. D.: Uncertainty estimates in regional and global observed temperature changes: A new data set from 1850, *Journal of Geophysical Research*, 111, D12106, 10.1029/2005JD006548, 2006.
- Capron, E., Govin, A., Stone, E. J., Masson-Delmotte, V., Mulitza, S., Otto-Bliesner, B., Rasmussen, T. L., Sime, L. C., Waelbroeck, C., and Wolff, E. W.: Temporal and spatial structure of multi-millennial temperature changes at high latitudes during the Last Interglacial, *Quatern Sci Rev*, 103, 116-133, 10.1016/j.quascirev.2014.08.018, 2014.
- 525 Capron, E., Govin, A., Feng, R., Otto-Bliesner, B. L., and Wolff, E. W.: Critical evaluation of climate syntheses to benchmark CMIP6/PMIP4 127 ka Last Interglacial simulations in the high-latitude regions, *Quatern Sci Rev*, 168, 137-150, 2017.
- Chadwick, M., Allen, C. S., Sime, L. C., and Hillenbrand, C. D.: Analysing the timing of peak warming and minimum winter sea-ice extent in the Southern Ocean during MIS 5e, *Quaternary Science Reviews*, 229, 106134, 10.1016/j.quascirev.2019.106134, 2020.
- 530 Clark, P. U., He, F., Gollledge, N. R., Mitrovica, J. X., Dutton, A., Hoffman, J. S., and Dendy, S.: Oceanic forcing of penultimate deglacial and last interglacial sea-level rise, *Nature*, 577, 660-664, 10.1038/s41586-020-1931-7, 2020.
- CLIMAP: The Last Interglacial ocean, *Quatern Res*, 21, 123-224, 1984.
- Cortese, G., Dunbar, G. B., Carter, L., Scott, G., Bostock, H., Bowen, M., Crundwell, M., Hayward, B. W., Howard, W., Martínez, J. I., Moy, A., Neil, H., Sabaa, A., and Sturm, A.: Southwest Pacific Ocean response to a warmer world: Insights from Marine Isotope Stage 5e, *Paleoceanography*, 28, 585-598, 10.1002/palo.20052, 2013.
- 535 Dakos, V., Scheffer, M., van Nes, E. H., Brovkin, V., Petoukhov, V., and Held, H.: Slowing down as an early warning signal for abrupt climate change, *Proceedings of the National Academy of Sciences*, 105, 14308-14312, 2008.
- DeConto, R. M., and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591-597, 10.1038/nature17145, 2016.
- 540 Dieckmann, G., Spindler, M., Lange, M. A., Ackley, S. F., and Eicken, H.: Antarctic sea ice: a habitat for the foraminifer *Neogloboquadrina pachyderma*, *Journal of Foraminiferal Research*, 21, 182-189, 1991.
- Doblin, M. A., and van Sebille, E.: Drift in ocean currents impacts intergenerational microbial exposure to temperature, *PNAS*, 113, 5700-5705, 10.1073/pnas.1521093113, 2016.
- 545 Dutton, A., Carlson, A., Long, A., Milne, G., Clark, P., DeConto, R., Horton, B., Rahmstorf, S., and Raymo, M.: Sea-level rise due to polar ice-sheet mass loss during past warm periods, *Science*, 349, 153, 2015.
- Elderfield, H., and Ganssen, G.: Past temperature and $\delta^{18}\text{O}$ of surface ocean waters inferred from foraminiferal Mg/Ca ratios, *Nature*, 405, 442-445, 10.1038/35013033, 2000.
- Esper, O., and Gersonde, R.: Quaternary surface water temperature estimations: New diatom transfer functions for the Southern Ocean, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 414, 1-19, 2014.
- 550

- Fischer, H., Meissner, K. J., Mix, A. C., Abram, N. J., Austermann, J., Brovkin, V., Capron, E., Colombaroli, D., Daniau, A.-L., Dyez, K. A., Felis, T., Finkelstein, S. A., Jaccard, S. L., McClymont, E. L., Rovere, A., Sutter, J., Wolff, E. W., Affolter, S., Bakker, P., Ballesteros-Cánovas, J. A., Barbante, C., Caley, T., Carlson, A. E., Churakova, O., Cortese, G., Cumming, B. F., Davis, B. A. S., de Vernal, A., Emile-Geay, J., Fritz, S. C., Gierz, P., Gottschalk, J., Holloway, M. D., Joos, F., Kucera, M., Loutre, M.-F., Lunt, D. J., Marcisz, K., Marlon, J. R., Martinez, P., Masson-Delmotte, V., Nehrbass-Ahles, C., Otto-Bliesner, B. L., Raible, C. C., Risebrobakken, B., Sánchez Goñi, M. F., Arrigo, J. S., Sarnthein, M., Sjolte, J., Stocker, T. F., Velasquez Álvarez, P. A., Tinner, W., Valdes, P. J., Vogel, H., Wanner, H., Yan, Q., Yu, Z., Ziegler, M., and Zhou, L.: Palaeoclimate constraints on the impact of 2 °C anthropogenic warming and beyond, *Nature Geoscience*, 11, 474-485, 10.1038/s41561-018-0146-0, 2018.
- 555
- 560 Fogwill, C. J., Turney, C. S. M., Meissner, K. J., Golledge, N. R., Spence, P., Roberts, J. L., England, M. H., Jones, R. T., and Carter, L.: Testing the sensitivity of the East Antarctic Ice Sheet to Southern Ocean dynamics: past changes and future implications, *J Quatern Sci*, 29, 91-98, 10.1002/jqs.2683, 2014.
- Fogwill, C. J., Phipps, S. J., Turney, C. S. M., and Golledge, N. R.: Sensitivity of the Southern Ocean to enhanced regional Antarctic ice sheet meltwater input, *Earth's Future*, 2015EF000306, doi: 000310.001002/002015EF000306, 10.1002/2015EF000306, 2015.
- 565 Galaaen, E. V., Ninnemann, U. S., Irvall, N., Kleiven, H. F., Rosenthal, Y., Kissel, C., and Hodell, D. A.: Rapid reductions in North Atlantic Deep Water during the peak of the Last Interglacial period, *Science*, 343, 1129-1132, 10.1126/science.1248667, 2014.
- Golledge, N. R., Kowalewski, D. E., Naish, T. R., Levy, R. H., Fogwill, C. J., and Gasson, E. G. W.: The multi-millennial Antarctic commitment to future sea-level rise, *Nature*, 526, 421-425, 10.1038/nature15706, 2015.
- 570 Govin, A., Capron, E., Tzedakis, P. C., Verheyden, S., Ghaleb, B., Hillaire-Marcel, C., St-Onge, G., Stoner, J. S., Bassinot, F., Bazin, L., Blunier, T., Combourieu-Nebout, N., El Ouahabi, A., Genty, D., Gersonde, R., Jimenez-Amat, P., Landais, A., Martrat, B., Masson-Delmotte, V., Parrenin, F., Seidenkrantz, M. S., Veres, D., Waelbroeck, C., and Zahn, R.: Sequence of events from the onset to the demise of the Last Interglacial: Evaluating strengths and limitations of chronologies used in climatic archives, *Quatern Sci Rev*, 129, 1-36, 10.1016/j.quascirev.2015.09.018, 2015.
- 575 Grant, K. M., Rohling, E. J., Ramsey, C. B., Cheng, H., Edwards, R. L., Florindo, F., Heslop, D., Marra, F., Roberts, A. P., Tamisiea, M. E., and Williams, F.: Sea-level variability over five glacial cycles, *Nature Comms*, 5, 5076, 10.1038/ncomms6076, 2014.
- Hansen, J. E.: A slippery slope: How much global warming constitutes 'dangerous anthropogenic interference'?, *Climatic Change*, 68, 269-279, 2005.
- 580 Hayes, C. T., Martínez-García, A., Hasenfratz, A. P., Jaccard, S. L., Hodell, D. A., Sigman, D. M., Haug, G. H., and Anderson, R. F.: A stagnation event in the deep South Atlantic during the last interglacial period, *Science*, 346, 1514-1517, 10.1126/science.1256620, 2014.
- Hellweger, F. L., van Sebille, E., Calfee, B. C., Chandler, J. W., Zinser, E. R., Swan, B. K., and Fredrick, N. D.: The Role of Ocean Currents in the Temperature Selection of Plankton: Insights from an Individual-Based Model, *PLOS ONE*, 11, e0167010, 10.1371/journal.pone.0167010, 2016.
- 585 Hoffman, J. S., Clark, P. U., Parnell, A. C., and He, F.: Regional and global sea-surface temperatures during the last interglaciation, *Science*, 355, 276-279, 10.1126/science.aai8464, 2017.
- Huang, B., Menne, M. J., Boyer, T., Freeman, E., Gleason, B. E., Lawrimore, J. H., Liu, C., Rennie, J. J., Schreck, C. J., III, Sun, F., Vose, R., Williams, C. N., Yin, X., and Zhang, H.-M.: Uncertainty Estimates for Sea Surface Temperature and Land Surface Air Temperature in NOAA GlobalTemp Version 5, *Journal of Climate*, 33, 1351-1379, 10.1175/jcli-d-19-0395.1, 2020.
- 590 Hönisch, B., Allen, K. A., Lea, D. W., Spero, H. J., Eggins, S. M., Arbuszewski, J., deMenocal, P., Rosenthal, Y., Russell, A. D., and Elderfield, H.: The influence of salinity on Mg/Ca in planktic foraminifers—Evidence from cultures, core-top sediments and complementary $\delta^{18}\text{O}$, *Geochim Cosmo Acta*, 121, 196-213, 2013.
- 595 IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp., 2013.
- Jones, R. T., Turney, C. S. M., Lang, B., Brooks, S. J., Rundgren, M., Hammarlund, D., Björck, S., and Fogwill, C. J.: Delayed maximum northern European summer temperatures during the Last Interglacial as a result of Greenland Ice Sheet melt, 600 *Geology*, 45, 23-26, 10.1130/g38402.1, 2017.

- Jonkers, L., Reynolds, C. E., Richey, J., and Hall, I. R.: Lunar periodicity in the shell flux of planktonic foraminifera in the Gulf of Mexico, *Biogeosciences*, 12, 3061-3070, 2015.
- Jonkers, L., Hillebrand, H., and Kucera, M.: Global change drives modern plankton communities away from the pre-industrial state, *Nature*, 570, 372-375, 10.1038/s41586-019-1230-3, 2019.
- 605 Kandiano, E. S., Bauch, H. A., and Müller, A.: Sea surface temperature variability in the North Atlantic during the last two glacial-interglacial cycles: comparison of faunal, oxygen isotopic, and Mg/Ca-derived records, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 204, 145-164, 2004.
- Kienast, S. S., Winckler, G., Lippold, J., Albani, S., and Mahowald, N. M.: Tracing dust input to the global ocean using thorium isotopes in marine sediments: ThoroMap, *Global Biogeochem Cycles*, 30, 1526-1541, doi:10.1002/2016GB005408, 2016.
- 610 Kim, S.-J., Crowley, T.J. and Stössel, A.: Local orbital forcing of Antarctic climate change during the Last Interglacial. In: *Science*, 1998.
- Kopp, R. E., Simons, F. J., Mitrovica, J. X., Maloof, A. C., and Oppenheimer, M.: Probabilistic assessment of sea level during the last interglacial stage, *Nature*, 462, 863-867, 2009.
- 615 Köhler, P., Nehrbass-Ahles, C., Schmitt, J., Stocker, T. F., and Fischer, H.: A 156 kyr smoothed history of the atmospheric greenhouse gases CO₂, CH₄, and N₂O and their radiative forcing, *Earth Syst. Sci. Data*, 9, 363-387, 10.5194/essd-9-363-2017, 2017.
- Lange, M., and van Sebille, E.: Parcels v0.9: prototyping a Lagrangian ocean analysis framework for the petascale age, *Geosci. Model Dev.*, 10, 4175-4186, 10.5194/gmd-10-4175-2017, 2017.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A. C. M., and Levrard, B.: A long-term numerical solution for the insolation quantities of the earth, *Astronomy & Astrophysics*, 428, 261-285, 10.1051/0004-6361:20041335, 2004.
- 620 Lenton, T. M., Held, H., Kriegler, E., Hall, J. W., Lucht, W., Rahmstorf, S., and Schellnhuber, H. J.: Tipping elements in the Earth's climate system, *PNAS*, 105, 1786-1793, 2008.
- Lisiecki, L. E., and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records, *Paleoceanography*, 20, doi:10.1029/2004PA001071, 2005.
- 625 Lunt, D. J., Abe-Ouchi, A., Bakker, P., Berger, A., Braconnot, P., Charbit, S., Fischer, N., Herold, N., Jungclauss, J. H., Khon, V. C., Krebs-Kanzow, U., Langebroeck, P. M., Lohmann, G., Nisancioglu, K. H., Otto-Bliesner, B. L., Park, W., Pfeiffer, M., Phipps, S. J., Prange, M., Rachmayani, R., Renssen, H., Rosenbloom, N., Schneider, B., Stone, E. J., Takahashi, K., Wei, W., Yin, Q., and Zhang, Z. S.: A multi-model assessment of last interglacial temperatures, *Clim Past*, 9, 699-717, 10.5194/cp-9-699-2013, 2013.
- 630 Martinson, D. G., Pisias, N. G., Hays, J. D., Imbrie, J., Moore, T. C., and Shackleton, N. J.: Age dating and the orbital theory of the Ice Ages: Development of a high-resolution 0 to 300,000-year chronostratigraphy, *Quaternary Research*, 27, 1-29, 1987.
- Masumoto, Y., Sasaki, H., Kagimoto, T., Komori, N., Ishida, A., Sasai, Y., Miyama, T., Motoi, T., Mitsudera, H., Takahashi, K., Sakuma, H., and Yamagata, T.: A fifty-year eddy-resolving simulation of the world ocean – Preliminary outcomes of OFES (OGCM for the Earth simulator), *Journal of the Earth Simulator*, 1, 35-56, 2004.
- 635 McKay, N. P., Overpeck, J. T., and Otto-Bliesner, B. L.: The role of ocean thermal expansion in Last Interglacial sea level rise, *GRL*, 38, L14605, 10.1029/2011gl048280, 2011.
- Mercer, J. H., and Emiliani, C.: Antarctic ice and interglacial high sea levels, *Science*, 168, 1605-1606, 10.1126/science.168.3939.1605-a, 1970.
- Mercer, J. H.: West Antarctic ice sheet and CO₂ greenhouse effect: a threat of disaster, *Nature*, 271, 321-325, 1978.
- 640 Miller, G. H., Alley, R. B., Brigham-Grette, J., Fitzpatrick, J. J., Polyak, L., Serreze, M. C., and White, J. W. C.: Arctic amplification: can the past constrain the future?, *Quaternary Science Reviews*, 29, 1779-1790, 10.1016/j.quascirev.2010.02.008, 2010.
- Monroy, P., Hernández-García, E., Rossi, V., and López, C.: Modeling the dynamical sinking of biogenic particles in oceanic flow, *Nonlin. Processes Geophys.*, 24, 293-305, 10.5194/npg-24-293-2017, 2017.
- 645 Müller, P. J., Kirst, G., Ruhland, G., von Storch, I., and Rosell-Melé, A.: Calibration of the alkenone paleotemperature index U37K' based on core-tops from the eastern South Atlantic and the global ocean (60°N-60°S), *Geochim Cosmo Acta*, 62, 1757-1772, 10.1016/S0016-7037(98)00097-0, 1998.
- NEEM Community Members: Eemian interglacial reconstructed from a Greenland folded ice core, *Nature*, 493, 489-494, , 2013.

- 650 Nooteboom, P. D., Bijl, P. K., van Sebille, E., von der Heydt, A. S., and Dijkstra, H. A.: Transport bias by ocean currents in sedimentary microplankton assemblages: Implications for paleoceanographic reconstructions, *Paleoceanography and Paleoclimatology*, doi: 10.1029/2019pa003606, 10.1029/2019pa003606,
 Nooteboom, P. D., Delandmeter, P., van Sebille, E., Bijl, P. K., Dijkstra, H. A., and von der Heydt, A. S.: Resolution
 655 10.1371/journal.pone.0238650, 2020.
 Otto-Bliesner, B. L., Rosenbloom, N., Stone, E. J., McKay, N. P., Lunt, D. J., Brady, E. C., and Overpeck, J. T.: How warm was the last interglacial? New model–data comparisons, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 371, doi: 10.1098/rsta.2013.0097, 10.1098/rsta.2013.0097, 2013.
- 660 Overpeck, J., Sturm, M., Francis, J. A., Perovich, D. K., Serreze, M. C., Benner, R., Carmack, E. C., Chapin, F. S. I., Gerlach, S. C., Hamilton, L. C., Hinzman, L. D., Holland, M., Huntington, H. P., Key, J., .R., Lloyd, A. H., MacDonald, G. M., McFadden, J., Noone, D., Prowse, T. D., Schlosser, P., and Vörösmarty, C.: Arctic system on trajectory to new, seasonally ice-free state, *Eos Transactions AGU*, 86, 309–313, 2005.
 Overpeck, J., Otto-Bliesner, B., Miller, G., Muhs, D., Alley, R., and Kiehl, J.: Paleoclimatic evidence for future ice-sheet instability and rapid sea-level rise, *Science*, 311, 1747-1750, 10.1126/science.1115159, 2006.
- 665 PAGES2k Consortium, Emile-Geay, J., McKay, N. P., Kaufman, D. S., von Gunten, L., Wang, J., Anchukaitis, K. J., Abram, N. J., Addison, J. A., Curran, M. A. J., Evans, M. N., Henley, B. J., Hao, Z., Martrat, B., McGregor, H. V., Neukom, R., Pederson, G. T., Stenni, B., Thirumalai, K., Werner, J. P., Xu, C., Divine, D. V., Dixon, B. C., Gergis, J., Mundo, I. A., Nakatsuka, T., Phipps, S. J., Routson, C. C., Steig, E. J., Tierney, J. E., Tyler, J. J., Allen, K. J., Bertler, N. A. N., Björklund, J., Chase, B. M., Chen, M.-T., Cook, E., de Jong, R., DeLong, K. L., Dixon, D. A., Ekaykin, A. A., Ersek, V., Filipsson, H.
 670 L., Francus, P., Freund, M. B., Frezzotti, M., Gaire, N. P., Gajewski, K., Ge, Q., Goosse, H., Gornostaeva, A., Grosjean, M., Horiuchi, K., Hormes, A., Husum, K., Isaksson, E., Kandasamy, S., Kawamura, K., Kilbourne, K. H., Koç, N., Leduc, G., Linderholm, H. W., Lorrey, A. M., Mikhalenko, V., Mortyn, P. G., Motoyama, H., Moy, A. D., Mulvaney, R., Munz, P. M., Nash, D. J., Oerter, H., Opel, T., Orsi, A. J., Ovchinnikov, D. V., Porter, T. J., Roop, H. A., Saenger, C., Sano, M., Sauchyn, D., Saunders, K. M., Seidenkrantz, M.-S., Severi, M., Shao, X., Sicre, M.-A., Sigl, M., Sinclair, K., St. George, S., St. Jacques, J.-M., Thamban, M., Kuwar Thapa, U., Thomas, E. R., Turney, C., Uemura, R., Viau, A. E., Vladimirova, D. O., Wahl, E. R.,
 675 White, J. W. C., Yu, Z., and Zinke, J.: A global multiproxy database for temperature reconstructions of the Common Era, *Scientific Data*, 4, 170088, 10.1038/sdata.2017.88, 2017.
 Past Interglacials Working Group of PAGES: Interglacials of the last 800,000 years, *Reviews of Geophysics*, 54, 162-219, 10.1002/2015RG000482, 2016.
- 680 Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J. M., Basile, I., Bender, M., Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M., Lipenkov, V. Y., Lorius, C., Pepin, L., Ritz, C., Saltzman, E., and Stievenard, M.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399, 429-436, 10.1038/20859, 1999.
 Pisias, N. G., and Mix, A. C.: Spatial and temporal oceanographic variability of the eastern equatorial Pacific during the late Pleistocene: Evidence from radiolaria microfossils, *Paleoceanography*, 12, 381-393, 1997.
- 685 Prahl, F. G., Sparrow, M. A., and Wolfe, G. V.: Physiological impacts on alkenone paleothermometry, *Paleoceanography*, 18, 10.1029/2002pa000803, 2003.
 Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *Journal of Geophysical Research: Atmospheres*, 108, 4407, doi:4410.1029/2002JD002670, 10.1029/2002JD002670, 2003.
- 690 Rembauville, M., Manno, C., Tarling, G. A., Blain, S., and Salter, I.: Strong contribution of diatom resting spores to deep-sea carbon transfer in naturally iron-fertilized waters downstream of South Georgia, *Deep Sea Research Part I: Oceanographic Research Papers*, 115, 22-35, 10.1016/j.dsr.2016.05.002, 2016.
- 695 Rohling, E. J., Cane, T. R., Cooke, S., Sprovieri, M., Bouloubassi, I., Emeis, K. C., Schiebel, R., Kroon, D., Jorissen, F. J., Lorre, A., and Kemp, A. E. S.: African monsoon variability during the previous interglacial maximum, *EPSL*, 202, 61-75, 2002.
 Rohling, E. J., Grant, K., Hemleben, C., Siddall, M., Hoogakker, B. A. A., Bolshaw, M., and Kucera, M.: High rates of sea-level rise during the last interglacial period, *Nature Geosci*, 1, 38-42, 10.1038/ngeo.2007.28, 2008.

- 700 Rohling, E. J., Hibbert, F. D., Williams, F. H., Grant, K. M., Marino, G., Foster, G. L., Hennekam, R., de Lange, G. J., Roberts, A. P., Yu, J., Webster, J. M., and Yokoyama, Y.: Differences between the last two glacial maxima and implications for ice-sheet, $\delta^{18}\text{O}$, and sea-level reconstructions, *Quatern Sci Rev*, 176, 1-28, 10.1016/j.quascirev.2017.09.009, 2017.
- Rohling, E. J., Hibbert, F. D., Grant, K. M., Galaasen, E. V., Irvall, N., Kleiven, H. F., Marino, G., Ninnemann, U., Roberts, A. P., Rosenthal, Y., Schulz, H., Williams, F. H., and Yu, J.: Asynchronous Antarctic and Greenland ice-volume contributions to the last interglacial sea-level highstand, *Nature Communications*, 10, 5040, 10.1038/s41467-019-12874-3, 2019.
- 705 Schellnhuber, H. J., Rahmstorf, S., and Winkelmann, R.: Why the right climate target was agreed in Paris, *Nature Climate Change*, 6, 649, 10.1038/nclimate3013, 2016.
- Schneider, R., Schmitt, J., Köhler, P., Joos, F., and Fischer, H.: A reconstruction of atmospheric carbon dioxide and its stable carbon isotopic composition from the penultimate glacial maximum to the last glacial inception, *Climate of the Past*, 9, 2507-2523, 2013.
- 710 Schneider, R. R., Müller, P. J., and Acheson, R.: Atlantic alkenone sea-surface temperature records, in: *Reconstructing Ocean History: A Window into the Future*, Kluwer Academic/Plenum Publishers, New York, 33-55, 1999.
- Segev, E., Castañeda, I. S., Sikes, E. L., Vlamakis, H., and Kolter, R.: Bacterial influence on alkenones in live microalgae, *Journal of Phycology*, 52, 125-130, 10.1111/jpy.12370, 2016.
- 715 Shackleton, S., Baggenstos, D., Menking, J. A., Dyonisius, M. N., Bereiter, B., Bauska, T. K., Rhodes, R. H., Brook, E. J., Petrenko, V. V., McConnell, J. R., Kellerhals, T., Häberli, M., Schmitt, J., Fischer, H., and Severinghaus, J. P.: Global ocean heat content in the Last Interglacial, *Nature Geosci*, 13, 77-81, 10.1038/s41561-019-0498-0, 2020.
- Sikes, E. L., O'Leary, T., Nodder, S. D., and Volkman, J. K.: Alkenone temperature records and biomarker flux at the subtropical front on the Chatham Rise, SW Pacific Ocean, *Deep Sea Research Part I: Oceanographic Research Papers*, 52, 721-748, 2005.
- 720 Spindler, M.: On the salinity tolerance of the planktonic foraminifer *Neogloboquadrina pachyderma* from Antarctic sea ice, *Proc. NIPR Symp. Polar Biol*, 1996, 85-91,
- Steffen, W., Rockström, J., Richardson, K., Lenton, T. M., Folke, C., Liverman, D., Summerhayes, C. P., Barnosky, A. D., Cornell, S. E., Crucifix, M., Donges, J. F., Fetzer, I., Lade, S. J., Scheffer, M., Winkelmann, R., and Schellnhuber, H. J.: Trajectories of the Earth System in the Anthropocene, *PNAS*, 115, 8252-8259, 10.1073/pnas.1810141115, 2018.
- 725 Sutter, J., Gierz, P., Grosfeld, K., Thoma, M., and Lohmann, G.: Ocean temperature thresholds for Last Interglacial West Antarctic Ice Sheet collapse, *GRL*, 43, 2675-2682, 10.1002/2016GL067818, 2016.
- Thomas, Z. A., Kwasniok, F., Boulton, C. A., Cox, P. M., Jones, R. T., Lenton, T. M., and Turney, C. S. M.: Early warnings and missed alarms for abrupt monsoon transitions, *Clim Past*, 11, 1621-1633, 10.5194/cp-11-1621-2015, 2015.
- 730 Thomas, Z. A.: Using natural archives to detect climate and environmental tipping points in the Earth System, *Quatern Sci Rev*, 152, 60-71, 10.1016/j.quascirev.2016.09.026, 2016.
- Thomas, Z. A., Jones, R. T., Turney, C. S. M., Golledge, N., Fogwill, C., Bradshaw, C. J. A., Meniel, L., McKay, N. P., Bird, M., Palmer, J., Kershaw, P., Wilmschurst, J., and Muscheler, R.: Tipping elements and amplified polar warming during the Last Interglacial, *Quaternary Science Reviews*, 233, 106222, 10.1016/j.quascirev.2020.106222, 2020.
- 735 Turney, C. S. M., and Jones, R. T.: Does the Agulhas Current amplify global temperatures during super-interglacials?, *J Quatern Sci*, 25, 839-843, 2010.
- Turney, C. S. M., and Jones, R. T.: Response to Comment on 'Does the Agulhas Current amplify global temperatures during super-interglacials?', *J Quatern Sci*, 26, 870-871, 10.1002/jqs.1556, 2011.
- 740 Turney, C. S. M., Jones, R., McKay, N., Van Sebille, E., Thomas, Z., Hillenbrand, C.-D., and Fogwill, C.: A global reconstruction of sea-surface temperatures for the Last Interglacial (129-116 kyr). In: *PANGAEA*, <https://doi.pangaea.de/10.1594/PANGAEA.904381>, 2019.
- Turney, C. S. M., Fogwill, C. J., Golledge, N. R., McKay, N. P., van Sebille, E., Jones, R. T., Etheridge, D., Rubino, M., Thornton, D. P., Davies, S. M., Ramsey, C. B., Thomas, Z. A., Bird, M. I., Munksgaard, N. C., Kohno, M., Woodward, J., Winter, K., Weyrich, L. S., Rootes, C. M., Millman, H., Albert, P. G., Rivera, A., van Ommen, T., Curran, M., Moy, A., Rahmstorf, S., Kawamura, K., Hillenbrand, C.-D., Weber, M. E., Manning, C. J., Young, J., and Cooper, A.: Early Last Interglacial ocean warming drove substantial ice mass loss from Antarctica, *PNAS*, 117, 3996-4006, 10.1073/pnas.1902469117, 2020.
- 745 Tzedakis, P. C., Drysdale, R. N., Margari, V., Skinner, L. C., Meniel, L., Rhodes, R. H., Taschetto, A. S., Hodell, D. A., Crowhurst, S. J., Hellstrom, J. C., Fallick, A. E., Grimalt, J. O., McManus, J. F., Martrat, B., Mokeddem, Z., Parrenin, F.,

- 750 Regattieri, E., Roe, K., and Zanchetta, G.: Enhanced climate instability in the North Atlantic and southern Europe during the Last Interglacial, *Nature Comms*, 9, 4235, doi: 4210.1038/s41467-41018-06683-41463, 10.1038/s41467-018-06683-3, 2018.
- van Sebille, E., England, M. H., Zika, J. D., and Sloyan, B. M.: Tasman leakage in a fine-resolution ocean model, *GRL*, 39, L06601, 10.1029/2012GL051004, 2012.
- 755 van Sebille, E., Scussolini, P., Durgadoo, J. V., Peeters, F. J. C., Biastoch, A., Weijer, W., Turney, C., Paris, C. B., and Zahn, R.: Ocean currents generate large footprints in marine palaeoclimate proxies, *Nature Comms*, 6, 6521, 10.1038/ncomms7521, 2015.
- Viebahn, J. P., Heydt, A. S., Le Bars, D., and Dijkstra, H. A.: Effects of Drake Passage on a strongly eddying global ocean, *Paleoceanography and Paleoclimatology*, 31, 564-581, 2016.
- Visser, K., Thunell, R., and Stott, L.: Magnitude and timing of temperature change in the Indo-Pacific warm pool during deglaciation, *Nature*, 421, 152-155, 2003.
- 760 Vogelsang, E., Sarnthein, M., and Pflaumann, U.: $\delta^{18}\text{O}$ stratigraphy, chronology, and sea surface temperatures of Atlantic sediment records (GLAMAP-2000 Kiel), Universität Kiel, Kiel, Germany, 244, 2001.
- von Gyldenfeldt, A.-B., Carstens, J., and Meincke, J.: Estimation of the catchment area of a sediment trap by means of current meters and foraminiferal tests, *Deep Sea Research Part II: Topical Studies in Oceanography*, 47, 1701-1717, 10.1016/S0967-0645(00)00004-7, 2000.
- 765 Waelbroeck, C., Frank, N., Jouzel, J., Parrenin, F., Masson-Delmotte, V., and Genty, D.: Transferring radiometric dating of the last interglacial sea level high stand to marine and ice core records, *EPSL*, 265, 183-194, 2008.
- Wang, Y. J., Cheng, H., Edwards, R. L., Kong, X. G., Shao, X. H., Chen, S. T., Wu, J. Y., Jiang, X. Y., Wang, X. F., and An, Z. S.: Millennial- and orbital-scale changes in the East Asian monsoon over the past 224,000 years, *Nature*, 451, 1090-1093, 10.1038/nature06692, 2008.
- 770 Wessel, P., Smith, W. H., Scharroo, R., Luis, J., and Wobbe, F.: Generic mapping tools: improved version released, *Eos, Transactions American Geophysical Union*, 94, 409-410, 2013.
- White, J. W. C.: Don't touch that dial, *Nature*, 364, 186, 1993.