



Interactive comment

Interactive comment on "Radiative forcing of climate change from the Copernicus reanalysis of atmospheric composition" by Nicolas Bellouin et al.

Anonymous Referee #1

Earth Syst. Sci. Data Discuss.,

https://doi.org/10.5194/essd-2019-251-RC1, 2020 © Author(s) 2020. This work is distributed under

the Creative Commons Attribution 4.0 License.

Received and published: 27 February 2020

This paper describes a data set of climate radiative forcing (2003-2016) calculated from the Copernicus reanalysis of atmospheric composition, including CO2, CH4, O3, and aerosol. Because these atmospheric constituents are constrained to some extent by observations through data assimilation approach, the derived radiative forcing is believed to be more reliable than free-run GCM simulations. The data set would be useful to several applications. The paper is generally well written. I would recommend the paper be published in ESSD after some clarifications (mostly technical nature).

1. in the abstract, I would suggest that they split ozone and aerosol radiative forcing numbers (line 25-26) into two components, i.e., tropospheric and stratospheric for Printer-friendly version

Discussion paper



ozone, ARI and ACI for aerosol. Nevertheless, it says that they are dealing with 6 forcing agents.

2. line 85-94: it is helpful to say something more about the CAMS, e.g., what kinds of observations have been used in the reanalysis. A table may be adequate.

3. line 100: "2003-2018" I found that all related figures are for 2003-2016.

4. line 146: neglecting aerosol scattering in the LW spectrum may introduce significant uncertainty. Maybe it is useful to discuss the uncertainty here.

5. line 210-211: "Anthropogenic fractions therefore peak in late summer in South America and southern Africa". I think it should be "boreal summer".

6. line 259-261: Do recent satellite-based estimates of above-cloud aerosol radiative effect justify their neglect of cloudy-sky radiative effect?

7. Line 425-427: Randles et al. (2013) assessed the uncertainty in RF_ARI associated with the two-stream approximation, which could be included here.

8. Line 519-520: "Interestingly, cloud masking of RFari is larger than RFaci". it is kind of confusing.

9. line 622: need to define "rate of change". Which two years are used to calculate the change, year n and n-1 or year n and n+1?

10. line 601-606: which satellite AOD data has been used in the reanalysis? can these trends be attributed to spurious trends in the satellite AOD?

11. Figure 7. you need a legend for those colored lines.

ESSDD

Interactive comment

Printer-friendly version





Interactive comment on Earth Syst. Sci. Data Discuss., https://doi.org/10.5194/essd-2019-251, 2020.