

“An update of IPCC climate reference regions for subcontinental analysis of climate model data: Definition and aggregated datasets” by Iturbide et al.

Response to Topical Editor

Topical Editor Decision: Publish subject to minor revisions (review by editor) (14 Aug 2020)
by David Carlson

Comments to the Author:

Please respond explicitly to all comments and suggestions in the most recent review.

Dear Editor, we have revised the document taking into account all comments and suggestions received. Below we provide point-by-point responses (in blue) for the comments (in black) and include the revised document including tracked changes.

In addition, please note:

Please write all zenodo DOI links in full one-click format, e.g <https://doi.org/10.5281/zenodo.3968318> (as you have it in the Iturbide et al. 2020 reference). Show DOI (permanent) links before (ephemeral) GitHub links.

Done

Give us page numbers or use a continuous sequence of line numbers.

We have include page numbers in the revised manuscript.

Copernicus journals use numbers rather letters to denote institutional affiliations. You might as well change these now to save the proofreaders one task.

Done

Section 3.1 line 6: something missing or mis-stated here.

Thank you. This has been fixed in the revised version.

Section 4 line 45: something wrong or extra here

Thank you. An extra word has been removed. Now it reads: *“...the ATLAS initiative presented here facilitates intercomparison of results and consistency checks for the reference climatic regions.”*

“An update of IPCC climate reference regions for subcontinental analysis of climate model data: Definition and aggregated datasets” by Iturbide et al.

Response to Referee 1

I would like to thank the authors for a very thorough response to my comment. I am generally satisfied with their work and the revised manuscript’s incorporation of those changes. Whilst I would be happy to see this manuscript now published, I have noticed a few minor additional things that I would commented on beforehand.

Dear referee. Thank you very much for the time devoted to the revision of our manuscript and the positive feedback received. Please find below a point-by point response (in blue) to your comments (in black).

- Can you put Giorgi in either italics or perhaps quotation marks in the phrase “Giorgi reference regions”? This differentiates the name of the regions from the name of the author.

Done

- Please change “larger number of smaller regions” to “higher number of smaller regions”. This avoids a directly opposite adjectives.

Done

- Are you sure that you want to label your new dataset “updated IPCC WGI reference regions”? Calling it “IPCC WGI reference regions, version 2” would allow for subsequent updates. This change would only need to be implemented in one or two places and not the title

We agree with the referee and have included version numbers to refer to the previous three versions (*Giorgi*, *SREX* and *AR5*) and the current update (version 4). This has been described in the introduction and in Figure 1.

- Please remove “used here” after GPC. It seems unnecessary

Done

- Thank you for providing more clarity about the interpolation. I’m a little unsure why you are interpolating the land and separately. It’s not what I would have done, but I doubt it is wrong. Firstly I would have interpolated the region and land/sea mask onto the data grid (more for computational efficiency rather than anything else). Certainly for interpolating temperature anomalies, I would expect the land to provide some useful additional information for the values over the ocean. I’m not sure that the manuscript needs any changes or discussion of this issues though.

In some cases, the analysis of land and sea are done separately, particularly in the case of temperatures, highly influenced by the land/sea contrast. Therefore, land/sea masks are

used in the interpolation process to make sure that the land/sea values provided in the final grid correspond to actual land/sea model values.

- In the old definitions, CAR was computed across both land and sea combined. Here you seem to redefine it as having separate land and sea values. Can you please document and justify this change?

The higher resolution of CMIP6 models provides higher information over land in this region. The number of land gridboxes increases from 1 to 19 when changing from 2 to 1 degrees reference grid (see Figures 5d and 5a). Therefore, this region now allows for both land and sea analysis. This has been documented in the new version of the manuscript.

- I appreciate that you've branched a region off for Madagascar. The manuscript text does not mention this however.

Thank you for pointing this out. The revised manuscript introduces this new region.

- Can you please adopt only 3 letter acronyms for NEAF etc? I suggest dropping the A from these regions.

We kept the two letters for Africa (AF) used in AR5 to avoid confusion with Asia (A). After analysing several options, finally we opted by treating equally all continents and use AF for Africa instead of dropping the first letter using "F" instead. The length of the code shouldn't be a problem as long as it is informative and concise enough.

- In the old definitions, SAS was computed twice having separate land and sea values. Now it is only land (as the neighbouring seas have been branched off). This is fine, but it needs documenting.

This has been documented in the revised manuscript.

- You may want to consider renaming BOB. It's an English man's name (unlike the others). Maybe use BGL instead

Bay of Bengal (BOB) is widely used for this region. We thank the referee for this suggestion, but would rather keep the original name thus avoiding changes (that are not critical) for names and regions at this stage.

- I suggest you find a different acronym for ARS. It's too close to the slight rude English word "arse" for comfort.

As in the previous case, we prefer to keep the original acronym and avoid non-critical changes in regions and/or names at this stage.

- In the old definitions, SEA was computed twice, having separate land and sea values. This convention has now been introduced for MED, but dropped for South East Asia. Not only is this choice not documented, I don't understand the reason for it.

We thank the referee for pointing out this issue. We have modified the regions to include SEA as both land and ocean, as in the case of CAR and MED.

- I appreciate the additional text you have added about the oceans and coast. I feel that a few sentence explaining the ocean regions would be helpful. Whilst they are mainly self-explanatory, I know that the definition of the Southern Ocean is somewhat fluid. Personally, I'm happy with your choice, but I still think that you should justify it.

We agree with the referee and have now added a paragraph describing the new oceanic regions and providing a reference for the definition of the Southern Ocean and reasoning for the choice of the extent of the equatorial oceanic regions (see below).

- The definition of the equatorial band of the oceans has expanded to 10oS to either 7.6oN or 7oN (according to the csv file on GitHub) from the conventional 5oS-5oN (used in the old regions and for Nino averages). I guess this is due Sri Lanka – but again you ought to justify and document this.

The definition of “equatorial” in version 3 as the region bounded by 5N/S probably relates to that used for El Nino indices. However, Indian Ocean Dipole indices use a region bounded by 10N/S so to include both within the version 4 regions the latter range was chosen.

- I would change “single run” to “one ensemble member per model” in the sentence about the aggregated-dataset directory.

Done

- Please change “evidences” to something like “project”. Evidence is too strong for something based on model projections.

Thank you. Done.

- Would it be possible to alter the acknowledgements sentence to acknowledge the Australian commentators and to note that I chose to waive my anonymity.

Thank you again. This is particularly relevant due to the time and effort you all spent on our paper. This has been included in the revised manuscript: *“The authors are also grateful to the two reviewers (Chris Brierley and one anonymous) who helped to improve the original manuscript with a detailed in deep revision providing constructive comments, and to Michael Grose and Jason Evans who provided useful comments during the interactive discussion.”*

- The caption for Fig. 6 discuss 55 regions. Please update.

Done.

1 **An update of IPCC climate reference regions for subcontinental analysis of climate model**
 2 **data: Definition and aggregated datasets**
 3

4 Maialen Iturbide¹, José Manuel Gutiérrez¹, Lincoln Muniz Alves², Joaquín Bedia³, Ruth
 5 Cerezo-Mota⁴, Ezequiel Cimorelli⁵, Antonio S. Cofiño³, Alejandro Di Luca⁵, Sergio
 6 Henrique Faria^{6,7}, Irina Gorodetskaya⁸, Mathias Hauser⁹, Sixto Herrera³, Kevin Hennessy¹⁰,
 7 Helene T. Hewitt¹¹, Richard G. Jones^{11,12}, Svitlana Krakovska^{13,14}, Rodrigo Manzanas^{15,3},
 8 Daniel Martínez-Castro^{16,17}, Gemma Teresa Narisma¹⁸, Intan S. Nurhati¹⁹, Izidine Pinto²⁰,
 9 Sonia I. Seneviratne⁹, Bart van den Hurk²¹, and Carolina S. Vera²²

Eliminado: ^{a...} José Manuel Gutiérrez ^{a...}, Lincoln Muniz Alves ^{b...}, Joaquín Bedia ^{c...}, Ruth Cerezo-Mota ^{d...}, Ezequiel Cimorelli ^{e...}, Antonio S. Cofiño ^{e...}, Alejandro Di Luca ^{e...}, Sergio Henrique Faria ^{f...f2...}, Irina Gorodetskaya ^{g...}, Mathias Hauser ^{h...}, Sixto Herrera ^{c...}, Kevin Hennessy ^{i...0}, Helene T. Hewitt ^{j...1}, Richard G. Jones ^{j...1,k...2}, Svitlana Krakovska ^{l...3,m...4}, Rodrigo Manzanas ^{n...5c...}, Daniel Martínez-Castro ^{o...6,p...7}, Gemma Teresa Narisma ^{q...8}, Intan S. Nurhati ^{r...9}, Izidine Pinto ^{s...0}, Sonia I. Seneviratne ^{h...}, Bart van den Hurk ^{t...1}, and Carolina S. Vera ^u ... [1]

10 ¹Grupo de Meteorología. Instituto de Física de Cantabria (CSIC-UC). Santander, Spain

Eliminado: ^a

11 ²National Institute for Space Research. Sa ã José dos Campos, Brazil

Eliminado: ^b

12 ³Grupo de Meteorología. Dpto. Matemática Aplicada y Ciencias de la Computación. University of Cantabria. Santander, Spain

Eliminado: ^c

14 ⁴Universidad Nacional Autónoma de México (UNAM). Mexico city, Mexico

Eliminado: ^d

15 ⁵Climate Change Research Centre and ARC Centre of Excellence for Climate Extremes, University of New South Wales. Sydney, Australia

Eliminado: ^e

17 ⁶Basque Centre for Climate Change (BC3), Leioa, Spain

Eliminado: ^f

18 ⁷IKERBASQUE, Basque Foundation for Science, Bilbao, Spain

Eliminado: ^{f2}

19 ⁸Centre for Environmental and Marine Studies Department of Physics, University of Aveiro. Aveiro, Portugal

Eliminado: ^g

21 ⁹Institute for Atmospheric and Climate Science, ETH Zurich. Zurich, Switzerland

Eliminado: ^h

22 ¹⁰CSIRO Oceans and Atmosphere. Melbourne, Australia

Eliminado: ^{i...0}CSIRO Oceans and Atmosphere. MelbourneCanberra... [2]

23 ¹¹Met Office Hadley Centre. Exeter, United Kingdom

Eliminado: ^j

24 ¹²School of Geography and Environment, University of Oxford. UK

Eliminado: ^k

25 ¹³Ukrainian Hydrometeorological Institute. Kyiv, Ukraine

Eliminado: ^l

26 ¹⁴State Institution National Antarctic Scientific Center. Kyiv, Ukraine

Eliminado: ^m

27 ¹⁵Intergovernmental Panel on Climate Change (IPCC), WGI-TSU, Université Paris-Saclay. Paris, France

Eliminado: ⁿ

28 ¹⁶Instituto de Meteorología de Cuba. La Habana, Cuba

Eliminado: ^o

29 ¹⁷Instituto Geofísico del Perú. Lima, Perú

Eliminado: ^p

30 ¹⁸Manila Observatory, Ateneo de Manila University campus. Quezon City, Philippines

Eliminado: ^q

31 ¹⁹Research Center for Oceanography. Indonesian Institute of Sciences. Jakarta, Indonesia

Eliminado: ^r

32 ²⁰Climate System Analysis Group (CSAG), University of Cape Town. South Africa

Eliminado: ^s

Con formato: Número de página

1 ²¹ DELTARES. Delft, The Netherlands

Eliminado: ^t

2 ²² Departamento de Ciencias de la Atmósfera y los Océanos, FCEyN-UBA. Centro de Investigaciones del
3 Mar y la Atmósfera (CIMA), Instituto Franco Argentino sobre Estudios de Clima y sus Impactos (UMI
4 IFAECI)/CNRS-CONICET. Buenos Aires, Argentina

Eliminado: ^u

5

6 **Abstract.** Several sets of reference regions have been used in the literature for the regional synthesis of
7 observed and modelled climate and climate change information. A popular example is the series of
8 reference regions used in the Intergovernmental Panel on Climate Change (IPCC) Special Report on
9 Managing the Risks of Extreme Events and Disasters to Advance Climate Adaptation (SREX). The
10 SREX regions were slightly modified for the 5th Assessment Report of the IPCC and used for reporting
11 sub-continental observed and projected changes over a reduced number (33) of climatologically
12 consistent regions encompassing a representative number of grid boxes. These regions are intended to
13 allow analysis of atmospheric data over broad land or ocean regions and have been used as the basis for
14 several popular spatially aggregated datasets, such as the seasonal mean temperature and precipitation in
15 IPCC regions for CMIP5.

Eliminado: set of

Eliminado: introduced

16 We present an updated version of the reference regions for the analysis of new observed and simulated
17 datasets (including CMIP6) which offer an opportunity for refinement due to the higher atmospheric
18 model resolution. As a result, the number of land and ocean regions is increased to 46 and ¹⁵
19 respectively, better representing consistent regional climate features. The paper describes the rationale for
20 the definition of the new regions and analyses their homogeneity. The regions are defined as polygons
21 and are provided as coordinates and shapefile together with companion R and Python notebooks to
22 illustrate their use in practical problems (e.g. calculating regional averages). We also describe the
23 generation of a new dataset with monthly temperature and precipitation, spatially aggregated in the new
24 regions, currently for CMIP5 and CMIP6, to be extended to other datasets in the future (including
25 observations). The use of these reference regions, dataset and code is illustrated through a worked
26 example using scatter plots to offer guidance on the likely range of future climate change at the scale of
27 the reference regions. The regions, datasets and code (R and Python notebooks) are freely available at the
28 ATLAS GitHub repository; <https://github.com/SantanderMetGroup/ATLAS>,
29 <https://doi.org/10.5281/zenodo.3998463> (Iturbide et al., 2020).

Eliminado: 14

30 **KEY WORDS:** *Regional climate change; Climatic regions; CMIP5; CMIP6; Climate change projections;*
31 *Reproducibility*

Eliminado: doi:10.5281/zenodo.3968318

32 *Copyright statement. The reference regions and the aggregated datasets derived from CMIP5 described in this paper*
33 *are made available in the ATLAS GitHub repository (<https://github.com/SantanderMetGroup/ATLAS>) under the*
34 *Creative Commons Attribution (CC-BY) 4.0 license, whereas the scripts and code are made available under the GNU*
35 *General Public License (GPL) v3.0. Additional products included in the ATLAS GitHub comply with the licenses of*
36 *the original datasets and are periodically updated (e.g. CMIP6 aggregated dataset).*

37 1 Introduction

38 Different sets of climate reference regions have been proposed in the literature for the regional synthesis of historical
39 trends and future climate change projections, and have been subsequently used in the different Assessment Reports of
40 the IPCC (we refer to these sets as IPCC WGI reference regions). The Giorgi reference regions (originally 23
41 rectangular regions proposed in Giorgi and Francisco, 2000; denoted here as version 1) were used in the third (AR3,
42 Giorgi et al., 2001) and fourth (AR4, Christensen et al., 2007) IPCC Assessment reports. These regions were
43 modified using more flexible polygons in the IPCC Special Report on Managing the Risks of Extreme Events and
44 Disasters to Advance Climate Adaptation (SREX, Seneviratne et al., 2012; version 2) and then slightly modified and
45 extended to 33 regions (by including island states, the Arctic and Antarctica) for the fifth Assessment Report (AR5,
46 van Oldenborgh et al., 2013; version 3), as shown in Figure 1a. The objective in these revisions was to improve the
47 climatic consistency of the regions so they represent sub-continental areas of greater climatic coherency. This process
48 typically resulted in a higher number of smaller regions, constrained by the relatively coarse resolution of the global
49 models, since each region should encompass a sufficient number of gridboxes. The AR5 reference regions

Eliminado: observed and model-projected climate change information...

Con formato: Fuente: Cursiva

Con formato: Fuente: Cursiva

Con formato: Fuente: Cursiva

Eliminado: squared

Con formato: Fuente: Cursiva

Eliminado: IPCC SREX special report (

Con formato: Fuente: Cursiva

Con formato: Fuente: Cursiva

Eliminado: larger

Eliminado: IPCC

Con formato: Número de página

1 (http://www.ipcc-data.org/guidelines/pages/ar5_regions.html; last access: 30 July 2020) were developed for reporting
2 sub-continental CMIP5 projections (with an average horizontal resolution greater than 2°) and were quickly adopted
3 by the research community as a basis for regional analysis in a variety of applications (Barring and Strandberg, 2018;
4 Madakumbura et al., 2019). Moreover, these regions have been used to generate popular spatially aggregated
5 datasets, such as the *seasonal mean temperature and precipitation in IPCC regions for CMIP5* (McSweeney et al.,
6 2015), which provides ready-to-use information from the CMIP5 models, suitable for regional analysis of climate
7 projections and their uncertainties. This dataset can be directly used by researchers and stakeholders for a variety of
8 purposes, including assessing the internal variability, model and scenario uncertainty components (Hawkins and
9 Sutton, 2009), or assisting in the comparison and selection of representative sub-ensembles for impact studies (e.g.,
10 Ruane and McDermaid, 2017).

11 The increasing availability of CMIP6 multi-model simulations (O'Neill et al., 2016; NCC editorial, 2019) offers an
12 opportunity to refine the AR5 reference regions —due to the higher atmospheric model resolution, typically around
13 1°— and also to produce ready-to-use aggregated regional information for the updated reference regions. This is a
14 timely task due to the great interest of the research community in the higher sensitivity of some CMIP6 models and
15 the potential implications for climate change studies (Forster et al., 2020). Here, we present the results of an initiative
16 carried out during the last year to achieve this goal. First, we present the updated regions (referred to as *IPCC WGI*
17 *reference regions, version 4*) and describe the rationale for the revision, which was guided by two basic principles: 1)
18 climatic consistency and better representation of regional climate features and 2) representativeness of model results
19 (sufficient number of model gridboxes per region). Climatic homogeneity is characterized in terms of mean
20 temperature and precipitation considering Köppen-Geiger climatic regions (Rudel and Kottek, 2010), the annual
21 cycle and projected changes over the reference regions. The resulting 46 land plus 15 ocean regions (see Figure 1b)
22 are provided as coordinates (in csv format) and also as a shapefile, with companion notebooks to illustrate their use in
23 R and Python.

24 Second, we describe the monthly regional temperature and precipitation dataset obtained by spatially aggregating the
25 model data over the reference regions (currently for CMIP5 and CMIP6, to be extended later to observations and
26 additional datasets). Finally, the use of these reference regions, datasets and code is illustrated through a reproducible
27 example which analyses the likely range of future temperature and precipitation changes that are expected for
28 different European regions using scatter plots.

29 Section 2 presents the data and methods used in this work. Section 3 describes the reference regions and their
30 rationale. The regionally aggregated CMIP5 dataset is presented in Section 4 and links are provided for additional
31 aggregated datasets (e.g. CMIP6, which are periodically updated); a reproducible illustrative example is described in
32 Section 5. Finally, conclusions and discussion are presented in Section 6.

33 2 Data and Methods

34 We use global gridded observations to characterize the regional climatological conditions at a sub-continental scale.
35 In particular, we use CRU TS (version 4.03; Harris et al., 2014; Harris and Jones, 2020) providing monthly
36 precipitation and temperature with a resolution of 0.5° over land for the period 1901-2017. Figure 2a-b shows the
37 annual mean temperature and precipitation climatology for the period 1981-2010. CRU TS does not cover Antarctica,
38 which is therefore infilled with an alternative dataset, namely the EWEMBI gridded observations (Lange, 2019).
39 Figure 2c shows the Köppen-Geiger climatic regions (Rudel and Kottek, 2010) computed from these datasets.
40 Quantifying the observational uncertainty is an increasing concern in climate studies, particularly for precipitation
41 (Kotlarski et al., 2019). Therefore, we use two additional observational datasets for precipitation in some parts of this
42 study: 1) Global Precipitation Climatology Centre (GPCC, v2018; Schneider et al., 2011) providing monthly land
43 precipitation values with 0.5° resolution for the period from 1891 to 2016, and 2) Global Precipitation Climatology
44 Project (GPCP; Monthly Version 2.3 gridded, merged satellite/gauge precipitation; Huffman et al., 2009), providing
45 monthly land and ocean precipitation values with a resolution of 2.5° for the period 1979-2018. We show results for
46 the current WMO climatological standard normal period 1981–2010 (WMO, 2017).

47 Global model scenario data were downloaded for CMIP5 (Taylor et al., 2012)/CMIP6 (O'Neill et al., 2016) models
48 for the historical (1850- 2005/1850-2014) and RCP2.6/SSP1-2.6, RCP4.5/SSP2-4.5 and RCP8.5/SSP5-8.5 future
49 scenarios (2006-2100/2015-2100). Data for CMIP5 (curated version used for IPCC-AR5) was downloaded from the
50 IPCC Data Distribution Center (https://www.ipcc-data.org/sim/gcm_monthly/AR5/index.html; last accessed, 31 Dec
51 2019) and for CMIP6 was downloaded from the Earth System Grid Federation (ESGF, Balaji et al., 2018); a
52 periodically updated inventory is available at the ATLAS GitHub repository (in the *AtlasHub-inventory* folder). All
53 model data have been interpolated to common 2° (for CMIP5) and 1° (CMIP6) grids —separately for land and ocean
54 gridboxes using conservative remapping (using CDO with the models and target land/sea masks; CDO, 2019),—

Eliminado: updated

Eliminado: of

Eliminado: 14

Eliminado: as

Eliminado:)

Eliminado: .

Eliminado: .

Eliminado: .

Eliminado: used here

Eliminado: was

Eliminado: s

Con formato: Número de página

1 which are typical model resolutions for CMIP5 and CMIP6 models, respectively. The common grids and land/sea
2 masks are available in the ATLAS GitHub repository (in the *reference-grids* folder).

3 In this paper we illustrate the results using the curated CMIP5 dataset and refer to the ATLAS GitHub repository for
4 similar results for CMIP6. Figure 2d-e shows the CMIP5 multi-model climate change signal for annual mean
5 temperature (in absolute terms) and precipitation (relative, in %) for RCP8.5 2081-2100 (w.r.t. the modern climate
6 baseline 1986-2005 used in AR5). This figure shows the typical spatial climate change patterns and is used to
7 illustrate the consistency of the regional signals in the climate reference regions.

8 3 Reference Regions: Rationale and Definition

9 The *Giorgi* reference regions were originally defined with the goal to represent consistent climatic regimes and
10 physiographic settings, while maintaining an appropriate size for model representation (thousands of kilometers, to
11 contain several model gridboxes), using some subjectivity in the final selection (Giorgi and Francisco, 2000). Here,
12 we are guided by the same basic principles to define a new version of the reference regions (see Figure 1b). Climatic
13 homogeneity is characterized in terms of mean temperature and precipitation considering Köppen-Geiger climatic
14 regions (see Figure 2) and also the annual precipitation cycle (Figures 3 and 4); in the latter case, observational
15 uncertainty is analysed using the three alternative datasets described in Sec. 2. Representativity of model results
16 (sufficient number of gridboxes per region) is analysed at the end of this section in Figure 5.

17 3.1 Definition of new regions

18 Here we describe the rationale for the new version of the reference regions presented in this paper (version 4, see
19 Figure 1b) which is based on the latest available version (version 3, see Figure 1a) that was used in AR5. In contrast
20 to the AR5 regions, the new version includes 16 oceanic regions suitable for the analysis of large-scale atmospheric
21 data. Many of the new land regions are defined by splitting the previous ones to increase climatic homogeneity as
22 described below.

23 In North America, the AR5 Polar Greenland-Iceland (GIC) region is divided in two, Northeastern North America
24 (NEN) and Greenland/Iceland (GIC), to better accommodate the subarctic and Polar climates, respectively (Figure
25 2c). The eastern and central Northern America regions (ENA and CNA) are maintained mostly unaltered while the
26 western part is reorganized to increase climate consistency. The new Northwestern region (NWN) includes mostly
27 the subarctic regions, whereas the modified western region (WNA) encompasses a variety of regional intermixed
28 climates (semiarid, Mediterranean, and continental; see Figure 2c) which are difficult to further separate due to the
29 complex orography.

30 The new North Central America (NCA) region includes the semiarid and arid climates of Northern Mexico,
31 separating them from the tropical climates in southern Central America which constitute a new region (SCA). The
32 Caribbean (CAR) region has been modified to fully include the Greater Antilles.

33 In South America, the AR5 northwestern Amazonia region is divided into three subregions to separate the Northern
34 South America (NSA) region from the western region (NWS) —which includes the northern Andes Mountains range,
35 — and the South America Monsoon (SAM) region. These regions represent sub-continental areas of greater climatic
36 coherency (Espinoza et al 2019), both in terms of climate and climate change signals (Figures 2c-e), and exhibit
37 characteristic seasonal precipitation cycles (Figure 3), with a rainy season from October to March in SAM and no
38 clear wet and dry seasons for NSA and NWS. The Northeastern region is maintained, but the name is changed to
39 Northeastern South America (NES). The old southern South America region is divided in two, separating the
40 northern (southeastern South America, SES) and southern (SAS) parts, the later encompassing the mostly cold desert
41 climates exhibited in this region (see Figure 2c).

42 The three European reference regions NEU, CEU (renamed Western and Central Europe, WCE) and MED have been
43 maintained unaltered since they encompass the main regional climates in Europe, from subarctic, to
44 oceanic/continental and to Mediterranean. However, an additional region has been introduced in Eastern Europe
45 (EEU), encompassing the continental climate on the western side of the Ural mountain range.

46 For Africa, the AR5 WAF region has been divided in two (WAF and CAF); although these regions have similar
47 Köppen-Geiger climates (see Figure 2c), they have very different annual cycles (Figure 4) and therefore should be
48 analysed independently (Diedhiou et al., 2018). A similar situation was found in the original EAF (Osima et al.,
49 2018) which was also divided in two, a Northern subregion (NEAF) which includes the arid region of the Horn of
50 Africa, and a Southern subregion (SEAF). These two regions also exhibit different precipitation seasonal cycles, with

Con formato: Fuente: Cursiva

Eliminado: the

Eliminado: revised

Eliminado: was

Eliminado: was

Eliminado: , and the

Eliminado: old

Eliminado: SAS

Eliminado: old

Eliminado: in divided

Eliminado: , see Figure 2b

Con formato: Número de página

1 different timing of the annual maximum (see Figure 4). Moreover, the South Africa region SAF was also divided in
2 subregions with different rainfall regimes (Maure et al., 2018), the western subregion (WSAF) including the arid
3 regional climates, and the Eastern region (ESAF). Additionally, the (sub)tropical region of Madagascar (MDG) was
4 split from the continent.

Eliminado: two

Eliminado: ted

5 In the case of Asia, Northern Asia is subdivided in a Northern subarctic region (RAR), two regions for Western
6 (WSB) and Eastern (ESB) Siberia and a region for the Russian far East (RFE). The original Western Asia region
7 (WAS) is divided in two regions, Western central Asia (WCA) and the Arabian Peninsula (ARP), the later with an
8 arid climate; these two sub-regions exhibit a distinct seasonal cycle (see Figure 4). The old Tibetan plateau (TIB)
9 region is divided in two subregions, separating the highland climate of the Tibetan plateau in the South (TIB) from
10 the northern arid subregion (Eastern Central Asia, ECA). The South Asia (SAS), East Asia (EAS) and Southeast Asia
11 (SEA) are maintained unaltered, with the exception of adjustments caused by changes in neighboring regions and the
12 definition of two ocean regions (the Arabian Sea and the Bay of Bengal) for the oceanic part of the original SAS.

Eliminado: (

Eliminado: s

Eliminado: b

Eliminado:)

Eliminado: (

Eliminado: ,

Eliminado: regions

Eliminado: 14

Eliminado: and

Eliminado: and

Eliminado: considering

Eliminado: ****RATIONALE FOR THE OCEAN
REGIONS***...

13 Regarding Australasia, the Southern region (SAU) is now further south, better differentiating the rainfall climatology,
14 (Fig. 2c) and separated from the oceanic New Zealand (NZ). The Northern region is divided in three subregions to
15 increase climatic consistency (CSIRO and Bureau of Meteorology, 2015; see Figure 1b) separating the northern
16 tropical region (NAU), the central arid region (CAU) and the subtropical east coast (EAU).

17 In contrast to the version 3 reference regions used in AR5, those defined in this paper also include 15 oceanic regions
18 (note that the Caribbean, the Mediterranean and South East Asia are considered both land and ocean regions, defined
19 using the land and sea masks, respectively). In version 3 only selected sub-domains of the Indian and tropical Pacific
20 ocean were designated as reference regions and the rest of the main oceanic regions were not represented. Version 4
21 includes representation of all major oceanic regions. The equatorial and northern and southern extents of each of the
22 main non-polar oceans are defined as separate regions with the added refinement of dividing the "northern Indian
23 ocean" into two, the Arabian Sea and the Bay of Bengal. The Arctic Ocean is defined as the region north of the main
24 Eurasian and North American landmass which then also defines the northern extent of the North Pacific and Atlantic
25 regions. The equatorial regions extend from 10S to 10N to include those regions used to define indices for both El
26 Nino and the Indian Ocean Dipole. The southern extent of the South Pacific, Atlantic and Indian regions are similar
27 to those defined by Durack and Wijffels (2010) with the remaining ocean region to the south defined as a single
28 Southern Oceans region.

29 Since these ocean regions largely exclude the coastal zones (which are often included in the "land" regions), they are
30 generally more suitable for the analysis of large-scale atmospheric data. Figures 2d and 2e demonstrate that in this
31 respect the ocean regions are a good addition to the AR5 definitions even though they were not developed with the
32 intention of defining ocean basin masks for zonal means used by oceanographers. However, we note that since the
33 coastal regions can be defined by applying a land-sea mask to the land boxes, it is possible to combine regions to
34 enable the more traditional ocean basin definitions used by oceanographers to be produced to a large extent (albeit
35 not exactly).

Eliminado: s

36 3.2 Representativeness of model results

37 The higher atmospheric resolution of CMIP6 yields better model representation on the reference regions (more
38 gridboxes per region) allowing a revision for better climatic consistency (e.g. dividing heterogeneous regions) while
39 preserving model representativeness. Figure 5 illustrates this, displaying the number of gridboxes (only land
40 gridboxes for land regions) in each of the AR5 (last column) and revised (first column) reference regions for the two
41 reference grids (1° and 2°), as well as for the CMIP6 model grids (representing the multi-model mean of gridbox
42 numbers). This figure shows that the 1° grid provides a good reference for CMIP6. Moreover, it shows that the new
43 reference regions are more representative than the AR5 ones due to the increase of model resolution (see Figures 5a
44 and 5d, corresponding to the cases of CMIP6 data on the updated reference regions, and to CMIP5 data in the original
45 AR5 regions, respectively). The regions with the smallest number of gridboxes correspond to three island regions:
46 The Caribbean (CAR), New Zealand (NZ), and Madagascar (MDG), with around 20-60 gridboxes per region. Note
47 that the updated regions are also suitable for the analysis of CMIP5 data (at 2° resolution, Fig. 5c) since all regions
48 encompass over ten land gridboxes, with the exception of the three above-mentioned regions, where results should be
49 interpreted with caution.

50 These updated regions are defined as polygons (the lines in Figure 1 are straight lines on a projected plane) and are
51 provided as coordinates and shapefile at the ATLAS GitHub (*reference-regions* folder); the reference grids and land-
52 sea masks can be found at the *reference-grids* folder. Moreover, companion R and Python notebooks are also

Con formato: Número de página

1 available (*reference-regions/notebooks*) to illustrate their use in practical problems (e.g. calculating regional
2 averages).

3 4 Regionally Aggregated CMIP Datasets

4 The seasonal mean temperature and precipitation in CMIP5 models averaged over the version 3 (AR5) reference
5 regions is a popular dataset, suitable for the regional analysis of climate projections and their uncertainties
6 (McSweeney et al., 2015). Here we extend this idea to the new regions and model data and compute aggregated
7 monthly results over the different reference regions (see Figure 1b) for all the CMIP5 simulations (and also the
8 available CMIP6 ones), considering land only, sea only, and land-sea gridboxes (the land/sea masks are available in
9 the ATLAS GitHub repository, *reference-grids*). Results are calculated for each model simulation and stored
10 individually as a text-csv file, with regions in columns (including the global results in the last column) and dates
11 (months) in rows; results for a one ensemble member per model are included directly in the ATLAS Github
12 repository (*aggregated-datasets* folder), and links are provided to the general dataset (full ensemble with all runs)
13 which allows for internal variability studies.

14 Whereas the aggregated CMIP5 dataset is final, results for CMIP6 will be regularly updated when new data become
15 available at ESGF; these two datasets constitute alternative lines of evidence for climate change studies and the
16 ATLAS initiative presented here facilitates intercomparison of results and consistency checks for the reference
17 climatic regions. Note that although the aggregated data provides summary climate information for each sub-
18 continental region which is useful for a broad spectrum of users, detailed climate information at local or regional
19 scales (in each sub-continental region) would be required for further regional analysis.

20 5 Illustrative Case Study

21 To demonstrate a potential application of the reference regions and the associated regionally-averaged CMIP data
22 (for temperature and precipitation), we show a simple case study illustrating the projected range of future
23 temperature/precipitation change. This can provide useful context information for a variety of impact and adaptation
24 studies. In particular, we use scatter plots to show the median, 10th, and 90th percentiles of the CMIP5 ensemble
25 change. We focus on three illustrative European regions (NEU, WCE and MED) with opposite climate change
26 signals for precipitation (see Figure 2e). The code and data needed to run this example (which can be extended to
27 other regions, or combination of regions, and datasets, e.g. CMIP6) are all available at the ATLAS Github repository
28 (*aggregated-datasets/scripts* folder) and can be run in a local R session accessing the Github data with no further
29 requirements.

30 Figure 6 shows the projected changes in annual mean temperature and precipitation resulting from the script
31 *scatterplots_TvsP.R*. In particular, results from RCP2.6, RCP4.5 and RCP8.5 scenarios for early (2021-2040), mid
32 (2041-2060 and 2061-2080) and late (2081-2100) 21st century — relative to the 1986-2005 baseline period — for
33 each of the three European subregions are displayed. This figure projects an increase of temperature in all European
34 domains —with similar warming in all regions for the different scenarios and future periods— and a consistent
35 meridional gradient of changes in precipitation, with a clear precipitation increase in NEU, non-changing conditions
36 in WCE (uncertainty range crossing the zero line), and reduced precipitation over MED. The same scripts can be
37 applied to the currently available CMIP6 dataset by changing two parameters to check the consistency of these results
38 for the updated models and scenarios.

39 Note that this illustrative example can be modified to serve different purposes. For instance, the same diagram can be
40 adapted to display the individual model values (or to select the subset of models spanning the uncertainty range) in
41 order to assist in the comparison and the selection of representative sub-ensembles for impact studies (e.g. Ruane and
42 McDermid, 2017). The calculation of the regional aggregated values is time consuming (computed offline and results
43 are provided in the Github repository); however, accessing the values and plotting the results is straightforward and
44 the scripts provided run in a few seconds.

45 6 Conclusions and Discussion

46 A new set of 46 land plus 15 ocean regions is introduced in this work updating the previous set of IPCC AR5-WGI
47 reference regions for the regional synthesis of model-projected climate change information (in particular for the new
48 CMIP6 simulations). The new regions increase the climatic consistency of the previous ones —by rearranging and
49 dividing regions exhibiting mixed regional climates— and have a suitable model representation (the minimum is in
50 the range 20-60 model gridboxes for three particular island regions: the Caribbean, New Zealand and Madagascar.

Eliminado: in IPCC regions for

Eliminado: McSweeney et al. (2015) is a popular dataset based on the IPCC AR5

Eliminado: ed

Eliminado: d

Eliminado: model runs

Eliminado: run

Eliminado: single run

Eliminado: s

Eliminado: allows

Eliminado: evidences

Eliminado: 14

Con formato: Número de página

1 This revision was guided by the basic principles of climatic consistency and model representativeness, but there is of
2 course some subjectivity in the final selection.

3 We also present a new dataset of monthly CMIP5/6 spatially aggregated information using the new reference regions
4 and the available CMIP5 (from the IPCC-DDC) and CMIP6 data (from ESGF, as of 30 September 2019), and
5 describe a worked example on how to use this dataset to inform regional climate change studies, in particular about
6 the likely range of future temperature/precipitation changes for the different European reference regions using scatter
7 plots.

Eliminado: -out

8 7 Code and data availability

9 The present work is part of the climate change ATLAS initiative (which is aligned with IPCC AR6 activities). The
10 definition of the regions, the code and the associated spatially-aggregated datasets are available at the GitHub
11 ATLAS repository: <https://github.com/SantanderMetGroup/ATLAS>, <https://doi.org/10.5281/zenodo.3998463>,
12 (Iturbide et al., 2020). The ATLAS project builds in the publicly available climate4R R framework (Iturbide et al.,
13 2019) (available under the GNU General Public License v3.0) and provides additional functions which may be
14 relevant for the users of the reference regions and aggregated datasets, such as the calculation of global warming
15 levels, thus enhancing the functionalities presented in this work. The python notebook is based on the *regionmask*
16 (Hauser, 2019) and xarray (Hoyer and Hamman, 2017) packages, among others. The results for CMIP5 are based on
17 the final curated dataset used for IPCC-AR5, but other datasets will be updated periodically when new data becomes
18 available (e.g. CMIP6, still in progress).

Eliminado: doi:10.5281/zenodo.3968318

Eliminado: open source projects

19 Regarding the original datasets used in this work, all of them are publicly available from the local providers — CRU
20 TS4.03 is distributed under the Open Database License, and EWEMBI and GPCCv2018 are distributed under the
21 Creative Commons Attribution 4.0 International License — and/or the Earth System Grid Federation (ESGF, Balaji
22 et al., 2018) — CMIP5 and CMIP6. Moreover, for the sake of reproducibility some datasets have been also
23 replicated at the Santander Climate Data Service which is transparently accessible from climate4R via the User Data
24 Gateway (registration is required to accept the terms of use of the original datasets; more information at
25 <http://meteo.unican.es/udg-wiki>, last access: 30 July 2020).

26 **Author contributions.** Gutiérrez J.M. and Iturbide M. conceived the study and wrote the code and the manuscript;
27 vandenHurk B. conceived the case study; Iturbide M. and Hauser M. implemented the R and Python companion
28 notebooks; all authors contributed to the definition of the regions, to the discussion and revised the text and the
29 results.

30 **Competing interests.** The authors declare that there is not any competing interest.

31

32

33

34

Acknowledgements

35 We acknowledge the World Climate Research Program's Working Group on Coupled Modelling, which
36 is responsible for CMIP, and we thank the climate modeling groups (listed in the Atlas GitHub) for
37 producing and making available their model output.

38 JMG and SH acknowledge support from the Spanish Government through the research and innovation
39 programme (project ref. PID2019-111481RB-I00) and the María de Maeztu excellence programme (Ref.
40 MdM-2017-0765). SHF acknowledges support from the Spanish Government through the María de
41 Maeztu excellence programme (Ref. MDM-2017-0714), and by the Basque Government through the
42 BERC 2018–2021 programme.

43 [The authors are also grateful to the two reviewers \(Chris Brierley and one anonymous\) who helped to](#)
44 [improve the original manuscript with a detailed in deep revision providing constructive comments, and to](#)
45 [Michael Grose and Jason Evans who provided useful comments during the interactive discussion.](#)

Eliminado: The authors are also grateful to anonymous reviewers who helped to improve the original manuscript.

46

47

REFERENCES

48

49

50

51

Balaji, V., Taylor, K. E., Juckes, M., Lawrence, B. N., Durack, P. J., Lautenschlager, M., Blanton, C.,
Cinquini, L., Denvil, S., Elkington, M., Guglielmo, F., Guilyardi, E., Hassell, D., Kharin, S.,
Kindermann, S., Nikonov, S., Radhakrishnan, A., Stockhouse, M., Weigel, T., and Williams, D.:

Con formato: Número de página

1 Requirements for a global data infrastructure in support of CMIP6, Geoscientific Model
2 Development, 11, 3659–3680, <https://doi.org/https://doi.org/10.5194/gmd-11-3659-2018>, 2018.

3 Barring, L. and Strandberg, G.: Does the projected pathway to global warming targets matter?,
4 Environmental Research Letters, 13, 024 029, <https://doi.org/10.1088/1748-9326/aa9f72>, publisher:
5 IOP Publishing, 2018.

6 CDO: Climate Data Operator Version 1.9.8. Max-Planck-Institute for Meteorology,
7 <https://code.mpimet.mpg.de/projects/cdo>, 2019.

8 CSIRO and Bureau of Meteorology 2015, Climate Change in Australia Information for Australia’s Natural
9 Resource Management Regions: Technical Report, CSIRO and Bureau of Meteorology, Australia.
10 <http://www.climatechangeinaustralia.gov.au>

11 Christensen, J., Hewitson, B., Busuioic, A., Chen, A., Gao, X., Held, I., Jones, R., Kolli, R., Kwon, W.-T.,
12 Laprise, R., Rueda, V. M., Mearns, L., Menez, C., Rn, J., Rinke, A., Sarr, A., and Whetton, P.:
13 Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth
14 Assessment Report of the Intergovernmental Panel on Climate Change [Solomon, S., D. Qin, M.
15 Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)], book section
16 Regional Climate Projections, pp. 847–940, Cambridge University Press, Cambridge, United
17 Kingdom and New York, NY, USA, 2007.

18 Diedhiou, A., Bichet, A., Wartenburger, R., Seneviratne, S. I., Rowell, D. P., Sylla, M. B., Diallo, I.,
19 Todzo, S., Touré, N. E., Camara, M., Ngatchah, B. N., Kane, N. A., Tall, L., and Affholder, F.:
20 Changes in climate extremes over West and Central Africa at 1.5°C and 2°C global warming,
21 Environmental Research Letters, 13, 065 020, <https://doi.org/10.1088/1748-9326/aac3e5>, 2018.

22 [Durack, P. J., and S. E. Wijffels, 2010: Fifty-Year Trends in Global Ocean Salinities and Their
23 Relationship to Broad-Scale Warming. J. Climate, 23, 4342–4362,
24 <https://doi.org/10.1175/2010JCLI3377.1>.](https://doi.org/10.1175/2010JCLI3377.1)

25 Espinoza, J.C., Ronchail, J., Marengo, J.A. et al. Contrasting North–South changes in Amazon wet-day
26 and dry-day frequency and related atmospheric features (1981–2017). Clim Dyn 52, 5413–5430,
27 2019. <https://doi.org/10.1007/s00382-018-4462-2>

28 Forster, P. M., Maycock, A. C., McKenna, C. M., and Smith, C. J.: Latest climate models confirm need
29 for urgent mitigation, Nature Climate Change, 10, 7–10, <https://doi.org/10.1038/s41558-019-0660-0>, 2020.

30

31 Giorgi, F. and Francisco, R.: Uncertainties in regional climate change prediction: a regional analysis of
32 ensemble simulations with the HADCM2 coupled AOGCM, Climate Dynamics, 16, 169–182,
33 <https://doi.org/10.1007/PL00013733>, 2000.

34 Giorgi, F., Hewitson, B., Christensen, J., Hulme, M., Storch, H. V., Whetton, P., Jones, R., Mearns, L.,
35 and Fu, C.: Climate Change 2001: The Scientific Basis [J. T. Houghton, Y. Ding, D. J. Griggs, M.
36 Noguera, P. J. van der Linden, X. Dai, K. Maskell, C. A. Johnson (eds.)], book section Regional
37 Climate Information?Evaluation and Projections, pp. 583–638, Cambridge University Press, 2001.

38 Harris, I. and Jones, P.: CRU TS4.03: Climatic Research Unit (CRU) Time-Series (TS) version 4.03 of
39 high-resolution gridded data of month-by-month variation in climate (Jan. 1901- Dec. 2018), Centre
40 for Environmental Data Analysis,
41 <https://doi.org/http://dx.doi.org/10.5285/10d3e3640f004c578403419aac167d82>, 2020.

42 Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H.: Updated high-resolution grids of monthly climatic
43 observations – the CRU TS3.10 Dataset, International Journal of Climatology, 34, 623–642,
44 <https://doi.org/10.1002/joc.3711>, 2014.

45 Hawkins, E. and Sutton, R.: The Potential to Narrow Uncertainty in Regional Climate Predictions, Bulletin
46 of the American Meteorological Society, 90, 1095–1108,
47 <https://doi.org/10.1175/2009BAMS2607.1>, 2009.

48 Hoyer, S. and Hamman, J., 2017. xarray: N-D labeled Arrays and Datasets in Python. Journal of Open
49 Research Software, 5(1), p.10. DOI: doi:10.5334/jors.148.

50 Huffman, G. J., Adler, R. F., Bolvin, D. T., and Gu, G.: Improving the global precipitation record: GPCP
51 Version 2.1, Geophysical Research Letters, 36, <https://doi.org/10.1029/2009GL040000>, 2009.

52 Iturbide, M., Bedia, J., Herrera, S., Baño-Medina, J., Fernández, J., Frías, M. D., Manzanar, R., San-
53 Martín, D., Cimadevilla, E., Cofiño, A. S., and Gutiérrez, J. M.: The R-based climate4R open

1 framework for reproducible climate data access and post-processing, *Environmental Modelling &*
2 *Software*, 111, 42–54, <https://doi.org/10.1016/j.envsoft.2018.09.009>, 2019.

3 Iturbide, M., Gutiérrez, J. M., Cimadevilla, E., Bedia, J., Hauser, M., and Manzanar, R.:
4 SantanderMetGroup/ATLAS GitHub (Version v1.6), <https://doi.org/10.5281/zenodo.3998463>,
5 2020.

6 Kotlarski, S., Szabó, P., Herrera, S., Ráty, O., Keuler, K., Soares, P. M., Cardoso, R. M., Bosshard, T.,
7 Pagé, C., Boberg, F., Gutiérrez, J. M., Isotta, F. A., Jaczewski, A., Kreienkamp, F., Liniger, M. A.,
8 Lussana, C., and Pianko-Kluczynska, K.: Observational uncertainty and regional climate model
9 evaluation: A pan-European perspective, *International Journal of Climatology*, 39, 3730–3749,
10 <https://doi.org/10.1002/joc.5249>, 2019.

11 Lange, S.: Earth2Observe, WFDEI and ERA-Interim data Merged and Bias-corrected for ISIMIP
12 (EWEMBI). V. 1.1. GFZ Data Services., <http://doi.org/10.5880/pik.2019.004>, 2019.

13 Madakumbura, G. D., Kim, H., Utsumi, N., Shioyama, H., Fischer, E. M., Seland, A., Scinocca, J. F.,
14 Mitchell, D. M., Hirabayashi, Y., and Oki, T.: Event-to-event intensification of the hydrologic cycle
15 from 1.5°C to a 2°C warmer world, *Scientific Reports*, 9, 1–7, <https://doi.org/10.1038/s41598-019-39936-2>, 2019.

16 Maure, G., Pinto, I., Ndebele-Murisa, M., Muthige, M., Lennard, C., Nikulin, G., Dosio, A., and Meque,
17 A.: The southern African climate under 1.5°C and 2°C of global warming as simulated by CORDEX
18 regional climate models, *Environmental Research Letters*, 13, 065 002,
19 <https://doi.org/10.1088/1748-9326/aab190>, 2018.

20 McSweeney, C. F., Jones, R. G., Lee, R. W., and Rowell, D. P.: Selecting CMIP5 GCMs for downscaling
21 over multiple regions, *Climate Dynamics*, 44, 3237–3260, <https://doi.org/10.1007/s00382-014-2418-8>, 2015.

22 NCC editorial: The CMIP6 landscape, *Nature Climate Change*, 9, 727–727,
23 <https://doi.org/10.1038/s41558-019-0599-1>, 2019.

24 O’Neill, B. C., Tebaldi, C., Vuuren, D. P. v., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler,
25 E., Lamarque, J.-F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario
26 Model Intercomparison Project (ScenarioMIP) for CMIP6, *Geoscientific
27 Model Development*, 9, 3461–3482, <https://doi.org/10.5194/gmd-9-3461-2016>, 2016.

28 Osima, S., Indasi, V. S., Zaroug, M., Endris, H. S., Gudoshava, M., Misiani, H. O., Nimusiima, A., Anyah,
29 R. O., Otieno, G., Ogwang, B. A., Jain, S., Kondowe, A. L., Mwangi, E., Lennard, C., Nikulin, G.,
30 and Dosio, A.: Projected climate over the Greater Horn of Africa under 1.5°C and 2°C global
31 warming, *Environmental Research Letters*, 13, 065 004, <https://doi.org/10.1088/1748-9326/aaba1b>,
32 2018.

33 Ruane, A. C. and McDermid, S. P.: Selection of a representative subset of global climate models that
34 captures the profile of regional changes for integrated climate impacts assessment, *Earth
35 Perspectives*, 4, 1, <https://doi.org/10.1186/s40322-017-0036-4>, 2017.

36 Rubel, F. and Kottek, M.: Observed and projected climate shifts 1901-2100 depicted by world maps of
37 the Köppen-Geiger climate classification, *Meteorologische Zeitschrift*, pp. 135–141,
38 <https://doi.org/10.1127/0941-2948/2010/0430>, 2010.

39 Schneider, U., Becker, A., Finger, P., Meyer-Christoffer, A., Rudolf, B., and Ziese, M.: GPCC Full Data
40 Reanalysis Version 6.0 at 0.5: Monthly Land-Surface Precipitation from Rain-Gauges built on
41 GTS-based and Historic Data., https://doi.org/10.5676/DWD_GPCC/FD_M_V7_050, 2011.

42 Seneviratne, S., Nicholls, N., Easterling, D., Goodess, C., Kanae, S., Kossin, J., Luo, Y., Marengo, J.,
43 McInnes, K., Rahimi, M., Reichstein, M., Sorteberg, A., Vera, C., and Zhang, X.: Managing the
44 Risks of Extreme Events and Disasters to Advance Climate Change Adaptation [Field, Barros,
45 Stocker et al. (eds.)], book section Changes in climate extremes and their impacts on the natural
46 physical environment, pp. 109–230, Cambridge University Press, Cambridge, United Kingdom and
47 New York, NY, USA, 2012.

48 Sun, Q., Miao, C., Duan, Q., Ashouri, H., Sorooshian, S., and Hsu, K.-L.: A Review of Global
49 Precipitation Data Sets: Data Sources, Estimation, and Intercomparisons, *Reviews of Geophysics*,
50 56, 79–107, <https://doi.org/10.1002/2017RG000574>, 2018.

Eliminado: 5

Eliminado: <http://doi.org/10.5281/zenodo.3968318>

1 Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An Overview of CMIP5 and the Experiment Design,
2 Bulletin of the American Meteorological Society, 93, 485–498, [https://doi.org/10.1175/BAMS-D-](https://doi.org/10.1175/BAMS-D-11-00094.1)
3 [11-00094.1](https://doi.org/10.1175/BAMS-D-11-00094.1), 2012.

4 van Oldenborgh, G.J., M. C., Arblaster, J., Christensen, J., Marotzke, J., Power, S., Rummukainen, M.,
5 and Zhou, T.: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I
6 to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F.
7 and Qin, D. and Plattner, G.-K. and Tignor, M. and Allen, S.K. and Boschung, J. and Nauels, A. and
8 Xia, Y. and Bex, V. and Midgley, P.M. (eds.)], book section Annex I: Atlas of Global and Regional
9 Climate Projections, pp. 1311–1394, Cambridge University Press, Cambridge, United Kingdom and
10 New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.029>,
11 www.climatechange2013.org, 2013.


12 WMO: WMO Guidelines on the Calculation of Climate Normals. WMO - No. 1203,
13 https://library.wmo.int/doc_num.php?explnum_id=4166, 2017.

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32


1
2

FIGURES

(a) IPCC WGI Reference Regions (v3, AR5)



(b) Updated IPCC WGI Reference Regions (v4)




1	GIC	Greenland/Iceland	23	SAH	Sahara	43	NAU	N. Australia
2	NWN	N.W. North-America	24	WAF	Western-Africa	44	CAU	C. Australia
3	NEN	N.E. North-America	25	CAF	Central-Africa	45	EAU	E. Australia
4	WNA	W. North-America	26	NEAF	N. Eastern-Africa	46	SAU	S. Australia
5	CNA	C. North-America	27	SEAF	S. Eastern-Africa	47	NZ	New-Zealand
6	ENA	E. North-America	28	WSAF	W. Southern-Africa	48	EAN	E. Antarctica
7	NCA	N. Central-America	29	ESAF	E. Southern-Africa	49	WAN	W. Antarctica
8	SCA	S. Central-America	30	MDG	Madagascar	50	ARO	Arctic-Ocean
9-10	CAR	Caribbean	31	RAR	Russian-Arctic	51	NPO	N. Pacific-Ocean
11	NWS	N.W. South-America	32	WSB	W. Siberia	52	EPO	Equatorial Pacific-Ocean
12	NSA	N. South-America	33	ESB	E. Siberia	53	SPO	S. Pacific-Ocean
13	NES	N.E. South-America	34	RFE	Russian-Far-East	54	NAO	N. Atlantic-Ocean
14	SAM	South-American-Monsoon	35	WCA	W. C. Asia	55	EAO	Equatorial Atlantic-Ocean
15	SWS	S.W. South-America	36	ECA	E. C. Asia	56	SAO	S. Atlantic-Ocean
16	SES	S.E. South-America	37	TIB	Tibetan-Plateau	57	ARS	Arabian-Sea
17	SSA	S. South-America	38	EAS	E. Asia	58	BOB	Bay-of-Bengal
18	NEU	N. Europe	39	ARP	Arabian-Peninsula	59	EIO	Equatorial Indian-Ocean
19	WCE	Western&Central-Europe	40	SAS	S. Asia	60	SIO	S. Indian-Ocean
20	EEU	E. Europe	41-42	SEA	S.E. Asia	61	SOO	Southern-Ocean
21-22	MED	Mediterranean						


3
4
5
6
7
8

Figure 1. Updated IPCC reference land (gray shading) and ocean (blue shading) regions; note that the Caribbean and the Mediterranean are considered both land and ocean regions (defined using the land and sea masks, respectively). Land masks are used to obtain land-only information for land regions (excluding the coastal white regions).

(a) IPCC AR5-1



(b) Updated




1	GIC	Greenland/Iceland	21	SAH	Sahara
2	NWN	N.W. North-America	22	WAF	Western-Africa
3	NEN	N.E. North-America	23	CAF	Central-Africa
4	WNA	W. North-America	24	NEAF	N. Eastern-Africa
5	CNA	C. North-America	25	SEAF	S. Eastern-Africa
6	ENA	E. North-America	26	WSAF	W. Southern-Africa
7	NCA	N. Central-America	27	ESAF	E. Southern-Africa
8	SCA	S. Central-America	28	MDG	Madagascar
9	CAR	Caribbean	29	RAR	Russian-Arctic
10	NWS	N.W. South-America	30	WSB	W. Siberia
11	NSA	N. South-America	31	ESB	E. Siberia
12	NES	N.E. South-America	32	RFE	Russian-Far-East
13	SAM	South-American-Monsoon	33	WCA	W. C. Asia
14	SWS	S.W. South-America	34	ECA	E. C. Asia
15	SES	S.E. South-America	35	TIB	Tibetan-Plateau
16	SSA	S. South-America	36	EAS	E. Asia
17	NEU	N. Europe	37	ARP	Arabian-Peninsula
18	WCE	Western&Central-Europe	38	SAS	S. Asia
19	EEU	E. Europe	39	SEA	S.E. Asia
20	MED	Mediterranean			

Eliminado:

Con formato: Centrado


Con formato: Número de página

1



2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23


Figure 2. (a) Global mean temperature, (b) accumulated precipitation and (c) Köppen-Geiger climate classification from the CRU-TS dataset for the period 1981-2010 (data for Antarctica is filled with the EWEMBI dataset). This information is used to characterize the regional climate consistency of the reference regions (solid lines). (d,e) Climate change projections for temperature and precipitation, respectively, from the CMIP5 curated dataset for RCP8.5 2081-2100 w.r.t. the AR5 modern climate baseline 1986-2005. Hatching indicates weak (less than 80%) model agreement on the sign of the change.



1
2
3
4
5
6
7
8
9
10


Figure 3. Observed annual cycle (1981-2010) for precipitation for the American reference regions from three different observational datasets (CRU-TS, GPCC, GPCP) and climate change signal (RCP8.5, 2081-2100 w.r.t. the AR5 modern climate baseline 1986-2005). The panel for each reference region shows the observed annual cycle (top, in monthly accumulated mm) and the monthly projected changes (bottom, in gray, as %); box and whiskers plots represent the spatial (gridbox) spread of monthly values over the region.

1
2
3




5
6
7
8
9
10
11
12

Figure 4. As figure 3 but for Europe, Asia and Australasia.



1
2
3 **Figure 5.** Number of gridboxes encompassed by the different reference regions for 1° (a,b) and
4 2° (c,d) resolution and for the CMIP6 model grids (e,f) –the multi-model mean is represented–,
5 considering the updated (a,c,e) and the original AR5 (b,d,f) reference regions. Colors indicate
6 regions with less than 250 gridboxes. The blue numbers in each of the regions show the number of
7 gridboxes (only land gridboxes for land regions).
8

1
2
3
4



5
6
7
8
9
10
11
12
13
14
15
16
17
18

Figure 6. Illustrative example of the use of reference regions and aggregated CMIP5 datasets: Regional mean changes in annual mean temperature and precipitation for three European regions (NEU, WCE and MED) for four future periods (2021-2040, 2041-2060, 2061- 2080, 2081-2100), as obtained from CMIP5 projections. Changes are absolute for temperature and relative for precipitation. Horizontal and vertical error bars represent ± 1 standard deviation from the mean calculated across the ensemble of included models. The script to generate this figure for all the 61 land and ocean regions (as well as the global results) from the aggregated and ready-to-use CMIP5 datasets is available at the ATLAS GitHub, and can be adapted to produce similar results for alternative datasets (e.g. CMIP6).

Eliminado: 55
Eliminado: including
Eliminado: the global region
Eliminado: aggregated

Con formato: Número de página

▼
▲
Página 1: [1] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:55:00**

▼
▲
Página 1: [1] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:55:00**

▼
▲
Página 1: [1] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:55:00**

▼
▲
Página 1: [1] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:55:00**

▼
▲
Página 1: [1] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:55:00**

▼
▲
Página 1: [1] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:55:00**

▼
▲
Página 1: [1] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:55:00**

▼
▲
Página 1: [1] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:55:00**

▼
▲
Página 1: [1] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:55:00**

▼
▲
Página 1: [2] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:57:00**

▼
▲
Página 1: [2] Eliminado **Fernandez Fernandez, Jesus** **23/8/20 17:57:00**