

We would like to thank both the reviewers for the careful and thorough reading of this manuscript and for the thoughtful comments and constructive suggestions they provided. They undoubtedly helped to improve the quality of this manuscript. In this response we are addressing the comments and feedback from the reviewers. We have numbered the reviewers comments (**R1.1, R1.2** etc. for Reviewer 1 and **R2.1, R2.2** etc. for Reviewer 2) in order to facilitate referencing to each comment. The pages and line numbers in our responses refer to the manuscript with the changes.

Anonymous Referee #1 (R1)

The manuscript by Athanasiou et al. introduces a new global dataset of nearshore slopes with the aim to address an important gap of global-scale coastal analyses. The manuscript is very well structured and the methods are presented in a comprehensive and understandable manner. From a methodological point of view, the work that the authors have carried out is very meticulous. Data processing has been conducted with high attention to detail and limitations of the work are explicitly acknowledged. Importantly, based on the description that is provided, the results are reproducible and the methods can be employed by other researchers when improved data become available.

I recommend that the manuscript is published. I have included below a small number of comments that the authors should address in order to clarify some methodological decisions and accordingly discuss/revise some small parts of the manuscript. I hope that the authors find the comments useful for improving the manuscript.

R1.1 - pg. 1 line28: the coast also comprises different land use/cover types, which also define its response to these events

As suggested by Reviewer 1 we included this missing part in our description:

pg.1 line28: “As the coastline comprises various different landforms (e.g. sandy coasts, rocky cliffs etc.) and land uses (e.g. heavily urbanized or natural), the response to these hazards can vary significantly both spatially and temporally.”

R1.2 - pg. 3 line 23: Based on my experience, the choice of the coastline is crucial for the analysis. According to the smoothing method described, a different coastline dataset (which could diverge considerably from the selected one) would influence the results (see also discussion by Lichter et al., 2010, in the context of exposure analysis). I believe that this is a point that the authors need to discuss (briefly) in the manuscript (there is mention of this in the limitations section, but not really addressing the point)

We did not explicitly test the sensitivity of the elevation profiles to the choice of coastline, but we considered other coastlines in the beginning of the study. The issue we encountered was that other datasets either lacked the required resolution (NOOA coastline) or did not have a global coverage and consistency (various national datasets). That is why we decided to use the OSM coastline. Nevertheless, we agree that this is an important issue that we now mention in the discussion:

Pg.10 line 28: “Although other available coastline datasets exist, either they have a coarser resolution, therefore not allowing for a correct representation of the separation between land and water, or do not have a global coverage. The importance of the coastline and DEM data has been previously highlighted with respect to statistics of exposure information (Lichter et

al., 2011). We expect that the choice of the coastline can have an effect in the elevation profiles, especially in areas with a high tidal range.”

R1.3 - pg. 4 lines 1-4: To my understanding, the smoothing process can considerably affect the calculation of the slope and generally produce much lower values. I am surprised that the authors do not discuss this point in the limitations. Also, doesn't this process essentially "remove" all cliffs? Finally, isn't this the reason why the calculated global slopes are milder than those observed? (see lines 5-8, pg. 8).

We understand the thinking of Reviewer 1 on this topic. With respect to the cliffs we believe that the calculated slopes may indeed be milder than the actual ones, but they can still be rather steep. This can be verified by the global distribution of slopes as well (Figure 5), where a high number of quite steep slopes exist (steeper than 1/2), related to the large length of coastline with cliff features. Since our study focused on non-cliffed coastlines we did not test this further. In relation to the second part of the question, we believe that, since the smoothing is applied in areas close to the shoreline, this will not have a substantial effect on the areas close to the depth of closure, and hence will not affect the estimates of slopes in areas where the slopes are not very steep. For example even in the Dutch coast where no smoothing was needed, there is a negative bias (Table 2). We believe that the negative bias is connected with general differences of the accuracy of the GEBCO bathymetry.

Nevertheless, to address the reviewer comment we have now included in the discussion section the following sentences:

Pg.11 line 5: “The smoothing procedure that was applied to the bathymetric and topographic data may have affected the nearshore slope values, especially at cliffed coastlines, where the computed slopes may be milder than the reality. Nevertheless, cliffed transects retained quite steep values as can be seen by the second peak in the global slope distribution (**Error! Reference source not found.**)”

R1.4 - pg. 4 line 14: I am a little confused with the interval (25m) chosen for the equidistant points along the transects, considering the horizontal resolution of the employed datasets. Please justify.

The choice of a 25m interval results from a number of sensitivity tests to ensure that 1) the cross-shore resolution is captured well and 2) the amount of data saved is manageable.

R1.5 - pg. 4 line16: what do "significant human interventions" include?

We are sorry for the confusion. Here we were mainly focusing on ports, when talking about "significant human interventions". We have now clarified this:

Pg.4 line 23: “...exclude transects located along coasts covered with ice, river inlets, or major ports, using the global transects of Luijendijk et al. (2018).”

R1.6 - pg. 5 line 18: how many of those not calculated were sandy beaches?

Using the same sandy beach detection dataset (Luijendijk et al. 2018), the estimated number of sandy transects for which the slope could not be calculated is ~33,000 of the 160,000 (~20%). We have now included this in the manuscript:

Pg.5 line 28: “The transects of which the nearshore slope could not be calculated were about 160,000 of the total 780,000. Almost 20% out of these 160,000 transects were identified as sandy.”

R1.7 - pg. 8 line 20: is it the wave climate affecting slope or vice versa? Also, swell waves often occur in coasts with mild slopes. I believe that the explanation here is rather weak.

In the long term (geological time scales) the waves can shape a coast and in the shorter term the coastal slopes affect the wave transformation. There are various processes that can also have a major effect in shaping the coastal profile like the sediment supply from rivers, the tectonics, the presence of submarine canyons etc. We have now tried to reformulate our discussion section with respect to these:

Pg.9 Line 13: “At sedimentary coasts the upper nearshore profile is mainly influenced by waves while the lower one is influenced by tectonics, sediment supply or major storm events.”

R1.8 - A general point: to my experience, the calculation of slope in a global application can be affected by the type of projection. Also, re-projecting datasets to different projections can be a complex (or even problematic) task in the context of global applications. I think that the authors would need to provide some additional information (also in the supplementary material if they prefer) regarding those choices and state what type of projections they employed and how slope/length were calculated for the entire world.

Actually, for our study we projected the OSM coastline to the local UTM projection and then created the transects. In this way we ensured that the spacing between the transects is consistent globally and that all the transects have the same (real) length. We have now added two sentences to the Methods sections supporting our choice:

Pg.4 line 13: “To ensure a globally consistent spacing and length of the transects, the coastline was re-projected to the local UTM projection, before creating the transects.”

Pg.4 line 20: “For each transect the longitude and latitude of the extraction points were used to sample from the global merged elevation grid, and create a continuous elevation profile.”

Anonymous Referee #2 (R2)

Is the article appropriate to support the publication of a data set? Yes, indeed. It describes the dataset, how data were compiled and/or produced and it also provides some examples of application.

Is the data set significant – unique, useful, and complete? The described and online provided dataset is significant, useful, worldwide distributed and it will help to fill a frequent commonly found gap in coastal data.

Is the data set itself of high quality? Due to its worldwide distribution, the data set can be considered of good quality. They have been validated against local data and bias and error measurements are provided. Is the data set publication, as submitted, of high quality? The data set publication is adequate to the importance and quality of the data provided. The length to describe the data is appropriated, and the publication itself has the right structure and cover the required aspects to be useful for readers and potential users of the data.

In what follows some specific comments are given.

R2.1 Page 2, line 8. The used definition of the depth-of-closure is not the standard one. The depth-of-closure defined by equation 1 is NOT the offshore limit where sediment transport is zero. The standard definition is “the depth seaward of which there is no significant change in bottom elevation”.

We appreciate the comment by Reviewer 2. We have now changed the definition to:

Pg.2 line 8: “...between the depth of closure d_c (i.e. the depth seaward of which there is no significant change in bottom elevation) and the shoreline (MSL).”

R2.2 Page 2, line 22. As it is written this sentence seems to indicate that in the presented study the depth of closure is going to be calculated in a different way of those mentioned in the previous sentence (empirical formulas using wave parameters).

We can understand that this sentence might send the wrong message to the reader. We have now changed it to:

Pg.2 line 21: “The most common approach to estimate d_c is using empirical formulae (Hallermeier, 1978; Birkemeier, 1985; Nicholls et al., 1998) that relate d_c to wave parameters. This has been employed in several studies both at the global scale (Hinkel et al., 2013) and regional scale (Brutsché et al., 2016; Toimil et al., 2017).”

R2.3 Page 5, line 1. According to the authors, d_c is calculated using the standard definition of wave height associated with a probability of occurrence of 12 hours/year but applied to a long (34 years) time series. Although used time series is long enough to be representative, it is not infrequent (Udo & Takeda reference provided by authors is an example) to assume that when d_c is going to be used in long-term processes characterization, this average wave climate can be substituted by other characteristic of extreme conditions (e.g. by using a wave height associated with a given return period). Which should be the expected impact of this approach in the dataset?

We think Reviewer 2 might have misunderstood our approach on this matter. We used the 34 year reanalysis of wave heights and periods to calculate the wave height exceeded only 12 hour in these 34 years. This means that we take the wave height with an exceedance probability of 0.137% in these 34 years, not the average of the yearly values. Nevertheless, as we understand that the current version might be confusing, we have now tried to make it more clear in the manuscript:

Pg.5 Line 13: “Following Nicholls et al. (1998), here we calculate $H_{e,t}$ as the wave height that is not exceeded more than 12 hours over the full 34 years’ time series. This approach is expected to lead to larger d_c values than taking the yearly statistics, as larger waves will be included in the time series.”

R2.4 Page 5, line 2. d_c is calculated using the Vousdoukas et al (2018b) wave data. However, there is no information on how good is the data set. In fact, in the original reference there is not too much mention to the calibration of hindcast against measured wave data. Since this parameter is used to define the profile length to compute the nearshore slope, which is the potential impact of the accuracy (or lack of) on the obtained values.

The reference we used previously did not include the validation of the wave model. We have now replaced the reference with the one, in which the detailed information of the validation can be found in the appendix:

Pg.5 Line 9: “Here we applied Eq. (1) using the full time series of significant wave heights H_s and peak wave periods T_p from the global reanalysis covering the period between 1980 and 2014 presented by Vousdoukas et al. (2017) . Waves were simulated using the third generation spectral wave model WW3. The model has been extensively validated and detailed information can be found in the reference provided.”

R2.5 Page 5, lines 5-6. Authors select d_c in an opposite manner as it has been traditionally defined, i.e. it is usually calculated as the most landward position whereas authors’ method results in the most seaward location.

We agree with Reviewer 2 that this is not the traditional way to define the depth of closure. Nevertheless, this would make a difference mainly in systems with pronounced sandbar systems, which cannot be explicitly captured with the present resolution. We have tested the sensitivity to this option and the differences in the calculated slopes were insignificant. We did these for the 8 validation sites that we use in our study and found that the error statistics had minor differences between the two methodologies (in the order of 2%).

R2.6 Page 6, line 23. This is not a validation of d_c computations. Authors are comparing their estimation with another estimation. Since in both cases they are using the same equation (no d_c data are available in the comparison), they are essentially comparing used wave data, i.e. they are checking whether Vousdoukas and WIS wave data are comparable (at statistical terms). Due to this, sentence in line 28 is not fully true. Differences can simply be explained by the difference in wave statistics at a given location among both wave data sources.

In order to do an explicit validation of d_c values the only option would be to compare with d_c values derived locally by numerous surveys to identify the location of no bed level change. This approach is quite challenging as long term local datasets are not always available. That is why we decided to at least validate our results with respect to previous studies which used

the same formulations. Nevertheless, we agree that this is not a direct validation of the d_c values. Therefore we have now used the word comparison in place of validation, as follows:

Pg.7 Line 4: “A thorough validation of the estimated depths of closure would demand various local bathymetrical surveys with good spatial resolution and sufficiently long temporal resolution and extent (i.e. 34 years of data). This kind of information is not readily available. Therefore, here we compare our results with”

R2.7 Due to the above-mentioned questions regarding d_c and its implications for calculating nearshore slope, it would be interesting which is the impact of changing d_c in different % (shallower and deeper) in the estimated nearshore slope. This will help to assess the robustness of estimated values as well as the real impact of the selection of the d_c -value on the data set due to the base data (shallow water bathymetry) accuracy and resolution.

We have re-calculated the nearshore slopes globally by adjusting the local depth of closure with a percentage of the originally estimated one. This adjustment was chosen to be -20%, -10%, +10% and +20% of the original value. We have now added the sensitivity test results to the manuscript and supplement:

Pg.12 Line 5: “Another important matter is the sensitivity of the estimated nearshore slopes to the depth of closure values used in this study. In order to test the robustness of our dataset, and how different wave statistics or depth of closure formulae could affect our estimations, we varied the previously calculated d_c values (Sect. **Error! Reference source not found.**). Changes of -20%, -10%, 10% and 20% of the originally estimated d_c were tested. The normalized differences of the estimated nearshore slopes had moderate sensitivity to the depth of closure variation (Fig. S12). When the depth of closure was increased and decreased by 10%, at almost 70% of the transects, the nearshore slope changed by the same percentage. On the other hand, with an increase and decrease of the d_c by 20%, the nearshore slope difference was in the order of 20% at about 75% of the transects. The number of transects with changes up to 20% was higher when only the sandy transects were considered. In general, the largest increase in the nearshore slope was observed when d_c was decreased. The spatial distribution of the relative nearshore slope changes showed that in most case the sensitivity was low. The most sensitivities areas to changes in d_c were areas with steep slopes (e.g. fjord areas in Norway, Chile and others) (Fig. S13).”

Figure S 12 (Fig. 1 in this response) : Histogram of normalized nearshore slope difference for all the transects (blue) and sandy transects (red) for four changes in the depth of closure (-20%, -10%, 10% and 20%). The data have been grouped in bars in increments of 10%.

Figure S 13 (Fig. 2 in this response): Global maps of normalized nearshore slope difference for all the transects for four changes in the depth of closure (-20%, -10%, 10% and 20% from top to bottom).

R2.8 Page 8, lines 20-23. The generally steeper slope along the West Americas coast is explained by using the role of swell as well as tectonics. Which should be the most dominant factor when dealing with the inner shelf slope? Is also sediment size any factor to be considered?

It is quite challenging to thoroughly address the question of which is the dominant factor across the entire West Americas coast. Nevertheless we have now tried to include all the factors that could play a role and reformulate our discussion:

Pg.9 Line 9: “Sediment grain size can play an important role as well on the shape and slope of the nearshore profile (Dean, 1991).”

Pg.9 Line 12: “The local slope values are ultimately the results of the combination of all the physical processes mentioned previously. At sedimentary coasts the upper nearshore profile is mainly influenced by waves while the lower one is influenced by tectonics, sediment supply or major storm events.”

R2.9 Section 5.2 apparently deals with possible improvements in estimation in SLR induced retreat. However, it is only a comparison between estimations using the provided dataset against the use of a common and single value worldwide (1/100). In this sense, it is just an improvement when comparing in those conditions. However, if these estimations are compared with many of the existing ones using local data, the improvement is not so evident.

As the reviewer points out, the improvement we highlight here is indeed that of using our dataset against the assumption of globally uniform nearshore slope (e.g. Hinkel et al. 2013). We agree with the reviewer that if local observations are available they would provide more accurate coastal retreat estimates locally. But for regional, continental or global studies that seek to quantify trends and hotspots, our dataset can provide an important improvement by incorporating the spatial variability of nearshore slopes versus a uniform slope assumption. In the manuscript we highlight the purpose of this comparison in the introduction and in the discussion sections (Pg. 2 Lines 5-7 and Pg. 9 Lines 13-17)

R2.10 Limitations is an important aspect to be considered. In this case, validation of the dataset needs to be improved by increasing local datasets to compare along world coastlines under different conditions and characteristics.

We agree with the comment of the reviewer. For this work, we have made use of most of the authors' contacts globally to get as much data as we could. The challenge is that local bathymetric data are sometime not publicly available. We expect that since our publication and dataset are open access, researchers in possession of local data could further validate our estimates against their measurements. We have additionally added a small part in the Discussion section that highlights the need for more data:

Pg.11 Line 25: “More long term coastal profile datasets, including coastal sites in diverse coastal environments, are needed to ensure a thorough validation of our dataset. The wealth of long term satellite data combined with the constantly evolving remote sensing techniques can be potentially an avenue to consider in the future.”

Global distribution of nearshore slopes with implications for coastal retreat

Panagiotis Athanasiou^{1,2}, Ap van Dongeren^{1,5}, Alessio Giardino¹, Michalis Voudoukas^{3,4}, Sandra Gaytan-Aguilar¹, Roshanka Ranasinghe^{5,2,1}

5 ¹ Deltares, Delft, The Netherlands

² Water Engineering and Management, Faculty of Engineering Technology, University of Twente, Enschede, The Netherlands

³ European Commission, Joint ~~European~~ Research Centre (JRC), Ispra, Italy

⁴ Department of Marine Sciences, University of the Aegean, Mitilene, Greece

10 ⁵ IHE Delft Institute for Water Education, Delft, The Netherlands

Correspondence to: Panagiotis Athanasiou (Panos.Athanasiou@deltares.nl)

Abstract

Nearshore slope, defined as the cross-shore gradient of the subaqueous profile, is an important input parameter which affects hydrodynamic and morphological coastal processes. It is used in both local and large-scale coastal investigations. However, due to unavailability of data, most studies, especially those that focus on continental or global scales, have historically adopted a uniform nearshore slope. This simplifying assumption could however have far reaching implications for predictions/projections thus obtained. Here, we present the first global dataset of nearshore slopes with a resolution of 1 km at almost 620,000 points along the global coastline. To this end, coastal profiles were constructed using global topobathymetric datasets. The results show that the nearshore slopes vary substantially around the world. An assessment of sea level rise (SLR) driven coastline recession (for an arbitrary 0.5 m SLR) with a globally uniform coastal slope of 1:100, as done in previous studies, and with the spatially variable coastal slopes computed herein shows that, on average, the former approach would under-estimate coastline recession by about 40%, albeit with significant spatial variation. The final dataset has been made publicly available at <https://doi.org/10.4121/uuid:a8297dcd-c34e-4e6d-bf66-9fb8913d983d>.

1 Introduction

25 Ten percent of the world's population lives in low-lying coastal areas; i.e. less than 10 m above the current mean sea level (MSL) (McGranahan et al., 2007). In the future, the population density in these areas is expected to increase even more due to high rates of population growth and urbanization (Neumann et al., 2015). At the same time, coastal areas are exposed to a number of marine hazards that can lead to flooding or erosion. As the coastline comprises various different landforms (e.g. sandy coasts, rocky cliffs etc.) and land uses (e.g. heavily urbanized or natural), the response to these hazards can vary significantly both spatially and temporally. One of the most vulnerable coastal types are sandy coasts which are highly

30

dynamic and can change in response to extreme marine events (McCall et al., 2010), long term trends in MSL (Zhang et al., 2004), natural gradients in alongshore sediment transport (Antolínez et al., 2018) or human interventions (Giardino et al., 2018a; Luijendijk et al., 2018; Mentaschi et al., 2018).

Local, regional or global studies that seek to quantify natural or human induced coastal change require high quality

5 nearshore bathymetry and sub-aerial topography data. However, as such data are rarely available, previous studies, especially at the regional or global scale, have had to rely on a number of limiting assumptions, such as globally uniform nearshore or beach slopes (Hinkel et al., 2013; Melet et al., 2018). The nearshore area is here defined as the part of the cross-shore profile between the depth of closure d_c (i.e. ~~the depth seaward of which there is no significant change in bottom elevation~~
~~the offshore limit where sediment transport is zero~~) and the shoreline (MSL). The nearshore slope (the ratio of the d_c over the
10 horizontal distance between the d_c and the shoreline) modulates the wave transformation (Battjes, 1974) and the total water levels (Serafin et al., 2019), while it is associated with geomorphological processes at various temporal scales (Bruun, 1962; Wright and Short, 1984; Dean, 1991). Therefore, a global assumption of uniform slope likely hides the spatial variability of coastal hydro- and morphodynamics.

Accurate information on the offshore limit of the nearshore area (i.e. the depth of closure d_c) and the nearshore slope is

15 important for numerous coastal engineering applications. For example, when behaviour-oriented models are employed, the depth of closure and the nearshore slope are crucial model inputs (Larson et al., 2004; Ruggiero et al., 2010; Hanson, 2014). At data poor locations, when bathymetric data are not available, an equilibrium bathymetric profile (Dean, 1991) is often assumed and the local d_c is needed to define the offshore profile limit (Udo and Takeda, 2017). Various assessments of future coastal recession due to sea level rise (SLR) have used the Bruun rule (Bruun, 1962) to quantify coastal retreat (Zhang
20 et al., 2004; Hinkel et al., 2013; Baron et al., 2014; Monioudi et al., 2017; Udo and Takeda, 2017; Ballesteros et al., 2018; Giardino et al., 2018b), an approach which is very sensitive to the nearshore slope. ~~Furthermore, several studies,~~
~~The most common approach to estimate d_c is using empirical formulae (Hallermeier, 1978; Birkemeier, 1985; Nicholls et al., 1998)~~
~~that relate d_c to wave parameters. This has been employed in several studies~~ both at the global scale (Hinkel et al., 2013)
and regional scale (Brutsché et al., 2016; Toimil et al., 2017), ~~have estimated the d_c using empirical formulae (Hallermeier,
25 1978; Birkemeier, 1985; Nicholls et al., 1998) that relate d_c to wave parameters.~~

Against the foregoing background, the present study sets out to provide the first ever global database of nearshore slopes and their associated depths of closure, which can be used in local, regional or global studies which aim to quantify coastal response to natural (i.e. marine extremes and climate change) or human induced phenomena. Our methods utilize available open-source global datasets of topography, bathymetry and shoreline location in order to create a seamless representation of
30 coastal morphology. Furthermore, global wave statistics are used to estimate d_c . The resulting dataset consists of an estimation of the local nearshore slopes at ~620.000 points along the global coastline.

This paper is structured as follows: Section 2 describes the methods used for the creation of the seamless coastal morphology map, the calculation of the global depths of closure, the definition of the coastal profiles and the calculation of the nearshore slopes. Section 3 presents the results of the analysis, consisting of a global dataset of depths of closure and nearshore slopes while Sect. 4 compares the results versus observed data at several locations around the world. The paper concludes with a discussion of key points, including a global analysis of the sensitivity of coastal recession estimates using the Bruun rule to the nearshore slopes, and the main conclusions.

2 Materials and methods

2.1 Workflow to obtain nearshore slopes

The estimation of nearshore slope required the following steps (Fig. 1). First, a seamless topo-bathymetric map was created (Sect. 2.2.1), then cross-shore transects were drawn with an accompanying elevation profile (Sect. 2.2.2). Using global wave data the depths of closure were determined (Sect. 2.3), after which the nearshore slopes could be determined (Sect. 2.4).

2.2 Coastal profiles

2.2.1 Global topo-bathymetric data

A cross-shore contiguous representation of the coastal profile is needed in order to accurately derive geometrical parameters such as the nearshore slope. Presently, elevation data are separated between topographic, i.e. representing the subaerial surface of the Earth, and bathymetric, i.e. representing the subaqueous seabed. Here we merge two of these datasets in order to create a seamless elevation layer with a smooth transition between land and water.

The topographic layer that was used is the MERIT digital elevation model (DEM) (Yamazaki et al., 2017), which is an improved version of the existing space borne DEMs SRTM3 v2.1 (Farr et al., 2007) and AW3D-30m v1 (Tadono et al., 2015), after removing multiple error components (i.e. absolute bias, stripe noise, speckle noise, and tree height bias). It represents the terrain elevations at a 3 arc-sec resolution (~90 m at the equator) and covers land areas between 90N-60S, referenced to EGM96 geoid. The bathymetric layer was obtained from the General Bathymetric Chart of the Oceans (<https://www.gebco.net>) which is based on ship track soundings with interpolation between soundings guided by satellite-derived gravity data. The version used was the GEBCO_2014 Grid (Weatherall et al., 2015), which has a 30 arc-sec resolution (~900 m at the equator) and it is generated by the aggregation of heterogeneous data types assuming all of them to be referred to MSL.

As the two datasets might overlap, the Open Street Map (OSM) (OpenStreetMap contributors, 2015) coastline was used to create a mask for land and water. For the final elevation dataset, a common grid was used, with a resolution corresponding to the higher resolution of the MERIT DEM dataset. Subsequently, the bathymetric data were interpolated only at the cells

seaward of the OSM coastline. This was performed by a linear interpolation of the values at neighbouring grid points in the x and y dimension. One of the key elements when merging bathymetry and elevation data is to create a smooth transition between land and water. We have applied a Low Pass Filter to decrease disparities between pixel values by averaging nearby pixels. The smoothing technique was applied only on the cells along the coastline. For each cell the vertical offset Δz was computed (as the difference in elevation with respect to its neighbouring cell). For the smoothing procedure different windows (i.e. number of grid cells around the grid cell to be smoothed) were used according to Δz . Smoothing window sizes of 3x3 were applied at areas where Δz was up to 10 m, 5x5 grid cells where Δz higher than 10m and 9X9 is few geographic areas where topography data present a big discrepancy with respect to the bathymetry.

2.2.2 Global elevation transects

The global coastline was defined using the OSM dataset of 2016. Since the full scale coastline was found to be too detailed for the purposes of the present work, the level 8 generalized version of the dataset was used, which is a smoothed version that removes small details and has been previously used in global coastal assessments (Luijendijk et al., 2018). About 1,000,000 cross-shore transects were defined along the OSM coastline, spaced at 1 km intervals. To ensure a globally consistent spacing and length of the transects, the coastline was re-projected to the local UTM projection, before creating the transects. The middle point of each transect was defined at the OSM location, with a 4km extension both to the landward and seaward direction. The profile length was chosen after testing different lengths to ensure that the active coastal profile is captured while minimizing data storage. The 1 km spacing was chosen in order to have a good coverage of the alongshore variability of the coastal profiles, while keeping the computational costs at a feasible level. Along each transect, a set of equidistant points was defined at an interval of 25 m, while flagging the locations where the elevation values were extracted from the global grid. For each transect the longitude and latitude of the extraction points were used to sample from the global merged elevation grid, and create a continuous elevation profile.

The global transects (~1,000,000) were first filtered to exclude transects located along coasts covered with ice, river inlets, or major ports and other significant human interventions, using the global transects of Luijendijk et al. (2018). This reduced the total number of transects to about 780,000 transects. Additionally, and specifically for the application described in Sect. 4.2, the sandy beach location data from Luijendijk et al. (2018) were used in order to identify the transects that were sandy, resulting in about 215,000 sandy transects. Note that gravel beaches are included in this dataset and are herein referred to as sandy.

2.3 Calculation of the depth-of-closure

The depth of closure was calculated using the formulation of Nicholls et al. (1998):

$$d_c = 2.28H_{e,t} - 68.5 \left(\frac{H_{e,t}^2}{gT_{e,t}^2} \right), \quad (1)$$

where $H_{e,t}$ is the significant wave height that is exceeded only 12 hours per t years, $T_{e,t}$ is the associated wave period, and g is the gravitational acceleration. The selection of the length of the time series t in this formula is directly associated with the temporal scale of the application of the depth of closure. This means that, for example, applications that consider temporal scales in the order of decades (i.e. coastline models, Bruun rule) should use a t that is consistent with the assessment scale.

5 Naturally, the temporal extent of the time series is dictated by the length of the available observations or modelling results. For example, Udo and Takeda (2017) applied Eq. (1) for SLR induced coastal recession assessments along the Japanese coast taking into account the effects of time scale for their 100-year Bruun rule application by using the maximum wave heights in their 5 years long wave record.

Here we applied Eq. (1) using the full time series of significant wave heights H_s and peak wave periods T_p from the global
10 reanalysis covering the period between 1980 and 2014 presented by Vousdoukas et al. (2017). Waves were simulated using the third generation spectral wave model WW3. The model has been extensively validated and detailed information can be found in the reference provided. The wave parameters were available every 3 hours at offshore locations covering the ice-free coasts. Following Nicholls et al. (1998) here we calculate $H_{e,t}$ as the wave height that is not exceeded more than 12 hours over the full 34 years' time series. This approach is expected to lead to larger d_c values than taking the yearly statistics, as
15 larger waves will be included in the time series.

2.4 Estimation of the nearshore slope

For each of the transects, the cross-shore location of the depth of closure d_c was identified using the elevation profile and the d_c value derived as described in this section. The location of the depth of closure was calculated by finding the first (i.e. moving from the most offshore point towards land) point which was shallower than the d_c and then linearly interpolating in
20 between the adjacent cross-shore points along the transect. Then the cross-shore location of the shoreline (MSL) point was estimated using a similar method. In case of multiple shoreline points (e.g. barrier islands) the point that was located closer to OSM defined point was chosen. The cross-shore distance between these two points was used as the horizontal length of the nearshore area L . The nearshore slope $\tan(\beta_{ns})$ was then calculated as the ratio $\frac{d_c}{L}$.

At locations in which the calculation could not be carried out as described above, an error code was assigned to describe the
25 cause of the calculation failure (Fig. S1-S2). For example, as the OSM coastline does not represent the open coast only, but includes a substantial number of areas with elevation that is far from the conventional profile presented in Fig. 1, the calculation of the slope in this way was not always successful. This was typically the case for coastlines dominated by fjord features (e.g. Norway, Chile). Furthermore, some transects did not have any offshore point in close proximity for the calculation of the depth of closure. The transects that of which the nearshore slope could not be calculated were about
30 160,000 of the total 780,000. Almost 20% out of these 160,000 transects were identified as sandy.

At some other locations (mostly embayed or protected areas) the depth of closure point was located seaward of the most offshore profile point. Therefore, the deepest point was taken as the depth of closure d_c . Additionally, due to the merging process of the bathymetric and topographic elevation data, elevation steps close to the shoreline were observed in some transects. These transects were determined by identifying abrupt changes of the slope close to the shoreline. A warning code
5 was also assigned to such transects (Fig. S1-S2).

3 Results

3.1 Global depths of closure

The resulting global distribution of d_c (Fig. 2) showed that on open ocean coasts, d_c generally increased with latitude because of the wave height, while d_c was smaller along the shores of more wave-sheltered seas. The global average value of
10 d_c was approximately 13 m, while it reached values of more than 20 m at some areas at higher latitudes (Fig. 3).

3.2 Nearshore slopes

The calculated nearshore slopes (Athanasίου, 2019) showed high spatial variability globally (Fig. 4 with continental zooms in Fig. S3-S8). The histogram of the calculated nearshore slopes (Fig. 5a) revealed that the most commonly computed nearshore slopes were around 0.005. Additionally, there was a high probability of occurrence of nearshore slopes larger than
15 0.2 due to the high number of transects at fjords and other steep coastlines. Furthermore, we detected several mildly sloping locations (i.e. nearshore slope lower than 0.002). Such mildly sloping coastlines are mostly characterized by high tidal range and/or muddy coastlines (e.g. Northwest Australia). The median nearshore slope worldwide was 0.007, while the 10th and 90th percentiles were 0.001 and 0.565 respectively (Fig. 5b). When only the sandy transects were considered (see Sect. 2.2.2), the most common nearshore slope shifted to 0.01, which corresponds with the globally uniform profile slope assumed
20 in previous global scale studies (Hinkel et al., 2013).

The nearshore slope value is dependent on various geological and hydrodynamic factors, such as wave characteristics, sediment size, sediment supply and large scale geological and tectonic processes, among others. For that reason we aggregated the nearshore slopes along various regions around the world, defined heuristically considering geographical proximity and their corresponding oceans or seas. A distinct variation of the nearshore slopes can be seen among regions
25 (Fig. 6). The three regions with the on average steeper slopes were the Pacific Islands, the Norwegian Sea and the Mediterranean Sea with median values of 0.025, 0.020 and 0.014 respectively. On the other hand, the three regions with the mildest nearshore slopes were North Australia, East U.S and Gulf of Mexico and Southeast Asia with median values of 0.001, 0.002 and 0.003 respectively. The variance of the aggregated nearshore slopes per region (defined as the logarithmic difference between the 5th and 95th percentile) varied, with West S.America and North Australia showing the highest and
30 lowest variance respectively.

4 Comparison with local observations

4.1 Depth of closure

Different methods exist to estimate the d_c and therefore the validation of this parameter can be quite challenging at a global scale. A thorough validation of the estimated depths of closure would demand various local bathymetrical surveys with good spatial resolution and sufficiently long temporal resolution and extent (i.e. 34 years of data). This kind of information is not readily available. Therefore, here we compare our results with a study at the U.S coast (Brutsché et al., 2016), which employed various formulas to calculate d_c , including Eq.(1), ~~to compare our results~~. We compared the d_c at the offshore locations of the U.S. coast (Fig. 2) with the d_c of the closest point from Brutsché et al. (2016), calculated using the same formula (Eq.(1)). Our predicted d_c showed skill in capturing the spatial variation as calculated by Brutsché et al. (2016) (Fig. S9), but presented a positive bias of 3.34 m (Fig. S10). This can be attributed to the different location and temporal scale of the wave statistics (i.e. Brutsché et al. (2016) used nearshore transformed wave time series of a 20 years hindcast).

4.2 Nearshore slopes

The computed slopes were validated both qualitatively and quantitatively. The former was performed regionally using coastal classifications of the European and the Mediterranean coastlines, while the latter employed observations at sandy beaches derived from local surveys at 8 study sites around the world.

4.2.1 Qualitative validation

Two coastal classification datasets were used to inspect the computed nearshore slopes grouped according to geomorphological coastal types. One was the Mediterranean Coastal Database (MCD) (Wolff et al., 2018), which classifies the Mediterranean coastline in four classes: 1) sandy beaches, 2) unerodible coasts, 3) muddy coastlines and 4) rocky coasts with pocket beaches. The other was the EUROSION geomorphological classification (EUROSION, 2004) of the counties of the European Union (EU), which includes 20 different coastal types. The latter was reclassified to the four classes of the MCD database in order to reduce the total number of classes (Fig. S11)

The calculated nearshore slopes followed the expected variability between the different coastal typologies for both classification datasets (Fig. 7). For example, muddy coastlines had relatively milder nearshore slopes (median values of 0.0024 and 0.0025 for MCD and EUROSION datasets respectively). Sandy coastlines had a median nearshore slope of 0.0107 and 0.0066 respectively for the MCD and EUROSION datasets, which relates to the steeper sandy beach slopes observed along the Mediterranean coastline. Unerodible and rocky coastlines had on average relatively steeper slopes.

4.2.2 Quantitative validation

The study sites for the quantitative validation were selected according to data availability and a criterion of having at least 20 transects overlapping with the derived global dataset to enable the computation of error statistics per site. The validation focussed only on sandy coastline transects that are not adjacent to morphologically complex areas (e.g. inlets, river mouths) determined using satellite imagery. The ground truth sites (Fig. 8; Table 1) represent different coastal geomorphologies with varying hydrodynamic forcing and span across different continents. We performed the validation on the basis of the nearshore slopes, using the d_c that were derived from the global analysis of wave data described in Sect. 2.2.2, for both local and global nearshore slope estimation. The local nearshore slope was calculated using the methodology described in Sect. 2.3.

10 When the elevation was represented by locally defined profiles, the comparison was performed using the closest globally derived transect. When point cloud data were available, they were first interpolated on a common grid and then the elevation was extracted along the global transects. The same process was performed when an elevation raster was available.

For a quantitative evaluation of the accuracy of the globally predicted slope values against the locally observed ones, various statistical parameters were calculated per site, as well as globally. This included the coefficient of determination R^2 , the normalized mean bias (NMB) and the normalized root mean square error (NRMSE). Note that there is no temporal coherence between the predicted and observed data, since the global elevation dataset is a merged product from temporally varying bathymetric sources (see Sect. 2.2.1) and the observed data were taken at different times. Therefore, it is expected that part of the calculated errors is due to potential temporal differences of the estimated and observed elevation data.

A regression analysis performed over all of the available global observations resulted in a normalized mean bias of almost -20 25% and an R^2 value of 0.44 (Fig. 9) implying that that extracted nearshore slopes captured the spatial variability of the ground truth data. The negative overall bias indicated that the calculated global slopes were on average milder than those observed.

The error statistics per validation site (Table 2) showed that overall, R^2 values were above 0.6 except for North California, Recife and South California where poorer agreement was found (i.e. 0.38, 0.25 and 0.3 respectively). At all validation sites (except Monterey Bay) a negative bias was observed. The NRMSE varied per study site, with values lower than 35% for five out of the eight validation sites. The high NRMSE at Emilia Romagna can be attributed to the quite high bias, but nevertheless the spatial variability was captured correctly ($R^2 = 0.7$). It should be noted that the number of overlapping sandy transects differed significantly between sites as the spatial extent of the available local surveys differed. For example, at Emilia Romagna and Monterey Bay there were only 25 and 27 overlapping transects, while at South California 382.

5 Discussion

5.1 Spatial variation of nearshore slopes

As discussed in the previous section the results of the present study revealed a quite pronounced spatial variation in nearshore slopes worldwide (Fig. 4). For example, the West coast of North and South America had on average steeper slopes in comparison to the East, which can be attributed to the swell dominated wave climate along the former (the same holds for the West and East coasts of Africa). The classification of the coasts with respect to the plate tectonics (Inman and Nordstrom, 1971) seems to be relevant to nearshore slopes distribution as well. Trailing edge coast with wider continental shelf width (e.g. North Sea, East Americas, North Australia) appeared to have milder nearshore slopes than leading edge coasts (e.g. West Americas and the biggest part of the Mediterranean). Sediment grain size can play an important role as well on the shape and slope of the nearshore profile (Dean, 1991). The coasts of the Gulf of Mexico, South and Southeast Asia showed on average mild nearshore slopes, which can be attributed to the large (historic) sediment supplies by some major rivers therein. The local slope values are ultimately the results of the combination of all the physical processes mentioned previously. At sedimentary coasts the upper nearshore profile is mainly influenced by waves while the lower one is influenced by tectonics, sediment supply or major storm events.

5.2 Improvements in estimations of SLR-induced retreat of sandy coasts

Similarly to the overall data set (including non-sandy coasts), the subset of nearshore slopes of sandy coasts displays a considerable spatial variation, even though the commonly-used 1:100 nearshore slope assumption (Hinkel et al., 2013) lies in the most probable range (Fig. 5). Using the Bruun rule (Bruun, 1962), we investigated the effect of applying the spatial variation of the nearshore slope on the estimation of shoreline retreat due to sea level rise, in comparison with applying a uniform 1:100 slope as commonly done for sandy beaches.

The Bruun rule is a simple two-dimensional mass conservation principle, which predicts a landwards and upwards response of sandy profiles as a response to SLR. It is expressed as (Bruun, 1962):

$$R = \frac{L}{d_c} S, \quad (2)$$

Where R is the horizontal coastal recession, d_c is the depth of closure, L is the horizontal length from the shoreline to the DoC and S is the SLR. While used extensively for lack of an efficient better method, its use has been a controversial matter in literature during the last years (Cooper and Pilkey, 2004; Ranasinghe and Stive, 2009), as it is based on a number of assumptions and simplifications (i.e. equilibrium profile existence, cross-shore sediment balance). Nevertheless it is the only method that can deal with coastal recession due to SLR at large scale in a computationally feasible manner. Since

$\tan(\beta_{ns}) = \frac{d_c}{L}$ this makes Eq.(2):

$$R = \frac{1}{\tan(\beta_{ns})} S, \quad (3)$$

Physically, there are limits of the nearshore slopes that can be encountered at sandy beaches related to the angle of repose of granular materials. For our dataset, the sandy transects (Luijendijk et al., 2018) were found to lie beyond these ranges in some cases. This can be attributed to local effects of geology, sediment budget, but also to data and methodological artefacts. Therefore, we applied upper and lower limits of 0.2 and 0.001 for the sandy nearshore slopes and constrained the slopes that lie beyond these limits to the limit values.

Following the linear relationship between the coastal recession R and the nearshore slope in Eq. (3), the differences in the coastal recession under a given SLR scenario can show significant spatial variations. In order to highlight this, the ratio between the coastal recession R_{calc} (i.e. using the nearshore slopes presented herein) and R_{100} (i.e. using a 1/100 nearshore slope globally) was estimated globally for all sandy transects (Fig. 10). The results indicate that for a large number of areas around the world the assumption of 1:100 nearshore slopes underestimates the potential coastal recession with a median $\frac{R_{calc}}{R_{100}}$ of 1.39. However, it should be noted that there are also several locations with steeper slopes, where the 1:100 assumption overestimates the SLR induced erosion.

In order to highlight the potential impact of this effect, a coastal recession estimate was computed for all transects of each region (see Fig. 6) using Eq. (3) for an arbitrary SLR of 0.5 m across the global coastline. The median value of the coastal recession of all transects per region was calculated as a robust description of its central tendency (Fig. 11) and compared with the value of 50 m (which is the recession that would be calculated for a SLR of 0.5 m using the 1:100 slope assumption). The results show that, in most of the defined regions the assumption of a 1:100 nearshore slope would result in the underestimation of coastal recession, especially in North Australia, South and Southeast Asia and the East U.S and Gulf of Mexico regions. However in some of the regions, such as the Mediterranean and the Norwegian Seas, and the West Americas the 1:100 slope assumption would lead to an overestimation due to the steeper nearshore slopes encountered on average there.

5.3 Limitations

The global scale of the present work introduced inevitable limitations related to data availability, computational efficiency and methodological constraints. Merging bathymetric and topographic data at a global scale can be quite challenging. This is associated with differences in spatial resolution, potential overlaps or differences in the vertical datum. Here we tried to resolve these issues by using the OSM coastline as the MSL to differentiate between sea and land and connect the elevation data. Although other available coastline datasets exist, either they have a coarser resolution, therefore not allowing for a correct representation of the separation between land and water, or do not have a global coverage. The importance of the coastline and DEM data has been previously highlighted with respect to statistics of exposure information (Lichter et al.,

2011). We expect that the choice of the coastline can have an effect in the elevation profiles, especially in areas with a high tidal range. Nevertheless

Additionally, GEBCO is a combination of bathymetric data derived from variable sources and different techniques and in some cases might have differences in the vertical datum (Weatherall et al., 2015). This might introduce regional differences on the trustworthiness of the results. The smoothing procedure that was applied to the bathymetric and topographic data may have affected the nearshore slope values, especially at cliffed coastlines, where the computed slopes may be milder than the reality. Nevertheless, cliffed transects retained quite steep values as can be seen by the second peak in the global slope distribution (Figure 5). Furthermore, we expect that at locations where the nearshore slopes are steep, the resolution of the bathymetry is not high enough to capture a large number of bathymetry points between the shoreline and the depth of closure, hiding the details of the profile. We believe that when bathymetrical dataset with better resolution and accuracy become available in the future, the presented technique can be applied to update the global nearshore slopes estimations.

The wave statistics that were used to determine the depth of closure across the global transects of the present study were taken from the closest offshore point of the wave reanalysis output (see Sect. 2.3). Ideally, the wave time series of the 28 years used here should be transformed to nearshore conditions at a location just offshore of the depth of closure using Snell's Law. This would require an assumption of a nearshore slope, then the calculation of d_c and then the re-calculation of the nearshore slope using an iterative approach until the slopes converge. This was deemed to be outside the scope of this study, considering the large number of transects. Nevertheless, we found that the nearshore slopes were not very sensitive to small changes in the depth of closure.

With respect to the comparison of the dataset compiled in the present study with available observations (Sect. 4), it should be mentioned that due to the scale of the study, it is quite challenging to perform a direct validation with the same spatiotemporal conditions. For example the comparison of the estimated d_c was performed against the results of a study that used the same formula, but employing wave statistics with different temporal extent and at different locations (Sect. 4.1). Additionally, the local surveys used for the quantitative comparison of the nearshore slopes had various sources, used different measuring techniques and were collected at different times (Sect. 4.2.2). Naturally, it is expected that these spatiotemporal differences had an impact on the error statistics computed herein. More long term coastal profile datasets, including coastal sites in diverse coastal environments, are needed to ensure a thorough validation of our dataset. The wealth of long term satellite data combined with the constantly evolving remote sensing techniques can be potentially an avenue to consider in the future.

In this study the nearshore slope was defined from the depth of closure until the shoreline. This could be a deviation from certain Bruun rule implementations that define the active profile slope from the depth of closure until the dune or berm crest (Zhang et al., 2004; Hinkel et al., 2013; Toimil et al., 2017; Udo and Takeda, 2017). Given that the available global topographic datasets lack the resolution to resolve dunes or berms (Vousdoukas et al., 2018b), their inclusion in the analysis

was not feasible, potentially resulting in an underestimation of the active profile slope (from the depth of closure to the top of the dune) along beaches with prominent dune systems. Nevertheless, since our dataset includes the nearshore slope and the depth of closure, one could calculate the slope from depth of closure to the dune or berm height if this information is available locally.

Another important matter is the sensitivity of the estimated nearshore slopes to the depth of closure values used in this study. In order to test the robustness of our dataset, and how different wave statistics or depth of closure formulae could affect our estimations, we varied the previously calculated d_c values (Sect. 2.3). Changes of -20%, -10%, 10% and 20% of the originally estimated d_c were tested. The normalized differences of the estimated nearshore slopes had moderate sensitivity to the depth of closure variation (Fig. S12). When the depth of closure was increased and decreased by 10%, at almost 70% of the transects, the nearshore slope changed by the same percentage. On the other hand, with an increase and decrease of the d_c by 20%, the nearshore slope difference was in the order of 20% at about 75% of the transects. The number of transects with changes up to 20% was higher when only the sandy transects were considered. In general, the largest increase in the nearshore slope was observed when d_c was decreased. The spatial distribution of the relative nearshore slope changes showed that in most case the sensitivity was low. The most sensitivities areas to changes in d_c were areas with steep slopes (e.g. fjord areas in Norway, Chile and others) (Fig. S13).

6 Data Availability

The final output data provide the location of each of the ~780,000 points along the global coastline together with: a) the depth of closure d_c , b) the nearshore slope $\tan(\beta_{sf})$ and c) an error /warning code in case the slope was not calculated or limitations in the profile were spotted. These data are given as a comma separated value file. The data can be downloaded via <https://doi.org/10.4121/uuid:a8297dcd-c34e-4e6d-bf66-9fb8913d983d> (Athanasίου, 2019).

7 Conclusions

A dataset of the global distribution of nearshore slopes at an alongshore resolution of 1 km is presented, using global elevation datasets and wave statistics. Depths of closure d_c were estimated worldwide using an empirical formulation and long-term wave statistics derived from 34-years wave reanalysis. The average d_c globally was almost 13m, attaining larger values at higher latitudes. The global median nearshore slope was 0.007, while most values were in the range 0.001 to 0.565 (10th and 90th percentile respectively). The most commonly computed nearshore slope was about 0.005, which increased to 0.01 when only sandy coasts were considered. However, the computed nearshore slopes exhibited substantial spatial variability, potentially corresponding with spatial variations in hydrodynamic forcing and geological conditions.

The dataset captured expected qualitative nearshore slope patterns when compared to available coastal classification datasets. Additionally, it captured the spatial variability of the nearshore slopes of sandy beaches at 8 validation sites from around the world, but with a negative bias of 25% (milder slopes than the observed ones).

5 The assessment of SLR driven coastline recession (for an arbitrary 0.5 m SLR) with globally uniform coastal slopes, as done in many previous studies, and the spatially variable nearshore slopes computed herein showed large differences between the recession amounts projected by the two approaches. Worldwide, the median coastline recession calculated with the nearshore slopes computed here was almost 40% larger than that computed by assuming a globally uniform 1:100 coastal slope, with the ratio between the two estimates varying substantially around the world. We believe that this dataset is a first step towards capturing the spatial variability of coastal profile characteristics and enabling a correct spatial representation of coastal impacts.

10

Author contribution

PA, AvD, AG, MV and RR designed the study. PA carried out the data processing and analysis. MV provided the offshore wave statistics. SGA composed the merged topobathymetric dataset. PA prepared the manuscript with contributions from all co-authors.

15 **Competing interests**

The authors declare that they have no conflict of interest.

Acknowledgements

This work has received funding from the EU Horizon 2020 Program for Research & Innovation, under grant agreement no 776613 (EUCP: “EUropean Climate Prediction system”; <https://www.eucp-project.eu>). Ap van Dongeren was funded in part by the Deltares Strategic Research Programme “Quantifying Flood Hazards and Impacts” while Alessio Giardino from the Research Programme “Flood Risk Strategies”. Roshanka Ranasinghe is supported by the AXA Research fund and the Deltares Strategic Research Programme “Coastal and Offshore Engineering”. We would like to acknowledge Arjen Luijendijk for providing the location of the sandy transects from their study. We would also like to acknowledge the map data copyrighted OpenStreetMap contributors for using the map data available from <https://www.openstreetmap.org>.

20

25

References

- Andrews, B. D., Defne, Z., Miselis, J. L. and Ganju, N. K.: Continuous terrain model for water circulation studies, Barnegat Bay, New Jersey, , <http://doi.org/http://dx.doi.org/10.5066/F7PK0D6B>, 2015.
- Antolínez, J. A. A., Murray, A. B., Méndez, F. J., Moore, L. J., Farley, G. and Wood, J.: Downscaling Changing Coastlines in a Changing Climate: The Hybrid Approach, *J. Geophys. Res. Earth Surf.*, 123(2), 229–251, <http://doi.org/10.1002/2017JF004367>, 2018.
- Athanasίου, P.: Global distribution of nearshore slopes, 4TU.Centre Res. Data, <http://doi.org/10.4121/uuid:a8297dcd-c34e-4e6d-bf66-9fb8913d983d>, 2019.
- Ballesteros, C., Jiménez, J. A., Valdemoro, H. I. and Bosom, E.: Erosion consequences on beach functions along the Maresme coast (NW Mediterranean, Spain), *Nat. Hazards*, 90(1), 173–195, <http://doi.org/10.1007/s11069-017-3038-5>, 2018.
- Barnard, P. L., van Ormondt, M., Erikson, L. H., Eshleman, J., Hapke, C., Ruggiero, P., Adams, P. N. and Foxgrover, A. C.: Development of the Coastal Storm Modeling System (CoSMoS) for predicting the impact of storms on high-energy, active-margin coasts, *Nat. Hazards*, 74(2), 1095–1125, <http://doi.org/10.1007/s11069-014-1236-y>, 2014.
- Baron, H. M., Ruggiero, P., Wood, N. J., Harris, E. L., Allan, J., Komar, P. D. and Corcoran, P.: Incorporating climate change and morphological uncertainty into coastal change hazard assessments, *Nat. Hazards*, 75(3), 2081–2102, <http://doi.org/10.1007/s11069-014-1417-8>, 2014.
- Battjes, J. A.: SURF SIMILARITY, in *Coastal Engineering Proceedings*, vol. 1, p. 26., 1974.
- Birkemeier, W. A.: Field Data on Seaward Limit of Profile Change, *J. Waterw. Port, Coastal, Ocean Eng.*, 111(3), 598–602, [http://doi.org/10.1061/\(asce\)0733-950x\(1985\)111:3\(598\)](http://doi.org/10.1061/(asce)0733-950x(1985)111:3(598)), 1985.
- Brutsché, K. E., Rosati, J., Pollock, C. E., McFall, B. C., Iii, J. R. and Cheryl, E.: Calculating depth of closure using WIS hindcast data., 2016.
- Bruun, P.: Sea-Level Rise as a Cause of Shore Erosion, in *Proceedings of the American Society of Civil Engineers.*, 1962.
- Cooper, J. A. G. and Pilkey, O. H.: Sea-level rise and shoreline retreat: Time to abandon the Bruun Rule, *Glob. Planet. Change*, 43(3–4), 157–171, <http://doi.org/10.1016/j.gloplacha.2004.07.001>, 2004.
- Dean, R. G.: Equilibrium Beach Profiles : Characteristics and Applications, *J. Coast. Res.*, <http://doi.org/10.2307/4297805>, 1991.
- EUROSION: Living with coastal erosion in Europe: Sediment and Space for Sustainability: PART I - Major findings and Policy Recommendations of the EUROSION project., 2004.
- Farr, T. G., Rosen, P. A., Caro, E., Crippen, R., Duren, R., Hensley, S., Kobrick, M., Paller, M., Rodriguez, E., Roth, L.,

- Seal, D., Shaffer, S., Shimada, J., Umland, J., Werner, M., Oskin, M., Burbank, D. and Alsdorf, D. E.: The shuttle radar topography mission, *Rev. Geophys.*, <http://doi.org/10.1029/2005RG000183>, 2007.
- Giardino, A., Schrijvershof, R., Nederhoff, C. M., de Vroeg, H., Brière, C., Tonnon, P. K., Caires, S., Walstra, D. J., Sosa, J., van Verseveld, W., Schellekens, J. and Sloff, C. J.: A quantitative assessment of human interventions and climate change on the West African sediment budget, *Ocean Coast. Manag.*, <http://doi.org/10.1016/j.ocecoaman.2017.11.008>, 2018a.
- Giardino, A., Nederhoff, C. and Vousdoukas, M. I.: Coastal hazard risk assessment for small islands: assessing the impact of climate change and disaster reduction measures on Ebeye(Marshall Islands), *J. Reg. Environ. Chang.*, 2018b.
- Hallermeier, R. J.: Uses for a calculated limit depth to beach erosion, in *Coastal Engineering*, pp. 1493–1512., 1978.
- Hanson, H.: Genesis: A Generalized Shoreline Change Numerical Model, *J. Coast. Res.*, 2014.
- 10 Hinkel, J., Nicholls, R. J., Tol, R. S. J., Wang, Z. B., Hamilton, J. M., Boot, G., Vafeidis, A. T., McFadden, L., Ganopolski, A. and Klein, R. J. T.: A global analysis of erosion of sandy beaches and sea-level rise: An application of DIVA, *Glob. Planet. Change*, 111, 150–158, <http://doi.org/10.1016/j.gloplacha.2013.09.002>, 2013.
- Inman, D. L. and Nordstrom, C. E.: On the Tectonic and Morphologic Classification of Coasts, *J. Geol.*, 79(1), 1–21, <http://doi.org/10.1086/627583>, 1971.
- 15 Larson, M., Erikson, L. and Hanson, H.: An analytical model to predict dune erosion due to wave impact, *Coast. Eng.*, 51(8–9), 675–696, <http://doi.org/10.1016/j.coastaleng.2004.07.003>, 2004.
- Lichter, M., Vafeidis, A. T. and Nicholls, R. J.: Exploring Data-Related Uncertainties in Analyses of Land Area and Population in the “Low-Elevation Coastal Zone” (LECZ), *J. Coast. Res.*, 27(4), 757, <http://doi.org/10.2112/jcoastres-d-10-00072.1>, 2011.
- 20 Luijendijk, A., Hagenaars, G., Ranasinghe, R., Baart, F., Donchyts, G. and Aarninkhof, S.: The State of the World’s Beaches, *Sci. Rep.*, 1–11, <http://doi.org/10.1038/s41598-018-24630-6>, 2018.
- McCall, R. T., Van Thiel de Vries, J. S. M., Plant, N. G., Van Dongeren, A. R., Roelvink, J. A., Thompson, D. M. and Reniers, A. J. H. M.: Two-dimensional time dependent hurricane overwash and erosion modeling at Santa Rosa Island, *Coast. Eng.*, <http://doi.org/10.1016/j.coastaleng.2010.02.006>, 2010.
- 25 McGranahan, G., Balk, D. and Anderson, B.: The rising tide: Assessing the risks of climate change and human settlements in low elevation coastal zones, *Environ. Urban.*, 19(1), 17–37, <http://doi.org/10.1177/0956247807076960>, 2007.
- Melet, A., Meyssignac, B., Almar, R. and Le Cozannet, G.: Under-estimated wave contribution to coastal sea-level rise, *Nat. Clim. Chang.*, 8(3), 234–239, <http://doi.org/10.1038/s41558-018-0088-y>, 2018.
- Mentaschi, L., Vousdoukas, M. I., Pekel, J. F., Voukouvalas, E. and Feyen, L.: Global long-term observations of coastal

- erosion and accretion, *Sci. Rep.*, 8(1), 1–11, <http://doi.org/10.1038/s41598-018-30904-w>, 2018.
- Monioudi, I. N., Velegrakis, A. F., Chatzipavlis, A. E., Rigos, A., Karambas, T., Vousdoukas, M. I., Hasiotis, T., Koukouroufli, N., Peduzzi, P., Manoutsoglou, E., Poulos, S. E. and Collins, M. B.: Assessment of island beach erosion due to sea level rise: The case of the Aegean archipelago (Eastern Mediterranean), *Nat. Hazards Earth Syst. Sci.*, 17(3), 449–466, <http://doi.org/10.5194/nhess-17-449-2017>, 2017.
- Neumann, B., Vafeidis, A. T., Zimmermann, J. and Nicholls, R. J.: Future coastal population growth and exposure to sea-level rise and coastal flooding - A global assessment, *PLoS One*, 10(3), <http://doi.org/10.1371/journal.pone.0118571>, 2015.
- Nicholls, R. J., Birkemeier, W. A. and Lee, G.: Evaluation of depth of closure using data from Duck , NC , USA, *Mar. Geol.*, 148, 179–201, 1998.
- 10 OpenStreetMap contributors: OSM coastline, [online] Available from: <https://planet.osm.org>, 2015.
- Ranasinghe, R. and Stive, M. J. F.: Rising seas and retreating coastlines, *Clim. Change*, 97(3), 465–468, <http://doi.org/10.1007/s10584-009-9593-3>, 2009.
- Rijkswaterstaat: Jarkus cross-shore profile measurements database, 2018.
- Ruggiero, P., Buijsman, M., Kaminsky, G. M. and Gelfenbaum, G.: Modeling the effects of wave climate and sediment supply variability on large-scale shoreline change, *Mar. Geol.*, 273(1–4), 127–140, <http://doi.org/10.1016/j.margeo.2010.02.008>, 2010.
- 15 Serafin, K. A., Ruggiero, P., Barnard, P. L. and Stockdon, H. .: The influence of shelf bathymetry and beach topography on extreme total water levels: Linking large-scale changes of the wave climate to local coastal hazards., *Coast. Eng.*, 150(May 2018), 1–17, <http://doi.org/10.1016/j.coastaleng.2019.03.012>, 2019.
- 20 Stevens, A. W., Logan, J. B., Snyder, A. G., Hoover, D. J., Barnard, P. L. and Warrick, J. A.: Beach topography and nearshore bathymetry of northern Monterey Bay, California: U.S. Geological Survey data release, , <http://doi.org/https://doi.org/10.5066/F76H4GCW>, 2017.
- Tadono, T., Takaku, J., Tsutsui, K., Oda, F. and Nagai, H.: Status of “ALOS World 3D (AW3D)” global DSM generation, *Int. Geosci. Remote Sens. Symp.*, 2015-Novem, 3822–3825, <http://doi.org/10.1109/IGARSS.2015.7326657>, 2015.
- 25 Toimil, A., Losada, I. J., Camus, P. and Díaz-Simal, P.: Managing coastal erosion under climate change at the regional scale, *Coast. Eng.*, 128(August), 106–122, <http://doi.org/10.1016/j.coastaleng.2017.08.004>, 2017.
- Udo, K. and Takeda, Y.: Projections of Future Beach Loss in Japan Due to Sea-Level Rise and Uncertainties in Projected Beach Loss, *Coast. Eng. J.*, 59(02), 1740006, <http://doi.org/10.1142/S057856341740006X>, 2017.
- Vousdoukas, M., Mentaschi, L., Voukouvalas, E., Kakoulaki, G., Pereira, P. S., Lino, A. P. and Feyen, L.: Assessing

compound flooding risk through dynamic simulations: application in Recife, Brazil, in *Geophysical Research Abstracts.*, 2018a.

Vousdoukas, M. I., Mentaschi, L., Voukouvalas, E., Verlaan, M. and Feyen, L.: Extreme sea levels on the rise along Europe's coasts, *Earth's Futur.*, 1–20, <http://doi.org/10.1002/ef2.192>, 2017.

- 5 Vousdoukas, M. I., Bouziotas, D., Giardino, A., Bouwer, L. M., Mentaschi, L. and Feyen, L.: Understanding epistemic uncertainty in large-scale coastal flood risk assessment for present and future climates, , (May), 1–27, 2018b.

Weatherall, P., Marks, K. M., Jakobsson, M., Schmitt, T., Tani, S., Arndt, J. E., Rovere, M., Chayes, D., Ferrini, V. and Wigley, R.: A new digital bathymetric model of the world's oceans, *Earth Sp. Sci.*, <http://doi.org/10.1002/2015EA000107>, 2015.

- 10 Wolff, C., Vafeidis, A. T., Muis, S., Lincke, D., Satta, A., Lionello, P., Jimenez, J. A., Conte, D. and Hinkel, J.: A Mediterranean coastal database for assessing the impacts of sea-level rise and associated hazards, *Sci. Data*, 5, 180044, <http://doi.org/10.1038/sdata.2018.44>, 2018.

Wright, L. D. and Short, A. D.: Morphodynamic variability of surf zones and beaches: A synthesis, *Mar. Geol.*, 56(1–4), 93–118, [http://doi.org/10.1016/0025-3227\(84\)90008-2](http://doi.org/10.1016/0025-3227(84)90008-2), 1984.

- 15 Yamazaki, D., Ikeshima, D., Tawatari, R., Yamaguchi, T., O'Loughlin, F., Neal, J. C., Sampson, C. C., Kanae, S. and Bates, P. D.: A high-accuracy map of global terrain elevations, *Geophys. Res. Lett.*, 44(11), 5844–5853, <http://doi.org/10.1002/2017GL072874>, 2017.

Zhang, K. Q., Douglas, B. C. and Leatherman, S. P.: Global warming and coastal erosion, *Clim. Change*, 64(1–2), 41–58, <http://doi.org/10.1023/b:clim.0000024690.32682.48>, 2004.

Table 1: Validation case studies description

Location (Country)	Data format	Resolution	Year	Source
Dutch Coast (The Netherlands)	Elevation profiles	Alongshore intervals of 250 m and cross-shore resolution of 5-10 m	2011	(Rijkswaterstaat, 2018)
Emilia Romagna (Italy)	Point cloud profiles	Alongshore intervals of ~500 m	2012	Topo-bathymetric surveys of beach profiles performed by Arpae-SIMC in the context of the Regional Topo-bathymetric Network
North California (U.S.)	Elevation profiles	Alongshore intervals of 100-200 m and cross-shore resolution of 2 m	Various (1929-2017)	(Barnard et al., 2014)
South California (U.S.)	Elevation profiles	Alongshore intervals of 100-200 m and cross-shore resolution of 2 m	Various (1930-2014)	(Barnard et al., 2014)
Monterey (U.S.)	Point cloud profiles	Alongshore intervals of 50-250 m	2017	(Stevens et al., 2017)
Long Island (U.S.)	Elevation raster	1/9 arc-sec cell size	2012	https://www.ngdc.noaa.gov/mgg/inundation/sandy/sandy_geoc.html
New Jersey (U.S.)	Elevation raster	10 m cell size	Various (1934-2013)	(Andrews et al., 2015)
Recife (Brazil)	Point cloud profiles	Alongshore intervals of ~150 m	Various (2012-2015)	(Vousdoukas et al., 2018a)

Table 2: Error statistics per validation site on the basis of the nearshore slopes $\tan(\beta_{ns})$.

Validation Site	Transects	Mean observed	NMB (%)	NRMSE (%)	R^2
nearshore slope					
Dutch Coast	83	0.0083	-18.0	24.4	0.62
Emilia Romagna	25	0.0062	-62.6	130.4	0.70
Long Island	139	0.0092	-31.4	33.3	0.80
Monterey Bay	27	0.0113	10.0	20.6	0.68
New Jersey	148	0.0100	-53.8	40.1	0.87
North California	71	0.0156	-11.2	18.4	0.38
Recife	62	0.0108	-83.6	65.0	0.25
South California	382	0.0204	-35.7	14.8	0.30

Figures




Figure 1: Work flow followed for the calculation of the nearshore slopes worldwide.




Figure 2: Depths of closure d_c along the global coastline using the formulation by Nicholls et al., (1998) (Eq. 1).




Figure 3: Histogram of the probability of occurrence of d_c globally (using bins of 0.5 m).




Figure 4: Global map of nearshore slopes. Red colours indicate steeper slopes while blue colours milder slopes. Note that in the colour scale the slopes have been grouped in non-equidistant increments in order to highlight the spatial differences.




Figure 5: (a) Histogram of the probability of occurrence of nearshore slopes. (b) Cumulative probability distribution of nearshore slopes. The graphs have been plotted for all transects (blue) and only for sandy transects (red). Note that the x axis is plotted in log scale.




Figure 6: (a) Nearshore slopes statistics per region. Dots indicate the median, while lines the range between the 5th and 95th percentiles. (b) Cumulative probability distribution of nearshore slopes per region. Dashed line indicates the 50% line. (c) Global map with the defined regions with their respective colour and id. Note that the sub-figures a and b use the colour scheme indicated by sub-figure c. The x-axis of sub-figures a and b are plotted at a log scale.




Figure 7: Cumulative probability distribution of nearshore slopes per coastal type for the (a) MCD dataset and (b) EUROSION dataset. The median values per coastal types are plotted with their respective colours as indicated in the legend. Note that the x axis is plotted in log scale.



Figure 8: Field sites used for validation of the estimated global nearshore slopes.




Figure 9: Linear regression between globally predicted nearshore slopes and nearshore slopes observed from local data for all study sites. Note that both axis are in log scale.




Figure 10: Global map of $\frac{R_{calc}}{R_{100}}$ where red colours indicate locations where the $\tan(\beta_{sf}) = 1:100$ assumption underestimates the coastal recession while blue colours locations where it overestimates it. Note that the colourbar uses a log scale.




Figure 11: Median coastal recession in meters, calculated per region for the sandy transects assuming 0.5 m of SLR across the global coastline. The vertical dashed line indicates the recession of 50 m, which would be the result if a 1/100 nearshore slope was assumed.