

# A multi-sensor satellite-based archive of the largest SO<sub>2</sub> volcanic eruptions since 2006

Pierre-Yves Tournigand<sup>1</sup>, Valeria Cigala<sup>2</sup>, Elzbieta Lasota<sup>1</sup>, Mohammed Hammouti<sup>3</sup>, Lieven Clarisse<sup>4</sup>, Hugues Brenot<sup>5</sup>, Fred Prata<sup>6</sup>, Gottfried Kirchengast<sup>7</sup>, Andrea K. Steiner<sup>7</sup>, Riccardo Biondi<sup>1</sup>

5 <sup>1</sup>Dipartimento di Geoscienze, Università degli Studi di Padova, Italy

<sup>2</sup>Ludwig-Maximilians-Universität München, Germany

<sup>3</sup>Politecnico di Milano, Italy

<sup>4</sup>Spectroscopy, Quantum Chemistry and Atmospheric Remote Sensing (SQUARES), Université Libre de Bruxelles, Belgium

10 <sup>5</sup>Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

<sup>6</sup>AIRES Pty Ltd: Mt Eliza, Victoria, Australia

<sup>7</sup>Wegener Center for Climate and Global Change, University of Graz, Austria

*Correspondence to:* Riccardo Biondi (riccardo@biondiriccardo.it)

**Abstract.** We present a multi-sensor archive collecting spatial and temporal information about volcanic SO<sub>2</sub> clouds generated by the eleven largest eruptions of this century. The detection and monitoring of volcanic clouds are an important topic for aviation management, climate issues and weather forecasts. Several studies focusing on single eruptive events exist, but no archive available at the moment combines quantitative data from as many instruments. We archived and collocated the SO<sub>2</sub> vertical column density estimations from three different satellite instruments (AIRS, IASI and GOME-2), atmospheric parameters as vertical profiles from the Global Navigation Satellite Systems (GNSS) Radio Occultations (RO) and the cloud top height and aerosol type from the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP). Additionally, we provide information about the cloud top height from three different algorithms, and the atmospheric anomaly due to the presence of the cloud. The dataset is gathering 206 days of SO<sub>2</sub> data, collocated with 44180 backscatter profiles and 64764 radio occultation profiles. The modular structure of the archive allows an easy collocation of the datasets according to the users' needs and the cross-comparison of the datasets shows different consistency of the parameters estimated with different sensors and algorithms, according to the sensitivity and resolution of the instruments. The data described here will be published with a DOI after final acceptance of this manuscript (Tournigand et al., 2020a, <http://doi.org/10.5880/figeo.2020.016>). During the discussion period, the data are accessible via this temporary link: <http://pmd.gfz-potsdam.de/panmetaworks/review/0f85d699707efcdc567765bd0dafaadf94b6df5a531f310167f7e974ea803bf>.

## 30 1. Introduction

Volcanoes around the world are a constant source of gaseous emissions. Both passively, during quiescent times, and actively, during eruptions (Robock, 2015; Carn et al., 2017). The most abundant gas species emitted are water (H<sub>2</sub>O), carbon dioxide (CO<sub>2</sub>) and sulfur dioxide (SO<sub>2</sub>), while less abundant ones include hydrogen (H<sub>2</sub>), hydrogen sulfur (H<sub>2</sub>S), hydrochloric acid (HCl) and carbon monoxide (CO) (Williams-Jones and Rymer, 2015). Once emitted into the atmosphere, some of these gases can react and transform, for example, SO<sub>2</sub> transforms into H<sub>2</sub>SO<sub>4</sub>. As with all volcanic eruption products, the gases emitted and the related aerosols can pose hazards to people as well as the environment (Williams-Jones and Rymer, 2015). Moreover, they can be responsible for regional and global climatic effects, depending on the latitude of the volcano, the altitude reached by the eruptive column and subsequent volcanic cloud (Robock, 2000; Robock, 2015; Williams-Jones and Rymer, 2015; Carn et al., 2016). In terms of global climatic impact, SO<sub>2</sub> injections into the stratosphere are of the greatest significance. The reason is that in the stratosphere the gas and subsequent aerosol can remain suspended for months to years and hence be transported around the globe affecting the absorption of both short- and longwave radiation (Robock, 2015). The duration and spatial spreading of emitted gases and aerosols in the atmosphere also depend on the erupted mass of volcanic material and the duration of the emission.

45 According to the Global Volcanism Program (GVP) of the Smithsonian Institute, an average of 55 to 88 eruptions  
(excluding permanent and semi-permanent degassing) have occurred worldwide each year since 1994. The eruptions  
display variability in their eruptive style (e.g., effusive or explosive), magma composition, the energy of the eruption,  
and the amount, type and size of the ejected material. To compare and characterize different eruptive events in an  
objective manner, the Volcanic Explosivity Index (VEI) was created. The VEI was introduced in 1982 by Newhall  
and Self (1982), drawing inspiration from the Richter scale for earthquake's magnitude, to provide a relative, semi-  
50 quantitative measure of the explosiveness of volcanic eruptions. The VEI classification, divided into categories from  
0 to 8, is based mainly on measures of magnitude, in terms of total ejecta volume, and/or intensity, in terms of mass  
flux or eruption plume height, depending on data availability (Newhall and Self, 1982; Houghton et al., 2013). The  
VEI has its limitations, nevertheless, it is being used extensively to provide an eruption descriptor that is  
understandable by researchers and policy-makers alike (Houghton et al., 2013).

55 The size of an eruption can, however, be significantly different when considering gas emissions only (Carn et al.,  
2016). Thus, a different eruption size classification can be outlined by the sulfur input into the atmosphere (Schnetzler  
et al., 1997; Robock, 2015; Carn et al., 2016). In 1997, Schnetzler et al. proposed the Volcanic Sulfur Dioxide Index  
(VSI) based on SO<sub>2</sub> retrievals performed with the Total Ozone Mapping Spectrometer (TOMS) onboard the Nimbus  
7 satellite. Medium VEI eruptions (e.g. VEI 4 events) can be characterized by the emission of a large quantity of  
60 tephra with respect to the quantity of SO<sub>2</sub> (e.g. the 2008 Chaiten eruption) or by the emission of a larger amount of  
SO<sub>2</sub> than tephra (e.g. the 2011 Nabro eruption) as shown in Carn et al. (2016). Using an improved SO<sub>2</sub> emissions  
retrieval approach and including less energetic events, Carn et al. (2016) found a broader range of SO<sub>2</sub> emission per  
VEI and a weaker first-order correlation. These findings suggest that the intensity and volcanic plume altitude are  
more relevant parameters for consideration in modelling SO<sub>2</sub> emissions and their climate impact (Robock, 2000; Carn  
65 et al., 2016).

The above-mentioned GVP provides the most extensive catalogue of historical eruptions with information related to  
both volcanoes and their eruptions. This catalogue is a first-hand source of information when starting an investigation  
of a given volcano and a given style of eruption. Similarly, the Large Magnitude Explosive Volcanic Eruptions  
(LaMEVE) dataset includes data such as the magnitude of the event, the bulk volume, the tephra fallout volume,  
70 column height of Quaternary (from 2.58 Ma ago to present) eruptions with VEI ≥ 4 (Brown et al., 2014). Other datasets  
are available including data on geochemical composition (e.g. Turner et al., 2009), or acoustic acquisitions (e.g. Fee  
et al., 2014), although these types of database are generally limited to ancillary information, a specific volcano, a  
specific time window or a specific instrument (e.g. de Moor et al., 2017).

75 Previous studies focusing on a single or a few eruptions are based on personal data collections or project collaborations  
and this makes it difficult for data comparison and studies with new techniques or algorithms. The eruptions of Okmok  
and Kasatochi in 2008 were the focus of a special issue (JGR Atmospheres, 2018) collecting 27 articles studying the  
clouds with a large number of the available remote sensing platforms and algorithms. The Sarychev Peak eruption in  
2009 was also well studied (e.g., Carn and Lopez, 2011; Kravitz et al., 2011; Rybin et al., 2011; Doeringer et al.,  
2012). The Eyjafjallajökull 2010 eruption affected the economy and social life of Europe and beyond, changing the  
80 rules of air traffic management and the volcanic cloud was subject of a number of studies (e.g., Flentje et al., 2010;  
Marenco et al., 2011; Prata and Prata, 2012; Stohl et al., 2012). The Grímsvötn eruption in 2011 was quite interesting  
from a scientific point of view because the SO<sub>2</sub> cloud was separated from the ash cloud (Moxnes et al., 2014) and  
different researchers studied it (e.g., Flemming and Inness, 2013; Marzano et al., 2013; Cook et al., 2014; Prata et al.,  
2017). The Nabro 2011 eruption was the subject of an interesting discussion regarding the direct intrusion to the  
85 stratosphere of the volcanic cloud (e.g., Bourassa et al., 2012; Clarisse et al., 2014; Fromm et al., 2014; Biondi et al.,  
2017) and the Puyehue Cordón Caulle, erupting in the same period, was of interest because the cloud moved around  
the globe (e.g., Bignami et al., 2014; Griessbach et al., 2014; Theys et al., 2014; Biondi et al., 2017). The Calbuco  
2015 eruption (Marzano et al., 2018; Lopes et al., 2019) was widely studied especially in connection to its impact on  
the Antarctic ozone hole (Ivy et al., 2017; Stone et al., 2017; Zhu et al., 2018; Zuev et al., 2018). The rest of the  
90 volcanic clouds, such as the ones produced by Merapi 2010 (Picquout et al., 2013), Tolbachik 2012 (Telling et al.,  
2015), and Kelut 2014 (Kristiansen et al., 2015; Vernier et al., 2016) also received some attention from the scientific  
community.

95 Considering SO<sub>2</sub> emissions, several datasets and inventories are available and updated over time, but generally include  
daily or yearly total emissions per volcano or per eruption. Ge et al. (2016) compiled an inventory for daily SO<sub>2</sub>  
emissions in the time frame 2005-2012 including global volcanic eruptions but also eight persistently degassing  
volcanoes retrieved by the Ozone Monitoring Instrument (OMI) onboard the Aura satellite. Carn et al. (2017)  
implemented it including OMI retrievals from 2005 to 2015 of emissions related to passive degassing. The most  
100 updated and complete dataset up to now is the Multi-Satellite Volcanic Sulfur Dioxide Database Long-Term L4 Global  
(MSVOLSO2L4) compiled by Carn (2019). The dataset provides “SO<sub>2</sub> mass loadings for all significant global  
volcanic eruptions detected from space since October 1978” to present (Carn, 2019). The MSVOLSO2L4 includes  
ancillary information about the volcanoes, such as the name and location of the volcano, as well as information about  
the eruptions, for example, start and end date, and VEI. This information is retrieved from the GVP database. The  
dataset also reports the observed plume altitude (in kilometres) where known. Otherwise, an estimated plume altitude  
105 above vent depending on eruption type, and the measured SO<sub>2</sub> mass in kilotons (= 1000 metric tons) is provided (Carn  
et al., 2016; Carn, 2019). The above-mentioned datasets provide important information for users mainly needing to  
assess the climatic impact of SO<sub>2</sub> from volcanic sources, however, none of them allows for mapping the SO<sub>2</sub> emissions  
and related altitude estimations in space and time and thus the direct testing and comparison of new models and  
techniques, like GNSS RO, for example. We think it is important to provide a complementary multi-satellite archive  
110 covering the largest eruptive events and their cloud development all around the world in order to facilitate the access  
to such data for future studies.

In this study, we present the first database predicated on satellite data, reporting:

1. SO<sub>2</sub> retrievals from Atmospheric Infrared Sounder (AIRS), Infrared Atmospheric Sounding Interferometer  
(IASI), and the Global Ozone Monitoring Experiment 2 (GOME-2). The data from the three sensors provide  
115 horizontal and temporal information on SO<sub>2</sub> concentrations;
2. SO<sub>2</sub> cloud altitude estimations from IASI, the Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP)  
backscatter and the Global Navigation Satellite System (GNSS) Radio Occultation (RO);
3. the cloud aerosol subtype information from CALIOP;
4. atmospheric properties such as temperature, pressure and humidity from GNSS RO profiles.

120 The information is provided for eruptions, after July 2008, classified by GVP as VEI 4 or larger and with an SO<sub>2</sub> total  
mass loading greater than 0.05 Tg. At the time of the archive preparation no eruption after 2016 had yet been classified  
as VEI 4 or greater. The selected volcanoes and relative eruptions are: Okmok 2008; Kasatochi 2008; Sarychev 2009;  
Eyjafjallajökull 2010; Merapi 2010; Grímsvötn 2011; Nabro 2011; Puyehue-Cordón Caulle 2011; Tolbachik 2012;  
Kelut 2014; Calbuco 2015. These are the most significant eruptions over the period 2006-2018. Further information  
on these eruptions can be found in Table 1.

125 To the best of our knowledge, there is no database to date collecting SO<sub>2</sub> maps and volcanic cloud altitude estimations  
from several instruments, cloud aerosols subtypes and atmospheric properties for explosive eruptions. Accurate  
knowledge on volcanic SO<sub>2</sub> cloud concentration and altitude as well as their spatial and temporal evolution, is of  
crucial importance in the investigation of an eruption’s climatic impact. Thus, we believe that the database presented  
here will help current and future investigations as well as support the development of more accurate retrieval  
130 methodologies.

## 2. Instrument and retrieval description

A summary of the instruments’ characteristics together with parameters provided in this work and references to the  
algorithms are reported in Table 1. In this archive, each sensor measuring SO<sub>2</sub> amounts measures the partial column  
density, due to their own limitations (see section 4.3) (Brenot et al., 2014). We here use the terms Vertical Column  
Density (VCD) to refer to this partial column density.  
135

## 2.1. AIRS

The Atmospheric Infrared Sounder (AIRS) is a cross-track hyperspectral instrument onboard the polar-orbiting satellite Aqua launched in June 2002 with an ascending orbit equator crossing local time at 13:30. AIRS completely covers the full globe two times per day with a swath of 1650 km and spatial resolution of 13.5 km x 13.5 km at nadir and 41 km x 21 km at high latitudes (Susskind et al., 2003). Each orbit is divided in granules, where a granule is a portion of AIRS orbit containing 6 minutes (2250 km x 1650 km) of data, which is officially defined by the National Aeronautics and Space Administration (NASA). The SO<sub>2</sub> pixels are identified using infrared channels centered at the 7.3 μm absorption peak relying on the correlation between the measured spectrum and a reference spectral shape. The amount of SO<sub>2</sub> in each pixel is computed by a least-squares procedure based on an off-line radiative transfer model. This technique performs well for SO<sub>2</sub> reaching high tropospheric altitudes or the stratosphere where the water vapor content is negligible. Comparisons with other techniques (Carn et al., 2016; Carn et al., 2017) show an agreement within 10–30% and a typical retrieval error for a single AIRS pixel of about 6 Dobson Unit (DU) (Prata and Bernardo, 2007).

## 2.2. IASI

The Infrared Atmospheric Sounding Interferometer (IASI) is a Fourier transform instrument onboard the near-polar sun-synchronous orbiting satellites Metop-A and Metop-B, respectively, launched in October 2006 and September 2012 with an ascending orbit equator crossing local time at 9:30. IASI completely covers the full globe two times per day with a swath of 2200 km and a spatial resolution of 12 km at nadir (Clerbaux et al., 2009). The SO<sub>2</sub> retrieval is based on a brightness temperature difference in the SO<sub>2</sub> ν<sub>3</sub> band centered at 7.3 μm (Clarisse et al., 2012) which is converted to SO<sub>2</sub> concentration integrated along the vertical axis the VCD using look-up tables and operational profiles of pressure, temperature and humidity. The retrieval of VCD assumes that all SO<sub>2</sub> is located at particular atmospheric layers (5, 7, 10, 13, 16, 19, 25 or 30 km above sea level) providing different estimations at different altitudes. It has a detection limit of around 0.5 DU at the tropopause, which increases for decreasing altitude (depending on the amount of water vapour in the atmosphere). For plumes above 500 hPa (about 5.5 km) the algorithm has a theoretical uncertainty between 3-6%. A second algorithm (Clarisse et al., 2014) is applied to compute the SO<sub>2</sub> cloud altitude with an accuracy of about 2 km for plumes below 20 km. The algorithm exploits the fact that the SO<sub>2</sub> ν<sub>3</sub> band interferes with strong water vapour absorptions, and that these interferences, by virtue of the vertical water vapour profile, have a strong dependency with height. Combining the two datasets, a single best-estimate VCD is obtained by interpolating the VCD columns of the first algorithm at the retrieved height.

## 2.3. GOME

The Global Ozone Monitoring Experiment 2 (GOME-2) is an ultraviolet-visible spectrometer, onboard of the Metop-A and Metop-B satellites, measuring solar light backscattered by the atmosphere or reflected by the Earth in nadir-viewing geometry with a swath of 1920 km and spatial resolution of 40 km x 80 km at nadir (Munro et al., 2006). The SO<sub>2</sub> VCD retrieval (Rix et al., 2012) is based on the strong SO<sub>2</sub> absorption between 240 and 400 nm and uses a differential optical absorption spectroscopy technique (Platt and Stutz, 2008). All measurements in the wavelength ranging from 315 to 326 nm are fitted to laboratory absorption data of SO<sub>2</sub> and converted to VCD with an air mass factor from radiative transfer models assuming hypothetical atmospheric layers representative of different scenarios of emissions. The SO<sub>2</sub> VCD is provided for 3 atmospheric layers representative of different scenarios of emissions: low troposphere (~2.5 km above the surface), upper troposphere (~6 km) and lower stratosphere (~15 km). The volcanic emission measurement is facilitated by large SO<sub>2</sub> columns generally at high altitudes (free-troposphere to lower stratosphere). However, for large SO<sub>2</sub> columns (typically >50 DU) the absorption tends to saturate leading to a general underestimation and directly affecting the product accuracy. For most volcanoes, there is no ground-based equipment to measure SO<sub>2</sub> during the eruption and the validation approach is usually a cross-comparisons with other satellite products. The O3M SAF validation report (Theys and Koukouli, 2015) shows that GOME-2 SO<sub>2</sub> product reaches the target/optimal accuracy of 50%/30% respectively. It is important to notice that the SO<sub>2</sub> retrievals from GOME-2 are also affected by clouds and instrumental noise especially at high solar zenith angles. These limitations have been filtered in the data used in this work, according to the criteria shown by Brenot et al. (2014).

## 2.4. CALIOP

185 The Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) is an instrument onboard the polar-orbiting Cloud-  
Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO). To estimate the Volcanic Cloud (VC) top  
altitude and validate the cloud top estimation from GNSS RO, we have used the Level 1 total attenuated backscatter  
at 532 nm (CAL\_LID\_L1 Version 4) with a vertical resolution of 60 m and horizontal resolution of 1 km between 10  
and 20 km of altitude, and a vertical resolution of 180 m and horizontal resolution of 1.67 km above 20 km (Winker  
190 et al., 2009). To extract the corresponding aerosol type we have used the Level 2 Vertical Feature Mask (Winker et  
al., 2009) product version 4.20. The CALIOP does not allow SO<sub>2</sub> measurements or estimation (it provides estimations  
of dust, elevated smoke, volcanic ash and sulfate) and the CALIOP classification algorithm do not include the volcanic  
ash type below the tropopause level (Kim et al. 2018) making difficult to distinguish the volcanic ash from other  
aerosol types in the lower troposphere. For these reasons, the selected CALIOP backscatter is collocated with the SO<sub>2</sub>  
estimation from AIRS, IASI and GOME-2 and this combination provides a complete information on the content and  
195 vertical structure of the cloud.

## 2.5. GNSS RO

The Global Navigation Satellite Systems (GNSS) Radio Occultation (RO) is an active limb sounding technique which  
uses radio signals transmitted by a GNSS satellite and received by a Low Earth Orbit satellite, where the atmosphere  
is vertically scanned due to the relative motion of the two satellites. The signal, travelling through the atmosphere,  
200 is refracted and bent due to the vertical gradient of atmospheric density. The effect of the atmosphere is represented by  
a bending angle, from which refractivity and density is retrieved. Refractivity in the neutral atmosphere depends  
mainly on temperature, pressure, and water vapour pressure. Information about the vertical structure of the troposphere  
and stratosphere is provided (Kursinski et al., 1997). The vertical resolution of the RO ranges between 100 m in the  
upper troposphere to about 500 m in the lower stratosphere at low/mid-latitudes (Zeng et al., 2019), while the  
205 horizontal resolution can range from about 50 km in the troposphere to 200-300 km in the stratosphere (Kursinski et  
al., 1997). In this archive we use the RO bending angle, refractivity, temperature, pressure and specific humidity  
profiles processed by the Wegener Center for Climate and Global Change (WEGC) with the Occultation Processing  
System (OPS) version 5.6 (EOPAC Team, 2019). We also provide the bending angle anomaly which is proven to be  
an efficient parameter to reveal the impact of the VC on the atmospheric structure (Biondi et al., 2017; Cigala et al.,  
210 2019; Stocker et al., 2019) because perturbations in the vertical structure are seen in the bending angle profile as  
anomalous peaks, specifically at the volcanic cloud top. The way this anomaly is computed is detailed in section 4.1.  
The RO profiles are obtained using a combination of geometric optics and wave optics retrieval (Angerer et al., 2017),  
with transition below the tropopause. The retrieval is based on orbit information and amplitude and phase data from  
the University Corporation of Atmospheric Research/COSMIC Data Analysis and Archive Center (UCAR/CDAAC)  
215 collected from the following RO missions: the CHALLENGING Minisatellite Payload (CHAMP) (Wickert et al., 2001),  
the Satélite de Aplicaciones Científicas (SAC-C) (Hajj et al., 2004), the Gravity Recovery And Climate Experiment  
(GRACE-A) (Beyerle et al., 2005), the FORMOSAT-3/COSMIC (Anthes et al., 2008), and the EUMETSAT/METOP  
missions (Luntama et al., 2008). The accuracy of the GNSS RO is 0.2 K in terms of temperature and 0.1% in terms of  
refractivity and the data from the different mission are very consistent (Scherllin-Pirscher et al., 2011), so there is no  
220 need of inter-calibration or homogenization (Foelsche et al., 2011).

## 3. Data description

The archive (Tournigand et al., 2020a) consists of two sets of files, the daily files and the eruption files, i.e., one file  
per eruption including all collected information. Thus, for each eruption listed in Table 2 the user can choose to access  
one single day or to the whole eruptive period depending on the user's demand. The number of days covered by the  
225 archive for each eruption depends on the SO<sub>2</sub> detection availability from AIRS, IASI and GOME-2. Also, the variables  
available from one day to another may differ according to SO<sub>2</sub> detection results and instruments availability. Each file

is in NetCDF-4 format and file names are self-explanatory with daily files following the format *volcanoname\_year\_month\_day* and the eruption files following the format *volcanoname*. As an example, a user who wishes to access all available data corresponding to the Sarychev volcano on 12 of June 2009 will have to look for the file *Sarychev\_2009\_06\_12.nc*. The organization of both file types is described hereafter for each instrument, the main information is provided in Table 3 and all the details are provided in the supplementary material including the geographical coverage of each VC (Figures S1-S11).

### 3.1. AIRS

AIRS data are organized in the same way in both file types. It consists of 4 variables namely AIRS\_lat, AIRS\_lon, AIRS\_date and AIRS\_SO2 respectively containing the latitude (°N), longitude (°E), date and time (POSIX time, number of seconds elapsed since 00:00:00 UTC 1st of January 1970) of each granule and their SO<sub>2</sub> VCD (DU). The variables AIRS\_lat, AIRS\_lon and AIRS\_SO2 are matrices with columns corresponding to the different granules. By selecting one column, the user can find each data point of the corresponding granule. The AIRS\_date variable, on the other hand, is a line vector with elements reporting the date and time of each granule. Only data points with SO<sub>2</sub> values higher than 0 DU have been included in the archive thus explaining the different amount of points from one granule to another.

### 3.2. IASI

IASI data are organized in the same way for both file types and are composed of 5 variables, IASI\_lat, IASI\_lon, IASI\_date, IASI\_SO2 and IASI\_height respectively containing the latitude (°N), longitude (°E), date and time (POSIX time), SO<sub>2</sub> (DU) and cloud altitude (m). The date variable consists of a line vector with elements corresponding to each scanning line. Similarly, the other variables are matrices with columns corresponding to the different scanning lines and rows to the data points of the given scanning line having an SO<sub>2</sub> content higher than 0 DU.

### 3.3. GOME-2

GOME-2 data's organization is identical in both file types. GOME-2 is composed of 6 variables, GOME\_lat, GOME\_lon, GOME\_date, GOME\_SO2\_1, GOME\_SO2\_2, GOME\_SO2\_3 respectively corresponding to the latitude (°N), longitude (°E), date and time (POSIX time), low troposphere SO<sub>2</sub> vertical column density (DU), mid-troposphere SO<sub>2</sub> vertical column density (DU) and low stratosphere SO<sub>2</sub> vertical column density (DU). As for AIRS and IASI, the date variable corresponds to a line vector providing each scanning line's date. The rest of the variables correspond to matrices with scanning lines also separated in columns and the data points of those scanning lines distributed in rows. In the case of GOME-2, only data points having their three SO<sub>2</sub> vertical columns contents higher than 0 DU were included. Pixel with high Solar Zenith Angle (SZA) has also been filtered.

### 3.4. CALIOP

CALIOP's section of the archive contains 6 variables, CALIOP\_lat, CALIOP\_lon, CALIOP\_date, CALIOP\_filename, CALIOP\_height, CALIOP\_type respectively corresponding to the latitude (°N), longitude (°E), date and time (POSIX time), name of CALIOP file, estimated VC top altitude (m) and aerosol type. The latitude, longitude, date and file name variables are column vectors with each row corresponding to latitude, longitude and date data of the designated file in CALIOP\_filename variable. CALIOP\_height variable contains all the cloud top altitude estimations based on CALIOP L1 532 nm version 4.10 backscatter product. This variable is a matrix with each row corresponding to a CALIOP file and the three columns indicating to which instrument the CALIOP file is collocated (at ±0.2° and ±1h) with, AIRS (column 1), IASI (column 2) or GOME (column 3).

CALIOP\_type variable is read as a string containing the type of aerosol retrieved from the L2 Vertical Feature Mask (VFM) version 4.20. The L2 VFM CALIOP product subdivides the aerosols into 10 types. Four of those types are of interest for this archive: type 2, 6, 9 and 10 respectively corresponding to dust, elevated smoke, volcanic ash and sulfate/other. This variable is subdivided into as many rows as there are CALIOP files, 16 columns containing the aerosols values and three sections corresponding to three levels of altitude -0.5 to 8.2 km, 8.2 to 20.2 km and 20.2 to

30.1 km. In each altitude level, the presence of one or several cloud types is indicated by the presence of their reference number.

### 3.5. GNSS RO

275 GNSS RO data are organized in different ways in the two file types. In the daily files, the RO data are separated in  
different variables according to the instrument they are collocated with (AIRS, IASI or GOME-2) at  $\pm 0.2^\circ$  spatially  
and  $\pm 12$ h temporally. For each set of RO data collocated with a given instrument 10 variables are provided: latitude  
( $^\circ$ N), longitude ( $^\circ$ E), date (POSIX time), bending angle (rad), bending angle anomaly (%), temperature (K), pressure  
(Pa), refractivity (N-unit), specific humidity ( $\text{kg}\cdot\text{kg}^{-1}$ ) and volcanic cloud top altitude (m). Each variable is a matrix  
280 with each column corresponding to a RO profile and rows to the latitude dimension. Only the variable containing  
volcanic cloud altitude is a line vector with each element corresponding to a different RO profiles.

In the files containing the whole eruptive period, the RO data are not separated according to the instrument they are  
collocated with but compiled all together. Thus, the same 10 variables are provided as in daily files, each containing  
the totality of the RO profiles.

## 4. Quality control and data processing

### 285 4.1. RO data

The RO profiles included in this archive are collocated spatially at  $\pm 0.2^\circ$  and temporally at  $\pm 12$ h with data points from  
the volcanic aerosol maps provided by AIRS, IASI, and GOME-2 acquisitions.

#### 4.1.1 Climatology

290 The RO reference climatology for each area of interest is calculated based on  $5^\circ$  latitude bands using our dataset of  
RO profiles covering a period from 2001 to 2017. The averaging of all available RO profiles present within each  
latitude band provides the RO reference climatology.

#### 4.1.2 Anomaly calculation

295 The bending angle (BA) anomaly integrated into this archive is calculated by subtracting the RO reference climatology  
profile from the individual RO BA profile and normalizing with respect to the reference climatology profile (Eq 1),  
following the methodology described in Biondi et al. (2017). The resulting anomaly displays variations when the  
bending angle differs from the climatology. Such variations indicate a change of atmospheric properties and are used  
to identify related atmospheric features. The presence of volcanic clouds in the atmosphere generates a prominent  
peak in the BA anomaly profile.

$$\alpha = \left( \frac{BA - BA_{clim}}{BA_{clim}} \right) \cdot 100 \quad (1)$$

300 Where  $\alpha$  is the bending angle anomaly, BA the individual bending angle profile and  $BA_{clim}$  the BA climatology profile.

#### 4.1.3 Peak detection

305 The peak detection of bending angle anomaly profiles was automatically done using a customized Matlab algorithm.  
This algorithm, further developed after Cigala et al. (2019), identifies all the peaks displaying a variation larger than  
4.5% in the bending angle anomaly profile with respect to local minimums. Only the peaks having their maximum  
value between 10 and 22 km of altitude are kept. Peaks vertically spreading over more than 8 km have been removed.  
Finally, amongst the remaining peaks, the lowest altitude one is selected as a cloud top altitude.

## 4.2. CALIOP

### 4.2.1 Cloud top automatic detection

310 For each L1 532 nm version 4.10 CALIOP backscatter product collocated with RO profiles, an automatic cloud top  
detection was performed using a customized Matlab algorithm. The collocation thresholds were kept at  $\pm 0.2^\circ$  and  
 $\pm 12$ h between RO profiles and CALIOP backscatter products to provide cloud top altitudes consistent between the  
instruments. The first step of the cloud top detection procedure consists of cropping the CALIOP backscatter image  
315 according to the collocated RO profile position at  $\pm 14^\circ$  in latitude and  $\pm 80^\circ$  in longitude. The objective of this first  
step is to focus the processing on a restricted zone around the position of the collocated RO profile in order to save  
computational time. These latitude and longitude ranges are based on series of tests performed on backscatter images  
and correspond to the best compromise between image size reduction and loss of volcanic cloud information.  
Threshold backscatter values are then implemented to remove the noise outside of the range  $3 \times 10^{-2}$  -  $7 \times 10^{-4}$   $\text{km}^{-1} \cdot \text{sr}^{-1}$   
320 to which volcanic clouds correspond. One median filter (4x3 pixels) and two Wiener filters (4x3 and 2x2 pixels) are  
then successively applied to the resulting backscatter image to reduce the noise within the threshold range. Below 10  
km altitude, the RO bending angle anomaly is noisy due to the presence of moisture. Thus we only included in this  
archive volcanic clouds with a top altitude above 10 km. The following step in the volcanic cloud top determination  
is then to remove image information below 10 km which also removes a significant part of meteorological clouds  
increasing the volcanic cloud top altitude detection accuracy. The next step of CALIOP data processing is to identify  
325 remaining groups of pixels (or clusters) within the image. The Matlab connected components finder is set in this  
customized algorithm to keep only the clusters combining more than 300 pixels. Amongst these selected clusters the  
nine biggest ones are kept for the final processing stage. Based on all the collocations between CALIOP and the GNSS  
RO, a statistical analysis of volcanic clouds defined by the CALIOP cloud mask and collocated with the RO within 2  
hours, has shown that the volcanic clouds are usually thinner than meteorological clouds with an aspect ratio lower  
330 than 0.09. According to this result, the final stage of the algorithm consists of distinguishing clusters corresponding  
to volcanic features from the ones corresponding to meteorological clouds setting the higher limit of the aspect ratio  
to 0.09. Finally, the remaining clusters' top altitudes are measured and an average value calculated and saved in the  
archive as an estimate of the volcanic cloud top altitude.

### 4.2.2 Cloud type detection

335 The cloud type detection was performed using the L2 Vertical Feature Mask (VFM) CALIOP product. These VFM  
products were collocated at  $\pm 0.2^\circ$  and  $\pm 1$ h with AIRS, IASI and GOME-2 for the purpose of such data is to confirm  
the presence of certain aerosols types simultaneously with  $\text{SO}_2$ . Among the different cloud types available in VFM  
products the types 2, 6, 9 and 10 were of particular interest for this archive since they respectively correspond to dust,  
elevated smoke, volcanic ash and sulfate/other. For each CALIOP file of the archive, the aerosol subtype was extracted  
340 using a customized Matlab routine. This Matlab algorithm reads the VFM data, detects the matching latitude/longitude  
points of all the CALIOP track collocated with AIRS, IASI and GOME-2 and subsets the latitude and longitude array  
data based on the chosen spatial window of  $2^\circ$ . The algorithm then extracts the feature sub-type of interest as a function  
of the altitude. The final output is subdivided in three levels of altitude -0.5 to 8.2 km, 8.2 to 20.2 km and 20.2 to  
30.1km.

## 345 4.3. Uncertainties

This archive combines five different approaches of volcanic cloud detection and each type of measurement/instrument  
has its own uncertainties. The comparison between different instruments, always faces uncertainties due to the spatial  
and temporal collocation (section 3) and to their spatial resolution (section 2). The difference in cloud top estimations  
can be partly explained by the different sensitivities and vertical resolution of the instruments. In addition, the number  
350 of collocations between RO and CALIOP is much smaller than for RO-IASI and IASI-CALIOP, respectively. The  
cloud top height estimation for eruptions with a large number of collocations (Calbuco, Kasatochi, Nabro and Sarychev  
Peak) is in general consistent within the techniques. For AIRS, IASI and GOME-2 the uncertainty depends on many  
parameters, such as thickness of the volcanic cloud, amount of aerosols and one of the most important the unknown



355 volcanic cloud altitude. Thus, the error is case dependent and a general value of measurement uncertainty cannot be provided. Furthermore, the measurement noise of instruments increases over time due to instrument degradation (Lang et al., 2009; Dikty et al., 2010). However, error budgets of AIRS and IASI can be respectively found in the studies by Prata and Bernardo (2007) and Clarisse et al. (2012), while an uncertainty analysis of GOME-2 is provided by Rix et al. 2012 in the case of the 2010 Eyjafjallajökull's volcanic eruption.

## 5. Data cross-comparison

360 Over the past decades, satellite data have proven efficient in volcanic cloud detection through a variety of techniques. Those data are essential in the study of the spreading of such clouds on a global scale but are scattered between the different agencies in charge of processing them. This archive gathers satellite data covering 10 VEI 4 and 1 VEI 5 eruptions from 2008 to 2016 with a total of 223 days of data coverage (Table 4).

365 This archive is organized in different sections (Figure 1) with each instrument estimation separated from each other. Several parameters are included within the dataset, such as SO<sub>2</sub> VCD and cloud top altitude, to allow cross-correlation between the different retrieval algorithms. The database allows the quick visualization of AIRS, IASI, GOME-2, CALIOP and RO data at a given date and time as well as the collocation of any instrument data points with another one. In order to illustrate the use of this archive, we extracted two test cases (Figure 2). The first case (Figure 2a) is the 2008 eruption of Kasatochi volcano for which we selected the 9<sup>th</sup> of August as reference. The second case (Figure 370 2b) is the 2009 Sarychev Peak eruption for which we selected the 12<sup>th</sup> of June as reference. In both cases we considered SO<sub>2</sub> values larger than 3 DU from AIRS, IASI and GOME-2 for 24 hours, RO profiles collocated within  $\pm 0.2^\circ$  and  $\pm 12$ h and CALIOP tracks collocated within  $\pm 12$ h.

375 In the case of Kasatochi, we selected 4178 AIRS, 1241 IASI and 56 GOME-2 data points with SO<sub>2</sub> VCD larger than 3 DU, 379 CALIOP profiles from 7 different tracks (Figure 2a, blue circles) within  $\pm 12$ h and 100 RO profiles (Figure 2a, red circles) collocated within  $\pm 0.2^\circ$  and  $\pm 12$ h. In the case of Sarychev, 1070 AIRS, 209 IASI and 41 GOME-2 data points, 261 CALIOP profiles from 3 different tracks and 54 RO profiles have been selected with the same criteria. Due to the modular archive structure reported in Figure 1, the user can easily select different time frames, different SO<sub>2</sub> thresholds and collocation period range to be adapted to any purpose.

380 We have manually verified the correct functioning of the algorithm which collocates the different instruments. We randomly selected several days from different eruptions and compared the date, time and coordinates of the acquisitions, then compared the results with those ones automatically provided by the algorithm. Additionally, we have used a visual validation method for all the samples plotting the SO<sub>2</sub> cloud superimposed to the CALIOP tracks and the RO tangent points. As for the cloud top height, we have collocated the RO and CALIOP estimations with the closest IASI pixel and compared the corresponding values. In Table 5 we report the number of collocations per pairs 385 of instruments with the averaged difference between the estimation. Depending on the eruption the different techniques can give variable performances, for example, the estimations of RO and CALIOP for the Eyjafjallajökull, Kasatochi and Grímsvötn were very close (average difference of 0.3 km, 0.9 km, 1.3 km respectively) while they were large for Calbuco (4.2 km). The difference in cloud top estimations can be partly explained by the sensitivity of the RO to the density of the atmosphere, denser clouds can be detected more likely than less dense clouds (Tournigand et al., 2020b). This reason, summed up to the uncertainties reported in the section 4.3, can contribute to the different 390 biases of the cross-comparisons. In general, the cloud top height estimation for eruptions with a large number of collocations (Calbuco, Kasatochi, Nabro and Sarychev Peak) are consistent within the techniques.

## 6. Results

395 We have collected 4535062 GOME-2 scanning lines covering 182 days, 336399 IASI scanning lines covering 172 days, 865 AIRS granules covering 122 days, 44180 CALIOP profiles covering 152 days, and 64764 RO profiles covering 194 days collocated to the VC emitted by 11 different eruptions with VEI  $\geq 4$ . The Kasatochi eruption has the best data coverage, Sarychev has the longest period of coverage (36 days), and Puyehue-Cordón Caulle is the only one not being covered by CALIOP (due to a technical issues on that period).

400 The archive allows to collocate data from five instruments working at different frequencies (Table 1), three of them (IASI, CALIOP and RO) able to provide information to develop an algorithm for the cloud top height estimation (Table 5). Part of this archive has been used in the past to develop an algorithm estimating the cloud top height by using the RO bending angle anomaly (Biondi et al., 2017; Cigala et al., 2019) and to understand the possible overshooting of the Nabro eruption in the stratosphere (Biondi et al., 2017).

405 The user is free to compare the SO<sub>2</sub> estimation from three different algorithms, to check the cloud structure by downloading the collocated CALIOP sub-tracks, and to analyse the impact of the volcanic cloud on the atmospheric vertical structure with the RO profiles. An example of use of this archive is shown in Figure 3. We collocated, the IASI SO<sub>2</sub> estimation the 12<sup>th</sup> of August 2008 (Figure 3e) and the AIRS SO<sub>2</sub> estimation the 11<sup>th</sup> of August 2008 (Figure 3l) of the Kasatochi VC, with the RO and CALIOP profiles. The map visually provides the SO<sub>2</sub> distribution and the position of the RO and CALIOP profiles. Panels d) and i) show the vertical distribution of the aerosol according to the CALIOP algorithm: on the 12<sup>th</sup> of August there was volcanic ash together with sulphate up to 13 km of altitude, while on the 11<sup>th</sup> of August the sulphate was prevailing and the cloud top was slightly lower (about 12 km). We then report the vertical profiles of temperature anomaly (c and h) and water vapour anomaly (a and f) when compared to the climatological values of the area. The behaviour of the two parameters are similar, but on the 12<sup>th</sup> of August the temperature anomaly in the lower troposphere is colder, and the water vapour anomalies are larger in the lower troposphere and at the cloud top layer. Finally, the bending angle anomaly (b and g) according to the algorithm reported in section 4.1.3, shows a cloud top height of 11.2 km (the 12<sup>th</sup> of August) against about 13 km from CALIOP, and a cloud top height of 11.9 km (the 11<sup>th</sup> of August) slightly lower than the detection from CALIOP. This comparison shows that the cloud structure was steady over the time with similar characteristics confirmed by four different instruments.

## 420 7. Data availability

The raw CALIOP data can be found at <https://urs.earthdata.nasa.gov/>.

425 The archive consists of daily files and “eruption” files. For each eruption, we provide access to single daily files or to one file for the whole eruptive period. The files of any eruption are compressed (.zip) NetCDF-4 format (including the daily and whole eruptive period) together with two pdfs (Supplement) describing the file structure. The file names are self-explanatory with daily files following the format *volcanoname\_year\_month\_day.nc* and the eruption files following the format *volcanoname.nc*. As an example, a user who wishes to access the data corresponding to Kasatochi volcano on 11<sup>th</sup> of August 2008 will have to look for the file *Kasatochi\_2008\_08\_11.nc*. In case the user wants all the available data for the Kasatochi eruption, they will have to look for the file *Kasatochi.nc*. The data structure of daily files and volcano files is reported in the Supplement. The data described here will be published with a DOI after final acceptance of this manuscript (Tournigand et al., 2020a, <http://doi.org/10.5880/figeo.2020.016>). During the discussion period, the data are accessible via this temporary link: <http://pmd.gfz-potsdam.de/panmetaworks/review/0f85d699707efcdc567765bd0dafaaadf94b6df5a531f310167f7e974ea803bf>.

## 430 8. Summary and conclusions

435 This paper presents the first comprehensive archive with quantitative information on large SO<sub>2</sub> volcanic clouds since 2006. We collected three different datasets of volcanic SO<sub>2</sub> detection from AIRS, IASI and GOME-2 instruments and co-located the detected pixels with the CALIOP and the GNSS RO products to get information about the cloud vertical structure. The archive provides information about the SO<sub>2</sub> detection and retrieval (with 3 different algorithms), the cloud top height (with 3 different algorithms), the cloud aerosol type (CALIOP vertical mask feature reference), the atmospheric parameters (bending angle, refractivity, temperature, pressure and specific humidity) and the atmospheric change due to the presence of the volcanic cloud (bending angle anomaly). At present, there are no public archives of volcanic clouds which can be used as a reference for further studies and all the information is scattered in different locations and available under different conditions. The aim of this archive and this paper is to provide the users with a complete set of state-of-the-art data. The interest in volcanic clouds detection and monitoring is high and there are still some challenges like the accurate determination of the cloud top height and cloud density to be faced. This archive

445 will make available to the scientific community a relevant number of cases to develop and test new algorithms on, thereby contributing to improving the accuracy on the estimation of fundamental volcanic clouds parameters. The modular structure of the archive can be easily extended in the future to smaller eruptions (VEI<4) and to other SO<sub>2</sub> estimations, facilitating the inter/cross-comparison between algorithms, allowing the reconstruction of the cloud structure and dynamical characteristics and supporting the development of cloud dispersion models.

450 **Author contributions.** PYT and VC collected the data. LC provided the IASI dataset. HB provided the GOME-2 dataset. FP provided the AIRS dataset. GK and AS provided the WEGC GNSS RO dataset. PYT, VC and EL conceived the algorithm approach and wrote the code. MH elaborated the CALIOP VFM data. PYT, VC and RB structured and wrote the manuscript. RB conceived the idea, coordinated the team, supervised the project and acquired the funding. EL, MH, LC, HB, FP, GK and AS reviewed the manuscript.

455 **Acknowledgement.** The work is accomplished in the frame of the VESUVIO project funded by the Supporting Talent in ReSearch (STARS) grant at Università degli Studi di Padova, IT. We would like to thank Armin Leuprecht for the support and all the suggestions to make the archive technically correct.

## References

Global Volcanism Program accessible at <https://volcano.si.edu/>

460 The 2008 Eruptions of Okmok and Kasatochi Volcanoes, Alaska, *Journal of Geophysical Research: Atmospheres*, Special issue, 7 February 2018.

Angerer, B., Ladstädter, F., Scherllin-Pirscher, B., Schwärz, M., Steiner, A.K., Foelsche, U., Kirchengast, G.: Quality aspects of the Wegener Center multi-satellite GPS radio occultation record OPSv5.6, *Atmospheric Measurement Techniques*, 10, 4845–4863, <https://doi.org/10.5194/amt-10-4845-2017>, 2017.

465 Anthes, R. A., Bernhardt, P. A., Chen, Y., Cucurull, L., Dymond, K. F., Ector, D., Healy, S. B., Ho, S., Hunt, D. C., Kuo, Y., Liu, H., Manning, K., McCormick, C., Meehan, T.K., Randel, W. J., Rocken, C., Schreiner, W. S., Sokolovskiy, S. V., Syndergaard, S., Thompson, D. C., Trenberth, K. E., Wee, T., Yen, N. L., Zeng, Z.: The COSMIC/Formosat/3 mission: early results, *Bull. Am. Meteorol. Soc.*, 89, 313–333. <http://dx.doi.org/10.1175/BAMS-89-3-313>, 2008.

470 Beyerle, G., Schmidt, T., Michalak, G., Heise, S., Wickert, J., Reigber, C.: GPS radio occultation with GRACE: atmospheric profiling utilizing the zero difference technique, *Geophys. Res. Lett.*, 32, L13806. <http://dx.doi.org/10.1029/2005GL023109>, 2005.

475 Bignami, C., Corradini, S., Merucci, L., de Michele, M., Raucoules, D., De Astis, G., ... & Piedra, J: Multisensor satellite monitoring of the 2011 Puyehue-Cordón Caulle eruption, *IEEE journal of selected topics in applied earth observations and remote sensing*, 7, 2786–2796, <https://doi.org/10.1109/JSTARS.2014.2320638>, 2014.

Biondi, R., Steiner, A. K., Kirchengast, G., Brenot, H., & Rieckh, T.: Supporting the detection and monitoring of VCs: A promising new application of Global Navigation Satellite System radio occultation, *Adv. Space Res.*, 60, 2707–2722, <https://doi.org/10.1016/j.asr.2017.06.039>, 2017.

480 Bourassa, A. E., Robock, A., Randel, W. J., Deshler, T., Rieger, L. A., Lloyd, N. D., ... & Degenstein, D. A: Large volcanic aerosol load in the stratosphere linked to Asian monsoon transport, *Science*, 337, 78–81, <https://doi.org/10.1126/science.1219371>, 2012.

Brenot, H., Theys, N., Clarisse, L., van Geffen, J., van Gent, J., Van Roozendaal, M., et al.: Support to Aviation Control Service (SACS): an online service for near-real-time satellite monitoring of volcanic plumes, *Natural Hazards and Earth System Sciences*, 14(5), 1099–1123. <https://doi.org/10.5194/nhess-14-1099-2014>, 2014.

- 485 Brown, S. K., Crosweller, H. S., Sparks, R. S. J., Cottrell, E., Deligne, N. I., Guerrero, N. O., ... Takarada, S.: Characterisation of the Quaternary eruption record: Analysis of the Large Magnitude Explosive Volcanic Eruptions (LaMEVE) database, *Journal of Applied Volcanology*, 3(1), 1–22. <https://doi.org/10.1186/2191-5040-3-5>, 2014.
- Carboni, E., Grainger, R. G., Mather, T. A., Pyle, D. M., Thomas, G. E., Siddans, R., Smith, A. J. A., Dudhia, A., Koukoulis, M. E., Balis, D.: The vertical distribution of volcanic SO<sub>2</sub> plumes measured by IASI, *Atmospheric Chemistry and Physics*, 16(7), 4343–4367. <https://doi.org/10.5194/acp-16-4343-2016>, 2016.
- 490 Carn, S. A., & Lopez, T. M.: Opportunistic validation of sulfur dioxide in the Sarychev Peak volcanic eruption cloud, *Atmospheric Measurement Techniques*, 4(9), 1705, <https://doi.org/10.5194/amt-4-1705-2011>, 2011.
- Carn, S. A., Clarisse, L., & Prata, A. J.: Multi-decadal satellite measurements of global volcanic degassing, *Journal of Volcanology and Geothermal Research*, 311, 99–134. <https://doi.org/10.1016/j.jvolgeores.2016.01.002>, 2016.
- 495 Carn, S. A., Fioletov, V. E., Mclinden, C. A., Li, C., & Krotkov, N. A.: A decade of global volcanic SO<sub>2</sub> emissions measured from space, *Scientific Reports*, 7, 1–12. <https://doi.org/10.1038/srep44095>, 2017.
- Carn, S. A.: Multi-Satellite Volcanic Sulfur Dioxide L4 Long-Term Global Database V3, Greenbelt, MD, USA, Goddard Earth Science Data and Information Services Center (GES DISC), 10.5067/MEASURES/SO2/DATA404, 2019.
- 500 Cigala, V., Biondi, R., Prata, A. J., Steiner, A. K., Kirchengast, G., & Brenot, H.: GNSS Radio Occultation Advances the Monitoring of Volcanic Clouds: The Case of the 2008 Kasatochi Eruption, *Remote Sensing*, 11, 2199, <https://doi.org/10.3390/rs11192199>, 2019.
- Clarisse, L., Hurtmans, D., Clerbaux, C., Hadji-Lazaro, J., Ngadi, Y., and Coheur, P.-F.: Retrieval of sulphur dioxide from the infrared atmospheric sounding interferometer (IASI), *Atmos. Meas. Tech.*, 5, 581–594, <https://doi.org/10.5194/amt-5-581-2012>, 2012.
- 505 Clarisse, L., Coheur, P.-F., Theys, N., Hurtmans, D., and Clerbaux, C.: The 2011 Nabro eruption, a SO<sub>2</sub> plume height analysis using IASI measurements, *Atmos. Chem. Phys.*, 14, 3095–3111, <https://doi.org/10.5194/acp-14-3095-2014>, 2014.
- Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., Wespes, C., Coheur, P.-F.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmos. Chem. Phys.*, 9, 6041–6054, <https://doi.org/10.5194/acp-9-6041-2009>, 2009.
- Cooke, M. C., Francis, P. N., Millington, S., Saunders, R., & Witham, C.: Detection of the Grímsvötn 2011 volcanic eruption plumes using infrared satellite measurements, *Atmospheric Science Letters*, 15, 321–327, <https://doi.org/10.1002/asl2.5062>, 014.
- 515 de Moor, J. M., Kern, C., Avar, G., Muller, C., Aiuppa, A., Saballos, A., ... Fischer, T. P.: A New Sulfur and Carbon Degassing Inventory for the Southern Central American Volcanic Arc: The Importance of Accurate Time-Series Data Sets and Possible Tectonic Processes Responsible for Temporal Variations in Arc-Scale Volatile Emissions, *Geochemistry, Geophysics, Geosystems*, 18, 4437–4468. <https://doi.org/10.1002/2017GC007141>, 2017.
- Doeringer, D., Eldering, A., Boone, C. D., González Abad, G., & Bernath, P. F.: Observation of sulfate aerosols and SO<sub>2</sub> from the Sarychev volcanic eruption using data from the Atmospheric Chemistry Experiment (ACE), *Journal of Geophysical Research: Atmospheres*, 117, D3, <https://doi.org/10.1029/2011JD0165562>, 012.
- 520 Dikty, S., Richter, A., Bovensmann, H., Wittrock, F., Weber, M., Noël, S., Burrows, J. P., Munro, R., and Lang, R.: GOME-2 level 2 products at IUP Bremen and first results on the quantification of the effects of optical degradation, paper presented at Meteorological Satellite Conference, Eur. Organ. for the Exploit. of Meteorol. Satell., Cordoba, Spain, 20–24 Sept, 2010.
- 525 EOPAC Team, GNSS Radio Occultation Record (OPS 5.6 2001–2018), University of Graz, Austria, <https://doi.org/10.25364/WEGC/OPS5.6:2019.1>, 2019.

- 530 Fee, D., Yokoo, A., & Johnson, J. B.: Introduction to an open community infrasound dataset from the actively erupting Sakurajima Volcano, Japan, *Seismological Research Letters*, 85, 1151–1162. <https://doi.org/10.1785/0220140051>, 2014.
- Flemming, J., & Inness, A.: Volcanic sulfur dioxide plume forecasts based on UV satellite retrievals for the 2011 Grímsvötn and the 2010 Eyjafjallajökull eruption, *Journal of Geophysical Research: Atmospheres*, 118, 10-172, <https://doi.org/10.1002/jgrd.507532>, 013.
- 535 Flentje, H., Claude, H., Elste, T., Gilge, S., Köhler, U., Plass-Dülmer, C., ... & Fricke, W.: The Eyjafjallajökull eruption in April 2010-detection of volcanic plume using in-situ measurements, ozone sondes and lidar-ceilometer profiles, *Atmospheric Chemistry and Physics*, 10, 10085, <https://doi.org/10.5194/acp-10-10085-2010>, 2010.
- Foelsche, U., Scherllin-Pirscher, B., Ladstädter, F., Steiner, A.K., Kirchengast, G.: Refractivity and temperature climate records from multiple radio occultation satellites consistent within 0.05%, *Atmos. Meas. Tech.*, 4, 2007–2018, <http://dx.doi.org/10.5194/amt-4-2007-2011>, 2011.
- 540 Fromm, M., Kablick III, G., Nedoluha, G., Carboni, E., Grainger, R., Campbell, J., & Lewis, J.: Correcting the record of volcanic stratospheric aerosol impact: Nabro and Sarychev Peak, *Journal of Geophysical Research: Atmospheres*, 119, 10-343, <https://doi.org/10.1002/2014JD0215072>, 2014.
- 545 Ge, C., Wang, J., Carn, S., Yang, K., Ginoux, P., & Krotkov, N.: Satellite-based global volcanic SO<sub>2</sub> emissions and sulfate direct radiative forcing during 2005-2012, *Journal of Geophysical Research: Atmospheres*, 121, 3446–3464. <https://doi.org/10.1002/2015JD023134>, 2016.
- Griessbach, S., Hoffmann, L., Spang, R., & Riese, M.: Volcanic ash detection with infrared limb sounding: MIPAS observations and radiative transfer simulations, *Atmos. Meas. Tech.*, 7, 1487-1507, <https://doi.org/10.5194/amt-7-1487-2014>, 2014.
- 550 Hajj, G. A., Ao, C. O., Iijima, B. A., Kuang, D., Kursinski, E. R., Mannauci, A.J., Meehan, T. K., Romans, L. J., de la Torre Juarez, M., Yunck, T. P.: CHAMP and SAC-C atmospheric occultation results and intercomparisons, *J. Geophys. Res.*, 109, D06109. <http://dx.doi.org/10.1029/2003JD003909>, 2004.
- Houghton, B. F., Swanson, D. A., Rausch, J., Carey, R. J., Fagents, S. A., & Orr, T. R.: Pushing the volcanic explosivity index to its limit and beyond: Constraints from exceptionally weak explosive eruptions at Kilauea in 2008, *Geology*, 41, 627–630. <https://doi.org/10.1130/G34146.1>, 2013.
- 555 Ivy, D. J., Solomon, S., Kinnison, D., Mills, M. J., Schmidt, A., & Neely, R. R.: The influence of the Calbuco eruption on the 2015 Antarctic ozone hole in a fully coupled chemistry-climate model, *Geophysical Research Letters*, 44(5), 2556-2561, <https://doi.org/10.1002/2016GL071925>, 2017.
- 560 Kim, M. H., Omar, A. H., Tackett, J. L., Vaughan, M. A., Winker, D. M., Trepte, C. R., ... & Kar, J.: The CALIPSO version 4 automated aerosol classification and lidar ratio selection algorithm, *Atmospheric Measurement Techniques*, 11, 6107, <https://doi.org/10.5194/amt-11-6107-2018>, 2018.
- Kravitz, B., Robock, A., Bourassa, A., Deshler, T., Wu, D., Mattis, I., ... & Duck, T. J.: Simulation and observations of stratospheric aerosols from the 2009 Sarychev volcanic eruption, *Journal of Geophysical Research: Atmospheres*, 116, D18, <https://doi.org/10.1029/2010JD015501>, 2011.
- 565 Kristiansen, N. I., Prata, A. J., Stohl, A., Carn S. A.: Stratospheric volcanic ash emissions from the 13 February 2014 Kelut eruption, *Geophysical Research Letters*, 42, 588-596, <https://doi.org/10.1002/2014GL062307>, 2015.
- Kursinski, E. R., Hajj, G. A., Schofield, J. T., Linfield, R. P.: Observing Earth's atmosphere with radio occultation measurements using the Global Positioning System, *J. Geophys. Res.*, 102(D19), 23,423-429,465. <https://doi.org/10.1029/97JD01569>, 1997.

- 570 Lang, R., Munro, R., Livschitz, Y., Dyer, R., and Lacan, A.: GOME-2 FM3 long-term in-orbit degradation—Basic signatures after 2nd throughput test, Tech Rep. EUM.OPS-EPS.DOC.09.0464, Eur. Organ. for the Exploit. of Meteorol. Satell., Darmstadt, Germany, 2009.
- JS Lopes, F., Silva, J. J., Antuña Marrero, J. C., Taha, G., & Landulfo, E.: Synergetic aerosol layer observation after the 2015 Calbuco volcanic eruption event, *Remote Sensing*, 11, 195, <https://doi.org/10.3390/rs11020195>, 2019.
- 575 Luntama, J. P., Kirchengast, G., Borsche, M., Foelsche, U., Steiner, A., Healy, S., von Engel, A., O'Clérigh, E., Marquardt, C.: Prospects of the EPS GRAS mission for operational atmospheric applications, *Bull. Am. Meteorol. Soc.*, 89(12), 18631875. <https://doi.org/10.1175/2008BAMS2399.1>, 2008.
- Marenco, F., Johnson, B., Turnbull, K., Newman, S., Haywood, J., Webster, H., & Ricketts, H.: Airborne lidar observations of the 2010 Eyjafjallajökull volcanic ash plume, *Journal of Geophysical Research: Atmospheres*, 116, D20, <https://doi.org/10.1029/2011JD016396>, 2011.
- 580 Marzano, F. S., Lamantea, M., Montopoli, M., Herzog, M., Graf, H., & Cimini, D.: Microwave remote sensing of the 2011 Plinian eruption of the Grímsvötn Icelandic volcano, *Remote sensing of Environment*, 129, 168-184, <https://doi.org/10.1016/j.rse.2012.11.005>, 2013.
- 585 Marzano, F. S., Corradini, S., Mereu, L., Kylling, A., Montopoli, M., Cimini, D., ... & Stelitano, D.: Multisatellite Multisensor Observations of a Sub-Plinian Volcanic Eruption: The 2015 Calbuco Explosive Event in Chile, *IEEE Transactions on Geoscience and Remote Sensing*, 56, 2597-2612, <https://doi.org/10.1109/TGRS.2017.2769003>, 2018.
- Munro, R., Eisinger, M., Anderson, C., Callies, J., Corpaccioli, E., Lang, R., Lefebvre, A., Livschitz, Y., Pérez Albiñana, A.: GOME-2 on MetOp, *Proc. of The 2006 EUMETSAT Meteorological Satellite Conference*, Helsinki, Finland, p. 48, 2006.
- 590 Newhall, C. G., & Self, S.: The volcanic explosivity index (VEI) an estimate of explosive magnitude for historical volcanism, *Journal of Geophysical Research*, 87(C2), 1231. <https://doi.org/10.1029/JC087iC02p01231>, 1982.
- Pardini, F., Burton, M., Arzilli, F., La Spina, G., & Polacci, M.: SO<sub>2</sub> emissions, plume heights and magmatic processes inferred from satellite data: The 2015 Calbuco eruptions, *Journal of Volcanology and Geothermal Research*, 361, 12–24. <https://doi.org/10.1016/j.jvolgeores.2018.08.001>, 2018.
- 595 Picquout, A., Lavigne, F., Mei, E. T. W., Grancher, D., Noer, C., Vidal, C. M., & Hadmoko, D. S.: Air traffic disturbance due to the 2010 Merapi volcano eruption, *Journal of volcanology and geothermal research*, 261, 366-375, <https://doi.org/10.1016/j.jvolgeores.2013.04.005>, 2013.
- Platt, U. and Stutz, J.: *Differential optical absorption spectroscopy: Principles and Applications.*, Springer-Verlag Berlin Heidelberg, 2008.
- 600 Prata, A. J., & Bernardo, C.: Retrieval of volcanic SO<sub>2</sub> column abundance from Atmospheric Infrared Sounder data, *Journal of Geophysical Research: Atmospheres*, 112(D20), <https://doi.org/10.1029/2006JD007955>, 2007.
- Prata, A. J., & Prata, A. T.: Eyjafjallajökull volcanic ash concentrations determined using Spin Enhanced Visible and Infrared Imager measurements, *Journal of Geophysical Research: Atmospheres*, 117, D20, <https://doi.org/10.1029/2011JD016800>, 2012.
- 605 Prata, A. J., Gangale, G., Clarisse, L., & Karagulian, F.: Ash and sulfur dioxide in the 2008 eruptions of Okmok and Kasatochi: Insights from high spectral resolution satellite measurements, *Journal of Geophysical Research*, 115(22), D00L18, <https://doi.org/10.1029/2009JD013556>, 2010.
- Prata, F., Woodhouse, M., Huppert, H. E., Prata, A., Thordarson, T., & Carn, S.: Atmospheric processes affecting the separation of volcanic ash and SO<sub>2</sub> in volcanic eruptions: Inferences from the May 2011 Grímsvötn eruption, *Atmospheric Chemistry and Physics*, 17(17), 10709–10732. <https://doi.org/10.5194/acp-17-10709-2017>, 2017.

- 610 Rix, M., Valks, P., Hao, N., Loyola, D. G., Schlager, H., Huntrieser, H. H., Flemming, J., Koehler, U., Schumann, U., and Inness, A.: Volcanic SO<sub>2</sub>, BrO and plume height estimations using GOME-2 satellite measurements during the eruption of Eyjafjallajökull in May 2010, *J. Geophys. Res.*, 117, D00U19, <https://doi.org/10.1029/2011JD016718>, 2012.
- 615 Robock, A.: Climatic impacts of volcanic eruptions, *The encyclopedia of volcanoes* (pp. 935-942). Academic Press, 2015.
- Rybin, A., Chibisova, M., Webley, P., Steensen, T., Izbekov, P., Neal, C., & Realmuto, V.: Satellite and ground observations of the June 2009 eruption of Sarychev Peak volcano, Matua Island, Central Kuriles, *Bulletin of Volcanology*, 73, 1377-1392, <https://doi.org/10.1007/s00445-011-0481-0>, 2011.
- 620 Scherllin-Pirscher, B., Steiner, A.K., Kirchengast, G., Kuo, Y.-H., Foelsche, U.: Empirical analysis and modeling of errors of atmospheric profiles from GPS radio occultation, *Atmos. Meas. Tech.*, 4, 1875–1890. <http://dx.doi.org/10.5194/amt-4-1875-2011>, 2011.
- Schnetzler, C. C., Bluth, G. J. S., Krueger, A. J., & Walter, L. S.: A proposed volcanic sulfur dioxide index (VSI), *Journal of Geophysical Research: Solid Earth*, 102, 20087-20091, <https://doi.org/10.1029/97JB01142>, 1997.
- 625 Stohl, A., Prata, A. J., Eckhardt, S., Clarisse, L., Durant, A., Henne, S., ... & Stebel, K.: Determination of time-and height-resolved volcanic ash emissions and their use for quantitative ash dispersion modeling: the 2010 Eyjafjallajökull eruption, *Atmospheric Chemistry and Physics*, 11, 4333-4351, <https://doi.org/10.5194/acp-11-4333-2011>, 2011.
- 630 Stocker, M., Ladstädter, F., Wilhelmssen, H., & Steiner, A. K.: Quantifying Stratospheric Temperature Signals and Climate Imprints From Post-2000 Volcanic Eruptions, *Geophysical Research Letters*, 46, 12486-12494, <https://doi.org/10.1029/2019GL084396>, 2019.
- Stone, K. A., Solomon, S., Kinnison, D. E., Pitts, M. C., Poole, L. R., Mills, M. J., ... & Vernier, J. P.: Observing the impact of Calbuco volcanic aerosols on South Polar ozone depletion in 2015, *Journal of Geophysical Research: Atmospheres*, 122(21), 11-862, <https://doi.org/10.1002/2017JD0269872>, 017.
- 635 Surono, Jousset, P., Pallister, J., Boichu, M., Buongiorno, M. F., Budisantoso, A., ... Lavigne, F.: The 2010 explosive eruption of Java's Merapi volcano—A '100-year' event, *Journal of Volcanology and Geothermal Research*, 241–242, 121–135, <https://doi.org/10.1016/j.jvolgeores.2012.06.018>, 2012.
- Susskind, J., Barnet, C. D., & Blaisdell, J. M.: Retrieval of atmospheric and surface parameters from AIRS/AMSU/HSB data in the presence of clouds, *IEEE Transactions on Geoscience and Remote Sensing*, 41, 390-409, <https://doi.org/10.1109/TGRS.2002.808236>, 2003.
- 640 Telling, J., Flower, V. J. B., & Carn, S. A.: A multi-sensor satellite assessment of SO<sub>2</sub> emissions from the 2012–13 eruption of Plosky Tolbachik volcano, Kamchatka, *Journal of Volcanology and Geothermal Research*, 307, 98-106, <https://doi.org/10.1016/j.jvolgeores.2015.07.010>, 2015.
- 645 Theys, N., Campion, R., Clarisse, L., Brenot, H., van Gent, J., Dils, B., Corradini, S., Merucci, L., Coheur, P.-F., Van Roozendael, M., Hurtmans, D., Clerbaux, C., Tait, S., and Ferrucci, F.: Volcanic SO<sub>2</sub> fluxes derived from satellite data: a survey using OMI, GOME-2, IASI and MODIS, *Atmos. Chem. Phys.*, 13, 5945–5968, <https://doi.org/10.5194/acp-13-5945-2013>, 2013.
- Theys, N., De Smedt, I., Van Roozendael, M., Froidevaux, L., Clarisse, L., & Hendrick, F.: First satellite detection of volcanic OClO after the eruption of Puyehue-Cordón Caulle, *Geophysical Research Letters*, 41, 667-672, <https://doi.org/10.1002/2013GL058416>, 2014.
- 650 Tournigand, P.-Y., Cigala, V., Lasota, E., Hammouti, M., Clarisse, L., Brenot, H., Prata, F., Kirchengast, G., Steiner, A., Biondi, R.: A comprehensive archive of large SO<sub>2</sub> volcanic clouds in 2000s, *GFZ Data Services*. <http://doi.org/10.5880/figeo.2020.016>, 2020a.



Tournigand, P.-Y., Cigala, V., Prata, F., Steiner, A., Kirchengast, G., Brenot, H., Clarisse, L., Biondi, R.: the 2015 Calbuco volcanic cloud detection using GNSS radio occultation and satellite lidar, IGARSS 2020 Proceedings, 2020b.

655 Turner, M. B., Bebbington, M. S., Cronin, S. J., & Stewart, R. B.: Merging eruption datasets: Building an integrated Holocene eruptive record for Mt Taranaki, New Zealand, *Bulletin of Volcanology*, 71(8), 903–918, <https://doi.org/10.1007/s00445-009-0274-x>, 2009.

Vernier, J. P., Fairlie, T. D., Deshler, T., Natarajan, M., Knepp, T., Foster, K., ... & Trepte, C.: In situ and space-based observations of the Kelud volcanic plume: The persistence of ash in the lower stratosphere, *Journal of Geophysical Research: Atmospheres*, 121, 11-104, <https://doi.org/10.1002/2016JD025344>, 2016.

660 Wickert, J., Reigber, C., Beyerle, G., König, R., Marquardt, C., Schmidt, T., Grunwaldt, L., Galas, R., Meehan, T. K., Melbourne, W. G., Hocke, K.: Atmosphere sounding by GPS radio occultation: First results from CHAMP, *Geophysical research letters*, 28(17), 3263-3266, <https://doi.org/10.1029/2001GL013117>, 2001.

Williams-Jones, G., & Rymer, H.: Hazards of volcanic gases, *The Encyclopedia of Volcanoes* (pp. 985-992), Academic Press, 2015.

665 Winker, D. M., Vaughan, M. A., Omar, A., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., Young, S. A.: Overview of the CALIPSO mission and CALIOP data processing algorithms, *J. Atmos. Oceanic Technol.*, 26, 2310–2323. <http://dx.doi.org/10.1175/2009JTECHA1281.1>, 2009.

670 Yang, K., Liu, X., Bhartia, P. K., Krotkov, N., Carn, S., Hughes, E., Krueger, A., Spurr, R., and Trahan, S.: Direct retrieval of sulfur dioxide amount and altitude from spaceborne hyperspectral UV measurements: Theory and application, *J. Geophys. Res.*, 115, D00L09, <https://doi.org/10.1029/2010JD013982>, 2010.

Zeng, Z., Sokolovskiy, S., Schreiner, W. S., Hunt, D.: Representation of Vertical Atmospheric Structures by Radio Occultation Observations in the Upper Troposphere and Lower Stratosphere: Comparison to High-Resolution Radiosonde Profiles, *J. Atmos. Ocean. Technol.*, 36(4), 655–670. <http://dx.doi.org/10.1175/JTECH-D-18-0105.1>, 2019.

675 Zhu, Y., Toon, O. B., Kinnison, D., Harvey, V. L., Mills, M. J., Bardeen, C. G., ... & Jégou, F.: Stratospheric aerosols, polar stratospheric clouds, and polar ozone depletion after the Mount Calbuco eruption in 2015, *Journal of Geophysical Research: Atmospheres*, 123(21), 12-308, <https://doi.org/10.1029/2018JD0289742>, 018.

680 Zuev, V. V., Savelieva, E. S., & Parezheva, T. V.: Study of the Possible Impact of the Calbuco Volcano Eruption on the Abnormal Destruction of Stratospheric Ozone over the Antarctic in Spring 2015, *Atmospheric and Oceanic Optics*, 31(6), 665-669, <https://doi.org/10.1134/S1024856018060192>, 2018.

685

**Table 1. Summary of data used to build the archive.**

Sensor	Satellite(s)	Vertical resolution	Spatial resolution	Estimation	Wavelength	Algorithm reference
<i>AIRS</i>	Aqua (A-Train)	NA	13.5 km x 13.5 km	SO <sub>2</sub> VCD	7.3 μm	Prata and Bernardo, 2007
<i>IASI</i>	MetOp-A/B	NA	12 km diameter	SO <sub>2</sub> VCD		Clarisse et al., 2012



				Cloud top height	3.62 - 15.5 $\mu\text{m}$	Clarisse et al., 2014
<b>GOME-2</b>	MetOp-A/B	NA	40 km x80 km	SO <sub>2</sub> VCD	240 - 400 nm	Rix et al., 2012
<b>CALIOP</b>	CALIPSO	60 m below 20 km 180 m above 20 km	1 km 1.667 km	Cloud top height Cloud type	532 nm 1064 nm	Winker et al., 2009
<b>GNSS RO</b>	CHAMP COSMIC C/NOFS SAC-C GRACE-A Met-Op	100 m in the troposphere 600 m in the stratosphere	50 km in the troposphere 200–300 km in the stratosphere	Bending angle Bangle anomaly Refractivity Temperature Pressure Specific Humidity Cloud top height	19.05*10 <sup>4</sup> $\mu\text{m}$ 24.40*10 <sup>4</sup> $\mu\text{m}$	Angerer et al., 2017 Cigala et al., 2019

690

**Table 2. Summary of the volcanoes and related eruption selected for the database. The following information is provided: the name of each volcano; the eruption start date as provided by the GVP; the spatial location of the volcano in latitude and longitude; the plume/cloud height as a range estimated from IASI, CALIOP and GNSS RO (see details below) and the SO<sub>2</sub> mass loading in Tg as reported in the literature.**

Volcano name	Main eruptive event date	VEI	Location Latitude/Longitude (degree)	Archive start/end date	Cloud average height (km) - Sensor	SO <sub>2</sub> mass loading (Tg) (Reference) Sensor
Okmok (Figure S1)	12.07.2008	4	53.397/-168.166	12.07.2008 06.08.2008	12.6 - IASI 12.0 - CALIOP 12.5 - RO	0.12 (Spinei et al., 2010) OMI 0.3 (Prata et al. 2010) AIRS 0.09 (Carn et al., 2016) IASI
Kasatochi (Figure S2)	07.08.2008	4	52.172/-175.509	07.08.2008 29.08.2008	11.7 - IASI 12.0 - CALIOP 12.4 - RO	2.7 (Corradini et al., 2010) MODIS 1.2 (Prata et al., 2010) AIRS 2.2 (Krotkov et al., 2010) OMI 2.0 (Yang et al., 2010) OMI 1.7 (Karagulian et al.) IASI 1.7 (Kristiansen et al., 2010) Multi-sensor 1.6 (Clarisse et al., 2012) IASI 1.6 (Nowlan et al., 2011) GOME-2
Sarychev (Figure S3)	14.06.2009	4	48.092/153.200	11.06.2009 16.07.2009	12.2 - IASI 12.9 - CALIOP 12.3 - RO	1.2 (Haywood et al., 2010) IASI
Eyjafjallajökull (Figure S4)	20.03.2010	4	63.633/19.633	05.05.2010 21.05.2010	8.0 - IASI 12.2 - CALIOP 12.3 - RO	0.17 (Boichu et al., 2013) IASI 1.2 (Rix et al., 2012) GOME-2 0.18 (Carboni et al., 2012) IASI 0.06 (Pugnaghi et al., 2016) MODIS

Merapi (Figure S5)	04.11.2010	4	-7.542/110.442	26.10.2010 11.11.2010	12.4 - IASI 14.5 - CALIOP 16.1 - RO	0.44 (Surono et al., 2012) Multi-sensor
Grímsvötn (Figure S6)	21.05.2011	4	64.416/-17.316	22.05.2011 18.06.2011	10.8 - IASI 12.2 - CALIOP 11.7 - RO	0.24 (Prata et al., 2017) 0.38 (Carn et al., 2016) 0.4 (Sigmarsson et al., 2013) OMI+IASI 0.61 (Moxnes et al., 2014)
Nabro (Figure S7)	12.06.2011	4	13.370/41.700	31.05.2011 25.06.2011	12.2 - IASI 14.3 - CALIOP 15.3 - RO	4.5 (Theys et al., 2013) Multi-sensor
Puyehue- Cordón Caulle (Figure S8)	04.06.2011	5	-40.59/-72.117	07.06.2011 18.06.2011	12.2 - IASI NA - CALIOP 12.5 - RO	0.2 (Theys et al., 2013) IASI
Tolbachik (Figure S9)	27.11.2012	4	55.832/160.326	27.11.2012 03.12.2012	8.9 - IASI 11.4 - CALIOP 11.7 - RO	0.2 (Telling et al. 2015) Multi-sensor 0.09 (Carn et al., 2016) IASI
Kelut (Figure S10)	13.02.2014	4	-7.939/112.307	17.02.2014 18.02.2014	17.6 - IASI NA - CALIOP 16.9 - RO	0.2 (Carn et al., 2016) OMI 0,19 (Carn et al., 2016) IASI
Calbuco (Figure S11)	22.04.2015	4	-41.328/-72.607	24.04.2015 24.05.2015	16.4 - IASI 12.4 - CALIOP 14.8 - RO	0.3 (Pardini et al., 2018)

**Table 3. General description of all the variables contained in the archive**

Variable name	Content	Dimension (rows, columns)	Type	Unit
AIRS_lat	Latitude data, each column corresponds to a granule and each row to one data point in a granule.	AIRS_lat, date_AIRS	double	degrees north
AIRS_lon	Longitude data, each column corresponds to a granule and each row to one data point in a granule.	AIRS_lat, date_AIRS	double	degrees east
AIRS_date	Date of granule contained in each column.	1, date_AIRS	int	seconds since 1970-01-01 00:00:0.0
AIRS_SO2	SO2 data, each column corresponds to a granule and each row to one data point in a granule.	AIRS_lat, date_AIRS	double	DU
IASI_lat	Latitude data, each column corresponds to a granule and each row to one data point in a granule.	IASI_lat, date_IASI	double	degrees north
IASI_lon	Longitude data, each column corresponds to a granule and each row to one data point in a granule.	IASI_lat, date_IASI	double	degrees east
IASI_date	Date of granule contained in each column.	1, date_IASI	int	seconds since 1970-01-01 00:00:0.0

IASI_SO2	SO2 data, each column corresponds to a granule and each row to one data point in a granule.	IASI_lat, date_IASI	double	DU
IASI_height	Cloud top height estimated with IASI	IASI_lat, date_IASI	double	m
GOME_lat	Latitude data, each column corresponds to a granule and each row to one data point in a granule.	GOME_lat, date_GOME	double	degrees north
GOME_lon	Longitude data, each column corresponds to a granule and each row to one data point in a granule.	GOME_lat, date_GOME	double	degrees east
GOME_date	Date of granule contained in each column.	1, date_GOME	int	seconds since 1970-01-01 00:00:0.0
GOME_SO2_1	SO2 data, each column corresponds to a granule and each row to one data point in a granule.	GOME_lat, date_GOME	double	DU
GOME_SO2_2	SO2 data, each column corresponds to a granule and each row to one data point in a granule.	GOME_lat, date_GOME	double	DU
GOME_SO2_3	SO2 data, each column corresponds to a granule and each row to one data point in a granule.	GOME_lat, date_GOME	double	DU
CALIOP_lat	Latitude data, each row corresponds to one point of a CALIOP track.	CALIOP_lat, 1	double	degrees north
CALIOP_lon	Longitude data, each row corresponds to one point of a CALIOP track.	CALIOP_lat, 1	double	degrees east
CALIOP_date	Date and time, each row corresponds to one point of a CALIOP track.	CALIOP_lat, 1	int	seconds since 1970-01-01 00:00:0.0
CALIOP_filename	Filename, each row provides the filename of the given data point.	CALIOP_lat, CALIOP_char	char	n.a.
CALIOP_height	Cloud top altitude data, each row corresponds to one point of a CALIOP track and each column to a collocated sensor.	CALIOP_lat, Sensors	double	m
CALIOP_type	Cloud type data, each row corresponds to one point of a CALIOP track, three columns corresponding to three levels of altitude -0.5 to 8.2 km, 8.2 to 20.2km and 20.2 to 30.1km	CALIOP_lat, CALIOP_char2, CALIOP_type	double	n.a.
<b>Only volcano files</b>				
RO_lat	Latitude data, each row corresponds to one profile point and each column to a ro profile.	RO_lat, RO_profile	double	degrees north
RO_lon	Longitude data, each row corresponds to one profile point and each column to a ro profile.	RO_lat, RO_profile	double	degrees east
RO_date	Date and time data, each row corresponds to one profile point and each column to a ro profile.	RO_lat, RO_profile	int	seconds since 1970-01-01 00:00:0.0
RO_bending_angle	Bending angle data, each row corresponds to one profile point and each column to a ro profile.	RO_lat, RO_profile	double	rad
RO_anomaly_bending_angle	Bending angle anomaly data, each row corresponds to one profile point and each column to a ro profile.	RO_lat, RO_profile	double	percent
RO_temperature	Temperature data, each row corresponds to one profile point and each column to a ro profile.	RO_lat, RO_profile	double	K
RO_pressure	Pressure data, each row corresponds to one profile point and each column to a ro profile.	RO_lat, RO_profile	double	Pa
RO_refractivity	Refractivity data, each row corresponds to one profile point and each column to a ro profile.	RO_lat, RO_profile	double	1
RO_specific_humidity	Specific humidity data, each row corresponds to one profile point and each column to a ro profile.	RO_lat, RO_profile	double	kg.kg-1
RO_heightVC	Cloud top altitude data, each column corresponds to a ro profile.	1, RO_profile	double	m
<b>Only daily files</b>				
RO_AIRS_lat	Latitude data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	double	degrees north

RO_AIRS_lon	Longitude data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	double	degrees east
RO_AIRS_date	Date and time data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	int	seconds since 1970-01-01 00:00:0.0
RO_AIRS_bending_angle	Bending angle data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	double	rad
RO_AIRS_anomaly_bending_angle	Bending angle anomaly data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	double	percent
RO_AIRS_temperature	Temperature data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	double	K
RO_AIRS_pressure	Pressure data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	double	Pa
RO_AIRS_refractivity	Refractivity data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	double	1
RO_AIRS_specific_humidity	Specific humidity data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	double	kg.kg-1
RO_AIRS_heightVC	Cloud top altitude data, each column corresponds to a ro profile.	1, RO_AIRS_profile	double	m
RO_IASI_lat	Latitude data, each row corresponds to one profile point and each column to a ro profile.	RO_IASI_lat, RO_IASI_profile	double	degrees north
RO_IASI_lon	Longitude data, each row corresponds to one profile point and each column to a ro profile.	RO_IASI_lat, RO_IASI_profile	double	degrees east
RO_IASI_date	Date and time data, each row corresponds to one profile point and each column to a ro profile.	RO_IASI_lat, RO_IASI_profile	int	seconds since 1970-01-01 00:00:0.0
RO_IASI_bending_angle	Bending angle data, each row corresponds to one profile point and each column to a ro profile.	RO_IASI_lat, RO_IASI_profile	double	rad
RO_IASI_anomaly_bending_angle	Bending angle anomaly data, each row corresponds to one profile point and each column to a ro profile.	RO_IASI_lat, RO_IASI_profile	double	percent
RO_IASI_temperature	Temperature data, each row corresponds to one profile point and each column to a ro profile.	RO_IASI_lat, RO_IASI_profile	double	K
RO_IASI_pressure	Pressure data, each row corresponds to one profile point and each column to a ro profile.	RO_IASI_lat, RO_IASI_profile	double	Pa
RO_IASI_refractivity	Refractivity data, each row corresponds to one profile point and each column to a ro profile.	RO_IASI_lat, RO_IASI_profile	double	1
RO_IASI_specific_humidity	Specific humidity data, each row corresponds to one profile point and each column to a ro profile.	RO_IASI_lat, RO_IASI_profile	double	kg.kg-1
RO_IASI_heightVC	Cloud top altitude data, each column corresponds to a ro profile.	1, RO_IASI_profile	double	m
RO_GOME_lat	Latitude data, each row corresponds to one profile point and each column to a ro profile.	RO_GOME_lat, RO_GOME_profile	double	degrees north
RO_GOME_lon	Longitude data, each row corresponds to one profile point and each column to a ro profile.	RO_GOME_lat, RO_GOME_profile	double	degrees east
RO_GOME_date	Date and time data, each row corresponds to one profile point and each column to a ro profile.	RO_GOME_lat, RO_GOME_profile	int	seconds since 1970-01-01 00:00:0.0
RO_GOME_bending_angle	Bending angle data, each row corresponds to one profile point and each column to a ro profile.	RO_GOME_lat, RO_GOME_profile	double	rad
RO_GOME_anomaly_bending_angle	Bending angle anomaly data, each row corresponds to one profile point and each column to a ro profile.	RO_GOME_lat, RO_GOME_profile	double	percent
RO_GOME_temperature	Temperature data, each row corresponds to one profile point and each column to a ro profile.	RO_GOME_lat, RO_GOME_profile	double	K
RO_GOME_pressure	Pressure data, each row corresponds to one profile point and each column to a ro profile.	RO_GOME_lat, RO_GOME_profile	double	Pa
RO_GOME_refractivity	Refractivity data, each row corresponds to one profile point and each column to a ro profile.	RO_GOME_lat, RO_GOME_profile	double	1

RO_GOME_specific_humidity	Specific humidity data, each row corresponds to one profile point and each column to a ro profile.	RO_GOME_lat, RO_GOME_profile	double	kg.kg-1
RO_GOME_heightVC	Cloud top altitude data, each column corresponds to a ro profile.	1, RO_GOME_profile	double	m
RO_AIRS_lat	Latitude data, each row corresponds to one profile point and each column to a ro profile.	RO_AIRS_lat, RO_AIRS_profile	double	degrees north

695 **Table 4. Number of days, granules, scanning lines and profiles covered by the archive for each volcano in alphabetical order**

Volcano	CALIOP		AIRS		IASI		GOME		RO	
	# of profiles	# of days covered	# of granules	# of days covered	# of scanning lines	# of days covered	# of scanning lines	# of days covered	# of profiles	# of days covered
Calbuco	12495	30	350	30	42740	30	20992	5	5362	31
Eyjafjallajökull	3569	16	76	16	3980	16	164369	16	2624	17
Grímsvötn	6268	21	147	8	49824	20	833541	21	6007	21
Kasatochi	12897	23	247	13	103622	21	650031	23	17045	23
Kelut	72	2	1	1	1313	1	2575	1	83	2
Merapi	1053	11	27	10	4919	15	80193	16	984	17
Nabro	2463	11	123	12	59359	11	638316	9	7131	14
Okmok	5678	23	32	11	2931	18	737981	26	13255	26
PCC	0	0	76	11	21528	11	369992	11	664	12
Sarychev	11563	34	127	17	83533	35	1035931	36	16522	36
Tolbachik	617	7	9	2	5390	5	22133	5	449	7

**Table 5. The average difference between cloud top height estimated with pairs of sensors for each volcano. For each pair we report the average difference (or simply the difference when there is only 1 collocation) of all the collocations and the number of collocations. When there are no collocations the value reported is “/”.**

Volcano	RO-CALIOP		RO-IASI		IASI-CALIOP	
	altitude average (km)	#	altitude average (km)	#	altitude average (km)	#
Calbuco	4.2	39	3.4	867	4.9	308
Eyjafjallajökull	0.3	1	3.7	30	3.5	29
Grímsvötn	0.9	5	3.7	136	2.4	75
Kasatochi	1.3	70	1.2	3855	1.6	997
Kelut	/	0	1.7	20	/	0
Merapi	1.5	1	2.7	127	2.2	70
Nabro	3.4	9	4.3	609	3.6	204
Okmok	3.3	2	1.8	143	2.5	22
Puyehue-Cordón Caulle	/	0	1.6	193	/	0
Sarychev	1.5	24	1.5	1519	2.8	227
Tolbachik	/	0	3.0	68	/	0




Figure 1. Archive schematic organization.





Figure 2. Example of data use and data collocation. (a) Kasatochi cloud on 9<sup>th</sup> of August 2008; (b) Sarychev Peak cloud on 12<sup>th</sup> of June 2009. The central panels show the SO<sub>2</sub> VCD 2-dimensional spreading estimated by AIRS, GOME-2 and IASI, the top right panel show the CALIOP tracks for which the total attenuated backscatter profile is available and the bottom-

710 left panel shows the RO profiles collocated with the SO<sub>2</sub>. For IASI, CALIOP and RO the average volcanic cloud top altitude for the considered day is indicated (VC h).



**Figure 3. Case study Kastochi 2008. RO profiles corresponding to the CALIOP aerosol types profile collocated with the SO<sub>2</sub> estimation from IASI (top panels, 12/08/2008) and AIRS (bottom panels, 11/08/2008).**