

A global viral oceanography database (gVOD)

Le Xie¹, Wei Wei¹, Lanlan Cai¹, Xiaowei Chen¹, Yuhong Huang¹, Nianzhi Jiao¹, Rui Zhang^{1,*}, Ya-Wei Luo^{1,*}

¹State Key Laboratory of Marine Environmental Science, College of Ocean and Earth Sciences, Xiamen University, Xiamen,

5 Fujian 361102, China

Correspondence to: Rui Zhang (ruizhang@xmu.edu.cn) and Ya-Wei Luo (ywluo@xmu.edu.cn)

Abstract. Virioplankton are a key component of the marine biosphere in maintaining diversity of microorganisms and stabilizing ecosystems. They also contribute greatly to nutrient cycles/cycling by releasing organic matter after lysis of hosts.

10 In this study, we constructed the first global viral oceanography database (gVOD) by collecting 10,931 viral abundance (VA) data and 727 viral production (VP) data, along with host and relevant oceanographic parameters when available. Most VA data were obtained in the North Atlantic (32%) and North Pacific Oceans (29%), while the Southeast Pacific and Indian Oceans were quite under sampled. The VA in the global ocean was $1.17(\pm 3.31) \times 10^7$ particles ml^{-1} . The lytic and lysogenic VP in the global ocean was $9.87(\pm 24.16) \times 10^5$ and $2.53(\pm 8.64) \times 10^5$ particles $\text{ml}^{-1} \text{h}^{-1}$, respectively. Average VA in coastal oceans was

15 higher than that in surface open oceans [$3.61(\pm 6.30) \times 10^7$ versus $0.73(\pm 1.24) \times 10^7$ particles ml^{-1}], while average VP in coastal and surface open oceans was close. Vertically, VA, lytic and lysogenic VP decreased from surface to deep oceans by about one order of magnitude. The total number of viruses in the global ocean estimated by bin-averaging and the random forest method was 1.56×10^{30} particles and 1.49×10^{30} particles, leading to an estimate of global ocean viral biomass at 35.9 and 34.4 Tg C, respectively. We expect that the gVOD will be a fundamental and very useful database for laboratory, field and modelling

20 studies in marine ecology and biogeochemistry. The full gVOD database (Xie et al., 2020) is stored in PANGAEA (<https://doi.org/10.1594/PANGAEA.915758>).

1 Introduction

Virioplankton are the most abundant biological entities and one of the largest genetic reservoirs in the ocean (Breitbart, 2012; Fuhrman, 1999). With an estimation of $\sim 10^{23}$ marine microbes being infected every second, viruses play important roles in

25 affecting microbial mortality, regulating community composition and impacting biogeochemical cycles (Suttle, 2005; Zhang et al., 2007). Viruses were estimated to kill ~ 20 –40% of marine bacterioplankton every day, a rate similar to that caused by zooplankton grazing (Fuhrman, 1999). In particular, virus-mediated cell lysis effectively 'shunts' approximately 25% of the photosynthetically fixed carbon, which otherwise would be transferred to higher trophic levels, to the dissolved organic matter (DOM) pool, partly forming the basis of the microbial loop and leading to the recycling of nutrients (Suttle, 2007; Wilhelm

30 and Suttle, 1999). Furthermore, viral lysis can contribute to biological pump through the release of sticky lysates that accelerate the aggregation and sink of carbon into the deep sea (Suttle, 2005).

Compilation of the observations of viral abundance and activity in the global ocean is very necessary and urgent in understanding spatiotemporal distributions of viruses, exploring the controlling factors of viral processes, qualitatively and quantitatively assessing virus-host interactions and viral functioning in marine ecosystems, and even improving predictions of

35 large-scale marine ecosystem and Earth system models. Previous two studies (Bar-On and Milo, 2019; Wigington et al., 2016) summarized viral abundance data in the ocean and estimated viral biomass as well as virus-to-prokaryote ratio. However, the lacking of host parameters such as bacterial production, and oceanographic parameters such as temperature, salinity, nutrient concentrations, limits the usage of these datasets in broader oceanographic contexts. More importantly, there is no public database of viral activity in the global ocean, which substantially hinders our understandings of the ecological and

40 biogeochemical roles of viroplankton on global scale. In addition, the ecological functions of the viruses are tightly linked to their life strategies, mainly including the lytic and the lysogenic infection (Wommack and Colwell, 2000). The significance of viruses in oceanic biogeochemistry is mainly reflected through the lytic infection, which results in cell lysis and the release of DOM. In contrast, other more temperate viruses choosing the lysogenic infection can influence microbial diversity and metabolism by transferring new genes to their hosts, altering the expression of host genes, and not killing hosts for many
45 generations until an environmental or cellular trigger causes them to enter the lytic cycle. The lysogenic infection hence serves as a molecular “time bomb” (Paul, 2008). Therefore, it is necessary to include the quantity and quality data of viral life strategies in a viral oceanographic database.

In this study, we construct the first global viral oceanography database, namely gVOD, by collecting data of viral abundance (VA), lytic and lysogenic viral production (VP), as well as other related viral, host and oceanographic metadata when available.
50 Based on the database, we estimate the total viral number and biomass in the global ocean. In addition, the data of VA and VP generated with different techniques were compared to provide references for evaluating possible technical biases.

2 Data and methods

2.1 Database summary

In the gVOD, direct measurements of three core parameters (VA, lytic and lysogenic VP), as well as accessory viral, prokaryotic and oceanographic parameters when available, were collected from published papers or acquired from lead authors
55 or principal investigators (Table 1). Sampling information including date, latitude, longitude, depth and methods was included for each data record. We used ocean depth shallower or deeper than 200 m as a criterion to identify coastal or open ocean samples. The open ocean samples were further separated into surface and deep samples that collected in 0–200 m and >200 m, respectively.

60 The quality-controlled database consists of 10,931 VA data points (Table S1), 608 lytic VP data points and 119 lysogenic VP data points (Table S2). Most of VA (99.2%) and lytic VP (98.4%) and all lysogenic VP data have accompanying data of prokaryotic abundance (Table 1). For some samples, the abundances of flagellate, picoeukaryotes, *Synechococcus* and *Prochlorococcus* are also available. Prokaryotic productivity measurements cover 22.1% of VA, 57.7% of lytic VP and 76.5% of lysogenic VP data. The most available environmental parameters are salinity and temperature, providing oceanographic
65 information for about half of VA, two-thirds of lytic VP and nearly all lysogenic VP data. Oxygen and chlorophyll *a* concentration data are also adequate particularly for VA. The concentrations of different types of nutrient, including nitrate, silicate and phosphate, are available for many samples. Other environmental parameters (pH, light intensity, dissolved organic carbon concentration) are relatively scarce. Moreover, given that the frequency of viral infected cells was calculated, independently or together with VP, usually to quantify the impact of viral infection within the microbial community (Chen et
70 al., 2019; Payet and Suttle, 2013; Weinbauer et al., 2003), the reported frequencies of lytic infection (n = 438) and lysogenic infection (n = 266) in the literature were also collected into the database to facilitate the future exploration of marine viral

activities. Lastly, we collected 83 viral decay rate data, 206 viral burst size data and 111 virus-mediated mortality data, which can be useful for certain studies.

2.2 Viral abundance

75 The viral abundance in this database was counted using one of the following three methods. In the first method, viruses were harvested by ultracentrifuging onto copper grids and stained with uranyl acetate, and then enumerated using a transmission electron microscopy (TEM) (Akaike, 1974). In the second method, viruses were collected onto 0.02- μm filters and stained with a nucleic acid-specific fluorescent dye (e.g., SYBR Green I), and then were counted under an epifluorescence microscope (EFM) (Noble and Fuhrman, 1998). The third method counted viruses by using flow cytometers (FCM), before which viruses
80 were stained with fluorescent dye (e.g., SYBR Green I or SYBR Gold), and identified on the basis of the green fluorescence versus side scatter signal (Brussaard, 2004; Marie et al., 1999). The details of these three approaches have been described elsewhere (Weinbauer, 2004).

2.3 Lytic viral activity

Lytic VP is paramount and widely employed to assess the activity of lytic viruses in community-level and the roles of viruses
85 in marine ecosystems. In this database, the lytic VP was estimated by one of the following five methods. The first method estimated VP by calculating expected viral release rates by multiplying fraction of viral infected cells (mainly prokaryotes), prokaryotic productivity (assuming equal prokaryotic mortality rate) and burst size obtained from TEM studies (Noble and Steward, 2001) or virus-dilution approach (Weinbauer et al., 2002). For notational simplicity, in this paper we label this method as FPB to represent the three variables (Fraction of viral infected cells, Prokaryotic productivity, and Burst size) listed above
90 and used in the estimation. In the second method, called radioactive incorporation approach (RIA), lytic VP was estimated by determining viral DNA synthesis rates using a labelled radiotracer (e.g., ^3H -, ^{32}P -, or ^{14}C -labeled thymidine or leucine) and a conversion factor to quantify the incorporated radiotracer into viral particles (Noble and Steward, 2001; Zimina et al., 1973). The third method estimated the lytic VP from the viral decay rates (VDR), assuming that the abundance of virus particles is in steady state and then the loss rate of virus particles should be balanced by the production rate (Heldal and Bratbak, 1991). The
95 fourth method used fluorescently labelled viral tracers (FLVT) to measure the dilution rates from the decay of labelled viruses and net changes of the non-labelled viruses in natural viral community (Noble and Fuhrman, 2000). The fifth method quantified the increase of viral abundance during time course incubation using a virus dilution or virus reduction approach (VRA) (Weinbauer et al., 2010; Winget et al., 2005), which effectively avoided new viral infection by reducing viral abundances using pore-size filters or tangential flow filtration systems.

100 2.4 Lysogenic viral activity

Lysogenic VP is generally measured by detecting the proviruses (temperate viruses) that choose lysogenic infection in the environment. Lysogenic VP in this database was estimated using VRA described above after the provirus induction by

Mitomycin C (Weinbauer et al., 2002). Hence, the lysogenic VP was estimated as the difference in viral abundance per unit time between the Mitomycin C treated and the control samples.

105 2.5 Quality control

We conducted quality control for the VA, lytic and lysogenic VP data of the database. A negative lytic VP (Wells and Deming, 2006) was removed. All zero-value (below detection limit) data were kept in the database but were not included in the following analyses. For those positive-value data, we applied the Chauvenet's criterion to their log-transformed values to identify outliers (Glover et al., 2011): A datum was treated as an outlier when its probability of deviating from the observed mean was lower
110 than $1/(2n)$, where n was the number of data samples. Outliers were marked in the database and not included in the following analyses.

2.6 Total number and biomass of viruses in global ocean

Based on the VA data of our database, we estimated the total number of viruses in the global ocean using two methods. In the first method, we separately estimated the total number of viruses in different ocean basins, including the North and South
115 Atlantic Ocean, the North and South Pacific Ocean, the Indian Ocean, the Arctic Ocean, and the Mediterranean and Baltic Seas. For each basin, viral numbers were calculated for coastal and open oceans separately. The mean VA in each depth bin (coastal oceans: bins separated at 5, 10, 25, 50 and 100 m; open oceans: bins separated at 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 500, 1000 and 2000 m) were multiplied by its water volume to calculate the number of viruses in that bin. Please note that due to the insufficient data, mean VA in deep waters of the Arctic Ocean, Mediterranean Sea and Baltic Sea were
120 substituted by the value of the North Atlantic Ocean. The total number of viruses in global ocean was then calculated by summing up the estimates of the coastal and open ocean regions of all the counted ocean basins.

The second method used the random forest (MATLAB machine learning toolbox) (Breiman, 2001) to construct a model of VA based on sampling latitude, longitude, months and depths. VA data were binned to $1^\circ \times 1^\circ$ with 44 vertical layers and the mean VA of each bin, if data available, was fed into the random forest. When implementing the random forest, 75% samples
125 were randomly selected for training the model while the rest data were used for model validation. The trained model was then used to predict VA for each bin and then to estimate the total viral number in the global ocean.

The viral biomass of the global ocean was calculated from the virus numbers using a conversion factor of 0.023 fg C per viral particle, which was based on an empirical relationship between carbon contents in heads of marine viruses (C_{head}) and their sizes (Jover et al., 2014):

$$130 \quad C_{head} = 41(r - 2.5)^3 + 130(7.5r^2 - 18.75r + 15.63), \quad (1)$$

where r was the radius of viral head for which an average of 26.3 nm from the Tara Ocean expedition data was used (Brum et al., 2015).

In this paper, all the uncertainties reported in parentheses after the means are standard deviations, except that the standard errors of the mean are reported for the estimates of total viral number and biomass of the global ocean, because the mean values are used in the estimates and therefore the uncertainties of the means are the most interested.

3 Results and discussion

3.1 Data distribution

Most VA data were collected in the north hemisphere (particularly in tropical and subtropical regions), while fewer data in the southern hemisphere (Figs. 1a–1c). In total, nearly two-thirds of the VA data were sampled in the North Atlantic Ocean (32%) and North Pacific Ocean (29%) (Fig. 2a). In addition, 6 long-term time-series of VA were included in the compilation (Figs. 1a & 1b): the Bermuda Atlantic Time-series Study (BATS) in 2000–2009, the San Pedro Ocean Time Series (SPOTS) Microbial Observatory in 2000–2011, the Bedford Basin Monitor (BBM) in 1996–2000, the Rivers Inlet (RI) in 2008–2010, the Saanich Inlet (SI) in 2010–2012 and 2014–2015, and the Guanabara Bay (GB) in 2011–2014. Weekly VA were measured at BBM, approximately monthly samples were collected at BATS, SPOTS, SI, and GB year round, and monthly samples were collected at RI only in spring and summer. Vertically, most VA data were sampled in the surface ocean (≤ 200 m, 71%) while fewer data in the deep ocean (> 200 m, 29%) particularly below 1,000 m (Fig. 3a). Summer VA samples were most abundant while the fewest data in winter (Fig. 4a).

Lytic VP data in the north hemisphere are much more than those in the south hemisphere (Figs. 1d–1f), with almost half of lytic VP data were sampled in the North Pacific Ocean (31%) and North Atlantic Ocean (18%) (Fig. 2b). A majority of lytic VP data (86%) was collected in the surface ocean (Fig. 3b), while the deep samples were mostly from the North Atlantic and the western and northeastern Pacific Oceans (Fig. 1e). There were seasonal biases in lytic VP data, most of which were sampled in summer while rarely sampled in autumn (Fig. 4b). More lytic VP data were sampled in open oceans (63%) than in coastal waters (37%) (Fig. 5). Almost every lytic and lysogenic VP data accompanied with VA measurements.

There were very limited number of lysogenic VP data in both surface and deep oceans (Figs. 1g & 1h), with those deep samples being even much fewer than the already scarce surface ones (Fig. 3c). The northern hemisphere had slightly more lysogenic VP data than the southern hemisphere (Fig. 1i), with most lysogenic VP data sampled in the North Pacific (29%), the North Atlantic (24%) and the South Atlantic Oceans (23%) (Fig. 2c). Similar to lytic VP data, lysogenic VP data were tended to be collected in spring and summer than in other seasons particularly winter (Fig. 4c). Lysogenic VP data in the open ocean (77%) were also much more than those in coastal waters (23%) (Fig. 5).

In summary, most viral data were sampled in North Atlantic and Northeast Pacific Oceans (Figs. 1 & 2), and more data in the surface than in the deep oceans (Fig. 3). Viral data also tended to be sampled in summer (Fig. 4). Although the total viral data in the coastal samples were fewer than the open ocean samples (Fig. 5), they were more concentrated in the coastal zones considering their relatively small area in the global ocean.

3.2 Viral abundance in the global ocean

165 In the surface oceans, VA (n=7,768) mostly varied in the order of 10^6 to 10^8 particles ml^{-1} , with mean VA in coastal waters $[3.61(\pm 6.3) \times 10^7 \text{ particles } \text{ml}^{-1}]$ about 5 times higher than that in the open oceans $[7.3(\pm 12.4) \times 10^6 \text{ particles } \text{ml}^{-1}]$ (Figs. 6a & 6b). VA in coastal South Atlantic Ocean and Mediterranean and Baltic Seas was higher than that in other coastal oceans (Fig. 6a). Although the VA across different surface open oceans distributed in similar ranges, the average VA in Pacific (particularly in its southern portion) was higher than those in other basins (Fig. 6b), a pattern previously found in another study (Lara et al., 170 2017) using fewer data than this study. VA decreased with depth, with those in the global deep ocean $[1.26(\pm 2.44) \times 10^6 \text{ particles } \text{ml}^{-1}, n=3,164]$ about one order of magnitude lower than those in the surface (Figs. 7a & 7b). The vertical profiles in different open ocean basins more clearly showed that the VA in the Pacific was higher than that in the Atlantic in surface 1,000 m, while the difference did not exist in deeper oceans (Fig. 7b).

In our database, most VA samples were measured using FCM (7,353, 67.26%) and EFM (3,465, 31.71%), while only 112 175 (1.03%) VA samples were counted using TEM (Table S1). Previous studies have showed that the VA counted using FCM, which became more popular in studies after 2014 (Table S1), had a strong correlation with those using EFM (Brussaard et al., 2010; Marie et al., 1999; Payet and Suttle, 2008). Our data demonstrated that the VA obtained by FCM and EFM methods has consistent results in similar environments. For deep open ocean samples, VA using TEM are substantially lower than those using the other two methods (Fig. 8). But considering much fewer VA data points using TEM than others (Fig. 8 & Table S1), 180 we cannot conclude TEM substantially underestimated VA in the deep water samples. Nevertheless, our database provides references for methodological comparison in the future.

The total number of global ocean viruses estimated by binning the VA data (Figs. 7a & 7b) is $1.56(\pm 0.2) \times 10^{30}$ particles (mean \pm s.e.), which is very close to the estimate of $1.49(\pm 0.14) \times 10^{30}$ particles (mean \pm s.e.) using the random forest method (Fig. S1). Both values are consistent to the previous estimates of 10^{30} (Suttle, 2007), 1.29×10^{30} (Cobian Guemes et al., 2016) 185 and 1.5×10^{30} (Bar-On and Milo, 2019) viral particles for the global ocean. Using a conversion factor of 0.023 fg C per viral particle (see Methods), our two values of total viral number give the estimates of total viral biomass in the global ocean at 35.9 ± 0.46 and 34.4 ± 0.32 Tg C, respectively, confirming a recent estimate of 30 Tg C (Bar-On and Milo, 2019).

3.3 Viral production

In surface ocean, lytic VP (n=523) varied greatly from 10^3 to 10^7 particles $\text{ml}^{-1} \text{ h}^{-1}$ in different ocean basins (Figs. 6c & 6d). 190 The overall mean and standard deviation of lytic VP in the global ocean were $9.87(\pm 24.16) \times 10^5$ (ranging in $0.00746 \times 10^5 - 350 \times 10^5$) particles $\text{ml}^{-1} \text{ h}^{-1}$. Lytic VP values in surface open Pacific Ocean were about one order of magnitude higher than those in surface open Atlantic Ocean (Fig. 6d), a pattern consistent to VA (Fig. 6d). Lytic VP in the surface Arctic Ocean was much lower than that in other basins, which was expected considering its much lower biological productivity (Figs. 6c & 6d). Although insufficient lytic VP data (n=82) were available for meaningful statistical analyses in the deep waters (Fig. 9), the

195 existing data showed a general trend that VP decreased by one order of magnitude from the surface to the deep open oceans (Fig. 7c). Unlike VA, average lytic VP in coastal waters was close to that in the surface open ocean (Fig. 6c).

Most of the lytic VP (84.4%) in this database was estimated by VRA, suggesting that VRA was widely utilized in literature and became a standard method to estimate VP across different marine environments. Several studies have tried to compare different approaches estimating the lytic VP, revealing that the VRA method was more reliable and less laborious, compared to the probable overestimation by FLVT approach and the potential underestimation by RIA method, though such comparisons were mainly constrained to the coastal ocean (Helton et al., 2005; Karuza et al., 2010; Rastelli et al., 2016; Winget et al., 2005). Additionally, although a meaningful comparison of reported lytic VP values between disparate marine ecosystems is complicated by the inherent variability among approaches, the lytic VP rates in this database might provide a tentative global-scale insight into methodological comparison. Our statistics showed that, in similar environments, the lytic VP rates determined by FLVT and VRA were higher than those measured by RIA. For coastal samples, such difference among methods was not obvious (Fig. 9). However, due to the limited number of samples using the methods other than VRA (Fig. 9 and Table S2), we did not have adequate data to tell if the difference in VP was caused by the measurement methods, or the randomness of the samples. Hence, more measurements of lytic VP using multiple approaches simultaneously will be certainly needed to better evaluate the differences among them.

210 The lysogenic VP data were too few (surface n=85, deep ocean n= 34) for meaningful comparisons across different ocean basins or between the surface and deep waters, although the results were plotted for readers' reference (Figs. 6e & 6f). The overall lysogenic VP in the global ocean was estimated at $2.53(\pm 8.64) \times 10^5$ (ranging in $0.00132 \times 10^5 - 68.8 \times 10^5$) particles $\text{ml}^{-1} \text{h}^{-1}$, which was about one third of the level of lytic VP, although more data will be needed to better compare the two types of VP.

215 4 Data availability

The gVOD database (Xie et al., 2020) can be downloaded from PANGAEA at <https://doi.org/10.1594/PANGAEA.915758>.

5 Conclusion

We constructed a global ocean viral database (gVOD) by compiling 10,931 VA data, 608 lytic VP data and 119 lysogenic VP data. This database may be useful for global-scale studies of viral processes and their roles in marine ecosystems and biogeochemical cycles. The VA, lytic and lysogenic VP data were greatly variable. Most VA were counted using flow cytometers and epifluorescence microscopes, while the virus reduction approach was the most popular method in estimating VP. The lytic VP is about 3 times higher than the lysogenic VP. The calculation using the database also confirms the previous estimates of viral numbers and biomass in the global ocean.

Our database shows that the current investigations have the limitation in spatiotemporal coverage. The VA dataset has a poor coverage in South Pacific and Indian Ocean. The lytic VP dataset does not have a good coverage in South Pacific, Northwest Pacific, Indian and South Atlantic Oceans. The lysogenic VP data are very few in the global ocean. Vertically, all viral data were sampled much less in mesopelagic and deep oceans than in the surface oceans. Thus, the measurements of viral parameters in these regions and depths should be given high priority. In addition, more viral data should be sampled in winter to avoid seasonal biases.

The database is stored in a public data repository (PANGAEA), and will be updated regularly when new data become available. We hope that the database will be valuable for field and modelling studies in marine ecology, biogeochemistry and other areas of oceanography.

Author contributions.

RZ and YWL conceived and designed structure of database and mathematical analyses of the data. LX, WW, LC, XC and YH collected the data and described the metadata. LX, NJ, RZ and YWL conducted quality control and analyses of the data. LX, RZ and YWL led the writing of the paper, with contribution from all the co-authors.

Competing interests.

The authors declare that they have no conflict of interest.

Acknowledgements.

We would like to thank all the scientists and crews who collected the historical data. This work was funded by National Natural Science Foundation of China (91951209, 41890802, 42076153) and the MEL internal research fund (MELRI2003).

Reference

- Akaike, H.: A new look at the statistical model identification, IEEE transactions on automatic control, 19, 716–723, https://doi.org/10.1007/978-1-4612-1694-0_16, 1974.
- Bar-On, Y. M. and Milo, R.: The biomass composition of the oceans: a blueprint of our blue planet, Cell, 179, 1451–1454, <https://doi.org/10.1016/j.cell.2019.11.018>, 2019.
- Breiman, L.: Random forests, Machine Learning, 45, 5–32, 10.1023/a:1010933404324, 2001.

- Breitbart, M.: Marine viruses: truth or dare, *Ann Rev Mar Sci*, 4, 425–448, <https://doi.org/10.1146/annurev-marine-120709-142805>, 2012.
- 250 Brum, J. R., Ignacio-Espinoza, J. C., Roux, S., Doulier, G., Acinas, S. G., Alberti, A., Chaffron, S., Cruaud, C., de Vargas, C., Gasol, J. M., Gorsky, G., Gregory, A. C., Guidi, L., Hingamp, P., Iudicone, D., Not, F., Ogata, H., Pesant, S., Poulos, B. T., Schwenck, S. M., Speich, S., Dimier, C., Kandels-Lewis, S., Picheral, M., Searson, S., Bork, P., Bowler, C., Sunagawa, S., Wincker, P., Karsenti, E., and Sullivan, M. B.: Patterns and ecological drivers of ocean viral communities, *Science*, 348, 1261498, <https://doi.org/10.1126/science.1261498>, 2015.
- 255 Brussaard, C. P.: Optimization of procedures for counting viruses by flow cytometry, *Appl Environ Microbiol*, 70, 1506-1513, 10.1128/aem.70.3.1506-1513.2004, 2004.
- Brussaard, C. P. D., Payet, J. P., Winter, C., and Weinbauer, M. G.: Quantification of aquatic viruses by flow cytometry. In: *Manual of aquatic viral ecology*, <https://doi.org/10.4319/mave.2010.978-0-9845591-0-7.102>, 2010.
- Chen, X., Wei, W., Wang, J., Li, H., Sun, J., Ma, R., Jiao, N., and Zhang, R.: Tide driven microbial dynamics through virus-
 260 host interactions in the estuarine ecosystem, *Water Res*, 160, 118–129, <https://doi.org/10.1016/j.watres.2019.05.051>, 2019.
- Cobian Guemes, A. G., Youle, M., Cantu, V. A., Felts, B., Nulton, J., and Rohwer, F.: Viruses as winners in the game of life, *Annu Rev Virol*, 3, 197–214, <https://doi.org/10.1146/annurev-virology-100114-054952>, 2016.
- Fuhrman, J. A.: Marine viruses and their biogeochemical and ecological effects, *Nature*, 399, 541-548, 10.1038/21119, 1999.
- Glover, D. M., Jenkins, W. J., and Doney, S. C.: *Modeling Methods for Marine Science*, Cambridge University Press,
 265 10.1017/cbo9780511975721, 2011.
- Heldal, M. and Bratbak, G.: Production and Decay of Viruses in Aquatic Environments, *Marine Ecology Progress Series*, 72, 205-212, DOI 10.3354/meps072205, 1991.
- Helton, R. R., Cottrell, M. T., Kirchman, D. L., and Wommack, K. E.: Evaluation of incubation-based methods for estimating virioplankton production in estuaries, *Aquatic Microbial Ecology*, 41, 209-219, DOI 10.3354/ame041209, 2005.
- 270 Jover, L. F., Effler, T. C., Buchan, A., Wilhelm, S. W., and Weitz, J. S.: The elemental composition of virus particles: implications for marine biogeochemical cycles, *Nat Rev Microbiol*, 12, 519–528, <https://doi.org/10.1038/nrmicro3289>, 2014.


- Karuzza, A., Del Negro, P., Crevatin, E., and Fonda Umani, S.: Viral production in the Gulf of Trieste (Northern Adriatic Sea): Preliminary results using different methodological approaches, J Exp Mar Biol Ecol, 383, 96–104, <https://doi.org/10.1016/j.jembe.2009.12.003>, 2010.
- Lara, E., Vaqu  D., S  E. L., Boras, J. A., Gomes, A., Borrull, E., D z-Vives, C., Teira, E., Pernice, M. C., Garcia, F. C., Forn, I., Castillo, Y. M., Peir  A., Salazar, G., Mor n, X. A. G., Massana, R., Catal  T. S., Luna, G. M., Agust  S., Estrada, M., Gasol, J. M., and Duarte, C. M.: Unveiling the role and life strategies of viruses from the surface to the dark ocean, Sci Adv, 3, e1602565, <https://doi.org/10.1126/sciadv.1602565>, 2017.
- Marie, D., Brussaard, C. P. D., Thyraug, R., Bratbak, G., and Vaultot, D.: Enumeration of marine viruses in culture and natural samples by flow cytometry, Appl Environ Microbiol, 65, 45-52, 10.1128/AEM.65.1.45-52.1999, 1999.
- Noble, R. T. and Fuhrman, J. A.: Rapid virus production and removal as measured with fluorescently labeled viruses as tracers, Appl Environ Microbiol, 66, 3790-3797, 10.1128/aem.66.9.3790-3797.2000, 2000.
- Noble, R. T. and Fuhrman, J. A.: Use of SYBR Green I for rapid epifluorescence counts of marine viruses and bacteria, Aquatic Microbial Ecology, 14, 113-118, DOI 10.3354/ame014113, 1998.
- Noble, R. T. and Steward, G. F.: Estimating viral proliferation in aquatic samples, Methods in Microbiology, Vol 30, 30, 67-82, Doi 10.1016/S0580-9517(01)30040-5, 2001.
- Paul, J. H.: Prophages in marine bacteria: dangerous molecular time bombs or the key to survival in the seas?, ISME J, 2, 579–589, <https://doi.org/10.1038/ismej.2008.35>, 2008.
- Payet, J. P. and Suttle, C. A.: Physical and biological correlates of virus dynamics in the southern Beaufort Sea and Amundsen Gulf, J Marine Syst, 74, 933–945, <https://doi.org/10.1016/j.jmarsys.2007.11.002>, 2008.
- Payet, J. P. and Suttle, C. A.: To kill or not to kill: The balance between lytic and lysogenic viral infection is driven by trophic status, Limnol Oceanogr, 58, 465–474, <https://doi.org/10.4319/lm.2013.58.2.0465>, 2013.
- Rastelli, E., Dell'Anno, A., Corinaldesi, C., Middelboe, M., Noble, R. T., and Danovaro, R.: Quantification of viral and prokaryotic production rates in benthic ecosystems: A methods comparison, Front Microbiol, 7, 1501, <https://doi.org/10.3389/fmicb.2016.01501>, 2016.
- Suttle, C. A.: Marine viruses – major players in the global ecosystem, Nat Rev Microbiol, 5, 801–812, <https://doi.org/10.1038/nrmicro1750>, 2007.

- Suttle, C. A.: Viruses in the sea, *Nature*, 437, 356-361, 10.1038/nature04160, 2005.
- Weinbauer, M., Rowe, J., and Wilhelm, S.: Determining rates of virus production in aquatic systems by the virus reduction approach. In: *Manual of Aquatic Viral Ecology*, <https://doi.org/10.4319/mave.2010.978-0-9845591-0-7.1>, 2010.
- Weinbauer, M. G.: Ecology of prokaryotic viruses, *FEMS Microbiol Rev*, 28, 127-181, 10.1016/j.femsre.2003.08.001, 2004.
- Weinbauer, M. G., Brettar, I., and Hofle, M. G.: Lysogeny and virus-induced mortality of bacterioplankton in surface, deep, and anoxic marine waters, *Limnology and Oceanography*, 48, 1457-1465, DOI 10.4319/lo.2003.48.4.1457, 2003.
- Weinbauer, M. G., Winter, C., and Hofle, M. G.: Reconsidering transmission electron microscopy based estimates of viral infection of bacterio-plankton using conversion factors derived from natural communities, *Aquatic Microbial Ecology*, 27, 103-110, DOI 10.3354/ame027103, 2002.
- Wells, L. E. and Deming, J. W.: Significance of bacterivory and viral lysis in bottom waters of Franklin Bay, Canadian Arctic, during winter, *Aquatic Microbial Ecology*, 43, 209-221, DOI 10.3354/ame043209, 2006.
- Wigington, C. H., Sonderegger, D., Brussaard, C. P. D., Buchan, A., Finke, J. F., Fuhrman, J. A., Lennon, J. T., Middelboe, M., Suttle, C. A., Stock, C., Wilson, W. H., Wommack, K. E., Wilhelm, S. W., and Weitz, J. S.: Re-examination of the relationship between marine virus and microbial cell abundances, *Nat Microbiol*, 1, 15024, <https://doi.org/10.1038/nmicrobiol.2015.24>, 2016.
- Wilhelm, S. W. and Suttle, C. A.: Viruses and Nutrient Cycles in the Sea - Viruses play critical roles in the structure and function of aquatic food webs, *Bioscience*, 49, 781-788, Doi 10.2307/1313569, 1999.
- Winget, D. M., Williamson, K. E., Helton, R. R., and Wommack, K. E.: Tangential flow diafiltration: an improved technique for estimation of virioplankton production, *Aquatic Microbial Ecology*, 41, 221-232, DOI 10.3354/ame041221, 2005.
- Wommack, K. E. and Colwell, R. R.: Virioplankton: viruses in aquatic ecosystems, *Microbiol Mol Biol Rev*, 64, 69-114, 10.1128/mmbr.64.1.69-114.2000, 2000.
- Xie, L., Wei, W., Cai, L., Chen, X., Huang, Y., Zhang, R., and Luo, Y.-W.: A global viral oceanography database (gVOD) from 1987 to 2018, *PANGAEA*, <https://doi.org/10.1594/PANGAEA.915758>, 2020.
- Zhang, R., Weinbauer, M. G., and Qian, P. Y.: Viruses and flagellates sustain apparent richness and reduce biomass accumulation of bacterioplankton in coastal marine waters, *Environ Microbiol*, 9, 3008-3018, <https://doi.org/10.1111/j.1462-2920.2007.01410.x>, 2007.

Zimina, I. D., Maksimov, A. I., and Svetsov, V. I.: Investigation of excitation of ammonia decomposition products in a high-
325 frequency discharge, Journal of Applied Spectroscopy, 18, 771-772, 10.1007/bf00614106, 1973.

Table 1. Number of accessory viral, host and oceanographic parameters associated with each of the core viral parameters (VA, lytic and lysogenic VP).

	VA (n=10,931)	Lytic VP (n=608)	Lysogenic VP (n=119)
<i>Accessory viral parameters</i>			
Frequency of lytic infection	405	142	36
Frequency of lysogenic infection	227	96	36
Viral decay rate	83	65	27
Burst size	206	96	-
Virus-mediated bacterial mortality	46	46	19
% cells lysed	55	53	17
<i>Accessory host parameters</i>			
Prokaryotic abundance	10,846	598	119
Prokaryotic productivity	2,425	352	91
Flagellate abundance	411	44	7
Picoeukaryotic abundance	1,554	68	15
<i>Synechococcus</i> abundance	1,700	80	42
<i>Prochlorococcus</i> abundance	1,567	73	42
<i>Accessory oceanographic parameters</i>			
Chlorophyll <i>a</i>	3,949	244	71
Temperature	6,253	399	119
Salinity	6,360	370	85
Oxygen	4,930	82	46
Nitrate	2,707	91	46
Phosphate	3,153	144	51
Silicate	2,638	106	36
pH	96	47	-
Light intensity	35	35	-
Dissolved organic carbon	97	13	-



330 **Figure 1. Collected viral abundance (a, b), lytic viral production (d, f) and lysogenic viral production (g, h) in surface (≤ 200 m) (a,**
d and g) and in deep (> 200 m) waters (b, e and h), binned on 1°x1° grids. Color of each grid codes the mean value of the
parameters, and the size of the circles represents number of samples in each bin. Also shown the number of samples in latitudinal
bands (c, f and i). Numbers in (a) represent long-term time-series observations of viral abundance: 1, Rivers Inlet; 2: Saanich
Inlet; 3: San Pedro Ocean Time Series Microbial Observatory; 4: Bedford Basin Monitor; 5: Bermuda Atlantic Time-series Study
335 **and 6: Guanabara Bay.**




Figure 2. The fraction of (a) viral abundance, (b) lytic viral production and (c) lysogenic viral production data points in different oceans (N. Atl: North Atlantic, S. Atl: South Atlantic, N. Pac: North Pacific, S. Pac: South Pacific, Ind: Indian ocean, Arc: Arctic ocean, M & B: Mediterranean Sea and Baltic Sea).




Figure 3. Depth distribution of the data for (a) viral abundance, (b) lytic viral production, and (c) lysogenic viral production.







Figure 4. Seasonal distributions of number of samples for (a) viral abundance, (b) lytic viral production and (c) lysogenic viral production.



350 **Figure 5. The fraction of viral abundance (VA), lytic viral production (VP) and lysogenic VP data points in coastal versus in open oceans.**




355 **Figure 6.** The range of (a, d) viral abundance, (b, e) lytic viral production, and (c, f) lysogenic viral production in the different
oceans (N. Atl: North Atlantic, S. Atl: South Atlantic, N. Pac: North Pacific, S. Pac: South Pacific, Ind: Indian ocean, Arc: Arctic
ocean, M & B: Mediterranean Sea and Baltic Sea), grouping in coastal ocean (a–c) and open ocean samples in surface 200 m (d–f).
All data are shown in logarithmic scales. The red diamonds mark the mean value. The central red lines indicate the median, and
the bottom and top edges of the box indicate the 25th and 75th percentiles of the data. Error bars extend to the 5th and 95th
360 percentiles and the remaining outliers are marked with red plus signs.



365

Figure 7. Vertical profiles of average viral abundance of each ocean basin in (a) coastal and (b) open ocean waters. Also shown the vertical profiles for open water lytic viral production (c), while those for coastal samples were not constructed because of limited data points. Error bars represent one standard error of the mean.



370

Figure 8. Box plots of distribution of viral abundance using different measurement methods (EFM, FCM and TEM, see text for more details) in coastal (a), surface open ocean (b) and deep open ocean (c) waters. See caption of Figure 6 for details of lines and symbols of the box plots.




Figure 9. Comparison of lytic viral production using different measurement methods including VRA, RIA, FPB, FLVT and VDR (FPB: calculated by multiplying fraction of viral infected cells, prokaryotic production and burst size; RIA: radioactive incorporation approach; FLVT: fluorescently labelled viral tracers method; VRA: virus reduction approach; VDR: estimated by viral decay rates. See text for details) in coastal (a), surface open ocean (b) and deep open ocean (c) waters. See caption of Figure 6 for details of lines and symbols of the box plots.