

# **A global viral oceanography database (gVOD)**

Le Xie<sup>1</sup>, Wei Wei<sup>1</sup>, Lanlan Cai<sup>2</sup>, Xiaowei Chen<sup>1</sup>, Yuhong Huang<sup>1</sup>, Nianzhi Jiao<sup>1</sup>, Rui Zhang<sup>1,\*</sup>, Ya-Wei Luo<sup>1,\*</sup>

<sup>1</sup>State Key Laboratory of Marine Environmental Science, College of Ocean and Earth Sciences, Xiamen University, Xiamen,  
5 Fujian 361102, China

<sup>2</sup>Department of Ocean Science, The Hong Kong University of Science and Technology, Clear Water Bay, Hong Kong, China

*Correspondence to:* Rui Zhang (ruizhang@xmu.edu.cn) and Ya-Wei Luo (ywluo@xmu.edu.cn)

**Abstract.** Virioplankton are a key component of the marine biosphere in maintaining diversity of microorganisms and stabilizing ecosystems. They also contribute greatly to nutrient cycles/cycling by releasing organic matter after lysis of hosts. In this study, we constructed the first global viral oceanography database (gVOD) by collecting 10,931 viral abundance (VA) data and 727 viral production (VP) data, along with host and relevant oceanographic parameters when available. Most VA data were obtained in the North Atlantic (32%) and North Pacific Oceans (29%), while the Southeast Pacific and Indian Oceans were quite under sampled. The VA in the global ocean was  $1.17(\pm 3.31) \times 10^7$  particles  $\text{ml}^{-1}$ . The lytic and lysogenic VP in the global ocean was  $9.87(\pm 24.16) \times 10^5$  and  $2.53(\pm 8.64) \times 10^5$  particles  $\text{ml}^{-1} \text{h}^{-1}$ , respectively. Average VA in coastal oceans was higher than that in surface open oceans [ $3.61(\pm 6.30) \times 10^7$  versus  $0.73(\pm 1.24) \times 10^7$  particles  $\text{ml}^{-1}$ ], while average VP in coastal and surface open oceans was close. Vertically, VA, lytic and lysogenic VP decreased from surface to deep oceans by about one order of magnitude. The total number of viruses in the global ocean estimated by bin-averaging and the random forest method was  $1.56 \times 10^{30}$  particles and  $1.49 \times 10^{30}$  particles, leading to an estimate of global ocean viral biomass at 35.9 and 34.4 Tg C, respectively. We expect that the gVOD will be a fundamental and very useful database for laboratory, field and modelling studies in marine ecology and biogeochemistry. The full gVOD database (Xie et al., 2020) is stored in PANGAEA (<https://doi.org/10.1594/PANGAEA.915758>).

## 1 Introduction

Virioplankton are the most abundant biological entities and one of the largest genetic reservoirs in the ocean (Breitbart, 2012; Fuhrman, 1999). With an estimation of  $\sim 10^{23}$  marine microbes being infected every second, viruses play important roles in affecting microbial mortality, regulating community composition and impacting biogeochemical cycles (Suttle, 2005; Zhang et al., 2007). Viruses were estimated to kill  $\sim 20$ – $40\%$  of marine bacterioplankton every day, a rate similar to that caused by zooplankton grazing (Fuhrman, 1999). In particular, virus-mediated cell lysis effectively 'shunts' approximately 25% of the photosynthetically fixed carbon, which otherwise would be transferred to higher trophic levels, to the dissolved organic matter (DOM) pool, partly forming the basis of the microbial loop and leading to the recycling of nutrients (Suttle, 2007; Wilhelm and Suttle, 1999). Furthermore, viral lysis can contribute to biological pump through the release of sticky lysates that accelerate the aggregation and sink of carbon into the deep sea (Suttle, 2005).

Compilation of the observations of viral abundance and activity in the global ocean is very necessary and urgent in understanding spatiotemporal distributions of viruses, exploring the controlling factors of viral processes, qualitatively and quantitatively assessing virus-host interactions and viral functioning in marine ecosystems, and even improving predictions of large-scale marine ecosystem and Earth system models. Previous two studies (Bar-On and Milo, 2019; Wigington et al., 2016) summarized viral abundance data in the ocean and estimated viral biomass as well as virus-to-prokaryote ratio. However, the lacking of host parameters such as bacterial production, and oceanographic parameters such as temperature, salinity, nutrient concentrations, limits the usage of these datasets in broader oceanographic contexts. More importantly, there is no public database of viral activity in the global ocean, which substantially hinders our understandings of the ecological and

biogeochemical roles of viroplankton on global scale. In addition, the ecological functions of the viruses are tightly linked to their life strategies, mainly including the lytic and the lysogenic infection (Wommack and Colwell, 2000). The significance of viruses in oceanic biogeochemistry is mainly reflected through the lytic infection, which results in cell lysis and the release of DOM. In contrast, other more temperate viruses choosing the lysogenic infection can influence microbial diversity and metabolism by transferring new genes to their hosts, altering the expression of host genes, and not killing hosts for many generations until an environmental or cellular trigger causes them to enter the lytic cycle. The lysogenic infection hence serves as a molecular “time bomb” (Paul, 2008). Therefore, it is necessary to include the quantity and quality data of viral life strategies in a viral oceanographic database.

In this study, we construct the first global viral oceanography database, namely gVOD, by collecting data of viral abundance (VA), lytic and lysogenic viral production (VP), as well as other related viral, host and oceanographic metadata when available. Based on the database, we estimate the total viral number and biomass in the global ocean. In addition, the data of VA and VP generated with different techniques were compared to provide references for evaluating possible technical biases.

## 2 Data and methods

### 2.1 Database summary

In the gVOD, direct measurements of three core parameters (VA, lytic and lysogenic VP), as well as accessory viral, prokaryotic and oceanographic parameters when available, were collected from published papers or acquired from lead authors or principal investigators (Table 1). Sampling information including date, latitude, longitude, depth and methods was included for each data record. We used ocean depth shallower or deeper than 200 m as a criterion to identify coastal or open ocean samples. The open ocean samples were further separated into surface and deep samples that collected in 0–200 m and >200 m, respectively.

The quality-controlled database consists of 10,931 VA data points (Appendix Table A1), 608 lytic VP data points and 119 lysogenic VP data points (Appendix Table A2). Most of VA (99.2%) and lytic VP (98.4%) and all lysogenic VP data have accompanying data of prokaryotic abundance (Table 1). For some samples, the abundances of flagellate, picoeukaryotes, *Synechococcus* and *Prochlorococcus* are also available. Prokaryotic productivity measurements cover 22.1% of VA, 57.7% of lytic VP and 76.5% of lysogenic VP data. The most available environmental parameters are salinity and temperature, providing oceanographic information for about half of VA, two-thirds of lytic VP and nearly all lysogenic VP data. Oxygen and chlorophyll *a* concentration data are also adequate particularly for VA. The concentrations of different types of nutrient, including nitrate, silicate and phosphate, are available for many samples. Other environmental parameters (pH, light intensity, dissolved organic carbon concentration) are relatively scarce. Moreover, given that the frequency of viral infected cells was calculated, independently or together with VP, usually to quantify the impact of viral infection within the microbial community (Chen et al., 2019; Payet and Suttle, 2013; Weinbauer et al., 2003), the reported frequencies of lytic infection ( $n = 438$ ) and lysogenic infection ( $n = 266$ ) in the literature were also collected into the database to facilitate the future exploration of marine

viral activities. Lastly, we collected 83 viral decay rate data, 206 viral burst size data and 111 virus-mediated mortality data, which can be useful for certain studies. The gVOD is a compilation of all the available data, to our best knowledge, by 2019.

75 We plan to update the database every 5 years.

## 2.2 Viral abundance

The viral abundance in this database was counted using one of the following three methods. In the first method, viruses were harvested by ultracentrifuging onto copper grids and stained with uranyl acetate, and then enumerated using a transmission electron microscopy (TEM) (Akaike, 1974). In the second method, viruses were collected onto 0.02- $\mu\text{m}$  filters and stained with a nucleic acid-specific fluorescent dye (e.g., SYBR Green I), and then were counted under an epifluorescence microscope (EFM) (Noble and Fuhrman, 1998). The third method counted viruses by using flow cytometers (FCM), before which viruses were stained with fluorescent dye (e.g., SYBR Green I or SYBR Gold), and identified on the basis of the green fluorescence versus side scatter signal (Brussaard, 2004; Marie et al., 1999). The details of these three approaches have been described elsewhere (Weinbauer, 2004).

## 85 2.3 Lytic viral activity

Lytic VP is paramount and widely employed to assess the activity of lytic viruses in community-level and the roles of viruses in marine ecosystems. In this database, the lytic VP was estimated by one of the following five methods. The first method estimated VP by calculating expected viral release rates by multiplying fraction of viral infected cells (mainly prokaryotes), prokaryotic productivity (assuming equal prokaryotic mortality rate) and burst size obtained from TEM studies (Noble and Steward, 2001) or virus-dilution approach (Weinbauer et al., 2002). For notational simplicity, in this paper we label this method as FPB to represent the three variables (Fraction of viral infected cells, Prokaryotic productivity, and Burst size) listed above and used in the estimation. In the second method, called radioactive incorporation approach (RIA), lytic VP was estimated by determining viral DNA synthesis rates using a labelled radiotracer (e.g.,  $^3\text{H}$ -,  $^{32}\text{P}$ -, or  $^{14}\text{C}$ -labeled thymidine or leucine) and a conversion factor to quantify the incorporated radiotracer into viral particles (Noble and Steward, 2001; Zimina et al., 1973). The third method estimated the lytic VP from the viral decay rates (VDR), assuming that the abundance of virus particles is in steady state and then the loss rate of virus particles should be balanced by the production rate (Heldal and Bratbak, 1991). The fourth method used fluorescently labelled viral tracers (FLVT) to measure the dilution rates from the decay of labelled viruses and net changes of the non-labelled viruses in natural viral community (Noble and Fuhrman, 2000). The fifth method quantified the increase of viral abundance during time course incubation using a virus dilution or virus reduction approach (VRA) (Weinbauer et al., 2010; Winget et al., 2005), which effectively avoided new viral infection by reducing viral abundances using pore-size filters or tangential flow filtration systems.

## 2.4 Lysogenic viral activity

Lysogenic VP is generally measured by detecting the proviruses (temperate viruses) that choose lysogenic infection in the environment. Lysogenic VP in this database was estimated using VRA described above after the provirus induction by Mitomycin C (Weinbauer et al., 2002). Hence, the lysogenic VP was estimated as the difference in viral abundance per unit time between the Mitomycin C treated and the control samples.

## 2.5 Quality control

We conducted quality control for the VA, lytic and lysogenic VP data of the database. A negative lytic VP (Wells and Deming, 2006) was removed. All zero-value (below detection limit) data were kept in the database but were not included in the following analyses. For those positive-value data, we applied the Chauvenet's criterion to their log-transformed values to identify outliers (Glover et al., 2011): A datum was treated as an outlier when its probability of deviating from the observed mean was lower than  $1/(2n)$ , where  $n$  was the number of data samples. Outliers were marked in the database and not included in the following analyses.

## 2.6 Total number and biomass of viruses in global ocean

Based on the VA data of our database, we estimated the total number of viruses in the global ocean using two methods. In the first method, we separately estimated the total number of viruses in different ocean basins, including the North and South Atlantic Ocean, the North and South Pacific Ocean, the Indian Ocean, the Arctic Ocean, and the Mediterranean and Baltic Seas. For each basin, viral numbers were calculated for coastal and open oceans separately. The mean VA in each depth bin (coastal oceans: bins separated at 5, 10, 25, 50 and 100 m; open oceans: bins separated at 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 500, 1000 and 2000 m) were multiplied by its water volume to calculate the number of viruses in that bin. Please note that due to the insufficient data, mean VA in deep waters of the Arctic Ocean, Mediterranean Sea and Baltic Sea were substituted by the value of the North Atlantic Ocean. The total number of viruses in global ocean was then calculated by summing up the estimates of the coastal and open ocean regions of all the counted ocean basins.

The second method used the random forest (MATLAB machine learning toolbox) (Breiman, 2001) to construct a model of VA based on sampling latitude, longitude, months and depths. VA data were binned to  $1^\circ \times 1^\circ$  with 44 vertical layers and the mean VA of each bin, if data available, was fed into the random forest. When implementing the random forest, 75% samples were randomly selected for training the model while the rest data were used for model validation. The trained model was then used to predict VA for each bin and then to estimate the total viral number in the global ocean.

The viral biomass of the global ocean was calculated from the virus numbers using a conversion factor of 0.023 fg C per viral particle, which was based on an empirical relationship between carbon contents in heads of marine viruses ( $C_{head}$ ) and their sizes (Jover et al., 2014):

$$C_{head} = 41(r - 2.5)^3 + 130(7.5r^2 - 18.75r + 15.63), \quad (1)$$

where  $r$  was the radius of viral head for which an average of 26.3 nm from the Tara Ocean expedition data was used (Brum et al., 2015).

135 In this paper, all the uncertainties reported in parentheses after the means are standard deviations, except that the standard errors of the mean are reported for the estimates of total viral number and biomass of the global ocean, because the mean values are used in the estimates and therefore the uncertainties of the means are the most interested.

### 3 Results and discussion

#### 3.1 Data distribution

140 Most VA data were collected in the north hemisphere (particularly in tropical and subtropical regions), while fewer data in the southern hemisphere (Figs. 1a–1c). In total, nearly two-thirds of the VA data were sampled in the North Atlantic Ocean (32%) and North Pacific Ocean (29%) (Fig. 2a). In addition, 6 long-term time-series of VA were included in the compilation (Figs. 1a & 1b): the Bermuda Atlantic Time-series Study (BATS) in 2000–2009, the San Pedro Ocean Time Series (SPOTS) Microbial Observatory in 2000–2011, the Bedford Basin Monitor (BBM) in 1996–2000, the Rivers Inlet (RI) in 2008–2010, 145 the Saanich Inlet (SI) in 2010–2012 and 2014–2015, and the Guanabara Bay (GB) in 2011–2014. Weekly VA were measured at BBM, approximately monthly samples were collected at BATS, SPOTS, SI, and GB year-round, and monthly samples were collected at RI only in spring and summer. Vertically, most VA data were sampled in the surface ocean ( $\leq 200$  m, 71%) while fewer data in the deep ocean ( $> 200$  m, 29%) particularly below 1,000 m (Fig. 3a). Summer VA samples were most abundant while the fewest data in winter (Fig. 4a).

150 Lytic VP data in the north hemisphere are much more than those in the south hemisphere (Figs. 1d–1f), with almost half of lytic VP data were sampled in the North Pacific Ocean (31%) and North Atlantic Ocean (18%) (Fig. 2b). A majority of lytic VP data (86%) was collected in the surface ocean (Fig. 3b), while the deep samples were mostly from the North Atlantic and the western and northeastern Pacific Oceans (Fig. 1e). There were seasonal biases in lytic VP data, most of which were sampled in summer while rarely sampled in autumn (Fig. 4b). More lytic VP data were sampled in open oceans (63%) than in coastal 155 waters (37%) (Fig. 5). Almost every lytic and lysogenic VP data accompanied with VA measurements.

There were very limited number of lysogenic VP data in both surface and deep oceans (Figs. 1g & 1h), with those deep samples being even much fewer than the already scarce surface ones (Fig. 3c). The northern hemisphere had slightly more lysogenic VP data than the southern hemisphere (Fig. 1i), with most lysogenic VP data sampled in the North Pacific (29%), the North Atlantic (24%) and the South Atlantic Oceans (23%) (Fig. 2c). Similar to lytic VP data, lysogenic VP data were 160 tended to be collected in spring and summer than in other seasons particularly winter (Fig. 4c). Lysogenic VP data in the open ocean (77%) were also much more than those in coastal waters (23%) (Fig. 5).

In summary, most viral data were sampled in North Atlantic and Northeast Pacific Oceans (Figs. 1 & 2), and more data in the surface than in the deep oceans (Fig. 3). Viral data also tended to be sampled in summer (Fig. 4). Although the total viral

data in the coastal samples were fewer than the open ocean samples (Fig. 5), they were more concentrated in the coastal zones  
165 considering their relatively small area in the global ocean.

### 3.2 Viral abundance in the global ocean

In the surface oceans, VA (n=7,768) mostly varied in the order of  $10^6$  to  $10^8$  particles  $\text{ml}^{-1}$ , with mean VA in coastal waters  
[ $3.61(\pm 6.3) \times 10^7$  particles  $\text{ml}^{-1}$ ] about 5 times higher than that in the open oceans [ $7.3(\pm 12.4) \times 10^6$  particles  $\text{ml}^{-1}$ ] (Figs. 6a &  
6b). VA in coastal South Atlantic Ocean and Mediterranean and Baltic Seas was higher than that in other coastal oceans (Fig.  
170 6a). Although the VA across different surface open oceans distributed in similar ranges, the average VA in Pacific (particularly  
in its southern portion) was higher than those in other basins (Fig. 6b), a pattern previously found in another study (Lara et al.,  
2017) using fewer data than this study. VA decreased with depth, with those in the global deep ocean [ $1.26(\pm 2.44) \times 10^6$   
particles  $\text{ml}^{-1}$ , n=3,164] about one order of magnitude lower than those in the surface (Figs. 7a & 7b). The vertical profiles in  
different open ocean basins more clearly showed that the VA in the Pacific was higher than that in the Atlantic in surface 1,000  
175 m, while the difference did not exist in deeper oceans (Fig. 7b).

In our database, most VA samples were measured using FCM (7,353, 67.26%) and EFM (3,465, 31.71%), while only 112  
(1.03%) VA samples were counted using TEM (Appendix Table A1). Previous studies have showed that the VA counted using  
FCM, which became more popular in studies after 2014 (Appendix Table A1), had a strong correlation with those using EFM  
(Brussaard et al., 2010; Marie et al., 1999; Payet and Suttle, 2008). Our data demonstrated that the VA obtained by FCM and  
180 EFM methods has consistent results in similar environments. For deep open ocean samples, VA using TEM are substantially  
lower than those using the other two methods (Fig. 8). But considering much fewer VA data points using TEM than others  
(Fig. 8 & Appendix Table A1), we cannot conclude TEM substantially underestimated VA in the deep water samples.  
Nevertheless, our database provides references for methodological comparison in the future.

The total number of global ocean viruses estimated by binning the VA data (Figs. 7a & 7b) is  $1.56(\pm 0.2) \times 10^{30}$  particles  
185 (mean  $\pm$  s.e.), which is very close to the estimate of  $1.49(\pm 0.14) \times 10^{30}$  particles (mean  $\pm$  s.e.) using the random forest method  
(Appendix Figure A1). Both values are consistent to the previous estimates of  $10^{30}$  (Suttle, 2007),  $1.29 \times 10^{30}$  (Cobian Guemes  
et al., 2016) and  $1.5 \times 10^{30}$  (Bar-On and Milo, 2019) viral particles for the global ocean. Using a conversion factor of 0.023 fg  
C per viral particle (see Methods), our two values of total viral number give the estimates of total viral biomass in the global  
ocean at  $35.9 \pm 0.46$  and  $34.4 \pm 0.32$  Tg C, respectively, confirming a recent estimate of 30 Tg C (Bar-On and Milo, 2019).

### 190 3.3 Viral production

In surface ocean, lytic VP (n=523) varied greatly from  $10^3$  to  $10^7$  particles  $\text{ml}^{-1} \text{h}^{-1}$  in different ocean basins (Figs. 6c & 6d).  
The overall mean and standard deviation of lytic VP in the global ocean were  $9.87(\pm 24.16) \times 10^5$  (ranging in  $0.00746 \times 10^5 -$   
 $350 \times 10^5$ ) particles  $\text{ml}^{-1} \text{h}^{-1}$ . Lytic VP values in surface open Pacific Ocean were about one order of magnitude higher than  
those in surface open Atlantic Ocean (Fig. 6d), a pattern consistent to VA (Fig. 6d). Lytic VP in the surface Arctic Ocean was  
195 much lower than that in other basins, which was expected considering its much lower biological productivity (Figs. 6c & 6d).

Although insufficient lytic VP data (n=82) were available for meaningful statistical analyses in the deep waters (Fig. 9), the existing data showed a general trend that VP decreased by one order of magnitude from the surface to the deep open oceans (Fig. 7c). Unlike VA, average lytic VP in coastal waters was close to that in the surface open ocean (Fig. 6c).

Most of the lytic VP (84.4%) in this database was estimated by VRA, suggesting that VRA was widely utilized in literature and became a standard method to estimate VP across different marine environments. Several studies have tried to compare different approaches estimating the lytic VP, revealing that the VRA method was more reliable and less laborious, compared to the probable overestimation by FLVT approach and the potential underestimation by RIA method, though such comparisons were mainly constrained to the coastal ocean (Helton et al., 2005; Karuza et al., 2010; Rastelli et al., 2016; Winget et al., 2005). Additionally, although a meaningful comparison of reported lytic VP values between disparate marine ecosystems is complicated by the inherent variability among approaches, the lytic VP rates in this database might provide a tentative global-scale insight into methodological comparison. Our statistics showed that, in similar environments, the lytic VP rates determined by FLVT and VRA were higher than those measured by RIA. For coastal samples, such difference among methods was not obvious (Fig. 9). However, due to the limited number of samples using the methods other than VRA (Fig. 9 and Appendix Table A2), we did not have adequate data to tell if the difference in VP was caused by the measurement methods, or the randomness of the samples. Hence, more measurements of lytic VP using multiple approaches simultaneously will be certainly needed to better evaluate the differences among them.

The lysogenic VP data were too few (surface n=85, deep ocean n= 34) for meaningful comparisons across different ocean basins or between the surface and deep waters, although the results were plotted for readers' reference (Figs. 6e & 6f). The overall lysogenic VP in the global ocean was estimated at  $2.53(\pm 8.64) \times 10^5$  (ranging in  $0.00132 \times 10^5 - 68.8 \times 10^5$ ) particles  $\text{ml}^{-1} \text{h}^{-1}$ , which was about one third of the level of lytic VP, although more data will be needed to better compare the two types of VP.

#### 4 Data availability

The gVOD database (Xie et al., 2020) can be downloaded from PANGAEA at <https://doi.org/10.1594/PANGAEA.915758>.

#### 5 Code availability

The MATLAB codes for calculating the total number of viruses can be found in the supplementary materials or be obtained by requesting the corresponding authors.



6 Conclusion

We constructed a global ocean viral database (gVOD) by compiling 10,931 VA data, 608 lytic VP data and 119 lysogenic VP data. This database may be useful for global-scale studies of viral processes and their roles in marine ecosystems and biogeochemical cycles. The VA, lytic and lysogenic VP data were greatly variable. Most VA were counted using flow cytometers and epifluorescence microscopes, while the virus reduction approach was the most popular method in estimating VP. The lytic VP is about 3 times higher than the lysogenic VP. The calculation using the database also confirms the previous estimates of viral numbers and biomass in the global ocean.

Our database shows that the current investigations have the limitation in spatiotemporal coverage. The VA dataset has a poor coverage in South Pacific and Indian Ocean. The lytic VP dataset does not have a good coverage in South Pacific, Northwest Pacific, Indian and South Atlantic Oceans. The lysogenic VP data are very few in the global ocean. Vertically, all viral data were sampled much less in mesopelagic and deep oceans than in the surface oceans. Thus, the measurements of viral parameters in these regions and depths should be given high priority. In addition, more viral data should be sampled in winter to avoid seasonal biases.

The database is stored in a public data repository (PANGAEA), and will be updated regularly when new data become available. We hope that the database will be valuable for field and modelling studies in marine ecology, biogeochemistry and other areas of oceanography.

Appendix Tables and Figures

**Appendix Table A1. Sources and methods of viral abundance data. EFM: counted by epifluorescence microscopes. FCM: counted by flow cytometer. TEM: counted by transmission election microscopy. Data marked by \* are those collected in a previous dataset (Wigington et al., 2016).**

Region	Number of data	Method	References
Canadian Arctic Shelf *	259	EFM	(Payet and Suttle, 2013)
Gulf of Alaska, Arctic *	292	EFM	(Balsom, 2003)
Franklin Bay (Arctic Ocean)	4	EFM	(Wells and Deming, 2006)
Greenland Sea, Arctic	79	FCM	(Boras et al., 2010a)
Greenland Sea *	124	EFM	(Wigington et al., 2016)
Arctic Ocean	56	FCM	(Finke et al., 2017)
Kora Sea (Arctic Ocean)	18	EFM	(Kopylov et al., 2019)

North Sea	16	EFM	(Weinbauer et al., 2002)
North Sea	9	EFM	(Winter et al., 2005)
North Sea	39	FCM	(Parada et al., 2008)
North Sea *	191	FCM	(Wigington et al., 2016)
North Sea *	95	FCM	(Wigington et al., 2016)
North Sea	23	FCM	(Winter et al., 2004)
North Atlantic	5	TEM	(Proctor and Fuhrman, 1990)
North Atlantic *	188	FCM	(Li and Dickie, 2001)
North Atlantic	11	EFM	(Auguet et al., 2005)
North Atlantic	6	FCM	(Parada et al., 2007)
North Atlantic	20	TEM	(Bettarel et al., 2008)
North Atlantic	31	EFM	(Rowe et al., 2008)
North Atlantic *	772	FCM	(De Corte et al., 2012)
North Atlantic	9	FCM	(Muck et al., 2014)
North Atlantic *	206	FCM	(Mojica et al., 2015)
North Atlantic	41	EFM	(Parsons et al., 2015)
North Atlantic	10	FCM	(Winter et al., 2018)
North Atlantic	39	FCM	(Finke et al., 2017)
Gulf of mexico	14	TEM	(Jiang and Paul, 1996)
Gulf of Mexico	9	EFM	(Weinbauer and Suttle, 1996)
Gulf of Mexico	12	TEM	(Cochran and Paul, 1998)
Gulf of Mexico	7	EFM	(Weinbauer and Suttle, 1999)
Gulf of Mexico	25	EFM	(Williamson et al., 2002)
Gulf of Mexico	28	EFM	(Long et al., 2008)
Chesapeake Bay	7	EFM	(Winget and Wommack, 2009)
Chesapeake Bay *	84	EFM	(Wang et al., 2011)
Sargasso Sea *	1382	EFM	(Parsons et al., 2012)
Tropical Atlantic	10	EFM	(Winter et al., 2008)
Tropical Atlantic	154	FCM	(De Corte et al., 2010)
Atlantic Ocean	426	FCM	(Lara et al., 2017)

South Atlantic	3	EFM	(Bettarel et al., 2011b)
South Atlantic	172	FCM	(Liang et al., 2014)
South Atlantic	3	FCM	(Gregoracci et al., 2015)
South Atlantic *	430	FCM	(De Corte et al., 2016)
Guanabana Bay (South Atlantic)	246	FCM	(Cabral et al., 2017)
Adriatic (Mediterranean)	35	TEM	(Weinbauer et al., 1993)
Mediterranean	25	EFM	(Bettarel et al., 2002)
Mediterranean and Baltic Sea	30	EFM	(Weinbauer et al., 2003)
Adriatic (Mediterranean)	4	EFM	(Bongiorni et al., 2005)
Mediterranean	48	EFM	(Magagnini et al., 2007)
Mediterranean	24	FCM	(Boras et al., 2009)
Mediterranean	3	EFM	(Motegi et al., 2009)
Mediterranean	43	FCM	(Winter et al., 2009)
Mediterranean	9	EFM	(Fonda Umani et al., 2010)
Mediterranean	6	FCM	(Bouvier and Maurice, 2011)
Mediterranean	10	EFM	(Maurice et al., 2011)
Mediterranean	338	FCM	(Magiopoulos and Pitta, 2012)
Mediterranean	45	EFM	(Maurice et al., 2013)
Mediterranean	1	FCM	(Motegi et al., 2014)
Mediterranean	2	FCM	(Thompson et al., 2014)
Mediterranean	21	EFM	(Ordulj et al., 2017)
Baltic Sea	6	FCM	(Holmfeldt et al., 2010)
Baltic sea	9	FCM	(Kostner et al., 2017)
Baltic sea	4	EFM	(Šulčius et al., 2018)
Indian Ocean *	52	EFM	(Wigington et al., 2016)
Indian Ocean	93	FCM	(Liang et al., 2014)

Cochin Estuary (Indian)	35	EFM	(Parvathi et al., 2013)
Cochin Estuary (Indian)	20	EFM	(Jasna et al., 2017)
Cochin Estuary (Indian)	39	EFM	(Jasna et al., 2018)
Indian Ocean	271	FCM	(Lara et al., 2017)
Indian Ocean	33	EFM	(Parvathi et al., 2018)
Red Sea	51	FCM	(Sabbagh et al., 2020)
Bering Sea	12	TEM	(Steward et al., 1996)
Bering Sea	15	FCM	(Finke et al., 2017)
Santa Monica Bay	7	EFM	(Noble and Fuhrman, 2000)
Japan sea	12	EFM	(Hwang and Cho, 2002)
Masan Bay	24	EFM	(Choi et al., 2003)
North Pacific	36	EFM	(Taylor et al., 2003)
North Pacific (HOT)	8	EFM	(Brum, 2005)
San Pedro Channel (North Pacific)	386	EFM	(Fuhrman et al., 2006)
North Pacific	11	EFM	(Hewson and Fuhrman, 2007)
Strait of Georgia (North Pacific) *	67	EFM	(Clasen et al., 2008)
Bach Dang Estuary (North Pacific)	15	TEM	(Bettarel et al., 2011a)
North Pacific *	355	FCM	(Yang et al., 2014)
Northwestern Pacific	39	FCM	(Li et al., 2014)
Pearl River Estuary (North Pacific)	19	FCM	(Ni et al., 2015)
North Pacific	4	EFM	(Pasulka et al., 2015)
Bohai sea	8	FCM	(Ma et al., 2016)
North Pacific *	9	EFM	(Wigington et al., 2016)
North Pacific	399	FCM	(Finke et al., 2017)

North Pacific	37	EFM	(Gainer et al., 2017)
Western Pacific	244	FCM	(Liang et al., 2017)
Western Pacific	222	FCM	(Zhao et al., 2020)
Jiulong Estuary (North Pacific)	27	FCM	(Chen et al., 2019)
South China Sea	751	FCM	unpublished
South China Sea	6	FCM	(Zhang et al., 2007)
South China Sea	13	FCM	(Chen et al., 2011)
South China Sea	5	EFM	(Nguyen-Kim et al., 2015)
Mariana Trench	6	FCM	(Li et al., 2018)
Pacific Ocean	171	FCM	(Liang et al., 2014)
Pacific Ocean	331	FCM	(Lara et al., 2017)
South Pacific *	12	EFM	(Wilhelm et al., 2003)
South Pacific	31	EFM	(Strzepek et al., 2005)
South Pacific	2	EFM	(Motegei C and Nagata T, 2007)
South Pacific *	24	EFM	(Rowe et al., 2012)
South Pacific	10	EFM	(Bouvy et al., 2012)
South Pacific *	16	EFM	(Matteson et al., 2012)
South Pacific *	542	FCM	(Yang et al., 2014)
Southern Ocean (Indian sector)	14	FCM	(Evans et al., 2009)
Southern Ocean (Indian sector)	10	FCM	(Malits et al., 2014)
Southern Ocean (Atlantic sector) *	33	FCM	(Evans and Brussaard, 2012)
Southern Ocean (Atlantic sector)	25	EFM	(Brum et al., 2016)
Southern Ocean (Pacific sector)	161	FCM	(Vaque et al., 2017)


**Appendix Table A2. Sources and methods of VP data. FPB: Calculated by multiplying fraction of viral infected cells, prokaryotic production and burst size; RIA: radioactive incorporation approach; FLVT: fluorescently labeled viral tracers method; VRA: virus reduction approach (see main text section 2.3 for details). MC: Virus reduction approach with addition of inducing agent such as Mitomycin C (see main text section 2.4 for details).**

Region	#lytic VP	Method of lytic VP	# lysogenic VP	Method of lysogenic VP	References
Canadian Arctic Shelf	14	VRA	-	-	(Payet and Suttle, 2013)
Franklin Bay (Arctic Ocean)	3	VRA	-	-	(Wells and Deming, 2006)
Greenland Sea, Arctic	17	VRA	7	MC	(Boras et al., 2010b)
North	9	VRA	-	-	(Winter et al., 2005)
North Sea	28	VRA	-	-	(Parada et al., 2008)
North Atlantic	26	VRA	-	-	(Rowe et al., 2008)
North Atlantic	18	VRA	10	MC	(De Corte et al., 2012)
North Atlantic	9	VRA	9	MC	(Muck et al., 2014)
North Atlantic	10	VRA	-	-	(Winter et al., 2018)
Chesapeake Bay	7	VRA	-	-	(Winget and Wommack, 2009)
Tropical Atlantic	9	VRA	8	MC	(De Corte et al., 2010)
Baltic Sea	6	VRA	-	-	(Holmfeldt et al., 2010)
Baltic Sea	9	VRA	-	-	(Kostner et al., 2017)
Baltic Sea	4	VRA	-	-	(Šulčius et al., 2018)
Mediterranean	24	VRA	11	MC	(Boras et al., 2009)
Adriatic (Mediterranean)	4	VRA	-	-	(Bongiorni et al., 2005)
Mediterranean	9	VRA	-	-	(Fonda Umani et al., 2010)

Mediterranean	1	VRA	-	-	(Motegi et al., 2014)
Mediterranean	21	VRA	-	-	(Ordulj et al., 2017)
Cochin Estuary (Indian)	35	RIA	-	-	(Parvathi et al., 2013)
Cochin Estuary (Indian))	20	VRA	-	-	(Jasna et al., 2017)
Indian Ocean	10	VRA	-	-	(Parvathi et al., 2018)
Bering Sea	12	FPB	-	-	(Steward et al., 1996)
Santa Monica Bay	7	FLVT	-	-	(Noble and Fuhrman, 2000)
Subtropics Atlantic, Pacific, Indian Ocean	30	VRA	27	MC	(Lara et al., 2017)
North Pacific (HOT)	4	VRA	-	-	(Brum, 2005)
North Pacific	9	VRA	-	-	(Hewson and Fuhrman, 2007)
North Pacific	18	VRA	-	-	(Gainer et al., 2017)
South Pacific	16	VRA	-	-	(Matteson et al., 2012)
Mariana Trench	6	VRA	-	-	(Li et al., 2018)
Western Pacific	38	VRA	-	-	(Li et al., 2014)
Japan Sea	12	FPB	-	-	(Hwang and Cho, 2002)
Masan Bay	24	FPB	-	-	(Choi et al., 2003)
Bach Dang Estuary (North Pacific)	7	VRA	-	-	(Bettarel et al., 2011a)
South China Sea	13	VRA	-	-	(Chen et al., 2011)
South China Sea	5	FLVT	-	-	(Nguyen-Kim et al., 2015)
Jiulong Estuary (North Pacific)	27	VRA	27	MC	(Chen et al., 2019)

Southwestern Pacific	9	VRA	-	-	(Rowe et al., 2012)
Southern Ocean (Indian sector)	14	VRA	-	-	(Evans et al., 2009)
Southern Ocean (Indian sector)	10	VRA	-	-	(Malits et al., 2014)
Southern Ocean (Pacific sector)	16	VRA	8	MC	(Vaque et al., 2017)
Southern Ocean (Atlantic sector)	5	VRA	-	-	(Weinbauer et al., 2009)
Southern Ocean (Atlantic sector)	33	VRA	12	MC	(Evans and Brussaard, 2012)





**Appendix Figure A1. Viral abundance projected using the random forest, showing average in (a) surface 200 m and**  
 250 **(b) deep (>200 m) oceans.**

#### **Author contributions.**

RZ and YWL conceived and designed structure of database and mathematical analyses of the data. LX, WW, LC, XC and YH collected the data and described the metadata. LX, NJ, RZ and YWL conducted quality control and analyses of the data. LX, RZ and YWL led the writing of the paper, with contribution from all the co-authors.

255 **Competing interests.**

The authors declare that they have no conflict of interest.

**Acknowledgements.**

We would like to thank all the scientists and crews who collected the historical data. This work was funded by National Natural Science Foundation of China (91951209, 41890802, 42076153) and the MEL internal research fund (MELRI2003).

260 **Reference**

Akaike, H.: A new look at the statistical model identification, IEEE transactions on automatic control, 19, 716–723, [https://doi.org/10.1007/978-1-4612-1694-0\\_16](https://doi.org/10.1007/978-1-4612-1694-0_16), 1974.

Auguet, J. C., Montanie, H., Delmas, D., Hartmann, H. J., and Huet, V.: Dynamic of virioplankton abundance and its environmental control in the Charente estuary (France), Microb Ecol, 50, 337–349, <http://doi.org/10.1007/s00248-005-0183-2>, 2005.

Balsom, A. L.: Macroinfaunal community composition and biomass, and bacterial and viral abundances from the gulf of Alaska to the Canadian Archipelago : a biodiversity study, Master of Science, University of Tennessee, 2003.

Bar-On, Y. M. and Milo, R.: The biomass composition of the oceans: a blueprint of our blue planet, Cell, 179, 1451–1454, <https://doi.org/10.1016/j.cell.2019.11.018>, 2019.

270 Bettarel, Y., Arfi, R., Bouvier, T., Bouvy, M., Briand, E., Colombet, J., Corbin, D., and Sime-Ngando, T.: Virioplankton distribution and activity in a tropical eutrophicated bay, Estuar Coast Shelf S, 80, 425–429, <https://doi.org/10.1016/j.ecss.2008.08.018>, 2008.

Bettarel, Y., Bouvier, T., Agis, M., Bouvier, C., Chu, T. V., Combe, M., Mari, X., Nghiem, M. N., Nguyen, T. T., Pham, T. T., Pringault, O., Rochelle-Newall, E., Torretton, J. P., and Tran, H. Q.: Viral distribution and life strategies in the Bach Dang  
275 Estuary, Vietnam, Microb Ecol, 62, 143–154, <https://doi.org/10.1007/s00248-011-9835-6>, 2011a.

Bettarel, Y., Bouvier, T., Bouvier, C., Carre, C., Desnues, A., Domaizon, I., Jacquet, S., Robin, A., and Sime-Ngando, T.: Ecological traits of planktonic viruses and prokaryotes along a full-salinity gradient, FEMS Microbiol Ecol, 76, 360–372, <https://doi.org/10.1111/j.1574-6941.2011.01054.x>, 2011b.

- Bettarel, Y., Dolan, J. R., Hornak, K., Lemee, R., Masin, M., Pedrotti, M. L., Rochelle-Newall, E., Simek, K., and Sime-  
 280 Ngando, T.: Strong, weak, and missing links in a microbial community of the N.W. Mediterranean Sea, *FEMS Microbiol Ecol*,  
 42, 451-462, <http://doi.org/10.1111/j.1574-6941.2002.tb01034.x>, 2002.
- Bongiorni, L., Magagnini, M., Armeni, M., Noble, R., and Danovaro, R.: Viral production, decay rates, and life strategies  
 along a trophic gradient in the North Adriatic Sea, *Appl Environ Microbiol*, 71, 6644-6650,  
<http://doi.org/10.1128/AEM.71.11.6644-6650.2005>, 2005.
- 285 Boras, J. A., Sala, M. M., Arrieta, J. M., Sà E. L., Felipe, J., Agustí S., Duarte, C. M., and Vaqu é D.: Effect of ice melting  
 on bacterial carbon fluxes channelled by viruses and protists in the Arctic Ocean, *Polar Biology*, 33, 1695-1707,  
<https://doi.org/10.1007/s00300-010-0798-8>, 2010a.
- Boras, J. A., Sala, M. M., Baltar, F., Ari'stegui, J., Duarte, C. M., and Vaque, D.: Effect of viruses and protists on bacteria in  
 eddies of the Canary Current region (subtropical northeast Atlantic), *Limnol Oceanogr*, 55, 885-898,  
 290 <https://doi.org/10.4319/lo.2009.55.2.0885>, 2010b.
- Boras, J. A., Sala, M. M., Vazquez-Dominguez, E., Weinbauer, M. G., and Vaque, D.: Annual changes of bacterial mortality  
 due to viruses and protists in an oligotrophic coastal environment (NW Mediterranean), *Environ Microbiol*, 11, 1181-1193,  
<https://doi.org/10.1111/j.1462-2920.2008.01849.x>, 2009.
- Bouvier, T. and Maurice, C. F.: A single-cell analysis of virioplankton adsorption, infection, and intracellular abundance in  
 295 different bacterioplankton physiologic categories, *Microb Ecol*, 62, 669-678, <https://doi.org/10.1007/s00248-011-9862-3>,  
 2011.
- Bouvy, M., Combe, M., Bettarel, Y., Dupuy, C., Rochelle-Newall, E., and Charpy, L.: Uncoupled viral and bacterial  
 distributions in coral reef waters of Tuamotu Archipelago (French Polynesia), *Mar Pollut Bull*, 65, 506-515,  
<https://doi.org/10.1016/j.marpolbul.2012.01.001>, 2012.
- 300 Breiman, L.: Random forests, *Machine Learning*, 45, 5-32, <https://doi.org/10.1023/A:1010933404324>, 2001.
- Breitbart, M.: Marine viruses: truth or dare, *Ann Rev Mar Sci*, 4, 425-448, <https://doi.org/10.1146/annurev-marine-120709-142805>, 2012.
- Brum, J. R.: Concentration, production and turnover of viruses and dissolved DNA pools at Stn ALOHA, North Pacific  
 Subtropical Gyre, *Aquatic Microbial Ecology*, 41, 103-113, <http://doi.org/10.3354/ame041103>, 2005.

- 305 Brum, J. R., Hurwitz, B. L., Schofield, O., Ducklow, H. W., and Sullivan, M. B.: Seasonal time bombs: dominant temperate viruses affect Southern Ocean microbial dynamics, *ISME J*, 10, 437-449, <https://doi.org/10.1038/ismej.2015.125>, 2016.
- Brum, J. R., Ignacio-Espinoza, J. C., Roux, S., Doulier, G., Acinas, S. G., Alberti, A., Chaffron, S., Cruaud, C., de Vargas, C., Gasol, J. M., Gorsky, G., Gregory, A. C., Guidi, L., Hingamp, P., Iudicone, D., Not, F., Ogata, H., Pesant, S., Poulos, B. T., Schwenck, S. M., Speich, S., Dimier, C., Kandels-Lewis, S., Picheral, M., Searson, S., Bork, P., Bowler, C., Sunagawa, S.,
- 310 Wincker, P., Karsenti, E., and Sullivan, M. B.: Patterns and ecological drivers of ocean viral communities, *Science*, 348, 1261498, <https://doi.org/10.1126/science.1261498>, 2015.
- Brussaard, C. P.: Optimization of procedures for counting viruses by flow cytometry, *Appl Environ Microbiol*, 70, 1506-1513, <http://doi.org/10.1128/AEM.70.3.1506-1513.2004>, 2004.
- Brussaard, C. P. D., Payet, J. P., Winter, C., and Weinbauer, M. G.: Quantification of aquatic viruses by flow cytometry. In:
- 315 Manual of aquatic viral ecology, <https://doi.org/10.4319/mave.2010.978-0-9845591-0-7.102>, 2010.
- Cabral, A. S., Lessa, M. M., Junger, P. C., Thompson, F. L., and Paranhos, R.: Virioplankton dynamics are related to eutrophication levels in a tropical urbanized bay, *Plos One*, 12, e0174653, <https://doi.org/10.1371/journal.pone.0174653>, 2017.
- Chen, X., Liu, H., Weinbauer, M., Chen, B., and Jiao, N.: Viral dynamics in the surface water of the western South China Sea in summer 2007, *Aquat Microb Ecol*, 63, 145-160, <https://doi.org/10.3354/ame01490>, 2011.
- 320 Chen, X., Wei, W., Wang, J., Li, H., Sun, J., Ma, R., Jiao, N., and Zhang, R.: Tide driven microbial dynamics through virus-host interactions in the estuarine ecosystem, *Water Res*, 160, 118–129, <https://doi.org/10.1016/j.watres.2019.05.051>, 2019.
- Choi, D. H., Hwang, C. Y., and Cho, B. C.: Comparison of virus- and bacterivory-induced bacterial mortality in the eutrophic Masan Bay, Korea, *Aquatic Microbial Ecology*, 30, 117-125, <http://doi.org/10.3354/ame030117>, 2003.
- Clasen, J. L., Brigden, S. M., Payet, J. P., and Suttle, C. A.: Evidence that viral abundance across oceans and lakes is driven
- 325 by different biological factors, *Freshwater Biol*, 53, 1090-1100, <https://doi.org/10.1111/j.1365-2427.2008.01992.x>, 2008.
- Cobian Guemes, A. G., Youle, M., Cantu, V. A., Felts, B., Nulton, J., and Rohwer, F.: Viruses as winners in the game of life, *Annu Rev Virol*, 3, 197–214, <https://doi.org/10.1146/annurev-virology-100114-054952>, 2016.
- Cochran, P. K. and Paul, J. H.: Seasonal abundance of lysogenic bacteria in a subtropical estuary, *Appl Environ Microbiol*, 64, 2308-2312, <http://doi.org/10.1128/AEM.64.6.2308-2312.1998>, 1998.

- 330 De Corte, D., Sintes, E., Winter, C., Yokokawa, T., Reinthaler, T., and Herndl, G. J.: Links between viral and prokaryotic communities throughout the water column in the (sub)tropical Atlantic Ocean, *ISME J*, 4, 1431-1442, <https://doi.org/10.1038/ismej.2010.65>, 2010.
- De Corte, D., Sintes, E., Yokokawa, T., Lekunberri, I., and Herndl, G. J.: Large-scale distribution of microbial and viral populations in the South Atlantic Ocean, *Environ Microbiol Rep*, 8, 305-315, <https://doi.org/10.1111/1758-2229.12381>, 2016.
- 335 De Corte, D., Sintes, E., Yokokawa, T., Reinthaler, T., and Herndl, G. J.: Links between viruses and prokaryotes throughout the water column along a North Atlantic latitudinal transect, *ISME J*, 6, 1566-1577, <https://doi.org/10.1038/ismej.2011.214>, 2012.
- Evans, C. and Brussaard, C. P.: Regional variation in lytic and lysogenic viral infection in the Southern Ocean and its contribution to biogeochemical cycling, *Appl Environ Microbiol*, 78, 6741-6748, <https://doi.org/10.1128/AEM.01388-12>,  
340 2012.
- Evans, C., Pearce, I., and Brussaard, C. P.: Viral-mediated lysis of microbes and carbon release in the sub-Antarctic and Polar Frontal zones of the Australian Southern Ocean, *Environ Microbiol*, 11, 2924-2934, <https://doi.org/10.1111/j.1462-2920.2009.02050.x>, 2009.
- Finke, J. F., Hunt, B. P. V., Winter, C., Carmack, E. C., and Suttle, C. A.: Nutrients and other environmental factors influence  
345 virus abundances across oxic and hypoxic marine environments, *Viruses*, 9, <https://doi.org/10.3390/v9060152>, 2017.
- Fonda Umani, S., Malisana, E., Focaracci, F., Magagnini, M., Corinaldesi, C., and Danovaro, R.: Disentangling the effect of viruses and nanoflagellates on prokaryotes in bathypelagic waters of the Mediterranean Sea, *Mar Ecol Prog Ser*, 418, 73-85, <https://doi.org/10.3354/meps08803>, 2010.
- Fuhrman, J. A.: Marine viruses and their biogeochemical and ecological effects, *Nature*, 399, 541-548,  
350 <https://doi.org/10.1038/21119>, 1999.
- Fuhrman, J. A., Hewson, I., Schwalbach, M. S., Steele, J. A., Brown, M. V., and Naeem, S.: Annually reoccurring bacterial communities are predictable from ocean conditions, *Proc Natl Acad Sci U S A*, 103, 13104-13109, <http://doi.org/10.1073/pnas.0602399103>, 2006.
- Gainer, P. J., Pound, H. L., Larkin, A. A., LeClerc, G. R., DeBruyn, J. M., Zinser, E. R., Johnson, Z. I., and Wilhelm, S. W.:  
355 Contrasting seasonal drivers of virus abundance and production in the North Pacific Ocean, *PLoS One*, 12, e0184371, <https://doi.org/10.1371/journal.pone.0184371>, 2017.

- Glover, D. M., Jenkins, W. J., and Doney, S. C.: Modeling Methods for Marine Science, Cambridge University Press, New York, <http://doi.org/10.1017/cbo9780511975721>, 2011.
- Gregoracci, G. B., Soares, A. C., Miranda, M. D., Coutinho, R., and Thompson, F. L.: Insights into the microbial and viral  
 360 dynamics of a coastal downwelling-upwelling transition, PLoS One, 10, e0137090, <https://doi.org/10.1371/journal.pone.0137090>, 2015.
- Heldal, M. and Bratbak, G.: Production and Decay of Viruses in Aquatic Environments, Marine Ecology Progress Series, 72, 205-212, <http://doi.org/10.3354/meps072205>, 1991.
- Helton, R. R., Cottrell, M. T., Kirchman, D. L., and Wommack, K. E.: Evaluation of incubation-based methods for estimating  
 365 virioplankton production in estuaries, Aquatic Microbial Ecology, 41, 209-219, <http://doi.org/10.3354/ame041209>, 2005.
- Hewson, I. and Fuhrman, J. A.: Covariation of viral parameters with bacterial assemblage richness and diversity in the water column and sediments, Deep Sea Research Part I: Oceanographic Research Papers, 54, 811-830, <https://doi.org/10.1016/j.dsr.2007.02.003>, 2007.
- Holmfeldt, K., Titelman, J., and Riemann, L.: Virus production and lysate recycling in different sub-basins of the northern  
 370 Baltic Sea, Microb Ecol, 60, 572-580, <https://doi.org/10.1007/s00248-010-9668-8>, 2010.
- Hwang, C. Y. and Cho, B. C.: Virus-infected bacteria in oligotrophic open waters of the East Sea, Korea, Aquatic Microbial Ecology, 30, 1-9, <http://doi.org/10.3354/ame030001>, 2002.
- Jasna, V., Parvathi, A., Pradeep Ram, A. S., Balachandran, K. K., Madhu, N. V., Nair, M., Jyothibabu, R., Jayalakshmy, K. V., Revichandran, C., and Sime-Ngando, T.: Viral-induced mortality of prokaryotes in a tropical monsoonal estuary, Front  
 375 Microbiol, 8, 895, <https://doi.org/10.3389/fmicb.2017.00895>, 2017.
- Jasna, V., Pradeep Ram, A. S., Parvathi, A., and Sime-Ngando, T.: Differential impact of lytic viruses on prokaryotic morphopopulations in a tropical estuarine system (Cochin estuary, India), Plos One, 13, e0194020, <https://doi.org/10.1371/journal.pone.0194020>, 2018.
- Jiang, S. C. and Paul, J. H.: Occurrence of lysogenic bacteria in marine microbial communities as determined by prophage  
 380 induction, Marine Ecology Progress Series, 142, 27-38, <http://doi.org/10.3354/meps142027>, 1996.
- Jover, L. F., Effler, T. C., Buchan, A., Wilhelm, S. W., and Weitz, J. S.: The elemental composition of virus particles: implications for marine biogeochemical cycles, Nat Rev Microbiol, 12, 519-528, <https://doi.org/10.1038/nrmicro3289>, 2014.

- Karuza, A., Del Negro, P., Crevatin, E., and Fonda Umani, S.: Viral production in the Gulf of Trieste (Northern Adriatic Sea): Preliminary results using different methodological approaches, J Exp Mar Biol Ecol, 383, 96–104,  
385 <https://doi.org/10.1016/j.jembe.2009.12.003>, 2010.
- Kopylov, A. I., Sazhin, A. F., Zabolotkina, E. A., Romanenko, A. V., Romanova, N. D., and Boltenkova, M. A.: Virioplankton  
of the Kara Sea and the Yenisei River estuary in early spring, Estuar Coast Shelf S, 217, 37-44,  
<https://doi.org/10.1016/j.ecss.2018.10.015>, 2019.
- Kostner, N., Scharnreiter, L., Jurgens, K., Labrenz, M., Herndl, G. J., and Winter, C.: High viral abundance as a consequence  
390 of low viral decay in the Baltic Sea redoxcline, PLoS One, 12, e0178467, <https://doi.org/10.1371/journal.pone.0178467>, 2017.
- Lara, E., Vaqu r D., S   E. L., Boras, J. A., Gomes, A., Borrull, E., D  z-Vives, C., Teira, E., Pernice, M. C., Garcia, F. C.,  
Forn, I., Castillo, Y. M., Peir   A., Salazar, G., Mor  n, X. A. G., Massana, R., Catal   T. S., Luna, G. M., Agust   S., Estrada,  
M., Gasol, J. M., and Duarte, C. M.: Unveiling the role and life strategies of viruses from the surface to the dark ocean, Sci  
Adv, 3, e1602565, <https://doi.org/10.1126/sciadv.1602565>, 2017.
- 395 Li, W. K. and Dickie, P. M.: Monitoring phytoplankton, bacterioplankton, and virioplankton in a coastal inlet (Bedford Basin)  
by flow cytometry, Cytometry, 44, 236-246, [http://doi.org/10.1002/1097-0320\(20010701\)44:3](http://doi.org/10.1002/1097-0320(20010701)44:3<236::aid-cyto1116>3.0.co;2-5)<236::aid-cyto1116>3.0.co;2-  
5, 2001.
- Li, Y., Luo, T., Sun, J., Cai, L., Liang, Y., Jiao, N., and Zhang, R.: Lytic viral infection of bacterioplankton in deep waters of  
the western Pacific Ocean, Biogeosciences, 11, 2531-2542, <https://doi.org/10.5194/bg-11-2531-2014>, 2014.
- 400 Li, Z. S., Wang, M., Luo, Z. X., Liu, L., Xia, J., Gong, Z., Jiang, Y., Shao, H. B., Chen, H. T., and Tian, J. W.: Vertical  
distribution and dynamic variation of virioplankton in the mariana trench. , Oceanologia et Limnologia Sinica, 49, 1251-1258,  
<https://doi.org/10.11693/hyhz20180400105>, 2018.
- Liang, Y., Li, L., Luo, T., Zhang, Y., Zhang, R., and Jiao, N.: Horizontal and vertical distribution of marine virioplankton: a  
basin scale investigation based on a global cruise, PLoS One, 9, e111634, <https://doi.org/10.1371/journal.pone.0111634>, 2014.
- 405 Liang, Y., Zhang, Y., Zhang, Y., Luo, T., Rivkin, R. B., and Jiao, N.: Distributions and relationships of virio- and picoplankton  
in the epi-, meso- and bathypelagic zones of the Western Pacific Ocean, FEMS Microbiol Ecol, 93,  
<https://doi.org/10.1093/femsec/fiw238>, 2017.

- Long, A., McDaniel, L. D., Mobberley, J., and Paul, J. H.: Comparison of lysogeny (prophage induction) in heterotrophic bacterial and *Synechococcus* populations in the Gulf of Mexico and Mississippi River plume, *ISME J*, 2, 132-144, <https://doi.org/10.1038/ismej.2007.102>, 2008.
- Ma, Y., Wang, M., Xia, J., Song, X., Liu, Q., Li, Y., Jiang, Y., Shao, H. B., and Zhang, Y. Y.: Studies on abundance and diversity of microplankton during brown tide around Qin-huangdao area, *Periodical of Ocean University of China*, 46, 142-150, <https://doi.org/10.16441/j.cnki.hdxh.20150426>, 2016.
- Magagnini, M., Corinaldesi, C., Monticelli, L. S., De Domenico, E., and Danovaro, R.: Viral abundance and distribution in mesopelagic and bathypelagic waters of the Mediterranean Sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 54, 1209-1220, <https://doi.org/10.1016/j.dsr.2007.05.006>, 2007.
- Magiopoulos, I. and Pitta, P.: Viruses in a deep oligotrophic sea: Seasonal distribution of marine viruses in the epi-, meso- and bathypelagic waters of the Eastern Mediterranean Sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 66, 1-10, <https://doi.org/10.1016/j.dsr.2012.03.009>, 2012.
- Malits, A., Christaki, U., Obernosterer, I., and Weinbauer, M. G.: Enhanced viral production and virus-mediated mortality of bacterioplankton in a natural iron-fertilized bloom event above the Kerguelen Plateau, *Biogeosciences*, 11, 6841-6853, <https://doi.org/10.5194/bg-11-6841-2014>, 2014.
- Marie, D., Brussaard, C. P. D., Thyrhaug, R., Bratbak, G., and Vaulot, D.: Enumeration of marine viruses in culture and natural samples by flow cytometry, *Appl Environ Microbiol*, 65, 45-52, <http://doi.org/10.1128/AEM.65.1.45-52.1999>, 1999.
- Matteson, A. R., Loar, S. N., Pickmere, S., DeBruyn, J. M., Ellwood, M. J., Boyd, P. W., Hutchins, D. A., and Wilhelm, S. W.: Production of viruses during a spring phytoplankton bloom in the South Pacific Ocean near of New Zealand, *FEMS Microbiol Ecol*, 79, 709-719, <https://doi.org/10.1111/j.1574-6941.2011.01251.x>, 2012.
- Maurice, C. F., Bouvier, C., de Wit, R., and Bouvier, T.: Linking the lytic and lysogenic bacteriophage cycles to environmental conditions, host physiology and their variability in coastal lagoons, *Environ Microbiol*, 15, 2463-2475, <https://doi.org/10.1111/1462-2920.12120>, 2013.
- Maurice, C. F., Mouillot, D., Bettarel, Y., De Wit, R., Sarmiento, H., and Bouvier, T.: Disentangling the relative influence of bacterioplankton phylogeny and metabolism on lysogeny in reservoirs and lagoons, *ISME J*, 5, 831-842, <https://doi.org/10.1038/ismej.2010.181>, 2011.



- Mojica, K. D. A., van de Poll, W. H., Kehoe, M., Huisman, J., Timmermans, K. R., Buma, A. G. J., van der Woerd, H. J.,  
 435 Hahn-Woernle, L., Dijkstra, H. A., and Brussaard, C. P. D.: Phytoplankton community structure in relation to vertical stratification along a north-south gradient in the northeast Atlantic Ocean, *Limnol Oceanogr*, 60, 1498-1521, <https://doi.org/10.1002/lno.10113>, 2015.
- Motegi C and Nagata T: Enhancement of viral production by addition of nitrogen or nitrogen plus carbon in subtropical surface waters of the South Pacific, *Aquat Microb Ecol*, 48, 27-34, <https://doi.org/10.3354/ame048027>, 2007.
- 440 Motegi, C., Kaiser, K., Benner, R., and Weinbauer, M. G.: Effect of P-limitation on prokaryotic and viral production in surface waters of the northwestern Mediterranean Sea, *J Plankton Res*, 37, 16-20, <https://doi.org/10.1093/plankt/fbu089>, 2014.
- Motegi, C., Nagata, T., Miki, T., Weinbauer, M. G., Legendre, L., and Rassoulzadegan, F.: Viral control of bacterial growth efficiency in marine pelagic environments, *Limnol Oceanogr*, 54, 1901-1910, <https://doi.org/10.4319/lo.2009.54.6.1901>, 2009.
- Muck, S., Griessler, T., Kostner, N., Klimiuk, A., Winter, C., and Herndl, G. J.: Fracture zones in the Mid Atlantic Ridge lead  
 445 to alterations in prokaryotic and viral parameters in deep-water masses, *Front Microbiol*, 5, 264, <https://doi.org/10.3389/fmicb.2014.00264>, 2014.
- Nguyen-Kim, H., Bettarel, Y., Bouvier, T., Bouvier, C., Doan-Nhu, H., Nguyen-Ngoc, L., Nguyen-Thanh, T., Tran-Quang, H., Brune, J., and Wommack, K. E.: Coral mucus is a hot spot for viral infections, *Appl Environ Microbiol*, 81, 5773-5783, <https://doi.org/10.1128/aem.00542-15>, 2015.
- 450 Ni, Z., Huang, X., and Zhang, X.: Picoplankton and virioplankton abundance and community structure in Pearl River estuary and Daya Bay, South China, *J Environ Sci (China)*, 32, 146-154, <https://doi.org/10.1016/j.jes.2014.12.019>, 2015.
- Noble, R. T. and Fuhrman, J. A.: Rapid virus production and removal as measured with fluorescently labeled viruses as tracers, *Appl Environ Microbiol*, 66, 3790-3797, <http://doi.org/10.1128/aem.66.9.3790-3797.2000>, 2000.
- Noble, R. T. and Fuhrman, J. A.: Use of SYBR Green I for rapid epifluorescence counts of marine viruses and bacteria, *Aquatic*  
 455 *Microbial Ecology*, 14, 113-118, <http://doi.org/10.3354/ame014113>, 1998.
- Noble, R. T. and Steward, G. F.: Estimating viral proliferation in aquatic samples, *Methods in Microbiology*, 30, 67-82, [http://doi.org/10.1016/S0580-9517\(01\)30040-5](http://doi.org/10.1016/S0580-9517(01)30040-5), 2001.
- Ordulj, M., Krstulović, N., Šantić, D., Jozić, S., and Šolić, M.: Viral dynamics in two trophically different areas in the central Adriatic Sea, *Helgoland Mar Res*, 71, <https://doi.org/10.1186/s10152-017-0502-2>, 2017.

- 460 Parada, V., Baudoux, A. C., Sintes, E., Weinbauer, M. G., and Herndl, G. J.: Dynamics and diversity of newly produced  
virio plankton in the North Sea, *ISME J*, 2, 924-936, <https://doi.org/10.1038/ismej.2008.57>, 2008.
- Parada, V., Sintes, E., van Aken, H. M., Weinbauer, M. G., and Herndl, G. J.: Viral abundance, decay, and diversity in the  
meso- and bathypelagic waters of the North Atlantic, *Appl Environ Microbiol*, 73, 4429-4438,  
<https://doi.org/10.1128/AEM.00029-07>, 2007.
- 465 Parsons, R. J., Breitbart, M., Lomas, M. W., and Carlson, C. A.: Ocean time-series reveals recurring seasonal patterns of  
virio plankton dynamics in the northwestern Sargasso Sea, *ISME J*, 6, 273-284, <https://doi.org/10.1038/ismej.2011.101>, 2012.
- Parsons, R. J., Nelson, C. E., Carlson, C. A., Denman, C. C., Andersson, A. J., Kledzik, A. L., Vergin, K. L., McNally, S. P.,  
Treich, A. H., and Giovannoni, S. J.: Marine bacterio plankton community turnover within seasonally hypoxic waters of a  
subtropical sound: Devil's Hole, Bermuda, *Environ Microbiol*, 17, 3481-3499, <https://doi.org/10.1111/1462-2920.12445>, 2015.
- 470 Parvathi, A., Jasna, V., Aparna, S., Pradeep Ram, A. S., Aswathy, V. K., Balachandran, K. K., Muraleedharan, K. R., Mathew,  
D., and Sime- Ngando, T.: High incidence of lysogeny in the oxygen minimum zones of the Arabian Sea (southwest coast of  
India), *Viruses*, 10, <https://doi.org/10.3390/v10110588>, 2018.
- Parvathi, A., Jasna, V., Haridevi, K. C., Jina, S., Greeshma, M., Breezy, J., and Nair, M.: Diurnal variations in bacterial and  
viral production in Cochin estuary, India, *Environ Monit Assess*, 185, 8077-8088, [https://doi.org/10.1007/s10661-013-3156-](https://doi.org/10.1007/s10661-013-3156-9)  
475 [9](https://doi.org/10.1007/s10661-013-3156-9), 2013.
- Pasulka, A. L., Samo, T. J., and Landry, M. R.: Grazer and viral impacts on microbial growth and mortality in the southern  
California Current Ecosystem, *J Plankton Res*, 37, 320-336, <https://doi.org/10.1093/plankt/fbv011>, 2015.
- Paul, J. H.: Prophages in marine bacteria: dangerous molecular time bombs or the key to survival in the seas?, *ISME J*, 2,  
579-589, <https://doi.org/10.1038/ismej.2008.35>, 2008.
- 480 Payet, J. P. and Suttle, C. A.: Physical and biological correlates of virus dynamics in the southern Beaufort Sea and Amundsen  
Gulf, *J Marine Syst*, 74, 933-945, <https://doi.org/10.1016/j.jmarsys.2007.11.002>, 2008.
- Payet, J. P. and Suttle, C. A.: To kill or not to kill: The balance between lytic and lysogenic viral infection is driven by trophic  
status, *Limnol Oceanogr*, 58, 465-474, <https://doi.org/10.4319/lo.2013.58.2.0465>, 2013.
- Proctor, L. M. and Fuhrman, J. A.: Viral Mortality of Marine-Bacteria and Cyanobacteria, *Nature*, 343, 60-62,  
485 <http://doi.org/10.1038/343060a0>, 1990.

- Rastelli, E., Dell'Anno, A., Corinaldesi, C., Middelboe, M., Noble, R. T., and Danovaro, R.: Quantification of viral and prokaryotic production rates in benthic ecosystems: A methods comparison, *Front Microbiol*, 7, 1501, <https://doi.org/10.3389/fmicb.2016.01501>, 2016.
- 490 Rowe, J. M., DeBruyn, J. M., Poorvin, L., LeClerc, G. R., Johnson, Z. I., Zinser, E. R., and Wilhelm, S. W.: Viral and bacterial abundance and production in the western Pacific Ocean and the relation to other oceanic realms, *FEMS Microbiol Ecol*, 79, 359-370, <https://doi.org/10.1111/j.1574-6941.2011.01223.x>, 2012.
- Rowe, J. M., Saxton, M. A., Cottrell, M. T., DeBruyn, J. M., Berg, G. M., Kirchman, D. L., Hutchins, D. A., and Wilhelm, S. W.: Constraints on viral production in the Sargasso Sea and North Atlantic, *Aquat Microb Ecol*, 52, 233-244, <https://doi.org/10.3354/ame01231>, 2008.
- 495 Sabbagh, E. I., Huete-Stauffer, T. M., Calleja, M. L. L., Silva, L., Viegas, M., and Moran, X. A. G.: Weekly variations of viruses and heterotrophic nanoflagellates and their potential impact on bacterioplankton in shallow waters of the central Red Sea, *FEMS Microbiol Ecol*, 96, <https://doi.org/10.1093/femsec/fiaa033>, 2020.
- Steward, G. F., Smith, D. C., and Azam, F.: Abundance and production of bacteria and viruses in the Bering and Chukchi Seas, *Marine Ecology Progress Series*, 131, 287-300, <http://doi.org/10.3354/meps131287>, 1996.
- 500 Strzepek, R. F., Maldonado, M. T., Higgins, J. L., Hall, J., Safi, K., Wilhelm, S. W., and Boyd, P. W.: Spinning the "Ferrous Wheel": The importance of the microbial community in an iron budget during the FeCycle experiment, *Global Biogeochemical Cycles*, 19, <http://dx.doi.org/10.1029/2005GB002490>, 2005.
- Šulčius, S., Reunamo, A., Paškauskas, R., and Leskinen, P.: Influence of environmental variation on the bacterioplankton community and its loss to viral lysis in the Curonian Lagoon, *Estuar Coast Shelf S*, 204, 76-85, <https://doi.org/10.1016/j.ecss.2018.02.029>, 2018.
- 505 Suttle, C. A.: Marine viruses – major players in the global ecosystem, *Nat Rev Microbiol*, 5, 801–812, <https://doi.org/10.1038/nrmicro1750>, 2007.
- Suttle, C. A.: Viruses in the sea, *Nature*, 437, 356-361, <http://doi.org/10.1038/nature04160>, 2005.
- Taylor, G. T., Hein, C., and Iabichella, M.: Temporal variations in viral distributions in the anoxic Cariaco Basin, *Aquatic Microbial Ecology*, 30, 103-116, DOI 10.3354/ame030103, 2003.
- 510


- Thompson, F., Winter, C., Garcia, J. A. L., Weinbauer, M. G., DuBow, M. S., and Herndl, G. J.: Comparison of deep-water viromes from the Atlantic Ocean and the Mediterranean Sea, *PLoS ONE*, 9, e100600, <https://doi.org/10.1371/journal.pone.0100600>, 2014.
- 515 Vaque, D., Boras, J. A., Torrent-Llagostera, F., Agusti, S., Arrieta, J. M., Lara, E., Castillo, Y. M., Duarte, C. M., and Sala, M. M.: Viruses and protists induced-mortality of prokaryotes around the Antarctic Peninsula during the austral summer, *Front Microbiol*, 8, 241, <https://doi.org/10.3389/fmicb.2017.00241>, 2017.
- Wang, K., Wommack, K. E., and Chen, F.: Abundance and distribution of *Synechococcus* spp. and cyanophages in the Chesapeake Bay, *Appl Environ Microbiol*, 77, 7459-7468, <https://doi.org/10.1128/AEM.00267-11>, 2011.
- Weinbauer, M., Rowe, J., and Wilhelm, S.: Determining rates of virus production in aquatic systems by the virus reduction approach. In: *Manual of Aquatic Viral Ecology*, <https://doi.org/10.4319/mave.2010.978-0-9845591-0-7.1>, 2010.
- 520 Weinbauer, M. G.: Ecology of prokaryotic viruses, *FEMS Microbiol Rev*, 28, 127-181, <http://doi.org/10.1016/j.femsre.2003.08.001>, 2004.
- Weinbauer, M. G., Brettar, I., and Hofle, M. G.: Lysogeny and virus-induced mortality of bacterioplankton in surface, deep, and anoxic marine waters, *Limnology and Oceanography*, 48, 1457-1465, <http://doi.org/10.4319/lo.2003.48.4.1457>, 2003.
- 525 Weinbauer, M. G., Fuks, D., and Peduzzi, P.: Distribution of Viruses and Dissolved DNA along a Coastal Trophic Gradient in the Northern Adriatic Sea, *Appl Environ Microbiol*, 59, 4074-4082, <http://doi.org/10.1128/AEM.59.12.4074-4082.1993>, 1993.
- Weinbauer, M. G., Jesus-Maria Arrieta, Christian Griebler, and Herndl, G. J.: Enhanced viral production and infection of bacterioplankton during an iron-induced phytoplankton bloom in the Southern Ocean, *Limnol Oceanogr*, 54, 774-784, <https://doi.org/10.4319/lo.2009.54.3.0774>, 2009.
- 530 Weinbauer, M. G. and Suttle, C. A.: Lysogeny and prophage induction in coastal and offshore bacterial communities, *Aquatic Microbial Ecology*, 18, 217-225, <http://doi.org/10.3354/ame018217>, 1999.
- Weinbauer, M. G. and Suttle, C. A.: Potential significance of lysogeny to bacteriophage production and bacterial mortality in coastal waters of the gulf of Mexico, *Appl Environ Microbiol*, 62, 4374-4380, <http://doi.org/10.1128/AEM.62.12.4374-4380.1996>, 1996.
- 535

- Weinbauer, M. G., Winter, C., and Hofle, M. G.: Reconsidering transmission electron microscopy based estimates of viral infection of bacterio-plankton using conversion factors derived from natural communities, *Aquatic Microbial Ecology*, 27, 103-110, <http://doi.org/10.3354/ame027103>, 2002.
- 540 Wells, L. E. and Deming, J. W.: Significance of bacterivory and viral lysis in bottom waters of Franklin Bay, Canadian Arctic, during winter, *Aquatic Microbial Ecology*, 43, 209-221, <http://doi.org/10.3354/ame043209>, 2006.
- Wigington, C. H., Sonderegger, D., Brussaard, C. P. D., Buchan, A., Finke, J. F., Fuhrman, J. A., Lennon, J. T., Middelboe, M., Suttle, C. A., Stock, C., Wilson, W. H., Wommack, K. E., Wilhelm, S. W., and Weitz, J. S.: Re-examination of the relationship between marine virus and microbial cell abundances, *Nat Microbiol*, 1, 15024, <https://doi.org/10.1038/nmicrobiol.2015.24>, 2016.
- 545 Wilhelm, S. W., Jeffrey, W. H., Dean, A. L., Meador, J., Pakulski, J. D., and Mitchell, D. L.: UV radiation induced DNA damage in marine viruses along a latitudinal gradient in the southeastern Pacific Ocean, *Aquatic Microbial Ecology*, 31, 1-8, <http://doi.org/10.3354/ame031001>, 2003.
- Wilhelm, S. W. and Suttle, C. A.: Viruses and Nutrient Cycles in the Sea - Viruses play critical roles in the structure and function of aquatic food webs, *Bioscience*, 49, 781-788, <http://doi.org/10.2307/1313569>, 1999.
- 550 Williamson, S. J., Houchin, L. A., McDaniel, L., and Paul, J. H.: Seasonal variation in lysogeny as depicted by prophage induction in Tampa Bay, Florida, *Appl Environ Microbiol*, 68, 4307-4314, <http://doi.org/10.1128/aem.68.9.4307-4314.2002>, 2002.
- Winget, D. M., Williamson, K. E., Helton, R. R., and Wommack, K. E.: Tangential flow diafiltration: an improved technique for estimation of virioplankton production, *Aquatic Microbial Ecology*, 41, 221-232, <http://doi.org/10.3354/ame041221>, 2005.
- 555 Winget, D. M. and Wommack, K. E.: Diel and daily fluctuations in virioplankton production in coastal ecosystems, *Environ Microbiol*, 11, 2904-2914, <https://doi.org/10.1111/j.1462-2920.2009.02038.x>, 2009.
- Winter, C., Herndl, G. J., and Weinbauer, M. G.: Diel cycles in viral infection of bacterioplankton in the North Sea, *Aquatic Microbial Ecology*, 35, 207-216, <http://doi.org/10.3354/ame035207>, 2004.
- 560 Winter, C., Křstner, N., Kruspe, C. P., Urban, D., Muck, S., Reinthaler, T., and Herndl, G. J.: Mixing alters the lytic activity of viruses in the dark ocean, *Ecology*, 99, 700-713, <https://doi.org/10.1002/ecy.2135>, 2018.

- Winter, C., Kerros, M.-E., and Weinbauer, M. G.: Seasonal and depth-related dynamics of prokaryotes and viruses in surface and deep waters of the northwestern Mediterranean Sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 56, 1972-1982, <https://doi.org/10.1016/j.dsr.2009.07.003>, 2009.
- 565 Winter, C., Moeseneder, M. M., Herndl, G. J., and Weinbauer, M. G.: Relationship of geographic distance, depth, temperature, and viruses with prokaryotic communities in the eastern tropical Atlantic Ocean, *Microb Ecol*, 56, 383-389, <https://doi.org/10.1007/s00248-007-9343-x>, 2008.
- Winter, C., Smit, A., Szoek-Denes, T., Herndl, G. J., and Weinbauer, M. G.: Modelling viral impact on bacterioplankton in the North Sea using artificial neural networks, *Environ Microbiol*, 7, 881-893, <http://doi.org/10.1111/j.1462-2920.2005.00768.x>, 2005.
- 570 Wommack, K. E. and Colwell, R. R.: Virioplankton: viruses in aquatic ecosystems, *Microbiol Mol Biol Rev*, 64, 69-114, <http://doi.org/10.1128/mmbr.64.1.69-114.2000>, 2000.
- Xie, L., Wei, W., Cai, L., Chen, X., Huang, Y., Zhang, R., and Luo, Y.-W.: A global viral oceanography database (gVOD) from 1987 to 2018, *PANGAEA*, <https://doi.org/10.1594/PANGAEA.915758>, 2020.
- Yang, Y., Yokokawa, T., Motegi, C., and Nagata, T.: Large-scale distribution of viruses in deep waters of the Pacific and Southern Oceans, *Aquat Microb Ecol*, 71, 193-202, <https://doi.org/10.3354/ame01677>, 2014.
- 575 Zhang, R., Weinbauer, M. G., and Qian, P. Y.: Viruses and flagellates sustain apparent richness and reduce biomass accumulation of bacterioplankton in coastal marine waters, *Environ Microbiol*, 9, 3008–3018, <https://doi.org/10.1111/j.1462-2920.2007.01410.x>, 2007.
- Zhao, Y., Zhao, Y., Zheng, S., Zhao, L., Li, X., Zhang, W., Gregori, G., and Xiao, T.: Virioplankton distribution in the tropical western Pacific Ocean in the vicinity of a seamount, *Microbiologyopen*, e1031, <https://doi.org/10.1002/mbo3.1031>, 2020.
- 580 Zimina, I. D., Maksimov, A. I., and Svetsov, V. I.: Investigation of excitation of ammonia decomposition products in a high-frequency discharge, *Journal of Applied Spectroscopy*, 18, 771-772, <http://doi.org/10.1007/bf00614106>, 1973.


585 **Table 1. Number of accessory viral, host and oceanographic parameters associated with each of the core viral parameters (VA, lytic and lysogenic VP).**

	VA (n=10,931)	Lytic VP (n=608)	Lysogenic VP (n=119)
<i>Accessory viral parameters</i>			
Frequency of lytic infection	405	142	36
Frequency of lysogenic infection	227	96	36
Viral decay rate	83	65	27
Burst size	206	96	-
Virus-mediated bacterial mortality	46	46	19
% cells lysed	55	53	17
<i>Accessory host parameters</i>			
Prokaryotic abundance	10,846	598	119
Prokaryotic productivity	2,425	352	91
Flagellate abundance	411	44	7
Picoeukaryotic abundance	1,554	68	15
<i>Synechococcus</i> abundance	1,700	80	42
<i>Prochlorococcus</i> abundance	1,567	73	42
<i>Accessory oceanographic parameters</i>			
Chlorophyll <i>a</i>	3,949	244	71
Temperature	6,253	399	119
Salinity	6,360	370	85
Oxygen	4,930	82	46
Nitrate	2,707	91	46
Phosphate	3,153	144	51
Silicate	2,638	106	36
pH	96	47	-
Light intensity	35	35	-
Dissolved organic carbon	97	13	-



**Figure 1. Collected viral abundance (a, b), lytic viral production (d, f) and lysogenic viral production (g, h) in surface ( $\leq 200$  m) (a, d and g) and in deep ( $> 200$  m) waters (b, e and h), binned on 1°x1° grids. Color of each grid codes the mean value of the parameters, and the size of the circles represents number of samples in each bin. Also shown the number of samples in latitudinal bands (c, f and i). Numbers in (a) represent long-term time-series observations of viral abundance: 1, Rivers Inlet; 2, Saanich Inlet; 3, San Pedro Ocean Time Series Microbial Observatory; 4, Bedford Basin Monitor; 5, Bermuda Atlantic Time-series Study and 6: Guanabara Bay.**





**Figure 2. The fraction of (a) viral abundance, (b) lytic viral production and (c) lysogenic viral production data points in different oceans (N. Atl: North Atlantic, S. Atl: South Atlantic, N. Pac: North Pacific, S. Pac: South Pacific, Ind: Indian ocean, Arc: Arctic ocean, M & B: Mediterranean Sea and Baltic Sea).**




Figure 3. Depth distribution of the data for (a) viral abundance, (b) lytic viral production, and (c) lysogenic viral production.






Figure 4. Seasonal distributions of number of samples for (a) viral abundance, (b) lytic viral production and (c) lysogenic viral production.



**Figure 5.** The fraction of viral abundance (VA), lytic viral production (VP) and lysogenic VP data points in coastal versus in open oceans.

610



**Figure 6.** The range of (a, d) viral abundance, (b, e) lytic viral production, and (c, f) lysogenic viral production in the different oceans (N. Atl: North Atlantic, S. Atl: South Atlantic, N. Pac: North Pacific, S. Pac: South Pacific, Ind: Indian ocean, Arc: Arctic ocean, M & B: Mediterranean Sea and Baltic Sea), grouping in coastal ocean (a–c) and open ocean samples in surface 200 m (d–f). All data are shown in logarithmic scales. The red diamonds mark the mean value. The central red lines indicate the median, and the bottom and top edges of the box indicate the 25th and 75th percentiles of the data. Error bars extend to the 5th and 95th percentiles and the remaining outliers are marked with red plus signs.




Figure 7. Vertical profiles of average viral abundance of each ocean basin in (a) coastal and (b) open ocean waters. Also shown the vertical profiles for open water lytic viral production (c), while those for coastal samples were not constructed because of limited data points. Error bars represent one standard error of the mean.





Figure 8. Box plots of distribution of viral abundance using different measurement methods (EFM, FCM and TEM, see text for more details) in coastal (a), surface open ocean (b) and deep open ocean (c) waters. See caption of Figure 6 for details of lines and symbols of the box plots.



**Figure 9. Comparison of lytic viral production using different measurement methods including VRA, RIA, FPB, FLVT and VDR (FPB: calculated by multiplying fraction of viral infected cells, prokaryotic production and burst size; RIA: radioactive incorporation approach; FLVT: fluorescently labelled viral tracers method; VRA: virus reduction approach; VDR: estimated by viral decay rates. See text for details) in coastal (a), surface open ocean (b) and deep open ocean (c) waters. See caption of Figure 6 for details of lines and symbols of the box plots.**