

A restructured and updated global soil respiration database (SRDB-V5)

Jinshi Jian¹, Rodrigo Vargas², Kristina Anderson-Teixeira^{3,4}, Emma Stell², Valentine Herrmann³, Mercedes Horn⁵, Nazar Kholod¹, Jason Manzon⁶, Rebecca Marchesi³, Darlin Paredes⁷, and Ben Bond-Lamberty^{1*}

Correspondence to: Ben Bond-Lamberty (bondlamberty@pnnl.gov)

¹ Pacific Northwest National Laboratory, Joint Global Change Research Institute at the University of Maryland–College Park, 5825 University Research Court, Suite 3500, College Park, MD 20740, USA

² Department of Plant and Soil Sciences, University of Delaware, Newark, DE, USA

³ Conservation Ecology Center; Smithsonian Conservation Biology Institute; Front Royal, VA, USA

⁴ Center for Tropical Forest Science-Forest Global Earth Observatory; Smithsonian Tropical Research Institute; Panama, Republic of Panama

⁵ University of Vermont, Rubenstein School of Environment and Natural Resources

⁶ University of Maryland

⁷ Georgetown University, School of Foreign Service

Abstract

Field-measured soil respiration (R_s , the soil-to-atmosphere CO_2 flux) observations were compiled into a global soil respiration database (SRDB) a decade ago, a resource that has been widely used by the biogeochemistry community to advance our understanding of R_s dynamics. Novel carbon cycle sciences questions require updated and augmented global information with better interoperability among datasets. Here, we restructured and updated the global R_s database to version SRDB-V5. The updated version has all previous fields revised for consistency and simplicity, and it has several new fields to include ancillary information (e.g., R_s measurement time, collar insertion depth, collar area). The new SRDB-V5 includes published papers through 2017 (800 independent studies) where total observations increased from 6633 in SRDB-V4 to 10366 in SRDB-V5. The SRDB-V5 features more R_s data published in Russian and Chinese scientific literature, has an improved global spatio-temporal coverage, and improved global climate-space representation. We also restructured the database so that it has stronger interoperability with other datasets related to carbon-cycle science. For instance, linking SRDB-V5 with an hourly timescale global soil respiration database (HGRsD) and an open community database for continuous soil respiration and other chamber flux data (COSORE) enables researchers to explore new questions. The updated SRDB-V5 aims to be a data framework for the scientific community to share seasonal to annual field R_s measurements, and it provides opportunities for the biogeochemistry community to better understand the spatial and temporal variability of R_s , its components, and the overall carbon cycle.

The database can be downloaded at <https://github.com/bpbond/srdb> and ORNL DAAC [Submitted].

All data and code to reproduce the results in this study can be found at: Jian, Jinshi, Bond-Lamberty, Ben. (2020). jinshijian/ESSD: SRDB-V5 first release (Version v1.0.0) [Data set]. Zenodo.

<http://doi.org/10.5281/zenodo.3876443>.

Keywords: SRDB, soil respiration, database, interoperability, reusability

1. Introduction

45 Soil respiration (R_s), the soil surface to atmosphere CO_2 flux, is one of the largest carbon fluxes between the
terrestrial land surface and atmosphere (Luo and Zhou, 2010). The majority of R_s is released by soil microbial/fauna
(heterotrophic respiration) and plant root respiration (autotrophic respiration). Soils hold a large amount (>2000 Pg
C to 1 m depth) of carbon, more than the total of carbon stock in the atmosphere and aboveground plants (Batjes,
2016; Tarnocai et al., 2009). Thus, its C efflux to the atmosphere has major implications for our understanding of
ecosystem- to global-scale biogeochemical cycling. For better monitoring soil carbon dynamics as well as to
50 investigate how soil carbon responds to global climate change, it is important to measure R_s across different
vegetation types and climate conditions.

Many field experiments have been conducted in recent decades to measure R_s in different climate conditions and
vegetation types (Bond-Lamberty and Thomson, 2010b; Davidson et al., 1998; Raich and Potter, 1995). However,
55 the resulting estimates of seasonal to annual R_s fluxes are scattered throughout the scientific literature in a variety of
formats. Therefore, compiling past R_s measurements together into a standardized data framework to support
synthesis analysis is very important to advance carbon cycle science.

Published site scale R_s measurements across the globe have been compiled and standardized into global soil
60 respiration databases to support synthesis studies, macro-to-global scale R_s estimates, and soil carbon response to
climate change investigation (Bond-Lamberty and Thomson, 2010a; Raich and Schlesinger, 1992). Schlesinger
(1977) compiled one of the earliest listings of R_s estimates from diverse ecosystems. Raich and Schlesinger (1992)
subsequently integrated R_s from published papers which covered 13 ecosystems, and developed a simple linear
model between R_s and climate factors (i.e., temperature and precipitation), estimating global R_s to be 68 ± 4 Pg C
65 yr^{-1} . Later, more R_s measurements (especially measured using Infra-Red Gas Analyzers (IRGA method) were added
and the global R_s was updated to 76-81 Pg C yr^{-1} (Raich et al., 2002; Raich and Potter, 1995). In 2010, Bond-
Lamberty and Thomson (2010a) compiled a comprehensive global soil respiration database (SRDB) and this
database was released for public usage. The SRDB contains annual and seasonal R_s measurements, ancillary carbon
pools and fluxes (e.g., gross primary production, net primary production, ecosystem respiration), response of R_s to
70 temperature and moisture (i.e., model parameters to describe the relationship between R_s and temperature and
moisture), and sites' background information (e.g., latitude, longitude, elevation, mean annual temperature, mean
annual precipitation) (Bond-Lamberty and Thomson, 2018, 2010a). With more IRGA-based R_s measurements added
and alkaline-based measurements excluded, Bond-Lamberty and Thomson (2010b) estimated the global R_s to be 98
 ± 12 Pg C yr^{-1} and estimated that global R_s was increasing at a rate of 0.1 Pg C yr^{-2} . The SRDB has been widely
75 used in the past decade since the first version was published (Bond-Lamberty and Thomson, 2010a), and to date it
has been cited 359 times (searched in Google Scholar on 5/20/2020) but its use continues to increase each year
(Figure 1).

The SRDB of Bond-Lamberty and Thomson (2010) however only recorded seasonal to annual R_s fluxes, hindering
80 analyses at finer temporal resolutions. Based on the SRDB, Jian et al. (2018c) collected SRDB studies reporting
diurnal R_s and compiled these into a global hourly soil respiration database (HGRsD). Similarly, Jian et al. (2018a)
further collected detailed monthly/daily time scale R_s measurements into a global monthly/daily soil respiration
database (MGRsD). More recently, Bond-Lamberty et al. (2020) have built a database (COSORE) of continuous
(typically half-hourly or hourly) datasets from globally-distributed sites. With these different-timescale databases,
85 R_s temporal variability, and its time-related driving processes and uncertainties, can be analyzed (Jian et al., 2018a,
2018b, 2018c). There is still a need to improve interoperability among R_s databases to expand available information,
improve database usage, and to advance our understanding of R_s dynamics across multiple spatial and temporal
scales.

90 In approaching a decadal reworking of the SRDB, we envisioned that it required improvements to increase its usage
across different disciplines. Some important information (e.g., collar area, collar insertion depth, R_s measure time,
soil temperature, soil moisture, soil temperature measure depth, and soil moisture measure depth) was not included
95 in the older versions (hereafter named SRDB-V1 to SRDB-V4), and thus important questions such as whether R_s
survey time (Cueva et al., 2017), collar insertion depth (Heinemeyer et al., 2011), and/or how collar cover area
affected R_s measurements accuracy could not be addressed. In addition, SRDB-V4 included data mainly published
in English (~98%), while data published in other languages (~2%) were rarely included (Epule, 2015). Some
metadata such as manipulation/treatments and measurement method were not standardized and thus were difficult to
100 use in subsequent meta-analyses. For instance, the attempt to link SRDB to the Forest Carbon Database (ForC)
showed that the old SRDB structure required modification before it can be linked with ForC (Anderson-Teixeira et
al., 2018a, 2018b). Finally, information about how heterotrophic (R_H) and autotrophic respiration (R_A) respond to
environmental conditions (i.e., temperature and soil moisture) was not included.

The older SRDB followed certain data integration principles, including inclusion criteria, database structure design,
and quality control (Bond-Lamberty and Thomson, 2010a), but improvements could be made. We have updated it to
105 a new version (hereafter named SRDB-V5) following FAIR protocols (i.e., Findable, Accessible, Interoperable, and
Reusable) (Wilkinson et al., 2016). This has been accomplished by 1) restructuring SRDB and improving its
interoperability so that data from SRDB-V5 can more easily be linked to external datasets; 2) separating the R_s , R_H ,
and R_A responses to temperature and soil moisture functions into a separate file to simplify the database and improve
its reusability; 3) adding collar area, collar insertion depth, and R_s measurement time information to SRDB-V5; 4)
110 collecting more R_s data published in Russian and Chinese scientific literature; 5) updating R_s records available
throughout the world from recently published literature (until 2017); and 6) improving the metadata description. We
hope that these efforts will significantly improve the future interoperability and reusability of SRDB-V5.

2. Methods

115 2.1 Soil respiration database restructuring

We restructured the SRDB for easier data collection and quality control. The previous global R_s database versions
(SRDB-V1 to SRDB-V4) mainly included 2 files: a “studies” file, which recorded the detailed metadata for all
published papers examined by the SRDB; and a “data” file, which stores all the R_s data, a variety of ancillary site,
soil, and carbon cycle data (e.g., GPP, NPP, ecosystem respiration), and related background information such as site
120 location, ecosystem type, and management (Bond-Lamberty and Thomson, 2010a). In SRDB-V5 the “studies” file
remains unchanged, but the “data” file is now separated into two files: “srdb-data” and “srdb-equations”. This
simplifies the structure of the former, while moving all the “Response of R_s to temperature and moisture” columns
in the SRDB to the latter. Note that the SRDB-V5 file format remains the same as the older versions, as comma
separated value data are easy to work with and universally readable by software.

125 2.2 Metadata

We standardized the background information of SRDB-V5. Most of the metadata are described by Bond-Lamberty
and Thomson (2010a), and here we only describe new added columns or metadata with updates (Table 1 to Table 3).
We added five columns (i.e., *Site_ID*, *Collar_height*, *Collar_depth*, *Chamber_area*, *Time_of_day*) in SRDB-V5.
130 Four columns (*Rs_max*, *Rs_maxday*, *Rs_min*, *Rs_minday*) were deleted (Table 1) because they were rarely reported
and had not been used by the community in the past ten years (according to our literature search, *Rs_max*,
Rs_maxday, *Rs_min*, *Rs_minday* have never been used). In the *Quality_flag* column, we added two more flags
related to R_s -temperature equations: Q15 means the equation was developed based on seasonal R_s data rather than

135 covering at least a whole year, and Q16 notes that there is a soil water content (SWC) component within the reported equation (Table 1).

140 For many analyses SRDB needs to be connected with other datasets, and a unique observation ID is essential for this process. In the SRDB-V5, we added a “*Site_ID*” column to guarantee a unique ID for each *R_s annual* observation within a study, enabling users to easily link SRDB-V5 records with external data such as MGRsD and HGRsD. The *Site_ID* is in the form of ‘CC-RC-IC’, where CC is the ISO Alpha-2 country code (https://www.nationsonline.org/oneworld/country_code_list.htm), RC is region code (state/province), and IC is identity code. Country code and region code are always present, but some studies report only one annual *R_s* value, and thus IC may or may not be present.

145 We standardized the coding of experimental manipulation, collapsing the previous *ad hoc* categories into a smaller set of standardized terms. This decreased the number of unique *Manipulation* field values from 689 to 276. We used the following criteria to simplify the manipulation in SRDB-V5: 1) Measurements from no-treatment (i.e., control) were categorized as “None”, 2) manipulation names were standardized (e.g., “clipping”, “clip”, and “clipped” are now all standardized as “Clip”), 3) we used the manipulation level to further describe the difference within a specific manipulation (e.g., “Litter manipulation” could have “double litter”, “50% litter removal”, “100% litter removal”).
150 With manipulation standardized, scientists can further analyze how manipulation affects *R_s*. For instance, comparing *R_s* measurements from the “CO₂” group (i.e., elevated CO₂ concentration treatment) with “None” (i.e., control) enables researchers to analyze how *R_s* responds to CO₂ concentration increase caused by CO₂ released from fossil fuel combustion. Similarly, data from the “Warm” and “Precipitation amount change” groupings will enable
155 scientists to more easily explore how soil carbon responds to global climate change. Barba et al. (2018) suggested that bias could arise from measurements made in “hot-spots” (i.e., areas with high values compared with the surrounding environment), and groupings such as “Ant mound” and “High N” facilitate data interpretation and analyses regarding “hot-spots”.

160 We also standardized the *R_s* measurement method (the *Meas_method*) and *R_s* partition method (*Partition_method*) fields. Measurement method was grouped into 9 types (Table 2) and the partition method was grouped into 8 types (Table 3). With these changes, scientists can more easily investigate whether different measure methods affect *R_s* results, as well as whether different partition methods affect *R_H* and *R_A* partitioning.

165 Latitude and longitude are key metadata as they can be used to link *R_s* measurements to spatial data (e.g., precipitation and air temperature). During the data collecting process, latitude and longitude values reported in the original paper were recorded in our database, generally to two significant digits. However, the precision of SRDB latitude and longitude can be affected by many factors: first, studies report latitude and longitude at different and sometimes uncertain levels of precision; second, studies use different methods for recording latitude and longitude;
170 and finally, some studies have multiple nearby sites, but report one general latitude and longitude for all those sites. However, it is unlikely that the error is very large, and in general we assume that linking *R_s* measurements to relatively coarse spatial data (e.g., 0.1°-0.5° resolution) should be unproblematic. When linking to high spatial resolution data (such as 30 m resolution remote sensing images), users should be aware that the variable and uncertain SRDB latitude and longitude precision may cause data quality issues. That said, SRDB-V5 was revised to
175 avoid unrealistic locations such as points in the ocean. Furthermore, the Latitude and Longitude fields should be within -90 to 90 and -180 to 180 degrees, respectively; whenever they are out of these ranges, a warning is raised.

2.3 Soil respiration database update

We updated the SRDB-V5 so that it has temporal coverage to 2017, and made an effort to collect *R_s* data published in Russian and Chinese literature to be more inclusive and expand its spatial coverage. Papers published in English
180 are the majority (~98%) of sources in SRDB, while papers published in other languages are rarely included (Bond-

185 Lamberty and Thomson, 2018, 2010a). This reflects the dominance of English as the language of international science, but there are some data available from the Russian-language literature, representing data from a large area (Russia represents ~11% of the terrestrial land surface) and a variety of climate types and vegetation types. In addition, in MGRsD and HGRsD, there were some Chinese-language papers or recently-published papers (103 studies, ~5% of the total studies in SRDB-V5), which were not included by SRDB. Now we have compiled data from those papers into SRDB-V5.

2.4 Data quality control

190 We developed an R (R Core Team, 2019) script to perform data quality and consistency checks. For example, the *Latitude* and *Longitude* fields have to be in specific ranges, otherwise a warning is raised. For details about the data constraints used to check each column in SRDB-V5 please see the ‘*srdb_check.R*’ script, which is available in the GitHub repository and as part of every release download (<https://github.com/bpbond/srdb/releases>). This script is also run on all *pull requests* to the Github repository, which enables us to flag data-quality problems before changes are made to the database.

2.5 Data coverage analysis

195 We compared mean annual temperature (MAT) and mean annual precipitation (MAP) of sites from SRDB with the global MAT and MAP to test the representation of the SRDB. We connected the sites from SRDB with external climate data (Willmott and Matsuura, 2001) through latitude and longitude, and obtained MAT and MAP. Barren area was masked according to the MODIS landcover (Friedl et al., 2002). Climate region was retrieved from the climate Köppen classification (Peel et al., 2007). We also obtained IGBP vegetation classification of the SRDB sites by connecting IGBP classification data (IGBP, 1990); vegetation was grouped into Agriculture, Arctic, Desert, 200 tropical forest (Tropic FOR), temperate & boreal forest (T&B FOR), Grassland, Savanna, Shrubland, Urban, and Wetland. If the MAT and MAP distribution of SRDB sites is similar to that of global MAT and MAP distribution, it should mean that the SRDB better represents the global flux R_s distribution as well. We also assume that as data sample size increases, the new database (e.g., SRDB-V5) should improve its representation compared with the older version (e.g., SRDB-V1). We tested the representation of sites in different vegetation types (IGBP, 1990). 205

Table 1. Summary of metadata updates in SRDB-V5 compared with the old version SRDB-V4.

Column	Description	Comments
<i>Site_ID</i>	CC-RC-IC (country code - region code - identity code)	Added in SRDB-V5
<i>Collar_height</i>	Total height of collar	Added in SRDB-V5
<i>Collar_depth</i>	Depth of collar inserted into soil (always < Collar_height)	Added in SRDB-V5
<i>Chamber_area</i>	Area of collar covering the surface	Added in SRDB-V5
<i>Time_of_day</i>	R_s survey time (e.g., 8to12 represents R_s measured from 8:00 to 12:00, local time; 0to24 stands for continuous measurement)	Added in SRDB-V5
<i>Rs_max</i>	Maximum R_s rate in a year	Deleted in SRDB-V5
<i>Rs_maxday</i>	Day of year R_s_{max} recorded	Deleted in SRDB-V5

<i>Rs_min</i>	Minimum R_s rate in a year	Deleted in SRDB-V5
<i>Rs_minday</i>	Day of year R_s_{min} recorded	Deleted in SRDB-V5
<i>Quality_flag</i>	Q15: equation simulated based on seasonal rather than annual data; Q16: Equation with SWC component	Updated in SRDB-V5
<i>Manipulation</i>	Decreased from 689 unique values to 276 after being standardized	Standardized in SRDB-V5
<i>Measure_method</i>	See Table 2	Standardized in SRDB-V5
<i>Partition_method</i>	See Table 3	Standardized in SRDB-V5

210

Table 2. Summary of standardized measurement method (*Meas_method*) in SRDB-V5.

<i>Meas_method</i>	Number of records (n)	Comments
IRGA	7734	Type of Infrared gas analyzer (e.g., LICOR 8100A)
Gas chromatography	1268	Take gas samples in the field, and measure CO ₂ concentration back in the laboratory to determine soil respiration rate
Alkali absorption	910	Using alkali absorption of CO ₂ to determine soil respiration rate
Not reported	238	Measure method not reported in the study
EC	88	Eddy covariance
Gradient	83	Measure CO ₂ concentration at different soil depth and calculate soil respiration rate based on gas diffusion law
Equation	15	Indirectly calculate soil respiration rate (e.g., through relationship between soil respiration and GPP)
Isotope	3	Determine soil respiration rate using isotope (e.g., C ¹³)
Unknown	27	None of above

Table 3. Summary of standardized partition method (*Partition_method*) in SRDB-V5.

<i>Partition_method</i>	Number of records (n)	Comments
Comparison	150	Separating soil respiration into heterotrophic and autotrophic components by comparing with e.g., bare, clear-cut, gap, or clip site
Exclusion	1121	Removing roots by trenching, deeply insert PVC pipe etc.

Extraction	180	Directly measure respiration from root to get autotrophic respiration
Girdling	23	Strips the stem bark to the depth of xylem, and measure respiration few months later to get the heterotrophic respiration
Isotope	68	Separating heterotrophic and autotrophic respiration through isotope labeling
Model	49	Separating heterotrophic and autotrophic respiration through a relationship (e.g., the relationship reported by (Bond-Lamberty et al., 2004))
TBCA	16	Determining heterotrophic and autotrophic respiration through total belowground carbon allocation calculation
Other	122	None of above

215 3. Results

220 The number of records of SRDB-V5 is much larger compared with older versions. Collecting R_s measurements from newly published literature (until 2017) greatly improves the total number of observations in the database (increased from 6,633 to 10,366) in SRDB-V5, but only somewhat improved its spatial coverage (Figure 2). The northern hemisphere mid-latitude regions, where SRDB-V4 has the most R_s sites, had the largest R_s increase in SRDB-V5 as well (blue dots in Figure 2). Adding literature in Chinese did not substantially improve the spatial coverage either, possibly because more and more R_s measurements in China have been published in English scientific literature. However, most sites in China are from the eastern part of the country, and measurements from western China, if available, will be important to include in future SRDB updates. We collected ~50 papers published in Russian, but only 14 of them (~0.7% of total studies of all languages in SRDB-V5) met the criteria [see (Bond-Lamberty and Thomson, 2010a) for details] and were included in the database. This small number of papers nonetheless substantially improved the database's spatial coverage of the Russian landmass (orange circles in Figure 2).

230 MAT and MAP distribution of SRDB sites are very similar to global distribution in Agriculture, Forest, and Grassland regions, indicating good representativeness of SRDB sites in these three vegetations (Figures 3 and 4). For Shrublands, sites in the oldest versions of the database (e.g., SRDB-V4) did not represent the global distribution well, but this distribution was greatly improved as more R_s measurements were included in SRDB-V5 (Figure 3). Sites from other vegetation types, however, were less representative of the corresponding global climate space, with barren lands were masked out (Figure 3, right panel). More specifically, Arctic sites in SRDB have relatively narrow MAT and MAP coverage compared with the global Arctic MAT and MAP distribution, probably because many regions in the Arctic are covered by snow all year round, and thus it is difficult to measure R_s in those sites (Virkkala et al., 2019). Desert SRDB sites have lower MAT but higher MAP than the global distribution, probably because: 1) the disproportionate amount of samples in temperate regions (Figure 2) means that most sample in deserts are likely from wetter deserts; 2) the Sahara has low MAP and high MAT and covers a large area of the world, but few studies were conducted there so that area of the world may simply represents the bias; and 3) many "deserts" that have been studied are in relatively close proximity with urban developments (e.g., southwestern USA, southern Europe) and those deserts are not as harsh nor extensive as the Sahara. Urban and Savanna sites in SRDB had lower MAT compared to their global distribution, probably because many tropical cities and savannas in South America, Asia, and Africa were rarely measured (Jian et al., 2020; Martin et al., 2012). We suggest that manuscripts

245 written in other languages, especially those in Portuguese, Spanish, and French could potentially increase the R_s measurements in South America and Africa.

250 Adding new measurements in SRDB-V5 has substantially increased total observations and the spatial coverage of sites was improved compared with SRDB-V4 (Figure 2). However, the distribution of annual R_s and seasonal R_s (growing, dry, wet, spring, summer, autumn, and winter season R_s) were similar in the SRDB-V5 compared to SRDB-V4 (Figure 5). We suspect that new R_s measurements are collected disproportionately from the same regions as previously sampled, and thus future studies should focus more on those regions with less data. For the future SRDB update, measurements from the Southern hemisphere, Desert, Arctic, and tropical forests, if available, will be important to include.

255






Figure 1. Summary of studies citing the global soil respiration database (SRDB) between 2010 and 2019. More and more studies are using SRDB since the first version (SRDB-V1) was published (Bond-Lamberty and Thomson, 2010a).

260




265

Figure 2. Spatial distribution of soil respiration (R_s) sites. The gray circles are R_s sites from the fourth version of global soil respiration database (SRDB-V4, $n=1584$); the red dots are sites from the literature published in Chinese and added in the fifth version of global soil respiration database (SRDB-V5, $n=41$), the orange dots represent sites from the literature published in Russian and added in SRDB-V5 ($n=16$); the blue dots are sites from the literature published in other languages (mainly in English) and added in the SRDB-V5 ($n=840$). The size of circles represents the sample size at each measurement site (i.e., bigger circles represent more data).



270 **Figure 3.** Comparison of mean annual temperature (MAT, °C) in the globe (in red) vs. MAT from the sites in the
 global soil respiration database (SRDB, in teal) by the vegetation types. SRDB-4 represents the older SRDB
 released in 2018 and SRDB-V5 represents the newest SRDB published in 2020. Data from SRDB cover ten
 275 vegetation types (Agriculture, Arctic, Desert, tropical forest (Tropic FOR), temperate and boreal forest (T&B FOR),
 Grassland, Savanna, Shrubland, Urban, and Wetland). Comparing the fourth version (SRDB-V4) to the newest
 version (SRDB-V5), MAT values of Agriculture, Forest, and Grassland sites generally well represent the global
 MAT; in contrast, MAT from Shrubland sites in the database did not well represent global means in the older
 SRDB-V4, but their representation significantly improved in the newest SRDB-V5; for other vegetation types
 (Arctic, Desert, Savanna, Urban, and Wetland (including peatland) in the right panel), the MAT of the database sites
 do not well represent the global MAT distribution. Note that the Barren region was masked using MODIS landcover
 280 data. The number within each panel represent number of records for each vegetation type.



285 **Figure 4.** Comparison of mean annual precipitation (MAP, mm) in the globe (in red) vs. MAP from the sites in the global soil respiration database (SRDB, in teal) by the vegetation types. SRDB-V4 is the older SRDB published in 2018 and SRDB-V5 is the newest SRDB published in 2020. Data from SRDB covered ten vegetation types (see Figure 3). Sites from Agriculture, Savanna, Forest, and Urban generally well represent the global MAP (left panel), while sites from Arctic, Desert, Grassland, Shrubland, and Wetland (including peatland) do not have a good MAP representation (right panel). Note that the Barren region was masked using MODIS landcover data. The number of records of each panel is the same as Figure 3.

290




Figure 5. Comparison of annual soil respiration (R_s) and seasonal R_s (growing, dry, and wet seasons, spring, summer, autumn, and winter) observations from SRDB-V4 vs. that from SRDB-V5. In summary, adding new measurements does not change the distribution of annual R_s or seasonal R_s in the databases.

295

4. Discussion

4.1 Forecasting global R_s , R_H , and R_A

The updated SRDB-V5 provides opportunities for constraining global R_s estimates in the future. Currently, estimated global R_s ranged from 68-101 Pg C yr⁻¹, with many uncertainties associated with measurements and propagation of errors evident when upscaling site-specific R_s measurements to regional and global scales (Bond-Lamberty and Thomson, 2010b; Jian et al., 2018a, 2018b; Raich et al., 2002; Raich and Potter, 1995; Raich and Schlesinger, 1992; Warner et al., 2019). For example, R_s has been usually measured during daylight hours, implicitly assuming that measurements during this period represent the mean daily R_s . In a water-limited ecosystem, however, Cueva et al. (2017) estimated a time-of-day bias ranging from -29 to +40%. On the global scale, based on the HGRsD, Jian et al. (2018c) found that not measuring R_s 24-hours continuously contributed less than 6% of bias when estimating diurnal R_s . Quantifying the amount of bias required detailed information about when R_s was measured and how long the measurement lasted (Jian et al., 2018c). In the SRDB-V5, we revised all the studies and

300

305

collected the “*Time_of_day*” information, which should enable future analyses of how R_s measurement bias is related to when R_s measurements were collected.

310

It is also widely accepted that chamber properties (e.g., volume, area) (Davidson et al., 2002) and collar insertion depth (Heinemeyer et al., 2011) affect the R_s measurement accuracy, but in global scale, this has not been quantitatively tested before to our knowledge. We added information in the SRDB-V5 to enable researchers to investigate whether chamber area (smaller chambers are more vulnerable to edge effects, while larger chambers may experience inadequate air mixing), collar height (which may affect air mixing in the chamber), and insertion depth (which may cut off roots) affect R_s measurement accuracy and bias at seasonal to annual scales.

315

Comparing SRDB-V1 through SRDB-V5, we found that the uneven spatial distribution of R_s sites has improved, but bias still remains, with measurements conducted unevenly around the world and in climate space (Figure 2-4). The reason for the spatially-uneven coverage of R_s sites is a combination of economy, national policy, environmental conditions, spatial heterogeneity, and many other issues. Most obviously, the northern hemisphere has much more data than the southern hemisphere, as the most economically developed and wealthiest countries tend to be in the middle latitude of the Northern hemisphere, and thus more funds, infrastructure, and a broader and deeper pool of students and technical experts are all available to support on-site R_s measurement in these regions.

320

Improving modelling frameworks may help mitigate the uneven spatial distribution of R_s sites. For example, Jian et al. (2018b) found that how R_s responds to temperature is significantly different among climate regions, and therefore climate-specific models may be more appropriate than a global single model to estimate global R_s . Alternatively, machine learning approaches that account for non-linearity and multiple potential combinations of environmental factors have been used to estimate global R_s (Warner et al., 2019). SRDB-V5 also significantly increased the R_s sample size, and analyses could be conducted to test whether the increasing sample size of R_s helps reduce uncertainty when upscaling from site to global scale R_s . We recognize that there are many other possible sources of bias, but it is nonetheless possible that the biogeochemistry community will be able to use SRDB-V5 to improve the confidence of global R_s modeling and constrain global carbon cycle estimates.

325

Linking SRDB-V5, MGRsD, HGRsD, and COSORE provides an opportunity for global R_H and R_A estimates. Soil respiration mainly consists of two parts, R_H and R_A , but it is difficult to separate these two components, and much less R_H and R_A data are available in the SRDB (Bond-Lamberty and Thomson, 2010a). Due to a lack of data, far fewer studies have analyzed R_H and R_A and estimated global R_H and R_A in the past decades. According to our knowledge, there are only four global R_H (or R_A) estimates based on the very limited extant data ($n < 500$) (Hashimoto et al., 2015; Konings et al., 2019; Tang et al., 2020; Warner et al., 2019). In the “srdbequations” file, response of R_H and R_A to temperature and moisture information will be recorded, which will inspire the study of R_H and R_A and how they respond to temperature and soil moisture in the future. Further, we argue that a big advantage of global soil respiration databases with finer temporal resolution (i.e., MGRsD, HGRsD, and COSORE) is that the sample size of R_H and R_A could be greatly increased (e.g., sample size could be ten-fold increased if using monthly time-scale). In addition, the spatial coverage of R_H and R_A data could also be improved, because sites not measured year-round R_H and R_A in SRDB (due to annual time-scale) could be compiled into MGRsD and HGRsD whenever R_H or R_A was measured. Based on the monthly R_H and R_A data and how they related to environmental conditions (such as temperature and precipitation), monthly global R_H and R_A products could be generated, which provide useful data products for the Earth System Models’ (ESMs) benchmarking. The disadvantages of the smaller timescale databases (MGRsD, HGRsD, and COSORE) is that those databases usually have much less spatial coverage; and much more data is available from the growing season than from the non-growing season. Therefore, spatial upscaling including time may result in additional bias and associated uncertainty that must be carefully investigated.

340

345

350

355 4.2 Perspective

The updated SRDB-V5 will further support the analysis of how different manipulations affect Rs. In the past decades, many field experiments have been conducted to study different questions, for example, how soil carbon responds to global climatic warming and changes in precipitation patterns (Vicca et al., 2014); or how human activities (forest management, agriculture cultivation, and pollution) affect terrestrial carbon cycling and soil carbon stock (Carrillo et al., 2014; Jasek et al., 2014). However, inconsistent results from different experiments have generated debate regarding the effects of environmental factors and manipulations in Rs. Now SRDB-V5 includes Rs measurements from both control and different kinds of treatments, providing opportunities for synthesis analysis of how manipulation affects Rs. However, these treatment data about Rs measurements were rarely used in the past decade, as the manipulation information in older versions of SRDB was not standardized and thus could not easily be used. The updated and standardized SRDB-V5 manipulation codes have the potential to enable manipulation-driven studies on the macro-to-global scale.

4.3 Future improvements

We made an effort to resolve some issues in the old versions of SRDB (V1-V4), but the database needs to be continuously improved in the future. There is much more potentially useful information that could be included in future SRDB updates, although it is important to remember that every additional piece of information comes with a never-ending cost (in terms of data entry time, quality assurance/quality control, etc). 1) *Number_of_collar*: The number of collars within a certain study area is important information to evaluate the representability of the Rs measurements; 2) Soil organic carbon (SOC) measured *in-situ* or from regional or global estimates (Guevara et al., 2020; Hengl et al., 2017); 3) currently, *Site_ID* in SRDB-V5 are only comparable with *Site_ID* of MGRsD and HGRsD, further updates to *Site_ID* so it can connect with more external datasets [e.g., FLUXNET, COSORE, and AmeriFlux and a global database of forest carbon stocks and fluxes (ForC) (Anderson-Teixeira et al., 2018b)]; 4) *Annual_soil_moisture*: including a mean value of soil moisture or intra-annual soil variability derived from remote sensing (Guevara and Vargas, 2019) when this variable was not measured at the site. In addition, some meta information can be improved. For example, there are still 276 manipulation types in the SRDB-V5, and many manipulation types (n=96 out of 276) with only 1 row of records. Efforts could be made in the next version of database update to further simplify the manipulation of SRDB. We recognize that with thousands of publications included in the SRDB, it is known that some entries are incorrect and some information may have been missed during literature collection. In the past years, users have pointed out many data input errors and missing data issues on the SRDB, we made a great effort to check and many corrections have been made. However, it is inevitable that mistakes and missing information still exist, therefore, there is a pressing need to continue with the development of quality assurance and quality control for each update.

4.4 Reducing interoperability barriers

High interoperability is needed to maximize the benefits of SRDB-V5 to improve our understanding of the global carbon cycle. Interoperability has been defined as an organized collective effort with the ultimate goal to maximize sharing and using information to produce knowledge; and high interoperability is achieved by reducing conceptual, technological, organizational and cultural barriers (Vargas et al., 2017). The improved SRDB-V5 has reduced conceptual barriers as it provides a standardized and replicable framework to organize global Rs information that has been used for over a decade (Bond-Lamberty and Thomson, 2010a). It has reduced technological barriers by improving standardization of data fields (see Tables 1-3), data formats compatible with other databases, and providing flexible R scripts (details please see section 2.4) in a Github repository for end users and potential data contributors. We recognize that measuring Rs has other technological barriers (e.g., standardization of instrumentation, electrical power supply) that limit the collection of new measurements in harsh environments or wide implementation in developing countries. Organizational barriers remain a challenge as this is a bottom-up effort in need of long-term support to continue improving the quality and developing the new versions of the SRDB.

400 Finally, we believe that cultural barriers have been reduced as the global scientific community has improved in recognizing the importance of standardized databases and data sharing following FAIR principles.

5. Data availability

405 Findability and Accessibility were well considered and described when SRDB-V1 was published (Bond-Lamberty and Thomson, 2010a). To summarize the updating progress, SRDB-V1 was the first availability of the full data set, released on 2010/05/28; SRDB-V2 was released on 2012/03/13, R_s date of publications from 2011 was integrated into the database; SRDB-V3 was released on 2014/08/04, R_s data of literature from 2012 was collected and added; SRDB-V4 was released on 2018/11/21, R_s data of literature through 2015 were collected and compiled into the database; and SRDB-V5 was released on 2020/04/24, R_s data of literature from 2017 was collected and added (Jian and Bond-Lamberty, 2020). The version release information was recorded at the Oak Ridge National Laboratory's Distributed Active Archive Center ORNL-DAAC (Submitted). All data and code to reproduce the results in this study can be found at: Jian, Jinshi, Bond-Lamberty, Ben. (2020). jinshijian/ESSD: SRDB-V5 first release (Version v1.0.0) [Data set]. Zenodo. <http://doi.org/10.5281/zenodo.3876443>.

Conclusion

415 A global soil respiration database (SRDB) was developed to integrate soil respiration measurements from the globe a decade ago. Since the first release in 2010 (SRDB-V1), it has been widely used to advance our understanding of carbon dynamic related questions. Here, we restructured SRDB to a new version (SRDB-V5) following FAIR principles. We show that the SRDB substantially improved its representativeness compared with the older versions (SRDB-V1 to SRDB-V4, Figure S1 and S2) and improved its spatial coverage. A primary goal of SRDB-V5 is to improve the interoperability and reusability, and make it possible for scientists to contribute in the future with the ultimate goal to improve our understanding of the global carbon cycle. With those goals in mind, the revised SRDB-V5 is now more user-friendly for the ecology, biogeochemistry, and modeling communities.

Acknowledgments

425 This research was supported by the US Department of Energy, Office of Science, Biological and Environmental Research as part of the Terrestrial Ecosystem Sciences Program. The Pacific Northwest National Laboratory is operated for DOE by Battelle Memorial Institute under contract DE-AC05-76RL01830. We would like to thank Dalei Hao for his help with the MODIS landuse data processing. R.V. acknowledges support from NASA CMS grant number 80NSSC18K0173.

430 Author contributions

B.B.-L and J.J. designed the new version global soil respiration database (SRDB-V5). B.B.-L searched and downloaded the new papers until 2017 and compiled the meta-information. B.B.-L, M.H., R.M., J.M., D.P. and J.J. contributed to data collection, N.K. collected data in Russian, K.A.T. and V.H. raised many useful suggestions while working to integrate with ForC, R.V. and E.S. provided feedback and insights in all phases. J.J. wrote the manuscript in close collaboration with all authors.

Competing interests

The authors declare no competing interests.

440 Using and citing SRDB-V5

SRDB-V5 can be used for individual, academic, research, commercial, and other usage, and can be repackaged without written permission. Research and non-research products using SRDB-V5 should cite this publication.

References

- 445 Anderson-Teixeira, K., ValentineHerr, Hua, M. W. M., Bond-Lamberty, B., isamcano, LeBauer, D. and Padial, A. J. S.: forc-db/ForC: Version for IPCC 2019 updated default tables for biomass growth of natural forests in the (sub)tropics., 2018a.
- Anderson-Teixeira, K. J., Wang, M. M. H., McGarvey, J. C., Herrmann, V., Tepley, A. J., Bond-Lamberty, B. and LeBauer, D. S.: ForC: a global database of forest carbon stocks and fluxes, *Ecology*, 99(6), 1507, doi:10.1002/ecy.2229, 2018b.
- 450 Barba, J., Cueva, A., Bahn, M., Barron-Gafford, G. A., Bond-Lamberty, B., Hanson, P. J., Jaimes, A., Kulmala, L., Pumpanen, J., Scott, R. L., Wohlfahrt, G. and Vargas, R.: Comparing ecosystem and soil respiration: Review and key challenges of tower-based and soil measurements, *Agric. For. Meteorol.*, 249, 434–443, doi:10.1016/j.agrformet.2017.10.028, 2018.
- 455 Batjes, N. H.: Harmonized soil property values for broad-scale modelling (WISE30sec) with estimates of global soil carbon stocks, *Geoderma*, 269, 61–68, doi:10.1016/j.geoderma.2016.01.034, 2016.
- Bond-Lamberty, B., Christianson, D. S., Malhotra, A., Pennington, S. C., Sihi, D., AghaKouchak, A., ... & Ataka, M. COSORE: A community database for continuous soil respiration and other soil-atmosphere greenhouse gas flux data. *Glob. Chang. Biol.*, <https://doi.org/10.1111/gcb.15353>, 2020.
- 460 Bond-Lamberty, B. and Thomson, A.: A global database of soil respiration data, *Biogeosciences*, 7(6), 1915–1926, doi:10.5194/bg-7-1915-2010, 2010a.
- Bond-Lamberty, B. and Thomson, A.: Temperature-associated increases in the global soil respiration record, *Nature*, 464(7288), 579–582, doi:10.1038/nature08930, 2010b.
- 465 Bond-Lamberty, B., Wang, C. and Gower, S. T.: A global relationship between the heterotrophic and autotrophic components of soil respiration?, *Glob. Chang. Biol.*, 10(10), 1756–1766, doi:10.1111/j.1365-2486.2004.00816.x, 2004.
- [Bond-Lamberty, B. and Thomson, A. M.: A Global Database of Soil Respiration Data, Version 4.0, ORNL DAAC \[online\] Available from: https://daac.ornl.gov/cgi-bin/download.pl?ds_id=1578&source=metadata](https://doi.org/10.1111/j.1365-2486.2004.00816.x), 2018.
- 470 Carrillo, Y., Dijkstra, F. A., Pendall, E., LeCain, D. and Tucker, C.: Plant rhizosphere influence on microbial C metabolism: the role of elevated CO₂, N availability and root stoichiometry, *Biogeochemistry*, 117(2-3), 229–240 [online] Available from: https://idp.springer.com/authorize/casa?redirect_uri=https://link.springer.com/article/10.1007/s10533-014-9954-5&casa_token=JPHHOot7m0sAAAAA:5hg3r6R12wfYUjNrszIxx713ilY57LARPd6AsFVi4qqzPKWdu2xDtKBKe9JkdbNkp7knorGO_E2SA6ylZg, 2014.
- 475 Cueva, A., Bullock, S. H., López-Reyes, E. and Vargas, R.: Potential bias of daily soil CO₂ efflux estimates due to sampling time, *Sci. Rep.*, 7(1), 11925, doi:10.1038/s41598-017-11849-y, 2017.
- Davidson, E. A., Belk, E. and Boone, R. D.: Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest, *Glob. Chang. Biol.*, 4(2), 217–227, doi:10.1046/j.1365-2486.1998.00128.x, 1998.
- 480 Davidson, E. A., Savage, K., Verchot, L. V. and Navarro, R.: Minimizing artifacts and biases in chamber-based measurements of soil respiration, *Agric. For. Meteorol.*, 113(1), 21–37, doi:10.1016/S0168-1923(02)00100-4, 2002.
- Epule, T. E.: A New Compendium of Soil Respiration Data for Africa, *Challenges*, 6(1), 88–97, doi:10.3390/challe6010088, 2015.

- 485 Friedl, M. A., McIver, D. K., Hodges, J. C. F., Zhang, X. Y., Muchoney, D., Strahler, A. H., Woodcock, C. E., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F. and Schaaf, C.: Global land cover mapping from MODIS: algorithms and early results, *Remote Sens. Environ.*, 83(1), 287–302, doi:10.1016/S0034-4257(02)00078-0, 2002.
- Guevara, M. and Vargas, R.: Downscaling satellite soil moisture using geomorphometry and machine learning, *PLoS One*, 14(9), e0219639, doi:10.1371/journal.pone.0219639, 2019.
- 490 Guevara, M., Arroyo, C. and Brunsell, N.: Soil Organic Carbon across Mexico and the conterminous United States (1991-2010), *Global* [online] Available from: https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019GB006219?casa_token=LWL2D4HFN4wAAAAA:pB-LbL0ifYvo83VtBTvbtUEBAPpiALnhv2mShJoEYGD0QJS_7_VHPuSkF-IYrRm0SxYkMuUuxXsekMzP, 2020.
- 495 Hashimoto, S., Carvalhais, N., Ito, A., Migliavacca, M., Nishina, K. and Reichstein, M.: Global spatiotemporal distribution of soil respiration modeled using a global database, *Biogeosciences*, 12, 4121–4132 [online] Available from: https://pure.mpg.de/pubman/faces/ViewItemOverviewPage.jsp?itemId=item_2161024, 2015.
- Heinemeyer, A., Di Bene, C., Lloyd, A. R., Tortorella, D., Baxter, R., Huntley, B., Gelsomino, A. and Ineson, P.: Soil respiration: implications of the plant-soil continuum and respiration chamber collar-insertion depth on measurement and modelling of soil CO₂ efflux rates in three ecosystems, *European Journal of Soil Science*, 62(1), 82–94, doi:10.1111/j.1365-2389.2010.01331.x, 2011.
- 500 Hengl, T., Mendes de Jesus, J., Heuvelink, B. M., Gerard, B. M., Heuvelink, B. M. G., Ruiperez Gonzalez, M., Kilibarda, M., Blagotic, A., Shangguan, W., Wright, M. N., Geng, X., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan, R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S. and Kempen, B.: SoilGrids250m: Global gridded soil information based on machine learning, *PLoS One*, 12(2), e0169748, doi:10.1371/journal.pone.0169748, doi:10.1371/journal.pone.0169748, 2017.
- 505 IGBP: The International Geosphere-Biosphere Programme: A Study of Global Change. The Initial Core Projects, IGBP Secretariat. [online] Available from: <https://play.google.com/store/books/details?id=s9m8zAEACAAJ>, 1990.
- Jasek, A., Zimnoch, M., Gorczyca, Z., Smula, E. and Rozanski, K.: Seasonal variability of soil CO₂ flux and its carbon isotope composition in Krakow urban area, Southern Poland, *Isotopes Environ. Health Stud.*, 50(2), 143–155, doi:10.1080/10256016.2014.868455, 2014.
- 510 Jian, J., Steele, M. K., Thomas, R. Q., Day, S. D. and Hodges, S. C.: Constraining estimates of global soil respiration by quantifying sources of variability, *Glob. Chang. Biol.*, 24(9), 4143–4159, doi:10.1111/gcb.14301, 2018a.
- Jian, J., Steele, M. K., Day, S. D. and Thomas, R. Q.: Future global soil respiration rates will swell despite regional decreases in temperature sensitivity caused by rising temperature, *Earth's Future* [online] Available from: <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018EF000937>, 2018b.
- 515 Jian, J., Steele, M. K., Day, S. D. and Thomas, R. Q.: Measurement strategies to account for soil respiration temporal heterogeneity across diverse regions, *Soil Biology and Biochemistry*, 125, 167–177 [online] Available from: <https://www.sciencedirect.com/science/article/pii/S0038071718302311>, 2018c.
- 520 Jian, J., Bond-Lamberty, B.: jinshijian/ESSD: SRDB-V5 first release (Version v1.0.0) [Data set]. Zenodo. <http://doi.org/10.5281/zenodo.3876443>, 2020.
- Jian, J., Bahn, M., Wang, C., Bailey, V. L. and Bond-Lamberty, B.: Prediction of annual soil respiration from its flux at mean annual temperature, *Agric. For. Meteorol.*, 287, 107961, doi:10.1016/j.agrformet.2020.107961, 2020.
- 525 Konings, A. G., Bloom, A. A., Liu, J., Parazoo, N. C., Schimel, D. S. and Bowman, K. W.: Global satellite-driven estimates of heterotrophic respiration, *Biogeosciences*, 16(11), 2269–2284, doi:10.5194/bg-16-2269-2019, 2019.
- Luo, Y. and Zhou, X.: *Soil Respiration and the Environment*, Elsevier. [online] Available from:

<https://play.google.com/store/books/details?id=BILt0bdU6AsC>, 2010.

Martin, L. J., Blossey, B. and Ellis, E.: Mapping where ecologists work: biases in the global distribution of terrestrial ecological observations, *Front. Ecol. Environ.*, 10(4), 195–201, doi:10.1890/110154, 2012.

530 Peel, M. C., Finlayson, B. L. and McMahon, T. A.: Updated world map of the Köppen-Geiger climate classification, [online] Available from: <https://hal.archives-ouvertes.fr/hal-00298818/>, 2007.

Raich, J. W. and Potter, C. S.: Global patterns of carbon dioxide emissions from soils, *Global Biogeochem. Cycles*, 9(1), 23–36, doi:10.1029/94GB02723, 1995.

535 Raich, J. W. and Schlesinger, W. H.: The global carbon dioxide flux in soil respiration and its relationship to vegetation and climate, *Tellus B Chem. Phys. Meteorol.*, 44(2), 81–99, doi:10.1034/j.1600-0889.1992.t01-1-00001.x, 1992.

Raich, J. W., Potter, C. S. and Bhagawati, D.: Interannual variability in global soil respiration, 1980-94, *Glob. Chang. Biol.*, 8(8), 800–812, doi:10.1046/j.1365-2486.2002.00511.x, 2002.

540 R Core Team: R: A Language and Environment for Statistical Computing, Version 3.6.1, [online] Available from: <https://www.R-project.org/>, 2019.

Schlesinger, W. H.: Carbon balance in terrestrial detritus, *Annual Reviews in Ecology and Systematics*, 8, 51–81, 1977.

545 Tang, X., Fan, S., Du, M., Zhang, W., Gao, S., Liu, S., Chen, G., Yu, Z., Yao, Y. and Yang, W.: Spatial-and temporal-pattern of global soil heterotrophic respiration in terrestrial ecosystems, *Earth System Science Data*, 12(2), 1037-1051, <https://doi.org/10.5194/essd-12-1037-2020>.

Vargas, R., Alcaraz-Segura, D., Birdsey, R., Brunzell, N. A., Cruz-Gaistardo, C. O., de Jong, B., Etchevers, J., Guevara, M., Hayes, D. J., Johnson, K., Loescher, H. W., Paz, F., Ryu, Y., Sanchez-Mejia, Z. and Toledo-Gutierrez, K. P.: Enhancing interoperability to facilitate implementation of REDD : case study of Mexico, *Carbon Management*, 8(1), 57–65, doi:10.1080/17583004.2017.1285177, 2017.

550 Vicca, S., Bahn, M., Estiarte, M., van Loon, E. E., Vargas, R., Alberti, G., Ambus, P., Arain, M. A., Beier, C., Bentley, L. P., Boroken, W., Buchmann, N., Collins, S. L., de Dato, G., Dukes, J. S., Escobar, C., Fay, P., Guidolotti, G., Hanson, P. J., Kahmen, A., Kröel-Dulay, G., Ladreiter-Knauss, T., Larsen, K. S., Lellei-Kovács, E., Lebrija-Trejos, E., Maestre, F. T., Marhan, S., Marshall, M., Meir, P., Miao, Y., Muhr, J., Niklaus, P. A., Ogaya, R., Peñuelas, J., Poll, C., Rustad, L. E., Savage, K., Schindlbacher, A., Schmidt, I. K., Smith, A. R., Sotta, E. D.,
555 Suseela, V., Tietema, A., van Gestel, N., van Straaten, O., Wan, S., Weber, U. and Janssens, I. A.: Corrigendum to “Can current moisture responses predict soil CO₂ efflux under altered precipitation regimes? A synthesis of manipulation experiments,” *Biogeosciences*, 11(12), 3307–3308, doi:10.5194/bg-11-3307-2014, 2014.

Virkkala, A.-M., Abdi, A. M., Luoto, M. and Metcalfe, D. B.: Identifying multidisciplinary research gaps across Arctic terrestrial gradients, *Environ. Res. Lett.*, 14(12), 124061, doi:10.1088/1748-9326/ab4291, 2019.

560 Warner, D. L., Bond-Lamberty, B., Jian, J., Stell, E. and Vargas, R.: Spatial Predictions and Associated Uncertainty of Annual Soil Respiration at the Global Scale, *Global Biogeochem. Cycles*, 7, 983, doi:10.1029/2019GB006264, 2019.

565 Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., Hoen, P. A. C., Hooft, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J. and Mons, B.: The FAIR Guiding Principles for scientific data

570 management and stewardship, *Sci Data*, 3, 160018, doi:10.1038/sdata.2016.18, 2016.

Willmott, C. J. and Matsuura, K.: Terrestrial air temperature and precipitation: Monthly and annual time series (1950--1999) Version 1.02, Center for Climatic Research, University of Delaware, Newark, 2001.