

The global forest above-ground biomass pool for 2010 estimated from high-resolution satellite observations

Maurizio Santoro¹, Oliver Cartus¹, Nuno Carvalhais^{2,3}, Danaë M. A. Rozendaal^{4,5,6}, Valerio Avitabile⁷, Arnau Araza⁴, Sytze de Bruin⁴, Martin Herold⁴, Shaun Quegan⁸, Pedro Rodríguez-Veiga^{9,10}, Heiko Balzter^{9,10}, João Carreiras⁸, Dmitry Schepaschenko^{11,12,13}, Mikhail Korets¹⁴, Masanobu Shimada¹⁵, Takuya Itoh¹⁶, Álvaro Moreno Martínez^{17,18}, Jura Cavlovic¹⁹, Roberto Cazzolla Gatti^{20,21}, Polyanna da Conceição Bispo^{9,22,23}, Nasheta Dewnath²³, Dewnath²², Nicolas Labrière²⁴, Labrière²³, Jingjing Liang²⁵, Liang²⁴, Jeremy Linsell^{26,27}, Linsell^{25,26}, Edward T.A. Mitchard²⁸, Mitchard²⁷, Alexandra Morel²⁹, Morel²⁸, Ana Maria Pacheco Pascagaza⁹, Casey M. Ryan²⁸, Ryan²⁷, Ferry Slik³⁰, Slik²⁹, Gaia Vaglio Laurin³¹, Laurin³⁰, Hans Verbeeck³², Verbeeck³¹, Arief Wijaya³³, Wijaya³², Simon Willeeck³⁴, Willcock³³

hat formatiert: Nicht Hochgestellt/ Tiefgestellt

¹Gamma Remote Sensing, 3073 Gümligen, Switzerland

²Max Planck Institute for Biogeochemistry, Hans Knöll Strasse 10, 07745 Jena, Germany

³Departamento de Ciências e Engenharia do Ambiente, DCEA, Faculdade de Ciências e Tecnologia, FCT, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

⁴Laboratory of Geo-Information Science and Remote Sensing, Wageningen University & Research, Droevendaalsesteeg 3, 6708 PB Wageningen, The Netherlands

⁵Plant Production Systems Group, Wageningen University & Research, P.O. Box 430, 6700 AK, Wageningen, The Netherlands

⁶Centre for Crop Systems Analysis, Wageningen University & Research, P.O. Box 430, 6700 AK, Wageningen, The Netherlands

⁷European Commission, Joint Research Centre, Ispra, Italy

⁸National Centre for Earth Observation (NCEO), University of Sheffield, Sheffield, S3 7RH, United Kingdom

⁹Centre for Landscape and Climate Research, School of Geography, Geology and the Environment, University of Leicester, LE1 7RH, United Kingdom

¹⁰National Centre for Earth Observation, University of (NCEO), Leicester, LE1 7RH, United Kingdom

¹¹International Institute for Applied Systems Analysis, Schlossplatz 1, A-2361 Laxenburg, Austria

¹²Center of Forest Ecology and Productivity of the Russian Academy of Sciences, Profsoyuznaya 84/32/14, Moscow, 117997, Russia

¹³Institute of Ecology and Geography, Siberian Federal University, 660041, Krasnoyarsk, 79 Svobodny Prospect, Russia

¹⁴Laboratory of Ecophysiology of Permafrost Systems, V.N. Sukachev Institute of Forest of the Siberian Branch of Russian Academy of Sciences – separated department of the KSC SB RAS, Krasnoyarsk, 660036, Russia

¹⁵Tokyo Denki University, School of Science and Engineering, Division of Architectural, Civil and Environmental Engineering

¹⁶Remote Sensing Technology Center of Japan, Tokyu Reit Toranomon Bldg, 3f, 3-17-1 Toranomon, Minato-Ku, Tokyo, 105-0001, Japan

¹⁷Image Processing Laboratory (IPL), Universitat de València, València, Spain

¹⁸Numerical Terradynamic Simulation Group (NTSG), University of Montana, Missoula, USA

¹⁹University of Zagreb, Faculty of Forestry and Wood Technology, Department of Forest Inventory and Management, Svetosimunska cesta 2523, 10000, Zagreb, Croatia

²⁰Biological Institute, Tomsk State University, 634050 Tomsk, Russia

²¹Konrad Lorenz Institute for Evolution and Cognition, 3400 Klosterneuburg, Austria

²²School²³ School of Environment, Education and Development/Department of Geography, University of Manchester, Oxford Road, M13 9PL Manchester, United Kingdom

²³Guyana²², Guyana Forestry Commission, 1 Water Street, Kingston, Georgetown, Guyana

hat formatiert: Deutsch (Schweiz)

hat formatiert: Deutsch (Schweiz)

hat formatiert: Deutsch (Schweiz)

hat formatiert: Englisch (Vereinigtes Königreich)

45 ²⁴Laboratoire²³Laboratoire Évolution et Diversité Biologique, UMR 5174 (CNRS/IRD/UPS), 31062 Toulouse Cedex 9, France
²⁵Department²⁴Department of Forestry and Natural Resources, Purdue University
²⁶A²⁵A Rocha International, Cambridge, United Kingdom
²⁷The²⁶The RSPB Centre for Conservation Science, Bedfordshire, United Kingdom
50 ²⁸University²⁷University of Edinburgh, School of GeoSciences, Crew Building, The King's Buildings, Edinburgh, EH9 3FF, United Kingdom
²⁹Department²⁸Department of Geography and Environmental Sciences, University of Dundee, United Kingdom
³⁰Faculty²⁹Faculty of Science, University Brunei Darussalam, Jln Tungku Link, Gadong, BE1410, Brunei Darussalam
amma Remote Sensing, 3073 Gümligen, Switzerland
55 ³¹Department³⁰Department for Innovation in Biological, Agro-Food and Forest Systems (DIBAF), University of Tuscia, 01100 Viterbo, Italy
³²CAVELab³¹CAVELab – Computational and Applied Vegetation Ecology, Department of Environment, Ghent University, Coupure Links 653, 9000 Gent, Belgium
mma Remote Sensing, 3073 Gümligen, Switzerland
60 ³³World³²World Resources Institute, Indonesia (WRI Indonesia), Department of Research, Data & Innovation, Wisma PMI, 3rd Floor, Jl. Wijaya I/63, Kebayoran Baru, South Jakarta, Indonesia
³⁴Schoel³³School of Natural Sciences, Bangor University, United Kingdom

Correspondence to: Maurizio Santoro (santoro@gamma-rs.ch)

Abstract. The terrestrial forest carbon pool is poorly quantified, in particular in regions with low forest inventory capacity. By combining multiple satellite observations of synthetic aperture radar (SAR) backscatter around the year 2010, we generated 65 a global, spatially explicit dataset of above-ground forest biomass (dry mass, AGB) with a spatial resolution of 1 ha. Using an extensive database of 110,897 AGB measurements from field inventory plots, we show that the spatial patterns and magnitude of AGB are well captured in our map with the exception of regional uncertainties in high carbon stock forests with AGB > 250 Mg ha⁻¹ where the retrieval was effectively based on a single radar observation. With a total global AGB of 522 Pg, our estimate of the terrestrial biomass pool in forests is lower than most estimates published in literature (426 - 571 Pg). 70 Nonetheless, our dataset increases knowledge on the spatial distribution of AGB compared to the global Forest Resources Assessment (FRA) by the Food and Agriculture Organization (FAO) and highlights the impact of a country's national inventory capacity on the accuracy of the biomass statistics reported to the FRA. We also reassessed previous remote sensing AGB maps, and identify major biases compared to inventory data, up to 120% of the inventory value in dry tropical forests, in the sub-tropics and temperate zone. Because of the high level of detail and the overall reliability of the AGB spatial patterns, 75 our global dataset of AGB is likely to have significant impacts on climate, carbon and socio-economic modelling schemes, and provides a crucial baseline in future carbon stock changes estimates. The dataset is available at:
<https://doi.pangaea.de/10.1594/PANGAEA.894711> (Santoro, 2018)(Santoro, 2018).

1 Introduction

Above-ground live biomass (AGB) is identified as one of 54 Essential Climate Variables (ECVs) by the Global Climate 80 Observing System (GCOS) because of its major role in the global carbon cycle. Biomass stores carbon removed from the atmosphere by photosynthesis in long-lived woody pools; and yields to carbon emissions to the atmosphere when disturbed.

Hence, accurate knowledge of its magnitude and spatial distribution is a key, and currently poorly constrained, part of the carbon cycle (Houghton, 2005) (Houghton, 2005). Information on forest biomass is required to quantify forest resources and determine their benefit in terms of ecosystem services (Reichstein and Carvalhais, 2019; Schepaschenko et al., 2015),
85 infer (Schepaschenko et al., 2015; Reichstein and Carvalhais, 2019), climate change mitigation and biodiversity conservation (Soto-Navarro et al., 2020). Biomass estimates allow inferring emissions from forest degradation (Houghton et al., 2009; Li et al., 2017) (Houghton et al., 2009; Li et al., 2017) and assist assisting with the derivation of emission factors (Herold et al., 2019; IPCC, 2006). This information (IPCC, 2006; Herold et al., 2019). Information on biomass also directly supports policy by quantifying national carbon stocks in the context of reducing emissions from deforestation and degradation (REDD+), the
90 Paris Agreement on Climate Change and the United Nations Sustainable Development Goals (Herold et al., 2019) (Gibbs et al., 2007; Herold et al., 2019). Finally, an improved knowledge on carbon stock patterns and dynamics from a better knowledge of forest biomass pools helps to constrain Earth System models (Bloom et al., 2016; Carvalhais et al., 2014; Ciais et al., 2014; Exbrayat et al., 2019; Bloom et al., 2016; Thurner et al., 2016; Baccini et al., 2017; Thum et al., 2017; Le Quéré et al., 2018; ThumExbrayat et al., 2017; Thurner et al., 2016 2019).

95 Plants Previous estimates have suggested that plants store about 80% of the live biomass forming the Earth's biosphere, with an estimated carbon pool of 450 GtC PgC (Bar-On et al., 2018). Around 320 GtC is in PgC were allocated to the AGB, representing approximately 70% of the overall pool, most of it stored in woody biomass (Bar-On et al., 2018). However, our knowledge of the terrestrial woody biomass stock is poorly known relatively uncertain (Houghton et al., 2009); global assessments of biomass in forests for the year 2007 reported 362 GtC (Houghton et al., 2009). This uncertainty is well illustrated by the variance among forest biomass estimates from inventory data. A global assessments of biomass in forests for the year 100 2007 reported 362 PgC based on a compilation of forest inventory resources (Pan et al., 2011) (Pan et al., 2011) and whereas approximately 300 GtC PgC were reported for the year 2010 based on the national contributions to the Food and Agriculture Organization (FAO) Forest Resources Assessment (FRA) (FAO, 2010). This uncertainty is a consequence of the uneven 105 characterization of AGB in terms of precision and timeliness of measurements (Ciais et al., 2014; Houghton et al., 2009), and the lack of a universal inventory system using a standard set of survey and reporting procedures (Houghton et al., 2009; Ciais et al., 2014), and the lack of a universal inventory system using a standard set of survey and reporting procedures. Most countries in the temperate and boreal zones have National Forest Inventories (NFIs) that use systematic regular sampling, albeit some national differences e.g., in the definition of forest area (Tompoo et al., 2010). In contrast, many of the tropical 110 countries have less developed inventory infrastructures or have only recently started to develop such infrastructure, often with the support of international initiatives (e.g., the UN REDD programme).

115 Remote sensing observations allow the estimation of global ecosystem properties and parameters (Schimel et al., 2015). No single observable derived measurement from remote sensing, however, directly measures AGB represents a direct measure of the forest AGB. Nonetheless, the demand on spatially explicit estimates of AGB and the wide range of satellite observations

hat formatiert: Deutsch
Feldfunktion geändert
hat formatiert: Deutsch
hat formatiert: Deutsch
hat formatiert: Deutsch
hat formatiert: Deutsch

collected in the last decades has fostered the development of a multitude of retrieval models based either on empirical regression techniques, physically-based mathematical models or machine learning algorithms (Lucas et al., 2015; Lu et al., 2016; Santoro and Cartus, 2018) (Lucas et al., 2015; Lu et al., 2016; Santoro and Cartus, 2018). The incapacity of remote sensing to measure biomass and the approximations in retrieval models cause inaccurate estimates of AGB at the pixel level.

120 Even the spatial distribution of AGB in global and biome-specific maps of remotely sensed AGB (Kindermann et al., 2008; Saatchi et al., 2011b; Baccini et al., 2012; Thurner et al., 2014; Liu et al., 2015; Avitabile et al., 2016; Hu et al., 2016) (Kindermann et al., 2008; Saatchi et al., 2011b; Baccini et al., 2012; Thurner et al., 2014; Liu et al., 2015; Avitabile et al., 2016; Hu et al., 2016) presents sometimes remarkable differences (Mitchard et al., 2013; Ometto et al., 2014; Schepaschenko et al., 2015; Rodríguez-Veiga et al., 2017), implying a strong variability of the global biomass pool estimate

125 (Table S1).

Feldfunktion geändert

hat formatiert: Deutsch

Global datasets of forest AGB from remote sensing observations mostly represent the period around the year stocks for a snapshot ranging between 2000 and 2010, and their coarse spatial resolution (≥ 500 m) hinders description of the fine-scale spatial variability of biomass, which, This aspect is of major importance when trying to capture changes in land use, natural disturbances and growth patterns (Houghton et al., 2009) (Houghton et al., 2009) or monitor management practices (Erb et al., 2017). Here, we assembled the currently widest set of publicly available RADAR, LiDAR and optical satellite observations suited to estimate forest biomass with the objective of generating a high-resolution, global map of spatially explicit estimates of AGB (Erb et al., 2018). Here, we assembled a wide set of publicly available RADAR, LiDAR and optical satellite observations suited to estimate forest variable with the objective of generating a high-resolution, global map of spatially explicit estimates of forest AGB (unit: Mg of dry mass per hectare) so to provide more recent, more detailed and possibly more accurate information on the spatial distribution of global AGB with respect to existing datasets. Our forest AGB map has a pixel size of 1 ha and is based on satellite remote sensing observations from around the year 2010. Here, we present the dataset together with an assessment of its validity using an extensive database of plot-level measurements of AGB. The significance of our map estimates is demonstrated in the context of biomass stock assessments by benchmarking with respect to the FAO

130 FRA country statistics. In addition, we compare our estimates and other published estimates of forest AGB derived from remote sensing observations with plot-based AGB measurements from inventory plots to prove illustrate the overall reliability of our estimates.

2 Material and Methods

2.1 Satellite data

145 The spatially explicit estimates of AGB were based on the radar backscattered intensity recorded by the Phase Phased Array-type L-band Synthetic Aperture Radar (PALSAR) instrument, onboard the Advanced Land Observing Satellite (ALOS) satellite, and the Advanced Synthetic Aperture Radar (ASAR) instrument operating at C-band, onboard the Environmental

Satellite (Envisat) (Supplement Section A.1). In addition, LiDAR-based metrics and surface reflectances were used throughout the process of biomass estimation.

150

ALOS PALSAR was an active microwave sensor operating at L-band (wavelength of 23 cm). Between 2006 and 2011, PALSAR acquired images in the Fine Beam ([FB Dual-polarisation \(FBD\)](#)) mode with [2520](#) m spatial resolution. Image acquisition followed a predefined observation scenario with the aim of achieving spatially and temporally consistent large-scale observational datasets. [Summer-time acquisitions from the FB \(Rosenqvist et al., 2007\)](#), [Summer-time acquisitions from the FBD](#) mode (mostly May to October) were used by the Japan Aerospace Exploration Agency (JAXA) to generate yearly mosaics of the radar backscatter for each year between 2007 and 2010 ([Shimada, 2010](#)). [Each image was orthorectified and radiometrically terrain corrected to gamma0 \(Shimada, 2010\)](#). The mosaics are [publically publicly](#) available and are provided in the form of image tiles of $1^\circ \times 1^\circ$ in latitude and longitude resampled to a grid with a pixel spacing of [0.00022000225](#). In this study, we used the mosaics of the co-polarized Horizontal-transmit Horizontal-receive (HH) and cross-polarized Horizontal-transmit Vertical-receive (HV) backscatter images. Images from 14,728 tiles were used to estimate biomass. 96 image tiles showing evident radiometric offsets with respect to adjacent ones, due for example to different environmental conditions (e.g., frozen vs. unfrozen conditions), were manually replaced with the corresponding image tile from the mosaic of 2009. This replacement ensured homogeneity of the L-band backscatter dataset across all landscapes.

160

Envisat ASAR was an active microwave sensor acquiring images at C-band (wavelength of 6 cm) between 2002 and 2012. ASAR operated in four different modes over land, with a spatial resolution of 30 m (Image Mode and Alternating Polarization Mode), 150 m (Wide Swath Mode) and 1,000 m (Global Monitoring Mode). Approximately 80% of the total number of observations consisted of GM observations. We processed the entire dataset of ASAR images of the SAR backscatter made available by the European Space Agency (ESA) through the Grid Processing on Demand platform to stacks of terrain geocoded,

170

pixel-area normalized and speckle-filtered images (Santoro et al., 2015b). Images acquired with the IM and WSM were geocoded to a pixel size of 0.0013888° in latitude and longitude, corresponding to an area on the ground of roughly 150 m \times 150 m at the Equator. Images acquired with the GMM were geocoded to a pixel size of 0.01° in latitude and longitude, corresponding to an area on the ground of approximately 1,000 m \times 1,000 m at the Equator. To obtain global full coverage, the IM and WSM images were further averaged and resampled to the pixel size of the GMM dataset to form a single 1 km

175

dataset of C-band backscatter observations. Each image was divided into tiles of $2^\circ \times 2^\circ$ in latitude and longitude. For this study, we used all Envisat ASAR images acquired in 2010 and 2011. [The density of the observations in time \(Fig. S1\) shows decreased from the polar latitudes with several observations per day to the tropical latitudes with approximately 200 observations on average over two years. Areas with a small number of observations correspond to regions seldom imaged during the lifetime of the Envisat mission \(e.g. New Zealand, Japan\) or imaged only at high resolution when the overlap of images from adjacent orbitals track was null \(e.g., California, western Amazon\).](#)

The parameterization of the biomass retrieval models relating biomass to SAR backscatter observations was supported by the GLA14 data product of the Geoscience Laser Altimeter System (GLAS) on board the Ice Cloud and Land Elevation Satellite (ICESat) that operated between 2003 and 2009. GLA14 represented the waveforms over land only in the form of the parameters
185 of a multi-Gaussian model fitted to the raw waveforms (Hoffon et al., 2000)(Hoffon et al., 2000), thus containing information about the vertical structure of vegetation. Because GLAS observations consisted of approximately 65 m large footprints acquired every 170 m along track with a distance between tracks of the order of 60 km, the GLA14 dataset was not dense enough to allow direct spatially explicit estimates of biomass. Here, we used the entire archive of GLA14 data products, provided by the National Snow & Ice Data Centre (NSIDC), to estimate forest height after filtering for footprints affected by
190 topography and various noise sources in the waveforms (Los et al., 2012; Simard et al., 2011)(Los et al., 2012; Simard et al., 2011). In addition, we computed an estimate of the canopy density for each footprint as the ratio of energy received from the canopy (i.e., returns from above the ground peak) to the total energy received. Our database of GLAS-based metrics consisted of 26.5 million footprints homogeneously distributed over all vegetated surfaces. While the SAR observations were used as predictors in the retrieval model (Section 3), the LiDAR observations supported the estimation of parameters of the retrieval
195 model that are time invariant such as forest transmissivity (Section 3). Herewith, the time difference between the SAR imagery (2010) and the LiDAR observations (2003-2009) did not impact the retrieval.

Global reflectances of Landsat 7 images (bands 3, 4, 5 and 7) acquired in 2010 were used to rescale biomass estimates from ASAR to the pixel size of the ALOS PALSAR dataset. The dataset was available in the form of a mosaic from the USGS
200 website (Hansen et al., 2013). The square pixels of the mosaic had a spacing of 0.00027°, i.e., roughly 30 m at the Equator. The dataset was downloaded from Google Earth Engine and resampled with nearest neighbour to the geometry of the ALOS PALSAR dataset.

Feldfunktion geändert

2.2 AGB estimation

Unlike investigations targeting estimation of that directly relate AGB from to the remote sensing data (Lu et al., 2016; Santoro and Cartus, 2018)(Lu et al., 2016; Santoro and Cartus, 2018), we estimated the density of the woody volume, referred to as growing stock volume (GSV, unit: $m^3 ha^{-1}$) from which AGB is then computed, for three reasons. First, the signal backscattered by a forest is primarily affected by the density, and to some degree the height, of the trees (Santoro et al., 2015a)(Santoro et al., 2015a). However, the short wavelength of the radar instruments means that only the upper part of the volume is seen so that an estimate of GSV (or AGB) would be the result of an inference from the SAR observation. Second, it has not been demonstrated that the SAR backscatter at C- and L-band is sensitive to the wood density of trees for a given level of GSV, thus not providing experimental support to a direct estimation of AGB. Without such evidence, it is preferable to estimate a forest structural parameter from the SAR backscatter and convert it to AGB using a separate layer combining the wood density and the stem-to-total biomass expansion factor, which does not depend on remote sensing observations. An open question is whether means exist that allow an unbiased characterization of wood density and stem-to-total biomass expansion globally.
210

215 Third, our approach to estimate AGB from remote sensing data mimics the approach based on forest field inventory data (Brown, 1987; Jenkins et al., 2003) where GSV is acknowledged to be the major predictor of AGB. The relevance of GSV to estimation of AGB is further emphasised by the country reports building up the FAO 2010 FRA (FAO, 2010). Of the 233 country reports, we identified 171 countries reporting numbers on AGB, and for roughly two thirds of these (111) the estimate of AGB was derived from an estimate of GSV, based on inventory or expert knowledge, using a scaling factor.

220

The AGB retrieval algorithm is outlined in the flowchart in Fig. 1 showing the interdependencies of datasets and retrieval models. It is referred to Supplement and the Algorithm Theoretical Basis Document (ATBD) compiled in the ESA funded GlobBiomass project as a detailed documentation of the processing steps, their justification, evaluation and experimental support (Quegan et al., 2017).

225




Figure 1. Flowchart of the AGB retrieval approach.

We applied a model-based approach known as BIOMASAR (Cartus et al., 2012b; Santoro et al., 2011)(Santoro et al., 2011;

230 Cartus et al., 2012b) separately to the ALOS PALSAR (Cartus et al., 2012b) and the Envisat ASAR radar backscatter datasets (Santoro et al., 2011, 2015a)(Santoro et al., 2011, 2015a) to obtain two independent, spatially explicit estimates of GSV. BIOMASAR inverts a Water Cloud Model (Pulliainen et al., 1994; Santoro et al., 2002)(Pulliainen et al., 1994; Santoro et al., 2002).

$$235 \quad \sigma_{for}^0 = \sigma_{gr}^0 e^{-\beta V} + \sigma_{veg}^0 (1 - e^{-\beta V}) \quad (1)$$

in which σ^0_{for} represents the forest backscatter, σ^0_{gr} and σ^0_{veg} represent the backscattering coefficients of the ground and vegetation layer, respectively. The exponential function, $e^{\beta V}$, represents the two-way forest transmissivity, where β is an empirically defined coefficient expressed in m^{-1} and V represents GSV. Eq. (1) neglects multiple scattering, which is acceptable for most forest conditions (Cartus et al., 2012a; Cartus and Santoro, 2019; Santoro et al., 2011).

σ^0_{for} represents the forest backscatter, i.e., the SAR backscatter observation from an ALOS PALSAR or an Envisat ASAR image (Section 2.1). σ^0_{gr} and σ^0_{veg} represent the backscattering coefficients of the ground and vegetation layer, respectively. The exponential function, $e^{\beta V}$, represents the two-way forest transmissivity, where β is an empirically defined coefficient expressed in m^{-1} and V represents GSV. Eq. (1) neglects multiple scattering, which is acceptable for most forest conditions (Santoro et al., 2011; Cartus et al., 2012a; Cartus and Santoro, 2019).




Figure 1. Flowchart of the AGB retrieval approach.

The model parameters β , σ^0_{gr} and σ^0_{veg} need to be estimated in order to invert the model and obtain an estimate of GSV from a measurement of the SAR backscatter. Estimates of the coefficient β are obtained with a model-based approach that relates canopy density and GSV observations through the transmissivity of the forest (Santoro et al., 2015a)(Santoro et al., 2015a). The estimation is stratified by the FAO Global Ecological Zones (Supplement Section A.1, Fig. S1S2 and Table S2). To estimate σ^0_{gr} and σ^0_{veg} , we rely on a self-calibration approach (Santoro et al., 2011) rather than using a set of reference measurements of the SAR backscatter and *in situ* GSV values. The limited availability of *in situ* information on biomass (e.g., from inventory plots or laser-based maps) prevents adaptive calibration of retrieval algorithms using conventional approaches. In many areas, particularly the tropics, the number of available plots is very limited so that models can only be calibrated using

reference information collected over large areas (Bouvet et al., 2018) with the risk of missing spatial variability in the backscatter. The model training approach is tailored to the radar wavelength in order to accommodate the different relationships of backscatter to biomass (BIOMASAR-C and BIOMASAR-L, see Supplement Section A.2 and A.3).

When N observations of the radar backscatter are available, a final estimate of GSV, with higher accuracy compared to the individual estimates, is obtained by means of a weighted linear combination of the individual estimates of GSV obtained by inverting Eq. (1) for each backscatter observation (Kurvanen et al., 1999; Santoro et al., 2011).
265 (1) for each backscatter observation (Santoro et al., 2011; Kurvanen et al., 1999).

$$V_{mt} = \frac{\sum_{i=1}^N w_i V_i}{\sum_{i=1}^N w_i} \quad (2)$$

270 In Eq. (2), the weights w_i are defined as the difference ($\sigma_{veg,i}^0 - \sigma_{gr,i}^0$) so that GSV estimates obtained from images with the strongest sensitivity to GSV are preferred to those obtained from images with no sensitivity to GSV (Santoro et al., 2011).

In this study, the BIOMASAR-C implementation was tailored to ingest ASAR data (Supplement Section A.2) and generated a GSV map at 1,000 m spatial resolution by combining individual estimates from the [hyper-temporal](#)-ASAR dataset. The
275 BIOMASAR-L implementation was tailored to ingest ALOS PALSAR data and generate a GSV data product at 25 m spatial resolution (Supplement Section A.3). The step in Eq. (2) became redundant because of the strong correlation of the ALOS PALSAR mosaics in time and the negligible weight attributed to the HH-polarized component when combined with the HV-polarized component ([Quegan et al., 2017](#)).[\(DUE GlobBiomass - Algorithm Theoretical Basis Document\)](#). As a result, the retrieved GSV with BIOMASAR-L was based on the single observation of the L-band SAR backscatter at HV-polarization in
280 the mosaic for 2010.

To merge estimates of GSV obtained at 25 m and 1,000 m, the latter were re-scaled to 25 m using a linear regression model (BIOMASAR-C+ in Fig. 1). The HH- and HV-polarized ALOS PALSAR backscatter (σ_{HH}^0 and σ_{HV}^0) and the Landsat bands 3,4,5 and 7 (B_3, B_4, B_5 and B_7) were used as predictors in the model in Eq. (3):

$$\log(V) = a_0 + a_1 \sigma_{HV}^0 + a_2 \sigma_{HH}^0 + a_3 B_3 + a_4 B_4 + a_5 B_5 + a_6 B_7 \quad (3)$$

The model was calibrated for each $1^\circ \times 1^\circ$ tile at a pixel size of 1,000 m to predict the BIOMASAR-C estimate of GSV at the 25 m scale. A bias correction had to be performed when retransforming the logarithmic GSV predictions to linear scale. The
290 bias was computed by differencing the original BIOMASAR-C GSV and the predictions from Eq. (3) aggregated to the 1,000 m pixel size. In spite of its simplicity, the multiple linear regression resulted in an overall superior performance when compared

Feldfunktion geändert

to results obtained with more sophisticated re-scaling methods ([Quegan et al., 2017 DUE GlobBiomass - Algorithm Theoretical Basis Document](#)).

295 The final estimate of GSV was obtained by weighting the rescaled ASAR-based estimates (V_{C+}) and PALSAR-based estimates (V_L) of GSV. The weighting scheme accounted for the different sensitivity of C- and L-band data to GSV, the number of observations used for estimating GSV, local errors in the model training and inversion and residual uncompensated topographic effects in the ALOS PALSAR mosaics (Supplement Section A.4). The GSV estimated with BIOMASAR-L was given more weight in areas of high GSV except in the case of steep terrain (Fig. [S2S3](#)). BIOMASAR-C+ GSV estimates were instead
 300 preferred in areas of low GSV and regions with rugged terrain (Fig. [S2S3](#)).

$$GSV = V_{C+} \cdot w_C + V_L \cdot w_L \quad (4)$$

To reduce the pixel-wise variability due to speckle in the radar data and amplified by the weak sensitivity of the C- and L-band
 305 backscatter to forest variables, spatial averaging using a 4 x 4 window was applied. This decreased the spatial resolution of the GSV estimates to 0.000888° in both latitude and longitude, corresponding to an area of approximately 1 ha at the Equator.

The conversion of GSV to AGB in Eq. (5) requires wood density (WD) and the biomass expansion factors (BEF), which give the allometric relationship between stem mass and whole aboveground mass, including branches and leaves.

310

$$AGB = GSV \cdot WD \cdot BEF \quad (5)$$

The spatial variations in WD and BEF result from biological processes that respond to local conditions, as has been demonstrated in regional studies showing environmental controls on the patterns of WD and BEF (Chave et al., 2009; Thurner
 315 et al., 2014). Towards a global assessment of WD patterns, we collected published databases based on inventory data where wood density (WD) is reported and explored machine learning methods to maximize the information content in relevant environmental variables (Supplement Section A.5). The final dataset of WD was obtained by integrating several individual predictions (Supplement Section A.5). To estimate the BEF, we used the generalized power-law function relating branch and leaf biomass to stem biomass (SB) (Thurner et al., 2014).

320

$$BEF = \frac{p_1 SB^{p_2 + SB}}{SB} \quad (6)$$

Because of the uneven distribution of samples for which biomass component measurements were available (Supplement Section A.5), the model in Eq. (6) was fitted to measurements of BEF and stem biomass for tropical and extra-tropical forests

325 only, each stratified by leaf type (broadleaves, evergreen conifers and deciduous conifers). All BEF models decreased rapidly
 for increasing stem biomass reaching an asymptote for low to medium stem biomass depending on ecoregion and leaf type
 (Quegan et al., 2017). All BEF models decreased rapidly for increasing stem biomass reaching an asymptote for low to medium
 stem biomass depending on ecoregion and leaf type (DUE GlobBiomass - Algorithm Theoretical Basis Document). The
 asymptotic BEF for tropical broadleaves species was modelled with a value of 1.36, higher than other forest types, which were
 330 characterized by values of 1.15-1.20.

Spatially explicit estimates of WD and BEF were obtained at 0.01° spatial resolution and resampled to the pixel size of the GSV dataset using bi-cubic interpolation. The AGB estimates resulting from the product of the GSV, WD and BEF estimates were obtained at a spatial resolution of 0.000888° in both latitude and longitude, i.e., with a pixel size of 1 ha at the Equator.

335 2.3 Uncertainty model

The uncertainty of the AGB estimates was quantified by their standard deviation. The standard deviation of the GSV estimates obtained with the BIOMASAR-C approach, δV_c , was quantified by propagating the standard error of the measured SAR backscatter, σ^0_{meas} , and the estimates of the forest backscatter model parameters σ^0_{gr} , σ^0_{df} , β and V_{df} (Santoro et al., 2015a)(Santoro et al., 2015a).

$$340 \quad \delta V_c = \sqrt{\left[(\delta \sigma^0_{meas})^2 \cdot \left(\frac{\partial V}{\partial \sigma^0_{meas}} \right)^2 + (\delta \sigma^0_{gr})^2 \cdot \left(\frac{\partial V}{\partial \sigma^0_{gr}} \right)^2 + (\delta \sigma^0_{df})^2 \cdot \left(\frac{\partial V}{\partial \sigma^0_{df}} \right)^2 + \right.} \\ \left. + (\delta \beta)^2 \cdot \left(\frac{\partial V}{\partial \beta} \right)^2 + (\delta V_{df})^2 \cdot \left(\frac{\partial V}{\partial V_{df}} \right)^2 \right] \quad (7)$$

The same approach was applied to the BIOMASAR-L procedure, in which case the error model also included components related to the average canopy density of dense forests, η_{df} , the average height of dense forests, h_{df} , and the two-way attenuation

345 coefficient, α . Following the results in (Los et al., 2012; Simard et al., 2011)(Simard et al., 2011; Los et al., 2012), which validated GLAS-based height estimates at boreal, temperate, sub-tropical, and tropical forest sites, we assumed standard errors for height estimates at the GLAS footprint-level between 4 m (boreal zone) and 10 m (tropical zone). As indicated by Garcia et al. (2012), the estimation error of canopy cover from ICESAT GLAS as the ratio of energy returned from the canopy to the total energy returned may be of the order of 15 to 20 %. We therefore assume a global error of 20%.

350 Differently than for the C-band case (Santoro et al., 2015a)(Santoro et al., 2015a), the standard deviation of the coefficient β was inferred from the relationship of the forest transmissivity, simulated with the aid of GLAS height and optical canopy density estimates, and GSV (Quegan et al., 2017)(DUE GlobBiomass - Algorithm Theoretical Basis Document). The 95% bounds of the estimates increased from $+/-0.002$ ha/m³ in the case of low values of β that are valid in boreal and subtropical

dry forests to +/-0.007 ha/m³ for the highest values of β that are applied in the tropics. For the two-way attenuation coefficient α , we assume a standard error deviation of 0.25 dB/m, which is roughly consistent with the range of values reported in the literature ([Quegan et al., 2017](#))([DUE GlobBiomass - Algorithm Theoretical Basis Document](#)).

$$\delta V_L = \sqrt{(\delta \sigma_{meas}^0)^2 \cdot \left(\frac{\partial V}{\partial \sigma_{meas}^0}\right)^2 + (\delta \sigma_{gr}^0)^2 \cdot \left(\frac{\partial V}{\partial \sigma_{gr}^0}\right)^2 + (\delta \sigma_{df}^0)^2 \cdot \left(\frac{\partial V}{\partial \sigma_{df}^0}\right)^2 + (\delta h_{df})^2 \cdot \left(\frac{\partial V}{\partial h_{df}}\right)^2 + (\delta \eta_{df})^2 \cdot \left(\frac{\partial V}{\partial \eta_{df}}\right)^2 + (\delta \alpha)^2 \cdot \left(\frac{\partial V}{\partial \alpha}\right)^2} \quad (8)$$

360 In the case of BIOMASAR-C, the standard deviation of the multi-temporal GSV estimate was modelled as a linear combination
of the single-image GSV standard deviations from Eq. (9) ([Santoro et al., 2015a](#))[\(Santoro et al., 2015a\)](#).

$$\delta V_{c,mt} = \sqrt{\sum_{i=1}^N w_i^2 \delta(V_{c,i})^2} \quad (9)$$

365 The uncertainty associated with the predictions of GSV obtained by rescaling the BIOMASAR-C GSV estimates was related
to the uncertainty in the coarse resolution GSV estimates and the scaling factors.

The relative standard deviation of the BIOMASAR-C GSV estimates ($\delta V_c/V_c$) was modelled as a function of the GSV estimates
(V_c) by means of an exponential model ([Quegan et al., 2017](#))([DUE GlobBiomass - Algorithm Theoretical Basis Document](#)).

370 In Eq. (10), the model coefficients a , b and c were estimated by means of a least squares regression for each of the FAO Global
Ecological Zones.

$$\delta V_c/V_c = ae^{bV_c} + c \quad (10)$$

375 To characterize the error associated with the rescaling model, we used the root mean square difference (RMSD) between the
original BIOMASAR-C GSV and the BIOMASAR-C+ GSV estimates aggregated to the pixel size of the former. A polynomial
function of the fourth order was found to adequately reproduce the relationship between the GSV sets across ecozones and
was used to fit the observed trend in GSV ([Quegan et al., 2017](#)[DUE GlobBiomass - Algorithm Theoretical Basis Document](#)).

Feldfunktion geändert

380 In the process of rescaling, we assumed that the standard deviation scaled with the pixel area of the GSV predictions. In Eq.
(11), the scaling factor between standard deviations was represented by the ratio between the pixel areas of the rescaled product
(A_{c+}) and the BIOMASAR-C data product (A_c).

$$\delta V_{C+} = \delta V_C \sqrt{\frac{w_{C+}}{A_c}}$$

(11)

385

The standard deviation of the GSV estimates was obtained with the same weighted linear combination of the standard deviations of the BIOMASAR-L and the BIOMASAR-C+ datasets:

$$\delta GSV = \delta V_{C+} \cdot w_C + \delta V_L \cdot w_L \quad (12)$$

390

The standard ~~error~~deviation of AGB was expressed in terms of partial derivatives of its components

$$\delta AGB = \sqrt{\left(\frac{\partial AGB}{\partial WD}\right)^2 \cdot \delta WD^2 + \left(\frac{\partial AGB}{\partial BEF}\right)^2 \cdot \delta BEF^2 + \left(\frac{\partial AGB}{\partial GSV}\right)^2 \cdot \delta GSV^2} \quad (13)$$

395 The standard deviation of the wood density estimates, δWD , was obtained by computing the variance of the predictions for each measurement of wood density and fitting a linear model (Fig. S3S4). The standard deviation of the BEF was expressed in terms of partial derivatives of its components

$$\delta BEF = \sqrt{\left(\frac{\partial BEF}{\partial p_1}\right)^2 \cdot \delta p_1^2 + \left(\frac{\partial BEF}{\partial p_2}\right)^2 \cdot \delta p_2^2 + \left(\frac{\partial BEF}{\partial SB}\right)^2 \cdot \delta SB^2} \quad (14)$$

400

In Eq. (14), δp_1 and δp_2 represent the standard deviations of the two coefficients of the BEF model.

2.4 Validation

To assess the accuracy of the AGB estimates, we used a reference dataset of AGB observations from 110,897 forest field inventory plots. The data were gathered from a variety of surveys undertaken by national forest inventories and research networks (Supplement Section A.6).

405 The opportunistic nature of our validation database led to an uneven spatial distribution of the reference samples (Fig. S4S5), as well as a variety of plot sizes, survey methods and allometric equations used (Table S3 and Table S4). The plots were mostly smaller than 1 ha (Table S3), implying that they often represented a small fraction of the area covered by a 1-ha map pixel. To 410 reduce the effect of random errors caused by the mismatch in resolution between the reference dataset and our map model, we aggregated our map and the plot data to 0.1° grid cells. This represented as a trade-off between capturing the local scale variability of AGB whilst allowing a number of plot measurements deemed sufficient to compute an average AGB representative of the area within the grid cell. In the end, our assessment cannot provide an indication of the validity of pixel-

based AGB estimates. Instead, it provides a measure of the accuracy of generalized spatial AGB patterns. This is, however, a
415 pragmatic approach when using measurements not designed for validation of estimates from remote sensing imagery but which often provide the only source of information on AGB in poorly inventoried regions or regions where national inventory data are not publicly available.

To provide a more comprehensive overview of the reliability of the spatial patterns in areas not covered by the database of plot
420 inventory measurements, the analysis was supplemented by a comparison of average GSV or AGB at the level of inventory reference units (polygons, counties, provinces and ecoregions). The scope of this analysis was primarily to identify systematic errors, on a large scale, that may not become evident when comparing at individual plot level. For a quantitative assessment of the retrieval at the scale of provincial and regional aggregates, we computed the RMSD between map and reference biomass averages relative to the average reference biomass and the bias between map and reference biomass averages. The RMSD was
425 computed as a weighted mean of the errors, where the weights corresponded to the ratio of the forest area to the total forest area. For Russia (Supplement Section A.6), GSV and AGB data were gathered for approximately 1,600 Forest Managements Units (FMU) ranging in size from 3,000 ha (e.g. intensive forestry or national parks in the European part) to 30,000,000 ha (remote territories in Siberia). For countries with a well-established national forest inventory that regularly publish regional statistics of forest biomass at the level of administrative or ecological units, we assembled a database of GSV and AGB
430 averages representative of the epoch 2010 (Supplement Section A.6).

2.5 Inter-comparison of AGB maps

The spatial distribution of AGB map estimates from our dataset were compared with biome and global AGB maps based on satellite remote sensing observations (Avitabile et al., 2016; Baccini et al., 2012; Liu et al., 2015; Saatchi et al., 2011b; Santoro et al., 2015a; Thurner et al., 2014) or ancillary datasets. The spatial distribution of AGB map estimates from our dataset were
435 compared with biome and global forest AGB maps based on satellite remote sensing observations (Saatchi et al., 2011b; Baccini et al., 2012; Thurner et al., 2014; Liu et al., 2015; Avitabile et al., 2016; Santoro et al., 2015a) or ancillary datasets (Kindermann et al., 2008). To do this, the datasets were first harmonized to a common geographic map projection, resampled to a pixel size of 0.01° and converted, when necessary, from AGB carbon units ($MgC\ ha^{-1}$) to AGB (Table S5). The datasets span a decade of input observations from 2000 to 2010 but could not be harmonized to reflect the conditions of a single epoch
440 due to the lack of information on growth rates. This, however, is not expected to have significant effects on the interpretation of the latitudinal profiles.

As a way of assessing the AGB patterns of each map, we compared latitudinal averages based on values from the 0.1° validation grid cells with corresponding values from the database of forest inventory plots. To obtain a homogeneous representation of
445 all latitudes, we grouped grid cells in 10° wide intervals.

3. Results

3.1 Global AGB dataset

The global AGB dataset (Fig. 2) was obtained by scaling global estimates of GSV (Fig. S5). While an additional assessment of our dataset against regional maps of AGB would further contribute to build confidence in the data product, it is felt that such

450 investigations require an own framework also making use of local reference data and expert knowledge as for example undertaken for Europe (Avitabile and Camia, 2018), the United States (Spawn et al., 2020) and Tanzania (Næsset et al., 2020). To do this, the datasets were first harmonized to a common geographic map projection and resampled to a pixel size of 0.01°. Datasets expressing AGB in carbon units (MgC ha^{-1}) were converted to AGB using a carbon fraction default value of 0.47 (IPCC, 2006) (Table S5). All datasets span a decade of input observations from 2000 to 2010 but could not be harmonized to 455 reflect the conditions of a single epoch due to the lack of information on growth rates. This, however, is not expected to have significant effects on the interpretation of the latitudinal profiles.




As a way of assessing the AGB patterns of each map, we compared latitudinal averages based on values from the 0.1° validation grid cells with corresponding values from the database of forest inventory plots. To obtain a homogeneous representation of


460 all latitudes, we grouped grid cells in 10° wide intervals.

3. Results

3.1 Global AGB dataset

The global AGB dataset (Fig. 2) was obtained by scaling global estimates of GSV (Fig. S6) using model-based estimates of wood density (Fig. S6S7) and stem-to-total biomass expansion factors (Fig. S7S8). The uncertainty of the AGB estimates is 465 reported as standard deviation (Fig. 2). The models for retrieving GSV and converting it to AGB were developed for all-woody vegetation, so we evaluated the estimates corresponding just to forest cover by regrouping the classes from the Climate Change Initiative Land Cover (CCI-LC) dataset of 2010 into forest and non-forest land (Supplement Section A.1, Table S6S7).





470

Figure 2. Map estimates of AGB (top panel) and AGB standard deviation expressed relative to the AGB (bottom panel). The colour bar of the AGB map has been truncated at 500 Mg ha⁻¹ to increase contrast. Similarly, the colour bar of the AGB relative standard

475 deviation has been truncated at 100%. The right hand panel shows the profile of average AGB along latitude (thick solid line) and the two-sided average standard deviation of AGB at a given latitude (horizontal bars).

At the 1-ha scale, the largest predicted value was 757 Mg ha⁻¹, corresponding to a GSV of 1,087 m³ ha⁻¹, in forests of the U.S. Pacific Northwest. However, for 99% of the world's forests, AGB was estimated to be less than 360 Mg ha⁻¹, and 90% was below 182 Mg ha⁻¹, which explains truncating the colour bar in Fig. 2 at 500 Mg ha⁻¹. The spatial distribution of AGB followed a clear latitudinal gradient (Fig. 2). In the northern hemisphere, AGB increased steadily with decreasing latitude across the boreal zone (between 75°N and 60°N), then remained fairly constant throughout the temperate (between 60°N and 40°N) and sub-tropical (between 40°N and 20°N) zones. AGB increased sharply as we enter and leave the tropical zone between 20°N and 20°S, though with a minimum at 13°N due to the large area of low biomass dry forests in the sub-Saharan region. The AGB of semi-tropical forests in the southern hemisphere between 20°S and 33°S was slightly lower than in the corresponding 485 latitude range in the northern hemisphere because of the larger proportion of low-density forest. The local maximum at 25°S


485


corresponded to the Atlantic forests of Brazil and dense sub-tropical forests along the east coast of Australia where biomass accumulation is favoured by higher precipitation. Temperate forests had higher AGB in the southern hemisphere (south of 33°S) than in the northern hemisphere because of the predominant highly productive evergreen and coniferous forest along the Chilean-Argentinean Andes, in south-eastern Australia and New Zealand. The peak at 42°S corresponds to the broadleaved

490 forests of Tasmania.

The AGB standard deviation, expressed in Fig. 2 relative to the AGB estimates, was on average 50% with an inter-quartile range of values between 44% and 61%. The rather constant relative uncertainty is also illustrated by the horizontal bars in the latitudinal profile of Fig. 2, which scale with the AGB level. The relative standard deviation was smaller than 100% for approximately 95% of the mapped pixels which explains truncating the colour bar of the AGB standard deviation map in Fig. 495 2 at 100%. The large majority of the AGB estimates for which the standard deviation exceeded the 100% level were below 20 Mg ha⁻¹ such as in sparsely vegetated regions corresponding to the transition to tundra in Canada and Alaska, the Siberian Lowlands and Far East Russia or in poorly stocked forests such as in northern China and west Madagascar.


500 The small-scale variability of AGB in forest landscapes is captured by the 1-ha pixel spacing of the dataset (Fig. 3 and Fig. 4). The example in Fig. 3 shows the region of the Bratsk Reservoir formed by the Angara River in Central Siberia, where forests are heavily managed for timber production. Clear-felling activities occur in polygons often larger than 10 ha. The map for a 1° × 1° area east of the reservoir shows forests with AGB above 100 Mg ha⁻¹, roughly corresponding to a GSV of at least 200 m³ ha⁻¹, interspersed with small rectangular white shapes corresponding to clear-cut areas (Fig. 3). Our estimates give more 505 detailed information on the spatial patterns of AGB in this region than the AGB product by Thurner et al. (Thurner et al., 2014) based on remote sensing data with a spatial resolution of 1,000 m.






510 **Figure 3.** AGB estimates from this study (left) and from Thurner et al. (Thurner et al., 2014) for a $1^\circ \times 1^\circ$ area in Central Siberia.

The example in Fig. 4 includes part of Trans-Amazonian Highway south of the Amazon River in the state of Pará, Brazil. This region is characterized by fishbone deforestation caused by lateral expansion of agriculture from the highway into pristine forest. The fishbone pattern is clear in our AGB map (Fig. 4, top left panel) and some isolated, small-scale patches of deforestation are also visible. For comparison, Fig. 4 shows AGB estimates by three pan-tropical maps based on remote sensing observations (Avitabile et al., 2016; Baccini et al., 2012; Saatchi et al., 2011b).
 515 Fig. 4 shows AGB estimates by three pan-tropical maps based on remote sensing observations (Saatchi et al., 2011b; Baccini et al., 2012; Avitabile et al., 2016). The deforestation patterns are less extended in the map by Saatchi et al. (2011b) because it was based on satellite data acquired around the year 2000, compared to the maps by Baccini et al. (2012), based on observations taken in 2007, and Avitabile et al. (2016), which
 520 merged the other two pan-tropical datasets. The level of detail in our map is much greater because of the 1-ha spatial resolution compared to 25 ha in Baccini et al. and nearly 100 ha in Avitabile et al. and Saatchi et al.






525 **Figure 4.** AGB estimates by this study (top left), Avitabile et al. (2016) (top right), Saatchi et al. (2011b) (bottom left) and Baccini et al. (2012) (bottom right) for a $1^\circ \times 1^\circ$ area in the state of Pará, Brazil.

3.2 Validity of AGB estimates

AGB averages were obtained for 6,456 0.1° grid cells including at least five inventory plots (Supplement Section A.6). These grid cells represented approximately 1% of the Earth's forest cover. The grid cell average AGB from the field inventory database ranged from 0 to 1,670 Mg ha^{-1} (median: 43 Mg ha^{-1} , mean: 60 Mg ha^{-1} , 99th percentile: 351 Mg ha^{-1}). The AGB histogram was skewed towards low values (Fig. 5a) because of the large proportion of measurements from the National Forest Inventories of Spain and Sweden (Table S4). The grid cell average AGB from the map ranged from 0 to 358 Mg ha^{-1} (median: 57 Mg ha^{-1} , mean: 67 Mg ha^{-1} , 99th percentile: 278 Mg ha^{-1}). The AGB histograms from the map (Fig. 5b) and the field inventory (Fig. 5a) were similar, with a mode at around 0, a decline to a shoulder, then a further decline to a long tail. In the field inventory the shoulder covered the range 25–50 Mg ha^{-1} and was followed by a slow decline, but for the map it extended to around 80 Mg ha^{-1} and then declined rapidly. This difference arises because the map tends to give higher values than inventory in the lower AGB range (Fig. 5c and Fig. S8S9). These trends have been reported for other pan-tropical and regional AGB studies (Avitabile et al., 2016; Rodriguez-Veiga et al., 2019)(Avitabile et al., 2016; Rodriguez-Veiga et al., 2019). The scatterplot of map against inventory values of AGB in Fig. 5c and the RMSD curve in Fig. S8S9 indicate an agreement in

540 trend between field inventory and map values up to about 250 Mg ha^{-1} . Above 250 Mg ha^{-1} , the map values rose as the field inventory value did, albeit more gently and with much greater variance. Disaggregating the data by major ecological domains, using the FAO Global Ecological Zones as reference, suggested slight differences in the agreement between map and inventory values in tropical, sub-tropical and temperate forests (Fig.s 5d-f).



545




Figure 5. Histograms of AGB from the field inventory database (a) and the map (b) for 0.1° grid cell values. (c) Scatterplot of map AGB against field inventory values for 0.1° grid cells (grey circles); the filled circles show the median AGB of the map values in each 10 Mg ha^{-1} wide interval of field inventory AGB values. The colour bar represents the number of grid cells within a given AGB interval. Similar scatterplots are given for the tropical zone (d), the sub-tropical zone (e), the temperate zone (f) and the boreal and polar zones (g) according to the FAO Global Ecological Zones. On each scatterplot, we report the root mean squared difference (RMSE) between map and field inventory AGB relative to the mean value of the reference AGB and the bias, i.e., the difference between mean values of the map AGB and the reference AGB. To improve presentation and because of the paucity of grid cells with AGB above 500 Mg ha^{-1} , axes are truncated at 500 Mg ha^{-1} .

550

555 3.3 Spatial distribution of AGB

Combining our AGB dataset with the CCI Land Cover dataset and the FAO ecological zones, we estimated a total AGB of 521 Pg for a forest area of 4,825 Million ha, corresponding to a global average forest AGB of 108 Mg ha^{-1} (Table 1). Using a carbon fraction default value of 0.47 (IPCC, 2006) we estimated a total above-ground carbon stock of 246 PgC. Tropical forests had the highest average AGB (147 Mg ha^{-1}), representing 64% of the total forest AGB and 47% of the total forest area. The second largest average AGB was found in temperate forests (102 Mg ha^{-1}), which accounted for 14% of the total AGB and 15% of the total forest area. Sub-tropical and boreal forests had similar average AGB (75 and 60 Mg ha^{-1} , respectively), but the area covered by the latter was almost three times larger. As a result, the total AGB of boreal forests was more than twice as large as that of sub-tropical forests and corresponded to 16% of the total AGB, thus being larger than the AGB pool in temperate forests. The total AGB of subtropical forests accounted for 560 7% of the global AGB. The contribution of polar forests to the global AGB pool was negligible (0.1%).

The highest values and range of biomass were found in looking at Fig. 6, the tropical rainforestsrainforest (TAr) ecozone hosted primarily high-density forests, with a median AGB of 238 Mg ha⁻¹ (Fig. 6) and the low-end of the interquartile range above 150 Mg ha⁻¹. Besides tropical rainforests, only the tropical mountain, temperate oceanic and temperate mountain ecozones (TM, TeDo and TeM, respectively) had median AGB value above 100 Mg ha⁻¹ (Fig. 6). For the tropical, sub-tropical and temperate ecozones, the AGB of forests in dry environments (shrubland, steppe and desert) was lower than in wet environments (rainforest, moist, humid, mountain, continental and oceanic). The AGB of boreal forests decreased with increasing latitude from the boreal coniferous (Ba) ecozone located at the southernmost edge of the boreal ecotone, through the boreal mountain (BM) ecozone, to the boreal tundra woodland (Bb) ecozone at the northernmost edge of the boreal zone.

575 The AGB standard deviation, here expressed relative to the AGB estimates, was fairly constant across most ecological zones (Fig. 6). The median value ranged between 44% and 57%, except for the tropical shrubland, tropical desert and polar ecozones (TBSH = 69%; TBWh = 92% and P = 84 %, respectively). Driven by the decreasing sensitivity of the radar backscatter to increasing GSV, the largest proportion of the AGB standard deviation was attributed to the uncertainty of the GSV estimates (Fig. S9), an effect further exacerbated in wet environments (Santoro et al., 2015a); S10. The uncertainty of the wood density estimates accounted for 7% to 20% (mean: 14%) of the AGB standard deviation (Fig. S9; S10), while the uncertainty in the BEF accounted for between 2% to 13% of the AGB standard deviation (mean: 7%) with the exception of the tropical desert zone (30%) (Fig. S9; S10). The uncertainty of the GSV was driven by the weak sensitivity of the radar backscatter to increasing GSV, an effect further exacerbated in wet environments (Santoro et al., 2015a), thus explaining the slightly higher uncertainty in the tropics and the sub-tropics. The larger uncertainty in sparsely forested regions compared to densely forested regions (Fig. 2) is a consequence of the substantially larger uncertainty of the σ_{gr}^0 parameter compared to the uncertainty of the σ_{veg}^0 in the GSV retrieval model in Eq. (1) (Fig. S11). In sparse forests or forests with low woody biomass stocks, the total backscatter from the forest is dominated by the ground scattering component, i.e., the term with σ_{gr}^0 in Eq. (1), thus being affected by larger uncertainty compared to the backscatter received from dense forests, for which the largest contribution to the measured backscatter is attributed to the scattering from the canopy, i.e. to the term with σ_{veg}^0 in Eq. (1).

595

Table 1. Total AGB, forest area and average AGB per major ecozone.

Ecozone

← Formatierte Tabelle

	Total AGB (Pg)	Forest area (10^6 ha)	Average AGB ($Mg\ ha^{-1}$)
Tropical	331.3	2251.7	147
Sub-tropical	36.2	483.0	75
Temperate	71.6	698.3	102
Boreal	81.2	1352.5	60
Polar	0.6	39.8	18
Total	521.0	4825.4	108

600




Figure 6. Box plot diagram of AGB (top) and its standard deviation by FAO global ecological zone. On each box, the central mark indicates the median, the bottom and top edges delimit the interquartile range, and the whisker delimits the 1-99 percentile range. Labels of global ecological zones: TAR: Tropical rain forest; TAwA: Tropical moist deciduous forest; TAwB: Tropical dry forest; TBSH: Tropical shrubland; TBWh: Tropical desert; TM: Tropical mountain systems; SCf: Subtropical humid forest; SCs: Subtropical dry forest; SBSH: Subtropical steppe; SBWh: Subtropical desert; SM: Subtropical mountain systems; TeDo: Temperate oceanic forest; TeDc: Temperate continental forest; TeBSk: Temperate steppe; TeBWk: Temperate desert; TeM: Temperate mountain systems; Ba: Boreal coniferous forest; Bb: Boreal tundra woodland; BM: Boreal mountain systems; P: Polar.

605

610 **3.4 Assessment of global forest biomass resources**

The most comprehensive summary of global forest resources and biomass pools is reported by the FAO in their quinquennial FRA. For the FRA, each country reports its values of forest area ~~and (in hectares)~~, total AGB (in Pg) and average AGB (in Mg ha⁻¹) according to their inventory capabilities. Forest area ~~is was~~ derived from inventory data or remote sensing data. AGB statistics ~~are were~~ derived either from measurements collected as part of national inventories, local inventories or estimates 615 reported in the literature. Adjustments ~~are were~~ applied by the FAO where necessary to ensure consistency with its own information sources on forest area and biomass resources.

While all 233 countries contributing to the FRA for 2010 reported their estimates of forest area, only 171 reported total AGB-
620 and average AGB. Of the remaining 62 countries, accounting for approximately 2.6% of the global land surface, 10 reported no forest cover, while 7 (Fiji, Eritrea, Uruguay, Ecuador, Paraguay, Japan and Venezuela) reported an estimate of forest area exceeding 10⁶ ha. Of the countries reporting on AGB, 111 derived their average AGB from values of the average GSV using one or multiple BCEF values. This was common practice for countries lacking a systematic national forest inventory. Among these, 79 relied on default numbers published by the Intergovernmental Panel on Climate Change (IPCC) (IPCC, 2006) (Table S7).

625

3.4.1 Global and continental statistics

The statistics on forest area, average AGB and total AGB from the FRA 2010 and from the combination of our AGB dataset with the CCI Land Cover dataset are reported by continent (Table 2). Our total AGB estimate of 522 Pg was 11% higher than the value of 468 Pg reported by the FRA. This difference is a consequence of the 23% larger forest area estimated from the 630 CCI Land Cover dataset (Table 2) despite the FRA reporting 9% higher global average AGB than our estimate (119 Mg ha⁻¹ vs. 108 Mg ha⁻¹, Table 2). Compared to Table 1 based on the FAO Global Ecological Zones dataset to delineate land surfaces, we estimate an additional 8 10⁶ ha of forest area and an additional 1.5 Pg of total AGB as a consequence of the more precise delineation of the land surface by the Database of Global Administrative Areas used as reference for the countries boundaries (GADM, <http://www.gadm.org>).

635

For the three continents spanning the tropics, we found the highest average AGB in South America, although our estimate (183 Mg ha⁻¹) was 11% lower than the corresponding value from the FRA (206 Mg ha⁻¹) (Table 2). South America also contained the largest total AGB pool. Although our estimate (155.9 Pg) was only 4% smaller than the FRA (162.3 Pg), the FRA did not provide AGB for approximately 10% of the forest area of South America (Table 2). We found a larger difference 640 between our results and those of FAO for Africa. The average AGB from our dataset was 24% lower than the FRA (108 Mg ha⁻¹ vs. 142 Mg ha⁻¹, Table 2) whilst the total AGB was only 11% smaller than in the FRA (84.8 Pg vs. 95.3 Pg) due to the

larger forest area we used from the CCI Land Cover dataset. In contrast, for Asia the average AGB from our dataset (115 Mg ha^{-1}) was 17% higher than in the FRA (98 Mg ha^{-1}) (Table 2) while our estimate of total AGB exceeded that from the FRA by 64% (89.4 Pg vs. 54.3 Pg) mainly because of the 40% larger forest area estimated from the CCI Land Cover dataset.

645

For the two continents spanning the northern boreal and temperate zones, the average AGBs from our estimates were well below 100 Mg ha^{-1} (Table 2). The average AGB estimate for Europe differed by less than 1% (Table 2) compared to the FRA whereas for North/Central America the difference was 15%. Because the forest area estimated from the CCI Land Cover dataset was larger than the FRA values, the total AGB estimated from our dataset was larger than the values reported in the 650 FRA by 20% for Europe and 14% for North/Central America.

Finally, the smallest of the continental pools of AGB was in Oceania, where our larger estimate of total AGB compared to the FRA was primarily explained by our estimate of AGB being on average almost 35% larger than the value derived from the FRA.

655

Table 2. Total AGB (unit: Pg), forest area (unit: ha) and average AGB (unit: Mg ha^{-1}) per continent from this study and from the FAO FRA 2010. The forest area column for the FRA does not account for countries reporting forest area but not AGB. For Asia, North/Central America and South America, 5%, 1% and 10% of the forest area did not contribute to the AGB to the FRA. For Africa, Europe and Russia, and Oceania, less than 1% of the forest area did not contribute to the FRA. The total forest area from 660 the FRA is $4.033 \cdot 10^6 \text{ ha}$.

Continent	This study			FAO FRA 2010		
	Total AGB (Pg)	Forest area (10^6 ha)	AGB (Mg ha^{-1})	Total AGB (Pg)	Forest area (10^6 ha)	AGB (Mg ha^{-1})
Africa	84.8	783.5	108	95.3	672.6	142
Asia	89.4	780.5	115	54.3	554.0	98
Europe and Russia	91.2	1268.3	72	73.7	1016.5	72
North and Central America	77.9	970.6	80	66.8	704.4	95
South America	155.9	850.0	183	162.3	788.9	206
Oceania	23.3	180.4	129	16.0	189.7	85
Total	522.5	4,833.4	108	468.5	3,926.1	119

← Formatierte Tabelle

3.4.2 National statistics

At the level of individual countries, the agreement between the total and average AGBs from our dataset and from the FRA differed depending on the continent (Fig. 7 and Fig. 8, Table S7). The largest difference between our countryaverage AGB and FRA AGB was in Africa (median difference: -60%, 51 countries), where for most countries the average AGB reported in the FRA exceeded the value from our dataset (Fig. 7). This is probably due to either AGB underestimation in countries dominated by high-density forests or to high biomass conversion and expansion factors (BCEF) used by countries with low-density forest when estimating values of AGB from their original measurements of GSV (Table S7). In addition, several countries used a small sample of plots for the calculations, as well as small-sized plots for heterogeneous forest areas, leading to large uncertainties in the values reported to the FRA. In Europe and South America, where the span of the countryaverage AGB was similar to Africa, we also saw underestimation patterns above 200 Mg ha⁻¹ (Fig. 7). Nonetheless, we did not seeidentify the trend in the range 0 - 200 Mg ha⁻¹ seen in Africa. For European and South American countries, the assessment of the reliability of our AGB estimatesaverages at country level was more meaningful because of the better developed national inventories (Fig. 8b) and the direct estimation of AGB from the inventory measurements, thus bypassing the use of a standard BCEF like in African countries (Table S7). Indeed, the smallest differences between the average AGB from the FRA and from our dataset were obtained in Europe (median difference: -8%, 42 countries) and South America (median difference: 2%, 9 countries) (Fig. 8a).

For North/Central America (19 countries), Asia (35 countries) and Oceania (9 countries), the median difference between our countryaverage AGB estimates and the FRA valuesnumbers was between -23% and -27% (Fig. 7, Table S7). The disagreement for North/Central America was largest for the Caribbean countries (Fig. 8a), for most of which the reference data used in the FRA havehad low to moderate quality and the NFI capacity was mostly low or limited.


The scatterplot for Asian countries shows data points of average AGB clustered AGB-estimates along the identity line (Fig. 7) but with two distinct regions. For the Asian Middle East stretching as far as Pakistan and the former Soviet countries of the Asian continent, the countryaverage AGB from our dataset was on average 70% smaller than the values reported in the FRA (Fig. 8a). The FRA country reports were based on highly stocked forest, which may not be representative of the true distribution ofaverage AGB. In contrast, the countryaverage AGB estimated from our dataset was on-averageapproximately 27% larger than the values reported in the FRA for the southern and eastern regions of the Asian continent (Fig. 8a). Several countries of Southeast Asia assumed their forests to be strongly degraded which justified the use of low countryreference values for the average GSV, and hence AGB, when reporting to ecompute theirthe FRA statistics.

Our estimates of average AGB for Australia (brown marker, Fig. 7, panel Oceania) and Papua New Guinea (green marker, Fig. 7, panel Oceania) exceed the values in the FRA, while being smaller for New Zealand (cyan marker, Fig. 7, panel Oceania).

Australia reported their biomass stock based on models calibrated with a small number of inventory measurements. For Papua New Guinea the FRA AGB was based on commercial volume for trees with a diameter at breast height of at least 50 cm, thus being a fraction of the true AGB. For New Zealand, the result was comparable to those obtained for European countries with large AGB, which are characterized by similar forest types and structures.

700

In this assessment, the impact of a country's NFI capacity (Romijn et al., 2015) on the quality of the values reported to the FRA is illustrated by the statistical parameters reported in Fig. 8 (panels c, d and e). The agreement between country AGB computed from our map and reported in the FRA increased with capacity level (correlation coefficient between 0.46 and 0.91, relative RMSD between 30% and 74%, mean difference between -42% and -12%). Several countries with mean AGB above 705 200 Mg ha⁻¹ and thus in the AGB range prone to underestimation (Fig. 5) were associated with an intermediate NFI capacity level (Fig. 8b), which partly explains the somewhat poor agreement between our estimatesaverage AGB and the average AGB reported in the FRA (Fig.s 8c, 8d and 8e).



Formatierte Tabelle

Forest area on continent

710 Figure 7. **CountryAverage AGB_per country** from the FAO FRA 2010 country reports and our map dataset. Countries have been grouped per continent. The size of each circle is proportional to the forest area of the country derived from the CCI Land Cover dataset. **For each continent, the (same scaling across all continents).** The colour ramp associated to the circles gives a graphical representation of the **min-max range of relative country forest areasarea** (different for each continent).





Figure 8. Difference between country AGB from our map and FRA, expressed relative to the FRA AGB (a), NFI capacities for the year 2010 (Romijn et al., 2015) (b), Pearson's correlation coefficient (c), relative RMSD (d) and mean difference (e) between map


estimates of country AGB and values from the FRA for NFI capacity level. Number of countries per NFI capacity level: 51 (low), 30 (limited), 26 (intermediate), 37 (good) and 23 (very good).

720

In terms of forest area, the orders of magnitude in the map and the FRA agreed but in most cases the area obtained from the CCI Land Cover dataset was larger than the country values reported to the FRA. The CCI value was on average 24% greater than in the FRA except for the countries of North/Central America, where the average difference was 81%. The discrepancy can be explained in terms of the different definitions of forest used in CCI when generating land cover maps and in the FRA 725 when using national data of different quality to generate the country estimates. However, the estimate of forest area obtained in this study is likely to be an underestimate, since it excluded land cover classes with a sparse tree and vegetation component that could be attributed to forest under less restrictive definitions of percentage tree cover (Mermoz et al., 2018).

As a result, our country estimates of total AGB were only slightly different from the FRA estimates (Fig. 9) because the lower 730 AGB densities from our dataset were compensated by the larger forest area values in the CCI Land Cover dataset (Table S7). For Asian and European countries, the difference was 7% and 5% on average, respectively. The difference was larger throughout the American continent where total AGB estimated in this study exceeded that in the FRA on average by 33% (North/Central America) and 23% (South America). Only for Africa and Oceania did we obtain a total AGB smaller than the 735 FRA, on average -37% for Africa and -29% for Oceania, because of the large difference between [country average](#) AGB values obtained in this study and [those](#) reported by FAO.





740 **Figure 9.** Total AGB from the FAO FRA 2010 country reports and from our dataset (see Table S7 for details).

3.5 Comparison of map estimates of AGB

All map datasets analysed in this study including ours (Table 3) showed similar latitudinal trends; (Fig. 10a), with the highest AGB in the southern hemisphere at 40°S and across the Equator and low AGB in the dry tropics at around 20°S and north of 745 10°N (Fig. 10a). However, there were some large differences between the maps (Fig. 10a), indicating illustrating the current uncertainty about the global forest carbon pools.

For the wet tropics between 15°S and 10°N, the AGB from our dataset was close to that from the pan-tropical dataset by Saatchi et al. (Fig. 10b) but is between 10% and 50% lower than the values by Baccini et al. and the GEOCARBON dataset.

750 The analysis of the latitudinal averages based on the 0.1° grid cell AGBs show an overestimate of AGB in these two datasets by about 10% to 60%. On the contrary, the average AGBs from our dataset were only by a few percent values higher than the corresponding values from the plot inventory database (Fig. 11a and 11b).

The differences between our dataset and each of the pan-tropical AGB datasets were much larger in tropical and sub-tropical regions north and south of the wet tropics. In the northern hemisphere, between 10°N and 30°N, Saatchi et al. and Baccini et 755 al. exceeded our estimates by between 40% and 110% (Fig. 9b). For the same latitude range, the GEOCARBON dataset differed from our values by 110% to -47% with increasing latitude (Fig. 10b). The latitudinal averages at the 0.1° grid cells

confirmed that all three maps were strongly biased while our dataset was closer to the values obtained from the plot inventory dataset (Fig. 11). The GEOCARBON dataset presented both positive and negative biases, being about 50% of the average AGB from the plot inventory dataset (Fig. 11a). The overestimate exceeded 100% of the average AGB from the plot inventory dataset in the case of Saatchi et al. and Baccini et al. (Fig. 11b). In the southern hemisphere, between 35°S and 15°S, the GEOCARBON dataset and Baccini et al. exceeded our AGB by 20% to 160%, whereas Saatchi et al. differed from our values by between +20% and -60% (Fig. 10b). The comparison at the level of the 0.1° grid cells indicates an overestimate by GEOCARBON and Baccini et al. between 20% and 60% of the average AGB from the plot inventory data (Fig.s 11a and 11b, respectively). The difference between the average AGB from our dataset and from Saatchi et al. with respect to values from the plot inventory dataset were instead mostly below 20%.

For the southern hemisphere temperate forests south of 35°S, our AGB exceeded the values from Saatchi et al. and the GEOCARBON dataset. The comparison was, however, of limited value because of the incomplete coverage by the former and the coarse resolution estimates used to fill gaps in the latter.

North of 30°N, we did not observe differences between our dataset, Thurner et al. and the GEOCARBON dataset. This was expected since all studies were based on the same Envisat ASAR dataset and the BIOMASAR algorithm. The only difference was visible at about 70°N (Fig. 9b and 6c) because of the tendency of ASAR to overestimate biomass in sparse tree vegetation.

This difference is, however, of minor importance in the context of global estimation of AGB stocks because ~~AGB hardly ever exceeded 30 Mg ha⁻¹ at these latitudes (Fig. 10a). Fig. 10a shows the AGB hardly ever exceeded 30 Mg ha⁻¹ at these latitudes in any of the datasets (Fig. 10a), this being consistent with values based on forest inventory measurements (Stolbovoi and Mc Callum, 2002; Gillis et al., 2005).~~ Figs 11a and 11c confirm that the three maps present similar spatial patterns of AGB; the difference with respect to the plot inventory dataset was about 20%-30% for all datasets except for GEOCARBON at 60°N where the difference was 60% of the plot inventory value. These results were attributed to the higher spatial resolution of our map. The coarse resolution AGB datasets by Liu et al. and Kindermann et al. agreed with our estimates in the large unbroken tracts of forest in the wet tropics between 10°S and 10°N and in the boreal zone around 60°N (Fig. 10d). However, the estimates of AGB by Liu et al. were up to 70% lower than our estimates in fragmented forest landscapes, e.g. between 50°S and 20°S and between 20°N and 50°N (Fig. 10d). The dataset by Kindermann et al. was better correlated with our estimates, the difference rarely exceeding 30% in absolute terms.

Table 3. Total AGB (Pg) for five latitude ranges roughly corresponding to the temperate forests of the southern hemisphere (60°S - 30°S), the humid and dry tropics of the southern hemisphere (30°S - 10°S), the wet tropics (10°S - 10°N), the humid and dry tropics of the northern hemisphere (10°N - 30°N), and the temperate and boreal forests of the northern hemisphere (30°N and 90°N). Values marked with superscript 1 indicate partial coverage of the latitude range by the corresponding map; n/a indicates not available in the latitude range. The forest area for each range is reported on the last line.


Source	Latitude range
--------	----------------

← Formatierte Tabelle


	60°S - 30°S	30°S - 10°S	10°S - 10°N	10°N - 30°N	30°N - 90°N
This study	12.5	49.8	253.9	37.3	160.8
(Saatchi et al., 2011a)	4.0 ¹	51.8	253.7	69.0 ¹	13.1 ¹
(Baccini et al., 2012)	n/a	55.8 ¹	278.8	39.4 ¹	n/a
GEOCARBON (Abitabile et al., 2016) for the tropics, (Santoro et al., 2015a) (Santoro et al., 2015a) for the northern hemisphere	10.3	39.9	248.0	19.3	108.2
(Thurner et al., 2014)	n/a	n/a	n/a	n/a	114.7
(Liu et al., 2015)	9.6	58.6	278.6	53.1	169.3
(Kindermann et al., 2008)	13.1	74.8	242.8	48.1	175.8
Forest area (10 ⁶ ha)	109.3	584.0	1366.1	497.8	2294.7

All AGB datasets yielded similar values of total AGB in the wet tropics between 10°S and 10°N (Table 3) with our estimate being intermediate to the others. For the latitude ranges corresponding to the humid and dry tropics, our total AGB was much lower than Saatchi et al. and Baccini et al. (87.1 Pg vs. 120.8 Pg and 95.2 Pg, respectively) because of their substantially higher estimates of AGB outside the rainforest region. It was also different from Liu et al. and Kindermann et al. (87.1 Pg vs. 111.7 Pg and 122.9 Pg, respectively) because the coarse spatial resolution of the data did not allow forest fragmentation to be accounted for. For boreal and temperate forests (Table 3, column: 30°N - 90°N) the total AGB from our dataset was larger than in Thurner et al. and the GEOCARBON dataset (160.8 Pg vs. 114.7 Pg and 108.2 Pg, respectively). In both these datasets, a more stringent definition of forest than the CCI Land Cover dataset was used resulting in a large number of AGB estimates equal to 0 Mg ha⁻¹ in areas labelled as forest by the CCI Land Cover dataset. Our estimate of total AGB was slightly lower than that from Liu et al. and Kindermann et al. because these datasets did not account for forest fragmentation. The values were difficult to compare for the latitude range including the temperate forests of the southern hemisphere (Table 3, column: 60°S - 30°S) because of the approximations underlying the AGB values obtained in the other maps (extrapolation from coarse

805 resolution estimates in GEOCARBON, statistics of large countries in Kindermann et al. and use of inversion models trained
in the tropics in Liu et al.).



810 **Figure 10.** Profiles of average AGB along latitude for each dataset listed in Table 3 (a), together with the corresponding relative
difference between the latitudinal AGB derived from the other datasets and this study (b, c and d), expressed in percentage values
relative to our estimates.






Figure 11. Relative difference of AGB from maps and plot inventory data averaged over the 0.1° grid cells validation cells (a, b and c). The number of grid cells stratified by 10° -wide latitude intervals is illustrated in (d).

4 Discussion and conclusions

Overall, our maps reproduced known spatial patterns of AGB (Fig. 5, Fig. S10 and Fig. S11) and GSV (Fig. S12) correctly, although with some systematic errors (Fig. S10, Table S8). The results indicate that our maps reproduced known spatial patterns of AGB (Fig. 5, Fig. S12 and Fig. S13) and GSV (Fig. S14) correctly, although with some systematic errors (Fig. S12, Table S8). Independent assessments confirmed these indications (Spawn et al., 2020; Schepaschenko et al., 2021). The dispersion of the AGB estimates about the identity line in Fig. 5 was often explained by regional biases (Fig. S10-S12) arising from approximations in the retrieval model and the use of remote sensing data with different spatial resolutions (Table S8). Hence, the global RMSD and bias statistics in Fig. 5c had limited informative value when trying to explain our results. Instead, values

at the level of continents or domains provided a more realistic indication of the quality of the AGB dataset since we could
825 identify a predominant error source in each (Table S8).

Interpreting Table S8, errors were partly a consequence of a simple model relating SAR observations to biomass through three
830 unknown parameters, whose estimates relied on several assumptions and generalization rules. In addition, to avoid unrealistic
model fits when relying on external data for model training, our model training procedure was forced to follow the average
relationship between SAR backscatter and biomass. This reduced the variability of the predicted backscatter with respect to
the measured backscatter, which translated to overestimation in the low AGB range and underestimation in the high AGB
range, as evidenced by Fig. 5. In spite of these weaknesses, we preferred such an approach to the traditional model training
based on a dataset of reference biomass measurements because of the paucity of reference measurements available globally,
which would lead to major AGB biases in regions not represented in the training dataset.

835

Underestimation of AGB in dense mature forests (Fig. 5 and Fig. S8S9) were further exacerbated by the weak sensitivity of
SAR backscatter to forest variables (Table S8). The range of AGB beyond 150 Mg ha⁻¹, in particular in wet biomes,
corresponded to a backscatter range at C- and L-band of less than 0.3 dB ([Cartus and Santoro, 2019; Santoro et al., 2015a](#))[\(Cartus and Santoro, 2019; Santoro et al., 2015a\)](#) and thus had substantial uncertainty. Combining estimates of biomass
840 from multiple C-band Envisat ASAR observations increased the accuracy of the AGB retrieval compared to individual
estimates (Santoro et al., 2011). However, the ALOS PALSAR L-band dataset consisted of a single observation so that any
contribution to the SAR backscatter not related to the forest itself translated into a bias. This was most relevant in the wet
tropics where AGB was estimated solely on such single observations. Imperfect modelling of the ALOS PALSAR backscatter
in sloping terrain caused an additional bias, which propagated through to the estimate of AGB (Fig. S13S14).

845

Characterization of the spatial variability of wood density and biomass expansion factors was critical. We acknowledge that
in less sampled regions, like savannas, the estimates of AGB might have been limited by the observations used to estimate
WD and BEF. Nonetheless, even if we could not quantify the impact of local BCEF biases on the AGB estimates because
GSV observations were unavailable for most of the inventory, our understanding of the retrieval errors (Table S8) and
850 uncertainties (Fig. S9S10) indicate that the main contributors to the errors were not associated to the BCEF layer. This
interpretation, however, is affected by the opportunistic dataset of plot inventory measurements available to assess the AGB
estimates. We agree that further research opportunities should focus on understanding the implications of local wood formation
strategies to AGB estimates at local scales and across environmental gradients.

855

Our estimate of the total AGB in forests increased from 522 Pg to 596 Pg when alsofurther accounting for the woody biomass
in non-forest CCI land cover classesrelated to non-woody natural vegetation. An additional 4 Pg was estimated in cropland

hat formatiert: Englisch (USA)

hat formatiert: Englisch (USA)

hat formatiert: Englisch (USA)

and non-vegetated land cover types, resulting in a terrestrial AGB pool of 600 Pg. The lack of AGB estimates for the Western Pacific Islands had a negligible effect on these values. The slight AGB underestimates woody AGB pool of 600 Pg, corresponding to approximately 282 PgC assuming the mean carbon fraction in woody vegetation of 0.47 (IPCC, 2006). Based on our dataset and accounting for additional maps of non-woody biomass and auxiliary datasets, a more comprehensive report of the global carbon stock both above- and below-ground was recently published (Spawn et al., 2020). The lack of AGB estimates for the Western Pacific Islands had a negligible effect on our total AGB values. AGB underestimates and substantial uncertainty in our dataset arose from having only a single observation of L-band backscatter and the inability of C-band observations to resolve levels of AGB in forests with high biomass stocks. This effectively led to use of a single radar observation to estimate biomass in dense tropical forests, causing our values to be at the low end of the range of estimates of the terrestrial AGB pool (Table 3 and Table S1). Although our estimates for the wet tropics were affected by bias, our dataset provides new insights into the spatial distribution and levels of AGB in tropical, sub-tropical and temperate forests (Table 3, Fig. 10) and provides overall more accurate spatial patterns of the terrestrial carbon pool than previous maps (Fig. 11). For the boreal domain, our estimates agreed with previous maps, which were reported to be unbiased (Table 3, Fig. 10), whilst providing a more detailed portrait of the spatial distribution of AGB (Fig. 3) because of the higher spatial resolution of the remote sensing datasets used in this study (1 ha vs. 100 ha).

At country level, we demonstrate the benefit of our map-based estimates of AGB and total AGB in the context of global reporting of biomass pools (Table 2, Fig. 7). Nonetheless, our dataset may not be sufficiently accurate for individual countries to use it as the only basis for reporting biomass and carbon pools, in particular for countries where the AGB estimates may be experiencing systematic retrieval errors (e.g., predominant high AGB, limited sensitivity of the backscatter to biomass, strong topography). However, it may provide support to national reporting in conjunction with a forest inventory system (Næsset et al., 2020). Our dataset agreed well with values reported in the 2010 FRA for countries with an established national forest inventory (Fig. 7, Table S7). However, for countries reporting AGB on the basis of expert knowledge, best guess estimates and default parameters, we identified clear regional patterns of either overestimation (e.g., in Africa, see Fig. 7) or underestimation (e.g., in Asia, see Fig. 7), which could be related to the capacity of national forest inventories (Fig. 8). Although the global average AGB in our map differs from the FRA value by only 8% (108 Mg ha⁻¹ vs. 119 Mg ha⁻¹, Table 2), the smaller forest area reported in the FRA implies that our total AGB estimate exceeded the value reported in the FRA by 11%.

Our dataset provides new insights on the spatial distribution and magnitude of terrestrial woody AGB as well as a valuable resource for climate and carbon modelling because of its completeness, standardized estimation procedure and high level of detail. The major caveat of past and current spaceborne remote sensing observations used to estimate biomass is their limited sensitivity to any forest parameter. Such limitations may be partly overcome by combining measurements and models, as demonstrated in this study. Nonetheless, it will only be during the 2020s that observations from space will allow accurate

hat formatiert: Deutsch

quantification of the terrestrial biomass pool. ~~with the~~, The availability of a wide range of observations from space, including LiDAR (Dubayah et al., 2020) and P-band SAR (Quegan et al., 2019), ~~which~~ are expected to provide more detailed information on the vertical ~~and horizontal~~ structure of forests, thus adding up to the information by backscatter measurements sensitive mostly to horizontal properties and thus of limited reliability in tall and dense forests.

895

Data availability

The AGB and GSV datasets can be downloaded from <http://globbiomass.org/products/global-mapping> or from the PANGAEA repository at <https://doi.org/10.1594/PANGAEA.894711> ([Santoro, 2018](#)) ([Santoro, 2018](#)).

Author contributions

900 M.Sa. and O.C. designed the study, developed the GSV retrieval algorithm, generated the GSV and AGB datasets, and interpreted the results. N.C. designed the GSV to AGB conversion and generated the conversion dataset. D.R., V.A., A.A., S.d.B. and M.H. designed the validation procedure of the AGB dataset, compiled the database of plot inventory data and computed the grid cell values from the reference database. S.Q. acted as scientific lead of the study. P.R.V., H.B. and J.Car. contributed to the validation of the AGB dataset in South America and Africa. D.S. and M.K. contributed to the validation of 905 the GSV dataset in Russia. M.Sh. and T.I. generated the ALOS PALSAR mosaics and provided support to their use. A.M.M. contributed to the estimation of the wood density dataset. J.Cav., R.C.G., P.d.C.B., N.D., N.L., J.Lia., J.Lin., E.T.A.M., A.M., A.P.P., C.M.R., F.S., G.V.L., H.V., A.W. and S.W. provided forest plot data. M.S. and O.C. wrote the manuscript. S.Q. overviewed the writing of the manuscript. All authors contributed to the drafting of the manuscript.

910 Competing interests

The authors declare they have no competing interests.

Disclaimer

The data is provided as is with no warranties.

915

Acknowledgments

The work was supported by the European Space Agency (ESA) within the Data User Element (DUE) GlobBiomass project (ESRIN contract No. 4000113100/14/I-NB). The National Centre for Earth Observation was supported by the UK Natural Environment Research Council. ~~The Russian Science Foundation supported data processing for the sample plots from Russia~~
920 ~~(project no. The FOS data collection for Russia was performed within the framework of the state assignment of the Center for~~

[Forest Ecology and Productivity of the Russian Academy of Sciences \(no. AAAA-A18-118052590019-7\), the Russian ground data preparation and pre-processing were financially supported by the Russian Science Foundation \(project no. 19-77-30015\).](#)

We are thankful to the GlobBiomass project team and F. M. Seifert (ESA) for valuable suggestions and stimulating scientific discussions. We are thankful to T. Tadono (JAXA EORC), M. Hayashi, (JAXA EORC), K. Kobayashi (RESTEC), A. Rosenqvist ([SoloEOsoloEO](#)) and J. Kellndorfer (EBD) for support with the use and interpretation of the ALOS PALSAR mosaics. Support by the CCI Land Cover project team, in particular, S. Bontemps, UCL, is greatly acknowledged. The help from Martin Jung (MPI-BGC) in feature selection and Ulrich Weber (MPI-BGC) for data processing for the GSV to AGB conversions is greatly acknowledged. Forest inventory data for the validation of the AGB map were made available among others by Ben de Jong, the Prince Edward Island Department of Communities, Land & Environment, Forests, Fish & Wildlife Division and the Nova Scotia Department of Natural Resources. Inventory Data were also provided by the Sustainable Landscapes Brazil project supported by the Brazilian Agricultural Research Corporation (EMBRAPA), the US Forest Service, and USAID, and the US Department of State. We thank [the two anonymous reviewers, contributors to the short comments](#), G. Hengeveld and J. Chave for reviewing and improving the manuscript.

Envisat ASAR data have been distributed and processed on ESA's Grid Processing On Demand (G-POD) platform under ESA's Category-1 Project IDs 6397 and 9209. Access to G-POD was obtained under ESA's Category-1 project nr. 9209. R. Cuccu, J. M. Delgado Blasco, S. Pinto and J. Farres, ESA/ESRIN, are acknowledged for implementation of SAR processing chain on G-POD and support.

940 References

- Avitabile, V. and Camia, A.: An assessment of forest biomass maps in Europe using harmonized national statistics and inventory plots, *For. Ecol. Manag.*, **409**, 489–498, <https://doi.org/10.1016/j.foreco.2017.11.047>, 2018.
- Avitabile, V., Herold, M., Heuvelink, G. B. M., Lewis, S. L., Phillips, O. L., Asner, G. P., Armston, J., Ashton, P. S., Banin, L., Bayol, N., Berry, N. J., Boeckx, P., de Jong, B. H. J., DeVries, B., Girardin, C. A. J., Kearsley, E., Lindsell, J. A., Lopez-Gonzalez, G., Lucas, R., Malhi, Y., Morel, A., Mitchard, E. T. A., Nagy, L., Qie, L., Quinones, M. J., Ryan, C. M., Ferry, S. J. W., Sunderland, T., Laurin, G. V., Gatti, R. C., Valentini, R., Verbeeck, H., Wijaya, A., and Wilcock, S.: An integrated pan-tropical biomass map using multiple reference datasets, *Global Glob. Change Biology-Biol.*, **22**(4), 1406–1420, <https://doi.org/10.1111/gcb.13139>, 2016.
- Baccini, A., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S., and Houghton, R. A.: Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps, *Nature Climate Nat. Clim. Change*, **2**(2), 182–185, <https://doi.org/10.1038/nclimate1354>, 2012.
- Baccini, A., Walker, W., Carvalho, L., Farina, M., Sulla-Menashe, D., and Houghton, R. A.: Tropical forests are a net carbon source based on aboveground measurements of gain and loss, *Science*, **358**, 230–234, <https://doi.org/10.1126/science.aam5962>, 2017.

- hat formatiert:** Schriftfarbe: Automatisch, Deutsch
Formatiert: Literaturverzeichnis2
hat formatiert: Schriftfarbe: Automatisch, Deutsch
hat formatiert: Schriftfarbe: Automatisch, Deutsch

- 955 Bar-On, Y. M., Phillips, R., and Milo, R.: The biomass distribution on Earth, *Proc. Natl. Acad. Sci. USA*, 115(25), 6506–6511, <https://doi.org/10.1073/pnas.1711842115>, 2018.
- Bloom, A. A., Exbrayat, J.-F., van der Velde, I. R., Feng, L., and Williams, M.: The decadal state of the terrestrial carbon cycle: Global retrievals of terrestrial carbon allocation, pools, and residence times, *Proceedings of the National Academy of Sciences, Proc. Natl. Acad. Sci.*, 113(5), 1285–1290, <https://doi.org/10.1073/pnas.1515160113>, 2016.
- 960 Bouvet, A., Mermoz, S., Le Toan, T., Villard, L., Mathieu, R., Naidoo, L., and Asner, G. P.: An above-ground biomass map of African savannahs and woodlands at 25 m resolution derived from ALOS PALSAR, *Remote Sensing of Environment, Sens. Environ.*, 206, 156–173, <https://doi.org/10.1016/j.rse.2017.12.030>, 2018.
- Brown, S.: Estimating biomass and biomass change of tropical forests: A primer, Food and Agriculture Organization, 55 pp., 1987.
- 965 Cartus, O. and Santoro, M.: Exploring combinations of multi-temporal and multi-frequency radar backscatter observations to estimate above-ground biomass of tropical forest, *Remote Sensing of Environment, Sens. Environ.*, 232, 111313, <https://doi.org/10.1016/j.rse.2019.111313>, 2019.
- Cartus, O., Kellndorfer, J., Rombach, M., and Walker, W.: Mapping canopy height and growing stock volume using airborne lidar, alos-palsarLiDAR, ALOS PALSAR, and landsat-landsat ETM+, *Remote Sensing, Sens. Environ.*, 4(11), 3320–3345, <https://doi.org/10.3390/rs4113320>, 2012a.
- 970 Cartus, O., Santoro, M., and Kellndorfer, J.: Mapping forest aboveground biomass in the Northeastern United States with ALOS PALSAR dual-polarization L-band, *Remote Sensing of Environment, Sens. Environ.*, 124, 466–478, <https://doi.org/10.1016/j.rse.2012.05.029>, 2012b.
- Carvalhais, N., Forkel, M., Khomik, M., Bellarby, J., Jung, M., Migliavacca, M., Mu, M., Saatchi, S., Santoro, M., Thurner, M., Weber, U., Ahrens, B., Beer, C., Cescatti, A., Randerson, J. T., and Reichstein, M.: Global covariation of carbon turnover times with climate in terrestrial ecosystems, *Nature*, 514(7521), 213–217, <https://doi.org/10.1038/nature13731>, 2014.
- 975 Chave, J., Coomes, D., Jansen, S., Lewis, S. L., Swenson, N. G., and Zanne, A. E.: Towards a worldwide wood economics spectrum, *Ecology Letters, Ecol. Lett.*, 12(4), 351–366, <https://doi.org/10.1111/j.1461-0248.2009.01285.x>, 2009.
- Ciais, P., Dolman, A. J., Bombelli, A., Duren, R., Peregon, A., Rayner, P. J., Miller, C., Gobron, N., Kinderman, G., Marland, G., Gruber, N., Chevallier, F., Andres, R. J., Balsamo, G., Bopp, L., Bréon, F.-M., Broquet, G., Dargaville, R., Battin, T. J., Borges, A., Bovensmann, H., Buchwitz, M., Butler, J., Canadell, J. G., Cook, R. B., DeFries, R., Engelen, R., Gurney, K. R., Heinze, C., Heimann, M., Held, A., Henry, M., Law, B., Luysaert, S., Miller, J., Moriyama, T., Moulin, C., Myneni, R. B., Nussli, C., Obersteiner, M., Ojima, D., Pan, Y., Paris, J.-D., Piao, S. L., Poultier, B., Plummer, S., Quegan, S., Raymond, P., Reichstein, M., Rivier, L., Sabine, C., Schimel, D., Tarasova, O., Valentini, R., Wang, R., van der Werf, G., Wickland, D., Williams, M., and Zehner, C.: Current systematic carbon-cycle observations and the need for implementing a policy-relevant carbon observing system, *Biogeosciences*, 11(13), 3547–3602, <https://doi.org/10.5194/bg-11-3547-2014>, 2014.
- Dubayah, R., Blair, J. B., Goetz, S., Fatoyinbo, L., Hansen, M., Healey, S., Hofton, M., Hurt, G., Kellner, J., Luthcke, S., Armston, J., Tang, H., Duncanson, L., Hancock, S., Jantz, P., Marselis, S., Patterson, P., Qi, W., and Silva, C.: The Global Ecosystem Dynamics Investigation: High-resolution laser ranging of the Earth's forests and topography, *Science of Sci. Remote Sensing, Sens.*, 100002, <https://doi.org/10.1016/j.srs.2020.100002>, 2020.
- Erb, K.-H., Kastner, T., Plutzar, C., Bais, A. L. S., Carvalhais, N., Fetzel, T., Gingrich, S., Haberl, H., Lauk, C., Niedertscheider, M., Pongratz, J., Thurner, M., and Luysaert, S.: Unexpectedly large impact of forest management and grazing on global vegetation biomass, *Nature*, 553(7686), 73–76, <https://doi.org/10.1038/nature25138>, 2017/2018.

- 995 Exbrayat, J.-F., Bloom, A. A., Carvalhais, N., Fischer, R., Huth, A., MacBean, N., and Williams, M.: Understanding the Land Carbon Cycle with Space Data: Current Status and Prospects, *Surv. Geophys.*, 40(4), 735–755, <https://doi.org/10.1007/s10712-019-09506-2>, 2019. ... [12]
- 000 FAO: Global Forest Resources Assessment 2010, Rome, [online] Available from: <http://www.fao.org/forestry/fra/fra2010/en/>, 2010. **hat formatiert:** Schriftfarbe: Automatisch, Deutsch
- 000 Gibbs, H. K., Brown, S., Niles, J. O., and Foley, J. A.: Monitoring and estimating tropical forest carbon stocks: making REDD a reality, *Environ. Res. Lett.*, 2, 045023, <https://doi.org/10.1088/1748-9326/2/4/045023>, 2007.
- 000 Gillis, M. D., Omule, A. Y., and Brierley, T.: Monitoring Canada's forests: The National Forest Inventory, *For. Chron.*, 81, 214–221, <https://doi.org/10.5558/tfc81214-2>, 2005.
- 005 Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thau, D., Stehman, S. V., Goetz, S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O., and Townshend, J. R. G.: High-resolution global maps of 21-st century forest cover change, *Science*, 342, 850–853, 2013, <https://doi.org/10.1126/science.1244693>, 2013. **hat formatiert:** Schriftfarbe: Automatisch, Deutsch
Formatiert: Literaturverzeichnis2
hat formatiert ... [13]
- 010 Herold, M., Carter, S., Avitabile, V., Espeso, A. B., Jonckheere, I., Lucas, R., McRoberts, R. E., Næsset, E., Nightingale, J., Petersen, R., Reiche, J., Romijn, E., Rosengqvist, A., Rozendaal, D. M. A., Seifert, F. M., Sanz, M. J., and De Sy, V.: The Role and Need for Space-Based Forest Biomass-Related Measurements in Environmental Management and Policy, *Surv. Geophys.*, 40(4), 757–778, <https://doi.org/10.1007/s10712-019-09510-6>, 2019. **hat formatiert** ... [14]
- 010 Hofton, M. A., Minster, J. B., and Blair, J. B.: Decomposition of laser altimeter waveforms, *IEEE Transactions on Geoscience and Trans. Geosci. Remote Sensing Sens.*, 38(4), 1989–1996, <https://doi.org/10.1109/36.851780>, 2000. **hat formatiert** ... [15]
- 010 Houghton, R. A.: Aboveground Forest Biomass-forest biomass and the Global Carbon Balance, *Globalglobal carbon balance, Glob. Change Biology Biol.*, 11(6), 945–958, <https://doi.org/10.1111/j.1365-2486.2005.00955.x>, 2005. **hat formatiert** ... [16]
- 015 Houghton, R. A., Hall, F., and Goetz, S. J.: Importance of forest biomass in the global carbon cycle, *Journal of Geophysical Research J. Geophys. Res.*, 114, G00E03, <https://doi.org/10.1029/2009JG000935>, 2009. **hat formatiert** ... [17]
- 015 Hu, T., Su, Y., Xue, B., Liu, J., Zhao, X., Fang, J., and Guo, Q.: Mapping Global Forest Aboveground Biomassglobal forest aboveground biomass with Spacebornespaceborne LiDAR, Optical Imageryoptical imagery, and Forest Inventory Dataforest inventory data, *Remote Sensing Sens.*, 8(7), 565, <https://doi.org/10.3390/rs8070565>, 2016. **hat formatiert** ... [18]
- 020 IPCC: 2006 IPCC guidelines for national greenhouse gas inventories, Volume 4: Agriculture, Forestry and Other Land Use, 2006. **hat formatiert:** Schriftfarbe: Automatisch, Deutsch
- 020 Jenkins, J. C., Chojnacky, D. C., Heath, L. S., and Birdsey, R. A.: National-scale biomass estimators for United States tree species, *Forest Science For. Sci.*, 49(1), 12–35, 2003. **hat formatiert** ... [19]
- 020 Kindermann, G. E., McCallum, I., Fritz, S., and Obersteiner, M.: A global forest growing stock, biomass and carbon map based on FAO statistics, *Silva Fennica Fenn.*, 42(3), 387–396, <https://doi.org/10.14214/sf.244>, 2008. **hat formatiert** ... [20]
- 025 Kurvonen, L., Pulliainen, J., and Hallikainen, M.: Retrieval of biomass in boreal forests from multitemporal ERS-1 and JERS-1 SAR images, *IEEE Transactions on Geoscience and Trans. Geosci. Remote Sensing Sens.*, 37(1), 198–205, <https://doi.org/10.1109/36.739154>, 1999. **hat formatiert** ... [21]
- Le Quéré, C., Andrew, R. M., Friedlingstein, P., Sitch, S., Hauck, J., Pongratz, J., Pickers, P. A., Korsbakken, J. I., Peters, G. P., Canadell, J. G., Arneth, A., Arora, V. K., Barbero, L., Bastos, A., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Doney, ...

- 030 S. C., Gkritzalis, T., Goll, D. S., Harris, I., Haverd, V., Hoffman, F. M., Hoppema, M., Houghton, R. A., Hurt, G., Ilyina, T., Jain, A. K., Johannessen, T., Jones, C. D., Kato, E., Keeling, R. F., Goldeijk, K. K., Landschützer, P., Lefèvre, N., Liemert, S., Liu, Z., Lombardozzi, D., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakao, S., Neill, C., Olsen, A., Ono, T., Patra, P., Pergon, A., Peters, W., Peylin, P., Pfeil, B., Pierrot, D., Poulet, B., Rehder, G., Resplandy, L., Robertson, E., Rocher, M., Rödenbeck, C., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Steinhoff, T., Sutton, A., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., Wright, R., Zachle, S., and Zheng, B.: Global Carbon Budget 2018, *Earth Syst. Sci. Data*, **10**(4), 2141–2194, <https://doi.org/10.5194/cssd-10-2141-2018>, 2018. hat formatiert ... [22]
- 035 Li, W., Ciais, P., Peng, S., Yue, C., Wang, Y., Thurner, M., Saatchi, S. S., Arneth, A., Avitabile, V., Carvalhais, N., Harper, A. B., Kato, E., Koven, C., Liu, Y. Y., Nabel, J. E. M. S., Pan, Y., Pongratz, J., Poulet, B., Pugh, T. A. M., Santoro, M., Sitch, S., Stocker, B. D., Viovy, N., Wiltshire, A., Yousefpour, R., and Zachle, S.: Land-use and land-cover change carbon emissions between 1901 and 2012 constrained by biomass observations, *Biogeosciences*, **14**(22), 5053–5067, <https://doi.org/10.5194/bg-14-5053-2017>, 2017. hat formatiert ... [23]
- 040 Liu, Y. Y., van Dijk, A. I. J. M., de Jeu, R. A. M., Canadell, J. G., McCabe, M. F., Evans, J. P., and Wang, G.: Recent reversal in loss of global terrestrial biomass, *Nature Climate Nat. Clim. Change*, **5**(5), 470–474, <https://doi.org/10.1038/nclimate2581>, 2015. hat formatiert ... [24]
- 045 Los, S. O., Rosette, J. A. B., Kljun, N., North, P. R. J., Chasmer, L., Suárez, J. C., Hopkinson, C., Hill, R. A., Van Gorsel, E., Mahoney, C., and Berni, J. A. J.: Vegetation height and cover fraction between 60° S and 60° N from ICESat GLAS data, *Geoscientific Geosci. Model Development Dev.*, **5**(2), 413–432, <https://doi.org/10.5194/gmd-5-413-2012>, 2012. hat formatiert ... [25]
- 050 Lu, D., Chen, Q., Wang, G., Liu, L., Li, G., and Moran, E.: A survey of remote sensing-based aboveground biomass estimation methods in forest ecosystems, *International Journal of Digital Int. J. Digit. Earth*, **9**(1), 63–105, <https://doi.org/10.1080/17538947.2014.990526>, 2016. hat formatiert ... [26]
- 055 Lucas, R. M., Mitchell, A. L., and Armston, J.: Measurement of Forest Above-Ground Biomass Using Active and Passive Remote Sensing at Large (Subnational to Global) Scales, *Current Forestry Reports: Curr. For. Rep.*, **1**(3), 162–177, <https://doi.org/10.1007/s40725-015-0021-9>, 2015. hat formatiert ... [27]
- 060 Mermoz, S., Bouvet, A., Toan, T. L., and Herold, M.: Impacts of the forest definitions adopted by African countries on carbon conservation, *Environ. Res. Lett.*, **13**(10), 104014, <https://doi.org/10.1088/1748-9326/aae3b1>, 2018. hat formatiert ... [28]
- Mitchard, E. T. A., Saatchi, S. S., Baccini, A., Asner, G. P., Goetz, S. J., Harris, N. L., and Brown, S.: Uncertainty in the spatial distribution of tropical forest biomass: A comparison of pan-tropical maps, *Carbon Balance and Management Manag.*, **8**(1), 1, <https://doi.org/10.1186/1750-0680-8-10>, 2013. hat formatiert ... [29]
- 065 Næsset, E., McRoberts, R. E., Pekkarinen, A., Saatchi, S., Santoro, M., Trier, Ø. D., Zahabu, E., and Gobakken, T.: Use of local and global maps of forest canopy height and aboveground biomass to enhance local estimates of biomass in miombo woodlands in Tanzania, *International Journal of Applied Int. J. Appl. Earth Observation and Obs. Geoinformation*, **89**, 102109, <https://doi.org/10.1016/j.jag.2020.102109>, 2020. hat formatiert ... [30]
- Ometto, J. P., Aguiar, A. P., Assis, T., Soler, L., Valle, P., Tejada, G., Lapola, D. M., and Meir, P.: Amazon forest biomass density maps: tackling the uncertainty in carbon emission estimates, *Climatic Clim. Change*, **124**(3), 545–560, <https://doi.org/10.1007/s10584-014-1058-7>, 2014. hat formatiert ... [31]
- Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S., and Hayes, D.: A hat formatiert: Schriftfarbe: Automatisch, Deutsch

- 070 [Large](#) and [Persistent Carbon Sink](#)[persistent carbon sink in the World's Forests](#)[world's forests](#), *Science*, 333(6045), 988–993, <https://doi.org/10.1126/science.1201609>, 2011. hat formatiert ... [32]
- 075 Pulliainen, J. T., Heiska, K., Hyppä, J., and Hallikainen, M. T.: Backscattering properties of boreal forests at the C- and X-bands, *IEEE Transactions on Geoscience and Trans. Geosci. Remote Sensing*, 32(5), 1041–1050, <https://doi.org/10.1109/36.312892>, 1994. hat formatiert ... [33]
- 080 Quegan, S., Rauste, Y., Bouvet, A., Carreiras, J., Cartus, O., Carvalhais, N., Le Toan, T., Mermoz, S. and Santoro, M.: DUE GlobBiomass - Algorithm Theoretical Basis Document. [online] Available from: <http://globbiomass.org/products/global-mapping/>, 2017. hat formatiert ... [34]
- 085 Quegan, S., Le Toan, T., Chave, J., Dall, J., Exbrayat, J.-F., Minh, D. H. T., Lomas, M., D'Alessandro, M. M., Paillou, P., Papathanassiou, K., Rocca, F., Saatchi, S., Scipal, K., Shugart, H., Smallman, T. L., Soja, M. J., Tebaldini, S., Ulander, L., Villard, L., and Williams, M.: The European Space Agency BIOMASS mission: Measuring forest above-ground biomass from space, *Remote Sensing of Environment*, 227, 44–60, <https://doi.org/10.1016/j.rse.2019.03.032>, 2019. hat formatiert ... [35]
- 090 Reichstein, M. and Carvalhais, N.: Aspects of Forest Biomass in the Earth System: Its Role and Major Unknowns, *Surv. Geophys.*, 40(4), 693–707, <https://doi.org/10.1007/s10712-019-09551-x>, 2019. hat formatiert ... [36]
- 095 Rodriguez-Veiga, P., Wheeler, J., Louis, V., Tansey, K., and Balzter, H.: Quantifying [Forest Biomass Carbon Stocks From Space](#), [Current Forestry Reports](#)[forest biomass carbon stocks from space](#), *Curr. For. Rep.*, 3(1), 1–18, <https://doi.org/10.1007/s40725-017-0052-5>, 2017. hat formatiert ... [37]
- 100 Rodriguez-Veiga, P., Quegan, S., Carreiras, J., Persson, H. J., Fransson, J. E. S., Hoscilo, A., Ziolkowski, D., Sterenekaz, K., Lohberger, S., Stängel, M., Berninger, A., Siegert, F., Avitabile, V., Herold, M., Mermoz, S., Bouvet, A., Le Toan, T., Carvalhais, N., Santoro, M., Cartus, O., Rauste, Y., Mathieu, R., Asner, G. P., Thiel, C., Pathé, C., Schmullius, C., Seifert, F. M., Tansey, K., and Balzter, H.: Forest biomass retrieval approaches from [earth observation in different biomes](#), [International Journal of Applied Earth Observation and Geoinformation](#)[in different biomes](#), *Int. J. Appl. Earth Obs. Geoinformation*, 77, 53–68, <https://doi.org/10.1016/j.jag.2018.12.008>, 2019. hat formatiert ... [38]
- Romijn, E., Lantican, C. B., Herold, M., Lindquist, E., Ochieng, R., Wijaya, A., Murdiyarso, D., and Verchot, L.: Assessing change in national forest monitoring capacities of 99 tropical countries, *Forest Ecology and Management*, [For. Ecol. Manag.](#), 352, 109–123, <https://doi.org/10.1016/j.foreco.2015.06.003>, 2015. hat formatiert ... [39]
- Rosenqvist, A., Shimada, M., Ito, N., and Watanabe, M.: ALOS PALSAR: A pathfinder mission for global-scale monitoring of the environment, *IEEE Trans Geosc Rem Sens Vol.* 45, 3307–3316, <https://doi.org/10.1109/TGRS.2007.901027>, 2007.
- Saatchi, S., Marlier, M., Chazdon, R. L., Clark, D. B., and Russell, A. E.: Impact of spatial variability of tropical forest structure on radar estimation of aboveground biomass, *Remote Sensing of Environment*, 115(11), 2836–2849, <https://doi.org/10.1016/j.rse.2010.07.015>, 2011a. hat formatiert Schriftfarbe: Automatisch, Deutsch Formatiert: Literaturverzeichnis2 ... [40]
- Saatchi, S. S., Harris, N. L., Brown, S., Lefsky, M., Mitchard, E. T. A., Salas, W., Zutta, B. R., Buermann, W., Lewis, S. L., Hagen, S., Petrova, S., White, L., Silman, M., and Morel, A.: Benchmark map of forest carbon stocks in tropical regions across three continents, *Proceedings of the National Academy of Sciences*, *Proc. Natl. Acad. Sci.*, 108(24), 9899–9904, <https://doi.org/10.1073/pnas.1019576108>, 2011b. hat formatiert ... [41]
- Santoro, M.: GlobBiomass - global datasets of forest biomass, <https://doi.org/10.1594/PANGAEA.894711>, 2018. hat formatiert ... [42]

- 105 Santoro, M. and Cartus, O.: Research Pathways of Forest Above Ground Biomass Estimation Basedforest above-ground biomass estimation based on SAR Backscatterbackscatter and Interferometricinterferometric, SAR Observationsobservations, Remote Sensing, Sens., 10(4), 608, <https://doi.org/10.3390/rs10040608>, 2018. ... [43]
- 110 Santoro, M., Askne, J., Smith, G., and Fransson, J. E. S.: Stem volume retrieval in boreal forests from ERS-1/2 interferometry, Remote Sensing of Environment, Sens. Environ., 81(1), 19–35, [https://doi.org/10.1016/S0034-4257\(01\)00329-7](https://doi.org/10.1016/S0034-4257(01)00329-7), 2002. ... [44]
- 115 Santoro, M., Beer, C., Cartus, O., Schmullius, C., Shvidenko, A., McCallum, I., Wegmüller, U., and Wiesmann, A.: Retrieval of growing stock volume in boreal forest using hyper-temporal series of Envisat ASAR ScanSAR backscatter measurements, Remote Sensing of Environment, Sens. Environ., 115(2), 490–507, <https://doi.org/10.1016/j.rse.2010.09.018>, 2011. ... [45]
- 120 Santoro, M., Beaudoin, A., Beer, C., Cartus, O., Fransson, J. E. S., Hall, R. J., Pathe, C., Schepaschenko, D., Schmullius, C., Shvidenko, A., Thurner, M., and Wegmüller, U.: Forest growing stock volume of the northern hemisphere: spatiallyspatially, explicit estimates for 2010 derived from Envisat ASAR data, Remote Sensing of Environment, Sens. Environ., 168, 316–334, <https://doi.org/10.1016/j.rse.2015.07.005>, 2015a. ... [46]
- 125 Santoro, M., Wegmüller, U., Lamarche, C., Bontemps, S., Defourny, P., and Arino, O.: Strengths and weaknesses of multi-year Envisat ASAR backscatter measurements to map permanent open water bodies at global scale, Remote Sensing of Environment, Sens. Environ., 171, 185–201, <https://doi.org/10.1016/j.rse.2015.10.031>, 2015b. ... [47]
- 130 Schepaschenko, D., Kraxner, F., See, L., Fuss, S., McCallum, I., Fritz, S., Perger, C., Shvidenko, A., Kindermann, G., Frank, S., Tum, M., Schmid, E., Balkovič, J., and Günther, K.: Global Biomass Information: From Data Generation to Application, in: Handbook of Clean Energy Systems, edited by J. Yan, pp. 1–23, John Wiley & Sons, Ltd, Chichester, UK, 1–23, <https://doi.org/10.1002/978111991978.hces173>, 2015. ... [48]
- 135 Schepaschenko, D., Molchanova, E., Fedorov, S., Karminov, V., Ontikov, P., Santoro, M., See, L., Kositsyn, V., Shvidenko, A., Romanovskaya, A., Korotkov, V., Lesiv, M., Bartalev, S., Fritz, S., Schepashchenko, M., and Kraxner, F.: Russian forest sequesters substantially more carbon than previously reported, Sci. Rep., 11:12825, 7, <https://doi.org/10.1038/s41598-021-92152-9>, 2021.
- 140 Schimel, D., Pavlick, R., Fisher, J. B., Asner, G. P., Saatchi, S., Townsend, P., Miller, C., Frankenberger, C., Hibbard, K., and Cox, P.: Observing terrestrial ecosystems and the carbon cycle from space, Global Glob. Change Biology, Biol., 21(5), 1762–1776, <https://doi.org/10.1111/gcb.12822>, 2015. ... [49]
- Shimada, M.: Ortho-rectification and slope correction of SAR data Usingusing DEM and its accuracy evaluation, IEEE Journal of Selected Topics in Applied Earth Observations and Applications, Remote Sensing, Sens., 3(4), 657–671, <https://doi.org/10.1109/JSTARS.2010.2027984>, 2010. ... [50]
- Simard, M., Pinto, N., Fisher, J. B., and Baccini, A.: Mapping forest canopy height globally with spaceborne lidar, Journal of Geophysical Research: LiDAR, J. Geophys. Res., 116(G4), <https://doi.org/10.1029/2011JG001708>, 2011. ... [51]
- Soto-Navarro, C., Ravilious, C., Arnell, A., de Lamo, X., Harfoot, M., Hill, S. L. L., Wearn, O. R., Santoro, M., Bouvet, A., Mermoz, S., Le Toan, T., Xia, J., Liu, S., Yuan, W., Spawn, S. A., Gibbs, H. K., Ferrier, S., Harwood, T., Alkemade, R., Schipper, A. M., Schmidt-Traub, G., Strassburg, B., Miles, L., Burgess, N. D., and Kapos, V.: Mapping co-benefits for carbon storage and biodiversity to inform conservation policy and action, Philos. Trans. R. Soc. B Biol. Sci., 375, 20190128, <https://doi.org/10.1098/rstb.2019.0128>, 2020.
- Spawn, S. A., Sullivan, C. C., Lark, T. J., and Gibbs, H. K.: Harmonized global maps of above and belowground biomass carbon density in the year 2010, Sci. Data, 7, 112, <https://doi.org/10.1038/s41597-020-0444-4>, 2020.

145 Stolbovoi, V. and Mc Callum, I.: Land Resources of Russia. International Institute for Applied Systems Analysis and Russian Academy of Science, Laxenburg, Austria, 2002.

Thum, T., MacBean, N., Peylin, P., Bacour, C., Santaren, D., Longdoz, B., Loustau, D., and Ciais, P.: The potential benefit of using forest biomass data in addition to carbon and water flux measurements to constrain ecosystem model parameters: Case studies at two temperate forest sites, *Agricultural and Forest Meteorology*, *Agric. For. Meteorol.*, 234–235, 48–65, <https://doi.org/10.1016/j.agrformet.2016.12.004>, 2017.

150 Thurner, M., Beer, C., Santoro, M., Carvalhais, N., Wutzler, T., Schepaschenko, D., Shvidenko, A., Kompter, E., Ahrens, B., Levick, S. R., and Schmullius, C.: Carbon stock and density of northern boreal and temperate forests, *Global Ecology and Biogeography*, *Glob. Ecol. Biogeogr.*, 23(3), 297–310, <https://doi.org/10.1111/geb.12125>, 2014.

155 Thurner, M., Beer, C., Carvalhais, N., Forkel, M., Santoro, M., Tum, M., and Schmullius, C.: Large-scale variation in boreal and temperate forest carbon turnover rate related to climate: Climate Related Forest C Turnover Rate, *Geophys. Res. Lett.*, 43(9), 4576–4585, <https://doi.org/10.1002/2016GL068794>, 2016.

Tomppo, E., Gschwantner, T., Lawrence, M., and McRoberts, R. E.: National Forest Inventories - Pathways for common reporting, Springer Science and Business Media, 612 pp., 2010.

hat formatiert: Schriftfarbe: Automatisch, Deutsch

Formatiert: Literaturverzeichnis2

hat formatiert: Schriftfarbe: Automatisch, Deutsch

Formatiert: Standard

Seite 43: [1] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [1] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [1] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [1] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [1] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [1] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [1] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [2] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [2] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [2] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [2] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [2] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [3] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [3] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 43: [3] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Schriftfarbe: Automatisch, Deutsch

Seite 43: [4] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [5] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [5] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [5] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [5] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [5] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [5] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [6] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [6] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [6] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [6] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [7] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [7] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [7] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [7] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
-------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 43: [8] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [8] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [8] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [9] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [9] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [9] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [9] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [10] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [10] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [10] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [10] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [11] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [11] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [11] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [11] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 43: [11] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [12] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [12] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [12] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [12] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [13] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [13] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [14] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [14] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [14] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [14] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [14] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [15] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 44: [15] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		

Schriftfarbe: Automatisch, Deutsch

Seite 44: [16] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [16] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [16] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [16] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [16] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [17] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [17] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [17] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [18] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [18] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [18] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [18] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [18] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [18] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [18] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [18] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 44: [19] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [19] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [20] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [20] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [20] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [20] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [20] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [21] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [21] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [21] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 44: [21] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [22] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [22] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [22] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [22] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [22] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [23] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [23] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [24] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [24] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [24] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [24] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [24] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [25] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [25] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [25] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [25] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [25] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [25] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [26] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [26] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 45: [26] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Schriftfarbe: Automatisch, Deutsch

Seite 45: [27] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [27] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [27] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [27] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [28] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [28] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [28] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [28] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [29] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [29] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [29] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [29] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [30] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [30] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [30] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 45: [31] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [31] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [31] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [31] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 45: [31] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [32] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [32] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [32] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [32] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [32] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [32] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [33] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [33] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [33] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [33] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [33] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 46: [34] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [35] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [35] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [35] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [35] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [36] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [36] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [36] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [36] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [36] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [37] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [37] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [37] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [37] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [37] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Seite 46: [38] hat formatiert Maurizio Santoro 17.06.21 15:43:00

Schriftfarbe: Automatisch, Deutsch

Schriftfarbe: Automatisch, Deutsch

Seite 46: [38] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [38] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [38] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [39] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [39] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [39] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [39] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [40] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [40] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [40] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [40] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [40] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [41] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [41] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [41] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [41] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 46: [42] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [43] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [43] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [43] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [43] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [43] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [43] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [43] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [43] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [43] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [44] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [44] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [45] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [45] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [45] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [45] hat formatiert Maurizio Santoro 17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch

Seite 47: [46] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [46] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [46] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [47] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [47] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [47] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [47] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [48] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [48] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [48] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [48] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [49] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [49] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [49] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		
Seite 47: [49] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
Schriftfarbe: Automatisch, Deutsch		

Schriftfarbe: Automatisch, Deutsch

Seite 47: [50] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [50] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [50] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [50] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [50] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [50] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [51] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [51] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [51] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [51] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch

Seite 47: [51] hat formatiert	Maurizio Santoro	17.06.21 15:43:00
--------------------------------------	-------------------------	--------------------------

Schriftfarbe: Automatisch, Deutsch