

DSCOVER/EPIC-derived global hourly/daily downward shortwave and photosynthetically active radiation data at $0.1^{\circ}\times 0.1^{\circ}$ resolution

Dalei Hao^{1,2,3}, Ghassem R. Asrar⁴, Yelu Zeng¹, Qing Zhu⁵, Jianguang Wen^{2,3}, Qing Xiao^{2,3}, Min Chen¹

¹Joint Global Change Research Institute, Pacific Northwest National Laboratory, College Park, MD 20740, USA

5 ²State Key Laboratory of Remote Sensing Science, Aerospace Information Research Institute, Chinese Academy of Sciences, Beijing 100101, China

³University of Chinese Academy of Sciences, Beijing 100049, China

⁴Universities Space Research Association, Columbia, MD 21046, USA

⁵Earth Science Division, Lawrence Berkeley National Lab, Berkeley, CA 94720, USA

10

Correspondence to: Min Chen (min.chen@pnnl.gov)

Abstract. Downward shortwave radiation (SW) and photosynthetically active radiation (PAR) play crucial roles in Earth system dynamics. Spaceborne remote sensing techniques provide a unique means for mapping accurate spatio-temporally-continuous SW/PAR, globally. However, any individual polar-orbiting or geostationary satellite cannot satisfy the desired high temporal resolution (sub-daily) and global coverage simultaneously, while integrating and fusing multi-source data from complementary satellites/sensors is challenging because of co-registration, inter-calibration, near real-time data delivery and the effects of discrepancies in orbital geometry. The Earth Polychromatic Imaging Camera (EPIC) onboard the Deep Space Climate Observatory (DSCOVER), launched in February 2015, offers an unprecedented possibility to bridge the gap between high temporal resolution and global coverage, and characterize the diurnal cycles of SW/PAR globally. In this study, we adopted a suite of well-validated data-driven machine-learning models to generate the first global land products of SW/PAR, from June 2015 to June 2019, based on DSCOVER/EPIC data. The derived products have high temporal resolution (hourly) and medium spatial resolution ($0.1^{\circ}\times 0.1^{\circ}$), and include estimates of the direct and diffuse components of SW/PAR. We used independently widely-distributed ground station data from the Baseline Surface Radiation Network (BSRN), the Surface Radiation Budget Network (SURFRAD), NOAA's Global Monitoring Division and the U.S. Department of Energy's Atmospheric System Research (ASR) program to evaluate the performance of our products, and further analyzed and compared the spatio-temporal characteristics of the derived products with the benchmarking Clouds and the Earth's Radiant Energy System Synoptic (CERES) data. We found both the hourly and daily products to be consistent with ground-based observations (e.g., hourly and daily total SWs have low biases of -3.96 and -0.71 W/m^2 and root mean square errors (RMSEs) of 103.50 and 35.40 W/m^2 , respectively). The developed products capture the complex spatio-temporal patterns well and accurately track substantial diurnal, monthly, and seasonal variations of SW/PAR when compared to CERES data. They provide a reliable and valuable alternative for solar photovoltaic applications worldwide and can be used to improve our understanding of the diurnal and seasonal variabilities of the terrestrial water, carbon and energy fluxes at various spatial scales. The products are freely available at <https://doi.org/10.25584/1595069> (Hao et al., 2020).

1 Introduction

Downward shortwave radiation (SW) and photosynthetically active radiation (PAR) profoundly affect the terrestrial environment (Wild et al., 2005), and are fundamental for global energy balance (Liang et al., 2010), carbon budget (Farquhar and Roderick, 2003), hydrological cycle (Roderick and Farquhar, 2002), and solar energy production and utilization (Sweerts et al., 2019). Partitioning total SW/PAR into their direct and diffuse components also is important for solar resource management and

photovoltaic power design (Khahro et al., 2015;Raptis et al., 2017), and terrestrial photosynthesis estimations (Mercado et al., 2009;Gu et al., 2002;Chen and Zhuang, 2014;Wang et al., 2018).

40 Satellite remote sensing has been widely used to map SW/PAR across various spatial and temporal scales (Pinker et al., 2005;Huang et al., 2019). Traditional ground-based observations have the required high accuracy but sparse point-specific distributions, and thus inadequate spatial representation (Korany et al., 2016), while numerical modelling with spatio-temporally-continuous mapping has relatively low spatial resolution and large errors and uncertainties (Zhao et al., 2013). In contrast, remote sensing offers a more reliable and efficient tool to estimate high-quality SW/PAR globally with high spatio-temporal resolution, 45 as it characterizes heterogeneous spatial distributions and captures the complex dynamic evolution of atmosphere, cloud and land surface processes at regional and even global scales (Huang et al., 2019;Li et al., 2020). Currently, a series of remote sensing-based SW/PAR datasets/products are available: 1) from polar-orbiting satellites, e.g., Breathing Earth System Simulator (BESS) (Ryu et al., 2018) and MODIS MCD18 products (Wang et al., 2020); 2) from geostationary satellites, e.g., Himawari-8/Advanced Himawari Imager (AHI) (Letu et al., 2020); and 3) from fusing multi-source data/observations, such as the Global LAnd Surface 50 Satellite (GLASS) (Zhang et al., 2014) and the Clouds and the Earth's Radiant Energy System Synoptic (CERES) (Rutan et al., 2015).

Global high-quality SW/PAR data at sub-daily scales are highly desired for investigating the diurnal variabilities of solar-induced fluorescence, photosynthesis (Damm et al., 2010) and evapotranspiration (Van Heerwaarden et al., 2010), and for solar energy utilization (Sweerts et al., 2019). However, accurately quantifying global SW/PAR is challenging based on current polar-orbiting or/and geostationary satellites/sensors, because: 1) sun-synchronous polar-orbiting satellites generally have high spatial 55 resolution but cannot capture the sub-daily variations of SW/PAR owing to low revisiting frequency; 2) geostationary satellites usually have high temporal resolution but limited geographical coverage (i.e. several different satellites systems for covering the entire Earth); 3) fusing multi-source data acquired from complementary satellites/sensors is challenging due to the issues of co-registration, inter-calibration, the effects of different orbital geometries and the difficulty of processing and delivering the final products in near-real time to users. In addition, most of the current remotely sensed SW/PAR estimations are conducted under the 60 assumption of an independent pixel approximation (IPA) and simply neglect the three-dimensional (3D) radiative effects caused by inhomogeneous cloud fields (Wyser et al., 2005). The 3D effects (e.g., nonlocal cloud shadows, reflections from cloud sides, and enhancement of downward radiation by photon diffusion from clouds) can significantly influence the accuracy and quality of high-temporal-resolution SW/PAR estimations and are perhaps the largest error source for SW/PAR retrievals (Wyser et al., 65 2005;Huang et al., 2019). Although several methods have been developed based on full 3D radiative transfer models (Liou et al., 2013), there is currently no practical and computationally-feasible approach to eliminate 3D radiative effects efficiently and completely (Huang et al., 2019).

The Deep Space Climate Observatory (DSCOVR), launched on February, 2015, leads a new era of monitoring the sun and Earth from deep space around the sun-Earth first Lagrange (L1) point (Burt and Smith, 2012). Its advanced Earth-facing camera, 70 Earth Polychromatic Imaging Camera (EPIC), onboard DSCOVR, views nearly the entire sunlit part of the Earth, from pole to pole, in near backscattering directions with 10 spectral bands from the ultra-violet to near-infrared wavelengths every 1~2 hours, giving EPIC a unique capability of monitoring and capturing the diurnal variation of ozone, clouds, aerosols, and vegetation properties (Marshak et al., 2018). DSCOVR/EPIC thus provides an unrivalled tool to capture the diurnal cycles of SW/PAR globally and overcomes some limitations of current remote sensing-based SW/PAR estimations. Compared to any individual polar-orbiting and geostationary satellite, DSCOVR/EPIC essentially bridges the gap between high revisiting frequency and global 75 coverage. Compared to the multi-source integration, the single DSCOVR/EPIC instrument avoids the compatibility and matching issues of using different sensors/satellites, and it is more suitable for processing and delivering the final products in real-time or

near real-time to users. Fortunately, DSCOVR/EPIC is also characterized by a nearly constant scattering phase angle (angle formed between the incident and scattered-to-satellite sunlight vectors) from 168.5° to 175.5°, which implies that DSCOVR/EPIC guarantees that the atmospheric column determining SW/PAR is nearly the same as that observed by the satellite. Therefore, DSCOVR/EPIC has the potential to reduce significantly the 3D radiative effects caused by the shift of the apparent position of clouds and their shadows which are related to the solar and viewing geometries.

The overarching goal of this study is to 1) develop, document and present DSCOVR/EPIC-derived SW/PAR products covering a period of about 4 years (from June, 2015 to June, 2019) based on a suite of well-validated machine learning methods (Hao et al., 2019) and 2) perform a systematic and comprehensive assessment of the accuracy, consistency and spatio-temporal patterns of these products against comparable but independently developed and published data/products; and 3) make the resulting dataset openly available for use by Earth system research and modelling, and for solar energy productions and use. The newly generated products are the first available SW/PAR products with high temporal frequency (hourly) and global coverage at a spatial resolution of 0.1°×0.1°, where the aggregated daily-scale data are available, and the direct and diffuse components of SW/PAR are also provided. We evaluate them against widely-distributed ground station data, analyze their spatio-temporal variations, and compared them to the widely-used CERES products. Finally, possible sources of uncertainties and potential improvements in the future study are discussed.

2 Material and methods

2.1 Remote sensing data

The DSCOVR/EPIC science team has routinely developed and published a suite of official Level 2 (L2) products from DSCOVR/EPIC observations (Marshak et al., 2018), including stratospheric ozone concentrations (Herman et al., 2018), sulfur dioxide (SO₂) from volcanic eruptions, atmospheric aerosols in the UV and visible spectral ranges, cloud parameters (Yang et al., 2019), atmospherically corrected land-surface reflectance and vegetation properties (Yang et al., 2017). These standard EPIC L2 products are publicly available from the NASA Langley Atmospheric Science Data Center and described in detail at https://eosweb.larc.nasa.gov/project/dscovr/dscovr_table. For this study, we obtained solar zenith angle, surface pressure, aerosol optical depth, cloud fraction from L2 aerosol product, cloud optical thickness and the most likely cloud phase from L2 cloud products, and total column ozone from L2 ozone product, as well as the available quality flags for these products. We re-projected all datasets into global latitude/longitude grids with a spatial resolution of 0.1°×0.1° using the nearest neighbourhood resampling method.


As a key component of the Earth Observing System (EOS) program, the CERES project has developed and published globally long-term observed top-of-the-atmosphere (TOA) and calculated surface fluxes for study of climate and cloud feedback (Wielicki et al., 1996). CERES Synoptic 1° (SYN1deg) Edition 4.1 products, released on August 22, 2019, contain global 1°×1° gridded monthly, daily, 3-hourly and hourly averaged TOA and surface fluxes (Rutan et al., 2015). In particular, the SYN1deg Edition 4.1 products provide diurnally complete SW/PAR and their direct and diffuse components. However, the SYN1deg Edition 4.1 products are not suitable for inferring long-term trends of surface fluxes, due to limited climate quality (https://ceres.larc.nasa.gov/documents/DQ_summaries/CERES_SYN1deg_Ed4A_DQS.pdf). The CERES Energy Balanced and Filled (EBAF) Edition 4.1 products, released on May 28, 2019, provide global 1°×1° gridded monthly averaged TOA and surface fluxes (Loeb et al., 2018;Kato et al., 2018). The CERES EBAF products are designed specifically for climate model evaluation and energy budget estimation, and are more suitable for long-term analysis of variability of SW/PAR (e.g. intra-seasonal and inter-annual changes). Both the SYN1deg and EBAF products are freely accessible via the CERES Visualization, Ordering and

Subsetting Tool (https://ceres.larc.nasa.gov/order_data.php). In this study, we used both hourly and daily CERES SYN1deg Edition 4.1 products as a reference to evaluate the spatio-temporal patterns of EPIC-derived SW/PAR products at both hourly and daily scales, and used the CERES EBAF Edition 4.1 products as a benchmark to evaluate the monthly and seasonal variations of EPIC-derived SW/PAR products.

120 2.2 Ground-based observation data

Ground-based measurements with high-quality instrumentation and long-term maintenance provide the most reliable and accurate SW/PAR data, which are generally deemed as the ground truth for evaluating the performance of remote sensing products. Since 1992, the international Baseline Surface Radiation Network (BSRN) under the World Climate Research Programme (WCRP) has provided high quality, high temporal resolution (1 min) ground-based radiation measurements of direct, diffuse and total SWs
125 (Driemel et al., 2018; Ohmura et al., 1998). The BSRN stations are placed strategically to be representative of their relatively large surrounding area, thus suitable for the evaluation of satellite data. The Surface Radiation Budget Network (SURFRAD) supported by the NOAA Climate Program Office is operating in climatologically diverse regions and measuring accurate, continuous, long-term surface radiation budget, and meteorological parameters routinely to support climate and weather studies over the United States (Augustine et al., 2000). The SURFRAD sites also measure and provide PAR data. The CERES/ARM Validation
130 Experiment (CAVE) collected 58 land surface sites from the BSRN, NOAA's Global Monitoring Division, SURFRAD, and the U.S. Department of Energy's Atmospheric System Research (ASR) program and some personal communications (Rutan et al., 2001;Rutan et al., 2015). In the CAVE, all original 1 min data were averaged to different temporal scales (i.e. hourly, daily and monthly) through the strict quality control and gap-filling using the linear interpolation. The CAVE dataset provides SW and its direct and diffuse components, but does not include PAR measurements. Further information on CAVE can be found at
135 <http://www-cave.larc.nasa.gov/>.

We used the CAVE datasets to evaluate the performance of EPIC-derived SW products, and used the SURFRAD datasets to evaluate the performance of EPIC-derived PAR products. The original SURFRAD data were first gap-filled using a linear interpolation technique and then temporally aggregated to both hourly and daily scales. Considering that some data from June, 2015 to December, 2016 were used to train and test the machine learning models (Hao et al., 2019), we used only all available data
140 from 43 CAVE and 7 SURFARAD sites from January, 2017 to June, 2019. **Fig. 1** shows the geographical distribution of ground-based observation stations for evaluation in the study. These sites are further classified into three groups of polar (Arctic or Antarctic), island or coastal, and continental sites.



145 **Figure 1.** Geographical distribution of *in situ* observation sites for used for evaluation of space-based products. The blue, red and magenta circles (with green border) denote the polar (Arctic or Antarctic), island or coastal, and continental sites, respectively; and the black triangle represents the SURFRAD sites.

2.3 Estimation of SW/PAR fluxes

We adopted the trained random forest models developed by Hao et al. (2019) to estimate global SW/PAR from DSCOVER/EPIC
 150 datasets. The evaluation conducted by (Hao et al., 2019) showed that the random forest models perform very well against the
 ground measurements from BRSN and SURFRAD sites. In the study, we acquired hourly driving data for estimating SW/PAR
 based on the collected DSCOVER/EPIC L2 products (introduced in **Section 2.1**). We used the nearest neighbour interpolation
 approach to fill any gaps, based on the assumption that the atmospheric and cloud conditions remain unchanged and only solar
 zenith angle varies during a short period (1~2 hours). Hourly SW/PAR estimates were then produced using the random forest
 155 models and were aggregated into daily products. However, we found that gaps existed in the derived SW/PAR products due to
 failed retrievals of atmospheric and cloud parameters, especially in Arctic and Antarctic polar regions. Therefore, we used the
 CERES SYN1deg Edition 4.1 products to fill these gaps, based on linear interpolation techniques. We assigned quality flags to the
 derived products to denote: 0: successfully estimated from DSCOVER/EPIC; 1: gap-filled based on CERES data; 2: missing data.

3 Results

160 3.1 Evaluation of estimated SW and PAR against ground-based observations


3.1.1 Overall performance of derived products

The hourly EPIC-derived diffuse, direct and total SW/PAR products match very well overall with the ground-based observations
 (Fig. 2). For diffuse SW, the bias, root mean square error (RMSE), relative RMSE (RMSE to mean value, RRMSE) and the
 coefficient of determination (R^2) are 9.8 W/m², 74.97 W/m², 55.21% and 0.60, respectively. Direct SW has negative bias of -16.39
 165 W/m², relatively large RMSE of 137.24 W/m² and RRMSE of 56.17%, and R^2 of 0.73. By contrast, both total SW and PAR have
 better performance with low biases (-3.96 and 7.31 W/m²), smaller RMSEs (103.50 and 50.44 W/m²) and RRMSEs (28.40% and

32.49%), and high R^2 values (0.87 and 0.83). These statistical metrics indicate that EPIC-based hourly SW/PAR estimates are comparable to or better than other remote sensing-based products, e.g., Himawari-8/AHI-derived total SW has similar RMSE of 101.86 W/m^2 and R^2 of 0.87 (Letu et al., 2020).

170 The daily SW/PAR estimates are well correlated with the ground-based observations (**Fig. 3**). Diffuse SW has a positive bias (5.25 W/m^2), a relatively small RMSE (25.25 W/m^2) but a large RRMSE (37.12%), and an R^2 of 0.65. By contrast, Direct SW has a negative bias (-6.09 W/m^2), a relatively large RMSE (45.46 W/m^2), a large RRMSE (39.49%) and a R^2 of 0.77. Total SW shows good performance with a low bias of -0.71 W/m^2 , a RMSE of 35.40 W/m^2 , a smaller RRMSE of 19.45%, and high R^2 of 0.87. Total PAR also shows good relationship with the ground-based data (positive bias of 4.08 W/m^2 , small RMSE of 16.80 W/m^2 and

175 RRMSE of 21.88% and high R^2 of 0.85). These results indicate that our daily products show comparable or better performance compared to other SW/PAR products, e.g., for total SW, MCD18 and GLASS products have similar RMSEs of 32.3 and 35.9 W/m^2 and higher biases of -7.8 and -7.6 W/m^2 (Wang et al., 2020).



180 **Figure 2.** Evaluation of EPIC-based hourly SW/PAR estimates against ground-based observations.





Figure 3. Evaluation of EPIC-based daily SW/PAR estimates against ground-based observations.

3.1.2 Temporal distribution of estimation errors of derived products

185 We first distinguish different sky conditions based on the ratio of diffuse to total SW ($skyl$): 1) clear: $skyl < 0.3$ during 70% time
of one day; 2) overcast: $skyl > 0.7$ during 70% time of one day); and 3) cloudy: all the other cases. **Fig. 4** shows the comparisons
of diurnal variations of both EPIC- and ground-based observed total SWs, averaged, during June, 2015-June, 2019 at 7 SURFRAD
sites under different sky conditions. For clear sky conditions, EPIC-based total SWs capture the diurnal variation well with small
RRMSEs ranging from 10.10%~16.24%. For cloudy sky, EPIC-based total SWs have better performance with RRMSEs smaller
190 than 10.02%. For overcast sky conditions, EPIC-based products overestimate the total SWs with RRMSEs larger than 29.20%. It
is noteworthy that EPIC-derived products show the worst performance for SURFRAD-BOS sites, likely caused by the rugged
terrain around this site. For diffuse SWs shown in **Fig. S1**, clear-sky EPIC-derived estimates have the largest RRMSEs. For direct
SWs, **Fig. S2** shows that overcast-sky EPIC-based estimates have large uncertainties due to their relatively small magnitude. **Fig.**
S3 shows that EPIC-based total PARs perform better than total SWs, especially for clear and cloudy sky conditions.



195

Figure 4. Diurnal variations of both EPIC- and ground-based total SWs, averaged, from June, 2015 to June, 2019 for different sky conditions at 7 SURFRAD sites.

We further analysed the accuracy of our products at both different local times and for different months. **Fig.5** shows that the accuracy of hourly SW/PAR estimates changes with the change in local time. The SW/PAR estimates for nearly local noon have negative biases, larger RMSEs but smaller RRMSEs, where those for early morning or later afternoon have positive biases, smaller RMSEs but larger RRMSEs. However, R^2 values of total SW and PAR estimates are generally larger than 0.7 for all local times. The daily SW/PAR estimates show good accuracy for all months. Total SW and PAR estimates from May to August have positive biases, larger RMSEs but smaller RRMSEs. The R^2 values of SW/PAR estimates show little monthly and seasonal dependency. These results confirm that both EPIC-based hourly and daily products have reliable accuracy, although the magnitudes of SW/PAR, and cloud and atmospheric conditions at different time (i.e. hour, day, and month) may affect the accuracy and uncertainties of these products.

205




Figure 5. Evaluation of EPIC-based hourly SW/PAR estimates at different local hours from 7:00 to 18:00.




Figure 6. Evaluation of EPIC-based daily SW/PAR estimates on different months during the study period.

3.1.3 Spatial distribution of estimated errors of derived products

The hourly and daily total SW estimates show similar spatial patterns in their accuracy statistics composed of biases, RMSEs, RRMSEs and R^2 values (**Figs. 7 and 8**). Polar regions have relatively small RMSE but large RRMSE, due to long-term or frequent ice/snow cover in these regions and a lack of proper accounting for land surface albedo in current products; the island and coastal regions show the worst performance with high bias, large RMSE and low R^2 ; and derived products for most of the continental sites perform well but show large spatial heterogeneities related to different land cover types, climate zones, surface topography, etc. The BSRN-IZA site, a high-mountain station located in Tenerife (Canary Islands, Spain), exhibit high negative bias and large errors and uncertainties, which can be explained by its geographic location in the Tenerife island within Teide volcano area (García et al., 2019), and the particular weather conditions for this area where the clouds affect the lower parts of the island (below 2000 m above sea level) while the sky for upper parts probably remains clear (Urraca et al., 2018). In general, EPIC-derived products have higher accuracy in continental regions with low bias and small RMSE and RRMSE, whereas island or coastal regions show very large bias, and large RMSE and RRMSE values (**Tables 3 and 4**), probably caused by the rapidly changing weather condition and the mixture of land and water in a grid cell (edge effects), which is identical with other studies (Boland et al., 2016; Wang and Pinker, 2009)





Figure 7. Spatial distributions of accuracy statistical metrics for EPIC-based hourly total SW at all ground-based sites. Circles with different colors indicate their different values.



230

Figure 8. Spatial distributions of accuracy statistical metrics for EPIC-based daily total SW at all ground-based sites. Circles with different colors indicate their different values.

Table 1. Evaluation of EPIC-based hourly SW estimates against the ground-based measurements for different regions.

Region	Parameter	Bias (W/m^2)	RMSE (W/m^2)	RRMSE (%)	R^2
Polar	Diffuse SW	4.11	60.22	53.78	0.68
	Direct SW	-11.70	85.37	104.54	0.63
	Total SW	-3.12	65.69	37.34	0.85
Island/Coastal	Diffuse SW	66.65	125.57	78.76	0.54
	Direct SW	-109.11	224.06	68.26	0.58
	Total SW	-32.25	134.13	28.90	0.83
Continental	Diffuse SW	4.43	69.05	50.80	0.62
	Direct SW	-9.27	130.53	50.14	0.76
	Total SW	-1.07	102.97	27.59	0.87

235

Table 2. Evaluation of EPIC-based daily SW estimates against the ground-based measurements for different regions.

Region	Parameter	Bias (W/m^2)	RMSE (W/m^2)	RRMSE (%)	R^2
Polar	Diffuse SW	2.54	26.79	37.45	0.82
	Direct SW	-2.57	40.28	90.98	0.71
	Total SW	0.28	35.96	31.65	0.88
Island/Coastal	Diffuse SW	34.58	48.33	61.66	0.45
	Direct SW	-46.72	84.87	56.68	0.50
	Total SW	-12.07	53.94	23.65	0.67
Continental	Diffuse SW	2.39	21.27	31.91	0.69
	Direct SW	-2.05	39.45	33.70	0.82
	Total SW	0.40	32.80	17.94	0.88

3.2 Globally spatio-temporal patterns of derived products

We investigated the spatial patterns of averaged total SWs during the three whole years of 2016-2018 for different seasons : 1) Spring, March, April, and May (MAM); 2) Summer, June, July, and August (JJA); 3) Autumn, September, October, and November, (SON); 4) Winter, December, January, and February (DJF). **Fig. 9(a-d)** show the EPIC-based products reflect the heterogeneous spatial distributions and track the globally seasonal variations that are mainly due to the sun angle variations. They also have a consistent pattern when compared with the CERES-derived products (**Fig. 9(e-f)**). **Fig. S4** shows that EPIC- and CERES-derived estimates have small differences over most regions, especially in Spring, Autumn and Winter, but some large discrepancies are observed in the Tibetan Plateau due to the frequent ice/snow cover and in the Congo basin due to the complex cloud and atmospheric conditions. However, our EPIC-derived products can reveal more spatial details than CERES-derived estimates due to their higher spatial resolution.






Figure 9. Global distributions of EPIC- and CERES-derived total SW fluxes for different seasons during the study period, 2016-2018.

Fig. 10 shows the temporal variations of both EPIC- and CERES-based monthly total SWs for the land surface of globe, northern and southern hemispheres during June, 2015 to June, 2019. All EPIC- and CERES-based products show similar temporal variations. From Autumn (SON) to next Spring (MAM), EPIC-based global SWs coincide well with CERES-derived ones, whereas in Summer (JJA), EPIC-based global SWs are lower than CERES-derived ones, due to the differences in northern hemisphere. The differences in spatial resolution, driving data, retrieval models/algorithms, etc. contribute to these discrepancies. **Fig. 11** displays the temporal variations of daily zonal averaged total SWs and PAR products. EPIC- and CERES-based estimates present highly consistent latitude-gradient distributions and temporal variations. **Fig. S5** shows the differences between EPIC- and CERES-based SW/PAR estimates. Overall, total SW and PAR and their direct and diffuse components have small differences, but the direct and diffuse components of SW show relatively large differences in the northern hemisphere, possibly due to the underestimation of CERES-based direct components and overestimation of CERES-derived diffuse components (Hao et al., 2019).



265

Figure 10. Temporal variations of EPIC- and CERES-based monthly total SW for the land surface of global, northern- and southern-hemispheres.



270

Figure 11. Temporal variations of EPIC- and CERES-based zonally-averaged daily total SW and PAR for global land areas. The vertical dark bars in (a, c) are due to the missing data.

4 Discussion

275

The proposed new SW/PAR products presented here make full use of the unique advantages of DSCOVR/EPIC, an instrument that observes nearly the entire sunlit areas of the Earth, from pole to pole, every 1~2 hours. These promising products have some unique attributes: 1) show high correlations with the ground station observations; 2) present reasonable and identical spatio-temporal SW/PAR patterns but more spatial details when compared to CERES products; and 3) capture accurately the diurnal cycles of SW/PAR. In addition, they are based on a single instrument, EPIC, and thus avoid the sensor-to-sensor differences and inconsistencies inherent in multi-source datasets. The DSCOVR/EPIC science team is continuing to refine their algorithms and upgrade the product quality, and we plan to update our products accordingly.

280

We expect and hope these products will have multiple uses in diverse earth science communities. When combined with the DSCOVR/EPIC-derived vegetation data, our globally high-accuracy products can be used to understand the diurnal variabilities and underlying processes of photosynthesis and evapotranspiration for terrestrial ecosystems. By virtue of the decomposition of the direct and diffuse PARs, our products are expected to improve the estimates of ecosystem photosynthesis capacity and primary production. Our products may also be valuable for site selection for solar power production and solar energy management.

Some sources of uncertainties probably affect the accuracy and reliability of our products (see **Figs 2 and 3**). There are some
285 geolocation, inter-calibration and misregistration issues in the current EPIC L1B version 2 products (Molina García et al., 2019),
which will be improved in the version 3 products (announced in the DSCOVR 2019 Science Team Meeting, Greenbelt, MD) in
future. The current EPIC L2 atmospheric and cloud products have large uncertainties or gaps in the ice/snow covered regions and
when the solar angle is large ($>70^\circ$) (Yang et al., 2019;Herman et al., 2018;Xu et al., 2017). We used the CERES data to fill the
gaps, especially in the polar regions. To allow users maximum flexibility, we also provided quality flags that indicate whether
290 pixels are successfully retrieved or gap-filled in the derived products. Current algorithms/models neglect the impact of water vapor
and land surface albedo, which could lead to some additional biases and uncertainties. This problem will be addressed through the
combination with high-quality EPIC-view water vapor products such as the EPIC-view Multi-Sensor Global Cloud and Radiance
Composites (Khlopenkov et al., 2017) and the development of land surface albedo products based on the EPIC Multi-Angle
Implementation of Atmospheric Correction (MAIAC) products (Hao et al., 2019). We also did not account for the impacts of
295 spatial mismatch between ground-based observation and EPIC-based data. In the future, we will collect long-term, high-quality
and widely-distributed ground-based datasets to improve and evaluate our products.

The spatial resolution of our current products is relatively coarse ($0.1^\circ \times 0.1^\circ$). The effective spatial resolution of the original
EPIC image is relevant to the observing angle, and is about 10km at nadir (near the centre of the image) and 20km at 60° (Marshak
et al., 2018). Higher spatial resolution SW/PAR products are desired for mapping of carbon and water fluxes as well as solar energy
300 assessment and operation, especially for islands and coastal regions (see **Tables 3 and 4**). Spatial scale mismatch between land
surface properties and derived SW/PAR estimates can limit the applications of our products (Ryu et al., 2018). We suggest that
spatial downscaling techniques can be used to improve our proposed products (Wang et al., 2019), especially for rugged terrain
(e.g. the Tibetan Plateau), where topographic effects (e.g. varied elevation, rotation of solar geometry, shadowing, terrain occlusion,
and multi-scattering) on SW/PAR should be considered and assessed (Zhang et al., 2019;Hao et al., 2018b;Hao et al., 2018a). We
305 believe such analysis/assessment can benefit greatly next generation mission of deep-space satellites/sensors such as
DSCOVR/EPIC, for optimizing/balancing the trade-off between data amount (spatial/temporal resolution) and transmission time,
perhaps leading to improved spatio-temporal resolution of future data products from such missions.

Finally, one shortcoming of current products is the relatively short period of the derived data records which cover only 4 years.
Such record length is not adequate to detect any globally long-term trends. A feasible solution is to merge DSCOVR/EPIC products
310 with reanalysis data/products to produce globally continuous, consistent and long-term SW/PAR datasets, through correcting the
reanalysis data based on satellite data (Feng and Wang, 2018). With the increasing record length of EPIC data, it is anticipated that
the temporal coverage of our proposed products will be also extended to support the long-term analysis in the future.

5 Data availability

Both the derived hourly and daily DSCOVR/EPIC-based global SW/PAR products are available at the DataHub
315 (<https://doi.org/10.25584/1595069>, Hao et al., 2020), Pacific Northwest National Laboratory (PNNL). The hourly data are grouped
by day in distinct NetCDF files, which are named as “EPIC_SW_PAR_Hourly_yyyymmdd.nc” where “yyyy”, “mm”, and “dd”
denote year, month, and day (UTC time). The daily data are grouped by month in distinct NetCDF files, which are named as
“EPIC_SW_PAR_Daily_yyyymm.nc” where “yyyy”, and “mm” denote year and month (UTC time). Each NetCDF file contains
latitude, longitude, time, diffuse SW, direct SW, diffuse PAR, direct PAR, and the corresponding quality flags which indicate
320 whether the pixel is gap-filled or not. The scale factor for the direct and diffuse SW/PAR is 0.1. The total SW/PAR estimates can

be calculated by combining the direct and diffuse components. The information about the version, creation date, reference, contact mails, and other comments are also included in the file.

6 Conclusions

This paper presents the first globally hourly and daily SW/PAR products, with a spatial resolution of $0.1^{\circ} \times 0.1^{\circ}$ for the period of
325 June, 2015~June, 2019 based on the DSCOVR/EPIC observations. The newly developed products are the first of their kind because
of high temporal frequency (hourly) and global coverage at a spatial resolution of $0.1^{\circ} \times 0.1^{\circ}$, only based on a single instrument,
DSCOVR/EPIC. We evaluated the EPIC-derived products against the globally-distributed ground-based data, and analysed and
compared the spatio-temporal variations of the proposed products with the well-characterized and widely-used CERES products.
330 EPIC-derived SW/PAR estimates and their direct and diffuse components show good consistencies with the globally-distributed
ground-based observations. The EPIC-derived products capture accurately the diurnal variabilities of SW/PAR under different sky
conditions. The comparisons with CERES data indicate that the developed products reflect complex spatial heterogeneities and
capture substantial seasonal variabilities of SW/PAR effectively with the same temporal resolution of hourly but higher spatial
resolution. The promising products offer an invaluable resource for solar photovoltaic applications and understanding and
exploring the diurnal cycles of terrestrial water, carbon, and energy fluxes at various temporal and spatial scales. We plan to update
335 our proposed products as additional EPIC observations become available, and with further improvements of the record length of
EPIC data and algorithm refinements that are planned by the EPIC/DSCVR science team in the future.

Author contributions

CM and HD designed the study. HD and CM produced the products, analyzed the results, and drafted the original paper. All authors
contributed to the analysis and interpretation of the results, and to improving this paper.

340 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

The CERES and CAVE data were obtained from the NASA Langley Research Center Atmospheric Science Data Center. We thank
the entire DSCOVR/EPIC team for the powerful L2 products and especially Dr. Pam Mlynczak for valuable information on CERES
345 Edition 4.1 products. We also thank Dr. Ben Bond-Lamberty for the valuable suggestions, and Emmanuel Bonilla for his help on
sharing the data on the DataHub of PNNL.

Financial support

This work was supported by a Laboratory Directed Research and Development project sponsored by the Pacific Northwest National
Laboratory (PNNL) of the U.S. Department of Energy. Dalei Hao was sponsored by China Scholarship Council. The research was
350 performed using resources available through Research Computing at PNNL. PNNL is operated by Battelle for the U.S. Department
of Energy under Contract DE-AC05-76RL01830.

References

- Augustine, J. A., DeLuise, J. J., and Long, C. N.: SURFRAD—A national surface radiation budget network for atmospheric research, *Bulletin of the American Meteorological Society*, 81, 2341-2358, 2000.
- 355 Boland, J., David, M., and Lauret, P.: Short term solar radiation forecasting: Island versus continental sites, *Energy*, 113, 186-192, 2016.
- Burt, J., and Smith, B.: Deep space climate observatory: The DSCOVR mission, 2012 IEEE Aerospace Conference, 2012, 1-13,
- Chen, M., and Zhuang, Q.: Evaluating aerosol direct radiative effects on global terrestrial ecosystem carbon dynamics from 2003 to 2010, *Tellus B: Chemical and Physical Meteorology*, 66, 21808, 2014.
- 360 Damm, A., Elbers, J., Eler, A., Gioli, B., Hamdi, K., Hutjes, R., Kosvancova, M., Meroni, M., Miglietta, F., and Moersch, A.: Remote sensing of sun-induced fluorescence to improve modeling of diurnal courses of gross primary production (GPP), *Global Change Biology*, 16, 171-186, 2010.
- Driemel, A., Augustine, J., Behrens, K., Colle, S., Cox, C., Cuevas-Agulló, E., Denn, F. M., Duprat, T., Fukuda, M., and Grobe, H.: Baseline Surface Radiation Network (BSRN): structure and data description (1992-2017), *Earth System Science Data*, 10, 1491-1501, 2018.
- Farquhar, G. D., and Roderick, M. L.: Pinatubo, diffuse light, and the carbon cycle, *Science*, 299, 1997-1998, 2003.
- 365 Feng, F., and Wang, K.: Merging satellite retrievals and reanalyses to produce global long-term and consistent surface incident solar radiation datasets, *Remote Sensing*, 10, 115, 2018.
- García, R. D., Cuevas, E., Ramos, R., Cachorro, V. E., Redondas, A., and Moreno-Ruiz, J. A.: Description of the Baseline Surface Radiation Network (BSRN) station at the Izaña Observatory (2009–2017): measurements and quality control/assurance procedures, *Geoscientific Instrumentation, Methods and Data Systems*, 8, 77-96, 2019.
- 370 Gu, L., Baldocchi, D., Verma, S. B., Black, T., Vesala, T., Falge, E. M., and Dowty, P. R.: Advantages of diffuse radiation for terrestrial ecosystem productivity, *Journal of Geophysical Research: Atmospheres*, 107, ACL 2-1-ACL 2-23, 2002.
- Hao, D., Wen, J., Xiao, Q., Wu, S., Lin, X., Dou, B., You, D., and Tang, Y.: Simulation and analysis of the topographic effects on snow-free albedo over rugged terrain, *Remote Sensing*, 10, 278, 2018a.
- Hao, D., Wen, J., Xiao, Q., Wu, S., Lin, X., You, D., and Tang, Y.: Impacts of DEM Geolocation Bias on Downward Surface Shortwave Radiation Estimation Over Clear-Sky Rugged Terrain: A Case Study in Dayekou Basin, China, *IEEE Geoscience and Remote Sensing Letters*, 16, 10-14, 2018b.
- 375 Hao, D., Asrar, G. R., Zeng, Y., Zhu, Q., Wen, J., Xiao, Q., and Chen, M.: Estimating hourly land surface downward shortwave and photosynthetically active radiation from DSCOVR/EPIC observations, *Remote Sensing of Environment*, 232, 111320, 2019.
- Hao, D., Chen, M., Asrar, G. R., Zeng, Y., Zhu, Q., Wen, J., and Xiao, Q.: A global DSCOVR/EPIC-based hourly/daily shortwave radiation/PAR dataset, DataHub for Pacific Northwest National Laboratory, <https://doi.org/10.25584/1595069>, 2020.
- 380 Herman, J., Huang, L., McPeters, R., Ziemke, J., Cede, A., and Blank, K.: Synoptic ozone, cloud reflectivity, and erythemal irradiance from sunrise to sunset for the whole earth as viewed by the DSCOVR spacecraft from the earth–sun Lagrange 1 orbit, *Atmos. Meas. Tech.*, 11, 177-194, 10.5194/amt-11-177-2018, 2018.
- Huang, G., Li, Z., Li, X., Liang, S., Yang, K., Wang, D., and Zhang, Y.: Estimating surface solar irradiance from satellites: Past, present, and future perspectives, *Remote Sensing of Environment*, 233, 111371, 2019.
- 385 Kato, S., Rose, F. G., Rutan, D. A., Thorsen, T. J., Loeb, N. G., Doelling, D. R., Huang, X., Smith, W. L., Su, W., and Ham, S.-H.: Surface irradiances of edition 4.0 clouds and the earth's radiant energy system (CERES) energy balanced and filled (EBAF) data product, *Journal of Climate*, 31, 4501-4527, 2018.
- 390 Khahro, S. F., Tabbassum, K., Talpur, S., Alvi, M. B., Liao, X., and Dong, L.: Evaluation of solar energy resources by establishing empirical models for diffuse solar radiation on tilted surface and analysis for optimum tilt angle for a prospective location in southern region of Sindh, Pakistan, *International Journal of Electrical Power & Energy Systems*, 64, 1073-1080, 2015.
- Khlopenkov, K., Duda, D., Thieman, M., Minnis, P., Su, W., and Bedka, K.: Development of multi-sensor global cloud and radiance composites for earth radiation budget monitoring from DSCOVR, *Remote Sensing of Clouds and the Atmosphere XXII*, 2017, 104240K,
- 395 Korany, M., Boraiy, M., Eissa, Y., Aoun, Y., Wahab, M. A., Alfaro, S., Blanc, P., El-Metwally, M., Ghedira, H., and Hungershoefer, K.: A database of multi-year (2004–2010) quality-assured surface solar hourly irradiation measurements for the Egyptian territory, 2016.
- Letu, H., Yang, K., Nakajima, T. Y., Ishimoto, H., Nagao, T. M., Riedi, J., Baran, A. J., Ma, R., Wang, T., and Shang, H.: High-resolution retrieval of cloud microphysical properties and surface solar radiation using Himawari-8/AHI next-generation geostationary satellite, *Remote Sensing of Environment*, 239, 111583, 2020.
- 400 Li, X., Al-Yaari, A., Schwank, M., Fan, L., Frappart, F., Swenson, J., and Wigneron, J.-P.: Compared performances of SMOS-IC soil moisture and vegetation optical depth retrievals based on Tau-Omega and Two-Stream microwave emission models, *Remote Sensing of Environment*, 236, 111502, 2020.
- Liang, S., Wang, K., Zhang, X., and Wild, M.: Review on estimation of land surface radiation and energy budgets from ground measurement, remote sensing and model simulations, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 3, 225-240, 2010.
- 405 Liou, K., Gu, Y., Leung, L., Lee, W., and Fovell, R.: A WRF simulation of the impact of 3-D radiative transfer on surface hydrology over the Rocky Mountains and Sierra Nevada, *Atmospheric Chemistry and Physics*, 13, 11709-11721, 2013.
- Loeb, N. G., Doelling, D. R., Wang, H., Su, W., Nguyen, C., Corbett, J. G., Liang, L., Mitrescu, C., Rose, F. G., and Kato, S.: Clouds and the earth's radiant energy system (CERES) energy balanced and filled (EBAF) top-of-atmosphere (TOA) edition-4.0 data product, *Journal of Climate*, 31, 895-918, 2018.
- 410 Marshak, A., Herman, J., Adam, S., Karin, B., Carn, S., Cede, A., Geogdzhayev, I., Huang, D., Huang, L.-K., and Knyazikhin, Y.: Earth observations from DSCOVR EPIC instrument, *Bulletin of the American Meteorological Society*, 99, 1829-1850, 2018.
- Mercado, L. M., Belloouin, N., Sitch, S., Boucher, O., Huntingford, C., Wild, M., and Cox, P. M.: Impact of changes in diffuse radiation on the global land carbon sink, *Nature*, 458, 1014, 2009.
- Molina García, V., Sasi, S., Efremenko, D. S., and Loyola, D.: Improvement of EPIC/DSCOVR Image Registration by Means of Automatic Coastline Detection, *Remote Sensing*, 11, 1747, 2019.
- 415 Ohmura, A., Dutton, E. G., Forgan, B., Fröhlich, C., Gilgen, H., Hegner, H., Heimo, A., König-Langlo, G., McArthur, B., and Müller, G.: Baseline Surface Radiation Network (BSRN/WCRP): New precision radiometry for climate research, *Bulletin of the American Meteorological Society*, 79, 2115-2136, 1998.

- Pinker, R., Zhang, B., and Dutton, E.: Do satellites detect trends in surface solar radiation?, *Science*, 308, 850-854, 2005.
- 420 Raptis, P., Kazadzis, S., Psiloglou, B., Kouremeti, N., Kosmopoulos, P., and Kazantzidis, A.: Measurements and model simulations of solar radiation at tilted planes, towards the maximization of energy capture, *Energy*, 130, 570-580, 2017.
- Roderick, M. L., and Farquhar, G. D.: The cause of decreased pan evaporation over the past 50 years, *science*, 298, 1410-1411, 2002.
- Rutan, D., Rose, F., Smith, N., and Charlock, T.: Validation data set for CERES surface and atmospheric radiation budget (SARB), *WCRP/GEWEX Newsletter*, 11, 11-12, 2001.
- 425 Rutan, D. A., Kato, S., Doelling, D. R., Rose, F. G., Nguyen, L. T., Caldwell, T. E., and Loeb, N. G.: CERES synoptic product: Methodology and validation of surface radiant flux, *Journal of Atmospheric and Oceanic Technology*, 32, 1121-1143, 2015.
- Ryu, Y., Jiang, C., Kobayashi, H., and Detto, M.: MODIS-derived global land products of shortwave radiation and diffuse and total photosynthetically active radiation at 5 km resolution from 2000, *Remote Sensing of Environment*, 204, 812-825, 2018.
- Sweerts, B., Pfenninger, S., Yang, S., Folini, D., Van der Zwaan, B., and Wild, M.: Estimation of losses in solar energy production from air pollution in China since 1960 using surface radiation data, *Nature Energy*, 4, 657-663, 2019.
- 430 Urraca, R., Huld, T., Gracia-Amillo, A., Martinez-de-Pison, F. J., Kaspar, F., and Sanz-Garcia, A.: Evaluation of global horizontal irradiance estimates from ERA5 and COSMO-REA6 reanalyses using ground and satellite-based data, *Solar Energy*, 164, 339-354, 2018.
- Van Heerwaarden, C. C., Vilà-Guerau de Arellano, J., Gounou, A., Guichard, F., and Couvreux, F.: Understanding the daily cycle of evapotranspiration: A method to quantify the influence of forcings and feedbacks, *Journal of Hydrometeorology*, 11, 1405-1422, 2010.
- 435 Wang, D., Liang, S., Zhang, Y., Gao, X., Brown, M. G., and Jia, A.: A New Set of MODIS Land Products (MCD18): Downward Shortwave Radiation and Photosynthetically Active Radiation, *Remote Sensing*, 12, 168, 2020.
- Wang, H., and Pinker, R.: Shortwave radiative fluxes from MODIS: Model development and implementation, *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- Wang, W., Yin, G., Zhao, W., Wen, F., and Yu, D.: Spatial Downscaling of MSG Downward Shortwave Radiation Product Under Clear-Sky Condition, *IEEE Transactions on Geoscience and Remote Sensing*, 2019.
- 440 Wang, X., Wu, J., Chen, M., Xu, X., Wang, Z., Wang, B., Wang, C., Piao, S., Lin, W., and Miao, G.: Field evidences for the positive effects of aerosols on tree growth, *Global change biology*, 24, 4983-4992, 2018.
- Wielicki, B. A., Barkstrom, B. R., Harrison, E. F., Lee III, R. B., Smith, G. L., and Cooper, J. E.: Clouds and the Earth's Radiant Energy System (CERES): An earth observing system experiment, *Bulletin of the American Meteorological Society*, 77, 853-868, 1996.
- 445 Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C. N., Dutton, E. G., Forgan, B., Kallis, A., Russak, V., and Tsvetkov, A.: From dimming to brightening: Decadal changes in solar radiation at Earth's surface, *Science*, 308, 847-850, 2005.
- Wyser, K., O'Hirok, W., and Gautier, C.: A simple method for removing 3-D radiative effects in satellite retrievals of surface irradiance, *Remote sensing of environment*, 94, 335-342, 2005.
- 450 Xu, X., Wang, J., Wang, Y., Zeng, J., Torres, O., Yang, Y., Marshak, A., Reid, J., and Miller, S.: Passive remote sensing of altitude and optical depth of dust plumes using the oxygen A and B bands: First results from EPIC/DSCOVR at Lagrange-1 point, *Geophysical Research Letters*, 44, 7544-7554, 2017.
- Yang, B., Knyazikhin, Y., Möttus, M., Rautiainen, M., Stenberg, P., Yan, L., Chen, C., Yan, K., Choi, S., and Park, T.: Estimation of leaf area index and its sunlit portion from DSCOVR EPIC data: Theoretical basis, *Remote sensing of environment*, 198, 69-84, 2017.
- Yang, Y., Meyer, K., Wind, G., Zhou, Y., Marshak, A., Platnick, S., Min, Q., Davis, A. B., Joiner, J., and Vasilkov, A.: Cloud products from the Earth Polychromatic Imaging Camera (EPIC): algorithms and initial evaluation, *Atmospheric Measurement Techniques*, 12, 2019.
- 455 Zhang, S., Li, X., She, J., and Peng, X.: Assimilating remote sensing data into GIS-based all sky solar radiation modeling for mountain terrain, *Remote Sensing of Environment*, 231, 111239, 2019.
- Zhang, X., Liang, S., Zhou, G., Wu, H., and Zhao, X.: Generating Global LAnd Surface Satellite incident shortwave radiation and photosynthetically active radiation products from multiple satellite data, *Remote Sensing of Environment*, 152, 318-332, 2014.
- 460 Zhao, L., Lee, X., and Liu, S.: Correcting surface solar radiation of two data assimilation systems against FLUXNET observations in North America, *Journal of Geophysical Research: Atmospheres*, 118, 9552-9564, 2013.