# Two decades of distributed global radiation time series across a mountainous semiarid area (Sierra Nevada, Spain)

3 Cristina Aguilar<sup>1</sup>, Rafael Pimentel<sup>1</sup>, María J. Polo<sup>1</sup>

<sup>1</sup>Fluvial Dynamics and Hydrology Research Group, Andalusian Institute of Earth System Research, University of Cordoba,
 5 Cordoba, Spain

6 *Correspondence to*: Cristina Aguilar (caguilar@uco.es)

7 Abstract. The main drawback of the reconstruction of high resolution distributed global radiation (Rg) time series in 8 mountainous semiarid environments is the common lack of station-based solar radiation registers. This work presents nineteen 9 years (2000-2018) of high spatial resolution (30 m) daily, monthly, and annual global radiation maps derived using the GIS-10 based model proposed by Aguilar et al. (2010) in a mountainous area in southern Europe: Sierra Nevada (SN) Mountain Range 11 (Spain). The model was driven by in situ daily global radiation measurements, from sixteen weather stations with historical 12 records in the area, a 30 m digital elevation model and 240 cloud-free Landsat images. The applicability of the modeling scheme was validated against daily global radiation records at the weather stations. Mean RMSE values of 2.63 MJ m<sup>-2</sup> dav<sup>-1</sup> 13 and best estimations on clear-sky days were obtained. Daily Rg at weather stations revealed greater variations in the maximum 14 15 values but no clear trends with altitude in any of the statistics. However, at the monthly and annual scales, there is an increase 16 in the high extreme statistics with the altitude of the weather station, especially above 1500 m a.s.l. Monthly R<sub>g</sub> maps showed significant spatial differences of up to 200 MJ m<sup>-2</sup> month<sup>-1</sup> that clearly followed the terrain configuration. July and December 17 were clearly the months with the highest and lowest values of  $R_{\sigma}$  received and the highest scatter in the monthly  $R_{\sigma}$  values was 18 19 found in the spring and fall months. The monthly  $R_{\nu}$  distribution was highly variable along the study period (2000-2018). Such 20 variability, especially in the wet season (October-May), determined the inter annual differences of up to 800 MJ m<sup>-2</sup> year<sup>-1</sup> in 21 the incoming global radiation in SN. The time series of the surface global radiation datasets here provided can be used to 22 analyze inter-annual and seasonal variation characteristics of the global radiation received in SN with high spatial detail (30 23 m). They can also be used as cross-validation reference data for other global radiation distributed datasets generated in SN 24 with different spatio-temporal interpolation techniques. Daily, monthly, and annual datasets in this study are available at 25 https://doi.pangaea.de/10.1594/PANGAEA.921012 (Aguilar et al., 2021).

- 26
- 27
- 28
- 29

# 30 1 Introduction

31 High mountain areas in semiarid environments present singular characteristics due to the continuous interaction of alpine 32 conditions in the summits with the surrounding semiarid climate. They play a key role as water providers during the warm and 33 dry season when they often constitute the only water source for many rivers. Here, water fluxes from the snowpacks show a 34 shift from the predominant partition between snowmelt and sublimation usually found in colder and wetter climates on an 35 annual and seasonal basis (Herrero and Polo, 2016). This shift is caused by the radiation balance that enhances sublimation 36 during cold and dry periods and intense snowmelt rates during late winter and spring in these areas (McDonell et al., 2013; 37 Liu et al., 2019). However, weather stations are not always equipped to monitor the global radiation nor their components and, 38 moreover, they are seldom found in high altitudes, especially over 1500 m a.s.l., which makes it difficult to accurately assess 39 not only the solar radiation temporal regime but also the spatial patterns of solar radiation fields in high mountain areas. This 40 impacts the availability of data for studies in mountains dealing with climate and hydrology, global warming, ecosystem 41 services provided by the snow areas, and environmental and social and economic impacts on-site and downstream (Yang et 42 al., 2010; Liu et al., 2012a; Tang et al., 2019). It is not surprising that many mountain regions are identified as biodiversity 43 hotspots around the world, with Mediterranean and other semiarid to arid regions being highly represented (Myers et al., 2000; 44 O'Farrell et al., 2010; Hewitt, 2011; Pauli et al., 2012).

45 There are several research papers on solar radiation estimations from routine ground-based observations in high altitude regions 46 (Dubayah and van Katwijk, 1992; Dubayah, 1994; Tovar et al., 1995; Oliphant et al., 2003; Tovar-Pescador et al., 2006; Yang 47 et al., 2006, 2010; Batllés et al., 2008; Bosch et al., 2008; Sheng et al., 2009; Aguilar et al., 2010; Mamassis et al., 2012; Chen 48 et al., 2013; Zhang et al., 2020). All of them insist on the need to consider topographic effects and advise of the errors that 49 simple interpolation/extrapolation techniques can create. Radiation data obtained from a dense and properly-maintained 50 weather station network in mountainous areas are rarely available and therefore, modeling techniques need to be applied. Liu 51 et al. (2012a) state that the most difficult issue in solar radiation modeling in data sparse regions is cloud accounting, due to 52 the rapid spatially and temporally changing weather conditions and the three-dimensional structure of clouds. This complexity 53 adds to the heterogeneity resulting from shadowing and reflection due to steep topography (Dubayah, 1992; Batllés et al., 54 2008; Mamassis et al., 2012; Chen et al., 2013; Zhang et al., 2019, 2020).

According to Dubayah and Rich (1995), as solar radiation models become more complex, they can be more difficult to use, mainly because of the requirement for additional input data. In fact, the complexity of physically-based solar radiation formulations for topography and the lack of the data needed to drive such formulations led in the past to the lack of suitable modeling tools (Dubayah, 1994). Thus, it is important that the models allow for some flexibility regarding the component of radiation calculated and the input data needed.

Excluding traditional interpolation methods there are two major methods for solar radiation modeling, namely, satellite-derived solar radiation estimates, and Geographic Information Systems (GIS)-based solar radiation models. Satellite-derived solar radiation models provide a wide spatial and temporal coverage, but low spatial resolution when dealing with pixels with a strong topographic gradient. By contrast, GIS-based models calculate the incoming solar radiation for each cell of a digital elevation model (DEM) and allow for higher spatial resolutions including topographic effects. In the past decades, several models based on GIS have been proposed (e.g., Dubayah and Rich, 1995; Fu and Rich, 2000a, 2002; Wilson and Gallant, 2000; Goldberg and Häntzschel, 2002; Sùri and Hofierka, 2004; Liu et al., 2012a; Zhang et al., 2019, 2020). Required input data include digital elevation values and atmospheric attenuation parameters that are commonly estimated from ground-based measurements and/or satellite data (Dubayah, 1994).

69 The aim of this study was to generate the spatiotemporal distribution of global solar radiation in a high mountain semiarid area 70 in southern Spain with a modeling scheme that reconstructs time map series from the usually available weather datasets. For 71 this purpose, a GIS-based topographic solar radiation model (Aguilar et al., 2010) was applied in Sierra Nevada (SN) (Spain), 72 a high mountain range running west-east parallel to the Mediterranean coastline with influence from both the sea and the 73 African continent to the South, and the continental conditions to the North. The accuracy of solar radiation estimates by the 74 model were evaluated in terms of the error in the approximation to observed data. This study site is a high-value environmental 75 area declared Biosphere Reserve by UNESCO in 1986 due to the exceptional presence of endemisms (Heywood, 1995; Blanca 76 et al., 1998; Anderson et al., 2011; Cañadas et al., 2014). Besides, SN is also included in the Global Change Observatories 77 Network given its singular location between two seas and two continents, and its extreme topographic gradients (Bonet-García 78 et al., 2015).

79 This paper presents 19 years of daily, monthly, and annual solar radiation maps with high resolution (30 m) over SN. The huge 80 number of members involved in the management of this area make this information valuable in different fields, such as: 81 hydrology, crucial role of energy budget in the hydrological cycle over this area; ecology, ecological communities' behaviour 82 and development clearly link with the amount of energy available; production systems downstream, as hydropower facilities 83 and traditional to tropical crop systems from the top to downhills. Besides, these data sets directly contribute, or are relevant 84 for many studies that could do so, to two of the 23 Unsolved Problems in Hydrology (UPH) recently posed by Blöschl et al. 85 (2019) in a participatory analytical discussion among the scientific community: UPH 16 "How can we use innovative technologies to measure surface and subsurface properties, states and fluxes at a range of spatial and temporal scales?" and 86 UPH 5 "What causes spatial heterogeneity and homogeneity in runoff, evaporation, subsurface water and material fluxes 87 88 (carbon and other nutrients, sediments), and in their sensitivity to their controls (e.g. snowfall regime, aridity, reaction 89 coefficients)?".

### 90 2 Study site

The Sierra Nevada mountain range (SN) is located 35 km north from the Mediterranean Sea (Fig. 1) and constitutes a mountainous area of the Natura 2000 network. Elevations rise up from 262 m a.s.l. to 3479 m a.s.l. in a 4583.72 km<sup>2</sup> area that runs parallel to the sea. High altitudinal gradients are representative of the area, with variation in elevation of about 3400 m in less than 40 km of horizontal distance and a mountain climate in the summits surrounded by Mediterranean climate in the 95 lower areas. Thus, the interaction of such conditions creates a strong heterogeneity in terms of soil types, landforms and 96 vegetation species that determine a complex hydrological response in the area and many endemic species (Heywood, 1995; 97 Blanca et al., 1998; Anderson et al., 2011). The rainfall regime is highly variable, even in consecutive years, with annual 98 cumulative values in the period (1960-2000) that range between 200 mm in dry years to 1000 mm in wet years, with an average 99 value of 510 mm (Pérez-Palazón et al., 2015). Temperature regime is also heterogeneous, with values of 26, 12.5 and 0.4 °C, 100 for maximum, mean, and minimum daily temperature in the same period.

101 The snow presence becomes relevant from November above 2000 m a.s.l. and extends up to spring with conditions that make 102 it possible the activity of a major ski resort in the area. However, in some winters, mild episodes can be found in January and 103 February that melt most of the snow much earlier than the mean end of the snow season in the area (Herrero et al., 2009; 104 Herrero and Polo, 2012). Because of its singular characteristics and fragile environment, Sierra Nevada receives international 105 recognition as a Biosphere Reserve (1986), a National Park (1999), an Important Bird Area (2003), a Special Area of 106 Conservation (2012) and one of the International Global Change Observatories in Mountain Areas. These environmental 107 protection figures together with the different and numerous members involved in the management of such a unique area have 108 determined the strong effort in data collection in the last years to advance in the knowledge of the different aspects that 109 determine the dynamics of this natural system. Moreover, global warming impacts threaten the environmental values of this 110 system but also the associated ecosystem services and social and economic activities due to the estimated shift of the snowfall 111 regime (Pérez-Palazón et al., 2018).



113 Figure 1. Location of the study site in southern Spain (left). Digital Elevation Model (DEM) and weather stations in Sierra Nevada

114 (SN) (right). The numbers correspond to the station codes.

115 3 Data

### 116 **3.1 Input data**

A digital elevation model (DEM) with 30 m spatial resolution and 1m vertical precision was used in this study (Fig. 1). The DEM was provided by the Andalusian regional administration and it was generated by digital stereo correlation of aerial photographs of the Spanish National Plan of Orthophotography. The DEM is used to calculate the slope, aspect, sky view factor and terrain configuration maps that are used in the modeling process (Dozier and Frew, 1990).

121 Meteorological input data are the longest available in-situ daily global radiation ( $R_{go}$ ) of 16 weather stations over the area (Fig. 122 1 and Table 1). The extent of the records in all weather stations ( $N_0$  in Table 1) was considered long enough to carry out the 123 evaluation process dating from February 2000 for the oldest station (608 in Table 1). 12 out of the 16 weather stations are 124 located above 1500 m a.s.l. and 7 of them above 2000 m a.s.l. (Fig. 1). The stations belong to four different organizations: The 125 Department of Agriculture, Fisheries and Environment of the Andalusian Government (601-608 in Table 1), the Water and 126 Environment Agency (1001 and 1002 in Table 1), the National Parks Organization (853-860 in Table 1) and the Guadalfeo 127 Network (802-804 in Table 1) described in Polo et al. (2019). Pyranometers used to collect the data were of different natures 128 but all of them with a characteristic range of around  $0.35 \sim 1.1$  um: Skye SP1110 (stations 601, 602, 604 and 608), Kipp & 129 Zonen SP-Lite pyranometer (station 802), HuksefluxLP02 (station 803), HuksefluxNR01 (stations 1001, 1002 and 804) and 130 Middleton Net Solar CNR1 (stations 853, 854, 855, 857, 858, 859 and 860).

131 In order to generate the complete global radiation data series for the whole-time span (01/02/2000-31/12/2018) we first apply 132 a quality-control check to the recorded data at the weather stations.

### 133 **3.2 Data quality control**

Numerous studies on quality control of measured solar radiation data can be found in the literature (Geiger et al., 2002; Younes et al., 2005; Moradi, 2009; Journée and Bertrand, 2011). Compared to other meteorological variables, solar radiation measurement is more prone to errors (Moradi, 2009). Younes et al. (2005) state two main sources of errors related to in situ measurement of solar radiation: those related to equipment and uncertainty and operational errors. Thus, prior to any computation two basic screenings were applied to recorded daily global radiation data to discard suspicious records associated with equipment and operational errors (Younes et al., 2005).

- Observed daily global radiation (R<sub>go</sub>) must be between the daily extraterrestrial radiation (R<sub>ext</sub>) and a minimum 3% of R<sub>ext</sub> (Geiger et al., 2002; Moradi, 2009).
- 142 2. Observed daily global radiation ( $R_{go}$ ) must be lower than the clear daily global radiation ( $R_{gcs}$ ) observed under a 143 highly transparent clear sky (Wu et al., 2007).  $R_{gcs}$  values were calculated with the model developed by Ineichen and
- Perez (2002) and the parameterization of Kasten and Young (1989) for the air mass. More detail regarding the equation as well as its parameters can be found in Aguilar et al. (2010).
- 146 The excluded values from these tests did not reach 1% of the data at any weather station.

147 A third quality control screening was applied following Younes et al. (2005) to detect erroneous data due to operational errors 148 related with particularities of weather stations in high altitudes (e.g., shadows, impacts of snow, mechanical failures, etc.). 149 They suggest a semi-automatic procedure that allows the creation of an expectancy envelope in the clearness index (CI)-diffuse 150 to global irradiance ratio (k) domain to reject data too obviously erroneous. The CI data range is divided into bands of equal 151 width, within which the mean and standard deviation of the k values,  $\mu_k$  and  $\sigma_k$ , are calculated. The top and bottom boundary 152 shapes are identified by fitting two polynomials through the points  $\mu_k \pm b\sigma_k$  limited between 0 and 1 to respect the physical 153 range of the CI. In this study b values between 2 and 3 were applied to limit both, the rejection of good data and the acceptance 154 of erroneous data to small percentages.

The CI was calculated with the observed data at each weather station. However, no measurements of daily diffuse radiation, R<sub>d</sub>, were available. Thus, the model proposed by Aguilar et al. (2010) was applied to generate daily diffuse radiation ( $R_{dp}$ ) at each weather station without considering the observed global data at such station. Obviously, this assumption depends on the validity of the model as well as on the quality of  $R_{go}$  datasets at the remaining weather stations. However, under the common lack of diffuse solar radiation measurements like the present one, modeling them can be an alternative (e.g., Yang et al., 2020) to reject erroneous  $R_g$  observations. This approach was proposed once the model had already been validated in a previous study (Aguilar et al., 2010) but keeping in mind the intrinsic limitations and assumptions previously stated.

After this quality test, the percentage of excluded values did not reach 10% at any weather station, with a mean value close to 2% when the whole set of stations was considered. Table 1 shows selected descriptors of the data sets at each station in this study after all the quality check process and Figure 2 shows the chronogram of the final input data availability per station (N in Table 1) used in this study.



167 Figure 2. Data availability in the analyzed period (01 Feb 2000 - 31 Dec 2018) for each weather station. Stations are sorted by 168 increasing altitude from the top to the bottom row.

**Table 1.** Information of the weather stations included in this study: elevation, z (m a.s.l.); code; data length, as initial and final dates of the time series; number of initially available daily records after the quality check. N (days); rate

170	time series; number of initially available daily records, $N_0$ (days); number of available daily records after the quality check, N (days); rate
171	of days for cloudy, N <sub>CI&lt;0.3</sub> (%), partially cloudy, N <sub>0.3<ci<0.6< sub=""> (%), and clear-sky conditions, N<sub>CI&gt;0.6</sub>, (%); and maximum, R<sub>go_max</sub> (MJ m<sup>-2</sup> day<sup>-</sup></ci<0.6<></sub>
172	<sup>1</sup> ), mean, R <sub>go_mean</sub> (MJ m <sup>-2</sup> day <sup>-1</sup> ), and minimum, R <sub>go_min</sub> (MJ m <sup>-2</sup> day <sup>-1</sup> ), daily global radiation observed values. The selected descriptors for
173	sky conditions and global radiation correspond to registered data after quality check

Z	Code	Initial date	Final date	No	N	N <sub>CI&lt;0.3</sub>	N <sub>0.6<ci<0.3< sub=""></ci<0.3<></sub>	N <sub>CI&gt;0.6</sub>	$R_{go\_max}$	Rgo_mean	$R_{\text{go}\_min}$
781	602	26/01/2001	31/12/2018	6521	6370	8	23	69	33.80	18.49	0.80
942	608	01/02/2000	31/12/2018	6883	6686	6	26	68	34.20	18.83	0.70
950	601	05/09/2000	31/12/2018	6600	6449	7	27	66	33.00	18.17	0.60
975	853	21/11/2007	29/12/2018	2833	2827	8	30	62	32.37	18.01	1.00
1212	604	05/09/2000	31/12/2018	6665	6485	7	29	64	33.00	18.09	0.70
1332	803	27/08/2009	31/12/2018	3407	3282	7	22	71	33.41	18.95	0.71
1530	854	26/10/2007	16/12/2018	3176	3169	10	28	62	32.91	17.97	1.10
1732	857	16/11/2007	29/12/2018	3042	3034	11	25	64	32.84	18.31	0.81
1735	859	23/01/2008	21/11/2018	2577	2573	11	23	66	33.67	19.11	0.59
2141	804	10/10/2012	31/12/2018	2272	2206	7	21	72	33.91	19.05	0.82
2155	855	02/01/2008	30/11/2017	2522	2519	13	30	57	33.64	17.64	0.78
2300	858	09/03/2008	20/09/2017	2385	2380	12	28	60	34.58	17.99	0.99
2325	1002	15/11/2008	29/10/2012	951	951	8	22	70	35.60	20.47	1.55
2510	802	04/11/2004	31/12/2018	5050	4849	6	19	75	36.29	20.28	0.69
2867	1001	16/11/2007	01/01/2014	1071	1071	6	28	66	33.70	18.06	1.68
3097	860	23/01/2008	09/09/2018	1858	1705	13	25	62	35.79	18.20	1.12

# 175 **3.3 Generation of global radiation maps**

The GIS-based solar radiation model proposed by Aguilar et al. (2010) that was previously implemented and validated in a small subwatershed located in the southwest of Sierra Nevada (Fig. 1) was extended to the whole area in this study. For validation purposes, data registered at weather stations are considered to represent the average values of the 30 m cell of the DEM on which they are located (Batllés et al., 2008; Martínez-Durbán et al., 2009).

180 The main equations and flowchart of the model are shown in Appendix A. The complete explanation of the algorithms as well 181 as the justification of the assumptions of the model can be found in detail in Aguilar et al. (2010).

182 The model was developed to be run using limited data but considering the agents that constitute the main sources of the spatial

183 and temporal variability of solar radiation. Results generated by the model include hourly maps of diffuse, beam and reflected

184 solar radiation values with minimum input data requirements as only topographic data, albedo estimations and measured daily

global radiation records ( $R_{go}$ ) at least at one weather station are required. As for the daily global radiation registers, even when

they are missing, their estimation from other more readily available meteorological data could always be a choice from the

187 literature (Hargreaves and Samani, 1982; Bristow and Campbell, 1984; Allen, 1997; Bechini et al., 2000; Winslow et al., 2001;

188 Donatelli et al., 2003, 2006; Yang and Koike, 2005; Diodato and Bellocchi, 2007; Wu et al., 2007; Ruiz-Arias et al., 2011; Liu

189 et al., 2012b; El Ouderni et al., 2013; Mullen et al., 2013).

190 The generation of global radiation maps with the model applied (Aguilar et al., 2010) requires a proper characterization of the

191 spatio-temporal patterns of albedo in the study site. 240 cloud-free Landsat imagery available for the study period from Landsat

192 5 TM (49 images), Landsat 7 ETM+ (141 images) and Landsat 8 OLI (50 images) with a 30 m spatial resolution were used.

193 Figure 3 shows the specific dates and sensors of the 240 images analyzed in this study. All images were first properly corrected,

and their reflectivity values computed (Pimentel et al., 2014). Albedo was then derived for each image following the same

195 procedure applied in Aguilar et al. (2010), which is based on the methodology described by Brest and Goward (1987), and

196 linearly interpolated on a daily time scale for the whole study period.



Figure 3. Dates and sensors of each Landsat image analyzed in the study period (01 Feb 2000 - 31 Dec 2018).

### 199 **3.4 Cross-validation at weather stations**

207

211

Once daily global radiation estimates were generated by the model a cross validation was applied at each weather station on the daily scale. This was carried out on a leave-one-out process, i.e., data from a weather station were removed from the input dataset to the model and predicted values ( $R_{gp}$ ) at that weather station were then compared to observed data ( $R_{gp}$ ).

203 Different indicators were computed to quantitatively evaluate the performance of the model (Muneer et al., 2007):

-The Root Mean Square Error (RMSE) (Eq. 1), where R<sub>gp</sub> and R<sub>go</sub> are the predicted and observed daily global radiation (MJ

 $m^{-2}$  day<sup>-1</sup>), respectively, and N is the number of observed daily data. It measures the difference between values predicted by the model and those which were observed.

$$RMSE = \sqrt{\frac{\sum \left(R_{gp} - R_{go}\right)^2}{N}}$$
(1)

-The deviation from the 1:1 line of observed vs. predicted daily solar radiation values. Linear fits forced through the origin were obtained (Eq. 2) and the slopes ( $\alpha$  in Eq. 2) are desired to be equal to 1. The coefficient of determination, R<sup>2</sup>, as the ratio of the explained variation to the total variation, was also computed.

$$R_{gp} = \alpha \cdot R_{go} \tag{2}$$

The RMSE values and linear fits were obtained for the whole dataset at each weather station, and also for different cloudiness levels to consider different atmospheric states that may condition the performance of the model according to previous studies (Batllés et al., 2008; Martínez-Durbán et al., 2009; Ruiz-Arias et al., 2009). Based on the cloudiness three types of weather conditions were analyzed: cloudy days (CI<0.3), partly cloudy days ( $0.3 \le CI \le 0.6$ ) and clear-sky days or cloudless days (CI $\ge 0.6$ ).

The cross-validation analysis was also carried out with deseasonalized daily data to remove the expected intra-annual course of global radiation data. The deseasonalization of the daily series was carried out applying a stable seasonal filter (Brockwell and Davis, 2002) as already done in a previous study with other hydrometeorological datasets (Aguilar et al., 2017). Besides, as the reliability of solar radiation estimates is conditioned by the availability of recorded data, the cross-validation analysis for the whole study period was also computed with limited data. Thus, global radiation estimated were generated with only the four stations (601, 602, 604 and 608 in Fig. 1) with the longest records (Figure 2) as inputs to the model. Results are shown in Appendix B.

Finally, to contrast the modeling scheme applied, another well-known GIS-based solar radiation model, Solar Analyst (SA) (Fu and Rich, 2000b) was also applied in the study site. Error values in the approximation to observed data and linear fits obtained in SN are shown in Appendix C.

227 The cross-validation assessment is summarized in Figure 4. With the global datasets (in black in Fig. 4), a very close 228 approximation of the model estimates to recorded data was obtained (mean  $\alpha$  value of 0.98 and mean R<sup>2</sup> values of 0.91). RMSE

- values varied for the different stations and ranged from 1.81 (station 804) to 3.76 (station 860) with a mean value of 2.63 MJ  $m^{-2}$  dav<sup>-1</sup>.
- 231 When the analysis was carried out in terms of the cloudiness level, a general overestimation by the model (e.g., a mean  $\alpha$  value 232 of 1.41) was always seen on cloudy days (CI $\leq$ 0.3). In contrast, on clear-sky days (CI>0.6) slopes were very close to 1 with a 233 mean  $\alpha$  value of 0.96. An intermediate behavior was found on partly cloudy days (0.3 < CI < 0.6) when the model slightly under 234 predicted (e.g., stations 854 and 608) or over predicted depending on the weather station. As for RMSE values, the lowest 235 values were always found for clear sky days, when the cloud influence is minimal and the attenuation is mostly explained by 236 changes in the atmospheric transmittance, followed by partly cloudy days with mean values of 2.07 and 3.07 MJ m<sup>-2</sup> day<sup>-1</sup>, respectively. The highest RMSE values were always found on cloudy days with mean values of 3.70 MJ m<sup>-2</sup> day<sup>-1</sup>. The high 237 proportion of clear-sky days (65%) and the low RMSE values on these days (2.07 MJ m<sup>-2</sup> day<sup>-1</sup>) revealed the general good 238 239 agreement of the model estimates with observed data. This is especially important in semiarid environments, where energy-240 limited hydrological processes (e.g., soil moisture depletion, evaporation, or snowmelt) are more relevant on clear-sky days 241 and they must be carefully computed in water and energy balance modeling, irrigation scheduling, etc. (Chen et al., 1999; 242 Mamassis et al., 2012).
- There is no clear pattern in the errors obtained with the elevation of the stations. The goodness of the model estimates was more affected by the interaction of the different characteristics of the weather station (e.g., slope, aspect, surrounding terrain configuration, orographic effects in the vertical development of clouds, etc.) than by the height of the station itself.
- With the deseasonalized time series (Fig. B1), differences were reduced among the different cloudiness levels. The most remarkable change was a significant improvement in the estimates of cloudy days in every station when the range of RMSE values shifted from 2.54-7.52 (in red in Fig. 4) to 1.72-5.16 MJ m<sup>-2</sup> day<sup>-1</sup> (in red in Fig. B1). Also, the range of the slopes significantly narrowed from 1.18-1.74 (red  $\alpha$  values in Fig. 4) to 0.92-1.09 (red  $\alpha$  values in Fig. B1). Thus, the comparison with deseasonalized data showed a higher accuracy of the model than the one obtained with the original datasets (Fig. 4).
- 251 The comparison with limited input datasets shown in Figure B2 confirmed the lower reliability of global radiation estimates
- in the first five years when datasets recorded at only four stations (601, 602, 604 and 608 in Fig. 1) were available in SN. Here,
- higher elevation stations are subjected to a slightly greater overestimation of solar radiation (1.34-2.04 in red in Fig. B2), especially during cloudy conditions when the RMSE values increased to 3.62-8.45 MJ m<sup>-2</sup> day<sup>-1</sup> (in red in Fig. B2).
- 255 The application of Solar Analyst (Table C.1) revealed in general worse approximations to observed data than those shown in
- Figure 4 with mean  $R^2$  values of 0.66 and RMSE values ranging from 3.59 (station 853) to 5.11 (station 859) with the global datasets.
- 258 The errors obtained in Figure 4 were within the order of magnitude of those found in previous studies in other mountainous
- areas (Yang et al., 2006; 2010; Zhang et al., 2020) and slightly improved those previously obtained on a small subarea (10 x 5
- 260 km<sup>2</sup>) in the north-eastern side of SN. Here, Tovar-Pescador et al. (2006) analyzed the application of SA in clear sky days with
- 261 a 168 global radiation dataset from 14 weather stations located at between 1091 and 1659 m.a.s.l. They obtained  $R^2$  values of
- 0.75, similar to the value here obtained with SA estimates in the whole SN area (0.77 in Table B.1) but lower than the R<sup>2</sup> equal

- to 0.99 obtained with the model (in orange in Figure 3). Then, Batllés et al. (2008) in another application of SA in the same
- area with a 2-year daily dataset obtained the best performances for clear-sky days. RMSE values obtained in clear-sky days in
- the present study, of 11.1 % (2.07 MJ m<sup>-2</sup> day<sup>-1</sup>); were the same as those obtained by Batllés et al. (2008) for clear sky days
- 266 (11%). Later, Ruiz-Arias et al. (2009) evaluated the application of four different GIS-based solar radiation models with a 523
- 267 global radiation dataset at the same study site. RMSE values for the global dataset ranged between 1.99 and 7.28 MJ m<sup>-2</sup> dav<sup>-</sup>
- 268 <sup>1</sup> depending on the model.
- 269 The order of magnitude of the errors (Figure 4) and its comparison with those obtained with more computationally and data
- 270 demanding GIS-based models in previous studies let us to conclude that the model is the best choice to generate global radiation
- data series in SN.
- Therefore, once the model was validated in the study site, daily  $R_g$  maps were generated and aggregated at the monthly and annual scales.
- 274





Figure 4. Cross validation analysis. Linear fits of daily predicted *vs.* observed  $R_g$  (MJ m<sup>-2</sup> day<sup>-1</sup>) at each of the selected stations for the global data (black), cloudy (CI<0.3 - red), partly cloudy (0.3<CI<0.6 - blue) and clear-sky days (CI>0.6 - orange). Stations are sorted by increasing altitude from left to right and from the top to the bottom row.

- 279
- 280
- 281

### 282 4 Results

Daily, monthly, and annual  $R_g$  datasets in SN are analyzed in this section at two spatial scales. First, the results at the weather station scale are presented. Thus, possible relationships between altitude and/or location of the weather station with the different  $R_g$  statistics and how this relation changes with the temporal scale of analysis can be assessed. Then, the  $R_g$  maps that can be downloaded as specified in section 6 are analyzed.

### 287 **4.1 Daily time series of global radiation in Sierra Nevada**

Figure 5 shows the statistical distribution of the daily  $R_g$  at each weather station ordered by increasing altitude and illustrates several questions. First, there is a very similar interquartile range among stations. Second, there are greater variations in the maximum daily  $R_g$  among the different stations with a mean value of 34.0 MJ m<sup>-2</sup> day<sup>-1</sup>. Third, even though a slight increase with altitude can be shown in the high extreme statistics of the daily  $R_g$  values (e.g., in the maximum or in the 90<sup>th</sup> percentile), there is not a clear trend. Therefore, other factors such as orientation, proximity to the sea or the terrain configuration in the surrounding terrain as suggested by Batllés et al. (2008) constitute relevant features in the study site.

Figure 6 shows an example of the spatial distribution in three representative days of cloudy, partially cloudy, and clear-sky conditions. Here the spatial distribution is clearly influenced by the topography of SN, especially in the clear sky day.



296

Figure 5. Statistical distribution of daily  $R_g$  (MJ m<sup>-2</sup> day<sup>-1</sup>) time series at each of the selected stations over the study area. The box shows 50% of the data, delimited by Q1 (lower) and Q3 (upper), the solid line represents the median, and whiskers show 10<sup>th</sup> and 90<sup>th</sup> percentiles. Brown, orange and yellow dots represent daily maximum, mean and minimum time series values. Stations are sorted by increasing altitude from left to right.



302

# 306 4.2 Monthly time series of global radiation in Sierra Nevada

The statistical distribution of monthly  $R_g$  per weather station (Fig. 7) shows that in every station: i) July and December constitute the months with the highest and the lowest values of  $R_g$ , respectively; ii) there is a quite linear increase in the monthly  $R_g$  values from January to July and a sudden drop in August with a slightly convex evolution till December; and iii) the interquartile range is significantly higher in the spring and fall, than in the summer and winter months.

The increase in the high extreme statistics of radiation with the altitude of the weather station becomes more apparent at the monthly scale (Fig. 7) than at the daily scale (Fig. 5) previously analyzed. Thus, maximum values of around 1000 MJ m<sup>-2</sup> month<sup>-1</sup> are reached in July in the highest stations (e.g., 1002, 802, 1001 and 860 in Fig. 7) whereas this value decreases to around 910 MJ m<sup>-2</sup> month<sup>-1</sup> in the four lowest stations except for station 608.

Monthly  $R_g$  maps show significant spatial differences of up to 200 MJ m<sup>-2</sup> month<sup>-1</sup> in both the mean monthly values (Fig. 8) that clearly follow the terrain configuration with summits and valleys receiving high and low solar radiation values, respectively. For example, the area in the north of SN that is highly shadowed by the highest peaks in the Iberian Peninsula (Mulhacen and Veleta with 3482 and 3396 m a.s.l., respectively) is easily visible, with the lowest relative levels of insolation

319 received within SN especially in the summer months (June, July, and August in Fig. 8).

320 Both, maps of the monthly mean and standard deviation of  $R_g$  (Fig. 8) and the statistical distribution of the monthly  $R_g$  in the

321 study site (Fig. 9), show the same behaviour as the one obtained at the weather stations regarding; i) July and December as the

months with the highest and lowest values of  $R_g$  received in SN; and ii) the highest scatter in the monthly  $R_g$  values in the spring and fall months.



325

Figure 7. Statistical distribution of monthly  $R_g$  (MJ m<sup>-2</sup> month<sup>-1</sup>) time series at each of the selected stations over the study area. The box shows 50% of the data, delimited by Q1 (lower) and Q3 (upper), the solid line represents the median, and whiskers show 10<sup>th</sup> and 90<sup>th</sup> percentiles. Brown, orange and yellow dots represent monthly maximum, mean and minimum time series values. Stations are sorted by increasing altitude from left to right and from the top to the bottom row.



 $\label{eq:stars} 333 \qquad \mbox{Figure 8. Monthly average and standard deviation of $R_g$ (MJ $m^{-2}$ month^{-1}$) in the study period (2000-2018) in $N$.}$ 





 $\begin{array}{ll} 336 \\ 336 \end{array} \mbox{Figure 9. Statistical distribution of the monthly $R_g$ (MJ m^{-2} month^{-1}) values throughout the study area. Whisker boxes represent the $10^{th}, 25^{th}, 50^{th}, 75^{th}$ and $90^{th}$ percentiles of each monthly map per year. } \end{array}$ 

- For the study period (2000-2018), there is a great heterogeneity in the statistical distribution of the monthly  $R_g$  in the study site (Fig. 9) especially in the incoming radiation along the months of the wet season (October-May). In this way, in the most insolated years in the study period (2005 and 2012), significantly higher monthly radiation values were found in certain months
- of the springtime (March and May 2012 and April 2005). In those months, the higher than usual rate of clear-sky over cloudy
   days finally determines the annual differences in the incoming global radiation in SN.
- When considering the temporal evolution of the distribution of  $R_g$  within the monthly maps in SN (Fig. 10), certain interannual differences can be observed along the study period, such as the existence of certain months in spring with unexpected low monthly radiation values (eg. 2001, 2004, 2007 and 2008), or two relative maximum monthly  $R_g$  values (e.g., 2009, 2010 and 2014). Moreover, Figure 10 shows a higher scatter in the monthly maximum (June-August) and minimum (November-January)
- 348 R<sub>g</sub> values in SN than when the analysis is carried out at each weather station (Fig. 7).
- 349



Figure 10. Evolution of the statistical distribution of monthly  $R_g$  (MJ m<sup>-2</sup> month<sup>-1</sup>) in the study period (2001-2018) throughout the study area. Grayscale colours represent the following percentiles: 5<sup>th</sup>, 10<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, 90<sup>th</sup> and 95<sup>th</sup>.

# 353 4.3 Annual times series of global radiation in Sierra Nevada

Unlike at the daily scale (Fig. 5), a great variability among the different weather stations in terms of the global radiation received at the annual temporal scale is found (Fig. 11). Thus, we find minimum annual  $R_g$  values from 5920 MJ m<sup>-2</sup> year<sup>-1</sup> in station 854 to around 6750 MJ m<sup>-2</sup> year<sup>-1</sup> in station 1002. This difference is even bigger in the maximum annual  $R_g$  values from 6700 to 7720 MJ m<sup>-2</sup> year<sup>-1</sup> in stations 854 and 802, respectively, and is also shown in the interquartile range.

- 358 When analyzing the influence of altitude, the weather stations above 1500 m a.s.l (854, 857, 859, 804, 855, 858, 1002, 802,
- 1001, 860 in Fig. 11) show their altitudinal gradient in all the statistics of the annual  $R_g$  values considered.

Annual  $R_g$  maps (Fig. 12) show the same spatial differences that follow the terrain configuration as those observed in the monthly time series (Fig. 8). For example, the area in the north of SN that is highly shadowed as previously mentioned corresponds to the area with the mean minimum annual values received in the study period, 4063 MJ m<sup>-2</sup> year<sup>-1</sup>, that only represents 63% the mean annual accumulated values in SN (6316 MJ m<sup>-2</sup> year<sup>-1</sup>).



364

369

Significant interannual differences can be easily shown with differences in the mean annual  $R_g$  value in the study area of up to 800 MJ m<sup>-2</sup> year<sup>-1</sup> between 2005 and 2018. Such years with particularly high and low annual incoming radiation also presented higher (6800 MJ m<sup>-2</sup> year<sup>-1</sup>) and lower median annual  $R_g$  values (6200 MJ m<sup>-2</sup> year<sup>-1</sup>), respectively, than the annual median for the whole study period in SN (6456 MJ m<sup>-2</sup> year<sup>-1</sup>) (Fig. 13). These results agree with the annual irradiation map obtained by Batllés et al. (2008) in the north-eastern part of SN. They reported maximum and minimum annual values of 7516 and 2342 MJ m<sup>-2</sup> year<sup>-1</sup> on the summits and in deep valleys, respectively, and thus, concluded that irradiation levels were more related to topographic characteristics than to altitude.

Figure 11. Statistical distribution of annual  $R_g$  (MJ m<sup>-2</sup> year<sup>-1</sup>) time series at each of the selected stations over the study area. The box shows 50% of the data, delimited by Q1 (lower) and Q3 (upper), the solid line represents the median, and whiskers show 10<sup>th</sup> and 90<sup>th</sup> percentiles. Brown, orange and yellow dots represent annual maximum, mean and minimum time series value. Stations are sorted by increasing altitude from left to right.



379 Figure 12. Annual global radiation (MJ m<sup>-2</sup> year<sup>-1</sup>) in the study period (2001-2018) in SN.



Figure 13. Evolution of the statistical distribution of annual  $R_g$  (MJ m<sup>-2</sup> year<sup>-1</sup>) in the study period (2001-2018) throughout the study area. Dashed lines represent the mean values of the percentiles analyzed.

### **5 Data availability**

381

385 The daily, monthly and annual global radiation maps derived in this study can be accessed and downloaded in .ncdf format 386 from: https://doi.pangaea.de/10.1594/PANGAEA.921012 (Aguilar et al., 2021). Besides, a .txt file containing the availability 387 (code 1) or gaps (code 0) in the daily  $R_g$  observations at each weather station has been added as a supplement to this paper. 388 Hourly datasets were also computed in this study but due to their large storing capacity requirements they have not been 389 included in the data repository specified above. Thus, hourly maps can be provided for certain dates upon request to the authors. 390 However, a validation of these hourly datasets like the one applied in the daily estimates at the weather stations has not been 391 specifically carried out in this study. Therefore, in case hourly maps are requested to the authors, these data should be taken 392 with caution as the only available validation in SN was carried out at one weather station (802 in Fig. 1) and for a shorter 393 period (2004-2010) in Aguilar et al. (2010).

# 394 6 Final remarks

This study presents nineteen years (2000-2018) of daily, monthly, and annual global radiation maps of high spatial resolution (30 m) in a high mountain Mediterranean site. In these areas the common lack of weather stations in high altitudes makes it difficult to accurately assess solar radiation spatial patterns.

398 A GIS-based modelling scheme based on measurements or estimations of incoming daily global radiation was applied and 399 validated in the sixteen weather stations available at this unique study site. Mean RMSE values ranged from 1.81 to 3.76 MJ 400  $m^{-2}$  day<sup>-1</sup>, depending on the weather station. The best estimations were always obtained on clear-sky days, when mean RMSE 401 values decreased to 2.07 MJ m<sup>-2</sup> dav<sup>-1</sup>. The largest errors were obtained on cloudy days, which constitute on average 10% of 402 the daily datasets, and, therefore, future research should be conducted in order to improve the estimations in these situations 403 keeping the minimum input data requirements (daily global radiation data) advantage of the model. However, the high 404 proportion (65%) of clear-sky days, and the low RMSE values on those days, allow one to conclude that there is a good 405 agreement between the model estimates and observed data in the study site.

Spatial differences of around 2000 MJ m<sup>-2</sup> yr<sup>-1</sup> were found within each year analyzed. In addition, significant differences were easily shown between the years in mean incoming values of up to 800 MJ m<sup>-2</sup> yr<sup>-1</sup>. Those differences were mostly due to the variability in the incoming radiation at the wet season (October-May), with higher rates of clear-sky days in the most insolated years (e.g., 2005).

Thus, we can affirm that the modeling scheme here applied is an efficient option in semiarid mountainous areas, where daily global radiation datasets constitute the only source of solar radiation data.

Time series of these surface global radiation datasets can be used to analyze inter-annual and seasonal variation characteristics of the global radiation received in SN with high spatial detail (30 m). The availability of long global radiation datasets allows to capture the annual variability within each cycle of the Sun activity, as reported in the literature (Scaffetta and Wilson, 2013),

415 and thus estimate its contribution to the annual variability of other climate variables in these semiarid mountainous areas.

416 Dense and properly-maintained weather station networks in mountainous areas are rarely available. Thus, these datasets can 417 also be used as cross-validation reference data for other global radiation distributed datasets generated in SN with different 418 spatio-temporal interpolation techniques.

These results can also assess the order of magnitude of different sources of spatial variability (altitude/slope/aspect gradients) as well as the seasonal range of variation at different time scales and their annual variability. This estimation may provide a first estimate of the order of magnitude of uncertainty of average calculations or spatial interpolation from a scarce number of weather stations in Mediterranean and semiarid mountain areas.

The correct assessment of the solar radiation regime is crucial to correctly determine the temporal evolution of energy-limited hydrological processes such as the snow layer dynamics, soil moisture depletion and evapotranspiration (Tomas-Burguera et al., 2019). Thus, as a key input parameter for the water and energy balance, these high spatial resolution solar radiation time series are useful not only for research on the snow domain and water planning in SN in the application of hydrological 427 modelling, but in many other applications. For example, within the agricultural sector in the estimations of evapotranspiration 428 for irrigation scheduling, ecology and biodiversity studies, stand-alone solar energy facilities designing and location, 429 recreational activities in the area that strongly rely on the hydro-meteorological conditions of SN, etc. Finally, this work 430 contributes to feed research related to some key questions in hydrology, as UPH 16 and UPH 5 identified by Blöschl et al. 431 (2019).

432

# 433 Author contributions

CA, in collaboration with MJP, conceived the research. CA processed the data, applied the quality control to the raw global radiation data, modelled global radiation datasets and developed the cross-validation algorithms. RP processed satellite data, generated albedo maps for the study period, prepared the final figures and the available datasets generated in the study. CA prepared the manuscript with contributions of MJP and RP; all authors discussed and revised the final text.

438

# 439 **Competing interests**

440 The authors declare that they have no conflict of interest.

441

# 442 Acknowledgements

443 This study was supported by the following research projects funded by Spanish Ministry of Science and Innovation - MICINN: 444 Research Project RTI2018-099043-B-I00, "Operability in hydrological management under snow torrentiality/drought 445 conditions in high mountain in semiarid watersheds"; and, by Spanish Ministry of Economy and Competitiveness - MINECO: 446 Research Project CGL 2014-58508R, "Global monitoring system for snow areas in Mediterranean regions: trends analysis and 447 implications for water resource management in Sierra Nevada", and Research Project CGL 2011-25632, "Snow dynamics in 448 Mediterranean regions and its modelling at different scales. Implication for water management". Moreover, the present work 449 was partially developed within the framework of the Panta Rhei Research Initiative of the International Association of 450 Hydrological Sciences (IAHS) (Working Groups Water and energy fluxes in a changing environment and Mountain 451 Hydrology). Rafael Pimentel acknowledges fundings by the modality 5.2 of the Programa Propio-2018 of the University of 452 Cordoba and the Juan de la Cierva Incorporación Programme of the Ministry of Science and Innovation (IJC2018-038093-I). 453 The continuous support of the Natural and National Park of Sierra Nevada has also been determinant for the development of 454 this line of research since 2002. Finally, tremendous appreciation is extended to all the weather station networks that maintain 455 and make accessible datasets to scientific research.

456

# 457 **References**

458 Aguilar, C. and Polo, M. J.: Generating reference evapotranspiration surfaces from the Hargreaves equation at watershed scale,

459 Hydrol. Earth Syst. Sci., 15, 2495-2508, <u>https://doi.org/10.5194/hess-15-2495-2011</u>, 2011.

- 460 Aguilar, C., Herrero, J., and Polo, M. J.: Topographic effects on solar radiation distribution in mountainous watersheds and
- their influence on reference evapotranspiration estimates at watershed scale, Hydrol. Earth Syst. Sci., 14, 2479–2494,
  https://doi.org/10.5194/hess-14-2479-2010, 2010.
- Aguilar, C., Montanari, A., and Polo, M.J.: Real-time updating of the flood frequency distribution through data assimilation,
  Hydrol. Earth Syst. Sci., 21, 3687–3700, https://doi.org/10.5194/hess-21-3687-2017, 2017.
- Aguilar, C., Pimentel, R., and Polo, M. J.: Time series of distributed global radiation data in Sierra Nevada (Spain) at different
   scales from historical weather stations. PANGAEA. https://doi.pangaea.de/10.1594/PANGAEA.921012, 2021.
- Allen, R.G.: Self-calibrating method for estimating solar radiation from air temperature, J. Hydrol. Eng., 2, 56-67,
  https://doi.org/10.1061/(ASCE)1084-0699(1997)2:2(56), 1997.
- 469 Anderson, R. S., Jiménez-Moreno, G., Carrión, J. S., and Pérez-Martínez, C.: Postglacial history of alpine vegetation, fire, and
- 470 climate from Laguna de Río Seco, Sierra Nevada, southern Spain, Quaternary Sci. Rev., 30, 1615–1629,
  471 https://doi.org/10.1016/i.quascirev.2011.03.005, 2011.
- 472 Batllés, J., Bosch, J.L., Tovar-Pescador, J., Martínez-Durbán, M., Ortega, R., and Miralles, I.: Determination of atmospheric
- 473 parameters to estimate global radiation in areas of complex topography: Generation of global irradiation map, Energy Convers.
- 474 Manage., 49, 336-345, https://doi.org/10.1016/j.enconman.2007.06.012, 2008.
- Bechini, L., Ducco, G., Donatelli, M. and Stein, A.: Modelling, interpolation and stochastic simulation in space and time of
  global solar radiation, Agric. Ecosyst. Environ., 81, 29-42, https://doi.org/10.1016/S0167-8809(00)00170-5, 2000.
- 477 Blanca, G., Cueto, M., Martínez-Lirola, M. J., and Molero-Mesa, J.: Threatened vascular flora of Sierra Nevada (Southern
- 478 Spain), Biological Conservation, 85(3), 269-285, doi:10.1016/S0006-3207(97)00169-9, 1998.

479 Blöschl, G. et al.: Twenty-three unsolved problems in hydrology (UPH) – a community perspective, Hydrol. Sci. J., 64 (1),

- 480 1141-1158, <u>https://doi.org/10.1080/02626667.2019.1620507</u>, 2019.
- 481 Bonet-García, F.J., Pérez-Luque, A.J., Moreno-Llorca, R.A., Pérez-Pérez, R., Puerta-Piñero, C., and Zamora, R.: Protected
- 482 areas as elicitors of human well-being in a developed region: a new synthetic (socioeconomic) approach, Biol. Conserv., 187
- 483 221-9, https://doi.org/10.1016/j.biocon.2015.04.027, 2015.
- Bosch, J.L., López, G., and Batlles, F.J.: Daily solar irradiation estimation over a mountainous area using artificial neural
  networks, Renew. Energy, 33, 1622-1628, https://doi.org/10.1016/j.renene.2007.09.012, 2008.
- Brest, C.L. and Goward, S.N.: Deriving surface albedo measurements from narrow band satellite data, Int. J. Remote Sens.,
- 487 8, 351-367, https://doi.org/10.1080/01431168708948646, 1987.
- 488 Bristow, K.L. and Campbell, G.S.: On the relationship between incoming solar radiation and daily maximum and minimum
- 489 temperature, Agric. For. Meteorol., 31, 159-166, https://doi.org/10.1016/0168-1923(84)90017-0, 1984.
- 490
- 491 Brockwell, P. J. and Davis, R. A.: Introduction to Time Series and Forecasting, Vol. 1, Taylor & Francis, 2002.

- Cañadas, E. M., Fenu, G., Peñas, J., Lorite, J., Mattana, E., and Bacchetta, G.: Hotspots within hotspots: 445 Endemic plant
  richness, environmental drivers, and implications for conservation, Biological Conservation, 170, 282-291,
  https://doi.org/10.1016/j.biocon.2013.12.007, 2014.
- Chen, J.M., Liu, J., Cihlar, J., and Goulden, M.L.: Daily canopy photosynthesis model through temporal and spatial scaling
  for remote sensing applications, Ecol. Modell., 124, 99-119, https://doi.org/10.1016/S0304-3800(99)00156-8, 1999.
- 497 Chen, X., Su, Z., Ma, Y., Yang, K., and Wang, B.: Estimation of surface energy fluxes under complex terrain of Mt.
- 498 Qomolangma over the Tibetan Plateau, Hydrol. Earth Syst. Sci., 17, 1607-1618, https://doi.org/10.5194/hess-17-1607-2013,
  499 2013.
- 500 Diodato, N. and Bellocchi, G.: Modelling solar radiation over complex terrains using monthly climatological data, Agric. For.
- 501 Meteorol. 144, 111-126, https://doi.org/10.1016/j.agrformet.2007.02.001, 2007.
- 502 Donatelli, M., Bellocchi, G., and Fontana, F.: RadEst3.00: software to estimate daily radiation data from commonly available 503 meteorological variables, European J. Agron., 18, 363-367, https://doi.org/10.1016/S1161-0301(02)00130-2, 2003.
- 504 Donatelli, M., Carlini, L., and Bellocchi. G.: A software component for estimating solar radiation, Environ. Modell. Software,
- 505 21(3), 411-416, https://doi.org/10.1016/j.envsoft.2005.04.002, 2006.
- 506 Dozier, J. and Frew, J.: Rapid calculation of terrain parameters for radiation modeling from digital elevation data, IEEE Trans.
- 507 Geosci. Remote Sens., 28, 963-969, https://doi.org/10.1109/36.58986, 1990.
- 508 Dozier, J., Bruno, J., and Downey, P.: A faster solution to the horizon problem, Comp. Geosci., 7, 145-151, 509 https://doi.org/10.1016/0098-3004(81)90026-1, 1981.
- 510 Dubayah, R.C.: Estimating net solar radiation using Landsat Thematic Mapper and digital elevation data, Water Resour. Res.,
- 511 28, 2469-2484, https://doi.org/10.1029/92WR00772, 1992.
- 512 Dubayah, R.C.: Modeling a solar radiation topo climatology for the Rio Grande river watershed, J. Veg. Sci. 5, 627-640, 513 https://doi.org/10.2307/3235879, 1994.
- 514 Dubayah, R.C. and van Katwijk, V.: The topographic distribution of annual incoming solar radiation in the Rio Grande river 515 basin, Geophys. Res. Lett., 19(22), 2231-2234, https://doi.org/10.1029/92GL02284, 1992.
- 516 Dubayah, R. and Rich, P.M.: Topographic solar radiation models for GIS, Int. J. Geogr. Inf. Syst., 9, 405-413, 517 https://doi.org/10.1080/02693799508902046, 1995.
- El Ouderni, A.R., Maatallah, T., El Alimi, S., and Ben Nassrallah, S.: Experimental assessment of the solar energy potential in the gulf of Tunis, Tunisia, Renew. Sustain. Energy Rev., 20, 155-168, https://doi.org/10.1016/j.rser.2012.11.016, 2013.
- 520 Fu, P. and Rich, P.M.: A geometric solar radiation model and its applications in agriculture and forestry. In: Proceedings of
- 521 the Second International Conference on Geospatial Information in Agriculture and Forestry, Lake Buena Vista, 10-12 January
- 522 2000, I-357-364, 2000a.
- 523 Fu, P. and Rich, P.M.: The Solar Analyst 1.0 Manual. Helios Environmental Modeling Institute (HEMI), USA, 2000b.
- 524 Fu, P. and Rich, P.M.: A Geometric Solar Radiation Model with Applications in Agriculture and Forestry, Comput. Electr.
- 525 Agric., 37, 25-35, https://doi.org/10.1016/S0168-1699(02)00115-1, 2002.

- 526 Geiger, M., Diabate, L., Menard, L., and Wald, L.: A web service for controlling the quality of measurements of global solar
- 527 irradiation, Sol. Energy, 73, 474-480, https://doi.org/10.1016/S0038-092X(02)00121-4, 2002.
- Goldberg, V. and Häntzschel, J.: Application of a radiation model for small-scale complex terrain in a GIS environment,
  Meteorol. Zeitschrift. 11(2), 119-128, https://doi: 10.1127/0941-2948/2002/0011-0119, 2002.
- Hargreaves, G.H. and Samani, Z.A.: Estimating potential evapotranspiration, J. Irrig. Drain. Engin. 108 IR3, 223-230, 1982.
- 531 Herrero, J. and Polo, M. J.: Parameterization of atmospheric longwave emissivity in a mountainous site for all sky conditions,
- 532 Hydrol. Earth Syst. Sci., 16, 3139–3147, https://doi.org/10.5194/hess-16-3139-2012, 2012.
- 533 Herrero, J. and Polo, M. J.: Evaposublimation from the snow in the Mediterranean mountains of Sierra Nevada (Spain), The
- 534 Cryosphere, 10, 2981–2998, https://doi.org/10.5194/tc-10-2981-2016, 2016.
- Herrero, J., Polo, M. J., Moñino, A., and Losada, M. A.: An energy balance snowmelt model in a Mediterranean site, J. Hydrol.,
- 536 371, 98–107, https://doi.org/10.1016/j.jhydrol.2009.03.021, 2009.
- 537 Hewitt, G.M.: Mediterranean Peninsulas: the evolution of hotspots. In: Zachos, F.E. and Habel, J.C. (eds), Biodiversity
- botspots: distribution and protection of conservation priority area, Springer, Berlin, pp. 123-147, 2011.
- Heywood, V. H.: The Mediterranean flora in the context of world diversity, Ecología Mediterránea, 21, 11-18, 1995.
- 540 Ineichen, P. and Pérez, R.: A new airmass independent formulation for the Linke Turbidity coefficient, Sol. Energy, 73, 151-
- 541 157, https://doi.org/10.1016/S0038-092X(02)00045-2, 2002.
- 542 Iqbal, M.: An introduction to solar radiation, Academic Press, Ontario, 1983.
- Jacovides, C.P., Hadjioannou, L., Pashiardis, S., and Stefanou, L.: On the diffuse fraction of daily and monthly global radiation
  for the island of Cyprus, Sol. Energy, 56, 565-572, https://doi.org/10.1016/0038-092X(96)81162-5, 1996.
- 545 Journée, M. and Bertrand, C.: Quality control of solar radiation data within the RMIB solar measurements network, Sol.
- 546 Energy, 85, 72-86, https://doi.org/10.1016/j.solener.2010.10.021, 2011.
- Kasten, F. and Young, A.T.: Revised optical air mass tables and approximation formula, App. Optics, 28(22), 4735-4738,
  https://doi.org/10.1364/AO.28.004735, 1989.
- 549 Liu, M., Bárdossy, A., Li, J., and Jiang, Y.: GIS-based modelling of topography-induced solar radiation variability in complex
- terrain for data sparse region, Int. J. Geogr. Inf. Sci., 26(7), 1281-1308, https://doi.org/10.1080/13658816.2011.641969, 2012a.
- 551 Liu, J.D., Liu, J.M., Linderholm, H.W., Chen, D.L., Yu, Q., Wu, D.R., and Haginoya, S.: Observation and calculation of the
- solar radiation on the Tibetan Plateau, Energy Convers. Manag., 57, 23-32, https://doi.org/10.1016/j.enconman.2011.12.007,
  2012b.
- Liu, Y., Zhang, P., Nie, L., Xu, J., Lu, X., and Li, S.: Exploration of the Snow Ablation Process in the Semiarid Region in
- 555 China by Combining Site-Based Measurements and the Utah Energy Balance Model-A Case Study of the Manas River Basin,
- 556 Water, 11, 1058, https://doi.org/10.3390/w11051058, 2019
- 557 MacDonell, S., Kinnard, C., Mölg, T., Nicholson, L., and Abermann, J.: Meteorological drivers of ablation processes on a cold
- 558 glacier in the semi-arid Andes of Chile, The Cryosphere, 7, 1513–1526, https://doi.org/10.5194/tc-7-1513-2013, 2013.

- 559 Mamassis, N., Efstratiadis, A., and Apostolidou, I.G.: Topography-adjusted solar radiation indices and their importance in 560 hvdrology, Hvdr. Sc. J., 57(4), 756-775, https://doi.org/10.1080/02626667.2012.670703, 2012.
- 561 Martínez-Durbán, M., Zarzalejo, L.F., Bosch, J.L., Rosiek, S., Polo, J., and Batlles, F.J.: Estimation of global daily irradiation
- 562 in complex topography zones using digital elevation models and meteosat images: Comparison of the results, Energy Convers.
- 563 Manage. 50, 2233-2238, https://doi.org/10.1016/j.enconman.2009.05.009, 2009.
- Moradi, I.: Quality control of global solar radiation using sunshine duration hours, Energy, 34, 1-6, https://doi.org/10.1016/j.energy.2008.09.006, 2009.
- Mullen, R., Marshall, L., and McGlynn, B.: A Beta Regression Model for Improved Solar Radiation Predictions, J. Appl.
  Meteor. Climatol., 52, 1923-1938, https://doi.org/10.1175/JAMC-D-12-038.1, 2013.
- Muneer, T., Younes, S., and Munawwar, S.: Discourses on solar radiation, Renew. Sust. Energy Rev., 11, 551-602,
  https://doi.org/10.1016/j.rser.2005.05.006, 2007.
- 570 Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B., and Kent, J.: Biodiversity hotspots for conservation
- 571 priorities, Nature, 403, 853-858, https://doi.org/10.1038/35002501, 2000.
- 572 O'Farrell, P.J., Reyers, B., Le Maitre, D.C., et al.: Multi-functional landscapes in semi arid environments: implications for
- 573 biodiversity and ecosystem services, Landscape Ecol., 25, 1231-1246, <u>https://doi.org/10.1007/s10980-010-9495-9</u>, 2010.
- Oliphant, A.J., Spronken-Smith, R.A., Sturman, A.P., and Owens, I.F.: Spatial Variability of Surface Radiation Fluxes in
  Mountainous Terrain, J. App. Meteorol., 42, 113-128, https://doi.org/10.1175/15200450(2003)042<0113:SVOSRF>2.0.CO;2, 2003.
- Pauli, H. et al.: Recent plant diversity changes on Europe's mountain summits, Science 336, 353-355,
  https://doi:10.1126/science.1219033, 2012.
- 579 Pérez-Palazón, M. J., Pimentel, R., Herrero, J., Aguilar, C., Perales, J. M., and Polo, M. J.: Extreme values of snow-related
- variables in Mediterranean regions: trends and long-term forecasting in Sierra Nevada (Spain), Proc. IAHS, 369, 157–162,
  https://doi.org/10.5194/piahs-369-157-2015, 2015.
- 582 Pérez-Palazón, M. J., Pimentel, R., and Polo, M. J.: Climate Trends Impact on the Snowfall Regime in Mediterranean Mountain
- 583 Areas: Future Scenario Assessment in Sierra Nevada (Spain), Water, 10, 720, https://doi.org/10.3390/w10060720, 2018.
- 584 Pimentel, R., Herrero, J., and Polo, M.J.: Estimating snow albedo patterns in a Mediterranean site from Landsat TM and ETM+
- 585 images, In: Proceedings of the SPIE, Remote Sensing for Agriculture, Ecosystems, and Hydrology XV, 88870L, Dresden,
- 586 Germany, 23-26 September 2013, doi:10.1117/12.2029064.
- 587 Pimentel, R., Herrero, J., and Polo, M.J.: Graphic user interface to preprocess Landsat TM, ETM+ and OLI images for
- 588 hydrological applications. In: Proceedings of the HIC 2014, 11th International Conference on Hydroinformatics, New York,
- 589 17-21 August 2014.
- 590 Polo, M. J., Herrero, J., Pimentel, R., and Pérez-Palazón, M. J.: The Guadalfeo Monitoring Network (Sierra Nevada, Spain):
- 591 14 years of measurements to understand the complexity of snow dynamics in semiarid regions, Earth Syst. Sci. Data, 11, 393–
- 592 407, https://doi.org/10.5194/essd-11-393-2019, 2019.

- Ruiz-Arias, J.A., Tovar-Pescador, J., Pozo-Vázquez, D., and Alsamamra, H.: A comparative analysis of DEM-based models
  to estimate the solar radiation in mountainous terrain, Int. J. Geogr. Inf. Sci., 23, 1049-1076,
  https://doi.org/10.1080/13658810802022806, 2009.
- 596 Ruiz-Arias, J.A., Alsamamra, H., Tovar-Pescador, J., and Pozo-Vázquez, D.: Proposal of a regressive model for the hourly 597 diffuse solar radiation under all sky conditions. Energy Convers. Manage.. 51. 881-893. 598 https://doi.org/10.1016/j.enconman.2009.11.024, 2010.

599 Ruiz-Arias, J.A., Pozo-Vázquez, D., Santos-Alamillos, F.J., Lara-Fanego, V., and Tovar-Pescador, J.A.: Topographic

600 geostatistical approach for mapping monthly mean values of daily global solar radiation: A case study in southern Spain, Agric.

601 For. Meteorol., 151, 1812-1822, https://doi.org/10.1016/j.agrformet.2011.07.021, 2011.

- 602 Scaffetta, N. and Willson, R.C.: Multiscale comparative spectral analysis of satellite total solar irradiance measurements from
- 603 2003 to 2013 reveals a planetary modulation of solar activity and its nonlinear dependence on the 11 yr solar cycle, Pattern
- 604 Recogn. Phys., 1, 123-133, doi: 10.5194/prp-1-123-2013, 2013.
- 605 Sheng, J., Wilson, J.P., and Lee, S.: Comparison of land surface temperature (LST) modeled with a spatially-distributed solar
- radiation model (SRAD) and remote sensing data, Environ. Modell. Software, 24(3), 436-443,
  https://doi.org/10.1016/j.envsoft.2008.09.003, 2009.
- Šùri, M. and Hofierka, J.: A new GIS-based solar radiation model and its application to photovoltaic assessments, Trans. GIS,
   8(2), 175-190, https://doi.org/10.1111/j.1467-9671.2004.00174.x, 2004.
- Tang, W., Yang, K., Qin, J., Li, X., and Niu, X.: A 16-year dataset (2000–2015) of high-resolution (3 h, 10 km) global surface
  solar radiation, Earth Syst. Sci. Data, 11, 1905–1915, https://doi.org/10.5194/essd-11-1905-2019, 2019.

Tomas-Burguera, M., Vicente-Serrano, S. M., Beguería, S., Reig, F., and Latorre, B.: Reference crop evapotranspiration
database in Spain (1961–2014), Earth Syst. Sci. Data, 11, 1917–1930, https://doi.org/10.5194/essd-11-1917-2019, 2019.

- 614 Tovar. J., Olmo, F.J., and Alados-Arboledas, L.: Local scale variability of solar radiation in a mountainous region, J. App.
- 615 Meteorol., 34, 2316-2322, https://doi.org/10.1175/1520-0450(1995)034<2316:LSVOSR>2.0.CO;2, 1995.
- 616 Tovar-Pescador, J., Pozo-Vázquez, D., Ruiz-Arias, J.A., Batllés, J., López, G., and Bosch, J.L.: On the use of the digital
- elevation model to estimate the solar radiation in areas of complex terrain, Meteorol. Appl., 13, 279-287,
  https://doi.org/10.1017/S1350482706002258, 2006.
- 619 Wilson, J.P. and Gallant, J.C.: Secondary topographic attributes. In: Wilson, J.P. and Gallant, J.C. (eds), Terrain Analysis:
- 620 Principles and Applications, John Wiley & Sons, New York, pp. 51-85, 2000.
- Winslow, J.C., Hunt, E.R., and Piper, S.C.: A globally applicable model of daily solar irradiance estimated from air temperature
   and precipitation data, Ecol. Modell., 143, 227-243, https://doi.org/10.1016/S0304-3800(01)00341-6, 2001.
- Wu, G.F., Liu, Y.L., and Wang, T.J.: Methods and strategy for modeling daily global solar radiation with measured meteorological data-A case study in Nanchang station, China, Energy Convers. Manage., 48, 2447-2452,
- 625 https://doi.org/10.1016/j.enconman.2007.04.011, 2007.

- 626 Yang, K. and Koike, T.: A general model to estimate hourly and daily solar radiation for hydrological studies., Water Resour.
- 627 Res. 41(10), W10403, doi: 10.1029/2005WR003976, https://doi.org/10.1029/2005WR003976, 2005.
- 428 Yang, K., Koike, T., and Ye, B.: Improving estimation of hourly, daily, and monthly solar radiation by importing global data
- 629 sets, Agric. For. Meteorol., 137, 43-55, https://doi.org/10.1016/j.agrformet.2006.02.001, 2006.
- 630 Yang. K., He, J., Tang, W.J., and Qin, J. and Cheng, C.C.K.: On downward shortwave and longwave radiations over high
- altitude regions: Observation and modeling in the Tibetan Plateau, Agric. For. Meteorol., 150, 38-46,
  https://doi.org/10.1016/i.agrformet.2009.08.004, 2010.
- Yang, L., Cao, Q., Yu, Y., and Liu, Y.: Comparison of daily diffuse radiation models in regions of China without solar radiation
  measurement, Energy, 191, 116571, https://doi.org/10.1016/j.energy.2019.116571, 2020.
- 635 Younes, S., Claywell, R., and Muneer, T.: Quality control of solar radiation data: present status and proposed new approaches,
- 636 Energy, 30, 1533-1549, https://doi.org/10.1016/j.energy.2004.04.031, 2005.
- Zhang, S., Li, X., She, J., and Peng, X. 2019. Assimilating remote sensing data into GIS-based all sky solar radiation modeling
  for mountain terrain, Remote Sens. Envir., 231, https://doi.org/10.1016/j.rse.2019.111239, 2019.
- 639 Zhang, M., Wang, B., Liu, D.L., Liu, J., Zhang, H., Feng, P., Kong, D., Cleverly, J., Yang, X., and Yu, Q.: Incorporating
- dynamic factors for improving a GIS-based solar radiation model. Trans. GIS 2020, <u>https://doi.org/10.1111/tgis.12607</u>, 2020.
   641

# 642 Appendix A. Solar radiation equations

The sequence followed by the model is summarized in Figure A1. Computations are classified at the point scale of weather stations (Point) and the distributed scale of grids of the Digital Elevation Model (DEM) (Distributed). The complete explanation of the algorithms and assumptions of the model can be found in detail in Aguilar et al. (2010).

Firstly, daily extraterrestrial radiation ( $R_{ext}$  in MJ m<sup>-2</sup> day<sup>-1</sup>) is computed by integrating the extraterrestrial radiation incident upon a horizontal surface relative to the sun's beams from sunrise to sunset (Eq. A1).

$$R_{ext} = E_o \cdot I_{SC} \cdot \cos(\theta_z)$$
(A1)

where  $I_{SC}$  is the solar constant (1367 W m<sup>-2</sup>),  $\theta_z$  is the zenith angle and  $E_o$ , the eccentricity factor. These variables were computed following the equations in Dozier et al. (1981).

Then, the daily clearness index (CI), as the ratio of observed daily global radiation ( $R_{go}$  in MJ m<sup>-2</sup> day<sup>-1</sup>) to the daily extraterrestrial radiation, is computed at each weather station (Eq. A2).

$$CI = \frac{R_{go}}{R_{ext}}$$
(A2)

654 CI is expressed in terms of two factors, CI<sub>CS</sub> and fCI<sub>cl</sub>. The first term represents the influence of atmosphere under clear-sky 655 conditions over solar radiation, while the second term includes the cloudiness effects that decrease the final incoming solar 656 radiation (Eq. A3). The approximation of Ineichen and Perez (2002) is used to compute the global radiation under clear-sky 657 conditions,  $R_{ecs}$ , and thus, distributed hourly  $R_{ecs}$  values are obtained from the sun elevation angle, the height of the cell, the 658 Linke turbidity factor (T<sub>L</sub>) and the atmospheric mass obtained following the parameterization of Kasten and Young (1989). 659 Thus, hourly CI<sub>CS</sub> values can be computed cell by cell and then the mean daily distributed values are generated. Once daily CI 660 and CI<sub>CS</sub> values are known, fCI<sub>cl</sub> is obtained at each weather station from Eq. A3 and spatially interpolated following the 661 inverse distance weighted (IDW) method. From daily CI<sub>CS</sub> and fCI<sub>cl</sub> maps, daily interpolated CI and R<sub>g</sub> values can be obtained 662 at cell scale from Eq. A3 and A4.

$$CI = CI_{CS} \cdot fCI_{cl} \tag{A3}$$

$$R_g = R_{ext} \cdot CI_{CS} \cdot fCI_{cl} \tag{A4}$$

Topographic effects need to be evaluated for the different sun positions during the day and thus, hourly values of the different components need to be derived. Two different procedures are currently available in the model. The first one proposed in Aguilar et al. (2010) applies Jacovides et al. (1996) (Eq. A5.1) to produce the daily diffuse ( $R_d$  in MJ m<sup>-2</sup> day<sup>-1</sup>) and daily beam values ( $R_b$  in MJ m<sup>-2</sup> day<sup>-1</sup>). The model finally computes hourly beam and diffuse values on horizontal surfaces ( $r_b$  and  $r_d$ , both in MJ m<sup>-2</sup> h<sup>-1</sup>), from the daily amounts and following the temporal pattern of extraterrestrial hourly radiation during the day.

$$\frac{R_d}{R_g} = \begin{cases} 0.992 - 0.0486CI & CI \le 0.1\\ 0.954 + 0.734CI - 3.806CI^2 + 1.703CI^3 & 0.1 < CI \le 0.71\\ 0.165 & CI > 0.71 \end{cases}$$
(A5.1)

The second approach uses the temporal pattern of extraterrestrial hourly radiation,  $r_{ext}$ , to generate hourly global values,  $r_g$ according to previous studies (Chen et al., 1999; Ruiz-Arias et al., 2011). Then, the hourly regressive model developed by Ruiz-Arias et al. (2010) is applied to estimate the hourly diffuse values (Eq. A5.2) from the hourly CI, CI<sub>h</sub>, as the ratio of  $r_g$  to  $r_{ext}$ . This model was implemented as it has been validated over Europe and USA using ground data from different sites, including some Spanish stations (Ruiz-Arias et al., 2010). Hourly beam values ( $r_b$ ) are thus obtained on a cell basis as the difference between global and diffuse hourly radiation distributed values.

670

$$\frac{r_d}{r_g} = 0.952 - 1.041e^{-\exp(2.3 - 4.702 \cdot CI_h)}$$
(A5.2)

First applications at the study site have shown negligible differences between both partitioning schemes. The differences with daily recorded data were insignificant in the second decimal place of error values. Thus, the results presented in this study

- 680 were obtained with the original scheme of Aguilar et al. (2010) (Eq. A5.1) while the authors continue working on the
- 681 improvement on the partitioning scheme of the model.
- 682 Then, the topographic correction is carried out and depending on the component, different procedures are applied.



### 684 Figure A1. Flow chart of the solar radiation model

Hourly beam radiation on a surface of slope β and orientation γ, ( $r_{b,\beta\gamma}$  in MJ m<sup>-2</sup> h<sup>-1</sup>), is calculated according to Eq. A6. in terms of  $r_b$ ,  $\theta_z$  and a new corrected zenith angle for the sloping surface,  $\theta$  (Iqbal, 1983). Then, the model checks the shading effects. Self-shading will occur if the angle between the normal to the surface and the solar vector is greater than 90 degrees. Finally, shading by nearby terrain takes place when the illumination angle is greater than the horizon angle in the same direction. The model previously obtains the horizons following the algorithms of Dozier et al. (1981) and Dozier and Frew (1990), by comparing the slopes between cells in the eight directions.

$$r_{b,\beta\gamma} = r_b \left( \frac{\cos\theta}{\cos\theta_z} \right) \tag{A6}$$

Hourly diffuse radiation on a surface of slope β and orientation  $\gamma$  ( $r_{d,\beta\gamma}$  in MJ m<sup>-2</sup> h<sup>-1</sup>), is calculated according to Eq. A7 in terms of  $r_d$  and SVF, the sky view factor, that modifies the incoming radiation incident on a flat surface to consider possibly obstruction effects on a sloping surface (Dubayah, 1992). Dozier and Frew (1990) obtained an analytical expression for the estimation of the SVF in terms of the different horizons in each direction considered assuming an isotropic sky.

$$r_{d,\beta\gamma} = r_d \cdot SVF \tag{A7}$$

Finally, hourly reflected radiation on a surface of slope β and orientation  $\gamma$  ( $r_{r,\beta\gamma}$  in MJ m<sup>-2</sup> h<sup>-1</sup>) and albedo ρ is calculated according to Dozier and Frew (1990) as expressed in Eq. A8.

$$r_{r,\beta\gamma} = \rho \cdot \left[\frac{1+\cos\beta}{2} - SVF\right] \cdot \left(r_d + r_b\right)$$
(A8)

Hourly global distributed radiation ( $r_{gp}$  in MJ m<sup>-2</sup> h<sup>-1</sup>) is obtained by addition of the three hourly components at each cell according to Eq. A9.

$$r_{gp} = r_{b,\beta\gamma} + r_{d,\beta\gamma} + r_{r,\beta\gamma} \tag{A9}$$

Finally, daily global distributed radiation ( $R_{gp}$  in MJ m<sup>-2</sup> day<sup>-1</sup>) is obtained as the summation of hourly global distributed radiation values (Eq. A10).

$$R_{gp} = \sum_{24h} r_{gp} \tag{A10}$$

# 706 Appendix B: Cross-validation with deseasonalized data series and limited input data

691



 707
 observed
 observed
 observed
 observed

 708
 Figure B1. Linear fits of daily deseasonalized predicted vs. observed Rg (MJ m<sup>-2</sup> day<sup>-1</sup>) at each one of the selected stations for the global data (black), cloudy (CI<0.3 - red), partly cloudy (0.3<CI<0.6 - blue) and clear-sky days (CI>0.6 - orange). Stations are sorted by increasing altitude from left to right and from the top to the bottom row.



Figure B2. Linear fits of daily predicted with limited input data *vs.* observed  $R_g$  (MJ m<sup>-2</sup> day<sup>-1</sup>) at each one of the selected stations for the global data (black), cloudy (CI<0.3 - red), partly cloudy (0.3<CI<0.6 - blue) and clear-sky days (CI>0.6 - orange). Stations are sorted by increasing altitude from left to right and from the top to the bottom row.

712

# 718 Appendix C: Comparison with Solar Analyst estimates

Solar Analyst (SA) is one of the most used GIS-based solar radiation models. It calculates the insolation across a landscape or
 for specific locations, based on the methods developed by Fu and Rich (2000a, 2000b, 2002). The total amount of radiation is
 given as global radiation and depends on the latitude of the site, topography, shadow cast and atmospheric attenuation. Global

radiation is computed in SA as the sum of direct and diffuse radiation. The equations and modeling scheme can be found in

detail in Fu and Rich (2000b).

Daily global radiation time series were generated in the study site with the Points Solar Radiation tool of SA at each weather station. Then a cross validation on a leave-one-out process was applied. The main inputs to the model, the diffuse fraction, k, and the atmospheric transmittivity,  $\tau$ , were estimated in the study site from observed global radiation data following Batlles et al. (2008).

728 SA underestimated observed daily values with a mean  $\alpha$  value of 0.78 in the global datasets (Table C.1). In terms of the 729 cloudiness level, a general underestimation by SA was always seen on cloudy (CI < 0.3) and clear-sky days (CI > 0.6) with slopes 730 of the fits significantly lower than 1 (mean  $\alpha$  values of 0.42 and 0.74 respectively). In contrast, a slight overestimation with a 731 mean  $\alpha$  value of 1.03 was found on partly cloudy days (0.3 < CI < 0.6). As for RMSE values, the lowest mean values were always found for cloudy days (1.59 MJ m<sup>-2</sup> day<sup>-1</sup>), also lower than those obtained in Figure 4 (3.70 MJ m<sup>-2</sup> day<sup>-1</sup>). However, despite 732 733 the lower RMSE values the deviation from the 1:1 linear fit in cloudy days with SA estimates was significant (mean  $\alpha$  value 734 of 0.42 and R<sup>2</sup> value of 0.39 in Table B.1). The highest RMSE values with SA estimates were always found on partly cloudy 735 days with a mean value of 5.35 MJ m<sup>-2</sup> day<sup>-1</sup> followed by clear-sky days with a mean RMSE value of 3.37 MJ m<sup>-2</sup> day<sup>-1</sup>, both considerably higher than those obtained in subsection 3.4 (3.07 and 2.07 MJ  $m^{-2}$  dav<sup>-1</sup> respectively). 736

Table C.1. Model performance with Solar Analyst: slope ( $\alpha$ ) and r<sup>2</sup> of the linear fit between daily predicted *vs.* observed R<sub>g</sub> and RMSE (MJ m<sup>-2</sup> day<sup>-1</sup>)

Station		Global data			CI≤0.3			0.3< CI≤0.6			CI>0.6		
z(m) Code		α	$\mathbb{R}^2$	RMSE	α	$\mathbb{R}^2$	RMSE	α	$\mathbb{R}^2$	RMSE	α	$\mathbb{R}^2$	RMSE
781	602	0.77	0.69	4.05	0.27	0.43	1	0.96	0.33	5.44	0.74	0.8	3
942	608	0.73	0.74	3.78	0.23	0.46	1.06	0.86	0.54	4.56	0.7	0.81	2.97
950	601	0.73	0.73	3.74	0.3	0.38	1.12	0.9	0.53	4.47	0.7	0.79	3.2
975	853	0.74	0.76	3.59	0.23	0.27	1.02	0.92	0.6	4.27	0.71	0.82	2.86
1212	604	0.78	0.7	4.11	0.29	0.37	1.36	0.98	0.5	4.89	0.74	0.79	3.23
1332	803	0.75	0.71	4.07	0.38	0.5	1.27	0.99	0.46	4.89	0.72	0.8	3.2
1530	854	0.79	0.7	4.22	0.39	0.59	1.34	1	0.49	5.12	0.74	0.8	3.11
1732	857	0.85	0.61	4.84	0.43	0.39	1.69	1.1	0.23	6.23	0.8	0.74	3.38
1735	859	0.83	0.55	5.11	0.55	0.45	1.49	1.2	0.13	6.44	0.78	0.77	3.34
2141	804	0.67	0.68	4.02	0.24	0.17	0.95	0.97	0.5	4.72	0.64	0.79	3.21
2155	855	0.78	0.6	4.96	0.55	0.45	1.93	1.14	0.48	5.37	0.73	0.77	3.56
2300	858	0.65	0.62	4.69	0.54	0.39	2	0.87	0.41	5.55	0.61	0.69	4.07
2325	1002	0.81	0.54	4.98	0.61	0.51	2.15	1.17	0.14	6.23	0.76	0.7	3.64

2510	802	0.79	0.65	4.67	0.57	0.49	2.17	1.09	0.41	5.64	0.75	0.78	3.45
2867	1001	0.88	0.63	4.8	0.5	0.34	2.47	1.16	0.37	5.69	0.84	0.75	3.75
3097	860	0.85	0.6	4.95	0.58	0.08	2.45	1.22	0.24	6.1	0.81	0.74	3.87
Me	ean	0.78	0.66	4.41	0.42	0.39	1.59	1.03	0.40	5.35	0.74	0.77	3.37

# 741 Appendix D: Nomenclature

742	Symbols
743	CI: daily clearness index
744	CI <sub>CS</sub> : daily clearness index in a cloudless atmosphere
745	CI <sub>h</sub> : hourly clearness index
746	E <sub>o</sub> : eccentricity factor
747	fCI <sub>cl</sub> : cloudiness effects factor
748	I <sub>SC</sub> : solar constant
749	k: diffuse to global irradiance ratio
750	N CI<0.3: rate of days for cloudy conditions
751	N 0.3 <ci<0.6: cloudy="" conditions<="" days="" for="" of="" partially="" rate="" td=""></ci<0.6:>
752	N CI>0.6: rate of days for clear-sky conditions
753	$N_{\rm o}\!\!:$ number of initially available daily records in the study period
754	N: number of available daily records after the quality check
755	Q <sub>1</sub> : Quartile 1
756	Q <sub>3</sub> : Quartile 3
757	R <sub>b</sub> : daily beam radiation
758	$R_{bp}$ : daily beam radiation predicted by the model
759	R <sub>d</sub> : daily diffuse radiation
760	$R_{dp}$ : daily diffuse radiation predicted by the model
761	Rext: daily extraterrestrial radiation
762	R <sub>g</sub> : global radiation
763	R <sub>gcs</sub> : global radiation under clear-sky conditions
764	$R_{\text{go}\_max}$ : maximum daily global radiation observed value
765	$R_{\text{go\_mean}}$ : mean daily global radiation observed value
766	$R_{\text{go}\_min}$ : minimum daily global radiation observed value
767	$R_{\rm gp}$ : daily global radiation predicted by the model
768	r <sub>b</sub> : hourly beam radiation on horizontal surfaces

769	$r_{b,\beta\gamma}$ : hourly beam radiation on a surface of slope $\beta$ and orientation $\gamma$
770	r <sub>d</sub> : hourly diffuse radiation on horizontal surfaces
771	$r_{d,\beta\gamma}$ : hourly diffuse radiation on a surface of slope $\beta$ and orientation $\gamma$
772	rext: hourly extraterrestrial radiation
773	$r_{r,\beta\gamma}$ : hourly reflected radiation on a surface of slope $\beta$ and orientation $\gamma$
774	rg: hourly global radiation on horizontal surfaces
775	rgp: hourly global radiation predicted by the model
776	R <sup>2</sup> : coefficient of determination
777	T <sub>L</sub> : Linke turbidity factor
778	z: elevation
779	
780	Abbreviations
781	DEM: Digital Elevation Model
782	IDW: Inverse Distance Weighted
783	RMSE: Root Mean Square Error
784	SA: Solar Analyst
785	SN: Sierra Nevada mountain range
786	SVF: Sky view factor
787	UPH: Unsolved Problems in Hydrology
788	
789	Greek symbols
790	$\alpha$ : slope of the fit between $R_{gp}$ and $R_{go}$
791	β: slope
792	γ: orientation
793	$\mu_k$ : mean of the diffuse to global irradiance ratio
794	ρ: albedo
795	$\sigma_k$ : standard deviation of the diffuse to global irradiance ratio
796	$\theta$ : corrected zenith angle for the sloping surface
797	$\theta_z$ : zenith angle
798	